

Uncertainty product of the spherical Abel-Poisson wavelet

I. Iglewska-Nowak*

December 25, 2021

Abstract

In the present paper, the variances in the space and momentum domains of the spherical Abel-Poisson wavelet, as well as the limit of the uncertainty product for $\rho \rightarrow 0$, where ρ is the scale parameter, are computed. The values of these quantities coincide with a certain accuracy with those of the spherical Poisson wavelet with $\frac{1}{2}$ substituted for the order parameter.

Key Words and Phrases: uncertainty product, time-frequency localization, Abel-Poisson wavelet

2010 MSC: 42C40

1 Introduction

Similarly as in physics (Heisenberg's uncertainty principle), there exist several uncertainty principles in mathematics. An uncertainty constant of a function is a measure for the trade-off between the spatial and frequency localization. For functions over the sphere, the *uncertainty product* was introduced by Narcowich and Ward in [13], see also [6] for a short history of the notion and a discussion about various nomenclature.

The present research is devoted to spherical wavelets (for the theory of wavelet transforms see [4]). The uncertainty product can be interpreted as one of indicators of 'goodness' of wavelets, because well-localized wavelets yield a well-localized wavelet transform, i.e., such that 'fuzziness' of the transform reflects mostly the 'fuzziness' of the analyzed signal. Some general results concerning the uncertainty product of a wide class of wavelets in limit $\rho \rightarrow 0$, where ρ denotes the scale parameter, can be found in [6]. The case of Poisson wavelets g_ρ^m , $m \in \mathbb{N}$ [3, 7] – a wavelet family which is the most popular one for applications because of their explicit representation as well as existence of discrete frames

*West Pomeranian University of Technology, Szczecin, School of Mathematics, al. Piastów 17, 70-310 Szczecin, Poland

[11, 10] – is investigated in [8], and a bound for the uncertainty constant of the spherical Gauss-Weierstrass wavelet is computed in [9].

In the present paper we investigate another wavelet family, namely the Abel-Poisson wavelet. The so-called variances in the space and momentum domains are computed explicitly and exactly. The uncertainty product of a function is the square root of the variances. In the case of the Abel-Poisson wavelet, its limit for $\rho \rightarrow 0$ is finite. Up to a certain order, the values of the variances and the uncertainty product coincide with those for Poisson wavelets [8] with $\frac{1}{2}$ substituted for the order parameter m (Poisson wavelets are defined for $m \geq 1$). The paper is organized as follows. After an introduction of the necessary notions and statements in Section 2, the main result of the paper, Theorem 3.1 is proven in Section 3.

2 Preliminaries

Let \mathcal{S}^n denote the n -dimensional unit sphere in $(n+1)$ -dimensional Euclidean space \mathbb{R}^{n+1} with spherical variables $(\vartheta, \vartheta_2, \dots, \vartheta_{n-1}, \varphi)$. Integrable zonal (rotation-invariant with respect to the x_1 -axis) functions over the sphere have the Gegenbauer expansion

$$f(x) = \sum_{l=0}^{\infty} \hat{f}(l) C_l^\lambda(\cos \vartheta)$$

with Gegenbauer coefficients

$$\hat{f}(l) = c(l, \lambda) \int_{-1}^1 f(t) C_l^\lambda(t) (1-t^2)^{\lambda-1/2} dt,$$

where λ is an index related to the space dimension by

$$\lambda = \frac{n-1}{2}$$

and c is a constant that depends on l and λ . C_l^λ , $l \in \mathbb{N}_0$, are the Gegenbauer polynomials of order $\lambda \in \mathbb{R}$ and degree $l \in \mathbb{N}_0$. Unless it leads to misunderstandings, zonal spherical functions will be identified with those of the first variable, i.e., we write $f(x)$ interchangeable with $f(\cos \vartheta)$.

The variances in the space and momentum domains of a $\mathcal{C}^2(\mathcal{S}^n)$ -function f with $\int_{\mathcal{S}^n} x |f(x)|^2 d\sigma(x) \neq 0$ are given by [12]

$$\text{var}_S(f) = \left(\frac{\int_{\mathcal{S}^n} |f(x)|^2 d\sigma(x)}{\int_{\mathcal{S}^n} x |f(x)|^2 d\sigma(x)} \right)^2 - 1$$

and

$$\text{var}_M(f) = - \frac{\int_{\mathcal{S}^n} \Delta^* f(x) \cdot \bar{f}(x) d\sigma(x)}{\int_{\mathcal{S}^n} |f(x)|^2 d\sigma(x)},$$

where Δ^* is the Laplace–Beltrami operator on \mathcal{S}^n . The quantity

$$U(f) = \sqrt{\text{var}_S(f)} \cdot \sqrt{\text{var}_M(f)}$$

is called the uncertainty product of f .

The uncertainty product of zonal functions may be computed from their Gegenbauer coefficients [5, Lemma 4.2] and according to the spherical uncertainty principle it is bounded from below by $\frac{n}{2}$, see [13, 14] and [1, formula (4.37)], [2, formula (12)].

Lemma 2.1 *Let a zonal square integrable and continuously differentiable function over \mathcal{S}^n be given by its Gegenbauer expansion*

$$f(\cos \vartheta) = \sum_{l=0}^{\infty} \widehat{f}(l) \mathcal{C}_l^\lambda(\cos \vartheta).$$

Its variances in space and momentum domain are equal to

$$\text{var}_S(f) = \left(\frac{\sum_{l=0}^{\infty} \frac{\lambda}{l+\lambda} \binom{l+2\lambda-1}{l} |\widehat{f}(l)|^2}{\sum_{l=0}^{\infty} \binom{l+2\lambda}{l} \frac{\lambda^2 \overline{\widehat{f}(l)} \widehat{f}(l+1) + \widehat{f}(l) \overline{\widehat{f}(l+1)}}{(l+\lambda)(l+\lambda+1)}} \right)^2 - 1, \quad (1)$$

$$\text{var}_M(f) = \frac{\sum_{l=1}^{\infty} \frac{l\lambda(l+2\lambda)}{l+\lambda} \binom{l+2\lambda-1}{l} |\widehat{f}(l)|^2}{\sum_{l=0}^{\infty} \frac{\lambda}{l+\lambda} \binom{l+2\lambda-1}{l} |\widehat{f}(l)|^2}, \quad (2)$$

whenever the series are convergent.

Theorem 2.2 *For $f \in \mathcal{L}^2(\mathcal{S}^n) \cap \mathcal{C}^1(\mathcal{S}^n)$, $U(f) \geq \frac{n}{2}$.*

The Abel-Poisson wavelet (with respect to the measure $\alpha(\rho) = \frac{1}{\rho}$) is given by

$$\Psi_\rho^A(x) = \sum_{l=0}^{\infty} \frac{\lambda+l}{\lambda} \sqrt{2l\rho} e^{-l\rho} \mathcal{C}_l^\lambda(\cos \vartheta),$$

where $\rho \in \mathbb{R}_+$ denotes the scale parameter, see [4, Section 3].

3 The uncertainty product of the Abel-Poisson wavelet

Theorem 3.1 *For the Abel-Poisson wavelet, the variances in space and momentum domain are given by*

$$\begin{aligned} \text{var}_S(\Psi_\rho^A) = & \left\{ \rho^n + 2 \left[2(n-1)(1 - e^{-2\rho})^n \alpha(\rho) \rho^2 - (4n^2 - 6n + 3) \rho^n \right] e^{2\rho} \right. \\ & \left. - \left[4(n-1)(1 - e^{-2\rho})^n \alpha(\rho) \rho^2 + (4n-5) \rho^n \right] e^{4\rho} \right\}^{-2} \\ & \cdot \left\{ 16(n-1)^2 \left[n-1 + (n+1) e^{2\rho} \right]^2 \rho^{2n} e^{2\rho} \right\} - 1, \end{aligned} \quad (3)$$

where α is a bounded function, and

$$\text{var}_M(\Psi_\rho^A) = \frac{n(n+1)[n+(n+3)e^{2\rho} + e^{4\rho}]e^{2\rho}}{[n-1+(n+1)e^{2\rho}](e^{2\rho}-1)^2}. \quad (4)$$

The uncertainty product equals in limit $\rho \rightarrow 0$

$$\lim_{\rho \rightarrow 0} U(\Psi_\rho^A) = \frac{1}{2} \sqrt{\frac{(n+1)(n+2)(n^2-3n+3)}{n(n-1)}}. \quad (5)$$

Proof. Substituting

$$\widehat{\Psi}_\rho^A(l) = \frac{\lambda+l}{\lambda} \sqrt{2\rho l} e^{-\rho l}$$

to the expressions (1) and (2) we obtain

$$\begin{aligned} \text{var}_S(\Psi_\rho^A) &= \left(\frac{\sum_{l=1}^{\infty} \frac{l+\lambda}{\lambda} \binom{l+2\lambda-1}{l} l e^{-2\rho l}}{\sum_{l=1}^{\infty} \binom{l+2\lambda}{l} \cdot 2\sqrt{l(l+1)} e^{-\rho(2l+1)}} \right)^2 - 1, \\ \text{var}_M(\Psi_\rho^A) &= \frac{\sum_{l=1}^{\infty} \frac{l(l+\lambda)(l+2\lambda)}{\lambda} \binom{l+2\lambda-1}{l} l e^{-2\rho l}}{\sum_{l=1}^{\infty} \frac{l+\lambda}{\lambda} \binom{l+2\lambda-1}{l} l e^{-2\rho l}}. \end{aligned}$$

Set

$$S_{n,m}(\rho) = \sum_{l=1}^{\infty} \binom{l+2\lambda-1}{l} l^m e^{-2\rho l}.$$

Then,

$$\text{var}_S(\Psi_\rho^A) = \left[\frac{e^\rho \cdot A(\rho)}{B(\rho)} \right]^2 - 1, \quad (6)$$

$$\text{var}_M(\Psi_\rho^A) = \frac{C(\rho)}{A(\rho)} \quad (7)$$

for

$$A(\rho) = \frac{1}{\lambda} S_{n,2}(\rho) + S_{n,1}(\rho), \quad (8)$$

$$C(\rho) = \frac{1}{\lambda} S_{n,4}(\rho) + 3S_{n,3}(\rho) + 2\lambda S_{n,2}(\rho), \quad (9)$$

and

$$B(\rho) = \sum_{l=0}^{\infty} \frac{l+2\lambda}{\lambda} \binom{l+2\lambda-1}{l} \sqrt{l(l+1)} e^{-2\rho l}.$$

In order to estimate $B(\rho)$ by $S_{n,m}(\rho)$ note that for $l \in \mathbb{N}$

$$l + \frac{1}{2} - \frac{1}{8l} \leq \sqrt{l(l+1)} \leq l + \frac{1}{2} - \frac{1}{8l} + \frac{1}{16l^2}.$$

Thus,

$$\frac{l+2\lambda}{\lambda} \sqrt{l(l+1)} \geq \frac{l^2}{\lambda} + \left(\frac{1}{2\lambda} + 2\right)l + \left(1 - \frac{1}{8\lambda}\right) - \frac{1}{4l}$$

and

$$\begin{aligned} \frac{l+2\lambda}{\lambda} \sqrt{l(l+1)} &\leq \frac{l}{\lambda} \left(l + \frac{1}{2} - \frac{1}{8l} + \frac{1}{16l^2}\right) + 2 \left(l + \frac{1}{2\lambda}\right) \\ &= \frac{l^2}{\lambda} + \left(\frac{1}{2\lambda} + 2\right)l + \left(1 - \frac{1}{8\lambda}\right) + \frac{1}{16\lambda l}. \end{aligned}$$

Consequently,

$$B(\rho) = \frac{1}{\lambda} S_{n,2}(\rho) + \left(\frac{1}{2\lambda} + 2\right) S_{n,1}(\rho) + \left(1 - \frac{1}{8\lambda}\right) S_{n,0}(\rho) + R(\rho) \quad (10)$$

with

$$|R(\rho)| \leq \frac{1}{4} \cdot S_{n,-1}(\rho).$$

The values of $S_{n,1}$, $S_{n,2}$, $S_{n,3}$, and $S_{n,4}$ are obtained from

$$S_{n,0}(\rho) = \frac{1}{(1 - e^{-2\rho})^{n-1}} - 1, \quad (11)$$

compare the sixth equation on p. 352 in [8] (note that $S_{n,0}$ in [8] is defined as a series starting from $l = 0$), by the recurrence relation

$$S_{n,m+1}(\rho) = -\frac{1}{2} S'_{n,m}(\rho). \quad (12)$$

In order to estimate $S_{n,-1}(\rho)$ note that

$$S_{n,-1}(\rho) = \sum_{l=1}^{\infty} \left[P_{n-3}(l) + \frac{(n-2)!}{l} \right] e^{-2\rho l},$$

where P_{n-3} is a polynomial of degree $n-3$. Thus, $S_{n,-1}$ can be expressed as a linear combination of $S_{2,m}$, $m = -1, 0, \dots, n-3$,

$$S_{n,-1}(\rho) = \sum_{m=-1}^{n-3} \alpha_{n,m} \cdot S_{2,m}(\rho). \quad (13)$$

It follows from (11) and (12) that for $m \in \mathbb{N}_0$

$$S_{2,m}(\rho) = \frac{Q(n, e^{-2\rho})}{(1 - e^{-2\rho})^{m+1}}, \quad (14)$$

where Q is a polynomial. Further,

$$S_{2,-1}(\rho) = \sum_{l=1}^{\infty} \frac{e^{-2\rho l}}{l}.$$

Since the function $t \mapsto \frac{e^{-2\rho t}}{t}$ is monotonously decreasing on $(0, \infty)$,

$$\sum_{l=2}^{\infty} \frac{e^{-2\rho l}}{l} \leq \int_1^{\infty} \frac{e^{-2\rho t}}{t} dt.$$

Thus,

$$0 \leq S_{2,-1}(\rho) \leq e^{-2\rho} + \int_1^{\infty} \frac{e^{-2\rho t}}{t} dt = e^{-2\rho} + \Gamma(0, 2\rho). \quad (15)$$

It follows from (13), (14), (15), and

$$\frac{1}{(1 - e^{-2\rho})^m} = \mathcal{O}\left(\frac{1}{\rho^m}\right) \quad \text{for } \rho \rightarrow 0$$

that

$$S_{n,-1}(\rho) = \mathcal{O}\left(\frac{1}{\rho^{n-2}}\right) \quad \text{for } \rho \rightarrow 0.$$

Therefore, the rest term in (10) can be written as

$$R(\rho) = \frac{\alpha(\rho)}{\rho^{n-2}}$$

with a bounded function α . Formulae (3) and (4) are obtained from (6) and (7) by substituting (8), (10), and (9) with the series $S_{n,m}$, $m = 1, 2, 3, 4$, computed from (11) via (12). In the computation, $S_{n,0}$ is replaced by $(1 - e^{-2\rho})^{1-n}$. The difference between the two expressions can be absorbed by the rest term $R(\rho)$, and the replacement simplifies the calculations significantly.

(5) follows directly from (3) and (4). \square

The variances of the Abel-Poisson wavelet can be written as series

$$\begin{aligned} \text{var}_S(\Psi_\rho^A) &= \frac{n^2 - 3n + 3}{n(n-1)} \cdot \rho^2 - \frac{2n^2 - 4n + 3 + 2^{n+1} n(n-1) \alpha(\rho)}{n^2(n-1)} \cdot \rho^3 + \mathcal{O}(\rho^4), \\ \text{var}_M(\Psi_\rho^A) &= \frac{n^2 + 3n + 2}{4\rho^2} + \frac{n^2 - 1}{2n\rho} + \mathcal{O}(1) \end{aligned}$$

for $\rho \rightarrow 0$. Note that the first term of the variance in the space domain as well as the first two terms of the variance in the momentum domain are equal to those computed for Poisson wavelets [8, Theorem 3.1] with $\frac{1}{2}$ substituted for m . Consequently, the limit of $U(\Psi_\rho^A)$ for $\rho \rightarrow 0$ coincides with that for a Poisson wavelet with $m = \frac{1}{2}$. Note that formally the Abel-Poisson wavelet is the Poisson wavelet of order $\frac{1}{2}$. Therefore, the results concerning the variances and the uncertainty product are not surprising.

References

- [1] S. S. Goh and T. N. T. Goodman, *Uncertainty principles and asymptotic behavior*, Appl. Comput. Harmon. Anal. 16 (2004), no. 1, 19–43.

- [2] T. N. T. Goodman and S. S. Goh, *Uncertainty principles on circles and spheres*, Advances in constructive approximation: Vanderbilt 2003, 207–218, Mod. Methods Math., Nashboro Press, Brentwood, TN, 2004.
- [3] M. Holschneider and I. Igłowska–Nowak, *Poisson wavelets on the sphere*, J. Fourier Anal. Appl. 13 (2007), no. 4, 405–419.
- [4] I. Igłowska–Nowak, *Continuous wavelet transforms on n -dimensional spheres*, Appl. Comput. Harmon. Anal. 39 (2015), no. 2, 248–276.
- [5] I. Igłowska–Nowak, *Multiresolution on n -dimensional spheres*, Kyushu J. Math., *in press*.
- [6] I. Igłowska–Nowak, *On the uncertainty product of spherical wavelets*, Kyushu J. Math., *to appear*.
- [7] I. Igłowska–Nowak, *Poisson wavelets on n -dimensional spheres*, J. Fourier Anal. Appl. 21 (2015), no. 1, 206–227.
- [8] I. Igłowska–Nowak, *Uncertainty of Poisson wavelets*, Kyushu J. Math., *to appear*.
- [9] I. Igłowska–Nowak, *Uncertainty product of the spherical Gauss–Weierstrass wavelet*, preprint, https://www.researchgate.net/publication/321724086_Uncertainty_product_of_the_spherical_Gauss-Weierstrass_wavelet.
- [10] I. Igłowska–Nowak, *Semi-continuous and discrete wavelet frames on n -dimensional spheres*, Appl. Comput. Harmon. Anal. 40 (2016), no. 3, 529–552.
- [11] I. Igłowska–Nowak and M. Holschneider, *Frames of Poisson wavelets on the sphere*, Appl. Comput. Harmon. Anal. 28 (2010), no. 2, 227–248.
- [12] N. Laín Fernández, *Polynomial bases on the sphere*, doctoral thesis, Universität zu Lübeck 2003.
- [13] F. J. Narcowich and J. D. Ward, *Nonstationary wavelets on the m -sphere for scattered data*, Appl. Comput. Harmon. Anal. 3 (1996), no. 4, 324–336.
- [14] M. Rösler and M. Voit, *An uncertainty principle for ultraspherical expansions*, J. Math. Anal. Appl. 209 (1997), no. 2, 624–634.