

A NOTE ON BASIS PROBLEM IN NORMED SPACES

VINICIUS COELHO, JOILSON RIBEIRO, AND LUCIANA SALGADO

CONTENTS

1. Introduction and statements of main results	2
2. Applications	4
2.1. Bessaga-Pełczyński's Selection Principle	4
2.2. Banach problem for normed spaces	5
2.3. Spectral Theorem on inner product spaces	5
3. Problems	5
3.1. Organization of the text	6
4. Auxiliar Results	6
4.1. Essential Schauder basis	6
5. Proof of applications of main results	8
5.1. Proof of Bessaga-Pełczyński's Selection Principle	8
5.2. Proof of Banach problem for normed spaces	8
5.3. Proof of Spectral Theorem	9
6. Proofs of main results	11
6.1. The Banach-Grunblum's criterion for normed spaces	11
6.2. The Bessaga-Pełczyński Theorem for normed spaces	13
References	14

ABSTRACT. We prove the well known criterion of Banach-Grunblum and the Bessaga-Pełczyński Theorem for normed spaces, not necessarily complete (Banach) one. As application of these results, we show the Principle of Selection of Bessaga-Pełczyński for normed spaces and the Spectral Theorem for compact self-adjoint operators on inner product spaces.

Date: March 20, 2023.

V.C. is supported by CAPES, L.S. is partially supported by FAPERJ-Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro Projects APQ1-E-26/211.690/2021 SEI-260003/015270/2021 and JCNE-E-26/200.271/2023 SEI-260003/000640/2023, by CAPES — Finance Code 001, Fapesb-JCB0053/2013, PRODOC/UFBA 2014, CNPq and INCTMat-CAPES. V.C., J.R. and L.S. are grateful to Instituto de Matemática da Universidade Federal da Bahia, whose research facilities were most useful in preparation of this paper.

2010 Mathematics Subject Classification Primary: 46B15; Secondary: 46B20, 46B28. **Keywords:** Normed spaces, Schauder basis, Spectral theorem.

1. INTRODUCTION AND STATEMENTS OF MAIN RESULTS

In 1933, S. Banach affirmed that *every infinitely dimensional Banach space contains an infinite dimensional subspace with a basis*, without proof. In 1958, Bessaga and Pełczyński [3] developed several generalizations and modifications of Banach's claim, and proved their well known Selection Principle in this setting.

In this paper, we work on Banach's problem for normed (not necessarily complete) spaces, giving an expected generalization of the previous results, but not formally proved yet, as long as we know. We also generalize Banach-Grubblum's criterion in this context, and give a Spectral Theorem on inner product spaces.

For any vector space, it is well known that there exists an algebraic basis (or Hamel basis). However, there is another notion of basis, due to J. Schauder [14, 15], defined as follows:

Definition 1. A sequence $(x_n)_{n=1}^\infty$ in a normed space X is called a *Schauder basis* for X if for each x in X there is a unique sequence (a_n) of scalars such that $x = \sum_{i=1}^\infty a_n x_n$.

The uniqueness of the representation allows us to consider the linear operator for each n in \mathbb{N} :

$$x_n^* : X \rightarrow \mathbb{K}, x_n^* \left(\sum_{i=1}^\infty a_j x_j \right) = a_n,$$

this operators are called *coefficient operators* (or *coordinates operator*).

Let $(x_n)_{n=1}^\infty$ be a Schauder basis in the normed space $(X, \|\cdot\|)$, and consider the linear space $\mathcal{L}_X = \{(a_n)_{n=1}^\infty \mid \sum_{i=1}^\infty a_n x_n \text{ is convergent}\}$. A computation shows that the function

$\eta : \mathcal{L}_X \rightarrow \mathbb{R}$ given by $\eta((a_n)_{n=1}^\infty) := \sup \left\{ \left\| \sum_{i=1}^n a_i x_i \right\| : n \in \mathbb{N} \right\}$ is a norm in \mathcal{L}_X .

In this paper, our purpose is to provide the criterion of Banach-Grubblum and the Bessaga-Pełczyński Theorem for normed spaces. To deduce these results, we start with the following broad notion of Schauder basis.

Definition 2. Let X be a normed space and $(x_n)_{n=1}^\infty$ be a Schauder basis in X . We say that $(x_i)_{i=1}^\infty$ is an *essential Schauder basis* for X if $T_X : \mathcal{L}_X \rightarrow X$ given by $T_X((a_n)_{n=1}^\infty) = \sum_{n=1}^\infty a_n x_n$ is an isomorphism.

It is not difficult to show that in Banach spaces, every Schauder basis is an essential Schauder basis. This identification is important because with it all space with Schauder basis can be seen as a space of sequences. In the original definition of Schauder, there was the requirement that coordinates functional should be continuous. However, in 1932 Banach [2, pag 111] showed that in Banach spaces this is always true. But, if the space is not complete, this is false (see [5, Example 12.5].)

A sequence $(x_n)_{n=1}^\infty$ may not be a Schauder basis for a normed space X because $\overline{[x_n : n \in \mathbb{N}]}$ does not reach all the space X . In this case, we say that:

Definition 3. A sequence $(x_n)_{n=1}^\infty$ in a normed space X is a *basic sequence* if the sequence $(x_n)_n$ is a Schauder basis for $\overline{[x_n : n \in \mathbb{N}]}$.

In Banach spaces theory, we have the following practical and useful criterion for deciding whether a given sequence is basic or not.

Theorem 1.1. (*Banach-Grunblum's Criterion*) A sequence $(x_n)_{n=1}^\infty$ of non null vector in a Banach space X is a basic sequence if, and only if, there exists $M \geq 1$ such that for all sequence of scalar $(a_n)_{n=1}^\infty$:

$$(1.1) \quad \left\| \sum_{i=1}^m a_i x_i \right\| \leq M \left\| \sum_{i=1}^n a_i x_i \right\|$$

whenever $n \geq m$.

Proof. See e.g. [1, Proposition 1.1.9] or [4, Theorem 10.3.13]. \square

One of the implications of the above theorem was proved by S. Banach in 1932 and the other by M. Grunblum in 1941. We generalize this result for normed spaces. For this, it will be necessary to introduce the concept of essential basic sequence.

Definition 4. A sequence $(x_n)_{n=1}^\infty$ in a normed space X is an *essential basic sequence* if it is an essential Schauder basis for $\overline{[x_n : n \in \mathbb{N}]}$.

Note that if X is a Banach space, then every basic sequence is an essential basic sequence. Let $(S, \|\cdot\|)$ be a normed space, we denote by \widehat{S} the completion of S such that S is dense in \widehat{S} .

Theorem A. (*Banach-Grunblum's criterion for normed spaces*) Let $(x_n)_{n=1}^\infty$ be a sequence of non null vector in a normed space X . Then the following conditions are equivalents.

- (i) $(x_n)_{n=1}^\infty$ is an essential Schauder basis for $\widehat{[x_n : n \in \mathbb{N}]} \subseteq \widehat{X}$;
- (ii) $(x_n)_{n=1}^\infty$ is an essential basic sequence in X ;
- (iii) there exists $M \geq 1$ such that for all sequence of scalar $(a_n)_{n=1}^\infty$:

$$(1.2) \quad \left\| \sum_{i=1}^m a_i x_i \right\| \leq M \left\| \sum_{i=1}^n a_i x_i \right\|$$

whenever $n \geq m$.

As we said before, in his classic book [2], Banach announced without proof that in every Banach space of infinite dimension there is an infinite dimensional subspace with Schauder basis. The proof of this result only appeared in the literature in 1958 in a celebrated article by Bessaga and Pełczyński [3] (in that same year Bernard R. Gelbaum [10] also presented another proof). The demonstration presented in [3] is a consequence of the main result of their work, which became known as the *selection's principle of Bessaga-Pełczyński*.

Here, we show the Bessaga-Pełczyński's Theorem for normed spaces, and Selection Principle for normed spaces as an application. We need the definition of *equivalent sequence* for Essential Schauder's Basis.

Definition 5. Let $(x_n)_{n=1}^\infty$ be an essential Schauder's basis in a normed space X and $(y_n)_{n=1}^\infty$ be an essential Schauder's basis in a normed space Y . We say that $(x_n)_{n=1}^\infty$ is *equivalent* to $(y_n)_{n=1}^\infty$, and, in this case we write $(x_n)_{n=1}^\infty \approx (y_n)_{n=1}^\infty$, if, for any scalar's sequence the series $\sum_{n=1}^\infty a_n x_n$ is convergent in $\overline{[x_n : n \in \mathbb{N}]} \subseteq X$ if, and only if, the series $\sum_{n=1}^\infty a_n y_n$ is convergent in $\overline{[y_n : n \in \mathbb{N}]} \subseteq Y$.

The Bessaga-Pełczyński Theorem for normed spaces is the following result.

Theorem B. Let X be a normed space, $(x_n)_{n=1}^\infty$ be an essential basic sequence in X , and $(x_n^*)_{n=1}^\infty$ be the functional coefficients. If $(y_n)_{n=1}^\infty$ is a sequence in X such that

$$(1.3) \quad 0 < \sum_{n=1}^\infty \|x_n - y_n\| \cdot \|x_n^*\| =: \lambda < 1$$

then $(y_n)_{n=1}^\infty$ is an essential basic sequence in X equivalent to $(x_n)_{n=1}^\infty$.

2. APPLICATIONS

2.1. Bessaga-Pełczyński's Selection Principle. Using Theorem A and B, we obtain the Bessaga-Pełczyński's Selection Principle and this Corollary for normed spaces.

Definition 6. Let $(x_n)_{n=1}^\infty$ be an essential Schauder's basis in a normed space X , and $(k_n)_{n=0}^\infty$ be a sequence strictly increasing of positive integers, with $k_0 = 0$. A sequence of non null vectors $(y_n)_{n=1}^\infty$ in X is called *essential block basic sequence relative to $(x_n)_{n=1}^\infty$* if

$$y_n = \sum_{i=k_{n-1}+1}^{k_n} b_i x_i$$

where $b_i \in \mathbb{K}$.

The Bessaga-Pełczyński's Selection Principle is the following.

Theorem 2.1. Let $(x_n)_{n=1}^\infty$ be an essential Schauder basis in a normed space X and $(x_n^*)_{n=1}^\infty$ be the functional coefficients. If $(y_n)_{n=1}^\infty$ is a sequence in X such that $\inf_n \|y_n\| > 0$ and

$$\lim_{n \rightarrow \infty} x_i^*(y_n) = 0 \text{ for all } i \in \mathbb{N}$$

then $(y_n)_{n=1}^\infty$ contains an essential basic subsequence equivalent to block basic sequence relative to $(x_n)_{n=1}^\infty$.

Corollary 2.2. Let X be a normed space, $(y_n)_{n=1}^\infty$ is a sequence in X such that $\inf_n \|y_n\| > 0$ and $y_n \rightarrow 0$ weakly. Then $(y_n)_{n=1}^\infty$ contains an essential basic subsequence.

2.2. Banach problem for normed spaces. As mentioned before, the Bessaga and Pełczyński's demonstration that in every Banach space of infinite dimension there is an infinite-dimensional subspace with Schauder basis is a consequence of the nowadays known as the Bessaga-Pełczyński selection principle. As we have obtained this result for normed spaces, it is possible to follow the same steps and to show the same result. We stress that this is a already known result (see [6]). But here, the proof is done in an elementary way, using only the Banach-Grünblum's Criterion for normed spaces (Theorem A).

Theorem 2.3. *Every normed space contains an infinite-dimensional closed subspace with Schauder basis in which the canonical projections $(P_n)_n$ are bounded operators. Moreover, $\sup_n \|P_n\| < +\infty$.*

2.3. Spectral Theorem on inner product spaces. Let $T : N \rightarrow N$ be a compact self-adjoint operator, and $\widehat{T} : \widehat{N} \rightarrow \widehat{N}$ be the compact self-adjoint operator such that \widehat{T} is the bounded linear extension of T .

The following is the Spectral Theorem on inner product spaces.

Theorem 2.4. *Let N be an inner product space, $T : N \rightarrow N$ be a compact self-adjoint operator such that $T \neq 0$, $\text{sp}(T)$ be the spectrum of T , and $\text{sp}(\widehat{T})$ be the spectrum of \widehat{T} .*

- (i) *If $\text{sp}(T)$ is an infinite set such that $\text{sp}(T) \subsetneq \text{sp}(\widehat{T})$, then for each $x \in N$, there exists $w_x \in N$ such that $T(x) = \sum_{i=1}^{\infty} x_i \lambda_i \langle x, x_i \rangle + w_x$;*
- (ii) *If $\text{sp}(T)$ is an infinite set such that $\text{sp}(T) = \text{sp}(\widehat{T})$, then for each $x \in N$, $T(x) = \sum_{i=1}^{\infty} x_i \lambda_i \langle x, x_i \rangle$;*
- (iii) *If the cardinality of $\text{sp}(T)$ is a non-null natural number, then there exists a natural number $s \in \mathbb{N} \setminus \{0\}$ such that for all $x \in N$, $T(x) = \sum_{i=1}^s x_i \lambda_i \langle x, x_i \rangle$*

where (λ_i) 's are the eigenvalues of T .

3. PROBLEMS

It is known that all Banach space with Schauder basis is separable (see [13, Proposition 4.1.10]). Banach himself questioned about the reciprocal, that is, does every separable Banach space have a Schauder basis? In [9], Enflo showed that this is false, exhibiting a separable Banach space that has no Schauder basis.

Problem 1. *(The basis problem for normed spaces) Let X be a normed space that it is not a Banach space. Does every separable X have an essential Schauder basis?*

Definition 7. Let X be a Banach space, and $(x_n)_{n=1}^{\infty}$ a Schauder basis in X . The basis $(x_n)_{n=1}^{\infty}$ is an unconditional basis if, for each x in X , there exists a unique expansion of the form

$$x = \sum_{n=1}^{\infty} a_n x_n$$

where the sum converges unconditionally.

Gowers and Maurey in [11] showed that there exists a Banach space that do not contain unconditional basis.

Definition 8. Let X be a normed space, and $(x_n)_{n=1}^\infty$ an essential Schauder basis in X . The basis $(x_n)_{n=1}^\infty$ is an essential unconditional basis if, for each x in X , there exists a unique expansion of the form

$$x = \sum_{n=1}^{\infty} a_n x_n$$

where the sum converges unconditionally.

Problem 2. *(The unconditional basic sequence problem for normed spaces) Let X be a normed space that it is not a Banach space. Does every X have an essential unconditional basis?*

3.1. Organization of the text. In Sections 1, 2 and 3, we provided preliminary definitions in order to present the statements of the main results together with some applications and problems. In Section 4 we state some auxiliary results and prove some useful properties of Essential Schauder Basis. In Section 5, we give the proofs of applications of main results, divided into three subsections 5.1, 5.2 and 5.3, one for each of the applications. In Section 6, we give the proofs of our theorems, divided into two subsections 6.1 and 6.2, one for each of the Main Theorems A and B, respectively.

4. AUXILAR RESULTS

4.1. Essential Schauder basis. In the original definition of Schauder, there was the requirement that coordinates functional should be continuous. However, in [2, pag 111] is proved that this condition holds for Banach spaces. But, if the space is not complete, this assertion is false (see [5, Example 12.5]). We observe that if there exists an essential Schauder basis, then we recover this important property, as follows.

Theorem 4.1. *Each coordinate functional associated to an essential Schauder basis $(x_n)_{n=1}^\infty$ is a bounded linear application.*

Proof. Fix $n \in \mathbb{N}$, and let $x \in X$ be an arbitrary element, so we can write $x = \sum_{i=1}^{\infty} a_i x_i$. We are going to show that $x_n^* : X \rightarrow \mathbb{K}$ is a bounded linear application. In fact,

$$\begin{aligned} \|x_n\| \cdot |x_n^*(x)| &= \|x_n x_n^*(x)\| = \left\| \sum_{i=1}^n x_i^*(x) x_i - \sum_{i=1}^{n-1} x_i^*(x) x_i \right\| \\ &\leq \left\| \sum_{i=1}^n x_i^*(x) x_i \right\| + \left\| \sum_{i=1}^{n-1} x_i^*(x) x_i \right\| \leq 2\eta((a_j)_{j=1}^\infty) \leq 2 \|(T_X)^{-1}\| \cdot \|x\|. \end{aligned}$$

□

Corollary 4.2. *Let $(x_n)_{n=1}^\infty$ be an essential Schauder basis for the normed space X . Then, for each $n \in \mathbb{N}$, the linear operator*

$$P_n : X \rightarrow X, P_n \left(\sum_{i=1}^\infty a_i x_i \right) = \sum_{i=1}^n a_i x_i$$

is bounded.

Proof. Note that $P_n(\cdot) = \sum_{i=1}^n x_i^*(\cdot) x_i$, so by Theorem (4.1), P_n is a bounded operator. \square

In the proof of the next result, we will use some arguments from the proof of Corollary 4.1.17 of [13].

Theorem 4.3. *Let $(x_n)_{n=1}^\infty$ be an essential Schauder basis for a normed space X and $(P_n)_n^\infty$ the canonical projections. Then $\sup_n \|P_n\| < \infty$.*

Proof. For each x in X such that $\|x\| \leq 1$, we have that

$$\|P_n(x)\| = \left\| \sum_{i=1}^n a_i x_i \right\| \leq \sup_n \left\| \sum_{i=1}^n a_i x_i \right\| = \eta((a_j)_{j=1}^\infty) \leq \|(T_X)^{-1}\| \cdot \|x\| \leq \|(T_X)^{-1}\|.$$

So, $\|P_n\| \leq \|(T_X)^{-1}\|$ for all n in \mathbb{N} and $\sup_n \|P_n\| \leq \|(T_X)^{-1}\|$, and we are done. \square

The number $K_{(x_n)_{n=1}^\infty} := \sup_n \|P_n\|$ is called *essential constant of basis* $(x_n)_{n=1}^\infty$. Note that $\|P_n\| \geq 1$ for all n , then $K_{(x_n)_{n=1}^\infty} \geq 1$.

Corollary 4.4. *Let $(x_n)_{n=1}^\infty$ be an essential Schauder basis for a normed space X and $(x_n^*)_{n=1}^\infty$ be the coefficient operators. Then, for each $k \in \mathbb{N}$,*

$$(4.1) \quad 1 \leq \|x_k^*\| \cdot \|x_k\| \leq 2K_{(x_n)_{n=1}^\infty}$$

Proof. First, we note that

$$(4.2) \quad 1 = x_k^*(x_k) = |x_k^*(x_k)| \leq \|x_k^*\| \cdot \|x_k\|.$$

Now, let x be an arbitrary element of X such that $x \neq 0$, so

$$\begin{aligned} \|x_k\| \cdot |x_k^*(x)| &= \|x_k x_k^*(x)\| = \left\| \sum_{i=1}^k x_i^*(x) x_i - \sum_{i=1}^{k-1} x_i^*(x) x_i \right\| \\ &\leq \left\| \sum_{i=1}^k x_i^*(x) x_i \right\| + \left\| \sum_{i=1}^{k-1} x_i^*(x) x_i \right\| = \|P_k(x)\| + \|P_{k-1}(x)\| \\ &\leq \|P_k\| \cdot \|x\| + \|P_{k-1}(x)\| \cdot \|x\| \leq 2K_{(x_n)_{n=1}^\infty} \|x\|. \end{aligned}$$

This implies that $\|x_k\| \cdot \|x_k^*\| \leq 2K_{(x_n)_{n=1}^\infty}$. \square

5. PROOF OF APPLICATIONS OF MAIN RESULTS

5.1. Proof of Bessaga-Pełczyński's Selection Principle. Using Theorem A, it is possible to show the following result.

Corollary 5.1. *Let $(x_n)_{n=1}^\infty$ be an essential basic sequence in a normed space X . Then $K_{(x_n)_{n=1}^\infty} = \inf\{M : M \text{ satisfies (1.2)}\}$.*

We are going to provide the definitions and results needed to show the Bessaga-Pełczyński's Selection Principle for normed spaces.

Lemma 5.2. *Let $(x_n)_{n=1}^\infty$ be an essential Schauder's basis in a normed space X and $(y_n)_{n=1}^\infty$ be an essential block basic sequence relative to $(x_n)_{n=1}^\infty$. Then $(y_n)_{n=1}^\infty$ is an essential basic sequence in X and $K_{(y_n)_{n=1}^\infty} \leq K_{(x_n)_{n=1}^\infty}$.*

Proof. Use the Banach-Grunblum's criterion for normed spaces, Theorem A, and Corollary 5.1. \square

Follow the proof given in Theorem 4.3.19 of [13] for Banach spaces, and use Corollary 4.4, Lemma 5.2 and Theorem B to obtain the Bessaga-Pełczyński's selection principle for normed spaces, Theorem 2.1.

In the proof of Corollary 2.2, we use the arguments from the proof of Corollary 10.4.9 of [4].

Proof of Corollary 2.2. Consider the subspace $\overline{[y_n : n \in \mathbb{N}]}$ of X , that is separable. By Banach-Mazur's Theorem, $\overline{[y_n : n \in \mathbb{N}]}$ is isometrically isomorphic to a subspace of $C[0, 1]$, so there exists a linear isometry $T : \overline{[y_n : n \in \mathbb{N}]} \rightarrow C[0, 1]$ such that $\overline{[y_n : n \in \mathbb{N}]}$ and $T(\overline{[y_n : n \in \mathbb{N}]})$ are isometrically isomorphic. Observe that $\inf_n \|T(y_n)\| = \inf_n \|y_n\| > 0$. Let $(x_n)_{n=1}^\infty$ be a Schauder basis for $C[0, 1]$, and $(x_n^*)_n$ be the coefficient operators. By boundedness of operator T , since $(y_n)_n$ converges to 0 in the weak topology, we obtain that $(T(y_n))_n$ converges to 0 in the weak topology. So

$$\lim_{n \rightarrow \infty} x_k^*(T y_n) = 0 \text{ for all } k \text{ in } \mathbb{N}.$$

By Bessaga-Pełczyński's selection principle for normed spaces (Theorem 2.1), there exists an essential basic subsequence $(T(y_{n_k}))_{k=1}^\infty$ of $(T(y_n))_n$. Using that $T^{-1} : T(\overline{[y_n : n \in \mathbb{N}]}) \rightarrow \overline{[y_n : n \in \mathbb{N}]}$ is an isometric isomorphism, we have that $(y_{n_k})_{k=1}^\infty = (T^{-1}(T(y_{n_k})))_{k=1}^\infty$ is an essential basic sequence. This concludes the proof of Corollary 2.2. \square

5.2. Proof of Banach problem for normed spaces.

Proof of Theorem 2.3. We will use the same sequence $(x_n)_{n=1}^\infty$ obtained in the traditional proof (see e.g. [7, 8]). By Banach-Grunblum's criterion for normed spaces, Theorem A, we obtain that $(x_n)_{n=1}^\infty$ is an essential basic sequence. In particular, $(x_n)_{n=1}^\infty$ is an essential Schauder basis. Then by Theorem (4.3) the canonical projections $(P_n)_n$ are bounded and $\sup_n \|P_n\| < +\infty$. \square

5.3. Proof of Spectral Theorem. We begin by proving a consequence of Theorem A.

Lemma 5.3. *Let N be an inner product space, $S = \{x_n : n \in \mathbb{N}\}$ be an orthonormal set of N . Then*

(i) *For all $x \in N$, and any $n \in \mathbb{N}$, we have that $\sum_{i=1}^n |\langle x, x_i \rangle|^2 \leq \|x\|^2$ (and then*

$$\sum_{i=1}^{\infty} |\langle x, x_i \rangle|^2 \leq \|x\|^2);$$

(ii) *$(x_n)_{n=1}^{\infty}$ is an essential Schauder basis for $\widehat{[x_n : n \in \mathbb{N}]} \subseteq \widehat{N}$;*

(iii) *$(x_n)_{n=1}^{\infty}$ is an essential Schauder basis for $\overline{[x_n : n \in \mathbb{N}]} \subseteq N$.*

Proof. Just note that $0 \leq \langle x - \sum_{i=1}^n \langle x, x_i \rangle x_i, x - \sum_{i=1}^n \langle x, x_i \rangle x_i \rangle$ to prove item (i). To obtain items (ii) and (iii), by Banach-Grunblum's criterion for normed spaces, Theorem (A), we are reduced to prove that for all sequence of scalar $(a_n)_{n=1}^{\infty}$ we have that $\left\| \sum_{i=1}^m a_i x_i \right\| \leq \left\| \sum_{i=1}^n a_i x_i \right\|$ whenever $n \geq m$. But it is clear by orthonormality of S . \square

We need these auxiliar results.

Lemma 5.4. *Let N be an inner product space and $T : N \rightarrow N$ be a compact self-adjoint operator. Then $\widehat{T} : \widehat{N} \rightarrow \widehat{N}$ is a compact self-adjoint operator where \widehat{T} is the bounded linear extension of T and \widehat{N} is the completion of N such that N is dense in \widehat{N} .*

Proof. The proof is straightforward. \square

Lemma 5.5. *Let N be an inner product space and $U : N \rightarrow N$ be a compact self-adjoint operator. If U is not the null-operator, there exist $x \in N \setminus \{0\}$ and $\lambda \in \mathbb{R} \setminus \{0\}$ such that $U(x) = \lambda x$.*

Proof. Use the same arguments present in the proof of this result for compact self-adjoint operators on Hilbert spaces. \square

Proof of Theorem 2.4. By Lemma 5.4, $\widehat{T} : \widehat{N} \rightarrow \widehat{N}$ is a compact self-adjoint operator on a Hilbert space. There exist $\alpha_i \in \mathbb{R} \setminus \{0\}$ eigenvalues of \widehat{T} for i in a not empty subset A of \mathbb{N} such that $\ker(\widehat{T} - \alpha_i \widehat{I})$ is a finite dimensional subspace where $I : N \rightarrow N$ is given by $I(x) = x$ and \widehat{I} is the extension of I to \widehat{N} . For each $n \in A$, note that $N \cap \ker(\widehat{T} - \alpha_n \widehat{I}) = \ker(T - \alpha_n I) \subseteq N$, define $t_n := \dim \ker(T - \alpha_n I)$ and $\widehat{t}_n := \dim \ker(\widehat{T} - \alpha_n \widehat{I})$. Let be J be the subset of \mathbb{N} given by $\{n \in A : N \cap \ker(\widehat{T} - \alpha_n \widehat{I}) \neq \{0\}\}$, we have that $1 \leq t_n \leq \widehat{t}_n$. So for each $n \in J$ there exist $v_{n,1}, \dots, v_{n,t_n}$ such that $T(v_{n,j}) = \alpha_n v_{n,j}$ for all $j \in \{1, \dots, t_n\}$ and $\{v_{n,j} : 1 \leq j \leq t_n\}$ is an orthonormal set. Then $S = \{v_{n,j} : n \in J \text{ and } j \in \{1, \dots, t_n\}\}$ is an orthonormal set of N . We may write the orthonormal set S as $\{x_\ell \in N : \ell \in \mathcal{J}\}$ where \mathcal{J} is a subset of \mathbb{N} , and $T(x_\ell) = \lambda_\ell x_\ell$ for each $\ell \in \mathcal{J}$.

For each $n \in A$, there exists an orthonormal set $\{v_{n,t_n+1}, \dots, v_{n,\widehat{t}_n}\}$ of $\ker(\widehat{T} - \alpha_n \widehat{I})$, and then $\{v_{n,1}, \dots, v_{n,\widehat{t}_n}\}$ is an orthonormal set of $\ker(\widehat{T} - \alpha_n \widehat{I})$. Then $R = \{v_{n,j} : n \in A \text{ and } j \in \{1, \dots, \widehat{t}_n\}\}$ is an orthonormal set of \widehat{N} .

$j \in \{t_n + 1, \dots, \widehat{t}_n\}$ is an orthonormal set of \widehat{N} . We may write the orthonormal set R as $\{y_\ell \in N : \ell \in \widetilde{A}\}$ where \widetilde{A} is a subset of \mathbb{N} , and $\widehat{T}(y_\ell) = \beta_\ell y_\ell$ for each $\ell \in \widetilde{A}$.

Suppose that $\text{sp}(T)$ is an infinite set, then $\mathcal{J} = \mathbb{N}$. We are going to prove items (i) and (ii).

Let \widehat{F} be the completion of $F = [x_\ell : \ell \in \mathcal{J}]$, \widehat{G} be the completion of $G = [y_\ell : \ell \in \widetilde{A}]$, \widehat{E} be the completion of $E = \left[\bigcup_{n=1}^{\infty} \ker(\widehat{T} - \alpha_n \widehat{I}) \right] = [F \cup G]$. From the Spectral Theorem for Hilbert Spaces, we have that $\widehat{N} = \widehat{E} \oplus \ker(\widehat{T})$, and observe that $\widehat{E} = \widehat{F} \oplus \widehat{G}$.

Since S is an orthonormal set of N , from Lemma 5.3, we have that $(x_\ell)_{\ell=1}^{\infty}$ is an essential Schauder basis for \widehat{F} . For any $a \in \widehat{F}$, there exists a sequence of scalar $(\xi_n)_{n=1}^{\infty}$ such that $a = \sum_{i=1}^{\infty} \xi_i x_i$ where $\xi_i = \langle a, x_i \rangle$ for each $i \in \mathbb{N}$, so $\widehat{T}(a) = \sum_{i=1}^{\infty} x_i \lambda_i \langle a, x_i \rangle$.

Since R is an orthonormal set of \widehat{N} , from Lemma 5.3, we have that $(y_\ell)_{\ell \in \widetilde{A}}$ is an essential Schauder basis for \widehat{G} . For any $b \in \widehat{G}$, there exists a sequence of scalar $(\gamma_n)_{n \in \widetilde{A}}$ such that $b = \sum_{i \in \widetilde{A}} \gamma_i y_i$ where $\gamma_i = \langle b, y_i \rangle$ for each $i \in \widetilde{A}$, so $\widehat{T}(b) = \sum_{i \in \widetilde{A}} y_i \beta_i \langle b, y_i \rangle$.

We need the following result.

Lemma 5.6. $\widehat{T}(\widehat{F}) \subseteq N \cap \widehat{F}$.

Proof of Lemma 5.6. Let x be an arbitrary element of \widehat{F} . Since S is an orthonormal set of N , from Lemma 5.3, we have that $(x_\ell)_{\ell=1}^{\infty}$ is an essential Schauder basis for \widehat{F} . There exists a sequence of scalar $(a_n)_{n=1}^{\infty}$ such that $x = \sum_{i=1}^{\infty} a_i x_i$ where $a_i = \langle x, x_i \rangle$ for each $i \in \widetilde{J}$.

Note that $\widehat{T}(x) = \sum_{i=1}^{\infty} a_i \lambda_i x_i = \lim_{n \rightarrow \infty} \sum_{i=1}^n a_i \lambda_i x_i$, so $\widehat{T}(x) \in \widehat{F}$.

Consider the sequence $(y_n)_{n=1}^{\infty}$ where $y_n = \sum_{i=1}^n x_i a_i \in N$ for each $n \in \mathbb{N}$. Using Lemma 5.3, we see that $\|y_n\|^2 = \sum_{i=1}^n (a_i)^2 = \sum_{i=1}^n |\langle x, x_i \rangle|^2 \leq \|x\|^2$. Then $(y_n)_{n=1}^{\infty}$ is a bounded sequence of N , by compactness of T , there exists a subsequence $(T(y_{n_k}))_{n=1}^{\infty}$ that converges to some point $y \in N$. We have that y_n converges to x , and then $\widehat{T}(y_n)$ converges to $\widehat{T}(x)$, so $\widehat{T}(x) = y \in N$. This completes the proof of Lemma 5.6. \square

So for each $x \in \widehat{N}$ there exist $a_x \in \widehat{F}$, $b_x \in \widehat{G}$ and $c_x \in \ker(\widehat{T})$ such that $x = a_x + b_x + c_x$, and then $\widehat{T}(x) = \widehat{T}(a_x) + \widehat{T}(b_x)$ where $\widehat{T}(a_x) = \sum_{i=1}^{\infty} x_i \lambda_i \langle a_x, x_i \rangle$ and $\widehat{T}(b_x) = \sum_{i \in \widetilde{A}} y_i \beta_i \langle b_x, y_i \rangle$.

By Lemma 5.6, we get that $\widehat{T}(a_x) \in N$.

Now, suppose that $x \in N$, so $T(x) = \widehat{T}(x) \in N$ and $\widehat{T}(x) = \widehat{T}(a_x) + \widehat{T}(b_x)$ with $\widehat{T}(a_x) \in N$, then $\widehat{T}(b_x) \in N$. Note that $\langle a_x, x_i \rangle = \langle x, x_i \rangle$ and $\langle b_x, y_i \rangle = \langle x, y_i \rangle$. For $x \in N$, we

obtain that $T(x) = \sum_{i=1}^{\infty} x_i \lambda_i \langle x, x_i \rangle + \sum_{i \in \tilde{A}} y_i \beta_i \langle x, y_i \rangle$ where $\sum_{i=1}^{\infty} x_i \lambda_i \langle x, x_i \rangle, \sum_{i \in \tilde{A}} y_i \beta_i \langle x, y_i \rangle \in N$. This proves item (i).

Suppose that $\text{sp}(T) = \text{sp}(\widehat{T})$, this implies that $\tilde{A} = \emptyset$, and then $T(x) = \sum_{i=1}^{\infty} x_i \lambda_i \langle x, x_i \rangle$ for all $x \in N$. The item (ii) is proved.

It remains to show item (iii). Now, $\text{sp}(T)$ is a natural number k . This implies that $S = \{v_{n,j} : n \in J \text{ and } j \in \{1, \dots, t_n\}\}$ is a finite orthonormal set of N such that $T(v_{n,j}) = \alpha_n v_{n,j}$ for all $j \in \{1, \dots, t_n\}$ for each $n \in J$. We may write the orthonormal set S as $\{x_\ell \in N : \ell \in \{1, \dots, s\}\}$ for some $s \in \mathbb{N} \setminus \{0\}$ where $T(x_\ell) = \lambda_\ell x_\ell$ for each $\ell \in \{1, \dots, s\}$.

Note that $F = [x_\ell : \ell \in \{1, \dots, s\}]$ is a Banach space since F is a finite dimensional space. This implies that $N = F \oplus F^\perp$ where $F^\perp = \{w \in N : \langle w, z \rangle = 0 \text{ for all } z \in F\}$.

We claim that $F^\perp = \ker(T)$. A trivial verification shows that $\ker(T) \subseteq F^\perp$. Using that $T(F) \subseteq F$, we have that $T(F^\perp) \subseteq F^\perp$. Define $U = T|_{F^\perp} : F^\perp \rightarrow F^\perp$, and note that U is a compact self-adjoint operator. Suppose that F^\perp is not contained in $\ker(T)$, there exists $q \in F^\perp$ such that $T(q) \neq 0$, then $U(q) = T(q) \neq 0$ and U is not null-operator. By Lemma 5.5, there exist $\lambda \in \mathbb{R}$ and $p \in F^\perp \setminus \{0\}$ such that $T(p) = U(p) = \lambda p$, so $p \in F$. We obtain that $p \in F \cap F^\perp$, so $p = 0$. This contradiction shows that U is the null operator, and the claim is proved.

For $x \in N = F \oplus \ker(T)$, there exist $a_x \in F$ and $b_x \in \ker(T)$ such that $x = a_x + b_x$. Note that for $a \in F$, we have that $a = \sum_{i=1}^s \langle a, x_i \rangle x_i$ and $T(a) = \sum_{i=1}^s \langle a, x_i \rangle \lambda_i x_i$. We obtain that $T(x) = T(a_x) = \sum_{i=1}^s \langle x, x_i \rangle \lambda_i x_i$ since $\langle a_x, x_i \rangle = \langle x, x_i \rangle$ for all $i \in \{1, \dots, s\}$, and we are done. \square

6. PROOFS OF MAIN RESULTS

6.1. The Banach-Grunblum's criterion for normed spaces.

Proof of Theorem A. It is clear that (i) implies (ii). We are going to show that (ii) proves (iii).

Suppose that $(x_n)_{n=1}^\infty$ is an essential Schauder basis for $\overline{[x_n : n \in \mathbb{N}]}$. Now, consider the canonical projections $(P_n)_{n=1}^\infty$ in $E = \overline{[x_n : n \in \mathbb{N}]}$. By Theorem (4.3), we know that

$$1 \leq K_{(x_n)_{n=1}^\infty} = \sup_n \|P_n\| < \infty$$

Given a sequence of scalar $(a_n)_{n=1}^\infty$, if $n \geq m$, then

$$(6.1) \quad \left\| \sum_{i=1}^m a_i x_i \right\| = \left\| P_m \left(\sum_{i=1}^n a_i x_i \right) \right\| \leq \|P_m\| \cdot \left\| \sum_{i=1}^n a_i x_i \right\| \leq K_{(x_n)_{n=1}^\infty} \left\| \sum_{i=1}^n a_i x_i \right\|$$

We are reduced to proving (i) from (iii). Suppose that (1.2) holds for $M \geq 1$. A straightforward calculation shows that the set $\{x_\ell : \ell \in \mathbb{N}\}$ is linearly independent.

For each $n \in \mathbb{N}$ consider the linear functional given by

$$\varphi_n : [x_\ell : \ell \in \mathbb{N}] \rightarrow \mathbb{K}, \varphi_n \left(\sum_{i=1}^k a_i x_i \right) = a_n$$

and the linear operator

$$T_n : [x_\ell : \ell \in \mathbb{N}] \rightarrow [x_\ell : \ell \in \mathbb{N}], T_n \left(\sum_{i=1}^k a_i x_i \right) = \sum_{i=1}^n a_i x_i$$

defining $a_{k+1} = \dots = a_n = 0$ if necessary.

It is clear that φ_n is a bounded linear operator for all n .

By (1.2), we have that

$$\left\| T_n \left(\sum_{i=1}^k a_i x_i \right) \right\| = \left\| \sum_{i=1}^n a_i x_i \right\| \leq M \left\| \sum_{i=1}^k a_i x_i \right\|$$

and T_n is bounded with $\|T_n\| \leq M$.

There exists a bounded linear extension $\Phi_n : \widehat{F} \rightarrow \mathbb{K}$ such that $\Phi_n|_F = \varphi_n$ and $\|\Phi_n\| = \|\varphi_n\|$ where $F = [x_\ell : \ell \in \mathbb{N}]$.

Consider the bounded linear operator $T_n : F \rightarrow F$, so we may consider this bounded linear operator $T_n : [x_\ell : \ell \in \mathbb{N}] \rightarrow \widehat{F}$. There exists a bounded linear extension $R_n : \widehat{F} \rightarrow \widehat{F}$ such that $R_n|_F = T_n$ and $\|R_n\| = \|T_n\|$.

Note that $T_n(z) = \sum_{i=1}^n a_i x_i = \sum_{i=1}^n \varphi_i(z) x_i$ for all $z \in F$.

For all $x \in \widehat{F}$ we have that

$$(6.2) \quad R_n(x) = \sum_{i=1}^n \Phi_i(x) x_i.$$

In fact, let $x \in \widehat{F}$ be an arbitrary element, so $x = \lim_{k \rightarrow \infty} y_k$ where $y_k \in [x_\ell : \ell \in \mathbb{N}]$. But R_n is a bounded linear operator, so

$$R_n(x) = \lim_{k \rightarrow \infty} R_n(y_k) = \lim_{k \rightarrow \infty} \sum_{i=1}^n \varphi_i(y_k) x_i = \lim_{k \rightarrow \infty} \sum_{i=1}^n \Phi_i(y_k) x_i = \sum_{i=1}^n \Phi_i(x) x_i.$$

We obtain that $R_n(x) = \sum_{i=1}^n \Phi_i(x) x_i$ for all x in \widehat{F} . Now, given $x \in \widehat{F}$ and $\varepsilon > 0$, there exists $y = \sum_{j=1}^m a_j x_j \in [x_\ell : \ell \in \mathbb{N}]$ for some $m \geq 1$ such that $\|x - y\| < \varepsilon$. For $n > m$, we have

$$\begin{aligned}\|x - R_n(x)\| &\leq \|x - y\| + \|R_n(y) - y\| + \|R_n(x) - R_n(y)\| \\ &\leq \|x - y\| + \|y - y\| + \|R_n\| \cdot \|x - y\| \leq (1 + M)\varepsilon.\end{aligned}$$

Then $x = \lim_{n \rightarrow \infty} R_n(x)$. Using (6.2), we obtain

$$x = \lim_{n \rightarrow \infty} R_n(x) = \lim_{n \rightarrow \infty} \sum_{i=1}^n \Phi_i(x)x_i = \sum_{i=1}^{\infty} \Phi_i(x)x_i.$$

The uniqueness of the above representation is clear. So $(x_n)_{n=1}^{\infty}$ is a Schauder basis for \widehat{F} .

Given $x \in \widehat{F}$ with $x = \sum_{i=1}^{\infty} a_i x_i$. By (1.2), for each $n \in \mathbb{N}$ we have that

$$(6.3) \quad \left\| \sum_{i=1}^n a_i x_i \right\| \leq M \left\| \sum_{i=1}^{\infty} a_i x_i \right\| = M\|x\|.$$

Then

$$(6.4) \quad \sup_n \left\| \sum_{i=1}^n a_i x_i \right\| \leq M \left\| \sum_{i=1}^{\infty} a_i x_i \right\| = M\|x\|$$

and note that

$$(6.5) \quad \|x\| = \left\| \sum_{i=1}^{\infty} a_i x_i \right\| = \lim_{n \rightarrow \infty} \left\| \sum_{i=1}^n a_i x_i \right\| \leq \sup_n \left\| \sum_{i=1}^n a_i x_i \right\|.$$

Using (6.4) and (6.5), we obtain that $T_{\widehat{F}} : \mathcal{L}_{\widehat{F}} \rightarrow \widehat{F}$ given by $T_{\widehat{F}}((a_n)_{n=1}^{\infty}) = \sum_{n=1}^{\infty} a_n x_n$ is a linear isomorphism. Then $(x_n)_{n=1}^{\infty}$ is an essential Schauder basis for \widehat{F} , and we are done. \square

6.2. The Bessaga-Pełczyński Theorem for normed spaces.

Proof of Theorem B. Given a sequence $(a_n)_{n=1}^{\infty}$ of scalars,

$$\begin{aligned}\left\| \sum_{i=1}^n a_i (x_i - y_i) \right\| &= \left\| \sum_{i=1}^n x_i^* \left(\sum_{j=1}^n a_j x_j \right) (x_i - y_i) \right\| \leq \sum_{i=1}^n \left| x_i^* \left(\sum_{j=1}^n a_j x_j \right) \right| \cdot \|x_i - y_i\| \\ &\leq \left\| \sum_{j=1}^n a_j x_j \right\| \left(\sum_{i=1}^n \|x_i^*\| \cdot \|x_i - y_i\| \right) \leq \lambda \left\| \sum_{i=1}^n a_i x_i \right\|,\end{aligned}$$

we obtain that

$$\left\| \sum_{i=1}^n a_i x_i \right\| - \left\| \sum_{i=1}^n a_i y_i \right\| \leq \left\| \sum_{i=1}^n a_i x_i - \sum_{i=1}^n a_i y_i \right\| \leq \lambda \left\| \sum_{i=1}^n a_i x_i \right\|.$$

So

$$(6.6) \quad (1 - \lambda) \left\| \sum_{i=1}^n a_i x_i \right\| \leq \left\| \sum_{i=1}^n a_i y_i \right\| \leq (1 + \lambda) \left\| \sum_{i=1}^n a_i x_i \right\|$$

for all n in \mathbb{N} . But $(x_n)_{n=1}^\infty$ is an essential basic sequence in X . By Banach-Grunblum's criterion for normed spaces, Theorem (A), there exists $M \geq 1$ such that if $m \geq n$ then

$$(6.7) \quad \left\| \sum_{i=1}^n a_i y_i \right\| \leq (1 + \lambda) \left\| \sum_{i=1}^n a_i x_i \right\| \leq (1 + \lambda) M \left\| \sum_{i=1}^m a_i x_i \right\| \leq \frac{(1 + \lambda) M}{(1 - \lambda)} \left\| \sum_{i=1}^m a_i y_i \right\|.$$

From Banach-Grunblum's criterion for normed spaces, Theorem (A), $(y_n)_{n=1}^\infty$ is an essential basic sequence in X .

We are going to show that the series $\sum_{n=1}^\infty a_n y_n$ converges if the series $\sum_{n=1}^\infty a_n x_n$ is convergent.

Define $F = [x_n : n \in \mathbb{N}]$ and $Y = \overline{[y_n : n \in \mathbb{N}]}$. Consider $T : F \rightarrow Y$ the linear operator given by $T\left(\sum_{i=0}^k x_i a_i\right) = \sum_{i=0}^k y_i a_i$ for $x = \sum_{i=0}^k x_i a_i$ in F . Note that if $x \neq 0$, we

have that $\left\| T\left(\frac{x}{\|x\|}\right) \right\| = \frac{\|T(x)\|}{\|x\|} = \frac{\left\| \sum_{i=1}^k a_i y_i \right\|}{\left\| \sum_{i=1}^k a_i x_i \right\|}$ for $x = \sum_{i=0}^k x_i a_i$ in F , and by (6.6) we obtain that

$$\left\| T\left(\frac{x}{\|x\|}\right) \right\| \leq (1 + \lambda).$$

So $T : F \rightarrow Y$ is a bounded linear operator. We may consider T as the following bounded operator $T : \overline{F} \subseteq X \rightarrow Y$. There exists a bounded linear extension $\hat{T} : \hat{F} \rightarrow \hat{Y}$ such that $\hat{T}|_{\overline{[x_n : n \in \mathbb{N}]}} = T$ with $\|\hat{T}\| = \|T\|$.

Now, suppose that $x = \sum_{n=1}^\infty a_n x_n \in \overline{F} \subseteq X$ is convergent. This implies that

$$T(x) = \hat{T}(x) = \hat{T}\left(\lim_{n \rightarrow \infty} \sum_{i=1}^n a_i x_i\right) = \lim_{n \rightarrow \infty} \hat{T}\left(\sum_{i=1}^n a_i x_i\right) = \lim_{n \rightarrow \infty} \sum_{i=1}^n a_i y_i,$$

then the series $T(x) = \hat{T}(x) = \sum_{i=1}^\infty a_i y_i \in Y \subseteq X$ is convergent.

Analogously, we can show that if the $\sum_{i=1}^\infty a_i y_i$ is convergent, then $\sum_{i=1}^\infty a_i x_i$ is convergent. \square

REFERENCES

- [1] F. Albiac. N.J. Kalton. Topics in Banach Space Theory. *Graduate Texts in Mathematics*, 233. New York: Springer-Verlag, 2006.
- [2] S. Banach. Théorie des opérations linéaires. Monografie Matematyczne 1. Warsaw, 1932.

- [3] C. Bessaga; A. Pełczyński. On bases and unconditional convergence of series in Banach spaces. *Studia Mathematica* (1958). Volume: 17, Issue: 2, page 151-164.
- [4] G. Botelho, D. Pellegrino, E. Teixeira. Fundamentos de análise funcional. *Colecao textos universitarios*. Rio de Janeiro: SBM, 2015.
- [5] Swartz, C. Elementary Functional Analysis. *World Scientific*. Singapore, 2010.
- [6] M.M. Day. On the Basis Problem in Normed Spaces. *Proceedings of the American Mathematical Society*. Vol. 13, No. 4 (Aug., 1962), pp. 655-658.
- [7] M. Daws. Introduction to Bases in Banach Spaces, 2005.
<https://www1.maths.leeds.ac.uk/mdaws/pubs/bases.pdf>
- [8] J. Diestel. Sequences and Series in Banach Spaces. *Graduate texts in mathematics; 92* New York: Springer-Verlag, 1984.
- [9] P. Enflo. A counterexample to the approximation problem in Banach spaces. *Acta Mathematica*. 130 (1): 309-317, 1973.
- [10] B. R. Gelbaum. Notes on Banach spaces and bases. *An. Acad. Bras. Cienc.* 30 (1958), 29-36.
- [11] W.T. Gowers, B. Maurey. The unconditional basic sequence problem. arXiv:math/9205204, 1992.
- [12] C. W. McArthur. Developments in Schauder basis theory. *Bull. Amer. Math. Soc.* 78 (1972), 877-908.
- [13] R.E. Megginson. An introduction do Banach space theory. Springer, 1998.
- [14] J. Schauder. Theorie stetiger Abbildungen in Funktionalräumen. *Math. Z.*, 26, 47-65, 1927.
- [15] J. Schauder. Eine eigenschaft des haarschen orthogonalsystems. *Math. Z.*, 28, 317-320, 1928.

(V.C.) UNIVERSIDADE FEDERAL DO OESTE DA BAHIA, CENTRO MULTIDISCIPLINAR DE BOM JESUS DA LAPA, AV. MANOEL NOVAIS, 1064, CENTRO, 47600-000 - BOM JESUS DA LAPA-BA-BRAZIL
Email address: `viniciuscs@ufob.edu.br`

(J.R.) UNIVERSIDADE FEDERAL DA BAHIA, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, AV. ADHEMAR DE BARROS, S/N , ONDINA, 40170-110 - SALVADOR-BA-BRAZIL
Email address: `joilsonor@ufba.br`

(L.S.) UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, INSTITUTO DE MATEMÁTICA, AVENIDA ATHOS DA SILVEIRA RAMOS 149 CIDADE UNIVERSITÁRIA, P.O. BOX 68530, 21941-909 RIO DE JANEIRO-RJ-BRAZIL
Email address: `lsalgado@im.ufjr.br`, `lucianasalgado@ufrj.br`