

# Notes on Loewy series of centers of $p$ -blocks

Yoshihiro Otokita\*

## Abstract

The present paper describes some results on the Loewy series of the center of a modular group algebra in order to solve a problem on the number of irreducible ordinary characters. For instance, we prove that a  $p$ -block of a finite group has at least  $p+2$  characters if its defect group contains non-elementary center.

## 1 Introduction

In this paper we consider a problem on the number of irreducible ordinary characters of a finite group through some studies of the center of a modular group algebra.

Let  $G$  be a finite group and  $F$  an algebraically closed field of characteristic  $p > 0$ . For a  $p$ -block  $B$  of the group algebra  $FG$  with defect  $d$ , we denote by  $k(B)$  the number of associated irreducible ordinary characters. In [4] Héthelyi and Külshammer conjectured that:

$$2\sqrt{p-1} \leq k(B) \text{ unless } d = 0?$$

This is a refinement of a result by Maróti [10] (see also [9]) and is still unsolved in general. In modular representation theory it is known that  $k(B)$  is equal to the dimension of the center  $ZB$  of  $B$ . We accordingly examine the algebraic structure of  $ZB$  in order to solve the problem above.

In the next section we prove the following results.

We first determine the Loewy structure of  $ZB$  for cyclic defect groups. More precisely, it is shown that each codimension of the Loewy series of  $ZB$  is equal to 1 or the inertial index of  $B$ . Moreover we refer to a remark on  $ZB$  related to the problem above.

---

\*Department of Mathematics and Informatics, Chiba University, Chiba-shi 263-8522, Japan. E-mail: otokita@chiba-u.jp

The second subsection deals with bounds for the Loewy length  $LL(ZB)$ , that is, the nilpotency index of the Jacobson radical of  $ZB$ . Some papers [7, 8, 11, 12] have discussed its upper bounds: for example, Okuyama has stated  $LL(ZB) \leq p^d$ . The main purpose of this section is to give a lower bound for  $LL(ZB)$ . As an immediate corollary to this bound, we conclude that  $B$  has at least  $p + 2$  characters provided its defect group contains non-elementary center.

Throughout this paper we use the following notations.

We fix a defect group  $D$  of order  $p^d$ . Let  $N_G(D, b)$  be the inertial subgroup of a root  $b$  of  $B$  in  $N_G(D)$ . Then the inertial index of  $B$  is  $e(B) = |N_G(D, b) : DC_G(D)|$ . As usual  $l(B)$  denotes the number of irreducible Brauer characters associated to  $B$ . For a finite-dimensional algebra  $A$  over  $F$  with Jacobson radical  $J(A)$ , its Loewy length  $LL(A)$  is defined to be the smallest positive integer  $t$  such that  $J(A)^t = 0$ . For an integer  $n \geq 1$  we define  $c(n)$  as the codimension of  $J(A)^n$  in  $J(A)^{n-1}$ . Then  $c(n) \neq 0$  for any  $1 \leq n \leq LL(A)$ .

## 2 Results

### 2.1 Cyclic defect groups

In this section we study  $c(n)$  for  $ZB$ . It always holds  $c(1) = 1$  as  $ZB$  is local. Moreover it is clear that  $c(2) = 0$  if and only if  $d = 0$ . We next see the case of  $c(2) = 1$  in the proposition below. This is a corollary to [7, Proposition 8] since it is easy to check  $c(n) \leq 1$  for all  $3 \leq n$ .

**Proposition 1.** *If  $d > 0$ , then the following are equivalent:*

- (1)  $c(2) = 1$ ;
- (2)  $ZB$  is uniserial;
- (3)  $B$  is nilpotent and  $D$  is cyclic.

In the next theorem we determine all the codimensions  $c(n)$  for cyclic defect groups. In the proof below we use the algebra of fixed points associated to  $B$ . As  $N_G(D, b)$  acts on  $Z(D)$  we define

$$F[Z(D)]^{N_G(D, b)} = \{a \in F[Z(D)] ; t^{-1}at = a \text{ for } t \in N_G(D, b)\}.$$

The last part of this theorem is due to [5, Corollary 2.8].

**Theorem 2.** *If  $D$  is cyclic, then*

$$c(n) = \begin{cases} 1 & (n = 1) \\ e(B) & (n = 2) \\ 1 & (3 \leq n \leq LL(ZB)) \end{cases}$$

and

$$LL(ZB) = \frac{p^d - 1}{e(B)} + 1.$$

*Proof.* Put  $I = N_G(D, b)/C_G(D)$ . Since  $B$  is perfectly isometric to its Brauer correspondent in  $N_G(D)$  we may assume  $B = FH$  where  $H = D \rtimes I$  by [6]. Let  $\Gamma$  be the ideal of  $ZFH$  spanned by all class sums of defect 0. Then  $ZFH = FD^{N_G(D, b)} \oplus \Gamma$  (cf. [3, Theorem 1.1]). Since  $\Gamma$  is contained in the socle of  $FH$  we have  $\Gamma^2 = 0$ . Thus  $J(ZFH) = J(FD^{N_G(D, b)}) \oplus \Gamma$  and  $J(ZFH)^2 = J(FD^{N_G(D, b)})^2$ . As  $FD^{N_G(D, b)}$  is uniserial (see the proof of [5, Corollary 2.8]), it follows that  $c(n) = 1$  for  $3 \leq n \leq LL(ZB)$ . Therefore  $c(2) = k(B) - 1 - \{LL(ZB) - 2\} = e(B)$  by Dade's theory.  $\square$

For cyclic defect groups  $l(B) = e(B)$  and

$$k(B) = \frac{p^d - 1}{e(B)} + e(B) \geq 2\sqrt{p^d - 1}.$$

In general the relations between  $c(n)$ ,  $l(B)$  and  $e(B)$  are as yet unknown. However it is possible to answer the question in Introduction under the conditions below inspired by Theorem 2.

We now assume  $d > 0$  and one of the following holds:

- (i)  $e(B) \leq c(n)$  for some  $2 \leq n \leq LL(ZB)$ ;
- (ii)  $e(B) \leq l(B)$ .

Then we obtain  $2\sqrt{\lambda - 1} \leq k(B)$  where  $\lambda = LL(F[Z(D)])$  as follows:  
By [5, Corollary 2.7],

$$\frac{\lambda - 1}{e(B)} + 1 \leq LL(F[Z(D)]^{N_G(D, b)}) \leq LL(ZB).$$

It is known that  $l(B)$  is equal to the dimension of the Reynolds ideal of  $ZB$ . Since it is contained in the socle of  $ZB$ ,

$$k(B) \geq 1 + e(B) + \{(\lambda - 1)/e(B) - 1\} \geq 2\sqrt{\lambda - 1}$$

in either case.

If  $Z(D)$  has type  $(p^{a_1}, \dots, p^{a_r})$ , then  $\lambda = p^{a_1} + \dots + p^{a_r} - r + 1$ . Therefore we deduce  $2\sqrt{p - 1} \leq k(B)$  in the assumptions.

## 2.2 Lower bounds

In this subsection we consider a lower bound for  $LL(ZB)$ . By a result of Broué [1, Proposition (III) 1.1] there exists an ideal  $K$  of  $ZB$  such that  $ZB/K$  is isomorphic to  $F[Z(D)]^{N_G(D,b)}$ . Thereby  $LL(F[Z(D)]^{N_G(D,b)}) \leq LL(ZB)$ . In the next theorem we give a different lower bound for the left side from [5, Corollary 2.7].

**Theorem 3.** *Let  $p^m$  be the exponent of  $Z(D)$ . Then*

$$\frac{p^m + p - 2}{p - 1} \leq LL(F[Z(D)]^{N_G(D,b)}) \leq LL(ZB).$$

*Proof.* We fix an orbit  $\mathcal{O}$  of an element in  $Z(D)$  with maximal order by the action of  $N_G(D, b)$ . Remark that  $|\mathcal{O}|$  divides  $e(B)$  since  $DC_G(D)$  acts trivially on  $Z(D)$ , namely  $|\mathcal{O}| \neq 0$  in  $F$ . We put  $a = |\mathcal{O}|1 - \sum_{u \in \mathcal{O}} u$  where 1 is the unit in  $G$ . As  $a$  is contained in  $J(F[Z(D)]^{N_G(D,b)})$  it suffices to prove that  $a^t \neq 0$  where  $t = 1 + p + \dots + p^{m-1}$ . For  $0 \leq i \leq m-1$ ,

$$a^{p^i} = |\mathcal{O}|^{p^i} 1 - \sum_{u \in \mathcal{O}} u^{p^i} = |\mathcal{O}| 1 - \sum_{u \in \mathcal{O}} u^{p^i}$$

by Fermat's theorem. Hence each term of  $a^t = a \cdot a^p \cdots a^{p^{m-1}}$  has the form

$$(-1)^{|I|} |\mathcal{O}|^{m-|I|} \prod_{i \in I} (u_i)^{p^i}$$

where  $I \subseteq \{0, 1, \dots, m-1\}$  and  $u_i \in \mathcal{O}$ . Suppose now that  $\prod_{i \in I} (u_i)^{p^i} = 1$  for some  $I \neq \emptyset$ . Since the order of  $u_i$  is  $p^m$  for any  $i \in I$ , we may assume  $|I| \geq 2$ . If  $r = \min\{I\}$  and  $s = \min\{I - \{r\}\}$ , we obtain

$$1 = (1)^{p^{m-s}} = \left\{ \prod_{i \in I} (u_i)^{p^i} \right\}^{p^{m-s}} = \prod_{i \in I} (u_i)^{p^{m-s+i}} = (u_r)^{p^{m-s+r}} \neq 1,$$

a contradiction. Thus the coefficient of 1 in  $a^t$  is  $|\mathcal{O}|^m \neq 0$ . Therefore  $a^t \neq 0$  as claimed.  $\square$

In Theorem 3 it seems that  $Z(D)$  is generally not replaced with  $D$  itself. Suppose that  $B$  is a  $p$ -block with defect group

$$M_{p^d} = \langle x, y ; x^{p^{d-1}} = y^p = 1, y^{-1}xy = x^{1+p^{d-2}} \rangle$$

where  $d \geq 4$ . As is well known,  $M_{p^d}$  has exponent  $p^{d-1}$  and has cyclic center of order  $p^{d-2}$ . By [8, Proposition 10],

$$LL(ZB) = \frac{p^{d-2} - 1}{l(B)} + 1 \leq p^{d-2} < \frac{p^{d-1} + p - 2}{p - 1}.$$

Furthermore we note that Theorem 3 is clear in this case since  $l(B) \leq p - 1$  (cf. [13, Theorem 8.1, 8.8]).

As a corollary to this bound and [12, Proposition 2.2], we deduce:

**Corollary 4.** *Let  $p^m$  be the exponent of  $Z(D)$ . Then*

$$\frac{p^m + p - 2}{p - 1} \leq k(B) - l(B) + 1 \leq k(B).$$

*In particular,  $p + 2 \leq k(B)$  provided  $m \geq 2$ .*

At the end of this section we reconsider the previous remark.

Here assume  $d > 0$  and one of the following holds:

- (iii)  $Z(D)$  is not elementary abelian;
- (iv)  $e(B) \leq mr \cdot c(n)$  for some  $2 \leq n \leq LL(ZB)$ ;
- (v)  $e(B) \leq mr \cdot l(B)$

where  $p^m$  and  $r$  are the exponent and the rank of  $Z(D)$ , respectively. Then  $2\sqrt{p-1} \leq k(B)$  by Corollary 4 and the methods in (i) and (ii).

### 3 Examples

We introduce two examples by Brough-Schwabrow [2] and Héthelyi-Külshammer [4] in order to understand our theorems. In the following let  $P$  be a Sylow  $p$ -subgroup of  $G$ .

- (1) Assume  $p = 3$  and  $G = {}^2G_2(q)$  is the simple Ree group where  $q$  is a power of 3 at least 27. Then  $FG$  decomposes into two 3-blocks: the principal block  $B_0$  with  $q + 7$  characters and simple block  $B_1$  with the Steinberg character. By [2, Theorem 1.1], we have  $LL(ZB_0) = 3$ . In this case Theorem 3 indicates that  $Z(P)$  has exponent 3. In fact

$$Z(P) = \{x(0, 0, v) ; v \in \mathbb{F}_q\}$$

by the notations [2, page 59] and

$$x(0, 0, v)^3 = x(0, 0, 3v) = x(0, 0, 0).$$

- (2) We denote by  $k(G)$  and  $l(G)$  the numbers of irreducible ordinary and Brauer characters of  $G$ , respectively and apply Corollary 4 to  $P$  as a defect group of the principal block. Then it follows that  $p + 2 \leq k(G)$

when  $Z(P)$  is non-elementary, otherwise it is likely that  $k(G) - l(G) + 1$  is "small". Now assume  $p \equiv 23 \pmod{264}$ . Let  $X$  be a cyclic group of order  $x = (p - 1)/22$  and  $H$  the unique 2-fold covering group of the four-degree symmetric group. Then  $H \times X$  acts freely on an elementary abelian group  $P$  of order  $p^2$  and the Frobenius group  $G = P \rtimes (H \times X)$  satisfies

$$k(G) = \frac{p^2 - 1}{48x} + 8x = \frac{217}{264}p + \frac{25}{264}$$

as mentioned in [4, Remark (iii)]. Since  $P$  is a normal Sylow  $p$ -subgroup of  $G$ ,

$$l(G) = k(H \times X) = 8x = \frac{4}{11}p - \frac{4}{11}$$

and

$$k(G) - l(G) + 1 = \frac{11}{24}p + \frac{35}{24}.$$

## References

- [1] M. Broué, *Brauer coefficients of  $p$ -subgroups associated with a  $p$ -block of a finite group*, J. Algebra **56** (1979), 365–383.
- [2] J. Brough and I. Schwabrow, *On centres of 3-blocks of the Ree groups  ${}^2G_2(q)$* , J. Algebra **492** (2017), 57–73.
- [3] J. Brough and I. Schwabrow, *On the Loewy length of the center of a block with elementary abelian defect groups*, Comm. Algebra **46** (2018), 829–833.
- [4] L. Héthelyi and B. Külshammer, *On the number of conjugacy classes of a finite solvable group*, Bull. London Math. Soc. **32** (2000), 668–672.
- [5] S. Koshitani, B. Külshammer and B. Sambale, *On Loewy lengths of blocks*, Math. Proc. Cambridge Philos. Soc. **156** (2014), 555–570.
- [6] B. Külshammer, *Crossed products and blocks with normal defect groups*, Comm. Algebra **13** (1985), 147–168.
- [7] B. Külshammer, Y. Otokita and B. Sambale, *Loewy lengths of centers of blocks II*, to appear in Nagoya Math. J.
- [8] B. Külshammer and B. Sambale, *Loewy lengths of centers of blocks*, to appear in Q. J. Math.

- [9] G. Malle and A. Maróti, *On the number of  $p'$ -degree characters in a finite group*, Int. Math. Res. Not. IMRN 2016, 6118–6132.
- [10] A. Maróti, *A lower bound for the number of conjugacy classes of a finite group*, Adv. Math. **290** (2016), 1062–1078.
- [11] T. Okuyama, *On the radical of the center of a group algebra*, Hokkaido Math. J. **10** (1981), 406–408.
- [12] Y. Otokita, *Characterizations of blocks by Loewy lengths of their centers*, Proc. Amer. Math. Soc. **145** (2017), 3323–3329.
- [13] B. Sambale, *Blocks of finite groups and their invariants*, Springer Lecture Notes in Math. Vol. 2127, Springer–Verlag, Cham, 2014.