

Modular-value-based metrology with spin coherent pointers

Le Bin Ho^{1,*} and Yasushi Kondo^{1,2}

¹*Department of Physics, Kindai University, Higashi-Osaka, 577-8502, Japan*

²*Interdisciplinary Graduate School of Science and Engineering,
Kindai University, Higashi-Osaka, 577-8502, Japan*

(Dated: March 16, 2022)

Modular values are quantities that described by pre- and postselected states of quantum systems like weak values but are different from them: The associated interaction is not necessary to be weak. We discuss an optimal modular-value-based measurement with a spin coherent pointer: A quantum system is exposed to a field in which strength is to be estimated through its modular value. We consider two cases, with a two-dimensional and a higher-dimensional pointer, and evaluate the quantum Fisher information. The modular-value-based measurement has no merit in the former case, while its sensitivity can be enhanced in the latter case. We also consider the pointer under a phase-flip error. Our study should motivate researchers to apply the modular-value-based measurements for quantum metrology.

PACS numbers: 03.65.Ta, 06.20.-f, 42.50.Lc

Introduction.—In the concept of quantum sensing, a physical quantity on a small scale can be measured indirectly via a quantum system, a quantum property, or a quantum phenomenon [1]. The principle of measurements is (i) preparing quantum systems, hereafter called *sensors*, of which number is L , (ii) exposing to a field of which strength is to be measured for a period of t , and (iii) obtaining a state change before and after the exposure. The change is a measure of the strength of the field. This procedure is repeated T/t times in a total measurement time T . If the sensors are independent, the effective total measurement number N is given as $N = LT/t$. For fixed T and t , the uncertainty of the estimation is proportional to $1/\sqrt{L}$, which is known as the standard quantum limit or the shot noise limit [2–5]. However, if the uncertainty scales as $1/L$ then it is called the Heisenberg limit [6] which is a fundamental limit.

There have been a number of attempts to improve measurement protocols to overcome the standard quantum limit. One of the pioneering attempts is to employ squeezed states to reduce noises [7–9]. The entanglement is also a resource for defeating the standard quantum limit [3, 4, 9–12]. For example, Jones *et al.* have used *NOON* states [11, 12]. Zaiser *et al.* have claimed that the sensitivity can be significantly enhanced by using a quantum memory [13]. Matsuzaki *et al.* have proposed a protocol with a teleportation [14].

These studies focused on sensors exposed to a field. We, here, draw attention a method how the sensors are measured. One approach in this direction is employing a weak value [15–18]. Measurements with weak values had been expected to enhance the sensitivity [19–24], but it turned out that they can only reach the standard quantum limit [19–24]. The measurements with weak values, however, can be dramatically improved when sensors are entangled [25, 26]. Furthermore, by employing non-classical pointer states, the sensitivity can reach the

Heisenberg limit [20, 27, 28].

In this Letter, we discuss modular-value-based measurements [29–33] with spin- j coherent pointers [34, 35]. They are different from the weak-value-based ones and can allow arbitrary strength interactions between the pointers and sensors. The pointer may be considered as a *measurement device* for extracting the field information from the sensors. In order to evaluate the sensitivity, we focus on the quantum Fisher information contained in the Cramér-Rao inequality [6, 36–38]. Its maximum provides the lower bound on the sensitivity for measuring the field. In this Letter, L is fixed and we compare the Fisher information of the measurements under various conditions. We first examine a measurement with a $j = \frac{1}{2}$ (qubit) pointer and then move our attention to that with a $j \geq 1$ pointer. We also consider the case when the qubit pointer is under a phase-flip error [39].

Modular-value-based measurement.— We assume that the sensor exposed to a field is a qubit and that its initial state is $|\psi_i\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ which evolves to $|\psi_i(\omega)\rangle = \frac{1}{\sqrt{2}}(|0\rangle + e^{-i\omega t}|1\rangle)$ during a period t . ω is a measure of the field strength to be *estimated*. We prepare the pointer in a spin- j coherent state

$$|\xi\rangle = \sum_{m=-j}^j \sqrt{\binom{2j}{j+m}} \left(\cos \frac{\theta}{2}\right)^{j+m} \left(\sin \frac{\theta}{2} e^{-i\varphi}\right)^{j-m} |j, m\rangle \quad (1)$$

in spherical representation [34, 35], where $\theta \in [0, \pi]$ and $\varphi \in [0, 2\pi]$. $|j, m\rangle$ is the standard angular momentum basis, for a fixed j , $m \in [-j, j]$.

The interaction Hamiltonian between the sensor and the pointer is assumed to be $\hat{H} = g(\tau)\sigma_z \otimes |j, +j\rangle\langle j, +j|$. After the interaction during $\tau = 0 \sim \tau_0$, the joint state of the sensor-pointer yields

$$|\Psi(\omega)\rangle = \frac{1}{\sqrt{2}} \left(|0\rangle \otimes |\xi_+\rangle + e^{-i\omega t} |1\rangle \otimes |\xi_-\rangle \right), \quad (2)$$

where $|\xi_{\pm}\rangle = \exp(\mp ig|j, +j\rangle\langle j, +j|)|\xi\rangle$ and $g = \int_0^{\tau_0} g(\tau)d\tau$ is the coupling strength. After postselecting the sensor onto a final state $|\psi_f\rangle = \cos\frac{\phi}{2}|0\rangle + \sin\frac{\phi}{2}|1\rangle$, the normalized pointer state becomes

$$\begin{aligned} |\eta\rangle &= \langle\psi_f|\hat{U}|\psi_i(\omega)\rangle \otimes |\xi\rangle / \sqrt{p(\omega)} \\ &= \frac{\langle\psi_f|\psi_i(\omega)\rangle \left[\sum_{m=-j}^{j-1} c_{jm}|j, m\rangle + \left(\cos\frac{\theta}{2}\right)^{2j} (\sigma_z)_m |j, j\rangle \right]}{\sqrt{p(\omega)}}. \end{aligned} \quad (3)$$

Here, $p(\omega) = |\langle\psi_f|\hat{U}|\psi_i(\omega)\rangle \otimes |\xi\rangle|^2$ is the probability of successful postselection, where $\|\cdot\|$ is the *norm* of $(*)$,

$$c_{jm} = \sqrt{\binom{2j}{j+m}} \left(\cos\frac{\theta}{2}\right)^{j+m} \left(\sin\frac{\theta}{2}e^{-i\varphi}\right)^{j-m}, \text{ and}$$

$$(\sigma_z)_m = \frac{\langle\psi_f|e^{-ig\sigma_z}|\psi_i(\omega)\rangle}{\langle\psi_f|\psi_i(\omega)\rangle}$$

is ω dependent modular value of the observable σ_z . Note that $(\sigma_z)_m$ is independent of the choice of the initial pointer state $|\xi\rangle$. Throughout this Letter, we fix $g = \pi/2$. Under this condition, the modulus of the modular value, $|(\sigma_z)_m|$, becomes the minimum when $|\psi_f\rangle = |\psi_i\rangle$ ($\phi = \pi/2$) and does the maximum when $|\psi_f\rangle$ is orthogonal to $|\psi_i\rangle$ ($\phi = 3\pi/2$).

The uncertainty $\delta\omega$ in the estimation of ω after N independent measurements is defined by $\delta\omega = \sqrt{\langle(\omega - \bar{\omega})^2\rangle}$, where $\bar{\omega}$ is the average of the measured ω 's. Its minimum is determined by the Cramér-Rao lower bound as $\delta\omega \geq 1/\sqrt{NF}$ [6, 36–38]. F is the Fisher information and is defined by $F = \sum_k [1/p_k(\omega)] [dp_k(\omega)/d\omega]^2$, where $p_k(\omega)$ is the probability distribution for obtaining the k 'th experimental result. The maximization of F over all possible measurements leads to the quantum Fisher information Q [38, 40, 41]. For a pure quantum state $|\psi(\omega)\rangle$, Q is given by [40]

$$Q = 4 \left[\frac{d\langle\psi(\omega)|}{d\omega} \frac{d|\psi(\omega)\rangle}{d\omega} - \left| \frac{d\langle\psi(\omega)|}{d\omega} |\psi(\omega)\rangle \right|^2 \right]. \quad (4)$$

In order to obtain the best sensitivity, we have to maximize Q when N is fixed.

We define the quantum Fisher information of the sensor of which state is $|\psi_i(\omega)\rangle$ as Q_0 . By following Eq. (4), we obtain $Q_0 = t^2$. Q_0 is also obtained in the case of Ramsey sensing [2, 4]. Therefore, we call these measurement protocols which give $Q = t^2$ as the conventional measurement. Q_0 is employed as the reference when comparing the Fisher information in various measurements in this Letter.

In measurement protocols with postselection, Q is given by the sum of Q_m and F_p [19–21, 42]. Q_m is called the *measured quantum Fisher information* which is a product of the Fisher information of the final pointer state and the probability of successful postselection $p(\omega)$.

F_p is called the *postselected classical Fisher information* defined as [19, 21]

$$F_p = \frac{1}{p(\omega)[1-p(\omega)]} \left[\frac{dp(\omega)}{d\omega} \right]^2. \quad (5)$$

As shown in the Supplementary Material [43], $F_p = O(\omega^2)$ at $\omega t \ll 1$ for all j . Therefore, we can ignore F_p and we have to consider only Q_m . We rewrite the final pointer state Eq. (3) as

$$|\eta\rangle = \frac{1}{\sqrt{2}} \left(\cos\frac{\phi}{2} |\xi_+\rangle + \sin\frac{\phi}{2} e^{-i\omega t} |\xi_-\rangle \right) / \sqrt{p(\omega)} \quad (6)$$

and then substitute it into Eq. (4). We will evaluate Q_m in two cases of $j = 1/2$ and $j \geq 1$. We introduce $Q_m^{(j)}$ which denotes the measured quantum Fisher information with the spin- j coherent pointer in our measurement protocol.

Two-dimensional pointer.—We evaluate $Q_m^{(1/2)}$. According to the definition of Q_m , we substitute Eq. (6) into Eq. (4) and make a product with $p(\omega)$ [43], we obtain

$$Q_m^{(1/2)} = \frac{t^2 \sin^2 \theta \sin^2 \phi}{2(1 - \cos \theta \sin \phi)} + O(\omega^2). \quad (7)$$

Note that $Q_m^{(1/2)}$ depends on both the postselected state (via ϕ) and the initial pointer state (via θ), as shown in Fig. 1. We can immediately observe $Q_m^{(1/2)}/Q_0 \leq 1$. The equality is satisfied when $(\phi, \theta) = (\pi/2, 0)$ and $(3\pi/2, \pi)$. This implies that modular-value-based measurements, like weak-value-based ones [19–24], cannot overcome the standard quantum limit. There is no advantage of the modular-value amplification in the qubit-pointer case.




FIG. 1. (Color online) $Q_m^{(1/2)}/Q_0$ as a function of θ and ϕ .

Higher-dimensional pointer.—We consider $Q_m^{(j)}$ for $j \geq 1$. Detailed calculations are shown in the Supplementary Material [43]. We obtain $Q_m^{(j)}/Q_m^{(1/2)} \leq 2j$. The equality is satisfied when $(\phi, \theta) = (3\pi/2, 0)$. In Fig. 2, $Q_m^{(j)}/Q_m^{(1/2)}$ is shown as a function of θ for $j = 1, 3/2$, and 2. When




FIG. 2. (Color online) $Q_m^{(j)}/Q_m^{(1/2)}$ for $j = 1, 3/2$ and 2 at $\phi = 3\pi/2$ where $|(\sigma_z)_m|$ is maximum. Its maximum is $2j$ and can be achieved at $\theta = 0$. Inset: the same as the main figure at $\phi = \pi/2$ where $|(\sigma_z)_m|$ is minimum.

$\phi = 3\pi/2$ ($|(\sigma_z)_m|$ is maximum), $Q_m^{(j)}/Q_m^{(1/2)}$ can be larger than 1. On the other hand, its maximum is 1 when $\phi = \pi/2$ ($|(\sigma_z)_m|$ is minimum). There is an advantage of the modular-value amplification in the case of $j \geq 1$.

Suppose that we have $L + 1$ qubits as a resource for measurement. We may employ them as L independent sensors and one qubit pointer. In this case, $\delta\omega$ scales $1/\sqrt{L}$ which corresponds to the standard quantum limit. In order to measure all sensors, we have to repeat measurements L times. On the other hand, we can employ them as one qubit sensor and a $j \geq 1$ pointer formed with L qubits. Note that $j = L/2$ [44]. Therefore, we expect the enhancement of the Fisher information as $Q_m^{(j)}/Q_m^{(1/2)} = L$ according to the above discussion. If we are allowed to measure L times as in the previous case, we may be able to expect another L enhancement in the Fisher information. In total, the enhancement of the Fisher information can scale L^2 . It implies $\delta\omega$ scales $1/L$. We claim that our modular-value-based measurement with a spin- j coherent pointer can approach the Heisenberg limit.

Measurements under noise.—So far, we assumed the ideal noiseless environments. Let us consider measurements under noisy environments where phase-flip errors occur on the pointer. The influence of the noise is described by the operator-sum representation as [39, 45],

$$\rho' = (1 - \nu)\rho + \nu\sigma_z\rho\sigma_z, \quad (8)$$

where ρ' is the density matrix of the pointer under the noise, ρ is that in the noiseless environment, and $0 < \nu < 1$ is the probability of the phase-flip. When $\nu = 1/2$, we totally lose the information of ω [45]. Hereafter, we only discuss a modular-value-based measurement with a qubit-pointer ($j = 1/2$) as a concrete example of noisy measurements.

By using the symmetric logarithmic derivative (SLD) operators defined by $\hat{L}_\lambda\rho + \rho\hat{L}_\lambda = 2\partial_\lambda\rho$ [38, 46], the




FIG. 3. (Color online) $(\nu - \nu^2) \nu Q_m^{(1/2)}$ as a function of θ and ϕ .

quantum Fisher information matrix ($H_{\alpha,\beta}$) is defined by [46–49]

$$H_{\alpha,\beta} = p(\omega) \text{Tr} \left[\rho' \frac{\hat{L}_\alpha \hat{L}_\beta + \hat{L}_\beta \hat{L}_\alpha}{2} \right], \quad (9)$$

where $\alpha, \beta = \omega$ or ν . $H_{\omega,\omega}$ is the quantum Fisher information associated with the estimation of ω and named as $\omega Q_m^{(1/2)}$, while $H_{\nu,\nu}$ is that with the estimation of ν and named as $\nu Q_m^{(1/2)}$. For $j = 1/2$, the final density matrix of the pointer is given by $\rho = |\eta\rangle\langle\eta|$, where $|\eta\rangle$ is given by Eq. (6). Then, we obtain

$$H = \begin{pmatrix} \frac{(1-2\nu)^2 t^2 \sin^2 \theta \sin^2 \phi}{2(1-\sin \theta \sin \phi)} & 0 \\ 0 & \frac{\sin^2 \theta \cos^2 \phi}{2(\nu-\nu^2)(1-\cos \theta \sin \phi)} \end{pmatrix} + O(\omega^2). \quad (10)$$

We observe $\omega Q_m^{(1/2)} = (1 - 2\nu)^2 Q_m^{(1/2)}$: This fact clearly illustrates that the noise degrades the measurements and that the θ - and ϕ -dependencies are inherited from $Q_m^{(1/2)}$ to $\omega Q_m^{(1/2)}$. It is in agreement with the general discussion under noise [28], too.

We show $(\nu - \nu^2) \nu Q_m^{(1/2)}$ as a function of θ and ϕ in Fig. 3. It is interesting to note that $\nu Q_m^{(1/2)} = 0$ when $\phi = \pi/2$ and $3\pi/2$ and that these ϕ 's give the maxima of $\omega Q_m^{(1/2)}$. Even more interesting, the parameter combinations of $\theta = \pi/2$ and $\phi = n\pi$ (n : integer) give the maxima of $\nu Q_m^{(1/2)}$ at the cost of $\omega Q_m^{(1/2)} = 0$. It implies that we can optimize the measurement of ν , or noise, by selecting θ and ϕ properly. It is very important to measure noise since noise often causes relaxation of quantum systems.

Conclusions.—We have investigated the modular-value-based measurements with spin- j coherent pointers. We discussed two cases with $j = 1/2$ and $j \geq 1$. From the viewpoint of quantum Fisher information, we first showed the modular-value-based measurements with a qubit pointer ($j = 1/2$) has no merit in the sensitivity enhancement as in a weak-value-based measurement with

a zero-mean Gaussian pointer. In contrast, the measurements with $j \geq 1$, the quantum Fisher information can become $2j$ times that of the qubit pointer. If L qubits consist a spin coherent pointer, the Fisher information in one measurement scales as L . By taking into account the time required for measurements, we claimed that the total Fisher information scales as L^2 and thus we have a chance to approach the Heisenberg limit. Our study in the presence of phase-flip errors shows that we can optimize a measurement for detecting noise.

Acknowledgments.—This work was supported by CREST(JPMJCR1774), JST.

* Electronic address: binho@kindai.ac.jp

- [1] C. L. Degen, F. Reinhard, and P. Cappellaro, *Rev. Mod. Phys.* **89**, 035002 (2017).
- [2] Y. Matsuzaki, S. C. Benjamin, and J. Fitzsimons, *Phys. Rev. A* **84**, 012103 (2011).
- [3] L. Pezzé and A. Smerzi, *Phys. Rev. Lett.* **102**, 100401 (2009).
- [4] S. F. Huelga, C. Macchiavello, T. Pellizzari, A. K. Ekert, M. B. Plenio, and J. I. Cirac, *Phys. Rev. Lett.* **79**, 3865 (1997).
- [5] W. M. Itano, J. C. Bergquist, J. J. Bollinger, J. M. Gilligan, D. J. Heinzen, F. L. Moore, M. G. Raizen, and D. J. Wineland, *Phys. Rev. A* **47**, 3554 (1993).
- [6] M. Zwierz, C. A. Pérez-Delgado, and P. Kok, *Phys. Rev. Lett.* **105**, 180402 (2010).
- [7] D. J. Wineland, J. J. Bollinger, W. M. Itano, F. L. Moore, and D. J. Heinzen, *Phys. Rev. A* **46**, R6797 (1992).
- [8] D. J. Wineland, J. J. Bollinger, W. M. Itano, and D. J. Heinzen, *Phys. Rev. A* **50**, 67 (1994).
- [9] V. Giovannetti, S. Lloyd, and L. Maccone, *Science* **306**, 1330 (2004).
- [10] V. Giovannetti, S. Lloyd, and L. Maccone, *Phys. Rev. Lett.* **96**, 010401 (2006).
- [11] J. A. Jones, S. D. Karlen, J. Fitzsimons, A. Ardavan, S. C. Benjamin, G. A. D. Briggs, and J. J. L. Morton, *Science* **324**, 1166 (2009).
- [12] S. Simmons, J. A. Jones, S. D. Karlen, A. Ardavan, and J. J. L. Morton, *Phys. Rev. A* **82**, 022330 (2010).
- [13] S. Zaiser, T. Rendler, I. Jakobi, T. Wolf, L. Sang-Yun, S. Wagner, V. Bergholm, T. Schulte-Herbrüggen, P. Neumann, and J. Wrachtrup, *Nat. Comm.* **7**, 12279 (2016).
- [14] Y. Matsuzaki, S. Benjamin, S. Nakayama, S. Saito, and W. J. Munro, *Phys. Rev. Lett.* **120**, 140501 (2018).
- [15] Y. Aharonov, D. Z. Albert, and L. Vaidman, *Phys. Rev. Lett.* **60**, 1351 (1988).
- [16] Y. Aharonov and D. Rohrlich, *Quantum Paradoxes: Quantum Theory for the Perplexed* (Wiley-VCH, 2005).
- [17] A. G. Kofman, S. Ashhab, and F. Nori, *Phys. Rep.* **520**, 43 (2012).
- [18] J. Dressel, M. Malik, F. M. Miatto, A. N. Jordan, and R. W. Boyd, *Rev. Mod. Phys.* **86**, 307 (2014).
- [19] G. B. Alves, A. Pimentel, M. Hor-Meyll, S. P. Walborn, L. Davidovich, and R. L. deMatos Filho, *Phys. Rev. A* **95**, 012104 (2017).
- [20] L. Zhang, A. Datta, and I. A. Walmsley, *Phys. Rev. Lett.* **114**, 210801 (2015).
- [21] G. B. Alves, B. M. Escher, R. L. de Matos Filho, N. Zagury, and L. Davidovich, *Phys. Rev. A* **91**, 062107 (2015).
- [22] S. Tanaka and N. Yamamoto, *Phys. Rev. A* **88**, 042116 (2013).
- [23] G. C. Knee, G. A. D. Briggs, S. C. Benjamin, and E. M. Gauger, *Phys. Rev. A* **87**, 012115 (2013).
- [24] G. C. Knee and E. M. Gauger, *Phys. Rev. X* **4**, 011032 (2014).
- [25] S. Pang, J. Dressel, and T. A. Brun, *Phys. Rev. Lett.* **113**, 030401 (2014).
- [26] S. Pang and T. A. Brun, *Phys. Rev. A* **92**, 012120 (2015).
- [27] S. Pang and T. A. Brun, *Phys. Rev. Lett.* **115**, 120401 (2015).
- [28] A. N. Jordan, J. Tollaksen, J. E. Troupe, J. Dressel, and Y. Aharonov, *Quantum Studies: Mathematics and Foundations* **2**, 5 (2015).
- [29] Y. Kedem and L. Vaidman, *Phys. Rev. Lett.* **105**, 230401 (2010).
- [30] L. B. Ho and N. Imoto, *Phys. Lett. A* **380**, 2129 (2016).
- [31] L. B. Ho and N. Imoto, *Phys. Rev. A* **95**, 032135 (2017).
- [32] L. B. Ho and N. Imoto, *J. Math. Phys.* **59**, 042107 (2018).
- [33] L. B. Ho and N. Imoto, *Phys. Rev. A* **97**, 012112 (2018).
- [34] J. P. Amiet and M. B. Cilibs, *J. Phys. A: Math. Gen.* **24**, 1515 (1991).
- [35] F. Bohnet-Waldraff, D. Braun, and O. Giraud, *Phys. Rev. A* **93**, 012104 (2016).
- [36] C. R. Rao, *Linear Statistical Inference and its Applications*, 2nd ed. (Wiley, New York, 1973).
- [37] H. F. Hofmann, M. E. Goggin, M. P. Almeida, and M. Barbieri, *Phys. Rev. A* **86**, 040102 (2012).
- [38] S. L. Braunstein and C. M. Caves, *Phys. Rev. Lett.* **72**, 3439 (1994).
- [39] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, (2000).
- [40] C. W. Helstrom, *Quantum Detection and Estimation Theory* (Academic Press, New York, 1976).
- [41] A. S. Holevo, *Probabilistic and Statistical Aspects of Quantum Theory* (North-Holland, Amsterdam, 1982).
- [42] J. Combes, C. Ferrie, Z. Jiang, and C. M. Caves, *Phys. Rev. A* **89**, 052117 (2014).
- [43] See supplementary material for the detailed calculation of the measured quantum Fisher information.
- [44] O. Giraud, D. Braun, D. Baguette, T. Bastin, and J. Martin, *Phys. Rev. Lett.* **114**, 080401 (2015).
- [45] E. Desurvire, *Classical and Quantum Information Theory An Introduction for the Telecom Scientist*, Cambridge University Press, (2009).
- [46] M. G. A. Paris, *Int. J. Quantum Inf.* **7**, 125 (2009).
- [47] A. Fujiwara and H. Nagaoka, *Phys. Lett. A* **201**, 119 (1995).
- [48] K. Matsumoto, *J. Phys. A* **35**, 3111 (2002).
- [49] C. W. Helstrom, *Phys. Lett. A* **25**, 101 (1967).

Supplementary Material for “Modular-value-based metrology with spin coherent pointers”

Quantum Fisher information in the case of $j = 1/2$: $Q_m^{(1/2)}$

We show the detailed calculation of the measured quantum Fisher information in the case of modular-value-based measurement with the spin- j coherent state when $j = 1/2$. Let remind us the final pointer state

$$|\eta\rangle = \frac{1}{\sqrt{2}} \left(\cos \frac{\phi}{2} |\xi_+\rangle + \sin \frac{\phi}{2} e^{-i\omega t} |\xi_-\rangle \right) / \sqrt{p(\omega)}, \quad (1)$$

where,

$$|\xi_{\pm}\rangle = e^{-i\varphi} \sin \frac{\theta}{2} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle \mp i \cos \frac{\theta}{2} \left| \frac{1}{2}, +\frac{1}{2} \right\rangle, \quad (2)$$

$$p(\omega) = (1 - \cos \theta \sin \phi \cos \omega t) / 2, \quad (3)$$

$$\langle \xi_{\pm} | \xi_{\pm} \rangle = 1, \quad \langle \xi_- | \xi_+ \rangle = \langle \xi_+ | \xi_- \rangle = -\cos \theta. \quad (4)$$

Taking the derivative of the final pointer state, we obtain

$$\frac{d|\eta\rangle}{d\omega} = \frac{-it}{\sqrt{2p(\omega)}} \left(\sin \frac{\phi}{2} e^{-i\omega t} |\xi_-\rangle \right). \quad (5)$$

Then, we have

$$\frac{d\langle\eta|}{d\omega} \frac{d|\eta\rangle}{d\omega} = \frac{t^2}{2p(\omega)} \sin^2 \frac{\phi}{2}, \text{ and} \quad (6)$$

$$\frac{d\langle\eta|}{d\omega} |\eta\rangle = \frac{it}{2p(\omega)} \left[\sin^2 \frac{\phi}{2} - \frac{1}{2} \cos \theta \sin \phi e^{i\omega t} \right]. \quad (7)$$

And thus, the measured quantum Fisher information yields

$$\begin{aligned} Q_m^{(1/2)} &= 4p(\omega) \left[\frac{d\langle\eta|}{d\omega} \frac{d|\eta\rangle}{d\omega} - \left| \frac{d\langle\eta|}{d\omega} |\eta\rangle \right|^2 \right] \\ &= \frac{t^2 \sin^2 \theta \sin^2 \phi}{2(1 - \cos \theta \sin \phi)} + O(\omega^2). \end{aligned} \quad (8)$$

Equation (7) in the main text is obtained.

To calculate the postselected classical Fisher information F_p , we substitute

$$\frac{dp(\omega)}{d\omega} = \frac{t}{2} \cos \theta \sin \phi \sin \omega t \quad (9)$$

to Eq. (5), then we obtain $F_p = O(\omega^2)$.

Quantum Fisher information in the case of $j \geq 1$: $Q_m^{(j)}$

The final pointer state is the same as Eq. (1), where

$$\begin{aligned} |\xi_{\pm}\rangle &= \exp \left(\mp i \frac{\pi}{2} |j, +j\rangle \langle j, +j| \right) |\xi\rangle \\ &= \sum_{m=-j, j \geq 1}^{j-1} c_{jm} |j, m\rangle \mp i \left(\cos \frac{\theta}{2} \right)^{2j} |j, +j\rangle, \end{aligned} \quad (10)$$

and

$$c_{jm} = \sqrt{\binom{2j}{j+m}} \left(\cos \frac{\theta}{2} \right)^{j+m} \left(\sin \frac{\theta}{2} e^{-i\varphi} \right)^{j-m}, \quad -j \leq m \leq j-1. \quad (11)$$

Then, we obtain

$$\langle \xi_{\pm} | \xi_{\pm} \rangle = 1, \text{ and } \langle \xi_{-} | \xi_{+} \rangle = \langle \xi_{+} | \xi_{-} \rangle = \sum_{m=-j}^{j-1} c_m^2 - (\cos \frac{\theta}{2})^{4j}, \quad (12)$$

as in the main text. Similar to the case of $j = 1/2$, we calculate the derivative of the final pointer state as

$$\frac{d|\eta\rangle}{d\omega} = \frac{-it}{\sqrt{2p(\omega)}} \left(\sin \frac{\phi}{2} e^{-i\omega t} |\xi_{-}\rangle \right). \quad (13)$$

Then, we have

$$\frac{d\langle \eta |}{d\omega} \frac{d|\eta\rangle}{d\omega} = \frac{t^2}{2p(\omega)} \sin^2 \frac{\phi}{2}, \text{ and } \frac{d\langle \eta |}{d\omega} |\eta\rangle = \frac{it}{2p(\omega)} \left\{ \sin^2 \frac{\phi}{2} + \frac{1}{2} \left[\sum_{m=-j}^{j-1} c_m^2 - (\cos \frac{\theta}{2})^{4j} \right] \sin \phi e^{i\omega t} \right\}. \quad (14)$$

The measured quantum Fisher information yields

$$\begin{aligned} Q_m^{(j)} &= 4p(\omega) \left[\frac{d\langle \eta |}{d\omega} \frac{d|\eta\rangle}{d\omega} - \left| \frac{d\langle \eta |}{d\omega} |\eta\rangle \right|^2 \right] \\ &= t^2 \left\{ 2 \sin^2 \frac{\phi}{2} - \frac{1}{p(\omega)} \left[\sin^2 \frac{\phi}{2} + \frac{1}{2} \left(\sum_{m=-j}^{j-1} c_m^2 - (\cos \frac{\theta}{2})^{4j} \right) \sin \phi e^{i\omega t} \right]^2 \right\} + O(\omega^2), \end{aligned} \quad (15)$$

where

$$p(\omega) = \frac{1}{2} \left[1 + \left(\sum_{m=-j}^{j-1} c_m^2 - (\cos \frac{\theta}{2})^{4j} \right) \cos \phi \cos \omega t \right]. \quad (16)$$

From this result, we can easily calculate the measured quantum Fisher information for any given j .