

# Remarks on some simple $C^*$ -algebras admitting a unique lower semicontinuous 2-quasitrace

Jacopo Bassi  
 Department of Mathematics  
 University of Tor Vergata  
 Via della Ricerca Scientifica 1  
 00133 Roma, Italy  
 E-mail: bassi@mat.uniroma2.it

## Abstract

Using different descriptions of the Cuntz semigroup and of the Pedersen ideal, it is shown that  $\sigma$ -unital simple  $C^*$ -algebras with almost unperforated Cuntz semigroup, a unique lower semicontinuous 2-quasitrace and whose stabilization has almost stable rank 1 are either stable or algebraically simple.

## 1 Introduction

The Cuntz semigroup was introduced by Cuntz in [C78] in order to study the existence of quasitraces on certain  $C^*$ -algebras. Since then, it has received attention in view of the counterexamples to the classification program of  $C^*$ -algebras by means of their Elliott invariant constructed by Toms in [T08], where it is proved the existence of simple, unital, nuclear, non-elementary  $C^*$ -algebras sharing the same Elliott invariant, but whose Cuntz semigroups differ. Despite the existence of many non-isomorphic nuclear, simple  $C^*$ -algebras with the same Cuntz semigroup, in [R12] it was proved that a modified version of the Cuntz semigroup classifies certain inductive limits of 1-dimensional  $NCCW$  (Non-Commutative CW) complexes.

---

2010 *Mathematics Subject Classification*: Primary 46L05.

*Key words and phrases*:  $C^*$ -algebras, Cuntz semigroup.

In [CEI08] it is shown that the Cuntz semigroup of a  $C^*$ -algebra can be equivalently described in terms of certain classes of Hilbert modules, and in [ORT11] another characterization in terms of classes of open projections in the double dual of the  $C^*$ -algebra is given. In both these papers it is observed that, under the assumption of almost stable rank 1, Cuntz comparison can be reduced to a simpler relation. Also, as shown in [ERS11], [TT12] and [T], in some cases, the Cuntz semigroup of a  $C^*$ -algebra can be reconstructed from the lower semicontinuous functions on the cone of lower semicontinuous (2-quasi)traces on the  $C^*$ -algebra and the Murray-von Neumann semigroup.

The Cuntz semigroup was recently used in [APRT18] and [T] in order to give answers to important open problems in the field of  $C^*$ -algebras, such as a conjecture by Blackadar and Handelman on dimension functions, the Global Glimm Halving Problem, and questions on the equivalence of certain different comparability properties.

Specializing to  $C^*$ -algebras with suitable properties, in the present article it is given a description of the Pedersen ideal in terms of Cuntz comparison, and the above correspondence between different descriptions of the Cuntz semigroup is employed in order to establish a connection between algebraic simplicity/stability and related finiteness/infiniteness properties.

## 1.1 Notation

If  $A$  is a  $C^*$ -algebra, we denote by  $\tilde{A}$  its minimal unitization; if  $A$  is unital, we denote by  $GL(A)$  the group of invertible elements in  $A$ . The  $C^*$ -algebra of compact operators on a separable infinite-dimensional Hilbert space is denoted by  $\mathbb{K}$ .

If  $A$  is a  $C^*$ -algebra and  $x$  is a positive element in  $A$ , we denote by  $E_x = \overline{xA}$  the corresponding right ideal, viewed as a Hilbert module. If  $x$  is strictly positive for  $A$ , we identify  $E_x$  with  $A$ . The hereditary  $C^*$ -algebra generated by an element  $y \in A$  is denoted by  $\text{her}(y)$ .

## 2 The Cuntz semigroup

This section contains definitions and preliminary results needed in the forthcoming part. Let  $A$  be a  $C^*$ -algebra and  $a, b$  two positive elements in  $A$ ; then  $a$  is *Cuntz subequivalent* to  $b$  ( $a \precsim b$ ) if there is a sequence  $\{x_n\}$  of elements in  $A$  such that  $a = \lim x_n b x_n^*$ ; and  $a$  is *Cuntz equivalent* to  $b$  ( $a \sim b$ ) if  $a \precsim b$  and  $b \precsim a$ . If  $A$  is a  $C^*$ -algebra, its *Cuntz semigroup*  $(Cu(A), \oplus, \leq) = Cu(A)$  is the partially ordered abelian semigroup whose elements are Cuntz classes of positive elements in  $A \otimes \mathbb{K}$ , where the semigroup structure is

obtained by composing any  $*$ -isomorphism  $M_2(\mathbb{K}) \otimes A \rightarrow \mathbb{K} \otimes A$  with the orthogonal sum  $\mathbb{K} \otimes A \times \mathbb{K} \otimes A \rightarrow M_2(\mathbb{K}) \otimes A$ . The partial order is induced by Cuntz subequivalence: if  $[a], [b] \in Cu(A)$  are the Cuntz classes of elements  $a$  and  $b$  respectively, then  $[a] \leq [b]$  if and only if  $a \precsim b$ . This is the same as  $W(A \otimes \mathbb{K})$  in the notation of [APT11].

If  $A$  is a  $C^*$ -algebra, every increasing sequence in  $Cu(A)$  admits a supremum ([APT11] Proposition 4.17). Let  $[a]$  and  $[b]$  be elements in  $Cu(A)$ , we say that  $[a]$  is *compactly contained in*  $[b]$  ( $[a] \ll [b]$ ) if for every increasing sequence  $\{[b_n]\} \subset Cu(A)$  such that  $[b] \leq \sup_n [b_n]$ , there is a  $k \in \mathbb{N}$  such that  $[a] \leq [b_k]$  (this is the way below relation in [APT11] Definition 4.1).

**Definition 2.1** ([APT18] Section 2.1 and Definition 5.3.1). Let  $A$  be a  $C^*$ -algebra and  $[a] \in Cu(A)$ .  $[a]$  is *compact* if  $[a] \ll [a]$ .  $[a]$  is *soft* if for every  $[b] \in Cu(A)$  satisfying  $[b] \ll [a]$  there is  $n \in \mathbb{N}$  such that  $(n+1)[b] \leq n[a]$ .

If  $A$  is a  $C^*$ -algebra, the purely non-compact elements in  $Cu(A)$ , defined in [CEI08] pag. 27, are soft by [APT18] Proposition 5.3.5.

**Definition 2.2** ([ERS11], Introduction). A *2-quasitrace* on a  $C^*$ -algebra  $A$  is a map  $\tau : (A \otimes \mathbb{K})_+ \rightarrow [0, \infty]$  that sends 0 to 0, is linear on commuting elements and satisfies the trace identity:  $\tau(x^*x) = \tau(xx^*)$  for every  $x \in A \otimes \mathbb{K}$ .

Denote the set of lower semicontinuous 2-quasitraces on  $A$  by  $QT_2(A)$ .

**Definition 2.3** ([ERS11] pag. 17). A *functional* on the Cuntz semigroup of a  $C^*$ -algebra  $A$  is a map  $Cu(A) \rightarrow [0, \infty]$  that preserves the 0 element, sum, partial order, and suprema of increasing sequences.

We denote the set of functionals on  $Cu(A)$  by  $F(Cu(A))$ . As usual, if  $\lambda$  is a functional on  $Cu(A)$ , we say that  $\lambda$  is *trivial* if it only takes values 0 and  $\infty$  and is *nontrivial* otherwise. The same terminology is used for the lower semicontinuous 2-quasitraces. There is a well known bijection  $QT_2(A) \simeq F(Cu(A))$  ([ERS11] Proposition 4.2).

If  $A$  is a simple  $C^*$ -algebra admitting a non-trivial lower semicontinuous 2-quasitrace, every non-zero element in  $Cu(A)$  is either compact or soft by [APT18] Proposition 5.3.16.

Let  $A$  be a  $C^*$ -algebra and consider the trivial lower semicontinuous 2-quasitraces on  $A$  defined by

$$\tau_0(a) = 0 \quad \text{for every } a \in (A \otimes \mathbb{K})_+, \quad \tau_\infty(a) = \begin{cases} 0 & \text{for } a = 0 \\ \infty & \text{for } a \in (A \otimes \mathbb{K})_+ \setminus \{0\}. \end{cases}$$

In the same way, we define trivial functionals on  $Cu(A)$  by

$$\lambda_0([a]) = 0 \quad \text{for all } [a] \in Cu(A), \quad \lambda_\infty([a]) = \begin{cases} 0 & \text{for } [a] = 0 \\ \infty & \text{for } [a] \in Cu(A) \setminus \{0\} \end{cases}.$$

**Lemma 2.4.** *Let  $A$  be a simple  $C^*$ -algebra. Then the only possible trivial lower semicontinuous 2-quasitraces on  $A$  are  $\tau_0$  and  $\tau_\infty$ ; the only possible trivial functionals on  $Cu(A)$  are  $\lambda_0$  and  $\lambda_\infty$ .*

*Furthermore,  $A$  admits a unique nontrivial lower semicontinuous 2-quasitrace (up to a scalar) if and only if  $Cu(A)$  admits a unique nontrivial functional (up to a scalar).*

*Proof.* Let  $\lambda$  be a trivial functional on  $Cu(A)$ . It follows from the proof of [KR00] Lemma 3.12 that the set  $N_\lambda := \{a \in A \otimes \mathbb{K} : \lambda([|a|]) = 0\}$  is an algebraic ideal. By simplicity it is either the set  $\{0\}$  or is dense in  $A \otimes \mathbb{K}$ .

If  $N_\lambda = \{0\}$ , then  $\lambda = \lambda_\infty$ ; hence suppose  $N_\lambda \neq 0$ . Let  $[a]$  be an element of  $Cu(A)$  and note that  $\lambda([a]) = \sup_{\epsilon > 0} \lambda([(a - \epsilon)_+])$ . Since  $N_\lambda$  is dense,  $\text{Ped}(A \otimes \mathbb{K}) \subset N_\lambda$  and so  $\lambda([(a - \epsilon)_+]) = 0$  for every  $\epsilon > 0$ . Thus  $\lambda([a]) = 0$  and  $\lambda = \lambda_0$ .

Using the bijective correspondence between functionals on the Cuntz semigroup and lower semicontinuous 2-quasitraces on the corresponding  $C^*$ -algebra, the 2-quasitrace associated to a functional  $\lambda$  is  $\tau_\lambda(a) = \int_0^\infty \lambda([(a - t)_+])$  for  $a \in A_+$ . Hence the functional  $\lambda_0$  corresponds to the lower semicontinuous 2-quasitrace  $\tau_0$  and  $\lambda_\infty$  corresponds to  $\tau_\infty$ .

It follows that if  $A$  admits a unique non-trivial lower semicontinuous 2-quasitrace up to scalars, the multiples of it correspond to the same multiples of the associated non-trivial functional on  $Cu(A)$ .  $\square$

Another property of the Cuntz semigroup of a  $C^*$ -algebra which turns out to be important, in particular for its connection with the Toms-Winter conjecture, is the so called *almost unperforation*:

**Definition 2.5** ([ERS11] pag. 26). A partially ordered semigroup  $(S, +, \leq)$  is said to be almost unperforated if, whenever there are elements  $s, t$  in  $S$  and  $n \in \mathbb{N}$  such that  $(n + 1)s \leq nt$ , then  $s \leq t$ .

**Theorem 2.6** ([APT18] Theorem 5.3.12). *Let  $A$  be a  $C^*$ -algebra such that  $Cu(A)$  is almost unperforated. If  $[a]$  and  $[b]$  are elements in  $Cu(A)$  such that  $[a]$  is soft and  $\lambda([a]) \leq \lambda([b])$  for every functional  $\lambda \in F(Cu(A))$ , then  $[a] \leq [b]$ .*

As already mentioned in the Introduction,  $C^*$ -algebras with almost stable rank 1 have received a particular attention in [CEI08] and [ORT11], since for these  $C^*$ -algebras Cuntz subequivalence can be described in terms of other relations. This fact will be an important ingredient in the following section. We recall this concept.

**Definition 2.7** ([R16] Definition 3.1). Let  $A$  be a  $C^*$ -algebra. We say that  $A$  has *almost stable rank 1* if for every hereditary  $C^*$ -subalgebra  $B \subset A$ , we have  $B \subset \overline{GL(\tilde{B})}$ .

Note that almost stable rank 1 is not a stable property ([T] Example 6.6).

### 3 Algebraic simplicity and stability

If  $A$  is a  $C^*$ -algebra, we denote by  $\text{Ped}(A)$  its Pedersen ideal, i.e. the dense ideal that is contained in every other dense ideal and refer to [P79] 5.6 for its construction and properties. We will give two descriptions of the Pedersen ideal for  $C^*$ -algebras belonging to a certain class and use these for the proof of Theorem 3.5.

The first characterization of the Pedersen ideal is in terms of compact containment for a certain class of  $C^*$ -algebras. A more general result, that appears as a Remark on page 4 of [TT12], was suggested to hold by Elliott without the assumption that the stabilization of the  $C^*$ -algebra has almost stable rank 1.

**Proposition 3.1.** *Let  $A$  be a  $C^*$ -algebra such that  $A \otimes \mathbb{K}$  has almost stable rank 1. Then*

$$\text{Ped}(A) = \{a \in A : \exists b \in M_\infty(A)_+ \text{ such that } [a] \ll [b]\}.$$

*Proof.* Let  $a$  be an element of  $A$  and write  $a = u|a|$  for its polar decomposition in  $A^{**}$ . If  $|a|$  belongs to  $\text{Ped}(A)$ , then by [P79] Proposition 5.6.2  $|a|^\alpha$  belongs to  $\text{Ped}(A)$  for every  $\alpha > 0$ ; note now that  $u|a|^{1-\alpha}$  belongs to  $A$  for every  $0 < \alpha < 1$  and so  $u|a| = u|a|^{1-\alpha}|a|^\alpha$  belongs to  $\text{Ped}(A)$ . Thus we are left to prove that the set

$$J_+ := \{a \in A_+ : \exists b \in M_\infty(A)_+ \text{ such that } [a] \ll [b]\}$$

is contained in  $\text{Ped}(A)$ , that

$$J := \{a \in A : \exists b \in M_\infty(A)_+ \text{ such that } [a] \ll [b]\}$$

is an ideal and that it is dense.

Let  $a$  be a positive element in  $A$  and suppose there is  $n \in \mathbb{N}$  and a positive element  $b$  in  $M_n(A)$  such that  $[a] \ll [b]$ . Then there is  $\epsilon > 0$  such that  $a \precsim (b - \epsilon)_+$  in  $M_n(A)$ . Since  $A \otimes \mathbb{K}$  has almost stable rank 1, by the proof of [CEI08] Theorem 3 and by Proposition 4.6 of [ORT11], there is an element  $y$  in  $M_n(A)$  such that  $\text{her}(a) = \text{her}(yy^*)$ ,  $y^*y \in \text{her}((b - \epsilon)_+)$ ; in particular by [P79] Proposition 5.6.2  $y^*y$  belongs to  $\text{Ped}(M_n(A))$  and the same is true for  $yy^*$ .

Now we want to show that for any  $n \in \mathbb{N}$ ,  $\text{Ped}(M_n(A)) = \text{Ped}(A) \otimes_{\text{alg}} M_n$ . Notice that  $\text{Ped}(A) \otimes_{\text{alg}} M_n$  is a dense ideal in  $M_n(A)$  and so  $\text{Ped}(M_n(A)) \subset \text{Ped}(A) \otimes_{\text{alg}} M_n$ . On the other hand, if  $x = (x_{i,j})$  is an element in  $M_n(A)$  such that  $x_{i,j}$  belongs to  $\text{Ped}(A)$  for every  $1 \leq i, j \leq n$ , we can write each  $x_{i,j}$  as a finite linear combination of positive

elements  $y_{i,j}^{(k)}$ , for each of which there are a natural number  $m_{i,j}^{(k)}$  and elements  $z_{i,j}^{(k,l)}, \tilde{z}_{i,j}^{(k,l)}$  such that  $y_{i,j}^{(k)} \leq \sum_{l=1}^{m_{i,j}^{(k)}} z_{i,j}^{(k,l)}$  and  $z_{i,j}^{(k,l)} \tilde{z}_{i,j}^{(k,l)} = z_{i,j}^{(k,l)}$ . Denoting by  $e_{i,j}$  the matrix units for  $M_n$ , it follows that the same is true if we replace the elements  $x_{i,j}, y_{i,j}^{(k)}, z_{i,j}^{(k,l)}$  and  $\tilde{z}_{i,j}^{(k,l)}$  with  $x_{i,j} \otimes e_{i,j}, y_{i,j}^{(k)} \otimes e_{i,j}, z_{i,j}^{(k,l)} \otimes e_{i,j}$  and  $\tilde{z}_{i,j}^{(k,l)} \otimes e_{j,j}$  respectively. It follows that for every  $1 \leq i, j \leq n$   $x_{i,j} \otimes e_{i,j}$  belongs to  $\text{Ped}(M_n(A))$  and so  $x \in \text{Ped}(M_n(A))$ . Thus  $\text{Ped}(M_n(A)) = M_n(\text{Ped}(A))$ . In particular, considering  $A$  as a  $C^*$ -subalgebra of  $M_n(A)$  via the embedding  $a \mapsto a \otimes e_{1,1}$ , it follows that  $A \cap \text{Ped}(M_n(A)) = A \cap (M_n(\text{Ped}(A))) = \text{Ped}(A)$ . Thus  $\text{her}(a) \subset A \cap M_n(\text{Ped}(A)) = \text{Ped}(A)$  and so  $a$  belongs to  $\text{Ped}(A)$ .

The proof that  $J$  is an ideal goes as in Lemma 3.12 of [KR00] and we omit it.

Since every element in  $A$  is a linear combination of positive elements and  $J$  is a linear space, we just have to prove that  $J_+$  is dense in  $A_+$ . Let  $a \in A_+$ , then  $\|a - (a - \epsilon)_+\| < \epsilon$  and  $[(a - \epsilon)_+] \ll [a]$ .

Hence  $J$  is a dense two-sided ideal and has to be equal to  $\text{Ped}(A)$ .  $\square$

**Corollary 3.2.** *Let  $A$  be a  $\sigma$ -unital stable  $C^*$ -algebra with almost stable rank 1 and let  $h$  be a strictly positive element for  $A$ . Then*

$$\begin{aligned} \text{Ped}(A) &= \{a \in A : \exists b \in A_+ \text{ such that } [a] \ll [b]\} \\ &= \{a \in A : [a] \ll [h]\}. \end{aligned}$$

*Proof.* Since  $A$  is stable, by [R04] Lemma 2.6, it is large in  $A \otimes \mathbb{K}$  in the sense of [R04] Definition 2.4. Suppose there are  $a \in A, b \in M_n(A)_+$  such that  $[a] \ll [b]$ , then there is  $\epsilon > 0$  such that  $[a] \leq [(b - \epsilon)_+] \ll [(b - \epsilon/2)_+]$ ; by largeness, there is  $a_{\epsilon/2} \in A_+$  such that  $[(b - \epsilon/2)_+] = [a_{\epsilon/2}]$  and thus  $[a] \ll [a_{\epsilon/2}]$ . The result follows from Proposition 3.1.  $\square$

**Proposition 3.3.** *Let  $A$  be a simple  $C^*$ -algebra with almost unperforated Cuntz semigroup such that  $A \otimes \mathbb{K}$  has almost stable rank 1 and assume that  $A$  admits a unique lower semi-continuous nontrivial 2-quasitrace  $\tau$  (up to scalar multiples). Let  $d_\tau$  be the corresponding functional on  $Cu(A)$ . Then*

$$\text{Ped}(A) = \{a \in A : d_\tau([a]) < \infty\}.$$

*Proof.* By Lemma 2.4  $d_\tau$  is a nontrivial functional on  $Cu(A)$  and is the unique such functional, up to scalars. Let  $J' := \{a \in A : d_\tau([a]) < \infty\}$ .

First of all notice that  $J'$  is an ideal (again, the proof goes as in Lemma 3.12 of [KR00], using the fact that  $d_\tau$  is a functional on  $Cu(A)$ ) and it is obviously dense since  $A$  is simple. By simplicity, every element in  $Cu(A)$  is either compact or soft.

Let  $a \in A$  be such that  $0 \neq d_\tau([a]) < \infty$ ; if  $[a]$  is compact, then  $a$  belongs to  $\text{Ped}(A)$  by Proposition 3.1, thus we can suppose that  $[a]$  is soft.

Since  $0 < d_\tau([a]) < \infty$  and  $d_\tau([a]) = \sup_n d_\tau([(a| - 1/n)_+])$ , there exists  $m \in \mathbb{N}$  such

that  $0 < d_\tau([( |a| - 1/m)_+]) < \infty$  and so there is a  $k \in \mathbb{N}$  such that  $kd_\tau([( |a| - 1/m)_+]) \geq d_\tau([|a|])$ . Using comparison and the fact that the only possible functionals on  $Cu(A)$  are  $\lambda_0$ ,  $\lambda_\infty$  and  $d_\tau$  by Lemma 2.4, we see that  $[|a|] \leq [( |a| - 1/m)_+ \otimes 1_k] \ll [( |a| - 1/2m)_+ \otimes 1_k]$  and so  $a$  belongs to  $\text{Ped}(A)$  by Proposition 3.1. The proof is complete.  $\square$

Let  $A$  be a  $C^*$ -algebra and  $[a] \in Cu(A)$ . Then  $[a]$  is finite if  $[a] + [b] = [a]$  implies  $[b] = 0$ , and  $[a]$  is infinite if there is a non-zero element  $[b] \in Cu(A)$  such that  $[a] + [b] = [a]$  (cfr. [APT18] Section 5.2.2).  $[a]$  is properly infinite if  $2[a] = [a]$ . If  $A$  is simple, then  $[a] \in Cu(A)$  is infinite if and only if it is properly infinite.

**Proposition 3.4.** *Let  $A$  be a simple  $C^*$ -algebra with almost unperforated Cuntz semigroup such that  $A \otimes \mathbb{K}$  has almost stable rank 1 and suppose that  $A$  admits a unique lower semicontinuous 2-quasitrace  $\tau$  (up to scalar multiples). Then*

$$\text{Ped}(A) = \{a \in A \mid [|a|] \text{ is finite} \}.$$

*Proof.* Let  $a$  be a positive element in  $A$ . We distinguish different cases.

If  $[|a|]$  is compact, then  $a$  belongs to  $\text{Ped}(A)$  by Proposition 3.1 and, since  $Cu(A)$  admits a non-trivial functional, it cannot contain infinite compact elements by simplicity.

Suppose that  $[|a|]$  is soft. By simplicity, any infinite element is properly infinite. Thus, if  $d_\tau([|a|]) < \infty$ , then  $[|a|]$  is finite.

If  $d_\tau([|a|]) = \infty$ , it follows from Proposition 3.3 that  $a$  does not belong to  $\text{Ped}(A)$  and by comparison that  $|a| \sim |a| \oplus |a|$ ; hence  $[|a|]$  is properly infinite.  $\square$

In [CEI08] it is defined a notion of Cuntz subequivalence and compact containment for countably generated Hilbert modules of a  $C^*$ -algebra and is proved that the induced partially ordered semigroup coincides with the Cuntz semigroup, in the sense that the two constructions are naturally equivalent ([CEI08] Section 6). Given two countably generated Hilbert modules  $X$  and  $Y$  over a  $C^*$ -algebra, we write  $X \precsim Y$  if  $X$  is Cuntz subequivalent to  $Y$ , and  $X \Subset Y$  if  $X$  is compactly contained in  $Y$ ; we refer to [CEI08] Section 1 for the definitions.

**Theorem 3.5.** *Let  $A$  be a  $\sigma$ -unital stable simple  $C^*$ -algebra with almost unperforated Cuntz semigroup, with almost stable rank 1 and admitting a unique lower semicontinuous nontrivial 2-quasitrace  $\tau$ . Let  $x$  be a positive element in  $A$ , then we have*

$$[x] \text{ is infinite} \Leftrightarrow \text{her}(x) \text{ is stable} \Leftrightarrow E_x \simeq A$$

and

$$[x] \text{ is finite} \Leftrightarrow \text{her}(x) \text{ is algebraically simple} \Leftrightarrow E_x \Subset A.$$

Furthermore, if  $y$  is a positive element in  $A$  such that  $[x] < [y]$ , then  $[x] \ll [y]$  and this happens if and only if there exist elements  $z \in A$  and  $e$  self adjoint in  $\text{her}(y)$  such that

$\text{her}(x) = \text{her}(z^*z)$ ,  $\text{her}(zz^*) \subset \text{her}(y)$ ,  $ea = a$  for every  $a \in \text{her}(zz^*)$ . This condition is also equivalent to the existence of a Hilbert module  $E$  over  $A$  such that  $E_x \simeq E \subseteq E_y$ .

*Proof.* If  $A$  is simple, the Cuntz class of a positive element  $x \in A$  is infinite if and only if it is properly infinite. Now, if  $[x]$  is properly infinite, then it defines the same Cuntz class of any strictly positive element for  $A$ ; by [CEI08] Theorem 3 and [ORT11] Proposition 4.3 it follows that, since  $A$  has almost stable rank 1,  $\text{her}(x) \simeq A$  and  $E_x \simeq A$  as countably generated Hilbert modules.

Conversely, the Cuntz class of a strictly positive element in a stable  $C^*$ -algebra is properly infinite, as well as the generator of a Hilbert module that is isomorphic to  $A$  and so the characterization of infinite elements in the Cuntz semigroup in terms of hereditary subalgebras and Hilbert modules is complete.

By Proposition 3.4  $[x]$  is finite if and only if  $x$  belongs to  $\text{Ped}(A)$  and in this case  $\text{her}(x)$  is algebraically simple by [P79] Proposition 5.6.2. Note now that if  $\text{her}(x)$  is algebraically simple and  $x$  does not belong to  $\text{Ped}(A)$ , then by Proposition 3.3  $d_\tau([x]) = \infty$ , then it follows that  $\text{her}(x)$  only admits the trivial lower semicontinuous 2-quasitrace  $\tau_\infty$ , contradicting the hypothesis. Hence, under our assumptions,  $\text{her}(x)$  is algebraically simple if and only if  $[x]$  is finite.

By Corollary 3.2,  $x$  belongs to  $\text{Ped}(A)$  if and only if  $[x] \ll [h]$  for any strictly positive element  $h$  for  $A$ . From the equivalence between compact containment for classes of Hilbert modules and the way below relation (see [CEI08] Theorem 1) it follows that  $[x] \ll [h]$  if and only if  $E_x \subseteq A$  up to isomorphism of Hilbert modules.

Suppose now that there is a positive element  $y \in A$  such that  $[x] < [y]$ . If either  $[x]$  or  $[y]$  is compact, then  $[x] \ll [y]$ , hence we can suppose  $[x]$  and  $[y]$  are soft. Note that  $d_\tau([(y-\epsilon)_+]) < d_\tau([y])$  for every  $\epsilon > 0$ , otherwise  $[y]$  would be compact by comparison and so, since  $d_\tau([y]) = \sup_{\epsilon > 0} d_\tau([(y-\epsilon)_+])$ , there exists  $\eta > 0$  such that  $d_\tau([x]) \leq d_\tau([(y-\eta)_+])$ , entailing  $[x] \ll [y]$ . From the equivalence between compact containment in the concrete and in the abstract sense proved in [CEI08] Theorem 1, this is the same as the existence of a Hilbert module  $E$  such that  $E_x \precsim E \subseteq E_y$ . Thus, in the case  $A$  has almost stable rank 1, it follows from Theorem 3 of [CEI08] that there exists a Hilbert module  $E'$  such that  $E_x \simeq E' \subset E \subseteq E_y$ , hence in particular  $E_x \simeq E' \subseteq E_y$ . Denote by  $\Phi : E_x \rightarrow E'$  an isomorphism of Hilbert modules; since  $A$  is  $\sigma$ -unital,  $x$  belongs to  $E_x$  and  $E' = \overline{\Phi(x)A}$ , with  $\Phi(x) \in A$  since  $E' \subset E_y$ ; it follows then from Proposition 4.3 of [ORT11] that there is an element  $z \in A$  such that  $\text{her}(x) = \text{her}(z^*z)$  and  $\text{her}(\Phi(x)) = \text{her}(zz^*)$ . If now  $e \in \mathbb{K}(E_y) = \text{her}(y)$  is a self adjoint element such that  $e|_{E_{\Phi(x)}} = \text{id}|_{E_{\Phi(x)}}$ , it follows that  $ea = a$  for every  $a \in \text{her}(\Phi(x))$ . Suppose now that there are an element  $z \in A$  and  $e$  selfadjoint in  $\text{her}(y)$  such that  $\text{her}(x) = \text{her}(z^*z)$ ,  $\text{her}(zz^*) \subset \text{her}(y)$  and  $ea = a$  for every



$a \in \text{her}(zz^*)$ . Then again by Proposition 4.3 of [ORT11] it follows that  $E_x \simeq E_{zz^*}$  and since  $zz^*$  belongs to  $\text{her}(zz^*)$ , it follows that  $ezz^* = zz^*$ , which implies  $e|_{E_{zz^*}} = \text{id}|_{E_{zz^*}}$  and hence  $E_x \simeq E_{zz^*} \subseteq E_y$ . In this case, it follows again by [CEI08] Theorem 1 that  $[x] \ll [y]$ .  $\square$

**Remark 3.6.** There are examples of non-algebraically simple, non-stable simple  $\sigma$ -unital  $C^*$ -algebras with stable rank 1, admitting non-trivial lower semicontinuous 2-quasitraces and whose Cuntz semigroup is almost unperforated. They were constructed by Blackadar in [B80]; namely, in Corollary 4.5 it is proved that for any simple unital  $AF$ -algebra  $A$  and any  $F_\sigma$  subset of its space of bounded traces, it is possible to find an hereditary  $AF$ -subalgebra  $B$  of  $A \otimes \mathbb{K}$  containing  $A$  such that the bounded traces on  $B$  are exactly the extensions coming from the  $F_\sigma$  set chosen. It is clear that this construction does not work if  $A$  has just one trace.

**Example 3.7.** The following example is well known. The  $C^*$ -algebra  $\mathbb{K}$  of compact operators on a separable Hilbert space has stable rank 1 and a unique nontrivial trace, up to scalars. Its Cuntz semigroup is  $Cu(\mathbb{K}) = \bar{\mathbb{N}}$ , which is almost unperforated. By Theorem 3.5 any hereditary  $C^*$ -subalgebra of  $\mathbb{K}$  is either algebraically simple (actually unital) or isomorphic to  $\mathbb{K}$ .

**Example 3.8.** The Jiang-Su algebra  $\mathcal{Z}$  (see [JS99]) has stable rank 1, a unique trace and almost unperforated Cuntz semigroup equal to  $Cu(\mathcal{Z}) = \mathbb{N} \sqcup (0, \infty]$ . The hereditary  $C^*$ -subalgebras of  $\mathcal{Z} \otimes \mathbb{K}$  generated by compact elements (corresponding to  $\mathbb{N} \subset Cu(A)$ ) are unital; the ones corresponding to elements in  $(0, \infty)$  are nonunital and algebraically simple.

**Example 3.9.** The  $C^*$ -algebra  $\mathcal{W}$  introduced by Jacelon in [J13] has stable rank 1, a unique trace and its Cuntz semigroup is  $Cu(\mathcal{W}) = [0, \infty]$ , which is almost unperforated (one could also just notice that  $\mathcal{W}$  is  $\mathcal{Z}$ -stable). Thus any  $\sigma$ -unital  $C^*$ -algebra that is Morita-equivalent to  $\mathcal{W}$  is either algebraically simple (in correspondence of elements in  $(0, \infty) \subset Cu(\mathcal{W})$ ) or stable.

**Example 3.10.** If  $X$  is a compact manifold with finite covering dimension foliated by a free minimal action of  $\mathbb{R}$  admitting a unique invariant transverse measure, then  $C(X) \rtimes \mathbb{R}$  satisfies the hypothesis of Theorem 3.5 in the case it is projectionless. This is a consequence of [MS06] Theorem 6.30, [HSWW17] Corollary 9.2 and [R16] Corollary 3.2.

## Acknowledgements

This research was partially funded by I.N.d.A.M. (grant n. U-UFMBAZ-2018-000926) and the University of Tor vergata (grant n. 669240 CUP: E52I15000700002). The author thanks Prof. Longo for the hospitality at the Università degli Studi di Roma Tor Vergata

for the period of this research. Special thanks go to André Schemaitat and Dominic Enders for many interesting discussions. In particular, Proposition 3.1 is the result of fruitful discussions with the colleague and friend André Schemaitat.

## References

- [APRT18] R. Antoine, F. Perera, L. Robert and H. Thiel,  $C^*$ -algebras of stable rank one and their Cuntz semigroups, preprint: arXiv:1809.03984 (2018).
- [APT18] R. Antoine, F. Perera and H. Thiel, *Tensor products and regularity properties of Cuntz semigroups*, no. 6 in Mem. Amer. Math. Soc., vol. 251, no. 1199, American Mathematical Society, Providence, Rhode Island, 2018.
- [APT11] P. Ara, F. Perera and A. S. Toms, *K-theory for operator algebras. Classification of  $C^*$ -algebras*, page 1–71 in: *Aspects of Operator Algebras and Applications*. P. Ara, F. Lledó and F. Perera (eds.). Contemporary Mathematics vol. 534, Amer. Math. Soc., Providence RI, 2011.
- [B80] B. Blackadar, Traces on simple AF- $C^*$ -algebras, J. Funct. Anal., **38**(1980), no. 2, 156–168.
- [CEI08] K. T. Coward, G. A. Elliott and C. Ivanescu, The Cuntz semigroup as an invariant for  $C^*$ -algebras, J. Reine Angew. Math., **623**(2008), 161–193.
- [C78] J. Cuntz, Dimension functions on simple  $C^*$ -algebras, Math. Ann., **233**(1978), 145–153.
- [ERS11] G. A. Elliott, L. Robert and L. Santiago, The cone of lower semicontinuous traces on a  $C^*$ -algebra, Amer. J. Math., **133**(2011), no. 4, 969–1005.
- [HSWW17] I. Hirshberg and G. Szabó and W. Winter and J. Wu, Rokhlin dimension for flows, Comm. Math. Phys., **353**(2017), no. 1, 253–316.
- [J13] B. Jacelon, A simple, monotracial, stably projectionless  $C^*$ -algebra, J. Lond. Math. Soc. (2), **87**(2013), no. 2, 365–383.
- [JS99] X. Jiang and H. Su, On a simple unital projectionless  $C^*$ -algebra, Amer. J. Math., **121**(1999), no. 2, 359–413.
- [KR00] E. Kirchberg and M. Rørdam, Non-simple purely infinite  $C^*$ -algebras, Amer. J. Math., **122**(2000), no. 3, 637–666.
- [MS06] C. C. Moore and C. L. Schochet, Global analysis on foliated spaces, 2nd edition in Math. Sci. Res. Inst. Publ., vol. 9, Cambridge University Press, New York, 2006.

- [ORT11] E. Ortega, M. Rørdam and H. Thiel, The Cuntz semigroup and comparison of open projections, *J. Funct. Anal.*, **260**(2011), no. 12, 3474–3493.
- [P79] G. K. Pedersen,  *$C^*$ -algebras and their automorphism groups*, no. 14 in London Math. Soc. Monogr. Ser., Academic Press, New York, 1979.
- [R12] L. Robert, Classification of inductive limits of 1-dimensional *NCCW* complexes, *Adv. Math.*, **231**(2012), no. 5, 2802–2836.
- [R16] ———, Remarks on  $\mathcal{Z}$ -stable projectionless  $C^*$ -algebras, *Glasg. Math. J.*, **58**(2016), no. 2, 273–277.
- [R04] M. Rørdam, Stable  $C^*$ -algebras, *Adv. Stud. Pure Math.*, Operator Algebras and Applications, **38**(2004), 1–23.
- [T] H. Thiel, Ranks of operators in simple  $C^*$ -algebras with stable rank one, *Commun. Math. Phys.* (to appear).
- [TT12] A. Tikuisis and A. Toms, On the structure of Cuntz semigroups in (possibly) nonunital  $C^*$ -algebras, *Canad. Math. Bull.*, **58**(2012), no. 2, 402–414.
- [T08] A. Toms, On the classification problem for nuclear  $C^*$ -algebras, *Ann. of Math.* (2), **167**(2008), 1029–1044.