arXiv:1806.08335v2 [math.NT] 29 Aug 2018

Four New Generalized Fibonacci Number Summation Identities

M.J. Kronenburg

Abstract

Two new generalized Fibonacci number summation identities are stated and proved, and two other new generalized Fibonacci number summation identities are derived from these, of which two special cases are already known in literature.

Keywords: Fibonacci number, Lucas number, generalized Fibonacci number.
MSC 2010: 11B39

1 Four New Generalized Fibonacci Number Summation Identities

Let F_n be the Fibonacci number, L_n be the Lucas number and G_n be the generalized Fibonacci number, for which $G_{n+2} = G_{n+1} + G_n$ with any seed G_0 and G_1 [1, 2, 4]. The two new generalized Fibonacci number summation identities are:

$$\sum_{k=0}^{n} \binom{n}{k} (-1)^k F_m^k F_{m+q}^{n-k} G_{p+qk} = (-1)^{nm} F_q^n G_{p-nm} \tag{1.1}$$

$$\sum_{k=0}^{n} \binom{n}{k} (-1)^k L_m^k L_{m+q}^{n-k} G_{p+qk} = 5^{\lfloor n/2 \rfloor} (-1)^{n(m+1)} F_q^n [G_{p-nm+1} - (-1)^n G_{p-nm-1}] \tag{1.2}$$

where $\lfloor x \rfloor$ is the greatest integer less than or equal to x, also called the floor of x [3]. Replacing m by $-m$ and q by $-q$ and using $F_{-n} = (-1)^{n+1} F_n$ and $L_{-n} = (-1)^n L_n$ these two identities transform into the following two identities:

$$\sum_{k=0}^{n} \binom{n}{k} (-1)^{(q+1)k} F_m^k F_{m+q}^{n-k} G_{p-qk} = F_q^n G_{p+nm} \tag{1.3}$$

$$\sum_{k=0}^{n} \binom{n}{k} (-1)^{(q+1)k} L_m^k L_{m+q}^{n-k} G_{p-qk} = 5^{\lfloor n/2 \rfloor} F_q^n [G_{p+nm+1} - (-1)^n G_{p+nm-1}] \tag{1.4}$$

The special cases $q = 1$ and $q = -1$ of (1.3) were already known in literature [1, 2, 4].

2 Proof of the New Fibonacci Summation Identities

The Fibonacci number summation identities (1.1) and (1.2) are proved for $G = F$ by first proving the $n = 1$ case, then proving a recurrence relation for the left side, and then showing that the right side fulfills the same recurrence relation, thus proving these identities by induction on n. Then the identities are generalized to G instead of F. The following Binet formulas for F_n and L_n are well known [1, 2, 3, 4]:

$$\phi = \frac{1}{2}(1 + \sqrt{5}) \tag{2.1}$$

$$F_n = \frac{1}{\sqrt{5}}(\phi^n - (1-\phi)^n) \tag{2.2}$$

$$L_n = \phi^n + (1-\phi)^n \tag{2.3}$$

First the following well known Fibonacci identity is needed [1, 2, 3, 4].

Theorem 2.1.

$$F_{m+1}F_n + F_m F_{n-1} = F_{n+m} \tag{2.4}$$

Proof. Substituting formula (2.2) and using $\phi(1-\phi) = -1$ and $L_{n-1} + L_{n+1} = 5F_n$:

$$\begin{aligned}
&\frac{1}{5}[(\phi^{m+1} - (1-\phi)^{m+1})(\phi^n - (1-\phi)^n) \\
&+ (\phi^m - (1-\phi)^m)(\phi^{n-1} - (1-\phi)^{n-1})] \\
=&\frac{1}{5}[\phi^{n+m+1} + (1-\phi)^{n+m+1} - \phi^{m+1}(1-\phi)^n - (1-\phi)^{m+1}\phi^n \\
&+ \phi^{n+m-1} + (1-\phi)^{n+m-1} - \phi^m(1-\phi)^{n-1} - (1-\phi)^m\phi^{n-1}] \\
=&\frac{1}{5}[L_{n+m+1} + L_{n+m-1} - (-1)^{m+1}L_{n-m-1} - (-1)^m L_{n-m-1}] \\
=&F_{n+m}
\end{aligned} \tag{2.5}$$

□

Substituting $-n$ for n and using $F_{-n} = (-1)^{n+1}F_n$ this identity becomes:

$$F_{n+1}F_m - F_n F_{m+1} = (-1)^n F_{m-n} \tag{2.6}$$

Adding and subtracting these two identities and using $F_{n-1} + F_{n+1} = L_n$ and $F_{n-1} - F_{n+1} = -F_n$ and $F_{m+1} - F_m = F_{m-1}$ and $F_{m-1} + F_{m+1} = L_m$ yields:

$$F_{m+n} + (-1)^n F_{m-n} = L_n F_m \tag{2.7}$$

$$F_{m+n} - (-1)^n F_{m-n} = F_n L_m \tag{2.8}$$

Adding the identity with $m-1$ and $m+1$ and using $F_{m-1} + F_{m+1} = L_m$ and $L_{m-1} + L_{m+1} = 5F_m$ yields two similar identities:

$$L_{m+n} + (-1)^n L_{m-n} = L_n L_m \tag{2.9}$$

$$L_{m+n} - (-1)^n L_{m-n} = 5F_n F_m \tag{2.10}$$

These last four identities were already known in literature [1, 2, 4]. Equations (2.8) and (2.10) are used in the proof below.

Theorem 2.2. *The following identity is the $n = 1$ case of (1.1):*

$$F_{m+q}F_p - F_mF_{p+q} = (-1)^m F_q F_{p-m} \tag{2.11}$$

Proof. Substituting formula (2.2) and using $\phi(1-\phi) = -1$ and (2.10) and $F_{-n} = (-1)^{n+1}F_n$:

$$\begin{aligned}
&\frac{1}{5}[(\phi^p - (1-\phi)^p)(\phi^{m+q} - (1-\phi)^{m+q}) \\
&- (\phi^{p+q} - (1-\phi)^{p+q})(\phi^m - (1-\phi)^m)] \\
=&\frac{1}{5}[\phi^{m+p+q} + (1-\phi)^{m+p+q} - \phi^p(1-\phi)^{m+q} - (1-\phi)^p\phi^{m+q} \\
&- (\phi^{m+p+q} + (1-\phi)^{m+p+q} - \phi^{p+q}(1-\phi)^m - (1-\phi)^{p+q}\phi^m)] \\
=&-\frac{1}{5}[(-1)^p L_{m-p+q} - (-1)^{p+q}L_{m-p-q}] \\
=&-\frac{1}{5}(-1)^p[L_{m-p+q} - (-1)^q L_{m-p-q}] \\
=&-(-1)^p F_q F_{m-p} \\
=&(-1)^m F_q F_{p-m}
\end{aligned} \tag{2.12}$$

□

Theorem 2.3. *The following identity is the $n = 1$ case of (1.2):*

$$L_{m+q}F_p - L_mF_{p+q} = (-1)^{m+1} F_q L_{p-m} \tag{2.13}$$

Proof. Substituting formula (2.2) and (2.3) and using $\phi(1-\phi) = -1$ and (2.8) and $L_{-n} = (-1)^n L_n$:

$$\begin{aligned}
&\frac{1}{\sqrt{5}}[(\phi^p - (1-\phi)^p)(\phi^{m+q} + (1-\phi)^{m+q}) \\
&- (\phi^{p+q} - (1-\phi)^{p+q})(\phi^m + (1-\phi)^m)] \\
=&\frac{1}{\sqrt{5}}[\phi^{m+p+q} - (1-\phi)^{m+p+q} + \phi^p(1-\phi)^{m+q} - (1-\phi)^p\phi^{m+q} \\
&- (\phi^{m+p+q} - (1-\phi)^{m+p+q} + \phi^{p+q}(1-\phi)^m - (1-\phi)^{p+q}\phi^m)] \\
=&-[(-1)^p F_{m-p+q} - (-1)^{p+q}F_{m-p-q}] \\
=&-(-1)^p[F_{m-p+q} - (-1)^q F_{m-p-q}] \\
=&-(-1)^p F_q L_{m-p} \\
=&(-1)^{m+1} F_q L_{p-m}
\end{aligned} \tag{2.14}$$

□

There are variants of (2.11) and (2.13) obtained by adding the identity with $p-1$ and $p+1$ and using $F_{p-1} + F_{p+1} = L_p$ and $L_{p-1} + L_{p+1} = 5F_p$:

$$F_{m+q}L_p - F_mL_{p+q} = (-1)^m F_q L_{p-m} \tag{2.15}$$

$$L_{m+q}L_p - L_mL_{p+q} = 5(-1)^{m+1} F_q F_{p-m} \tag{2.16}$$

Theorem 2.4. *Let $S(n,m,p,q)$ be the left side of identity (1.1), then:*

$$S(n,m,p,q) = F_{m+q}S(n-1,m,p,q) - F_m S(n-1,m,p+q,q) \tag{2.17}$$

Proof.

$$\begin{aligned}
&F_{m+q}\sum_{k=0}^{n-1}\binom{n-1}{k}(-1)^k F_m^k F_{m+q}^{n-k-1}F_{p+qk}\\
&-F_m\sum_{k=0}^{n-1}\binom{n-1}{k}(-1)^k F_m^k F_{m+q}^{n-k-1}F_{p+q(k+1)}\\
=&F_{m+q}\sum_{k=0}^{n-1}\binom{n-1}{k}(-1)^k F_m^k F_{m+q}^{n-k-1}F_{p+qk}\\
&-F_m\sum_{k=1}^{n}\binom{n-1}{k-1}(-1)^{k-1} F_m^{k-1} F_{m+q}^{n-k}F_{p+qk}\\
=&\sum_{k=0}^{n-1}\binom{n-1}{k}(-1)^k F_m^k F_{m+q}^{n-k}F_{p+qk}\\
&+\sum_{k=1}^{n}\binom{n-1}{k-1}(-1)^k F_m^k F_{m+q}^{n-k}F_{p+qk}
\end{aligned} \tag{2.18}$$

For $0<k<n$ the binomial coefficient addition formula [3] is used:

$$\binom{n-1}{k}+\binom{n-1}{k-1}=\binom{n}{k} \tag{2.19}$$

and for $k=0$ and $k=n$ separately the theorem is proved. □

When $S(n,m,p,q)$ is the left side of identity (1.2) the theorem is:

$$S(n,m,p,q) = L_{m+q}S(n-1,m,p,q) - L_m S(n-1,m,p+q,q) \tag{2.20}$$

and the proof is similar. When $S(n,m,p,q)$ is the right side of identities (1.1) and (1.2) the same recurrence relations are fulfilled.

Theorem 2.5. *Let $S(n,m,p,q)$ be the right side of (1.1), then also:*

$$S(n,m,p,q) = F_{m+q}S(n-1,m,p,q) - F_m S(n-1,m,p+q,q) \tag{2.21}$$

Proof. Using equation (2.11):

$$\begin{aligned}
&F_{m+q}(-1)^{(n-1)m}F_q^{n-1}F_{p-(n-1)m} - F_m(-1)^{(n-1)m}F_q^{n-1}F_{p-(n-1)m+q}\\
=&(-1)^{(n-1)m}F_q^{n-1}(F_{m+q}F_{p-nm+m} - F_m F_{p-nm+m+q})\\
=&(-1)^{nm}F_q^n F_{p-nm}
\end{aligned} \tag{2.22}$$

□

For identity (1.2) the following is needed:

$$F_{p-nm+1} - (-1)^n F_{p-nm-1} = \begin{cases} F_{p-nm} & \text{if } n \text{ is even} \\ L_{p-nm} & \text{if } n \text{ is odd} \end{cases} \tag{2.23}$$

Theorem 2.6. *Let $S(n, m, p, q)$ be the right side of (1.2), then also:*

$$S(n, m, p, q) = L_{m+q} S(n-1, m, p, q) - L_m S(n-1, m, p+q, q) \tag{2.24}$$

Proof. When n is odd and $n-1$ is even, using (2.13) and (2.23):

$$\begin{aligned} & L_{m+q} 5^{\frac{n-1}{2}} F_q^{n-1} F_{p-(n-1)m} - L_m 5^{\frac{n-1}{2}} F_q^{n-1} F_{p-(n-1)m+q} \\ =& 5^{\frac{n-1}{2}} F_q^{n-1} (L_{m+q} F_{p-nm+m} - L_m F_{p-nm+m+q}) \\ =& (-1)^{m+1} 5^{\frac{n-1}{2}} F_q^n L_{p-nm} \end{aligned} \tag{2.25}$$

When n is even and $n-1$ is odd, using (2.16) and (2.23):

$$\begin{aligned} & L_{m+q} 5^{\frac{n-2}{2}} (-1)^{m+1} F_q^{n-1} L_{p-(n-1)m} - L_m 5^{\frac{n-2}{2}} (-1)^{m+1} F_q^{n-1} L_{p-(n-1)m+q} \\ =& 5^{\frac{n}{2}-1} (-1)^{m+1} F_q^{n-1} (L_{m+q} L_{p-nm+m} - L_m L_{p-nm+m+q}) \\ =& 5^{\frac{n}{2}} F_q^n F_{p-nm} \end{aligned} \tag{2.26}$$

□

Having proved the two idenitities for $G = F$, the following is used:

$$G_n = \frac{1}{2}[(G_{-1} + G_1) F_n + G_0 L_n] \tag{2.27}$$

which is easily proved by checking it for $n = 0$ and $n = 1$ and $G_{n+2} = G_{n+1} + G_n$. This means that G_n is a linear combination of F_n and L_n, so it only needs to be checked that the identities hold for $G = F$ and $G = L$. This is demonstrated by adding the identities for $p-1$ and $p+1$ and using $F_{p-1} + F_{p+1} = L_p$.

3 Some Examples

The identities (1.1) to (1.4) with $n = 1$ to $n = 4$ result in the following identities.
Let F_n be the Fibonacci number, L_n be the Lucas number and G_n be the generalized Fibonacci number, for which $G_{n+2} = G_{n+1} + G_n$ with any seed G_0 and G_1 [1, 2, 4].
For integer n, m, p:

$$F_{m+p} G_n - F_m G_{n+p} = (-1)^m F_p G_{n-m} \tag{3.1}$$

$$L_{m+p} G_n - L_m G_{n+p} = (-1)^{m+1} F_p (G_{n-m+1} + G_{n-m-1}) \tag{3.2}$$

$$F_{m+p} G_n - (-1)^p F_m G_{n-p} = F_p G_{n+m} \tag{3.3}$$

$$L_{m+p} G_n - (-1)^p L_m G_{n-p} = F_p (G_{n+m+1} + G_{n+m-1}) \tag{3.4}$$

Identities (3.1) and (3.2) with $G = F$ and $G = L$ are equivalent to identities (19a), (19b), (20a) and (20b) in [4], and the well known identity [1, 2, 3, 4]:

$$F_{m+1}F_n + F_mF_{n-1} = F_{n+m} \tag{3.5}$$

is identity (3.3) with $G = F$ and $p = 1$.
For integer n, m, p:

$$F_{m+p}^2G_n - 2F_mF_{m+p}G_{n+p} + F_m^2G_{n+2p} = F_p^2G_{n-2m} \tag{3.6}$$

$$L_{m+p}^2G_n - 2L_mL_{m+p}G_{n+p} + L_m^2G_{n+2p} = 5F_p^2(G_{n-2m+1} - G_{n-2m-1}) \tag{3.7}$$

$$F_{m+p}^2G_n - 2(-1)^pF_mF_{m+p}G_{n-p} + F_m^2G_{n-2p} = F_p^2G_{n+2m} \tag{3.8}$$

$$L_{m+p}^2G_n - 2(-1)^pL_mL_{m+p}G_{n-p} + L_m^2G_{n-2p} = 5F_p^2(G_{n+2m+1} - G_{n+2m-1}) \tag{3.9}$$

For integer n, m, p:

$$F_{m+p}^3G_n - 3F_mF_{m+p}^2G_{n+p} + 3F_m^2F_{m+p}G_{n+2p} - F_m^3G_{n+3p} = (-1)^mF_p^3G_{n-3m} \tag{3.10}$$

$$\begin{aligned} &L_{m+p}^3G_n - 3L_mL_{m+p}^2G_{n+p} + 3L_m^2L_{m+p}G_{n+2p} - L_m^3G_{n+3p} \\ &= 5(-1)^{m+1}F_p^3(G_{n-3m+1} + G_{n-3m-1}) \end{aligned} \tag{3.11}$$

$$\begin{aligned} &F_{m+p}^3G_n - 3(-1)^pF_mF_{m+p}^2G_{n-p} + 3F_m^2F_{m+p}G_{n-2p} - (-1)^pF_m^3G_{n-3p} \\ &= F_p^3G_{n+3m} \end{aligned} \tag{3.12}$$

$$\begin{aligned} &L_{m+p}^3G_n - 3(-1)^pL_mL_{m+p}^2G_{n-p} + 3L_m^2L_{m+p}G_{n-2p} - (-1)^pL_m^3G_{n-3p} \\ &= 5F_p^3(G_{n+3m+1} + G_{n+3m-1}) \end{aligned} \tag{3.13}$$

For integer n, m, p:

$$\begin{aligned} &F_{m+p}^4G_n - 4F_mF_{m+p}^3G_{n+p} + 6F_m^2F_{m+p}^2G_{n+2p} - 4F_m^3F_{m+p}G_{n+3p} \\ &+ F_m^4G_{n+4p} = F_p^4G_{n-4m} \end{aligned} \tag{3.14}$$

$$\begin{aligned} &L_{m+p}^4G_n - 4L_mL_{m+p}^3G_{n+p} + 6L_m^2L_{m+p}^2G_{n+2p} - 4L_m^3L_{m+p}G_{n+3p} \\ &+ L_m^4G_{n+4p} = 25F_p^4(G_{n-4m+1} - G_{n-4m-1}) \end{aligned} \tag{3.15}$$

$$\begin{aligned} &F_{m+p}^4G_n - 4(-1)^pF_mF_{m+p}^3G_{n-p} + 6F_m^2F_{m+p}^2G_{n-2p} \\ &- 4(-1)^pF_m^3F_{m+p}G_{n-3p} + F_m^4G_{n-4p} = F_p^4G_{n+4m} \end{aligned} \tag{3.16}$$

$$\begin{aligned} &L_{m+p}^4G_n - 4(-1)^pL_mL_{m+p}^3G_{n-p} + 6L_m^2L_{m+p}^2G_{n-2p} \\ &- 4(-1)^pL_m^3L_{m+p}G_{n-3p} + L_m^4G_{n-4p} = 25F_p^4(G_{n+4m+1} - G_{n+4m-1}) \end{aligned} \tag{3.17}$$

References

[1] A.T. Benjamin, J.J. Quinn, *Proofs that Really Count*, The Mathematical Association of America, 2003.

[2] P. Chandra, E.W. Weisstein, *Fibonacci Number*. From Mathworld - A Wolfram Web Resource. `http://mathworld.wolfram.com/FibonacciNumber.html`

[3] R.L. Graham, D.E. Knuth, O. Patashnik, *Concrete Mathematics, A Foundation for Computer Science*, 2nd ed., Addison-Wesley, 1994.

[4] S. Vajda, *Fibonacci and Lucas Numbers, and the Golden Section*, Dover Publications, 1989.