

STABILITY OF ψ -HILFER IMPULSIVE FRACTIONAL DIFFERENTIAL EQUATIONS

J. VANTERLER DA C. SOUSA, KISHOR D. KUCCHE¹, AND E. CAPELAS DE OLIVEIRA

ABSTRACT. In this paper, we investigate the sufficient conditions for existence and uniqueness of solutions and δ -Ulam-Hyers-Rassias stability of an impulsive fractional differential equation involving ψ -Hilfer fractional derivative. Fixed point approach is used to obtain our main results.

Keywords: Impulsive fractional differential equation, δ -Ulam-Hyers-Rassias, ψ -Hilfer fractional derivative, Banach fixed point.

MSC 2010 subject classifications. 26A33, 34A08, 34A12, 34A37, 34DXX.

1. INTRODUCTION

Impulsive differential equations are used to describe the evolutionary processes that abruptly change their state at a certain moment. This subject received great importance and remarkable attention from the researchers because of its rich theory [1] and applicability in various branches of science and technology. A natural framework for mathematical modeling of many physical phenomena appearing in the field of mechanics, ecology, medicine, biology and electrical engineering can be provided via impulsive differential equations.

Wang and Zhang [2] investigated the existence and uniqueness of solutions to differential equations with not instantaneous impulses in a $P\beta$ -normed space of the form:

$$\begin{cases} x'(t) = f(t, x(t)), & t \in (s_i, t_{i+1}], \quad i = 0, 1, \dots, m, \\ x(t) = g_i(t, x(t)), & t \in (t_i, s_i], \quad i = 1, 2, \dots, m. \end{cases}$$

Wang et al. in [3], considering the ordinary nonlinear differential equation with not instantaneous impulses obtained existence and uniqueness of solutions and introduced an interesting concept of stability viz. generalized β -Ulam-Hyers-Rassias. Zeng et al. [4] extended the above investigations to the class of impulsive integro-differential equations with not instantaneous impulses.

With the expansion of the fractional calculus [5, 6, 7], the impulsive fractional differential equations gained a much attention and began to be studied, mainly due to the variety of results, from the stability study, existence to uniqueness [8, 9, 10, 11, 12, 13]. Impulsive differential equations in the space of Banach spaces have been dealt in [14, 15, 16, 17].

Discussions on concept of solutions, existence and uniqueness results pertaining to impulsive fractional Cauchy differential equations (IFDE) can also be found in the survey paper [18].

Wang and Zhang [19] recently, introduced a new class of nonlinear differential equations with fractional integrable impulses and established existence and uniqueness results and introduced new concepts of Bielecki-Ulam's type stability.

As discussed above, in the last years there are many published works related to stability, especially on impulsive differential equations of arbitrary order. Thus, one of the mains purpose of this paper is to expand this range of works in the literature involving stability and thus contribute to the expansion of the area.

In this paper, we apply fixed point approach to study stability of the modified impulsive fractional differential equations

$$(1.1) \quad \begin{cases} {}^H\mathbb{D}_{0+}^{\alpha,\beta;\psi} x(t) &= f(t, x(t)), \quad t \in (s_i, t_{i+1}], \quad i = 0, 1, \dots, m \\ x(t) &= g_i(t, x(t_i^+)), \quad t \in (t_i, s_i], \quad i = 1, 2, \dots, m \end{cases}$$

where ${}^H\mathbb{D}_{0+}^{\alpha,\beta;\psi}(\cdot)$ is the ψ -Hilfer fractional derivative with $0 < \alpha \leq 1$, $0 \leq \beta \leq 1$ and $0 = t_0 = s_0 < t_1 \leq s_1 \leq t_2 < \dots < t_m \leq s_m < t_{m+1} = T$ are prefixed numbers, $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and $g_i : [t_i, s_i] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous for all $i = 1, 2, \dots, m$ which is not instantaneous impulses.

The motivation for the elaboration of this paper is the contribution in the stability of fractional differential equations, in particular of the impulsive type. In this sense, as the main purpose of this paper, we investigated the δ -Ulam-Hyers-Rassias stability of the impulsive fractional differential equation by employing the fixed point approach.

For the fundamental properties of ψ -Hilfer fractional derivative and the basic theory of fractional differential equation involving ψ -Hilfer fractional derivative, we refer the readers to the papers of Sousa and Oliveira [7, 20].

This paper is divided as follows: in Section 2, we present the concepts of weighted and piecewise weighted function spaces. We also recall the definitions of ψ -Riemann-Liouville fractional integral, ψ -Hilfer fractional derivative and define the concept of generalized δ -Ulam-Hyers-Rassias stability. In Section 3, we investigate through the Theorem 2 the generalized δ -Ulam-Hyers-Rassias stability of the fractional differential equation.

2. PRELIMINARIES

Definition 1. [3] Suppose E is a vector space over \mathbb{K} . A function $\|\cdot\|_\delta : E \rightarrow [0, \infty)$ ($0 < \delta \leq 1$) is called a δ -norm if and only if it satisfies :

- (1) $\|x\|_\delta = 0$, if and only if $x = 0$;
- (2) $\|\lambda x\|_\delta = |\lambda|^\delta \|x\|_\delta$ for all $\lambda \in K$ and all $x \in E$;
- (3) $\|x + y\|_\delta \leq \|x\|_\delta + \|y\|_\delta$.

Let $J = [0, T]$, $J' = (0, T]$ and $C(J, \mathbb{R})$ the space of continuous functions. The weighted space $C_{1-\gamma;\psi}(J, \mathbb{R})$ of functions x on J' is defined by [7]

$$C_{1-\gamma;\psi}(J, \mathbb{R}) = \{x \in C(J', \mathbb{R}); (\psi(t) - \psi(0))^{1-\gamma} x(t) \in C(J, \mathbb{R})\}, \quad 0 \leq \gamma < 1$$

with the norm

$$\|x\|_{C_{1-\gamma;\psi}, \delta} = \sup_{t \in J'} \{(\psi(t) - \psi(0))^{1-\gamma} \|x(t)\|_{\delta}\}.$$

Obviously the space $C_{1-\gamma;\psi}(J, \mathbb{R})$ is a Banach space.

The piecewise weighted space $PC_{1-\gamma;\psi}(J, \mathbb{R})$ of functions x on $C((t_k, t_{k+1}], \mathbb{R})$ is defined by

$$PC_{1-\gamma;\psi}(J, \mathbb{R}) = \left\{ \begin{array}{l} (\psi(t) - \psi(t_k) - \psi(0))^{1-\gamma} x(t) \in C((t_k, t_{k+1}], \mathbb{R}) \text{ and} \\ \lim_{t \rightarrow t_k} (\psi(t) - \psi(t_k) - \psi(0))^{1-\gamma} x(t), \text{ exists for } k = 1, 2, \dots, m \end{array} \right\}$$

with norm

$$\|x\|_{PC_{1-\gamma;\psi}, \delta} := \max_{k=1,2,\dots,m} \left\{ \sup_{t \in (t_k, t_{k+1}]} (\psi(t) - \psi(t_k) - \psi(0))^{1-\gamma} \|x(t)\|_{\delta} \right\}$$

and there exists $x(t_k^-)$ and $x(t_k^+)$, $k = 1, 2, \dots, m$ with $x(t_k^-) = x(t_k^+)$. The space $PC_{1-\gamma;\psi}(J, \mathbb{R})$ is also a Banach space.

Let $n - 1 < \alpha \leq n$ with $n \in \mathbb{N}$, $J = [a, b]$ be an interval such that $-\infty \leq a < b \leq +\infty$ and let $f, \psi \in C^n([a, b], \mathbb{R})$ be two functions such that ψ is increasing and $\psi'(t) \neq 0$, for all $t \in J$. The ψ -Hilfer fractional derivative denoted by [7]

$${}^H\mathbb{D}_{0+}^{\alpha,\beta;\psi} y(t) = I_{0+}^{\beta(n-\alpha);\psi} \left(\frac{1}{\psi'(t)} \frac{d}{dt} \right)^n I_{0+}^{(1-\beta)(n-\alpha);\psi} y(t),$$

where $I_{0+}^{\xi;\psi}(\cdot)$ ($0 < \xi \leq 1$) is the ψ -Riemann-Liouville fractional integral [7].

Banach's fixed-point theorem is fundamental in this study, so we will enunciate it below.

Theorem 1. [3, 21] *Let (X, d) be a generalized complete metric space. Assume that $\Omega : X \rightarrow X$ is a strictly contractive operator with the Lipschitz constant $L < 1$. If there exists a nonnegative integer k such that $d(\Omega^{k+1}, \Omega^k) < \infty$ for some $x \in X$, then the following are true:*

- (1) *The sequence $\{\Omega^k x\}$ converges to a fixed point x^* of Ω ;*
- (2) *x^* is the unique fixed point of Ω in $\Omega^* = \{y \in X/d(\Omega^k x, y) < \infty\}$;*
- (3) *If $y \in X^*$, then*

$$d(y, x^*) \leq \frac{1}{1-L} d(\Omega y, y).$$

Let the space of piecewise weighted space continuous functions

$$X = \{g : J \rightarrow \mathbb{R}/g \in PC_{1-\gamma;\psi}(J, \mathbb{R}), \quad 0 \leq \gamma < 1\},$$

with the generalized metric on X given by

(2.1)

$$d(g, h) = \inf \left\{ C_1 + C_2 \in [0, \infty] / |g(t) - h(t)|^\delta \leq (C_1 + C_2) (\varphi^\delta(t) + \varepsilon^\delta), \text{ for all } t \in J \right\}$$

where $C_1 \in \left\{ C \in [0, \infty] / |g(t) - h(t)|^\delta \leq C\varphi^\delta(t), \text{ for all } t \in (s_i, t_{i+1}], i = 0, 1, \dots, m \right\}$ and $C_2 \in \left\{ C \in [0, \infty] / |g(t) - h(t)|^\delta \leq C\varepsilon^\delta, \text{ for all } t \in (t_i, s_i], i = 1, \dots, m \right\}$. Note that that (X, d) is a complete generalized metric space.

The function $x \in \tilde{U} := PC_{1-\gamma;\psi}(J, \mathbb{R}) \bigcap_{i=0}^m C^1((s_i, t_{i+1}], \mathbb{R})$ is a solution of the impulsive fractional differential equations

$$\begin{cases} {}^H\mathbb{D}_{0+}^{\alpha,\beta;\psi} x(t) &= f(t, x(t)), \quad t \in (s_i, t_{i+1}], \quad i = 0, 1, \dots, m \\ x(t) &= g_i(t, x(t_i^+)), \quad t \in (t_i, s_i], \quad i = 1, 2, \dots, m \\ I_{0+}^{1-\gamma;\psi} x(0) &= x_0 \in \mathbb{R} \end{cases}$$

where $I_{0+}^{1-\gamma;\psi}(\cdot)$ is the ψ -Riemann-Liouville fractional integral with $\gamma = \alpha + \beta(1 - \alpha)$, if x satisfies $I_{0+}^{1-\gamma;\psi} x(0) = x_0$, $x(t) = g_i(t, x(t_i^+))$, $t \in (t_i, s_i]$, $i = 1, 2, \dots, m$ and

$$\begin{cases} x(t) &= \Psi^\lambda(t, 0)x_0 + \frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) f(s, x(s)) ds, \quad t \in [0, t_1] \\ x(t) &= g_i(s_i, x(t_i^+)) + \frac{1}{\Gamma(\alpha)} \int_{s_i}^t N_\psi^\alpha(t, s) f(s, x(s)) ds, \quad t \in (s_i, t_{i+1}], \quad i = 1, 2, \dots, m. \end{cases}$$

with $N_\psi^\alpha(s, t) := \psi'(s) (\psi(t) - \psi(s))^{\alpha-1}$ and $\Psi^\lambda(t, 0) = \frac{(\psi(t) - \psi(0))^{\gamma-1}}{\Gamma(\gamma)}$.

Let $0 < \delta \leq 1$, $\xi \geq 0$, $\varphi \in PC_{1-\gamma;\psi}(J, \mathbb{R}_+)$ is nondecreasing and

$$(2.2) \quad \begin{cases} \left| {}^H\mathbb{D}_{0+}^{\alpha,\beta;\psi} y(t) - f(t, y(t)) \right| &\leq \varphi(t), \quad t \in (s_i, t_{i+1}], \quad i = 0, 1, \dots, m \\ |y(t) - g_i(t, y(t_i^+))| &\leq \xi, \quad t \in (t_i, s_i], \quad i = 1, 2, \dots, m \end{cases}$$

Definition 2. The Eq.(1.1) is generalized δ -Ulam-Hyers-Rassias stable with respect to (φ, ξ) if there exists $C_{f,\delta,g_i,\varphi} > 0$ such that for each solution $y \in \tilde{U}$ of the inequality Eq.(2.2) there exists a solution $x \in \tilde{U}$ of the Eq.(1.1) with

$$|y(t) - x(t)|^\delta \leq C_{f,\delta,g_i,\varphi} (\xi^\delta + \varphi^\delta(t)), \quad t \in J.$$

A function $y \in \tilde{U}$ is a solution of the inequality Eq.(2.2) if and only if there is $G \in \bigcap_{i=0}^m C^1((s_i, t_{i+1}], \mathbb{R})$ and $g \in \bigcap_{i=0}^m C([t_i, s_i], \mathbb{R})$, such that:

$$(a) \quad |G(t)| \leq \varphi(t), \quad t \in \bigcup_{i=0}^m (s_i, t_{i+1}] \text{ and } |g(t)| \leq \xi, \quad t \in \bigcup_{i=0}^m (t_i, s_i];$$

$$(b) {}^H\mathbb{D}_{0+}^{\alpha,\beta;\psi} y(t) = f(t, y(t)) + G(t), \quad t \in (s_i, t_{i+1}], \quad i = 0, 1, \dots, m;$$

$$(c) y(t) = g_i(t, y(t_i^+)) + g(t), \quad t \in (t_i, s_i], \quad i = 1, 2, \dots, m.$$

Remark 1. If $y \in \tilde{U}$ is a solution of the inequality Eq.(2.2) the y is a solution of the fractional integral inequality:

$$(2.3) \quad |y(t) - g_i(t, y(t_i^+))| \leq \xi, \quad t \in (t_i, s_i], \quad i = 0, 1, \dots, m$$

and

$$\left| y(t) - \Psi^\lambda(t, 0)y(0) - \frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) f(s, y(s)) ds \right| \leq \frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) \varphi(s) ds, \quad t \in [0, t_1]$$

and

$$(2.4) \quad \begin{aligned} & \left| y(t) - g_i(s_i, y(t_i^+)) - \frac{1}{\Gamma(\alpha)} \int_{s_i}^t N_\psi^\alpha(t, s) f(s, y(s)) ds \right| \\ & \leq \xi + \frac{1}{\Gamma(\alpha)} \int_{s_i}^t N_\psi^\alpha(t, s) \varphi(s) ds, \quad t \in (s_i, t_{i+1}], \quad i = 1, 2, \dots, m. \end{aligned}$$

3. δ -ULAM-HYERS-RASSIAS STABILITY

In this section, we present the main result of this paper, the stability of the type generalized δ -Ulam-Hyers-Rassias for the impulsive fractional differential equation Eq.(1.1), by means Banach's the fixed point theorem.

Before investigating the main result in this paper, we list some essential conditions for proof of the theorem:

$$(H1) \quad f \in C_{1-\gamma;\psi}(J \times \mathbb{R}, \mathbb{R});$$

$$(H2) \quad \text{There exists a positive constant } L_f \text{ such that}$$

$$|f(t, u_1) - f(t, u_2)| \leq L_f |u_1 - u_2|, \quad \text{for } t \in J \text{ and } u_1, u_2 \in \mathbb{R};$$

$$(H3) \quad g_i \in C_{1-\gamma;\psi}([t_i, s_i] \times \mathbb{R}, \mathbb{R}) \text{ and there are positive constants } L_{g_i}, \quad i = 1, 2, \dots, m \text{ such that}$$

$$|g_i(t, u_1) - g_i(t, u_2)| \leq L_{g_i} |u_1 - u_2|, \quad \text{for } t \in [t_i, s_i] \text{ and } u_1, u_2 \in \mathbb{R};$$

$$(H4) \quad \text{Let } \varphi \in C_{1-\gamma;\psi}(J, \mathbb{R}_+) \text{ be a nondecreasing function. There exists } C_\varphi > 0 \text{ such that}$$

$$\frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) \varphi(s) ds \leq c_\varphi \varphi(t), \quad \text{for } t \in J.$$

Theorem 2. Assume the conditions (H1)-(H4) be satisfies. If there exists a function $y \in \widetilde{U}$ satisfying Eq.(2.2), then there exists a unique solution $y_0 : J \rightarrow \mathbb{R}$ such that

$$(3.1) \quad y_0(t) = \begin{cases} \Psi^\lambda(t, 0)x(0) + \frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) f(s, y_0(s)) ds, & t \in [0, t_1] \\ g_i(t, y_0(t_i^+)), & t \in (t_i, s_i], \quad i = 1, 2, \dots, m \\ g_i(s_i, y_0(t_i^+)) + \frac{1}{\Gamma(\alpha)} \int_{s_i}^t N_\psi^\alpha(t, s) f(s, y_0(s)) ds, & t \in (s_i, t_{i+1}] \\ i = 1, 2, \dots, m. \end{cases}$$

and

$$(3.2) \quad |y(t) - y_0(t)|^\delta \leq \frac{(1 + C_\varphi^\delta)(\varphi^\delta(t) + \xi^\delta)}{1 - \Phi}, \quad t \in J$$

where

$$(3.3) \quad \Phi := \max_{i=1,2,\dots,m} \{L_{g_i}^\delta + L_f^\delta C_\varphi^\delta\}.$$

Proof. For prove of this result, consider the operator $\Omega : X \rightarrow X$ given by

$$(3.4) \quad \Omega x(t) = \begin{cases} \Psi^\lambda(t, 0)x(0) + \frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) f(s, x_0(s)) ds, & t \in [0, t_1] \\ g_i(t, x_0(t_i^+)), & t \in (t_i, s_i], \quad i = 1, 2, \dots, m \\ g_i(s_i, x_0(t_i^+)) + \frac{1}{\Gamma(\alpha)} \int_{s_i}^t N_\psi^\alpha(t, s) f(s, x_0(s)) ds, & \\ t \in (s_i, t_{i+1}], \quad i = 1, 2, \dots, m. \end{cases}$$

for all $x \in X$ and $t \in [0, T]$. Note that, Ω is a well defined operator according with condition (H1).

In order to use Banach's fixed point theorem, we prove that Ω is strictly contractive on X , in three cases. Note that

$$|g(t) - h(t)|^\delta \leq \begin{cases} C_1 \varphi^\delta(t), & t \in (s_i, t_{i+1}], \quad i = 0, 1, \dots, m \\ C_2 \xi^\delta, & t \in (t_i, s_i], \quad i = 1, \dots, m \end{cases}$$

is equivalent to

$$(3.5) \quad |g(t) - h(t)| \leq \begin{cases} C_1^\delta \varphi(t), & t \in (s_i, t_{i+1}], \quad i = 0, 1, \dots, m \\ C_2^\delta \xi, & t \in (t_i, s_i], \quad i = 1, \dots, m \end{cases}$$

Using the definition of Ω in Eq.(3.4), (H2), (H3) and Eq.(3.5), we have the following cases:

Case 1: For $t \in [0, t_1]$, and by the hypothesis (H2), (H4) and Eq.(3.5), we get

$$\begin{aligned} |\Omega g(t) - \Omega h(t)|^\delta &\leq \left| \frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) f(s, g(s)) ds - \frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) f(s, h(s)) ds \right|^\delta \\ &\leq L_f^\delta \left(\frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) |g(s) - h(s)| ds \right)^\delta \leq L_f^\delta C_1 c_\varphi^\delta \varphi(t)^\delta. \end{aligned}$$

Case 2: For $t \in (t_i, s_i]$, and by the hypothesis (H3) and Eq.(3.4), we obtain

$$|\Omega g(t) - \Omega h(t)|^\delta \leq |g_i(t, g(t_i^+)) - g_i(t, h(t_i^+))|^\delta \leq (L_{g_i} |g(t_i^+) - h(t_i^+)|)^\delta \leq L_{g_i}^\delta C_2 \xi^\delta.$$

Case 3: For $t \in (s_i, t_{i+1}]$ and using the hypothesis (H1), (H2), (H3) and Eq.(3.4), we have

$$\begin{aligned} &|\Omega g(t) - \Omega h(t)|^\delta \\ &\leq |g_i(s_i, g(t_i^+)) - g_i(s_i, h(t_i^+))|^\delta + \left| \frac{1}{\Gamma(\alpha)} \int_{s_i}^t N_\psi^\alpha(t, s) (f(s, g(s)) - f(s, h(s))) ds \right|^\delta \\ &\leq L_{g_i}^\delta C_2 \xi^\delta + L_f^\delta C_1 \left(\frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) \varphi(s) ds \right)^\delta \leq (L_{g_i}^\delta + L_f^\delta c_\varphi^\delta) (C_1 + C_2) (\varphi^\delta(t) + \xi^\delta). \end{aligned}$$

Then, we obtain

$$\begin{aligned} |\Omega g(t) - \Omega h(t)|^\delta &\leq \max_{i=1,2,\dots,m} (L_{g_i}^\delta + L_f^\delta c_\varphi^\delta) (C_1 + C_2) (\varphi^\delta(t) + \xi^\delta) \\ &= \Phi (C_1 + C_2) (\varphi^\delta(t) + \xi^\delta), \quad t \in J. \end{aligned}$$

Hence, we get

$$d(\Omega g, \Omega h) \leq \Phi d(g, h)$$

for any $g, h \in X$ and since the condition Eq.(3.3).

Now, we take $g_0 \in X$ and from the piecewise continuous property of g_0 and Ωg_0 , then there exists a constant $0 < G_1 < \infty$ so that

$$\begin{aligned} |\Omega g_0(t) - g_0(t)|^\delta &= \left| \Psi^\lambda(t, 0)x(0) + \frac{1}{\Gamma(\alpha)} \int_0^t N_\psi^\alpha(t, s) f(s, g_0(s)) ds - g_0(t) \right|^\delta \\ &\leq G_1 \varphi^\delta(t) \leq G_1 (\varphi^\delta(t) + \xi^\delta), \quad t \in [0, t_1]. \end{aligned}$$

On the other hand, also G_2 and G_3 with $0 < G_2 < \infty$ and $0 < G_3 < \infty$, such that,

$$\begin{aligned} |\Omega g_0(t) - g_0(t)|^\delta &= |g_i(t, g_0(t_i^+)) - g_0(t)|^\delta \\ &\leq G_2 \xi^\delta \leq G_2 (\varphi^\delta(t) + \xi^\delta), \quad t \in (t_i, s_i], \quad i = 1, 2, \dots, m \end{aligned}$$

and

$$\begin{aligned} |\Omega g_0(t) - g_0(t)|^\delta &= \left| g_i(s_i, g_0(t_i^+)) + \frac{1}{\Gamma(\alpha)} \int_{s_i}^t N_\psi^\alpha(t, s) f(s, g_0(s)) ds - g_0(t) \right|^\delta \\ &\leq G_3(\varphi^\delta(t) + \xi^\delta), \quad t \in (s_i, t_{i+1}], \quad i = 1, 2, \dots, m \end{aligned}$$

since f, g_i and g_0 are bounded on J and $\varphi(\cdot) + \xi^\delta > 0$. In this sense, Eq.(2.1) implies that

$$d(\Omega g_0, g_0) < \infty.$$

Note that, exists a continuous function $y_0 : J \rightarrow \mathbb{R}$ such that $\Omega^n y_0 \rightarrow y_0$ in (X, d) as $n \rightarrow \infty$ and $\Omega y_0 = y_0$, that is y_0 satisfies Eq.(3.4) for every $t \in J$ (Banach's fixed point theorem).

For finally the proof this theorem, we check that $0 < C_g < \infty$ such that

$$|g_0(t) - g(t)|^\delta \leq C_g(\varphi^\delta(t) + \xi^\delta), \quad \text{for any } t \in J$$

and assuming that g, g_0 are bounded on J and $\min_{t \in J}(\varphi^\delta(t) + \xi^\delta) > 0$.

Then, we get $d(g_0, g) < \infty$ for all $g \in X$, that is $X = \{g \in X / d(g_0, g) < \infty\}$. Therefore, we obtain that y_0 is the unique solution continuous function with the property Eq.(3.4).

On the other hand, using the hypotheses (H1)-(H4), Eq.(2.3), Eq.(2.4) and Eq.(2.4) it follows that

$$(3.6) \quad d(y, \Omega y) \leq 1 + C_\varphi^\delta.$$

Thus, from Eq.(3.6), we have

$$d(y, y_0) \leq \frac{d(\Omega y, y)}{1 - \Phi} \leq \frac{1 + C_\varphi^\delta}{1 - \Phi},$$

which means that Eq.(3.2) is true for $t \in J$. □

REFERENCES

- [1] V. Lakshmikantham, P. S. Simeonov, Theory of impulsive differential equations, Vol. 6, World scientific, 1989.
- [2] J. Wang, Y. Zhang, Existence and stability of solutions to nonlinear impulsive differential equations in β -normed spaces, Electronic J. Diff. Equations 2014 (83) (2014) 1–10.
- [3] J. Wang, Z. Lin, Y. Zhou, On the stability of new impulsive ordinary differential equations, Topol. Methods Nonlinear Anal. 46 (1) (2015) 303–314.
- [4] Z. Lin, W. Wei, J. Wang, Existence and stability results for impulsive integro-differential equations, Facta Universitatis 29 (2) (2014) 119–130.
- [5] S. G. Samko, A. A. Kilbas, O. I. Marichev, Fractional integrals and derivatives, Theory and Applications, Gordon and Breach, Yverdon 1993 (1993) 44.
- [6] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, Theory and Applications of Fractional Differential Equations, Vol. 204, Elsevier, Amsterdam, 2006.
- [7] J. Vanterler da C. Sousa, E. Capelas de Oliveira, On the ψ -Hilfer fractional derivative, Commun. Nonlinear Sci. Numer. Simulat. 60 (2018) 72–91.

- [8] Q. Wang, D. Lu, Y. Fang, Stability analysis of impulsive fractional differential systems with delay, *Appl. Math. Lett.* 40 (2015) 1–6.
- [9] I. Stamova, Global stability of impulsive fractional differential equations, *Appl. Math. and Comput.* 237 (2014) 605–612.
- [10] K. Liu, W. Jiang, Stability of nonlinear Caputo fractional differential equations, *Appl. Math. Modelling* 40 (5-6) (2016) 3919–3924.
- [11] J. Wang, W. Wei, Y. Yang, On some impulsive fractional differential equations in Banach spaces, *Opuscula Mathematica* 30 (4) (2010) 507–525.
- [12] T. L. Guo, Nonlinear impulsive fractional differential equations in Banach spaces, *Topol. Methods Nonlinear Anal.* 42 (1) (2013) 221–232.
- [13] J. Vanterler da C. Sousa, D. S. Oliveira, E. Capelas de Oliveira, On the existence and stability for impulsive fractional integrodifferential equation, *arXiv:1806.01442*, (2018).
- [14] A. Sivasankari, A. Leelamani, Existence of mild solutions for an impulsive fractional neutral integro-differential equations with non-local conditions in Banach spaces., *Nonlinear Studies* 24 (3).
- [15] P. Kumar, R. Haloi, D. Bahuguna, D. N. Pandey, Existence of solutions to a new class of abstract non-instantaneous impulsive fractional integro-differential equations, *Nonl. Dyn. and Systems Theory* (2016) 73.
- [16] M. Benchohra, D. Seba, Impulsive fractional differential equations in Banach spaces, *Electron. J. Qual. Theory Differ. Equ* 8 (1).
- [17] H. Gou, B. Li, Local and global existence of mild solution to impulsive fractional semilinear integro-differential equation with noncompact semigroup, *Commun. Nonlinear Sci. Numer. Simulat.* 42 (2017) 204–214.
- [18] J. Wang, M. M. Feckan, Y. Zhou, A survey on impulsive fractional differential equations, *Fract. Calc. Appl. Anal.* 16 (4) (2016) 806–831.
- [19] J. Wang, Y. Zhang, A class of nonlinear differential equations with fractional integrable impulses, *Commun. Nonlinear Sci. Numer. Simulat.* 19 (2) (2014) 3001–3010.
- [20] J. Vanterler da C. Sousa, E. Capelas de Oliveira, A Gronwall inequality and the Cauchy-type problem by means of ψ -Hilfer operator, *arXiv:1709.03634*.
- [21] J. Vanterler da C. Sousa, E. Capelas de Oliveira, Ulam—Hyers stability of a nonlinear fractional Volterra integro-differential equation, *Appl. Math. Lett.* 81 (2018) 50–56.

DEPARTMENT OF APPLIED MATHEMATICS, INSTITUTE OF MATHEMATICS, STATISTICS AND SCIENTIFIC COMPUTATION, UNIVERSITY OF CAMPINAS – UNICAMP, RUA SÉRGIO BUARQUE DE HOLANDA 651, 13083–859, CAMPINAS SP, BRAZIL

E-MAIL: *vanterlermatematico@hotmail.com*, *capelas@ime.unicamp.br*

¹ DEPARTMENT OF MATHEMATICS, SHIVAJI UNIVERSITY, KOLHAPUR 416 004, MAHARASHTRA, INDIA

E-MAIL: *kdkucche@gmail.com*