

# Special cases : moons, rings, comets, trojans

Juan Cabrera, María Fernández Jiménez, Antonio García Muñoz, Jean Schneider

**Abstract** Non-planetary bodies provide valuable insight into our current understanding of planetary formation and evolution. Although these objects are challenging to detect and characterize, the potential information to be drawn from them has motivated various searches through a number of techniques. Here, we briefly review the current status in the search of moons, rings, comets, and trojans in exoplanet systems and suggest what future discoveries may occur in the near future.

## Introduction

It is not original admitting that it is complicated to make accurate predictions about the future. There is a quote, attributed to Napoleon Bonaparte, warning against insisting too much in having the full control of the present status of a problem and a detailed plan for the future developments before starting to do the work (*Celui qui, au départ, insiste pour savoir où il va, quand il part et par où il passe n'ira pas loin*). However, considering the importance of the investment required to answer certain scientific questions, it is mandatory to have a realistic idea of the likelihood

---

Juan Cabrera

Institut für Planetenforschung, Deutsches Zentrum für Luft- und Raumfahrt, Rutherfordstr. 2, Berlin, Germany, e-mail: [juan.cabrera@dlr.de](mailto:juan.cabrera@dlr.de)

María Fernández Jiménez

Institut für Planetenforschung, Deutsches Zentrum für Luft- und Raumfahrt, Rutherfordstr. 2, Berlin, Germany, e-mail: [maria.fernandez.jimene@gmail.com](mailto:maria.fernandez.jimene@gmail.com)

Antonio García Muñoz

Zentrum für Astronomie und Astrophysik, TU Berlin, Hardenbergstrae 36, D-10623 Berlin, Germany, e-mail: [garciamunoz@astro.physik.tu-berlin.de](mailto:garciamunoz@astro.physik.tu-berlin.de)

Jean Schneider

LUTH, Observatoire de Paris, CNRS, Université Paris Diderot; 5 place Jules Janssen, 92195 Meudon, France, e-mail: [jean.schneider@obspm.fr](mailto:jean.schneider@obspm.fr)

of success of the research. These considerations apply to the topic of this chapter: the search and characterization of moons, rings, comets, and trojans in exoplanetary systems.

We know from the solar system that satellites and ring systems, minor planets, and comets provide meaningful insights into the processes of planetary formation and evolution (de Pater and Lissauer 2015). Many of them are interesting objects of research on their own, in particular considering their prospects of habitability. However, in view of the difficulty of categorically proving, or ruling out, the existence of life on other worlds in our solar system (i.e. Waite et al. 2017), one can rightfully wonder about the possibilities of actually finding life in extrasolar systems (for a summary of the technical difficulties, see Schneider et al. 2010; for a recent review on habitability, see Cockell et al. 2016).

In the following Sections we briefly discuss what to expect from near future researches about extrasolar systems of moons, rings, comets, and trojan minor planets.

## Exomoons

In this volume there is an excellent review on exomoons and ring detections by R. Heller, therefore we have orientated this chapter towards complementary aspects. Additionally, we refer the reader to some recent reviews on the topic by Barr (2016); Kipping et al. (2014a); Schneider et al. (2015); Sinukoff et al. (2013). In order not to dwell long on topics already addressed by previous reviews, we will only briefly discuss processes of exomoon formation and evolution before addressing the present status of discoveries and our expectations for the future.

The research on the processes leading to the formation of exomoons has benefited from studies applied to the solar system (see Heller and Pudritz 2015; Miguel and Ida 2016; Ogihara and Ida 2012; Crida and Charnoz 2012). However, exomoons are expected to be found in different environments depending on the details of their evolution in the disk (Fujii et al. 2017), the outcome of scattering processes (Gong et al. 2013), capture (Ochiai et al. 2014) or collisions (Barr and Bruck Syal 2017), to name a few. Exotic situations where planets are ejected from the planetary system conserving their moons have been mentioned, though their detection is extremely challenging in this configuration (Laughlin and Adams 2000).

Moons exist between the Roche lobe and the Hill radius of their host planets (Murray and Dermott 2000). There are numerous studies researching the dynamical stability and the tidal evolution of moons (Adams and Bloch 2016; Barnes and O'Brien 2002; Debes and Sigurdsson 2007; Domingos et al. 2006; Donnison 2010; Hong et al. 2015; Namouni 2010; Payne et al. 2013; Sasaki et al. 2012; Sasaki and Barnes 2014). Consequently, their final configuration will depend on planetary processes like migration (Spalding et al. 2016), photoevaporation (Yang et al. 2016), or tidal interactions (Cassidy et al. 2009), all of them open to a certain degree of interpretation today. Rather than a disadvantage this is an encouragement to study moons, as they could provide useful measurable constraints. However, the expected

diversity requires observational support to understand the relative impact of the different processes proposed.

Habitability is another important reason to look for exomoons (Kaltenegger 2010; Lammer et al. 2014). There are interesting processes that are exclusive of these systems and that deserve specific attention. They include tidal interactions (Dobos et al. 2017; Forgan and Kipping 2013; Heller 2012; Scharf 2006), planetary illumination (Forgan and Yotov 2014), amount of volatiles depending on the formation and migration mechanisms (Heller and Pudritz 2015; Heller and Barnes 2015). Finally, the possible presence of moons might impact the interpretation of biosignatures (Rein et al. 2014; Li et al. 2016).

### ***Current status of detections***

There is no known reason preventing exomoons from existing and we expect them to be present in many different configurations. Therefore, all detection methods applied to exoplanets (Wright and Gaudi 2013) have also been extended to detect exomoons with a varying degree of predicted success rate. The different detection methods are excellently described in Heller’s review in this volume and we will refer the reader to that text for an overview.

Almost twenty years ago there were high expectations on the detection possibilities of exomoons with space-borne facilities like Hubble (Brown et al. 2001), *CoRoT* (Sartoretti and Schneider 1999), or *Kepler* (Szabó et al. 2006); but also with microlensing (Gaudi et al. 2003) or direct imaging (Cabrera and Schneider 2007).


However, there is no uncontroversial detection of an exomoon as we write this lines in August 2017. The situation might change soon, as we will see in the next section. Which have been the difficulties encountered?

Photometric detections during transit and occultation are challenging, even with the latest instrumentation (e.g. see Dobos et al. 2016), and have two main practical limitations: stellar activity and instrumental systematics. A paradigmatic example is the transit of TrEs-1b (Rabus et al. 2009), whose Hubble light curve can be interpreted as a two planet system or as the passage of the transiting planet over an active region on the stellar surface. The same difficulty has been encountered by other systematic studies (Lewis et al. 2015). More challenging is the occurrence of instrumental systematics mimicking the effects of moons, like the case of Kepler-90g (Kipping et al. 2015a). Instrumental systematics can be very difficult to eliminate, even in presence of large amounts of data with high photometric quality (see, for example, Gaidos et al. 2016).

Therefore, the problem is not necessarily the detection but rather the unique interpretation of the measurement as being caused by an exomoon. There is indeed a number of processes that can lead to comparable observational effects but do not involve exomoons. In this respect, the transit timing variation (TTV) method is considered promising, as it can solve part of the degeneracies intrinsic to photometric detections (see Lewis and Fujii 2014, and references therein). However, the transit

timing variations (TTVs) of Kepler-46b (KOI-872b) can be interpreted as an additional planet in the system or as a moon (Nesvorný et al. 2012). TTVs are also strongly affected by stellar activity (Barros et al. 2013; Lewis 2013) and systematics (Szabó et al. 2013). As a result of these limitations, there are presently many systems studied (Weidner and Horne 2010; Kipping et al. 2013b,a, 2014a, 2015b; Hippke 2015; Kane 2017), but no claimed detection of exomoon.

Microlensing surveys have suffered from similar difficulties in the interpretation of the observations with the added challenge of the reproducibility of the measurements (Bennett et al. 2014; Skowron et al. 2014).




**Fig. 1** Comparison between the astrometric measurements obtained for HR8799b by Wertz et al. (2017) and the expected photocenter motion for a Jupiter-Saturn binary planet observed at 39 pc. See Cabrera and Schneider (2007) for details on how has been calculated the expected motion of the photocenter.

Ten years ago, we estimated the possibility of detecting moons around direct imaged planets measuring the reflex motion of the moon around the planet considering photon noise (Cabrera and Schneider 2007). The required precision of the astrometric measurement of the position of the planet was in the range from microarcsec to few milliarcsec. Unfortunately, it is more likely that these observations are actually limited by speckle noise and systematic uncertainties in the astrometric position of the host star rather than photon noise. However, precisions of milliarcseconds are

currently within reach (Wertz et al. 2017), though no moon has been claimed yet (see Fig. 1).

### *Expectations for the future*



**Fig. 2** Planet/moon events. As a satellite orbits its host planet, there are different events that can be observed in the system, depending on the relative configuration of the planet, satellite, and observer.

The photometric detection of moons is theoretically possible with space-borne photometry delivered by missions such as *CoRoT* and *Kepler* or with microlensing surveys, but so far no detection has been secured. Soon the next generation of exoplanet space-borne facilities, including *CHEOPS* (Simon et al. 2015), *TESS* (Ricker et al. 2015), and *PLATO* (Rauer et al. 2014), will expand on the *CoRoT* and *Kepler* legacy. The difficulty in the detection of exomoons will be the same, but these missions will observe brighter stars, easier to characterize, and will benefit from the experience of the previous surveys. For example, one wonders about the follow-up of the TTVs of *Kepler* candidates with *PLATO*, collecting a baseline of observations longer than ten years. *PLATO* has a smaller collecting area than *Kepler*, but a higher cadence, so the TTV accuracy will be close enough to make meaningful compar-

isons. These new missions might be able to definitely settle the nature of some of the candidates proposed today.

An important limitation of transit photometry arises from the fact that a system is in transit only a very small fraction of its orbit (0.15% of the time for the Earth around the Sun). The relative scarcity of the data and the difficulty of reproducing the observations, as the moon changes its relative phase from transit to transit, will not improve for the new missions. A possibility that might still have a chance are binary planets, which have a very distinct transit signature of larger amplitude. Though we know they are not common, they are known to exist in certain configurations (Nielsen et al. 2013; Best et al. 2017; Han et al. 2017).

Despite all the mentioned difficulties, in July 2017 Teachey et al. (2018) announced the presence of a possible exomoon around the planet Kepler-1625b. If confirmed, it would be a sensational discovery culminating the efforts of the HEK (Hunt for Exomoons with Kepler) team (Kipping et al. 2012). However, the authors remain cautious about the nature of the candidate and warn the community about the limited amount of existing observational evidence on this target. There are additional observations scheduled end of October 2017 with the potential to confirm the presence of the candidate. Unfortunately, this text will have to go into publication before the results of these observations are known, but we keep our fingers crossed.

In the near future, another breakthrough might come though from directly imaged planets (see the review by Bowler 2016). As mentioned earlier, current facilities can reach the milliarcsecond precision required to start sampling the existence of massive moons (see Fig. 1). There are important synergies with missions like *Gaia* (Gaia Collaboration et al. 2016), which will probe the parameter space for giant planets beyond the snow-line, potentially observable with current high-contrast adaptive optics facilities. *Gaia* planet yield is expected to outnumber the current sample known (Casertano et al. 2008). The reflex motion of planets, which allows to measure the moon's mass, and the next step, studying planet-moon occultation events (see Schneider et al. 2015), are promising methods to characterize exomoons. The reflex motion is observable during the whole orbit and the planet-moon events occur up to 5 times per moon orbit (see Fig. 2), which is in the order of up to a few tenths of days, giving considerable advantage in comparison to photometric transits. Observing the 5 types of events shown in Fig. 2 is only possible if the relative orientation of orbit of the satellite is favorable to the observer. However, two of the events, when the satellite casts its shadow on the planet and when the satellite is occulted by the shadow of the planet, only depend on the relative orientation of the satellite orbit with the orbital plane of the planet. If the orbital plane of the satellites has a low inclination with respect to the ecliptic, these events will be visible, regardless of the orientation towards the observer.

Though elusive, exomoons are fundamental in the study of the processes of planetary formation and evolution and, furthermore, provide a rich ground for studies constraining the processes of planetary formation and evolution and provide a great opportunity to study habitable systems. Given the new facilities that will become available in the coming years and the expertise accumulated, there are good reasons

to remain optimistic and hope that the next 10 years will be more fruitful than the last two decades.

## Rings

All the giant planets in the solar system are surrounded by systems of rings, though they have very different properties. The processes that affect the stability and evolution of rings involve tidal forces, dynamical interactions with moons, resonances, spiral waves, radiation pressure, and interactions with charged particles (de Pater and Lissauer 2015), making them a very rich field for research.

We bring up in this section rings and disks around planets, but there are ring-structures everywhere in the Universe within an unimaginable range of sizes (see, for example, the review by Latter et al. 2017).

For the detection methods of rings around extrasolar planets, we will refer to the review in this very same volume by R. Heller. Regarding formation mechanisms, see Zanazzi and Lai (2017) and references therein.

### *Current status of detections*

In contrast to exomoons, there are several detections claimed in the literature, though not all of them completely undisputed.

One example is the indirect detection of a ring system around the brown dwarf G 196-3 B (Zakhohay et al. 2017). A ring system with properties resembling those around Jupiter or Neptune, but with a very different age and on different environment, can satisfactorily reproduce the colors of this target.

A hypothetical ring system that can make its *première* in 2017 would be the one around  $\beta$  Pictoris b (Lagrange et al. 2010), a giant planet orbiting a 10 million year old star with an orbital period of about 35 years. The large semi-major axis makes the transit probability meager, but it might actually be transiting (Lecavelier des Etangs and Vidal-Madjar 2016; Wang et al. 2016) and a campaign has been orchestrated to characterize the Hill sphere of the planet as it crosses the stellar disk from Earth (Kenworthy 2017).

And there is the unusual, from our solar system perspective, system of rings proposed around J1407 (1SWASP J140747.93-394542.6 Mamajek et al. 2012; Rieder and Kenworthy 2016), which is described in Heller's chapter in this volume. We will mention it again in the next section.

Despite the wealth of giant planets at large orbital periods, including several Jupiter-analogs, found by different surveys (Bedell et al. 2015; Díaz et al. 2016; Esposito et al. 2013; Foreman-Mackey et al. 2016; Kipping et al. 2014b, 2016; Uehara et al. 2016), space missions like *CoRoT* and *Kepler* have not yielded any report

of exorings so far (see Heising et al. 2015; Lecavelier des Etangs et al. 2017; Turner et al. 2016).

The paper by Aizawa et al. (2017) deserves special attention as it carefully shows that the limitations of the photometric method with current surveys are not dictated by the photon noise, but by the residuals of systematic noise sources and the interpretation of the results, as it was the case for exomoons previously described.

Overcoming the limitations of photometric searches, there are alternative techniques like high resolution spectroscopy (Santos et al. 2015) and direct imaging.

### ***Characterization of rings with direct imaging***

WFIRST-AFTA (Spergel et al. 2015) is an observatory of NASA devoted to the study of dark matter, infrared astrophysics, and extrasolar planets. It is currently in its Phase A, undergoing the study of mission requirements. The mission concept is based on a 2.4 m telescope with a large field-of-view and is equipped with a wide field instrument and a coronagraph. The on-board spectrograph will foreseeably provide a contrast of  $10^{-9}$  and an inner working angle of  $3\lambda/D$  at 430 nm. Such a performance will enable the characterization of the atmospheres of directly imaged exoplanets (Greco and Burrows 2015).

We have used a numerical model based on previous work by Arnold and Schneider (2004) to simulate the integrated light curve of ringed planets that could be observed with WFIRST. The exercise intends to elucidate the effects that exorings would have in the phase curve and spectra of an exoplanet observed via direct imaging, thereby drawing conclusions on the planet atmosphere and the planet size. The planets are assumed at orbital distances of 1-10 AU from their star and 10 pc away from the solar system.

The numerical model considers mutual shadow of the planet on the ring and the ring on the planet. The shadow of the planet on the ring and the occultation are also taken into account. However, mutual reflection and shadow of the ring on the planet are neglected. The code accepts elliptical orbits and rings with fixed inner and outer radius. The planet is assumed to scatter starlight as a Lambertian sphere (Lester et al. 1979), and the rings are assumed to be planar. At the ring, only single scattering is considered.

There are nine parameters that define the planet-ring system geometry of the model: the planetary radius  $R_p$ , the inclination of the orbital plane  $i$ , the wavelength dependent planetary albedo  $A_p$ , the rings optical thickness  $\tau_R$ , the ring's inner and outer radius  $R_{in}$  and  $R_{out}$ , the single scattering albedo of the ring  $\omega_o$ , the ring's plane inclination  $i_R$ , and the ring's plane intersection with the orbital plane  $\lambda_R$  (Arnold and Schneider 2004).

As a reference for the capabilities of WFIRST we have made simulations of a ring system like J1407. To facilitate the study of this system with the planned specifications of WFIRST's inner working angle, we assumed that the system is located



at 10 pc from Earth, rather than the actual 128 pc. The values of the parameters used in the simulation are shown in Table 1 and the results are shown in Figs. 3 and 4.

A system of rings as that proposed for J1407 reflects a significant amount of the stellar light and produces a signal several orders of magnitude larger than the planet itself. This unique case therefore opens the possibility of spectroscopically investigating exorings without the interfering effect of the planet. Less extreme situations such as enabled by Saturn-like exoplanets will show signals that blend the ring and planet contributions. This blending will dilute the main absorption features in the planet atmosphere, thereby complicating its analysis.


**Table 1** Parameters used in the simulation of the ring system.

Element	Parameter	Value
Star	mass	$0.9M_{\text{Sun}}$
	distance	10 pc
Planet	$R_p$	$1.46 R_{\text{Jupiter}}$
	semi-major axis	5 au
	eccentricity	0.65
	$i$	$89^\circ$
Ring	$A_p$	Jupiter
	$\tau_R$	0.5
	$R_{\text{in}}$	$0.25 R_{\text{Hill}}$
	$R_{\text{out}}$	$R_{\text{Hill}}$
	$i_R$	$13^\circ$
	$\lambda_R$	$70^\circ$


The albedo values of Jupiter as a function of the wavelength are taken from Karkoschka (1994).

## Comets

Comets are some of the largest structures in the solar system, if one accounts for the extension of their tails (i.e. Neugebauer et al. 2007). However, the very low density of the extended tails makes their detection and characterization challenging outside the solar system. Nevertheless, this difficulty didn't stop observers from trying to observe the extended, low density, exospheres of extrasolar planets soon after their discovery (Rauer et al. 2000; Vidal-Madjar et al. 2003; Lecavelier Des Etangs et al. 2010; Haswell et al. 2012; Ehrenreich et al. 2012, 2015; Poppenhaeger et al. 2013). The evaporation of giant planets has been followed by the detection of disintegrating small planets which display tails similar to comets (Rappaport et al. 2012, 2014; Sanchis-Ojeda et al. 2015; Vanderburg et al. 2015), which in some cases has allowed the characterization of the properties of the particles in the tail (van Lieshout et al. 2014, 2016; Zhou et al. 2016; Alonso et al. 2016; Rappaport et al. 2016). Of interest



**Fig. 3** Contrast vs. inner working angle for a J1407-like system compared to the WFIRST detection limits for a planet without rings (dark blue) and with rings (orange). The detectability improves moving from left to right and from bottom to top. The red horizontal line represents the planned  $10^{-9}$  contrast limit of WFIRST. Correspondingly, the red vertical line represents a tentative limit of the inner working angle possibilities.



**Fig. 4** Phase curves for a J1407-like system. Left: for a system without rings. Right: for a system with rings, note the change in the vertical scale. In the latter, the shape of the phase curve is dominated by the size of the rings and their relative orientation to the observer.

are also the transient signatures recently discovered around the young star RIK-210, though their interpretation is not so straightforward (David et al. 2017).

The discovery of comets with photometric transit surveys has been more difficult than what was originally expected (Lecavelier Des Etangs et al. 1999) for possibly the same reasons that have been already described above. There have been clear detections in the circumstellar disk of young stars (Kiefer et al. 2014b,a; Eiroa et al. 2016; Marino et al. 2017). But the first evidence for exocomets transiting in front of a star in visible light had to wait until August 2017. It comes from the discovery of possibly several comets around the star KIC 3542116 (Rappaport et al. 2018). This pioneering paper further analyses the possible properties of the comets based on the shape of the observed light curves, which are book examples of the expectations in (Lecavelier Des Etangs et al. 1999). The authors of the paper are optimistic that new examples will be found in future analyses of the *Kepler* data.

One disputed case is KIC 8462852 (Boyajian et al. 2016), a target observed by the *Kepler* mission that shows irregular flux drops that account for up to 20% of the stellar flux lasting several days. Some teams have invoked the possibility of comets (Bodman and Quillen 2016) to explain the observations, but the hypothesis has been recently challenged (Wright and Sigurdsson 2016). However, new observations in May 2017 (triggered by a *tweet* by T. Boyajian, @tsboyajian) suggest that the phenomenon has a characteristic time-scale of 700 days and its origin, whatever its nature, is indeed gravitationally linked to the star. It has been proposed that a ringed planet and a large swarm of trojan bodies could explain the features observed in the light curve and predict its future behaviour (Ballesteros et al. 2018). The semi-major axis of the system would be around 6 au and the orbital period 12 years, a large observational span, but certainly within reach in the near future.

## Trojans

In the solar system the term Trojans refer to a family of minor bodies that share the orbits of the giant planets like Jupiter and Neptune. They represent a special case of the co-orbital dynamics of the N-body problem (see, for example, Veras et al. 2016).

The stability of such configurations is not simple, nor their dynamical evolution as planets migrate (Nesvorný et al. 2013), but there is no known reason preventing their existence in extrasolar systems. Only that, if they have similar sizes to those in the solar system, their direct detectability with photometry is beyond reach for current and near future facilities.

Therefore, researchers have tried to infer the presence of trojans studying the perturbations introduced in the orbit of their larger, companion planet, both with radial velocity, transit timing variations, or both (Ford and Holman 2007; Dobrovolskis 2013; Haghighipour et al. 2013; Leleu et al. 2015, 2017; Nesvorný and Vokrouhlický 2016; Vokrouhlický and Nesvorný 2014). There have been several systematic searches that have not found any reliable candidate so far (Madhusudhan and Winn 2009; Janson 2013).

There have been claims in the literature with detections, but so far none of these claims has been confirmed by subsequent independent analysis, like the cases of HD 82943 and HD 128311 (Goździewski and Konacki 2006), but see McArthur et al. (2014) and Rein (2015), or the Kepler candidates by Hippke and Angerhausen (2015), which at least in the case of Kepler-91 b have not been confirmed by later studies (Placek et al. 2015). A recent claim on WASP-12 and HD 18733 by Kislyakova et al. (2016) remains to be confirmed.

In this case, the direct confirmation of the presence of a trojan suffers from the same limitations of reproducibility, credibility in presence of correlated noise, and degeneracy of interpretation as in the previous examples of moons and rings, with the detriment of the smaller size and mass of the researched object.

## Summary

We would like to close with a quote attributed to Napoleon Bonaparte dissuading from taking predictions too seriously (*Il faut toujours se réserver le droit de rire le lendemain de ses idées de la veille*). Or in the words of a famous astronomer, *you should listen to theorists but never take them too seriously*. The discovery and characterization of moons, rings, comets, and trojans has proven more challenging than expected, but the interest of the community has not decreased in the last 20 years. It is rather the opposite. And there are also new ideas coming out, like *synestias* (Lock and Stewart 2017). These are transient structures predicted by theoretical models produced during planetary formation processes. They have not been observed, or confirmed independently yet, but they are certainly welcome because of their interest.

The wealth of data from transit photometry has been carefully studied and most of the systematics are well understood, yet no undisputed detection has been definitely accepted by the community. However, the situation is quickly changing in a very positive way. TESS and PLATO will have their chance in the next decade, but it is time to think about different methods, in particular direct imaging and probably high resolution spectroscopy (for example, the serendipitous discovery of a moon or ring system via Rossiter-McLaughlin during planet characterization). With instruments like Gaia, ALMA, and E-ELT class telescopes it is difficult not to end with the conclusion that moons, rings, comets, and trojans will not only be detected in large numbers in the next decades, but also will contribute to our knowledge about planets and planetary systems, in our galaxy and in the solar system.

## Cross-References

- On the Detection of Extrasolar Moons and Rings

## References

- Adams FC Bloch AM (2016) The stability of tidal equilibrium for hierarchical star-planet-moon systems. *MNRAS*462:2527–2541
- Aizawa M, Uehara S, Masuda K, Kawahara H Suto Y (2017) Toward Detection of Exoplanetary Rings via Transit Photometry: Methodology and a Possible Candidate. *AJ*153:193
- Alonso R, Rappaport S, Deeg HJ Palles E (2016) Gray transits of WD 1145+017 over the visible band. *A&A*589:L6
- Arnold L Schneider J (2004) The detectability of extrasolar planet surroundings. I. Reflected-light photometry of unresolved rings. *A&A*420:1153–1162
- Ballesteros FJ, Arnalte-Mur P, Fernandez-Soto A Martínez VJ (2018) KIC 8462852: Will the Trojans return in 2021? *MNRAS*473:L21–L25
- Barnes JW O'Brien DP (2002) Stability of Satellites around Close-in Extrasolar Giant Planets. *ApJ*575:1087–1093
- Barr AC (2016) Formation of exomoons: a solar system perspective. *Astronomical Review* 12(1-4):24–52
- Barr AC Bruck Syal M (2017) Formation of massive rocky exomoons by giant impact. *MNRAS*466:4868–4874
- Barros SCC, Boué G, Gibson NP et al. (2013) Transit timing variations in WASP-10b induced by stellar activity. *MNRAS*430:3032–3047
- Bedell M, Meléndez J, Bean JL et al. (2015) The Solar Twin Planet Search. II. A Jupiter twin around a solar twin. *A&A*581:A34
- Bennett DP, Batista V, Bond IA et al. (2014) MOA-2011-BLG-262Lb: A Sub-Earth-Mass Moon Orbiting a Gas Giant Primary or a High Velocity Planetary System in the Galactic Bulge. *ApJ*785:155
- Best WMJ, Liu MC, Dupuy TJ Magnier EA (2017) The Young L Dwarf 2MASS J11193254-1137466 Is a Planetary-mass Binary. *ApJ*843:L4
- Bodman EHL Quillen A (2016) KIC 8462852: Transit of a Large Comet Family. *ApJ*819:L34
- Bowler BP (2016) Imaging Extrasolar Giant Planets. *PASP*128(10):102,001
- Boyajian TS, LaCourse DM, Rappaport SA et al. (2016) Planet Hunters IX. KIC 8462852 - where's the flux? *MNRAS*457:3988–4004
- Brown TM, Charbonneau D, Gilliland RL, Noyes RW Burrows A (2001) Hubble Space Telescope Time-Series Photometry of the Transiting Planet of HD 209458. *ApJ*552:699–709
- Cabrera J Schneider J (2007) Detecting companions to extrasolar planets using mutual events. *A&A*464:1133–1138
- Casertano S, Lattanzi MG, Sozzetti A et al. (2008) Double-blind test program for astrometric planet detection with Gaia. *A&A*482:699–729
- Cassidy TA, Mendez R, Arras P, Johnson RE Skrutskie MF (2009) Massive Satellites of Close-In Gas Giant Exoplanets. *ApJ*704:1341–1348
- Cockell CS, Bush T, Bryce C et al. (2016) Habitability: A Review. *Astrobiology* 16:89–117
- Crida A Charnoz S (2012) Formation of Regular Satellites from Ancient Massive Rings in the Solar System. *Science* 338:1196–
- David TJ, Petigura EA, Hillenbrand LA et al. (2017) A Transient Transit Signature Associated with the Young Star RIK-210. *ApJ*835:168
- de Pater I Lissauer JJ (2015) Planetary Sciences
- Debes JH Sigurdsson S (2007) The Survival Rate of Ejected Terrestrial Planets with Moons. *ApJ*668:L167–L170
- Díaz RF, Rey J, Demangeon O et al. (2016) The SOPHIE search for northern extrasolar planets. XI. Three new companions and an orbit update: Giant planets in the habitable zone. *A&A*591:A146
- Dobos V, Kereszturi Á, Pál A Kiss LL (2016) Possibility for albedo estimation of exomoons: Why should we care about M dwarfs? *A&A*592:A139
- Dobos V, Heller R Turner EL (2017) The effect of multiple heat sources on exomoon habitable zones. *A&A*601:A91

- Dobrovolskis AR (2013) Effects of Trojan exoplanets on the reflex motions of their parent stars. *Icarus* 226:1635–1641
- Domingos RC, Winter OC Yokoyama T (2006) Stable satellites around extrasolar giant planets. *MNRAS* 373:1227–1234
- Donnison JR (2010) The Hill stability of the possible moons of extrasolar planets. *MNRAS* 406:1918–1934
- Ehrenreich D, Bourrier V, Bonfils X et al. (2012) Hint of a transiting extended atmosphere on 55 Cancri b. *A&A* 547:A18
- Ehrenreich D, Bourrier V, Wheatley PJ et al. (2015) A giant comet-like cloud of hydrogen escaping the warm Neptune-mass exoplanet GJ 436b. *Nature* 522:459–461
- Eiroa C, Rebollido I, Montesinos B et al. (2016) Exocomet signatures around the A-shell star  $\phi$  Leonis? *A&A* 594:L1
- Esposito S, Mesa D, Skemer A et al. (2013) LBT observations of the HR 8799 planetary system. First detection of HR 8799e in H band. *A&A* 549:A52
- Ford EB Holman MJ (2007) Using Transit Timing Observations to Search for Trojans of Transiting Extrasolar Planets. *ApJ* 664:L51–L54
- Foreman-Mackey D, Morton TD, Hogg DW, Agol E Schölkopf B (2016) The Population of Long-period Transiting Exoplanets. *AJ* 152:206
- Forgan D Kipping D (2013) Dynamical effects on the habitable zone for Earth-like exomoons. *MNRAS* 432:2994–3004
- Forgan D Yotov V (2014) The effect of planetary illumination on climate modelling of Earth-like exomoons. *MNRAS* 441:3513–3523
- Fujii YI, Kobayashi H, Takahashi SZ Gressel O (2017) Orbital Evolution of Moons in Weakly Accreting Circumplanetary Disks. *AJ* 153:194
- Gaia Collaboration, Brown AGA, Vallenari A et al. (2016) Gaia Data Release 1. Summary of the astrometric, photometric, and survey properties. *A&A* 595:A2
- Gaidos E, Mann AW Ansdell M (2016) The Enigmatic and Ephemeral M Dwarf System KOI 6705: Cheshire Cat or Wild Goose? *ApJ* 817:50
- Gaudi BS, Chang HY Han C (2003) Probing Structures of Distant Extrasolar Planets with Microlensing. *ApJ* 586:527–539
- Gong YX, Zhou JL, Xie JW Wu XM (2013) The Effect of Planet-Planet Scattering on the Survival of Exomoons. *ApJ* 769:L14
- Goździewski K Konacki M (2006) Trojan Pairs in the HD 128311 and HD 82943 Planetary Systems? *ApJ* 647:573–586
- Greco JP Burrows A (2015) The Direct Detectability of Giant Exoplanets in the Optical. *ApJ* 808:172
- Haghighipour N, Capen S Hinse TC (2013) Detection of Earth-mass and super-Earth Trojan planets using transit timing variation method. *Celestial Mechanics and Dynamical Astronomy* 117:75–89
- Han C, Udalski A, Sumi T et al. (2017) OGLE-2016-BLG-1469L: Microlensing Binary Composed of Brown Dwarfs. *ApJ* 843:59
- Haswell CA, Fossati L, Ayres T et al. (2012) Near-ultraviolet Absorption, Chromospheric Activity, and Star-Planet Interactions in the WASP-12 system. *ApJ* 760:79
- Heising MZ, Marcy GW Schlichting HE (2015) A Search for Ringed Exoplanets Using Kepler Photometry. *ApJ* 814:81
- Heller R (2012) Exomoon habitability constrained by energy flux and orbital stability. *A&A* 545:L8
- Heller R Barnes R (2015) Runaway greenhouse effect on exomoons due to irradiation from hot, young giant planets. *International Journal of Astrobiology* 14:335–343
- Heller R Pudritz R (2015) Conditions for water ice lines and Mars-mass exomoons around accreting super-Jovian planets at 1–20 AU from Sun-like stars. *A&A* 578:A19
- Hippke M (2015) On the Detection of Exomoons: A Search in Kepler Data for the Orbital Sampling Effect and the Scatter Peak. *ApJ* 806:51
- Hippke M Angerhausen D (2015) A Statistical Search for a Population of Exo-Trojans in the Kepler Data Set. *ApJ* 811:1

- Hong YC, Tiscareno MS, Nicholson PD, Lunine JI (2015) Orbital instability of close-in exomoons in non-coplanar systems. *MNRAS*449:828–834
- Janson M (2013) A Systematic Search for Trojan Planets in the Kepler Data. *ApJ*774:156
- Kaltenegger L (2010) Characterizing Habitable Exomoons. *ApJ*712:L125–L130
- Kane SR (2017) Worlds without Moons: Exomoon Constraints for Compact Planetary Systems. *ApJ*839:L19
- Karkoschka E (1994) Spectrophotometry of the jovian planets and Titan at 300- to 1000-nm wavelength: The methane spectrum. *Icarus*111:174–192
- Kenworthy M (2017) Looking for rings and things. *Nature Astronomy* 1:0099
- Kiefer F, Lecavelier des Etangs A, Augereau JC et al. (2014a) Exocomets in the circumstellar gas disk of HD 172555. *A&A*561:L10
- Kiefer F, Lecavelier des Etangs A, Boissier J et al. (2014b) Two families of exocomets in the  $\beta$  Pictoris system. *Nature*514:462–464
- Kipping DM, Bakos GÁ, Buchhave L, Nesvorný D, Schmitt A (2012) The Hunt for Exomoons with Kepler (HEK). I. Description of a New Observational project. *ApJ*750:115
- Kipping DM, Forgan D, Hartman J et al. (2013a) The Hunt for Exomoons with Kepler (HEK). III. The First Search for an Exomoon around a Habitable-zone Planet. *ApJ*777:134
- Kipping DM, Hartman J, Buchhave LA et al. (2013b) The Hunt for Exomoons with Kepler (HEK). II. Analysis of Seven Viable Satellite-hosting Planet Candidates. *ApJ*770:101
- Kipping DM, Nesvorný D, Buchhave LA et al. (2014a) The Hunt for Exomoons with Kepler (HEK). IV. A Search for Moons around Eight M Dwarfs. *ApJ*784:28
- Kipping DM, Torres G, Buchhave LA et al. (2014b) Discovery of a Transiting Planet near the Snow-line. *ApJ*795:25
- Kipping DM, Huang X, Nesvorný D et al. (2015a) The Possible Moon of Kepler-90g is a False Positive. *ApJ*799:L14
- Kipping DM, Schmitt AR, Huang X et al. (2015b) The Hunt for Exomoons with Kepler (HEK): V. A Survey of 41 Planetary Candidates for Exomoons. *ApJ*813:14
- Kipping DM, Torres G, Henze C et al. (2016) A Transiting Jupiter Analog. *ApJ*820:112
- Kislyakova KG, Pilat-Lohinger E, Funk B et al. (2016) On the ultraviolet anomalies of the WASP-12 and HD 189733 systems: Trojan satellites as a plasma source. *MNRAS*461:988–999
- Lagrange AM, Bonnefoy M, Chauvin G et al. (2010) A Giant Planet Imaged in the Disk of the Young Star  $\beta$  Pictoris. *Science* 329:57
- Lammer H, Schiefer SC, Juvan I et al. (2014) Origin and Stability of Exomoon Atmospheres: Implications for Habitability. *Origins of Life and Evolution of the Biosphere* 44:239–260
- Latter HN, Ogilvie GI, Rein H (2017) Planetary rings and other astrophysical disks. In: *Planetary Ring Systems*
- Laughlin G, Adams FC (2000) The Frozen Earth: Binary Scattering Events and the Fate of the Solar System. *Icarus*145:614–627
- Lecavelier des Etangs A, Vidal-Madjar A (2016) The orbit of beta Pictoris b as a transiting planet. *A&A*588:A60
- Lecavelier Des Etangs A, Vidal-Madjar A, Ferlet R (1999) Photometric stellar variation due to extra-solar comets. *A&A*343:916–922
- Lecavelier Des Etangs A, Ehrenreich D, Vidal-Madjar A et al. (2010) Evaporation of the planet HD 189733b observed in H I Lyman- $\alpha$ . *A&A*514:A72
- Lecavelier des Etangs A, Hébrard G, Blandin S et al. (2017) Search for rings and satellites around the exoplanet CoRoT-9b using Spitzer photometry. *A&A*603:A115
- Leleu A, Robutel P, Correia ACM (2015) Detectability of quasi-circular co-orbital planets. Application to the radial velocity technique. *A&A*581:A128
- Leleu A, Robutel P, Correia ACM, Lillo-Box J (2017) Detection of co-orbital planets by combining transit and radial-velocity measurements. *A&A*599:L7
- Lester TP, McCall ML, Tatum JB (1979) Theory of planetary photometry. *JRASC*73:233–257
- Lewis KM (2013) Detecting exo-moons with photometric transit timing - I. Effect of realistic stellar noise. *MNRAS*430:1473–1485

- Lewis KM Fujii Y (2014) Next Generation of Telescopes or Dynamics Required to Determine if Exo-moons have Prograde or Retrograde Orbits. *ApJ*791:L26
- Lewis KM, Ochiai H, Nagasawa M Ida S (2015) Extrasolar Binary Planets II: Detectability by Transit Observations. *ApJ*805:27
- Li T, Tian F, Wang Y, Wei W Huang X (2016) Distinguishing a Hypothetical Abiotic Planet-Moon System from a Single Inhabited Planet. *ApJ*817:L15
- Lock SJ Stewart ST (2017) The structure of terrestrial bodies: Impact heating, corotation limits, and synestias. *Journal of Geophysical Research (Planets)* 122:950–982
- Madhusudhan N Winn JN (2009) Empirical Constraints on Trojan Companions and Orbital Eccentricities in 25 Transiting Exoplanetary Systems. *ApJ*693:784–793
- Mamajek EE, Quillen AC, Pecaui MJ et al. (2012) Planetary Construction Zones in Occultation: Discovery of an Extrasolar Ring System Transiting a Young Sun-like Star and Future Prospects for Detecting Eclipses by Circumsecondary and Circumplanetary Disks. *AJ*143:72
- Marino S, Wyatt MC, Panić O et al. (2017) ALMA observations of the  $\eta$  Corvi debris disc: inward scattering of CO-rich exocomets by a chain of 3–30  $M_{\oplus}$  planets? *MNRAS*465:2595–2615
- McArthur BE, Benedict GF, Henry GW et al. (2014) Astrometry, Radial Velocity, and Photometry: The HD 128311 System Remixed with Data from HST, HET, and APT. *ApJ*795:41
- Miguel Y Ida S (2016) A semi-analytical model for exploring Galilean satellites formation from a massive disk. *Icarus*266:1–14
- Murray CD Dermott SF (2000) *Solar System Dynamics*
- Namouni F (2010) The Fate of Moons of Close-in Giant Exoplanets. *ApJ*719:L145–L147
- Nesvorný D Vokrouhlický D (2016) Dynamics and Transit Variations of Resonant Exoplanets. *ApJ*823:72
- Nesvorný D, Kipping DM, Buchhave LA et al. (2012) The Detection and Characterization of a Nontransiting Planet by Transit Timing Variations. *Science* 336:1133–
- Nesvorný D, Vokrouhlický D Morbidelli A (2013) Capture of Trojans by Jumping Jupiter. *ApJ*768:45
- Neugebauer M, Gloeckler G, Gosling JT et al. (2007) Encounter of the Ulysses Spacecraft with the Ion Tail of Comet MCNaught. *ApJ*667:1262–1266
- Nielsen EL, Liu MC, Wahhaj Z et al. (2013) The Gemini NICI Planet-Finding Campaign: The Frequency of Giant Planets around Young B and A Stars. *ApJ*776:4
- Ochiai H, Nagasawa M Ida S (2014) Extrasolar Binary Planets. I. Formation by Tidal Capture during Planet-Planet Scattering. *ApJ*790:92
- Ogihara M Ida S (2012) N-body Simulations of Satellite Formation around Giant Planets: Origin of Orbital Configuration of the Galilean Moons. *ApJ*753:60
- Payne MJ, Deck KM, Holman MJ Perets HB (2013) Stability of Satellites in Closely Packed Planetary Systems. *ApJ*775:L44
- Placek B, Knuth KH, Angerhausen D Jenkins JM (2015) Characterization of Kepler-91b and the Investigation of a Potential Trojan Companion Using EXONEST. *ApJ*814:147
- Poppenhaeger K, Schmitt JHMM Wolk SJ (2013) Transit Observations of the Hot Jupiter HD 189733b at X-Ray Wavelengths. *ApJ*773:62
- Rabus M, Alonso R, Belmonte JA et al. (2009) A cool starspot or a second transiting planet in the TrES-1 system? *A&A*494:391–397
- Rappaport S, Levine A, Chiang E et al. (2012) Possible Disintegrating Short-period Super-Mercury Orbiting KIC 12557548. *ApJ*752:1
- Rappaport S, Barclay T, DeVore J et al. (2014) KOI-2700b A Planet Candidate with Dusty Effluents on a 22 hr Orbit. *ApJ*784:40
- Rappaport S, Gary BL, Kaye T et al. (2016) Drifting asteroid fragments around WD 1145+017. *MNRAS*458:3904–3917
- Rappaport S, Vanderburg A, Jacobs T et al. (2018) Likely transiting exocomets detected by Kepler. *MNRAS*474:1453–1468
- Rauer H, Bockelée-Morvan D, Coustenis A, Guillot T Schneider J (2000) Search for an exosphere around 51 Pegasi B with ISO. *A&A*355:573–580



- Rauer H, Catala C, Aerts C et al. (2014) The PLATO 2.0 mission. *Experimental Astronomy* 38:249–330
- Rein H (2015) Reanalysis of radial velocity data from the resonant planetary system HD128311. *MNRAS* 448:L58–L61
- Rein H, Fujii Y, Spiegel DS (2014) Some inconvenient truths about biosignatures involving two chemical species on Earth-like exoplanets. *Proceedings of the National Academy of Science* 111:6871–6875
- Ricker GR, Winn JN, Vanderspek R et al. (2015) Transiting Exoplanet Survey Satellite (TESS). *Journal of Astronomical Telescopes, Instruments, and Systems* 1(1):014003
- Rieder S, Kenworthy MA (2016) Constraints on the size and dynamics of the J1407b ring system. *A&A* 596:A9
- Sanchis-Ojeda R, Rappaport S, Pallè E et al. (2015) The K2-ESPRINT Project I: Discovery of the Disintegrating Rocky Planet K2-22b with a Cometary Head and Leading Tail. *ApJ* 812:112
- Santos NC, Martins JHC, Boué G et al. (2015) Detecting ring systems around exoplanets using high resolution spectroscopy: the case of 51 Pegasi b. *A&A* 583:A50
- Sartoretti P, Schneider J (1999) On the detection of satellites of extrasolar planets with the method of transits. *A&AS* 134:553–560
- Sasaki T, Barnes JW (2014) Longevity of moons around habitable planets. *International Journal of Astrobiology* 13:324–336
- Sasaki T, Barnes JW, O’Brien DP (2012) Outcomes and Duration of Tidal Evolution in a Star-Planet-Moon System. *ApJ* 754:51
- Scharf CA (2006) The Potential for Tidally Heated Icy and Temperate Moons around Exoplanets. *ApJ* 648:1196–1205
- Schneider J, Léger A, Fridlund M et al. (2010) The Far Future of Exoplanet Direct Characterization. *Astrobiology* 10:121–126
- Schneider J, Lainey V, Cabrera J (2015) A next step in exoplanetology: exo-moons. *International Journal of Astrobiology* 14:191–199
- Simon AE, Szabó GM, Kiss LL, Fortier A, Benz W (2015) CHEOPS Performance for Exomoons: The Detectability of Exomoons by Using Optimal Decision Algorithm. *PASP* 127:1084
- Sinukoff E, Fulton B, Scuderi L, Gaidos E (2013) Below One Earth: The Detection, Formation, and Properties of Subterrestrial Worlds. *Space Sci Rev* 180:71–99
- Skowron J, Udalski A, Szymański MK et al. (2014) New Method to Measure Proper Motions of Microlensed Sources: Application to Candidate Free-floating-planet Event MOA-2011-BLG-262. *ApJ* 785:156
- Spalding C, Batygin K, Adams FC (2016) Resonant Removal of Exomoons during Planetary Migration. *ApJ* 817:18
- Spergel D, Gehrels N, Baltay C et al. (2015) Wide-Field Infrared Survey Telescope-Astrophysics Focused Telescope Assets WFIRST-AFTA 2015 Report. *ArXiv e-prints*
- Szabó GM, Szatmáry K, Divéki Z, Simon A (2006) Possibility of a photometric detection of “exo-moons”. *A&A* 450:395–398
- Szabó R, Szabó GM, Dálya G et al. (2013) Multiple planets or exomoons in Kepler hot Jupiter systems with transit timing variations? *A&A* 553:A17
- Teachey A, Kipping DM, Schmitt AR (2018) HEK. VI. On the Dearth of Galilean Analogs in Kepler, and the Exomoon Candidate Kepler-1625b I. *AJ* 155:36
- Turner JD, Pearson KA, Biddle LI et al. (2016) Ground-based near-UV observations of 15 transiting exoplanets: constraints on their atmospheres and no evidence for asymmetrical transits. *MNRAS* 459:789–819
- Uehara S, Kawahara H, Masuda K, Yamada S, Aizawa M (2016) Transiting Planet Candidates Beyond the Snow Line Detected by Visual Inspection of 7557 Kepler Objects of Interest. *ApJ* 822:2
- van Lieshout R, Min M, Dominik C (2014) Dusty tails of evaporating exoplanets. I. Constraints on the dust composition. *A&A* 572:A76
- van Lieshout R, Min M, Dominik C et al. (2016) Dusty tails of evaporating exoplanets. II. Physical modelling of the KIC 12557548b light curve. *A&A* 596:A32

- Vanderburg A, Johnson JA, Rappaport S et al. (2015) A disintegrating minor planet transiting a white dwarf. *Nature* 526:546–549
- Veras D, Marsh TR Gänsicke BT (2016) Dynamical mass and multiplicity constraints on co-orbital bodies around stars. *MNRAS* 461:1413–1420
- Vidal-Madjar A, Lecavelier des Etangs A, Désert JM et al. (2003) An extended upper atmosphere around the extrasolar planet HD209458b. *Nature* 422:143–146
- Vokrouhlický D Nesvorný D (2014) Transit Timing Variations for Planets Co-orbiting in the Horse-shoe Regime. *ApJ* 791:6
- Waite JH, Glein CR, Perryman RS et al. (2017) Cassini finds molecular hydrogen in the enceladus plume: Evidence for hydrothermal processes. *Science* 356(6334):155–159, URL <http://science.sciencemag.org/content/356/6334/155>
- Wang JJ, Graham JR, Pueyo L et al. (2016) The Orbit and Transit Prospects for  $\beta$  Pictoris b Constrained with One Milliarcsecond Astrometry. *AJ* 152:97
- Weidner C Horne K (2010) Limits on the orbits and masses of moons around currently-known transiting exoplanets. *A&A* 521:A76
- Wertz O, Absil O, Gómez González CA et al. (2017) VLT/SPHERE robust astrometry of the HR8799 planets at milliarcsecond-level accuracy. Orbital architecture analysis with PyAstroFit. *A&A* 598:A83
- Wright JT Gaudi BS (2013) Exoplanet detection methods. In: Oswalt T, French LM Kalas P (eds) *Planets, Stars and Stellar Systems*, vol 3, pp 489–540
- Wright JT Sigurdsson S (2016) Families of Plausible Solutions to the Puzzle of Boyajians Star. *ApJ* 829:L3
- Yang M, Xie JW, Zhou JL, Liu HG Zhang H (2016) Global Instability of the Exo-moon System Triggered by Photo-evaporation. *ApJ* 833:7
- Zakhohay OV, Zapatero Osorio MR, Béjar VJS Boehler Y (2017) Spectral energy distribution simulations of a possible ring structure around the young, red brown dwarf G 196-3 B. *MNRAS* 464:1108–1118
- Zanazzi JJ Lai D (2017) Extended transiting discs and rings around planets and brown dwarfs: theoretical constraints. *MNRAS* 464:3945–3954
- Zhou G, Kedziora-Chudczer L, Bailey J et al. (2016) Simultaneous infrared and optical observations of the transiting debris cloud around WD 1145+017. *MNRAS* 463:4422–4432