

On the expected runtime of multiple testing algorithms with bounded error

Georg Hahn

Abstract

Consider testing multiple hypotheses in the setting where the p-values of all hypotheses are unknown and thus have to be approximated using Monte Carlo simulations. One class of algorithms published in the literature for this scenario provides guarantees on the correctness of their testing result through the computation of confidence statements on all approximated p-values. This article focuses on the expected runtime of such algorithms and derives a variety of finite and infinite expected runtime results.

Keywords: algorithm; bounded error; computational effort; finite expected runtime; multiple testing.

1 Introduction

Consider the testing of $m \in \mathbb{N}$ hypotheses H_{01}, \dots, H_{0m} in a scenario in which the p-values p_1, \dots, p_m corresponding to the m hypotheses are unknown and thus have to be approximated using Monte Carlo simulations, for instance through bootstrap or permutation tests. Several algorithms published in the literature are designed for this scenario, either using a truncation rule to reach fast decisions (Besag and Clifford, 1991; Davidson and MacKinnon, 2000; Andrews and Buchinsky, 2000, 2003; van Wieringen et al., 2008; Sandve et al., 2011; Silva and Assunção, 2013, 2018) or using a heuristic approach to minimize the computational effort without truncation (Lin, 2005; Silva et al., 2009; Gandy and Hahn, 2017).

In the aforementioned Monte Carlo scenario, the main focus of this article lies on the expected runtime of methods which aim to provide a guarantee of correctness on their decision (rejection or non-rejection) for each hypothesis through the computation of a sequence of confidence intervals on each p-value. For the special case of a single hypothesis, it is known that algorithms which compute (or rely on) the decision of one hypothesis with respect to a fixed threshold have an infinite expected runtime. This result is restated below in Section 1.1.

The novelty of this article consists in the generalisation of expected runtime results from single to multiple hypotheses. To be precise, under a simple and weak asymptotic condition on the length of the intervals produced by the confidence sequence, the article shows the following three main results. For applications relying on independent testing, meaning decisions of multiple hypotheses tested at a constant (Bonferroni-type) threshold, all but two hypotheses can be decided in finite expected runtime (Section 2). This result does not extend to applications which require full knowledge of all individual decisions, for instance step-up or step-down procedures, in which case no algorithm can guarantee even a single decision in finite expected runtime (Section 3). Simulations included in the conclusions (Section 4) show that in practice, however, the number of pending decisions is typically low.

Although unconsidered in their original publications, the expected runtimes derived in this article apply to a whole class of published algorithms in the literature. For instance, they apply to the algorithms of Guo and Peddada (2008); Gandy and Hahn (2014) which provide a guarantee on the correctness of the decision on each hypothesis through the computation of Clopper and Pearson (1934) confidence intervals. Expected runtimes also hold true for the

confidence sequences of Robbins (1970); Lai (1976) employed in the methods of Gandy and Hahn (2016); Ding et al. (2018), as they do for the confidence sequences of Darling and Robbins (1967a,b) and the binomial confidence intervals of Armitage (1958) employed in Fay et al. (2007); Gandy (2009). The results of this article do not apply to the *push-out design* of Fay and Follmann (2002) and the *B-value design* of Kim (2010), which both achieve a bounded resampling risk without confidence statements on the p-values.

1.1 A single decision requires infinite expected runtime

Following an argument similar to (Gandy, 2009, Section 3.1), computing the decision of a hypothesis with random p-value requires an infinite expected runtime. Let $m = 1$ hypothesis H_{01} be given having a random p-value p_1 . Assume a sequential algorithm A tests H_{01} at some given threshold α by approximating p_1 through the drawing of Monte Carlo simulations and gives a guarantee of $1 - \epsilon$ on the correctness of its decision, where $\epsilon \in (0, 1)$.

Computing a decision on H_{01} is equivalent to deciding whether $p_1 \leq \alpha$ or $p_1 > \alpha$. For some $p_0 > \alpha$, consider testing $H_0 : p_1 \leq \alpha$ against $H_1 : p_1 = p_0$. A test can be constructed by rejecting H_0 if and only if A does not reject H_{01} . Due to the guarantee of algorithm A , both the type 1 and type 2 errors of this test are ϵ . For such a sequential test, a lower bound on the runtime τ (the expected number of steps) is given by (Wald, 1945, equation (4.81)) as

$$\mathbb{E}(\tau|p = p_1) \geq \frac{\epsilon \log\left(\frac{\epsilon}{1-\epsilon}\right) + (1 - \epsilon) \log\left(\frac{1-\epsilon}{\epsilon}\right)}{p_1 \log\left(\frac{p_1}{\alpha}\right) + (1 - p_1) \log\left(\frac{1-p_1}{1-\alpha}\right)}. \quad (1)$$

The same bound (1) can be derived for the case $p_0 \leq \alpha$. Abbreviate the numerator of (1) by C and consider p_1 in a Bayesian setup such that the following condition is satisfied.

Condition 1. Assume p_1 has a distribution function $F(p_1)$ with derivative $F'(\alpha) > 0$, and that for a suitable $\gamma > 0$, there exists a constant $d > 0$ such that $F'(p_1) \geq d$ in $(\alpha, \alpha + \gamma)$.

Amongst others, Condition 1 is satisfied for the distributions of the exponential family. Under Condition 1, $\mathbb{E}(\tau)$ can be bounded by

$$\begin{aligned} \mathbb{E}(\tau) &= \int_0^1 \mathbb{E}(\tau|p = p_1) dF(p_1) \geq \int_\alpha^{\alpha+\gamma} \mathbb{E}(\tau|p = p_1) dF(p_1) \\ &\geq C \cdot d \cdot \int_\alpha^{\alpha+\gamma} \left(p \log\left(\frac{p}{\alpha}\right) + (1 - p) \log\left(\frac{1-p}{1-\alpha}\right) \right)^{-1} dp = \infty, \end{aligned}$$

as the integrand is proportional to $(p - \alpha)^{-2}$ as $p \rightarrow \alpha$. This proves an infinite expected runtime for the sequential test of H_0 and, equivalently, for algorithm A .

2 Bonferroni-type multiple testing in expected finite time

Assume the testing of H_{01}, \dots, H_{0m} is carried out by comparing each p_i to a threshold value $\alpha_i \in (0, 1)$, $i \in \{1, \dots, m\}$, as done in, for instance, step-up and step-down procedures (Gandy and Hahn, 2016). For this assume $\alpha_1 < \dots < \alpha_m$. In applications which rely on multiple testing at a constant (Bonferroni-type) threshold with a guarantee of correctness through confidence statements on the p-values, it will be shown that decisions on all but two hypotheses can be computed in expected finite time. The guarantee of correctness on all decisions is assumed to hold simultaneously for all hypotheses at $1 - \epsilon$ for some pre-specified $\epsilon \in (0, 1)$.

To be precise, a stronger statement is proven. For all but two hypotheses, it can be decided in expected finite time which of the intervals

$$\mathcal{I} = \{[\alpha_i, \alpha_{i+1}) : \alpha_i < \alpha_{i+1}, i = 1, \dots, m - 1\} \cup \{[0, \alpha_1), [\alpha_m, 1]\} \quad (2)$$

their p-values fall into.

As p-values are unknown, they are approximated through Monte Carlo simulations, and confidence statements are provided via confidence intervals. Let $g(n)$ be the length of the confidence interval for a p-value after drawing n Monte Carlo simulations, and \hat{p}_n be the maximum likelihood estimate of p based on n Monte Carlo simulations. Alternatively, any other estimate \hat{p}_n is permissible whose deviation from the mean can be bounded with a Hoeffding type inequality (Hoeffding, 1963) (see the proof of Theorem 1 in the Supplementary Material).

Condition 2. *Any confidence interval for p contains \hat{p}_n . Moreover, $g(n) \in o(n^\gamma)$ for some $-\frac{1}{2} < \gamma < -\frac{1}{3}$.*

Define $D = \min_{I \in \mathcal{I}} d(p, \partial I)$ for a (random) p , where $d(p, \partial I)$ is the distance of p to the boundary of I . If the distance of p to \hat{p}_n is less than $D/2$ and the confidence interval for p has length $g(n) \leq D/2$, the confidence interval for p will be entirely contained in some $I \in \mathcal{I}$ (assuming it contains \hat{p}_n required by Condition 2). Therefore, a decision on which interval $I \in \mathcal{I}$ contains p is obtained on reaching the stopping time $\tau = \inf\{n : |\hat{p}_n - p| < D/2, g(n) < D/2\}$.

Let τ_1, \dots, τ_m be the stopping times of the m p-values and $\tau_{(1)} \leq \dots \leq \tau_{(m)}$ be their order statistic.

Theorem 1. *Let the density of p be bounded above by some finite constant. Assume $m \geq 3$. Under Condition 2, $\mathbb{E}(\tau_{(m-s)}) < \infty$ for $2 \leq s < m$.*

The proof of Theorem 1 is found in the Supplementary Material. For the Bonferroni (1936) correction, determining a $I \in \mathcal{I}$ containing the confidence interval of each hypothesis implies determining if the p-value of a hypothesis is above or below the constant testing threshold (subject to the overall $1 - \epsilon$ error probability), thus giving a decision on all but two hypotheses in finite expected time by Theorem 1. This result does not extend to multiple testing applications which depend on all individual decisions, for instance step-up or step-down procedures, in which case no algorithm can guarantee even a single decision in finite expected time (Section 3).

Section 1 of the Supplementary Material contains two results which highlight the asymptotic length of popular confidence intervals. These results are now used to establish that Condition 2 is satisfied for a variety of confidence sequences and the algorithms they use:

1. The length of Clopper and Pearson (1934) confidence intervals is $g(n) \propto n^{-1/2}(-\log \rho_n)^{1/2}$ (see the proof in the Supplementary Material), where ρ_n is a sequence controlling how the overall risk ϵ is spent. If $-\log(\rho_n) \propto \log(n)$ as in Gandy (2009); Gandy and Hahn (2014), then $g(n) \in o(n^\gamma)$ for any $-\frac{1}{2} < \gamma < -\frac{1}{3}$. Since the Clopper and Pearson (1934) intervals contain \hat{p}_n , Condition 2 is satisfied. The intervals are employed in the algorithms of Guo and Peddada (2008); Gandy and Hahn (2014).
2. Confidence intervals produced by the binomial confidence sequences of Robbins (1970); Lai (1976) satisfy $g(n) \propto n^{-1/2}\{\log(n \log n)\}^{1/2}$ (see the proof in the Supplementary Material). In (Lai, 1976, Section 3(A)) it is shown that the intervals contain \hat{p}_n , thus satisfying Condition 2. These confidence sequences are employed in the methods of Gandy and Hahn (2016); Ding et al. (2018).
3. Intervals produced by the confidence sequences of Darling and Robbins (1967a,b) have length $g(n) \propto n^{-1/2}\{\log(\log n)\}^{1/2}$ (Darling and Robbins, 1967a, Section 1) and are centered around the empirical mean, thus satisfying Condition 2.
4. Binomial confidence intervals of Armitage (1958) are employed in the methods of Fay et al. (2007); Gandy (2009). Being binomial exact intervals as the ones of Clopper and Pearson (1934), they satisfy Condition 2.

5. Condition 2 is satisfied for (asymptotic) maximum likelihood confidence intervals, given those are computed with a normal approximation. To be precise, the normal approximation produces confidence intervals of length $g(n) \propto n^{-1/2}$, thus satisfying Condition 2.
6. Bootstrap confidence intervals for p can be written as $[\hat{p} + q_\alpha, \hat{p} + q_{1-\alpha}]$ with $\alpha \in (0, 1)$, where q_α and $q_{1-\alpha}$ are the empirical α and $1 - \alpha$ quantiles of some bootstrap distribution F^* approximating an underlying distribution F that p is drawn from. Although specific to the application under consideration, the results of Theorem 1 apply given it has been verified that $q_{1-\alpha} - q_\alpha \in o(n^\gamma)$ for some $-\frac{1}{2} < \gamma < -\frac{1}{3}$. In particular, this is true if a normal approximation is used as in the case of maximum likelihood confidence intervals.

3 Extension to infinite expected runtime for multiple testing

For multiple testing applications having the property that with non-zero probability, no decision on any hypothesis can be made until it is known whether a certain hypothesis is rejected or non-rejected (as it is the case, for instance, for step-up and step-down procedures), under reasonable assumptions on the distribution of p-values, an algorithm with both a guarantee of correctness and a finite expected runtime for any number of decisions cannot exist.

Consider testing the m hypotheses H_{01}, \dots, H_{0m} satisfying the following condition.

Condition 3. *The p-values $p = (p_1, \dots, p_m)$ corresponding to H_{01}, \dots, H_{0m} have a joint distribution with support $[0, 1]^m$ and multivariate density $f_p(p)$. For all $\delta > 0$ there exists a $\kappa > 0$ such that $f_p(p) > \kappa$ on $[\delta, 1 - \delta]^m$.*

This condition is satisfied for many common densities. Assume testing is carried out with a step-up procedure that compares each p_i to a threshold value α_i , $i \in \{1, \dots, m\}$, where $\alpha_1 < \dots < \alpha_m$ (cf. Section 2). For any $0 < \eta < (\alpha_m - \alpha_{m-1})/4$, define $A = [\alpha_{m-1} + 2\eta, \alpha_m - 2\eta]^{m-1} \times [\alpha_m - \eta, \alpha_m + \eta]$. Let $B = [\delta_B, 1 - \delta_B]^m$ and choose $\delta_B \in (0, 1)$ in such a way that $A \cap B \neq \emptyset$.

Assuming the distribution of $p = (p_1, \dots, p_m)$ satisfies Condition 3 with $\delta = \delta_B$, draw a vector $\tilde{p} = (\tilde{p}_1, \dots, \tilde{p}_m)$ from f_p conditional on being in A . By definition of A , \tilde{p}_m is the largest value in \tilde{p} and will thus be compared to the threshold value α_m . By properties of a step-up procedure, as $\alpha_{m-1} < \tilde{p}_i < \tilde{p}_m$ for all $i \neq m$, no decision on any hypothesis can be made unless it is known whether \tilde{p}_m lies below or above α_m . In the former case, all hypotheses are rejected, in the latter case, all hypotheses are non-rejected. Under the additional assumption that the marginal distribution of p_m satisfies Condition 1, by Section 1.1, deciding H_{0m} requires an infinite expected time. Thus on the event $\{p \in A\}$, the expected time to decide any number of hypotheses is also infinite, $\mathbb{E}(\tau | p \in A) = \infty$, where τ denotes the number of Monte Carlo simulations.

By the law of total expectation, the unconditional expected time can be bounded as $\mathbb{E}(\tau) \geq \mathbb{E}(\tau | p \in A) \cdot \mathbb{P}(p \in A) = \infty$, where it was used that $\mathbb{E}(\tau | p \in A) = \infty$ and that $f_p(p) > \kappa > 0$ on B implies $\mathbb{P}(p \in A) \geq \mathbb{P}(p \in A \cap B) > 0$.

The above consideration proves an infinite expected runtime for step-up procedures for any desired number of decisions. It holds true, for instance, for the step-up procedures of Simes (1986); Hochberg (1988); Rom (1990); Benjamini and Hochberg (1995); Benjamini and Yekutieli (2001). A similar construction proves the same result for step-down procedures (Sidak, 1967; Holm, 1979; Shaffer, 1986). Extensions to other testing applications in which obtaining any decision can be made dependent on the decision of a single hypothesis are possible but application-specific.

4 Conclusions

According to Theorem 1, it can be decided in expected finite time for all but two hypotheses which interval $I \in \mathcal{I}$ in (2) their p-values are contained in. For independent (Bonferroni-type)




Figure 1: Survival function of the number of undecided hypotheses tested with the Benjamini and Hochberg (1995) procedure with threshold $\alpha = 0.1$. $R = 10^4$ repetitions. Number of hypotheses m from 10^1 (solid), 10^2 (dashed), 10^3 (dotted), 10^4 (dash-dotted).

testing, this implies at most two pending decisions in finite expected time.

For step-up and step-down procedures, the situation is more involved. This is because the individual decision on a hypothesis need not matter in step-up (step-down) procedures so long as there exists another hypothesis with a larger (smaller) p-value for which a decision is available. As a consequence, having two undecided hypotheses in the sense of Theorem 1 is not always representative of the actual number of decisions. Although in Section 3, this fact was used to show that under conditions, the expected time to compute any decision is infinite, in practice the decisions on a large number of hypotheses can often be computed quickly. For the Benjamini and Hochberg (1995) procedure with threshold $\alpha = 0.1$, Fig. 1 displays the survival function of the number of undecided hypotheses which remain if, out of $m \in \{10^1, 10^2, 10^3, 10^4\}$ hypotheses, for only $m - 2$ hypotheses it can be decided which $I \in \mathcal{I}$ in (2) their p-values are contained in. The figure is based on $R = 1000$ repetitions using p-values generated from the mixture distribution of Sandve et al. (2011), consisting of a proportion $\pi_0 = 0.8$ drawn from a uniform distribution in $[0, 1]$ and the remaining proportion $1 - \pi_0$ drawn from a $\text{beta}(0.5, 25)$ distribution. As can be seen, often only a handful of hypotheses remain without decision. On this note, a related question addresses the optimal strategy (where optimality is defined with respect to the expected number of erroneously classified hypotheses) for allocating Monte Carlo simulations to multiple hypotheses in order to maximize the accuracy of the testing result. This has been addressed in the literature (Hahn, 2019).

A Appendix

In the following, two versions of Hoeffding's inequality (Hoeffding, 1963) are used in several places. Specifically, let X_1, \dots, X_n be independent random variables which are almost surely bounded, that is $\mathbb{P}(X_i \in [a_i, b_i]) = 1$ for some constants a_i, b_i and all $i \in \{1, \dots, n\}$. Then by

(Hoeffding, 1963, Theorem 2) the empirical mean $\bar{X} = (X_1 + \dots + X_n)/n$ satisfies

$$\begin{aligned}\mathbb{P}(\bar{X} - \mathbb{E}(\bar{X}) \leq \delta) &\leq \exp\left(-\frac{2\delta^2 n^2}{\sum_{i=1}^n (b_i - a_i)^2}\right), \\ \mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| \geq \delta) &\leq 2 \exp\left(-\frac{2\delta^2 n^2}{\sum_{i=1}^n (b_i - a_i)^2}\right).\end{aligned}$$

In all following proofs, both inequalities will always be applied to a sum of independent Bernoulli random variables, in which case $b_i = 1$ and $a_i = 0$ for all $i \in \{1, \dots, n\}$, and the denominator of the exponentials above can be simplified to $\sum_{i=1}^n (b_i - a_i)^2 = n$.

A.1 Auxiliary lemmas

Lemma 1. *The two-sided Clopper and Pearson (1934) confidence interval I_n with coverage probability $1 - \rho_n$ based on n Monte Carlo simulations has length $|I_n| \leq 2(2n)^{-1/2}(-\log \rho_n)^{1/2}$.*

Proof. Suppose $S < n$ exceedances are observed among n Monte Carlo simulations. Let $\xi = (2n)^{-1/2}(-\log \rho_n)^{1/2}$ and regard the following probabilities conditional on S and n . The upper limit p_u of the interval I_n is the solution to $\mathbb{P}(X \leq S | p = p_u) = \rho_n$, where $X \sim \text{Binomial}(n, p)$. If $p > S/n + \xi$, by Hoeffding's inequality,

$$\mathbb{P}(X \leq S) = \mathbb{P}\left(\frac{X}{n} - \mathbb{E}\left(\frac{X}{n}\right) \leq \frac{S}{n} - \mathbb{E}\left(\frac{X}{n}\right)\right) \leq \exp\left(-\frac{2(S/n - p)^2 n^2}{n}\right) < \rho_n,$$

where the first equality was obtained by dividing by n and subtracting $\mathbb{E}(X/n) = p$ on both sides inside the probability, and where the Binomial variable X was expressed as $X = X_1 + \dots + X_n$ for some Bernoulli variables X_i , $i \in \{1, \dots, n\}$. Thus $p_u \leq S/n + \xi$. If $S = n$ then $p_u = 1$, implying $p_u \leq S/n + \xi$. Similarly, the lower limit p_l of I_n satisfies $p_l \geq S/n - \xi$. Together, $|I_n| = p_u - p_l \leq 2\xi$. \square

The proof of Lemma 1 is analogous to the one of (Gandy and Hahn, 2014, Lemma 2).

Lemma 2. *The confidence interval I_n produced by the Robbins (1970) and Lai (1976) Binomial confidence sequence based on n Monte Carlo simulations has length $|I_n| \leq n^{-1/2}\{\log(4n \log n)\}^{1/2}$ for large enough n .*

Proof. In (Lai, 1976, equation (3)) it is shown that $\mathbb{P}(p \notin I_n) \propto (n \log n)^{-1/2}$. Since I_n need not be centered around the maximum likelihood estimate \hat{p}_n , let $I_n^s \supseteq I_n$ be the smallest symmetric interval around \hat{p}_n containing I_n . Denote $I_n^s = [\hat{p}_n - t_n, \hat{p}_n + t_n]$ for some $t_n > 0$. Write \hat{p}_n as the average of n independent Bernoulli(p) random variables. By Hoeffding's inequality, using $\mathbb{E}(\hat{p}_n) = p$, one obtains $\mathbb{P}(p \notin I_n^s) = \mathbb{P}(|\hat{p}_n - p| > t_n) \leq 2 \exp(-2nt_n^2)$. Since $\mathbb{P}(p \notin I_n^s) \leq \mathbb{P}(p \notin I_n)$, $\mathbb{P}(p \notin I_n^s)$ is at least of order $(n \log n)^{-1/2}$. Thus, $t_n \leq (4n)^{-1/2}\{\log(4n \log n)\}^{1/2}$ for large enough n . \square

A.2 Proof of Theorem 1

Proof. The cdf of $D = \min_{I \in \mathcal{I}} d(p, \partial I)$ is bounded above by

$$\mathbb{P}(D \leq t) = \mathbb{P}(\exists I \in \mathcal{I} : d(p, \partial I) \leq t) \leq \sum_{I \in \mathcal{I}} \int_0^t d(p, \partial I) f_p(p) dp \leq (m+1) \int_0^t U dp \leq (m+1)Ut,$$

where it was used that the cardinality of \mathcal{I} is at most $m+1$, that the distance function d is bounded above by 1, and that the density f_p of the p-values is bounded above by a constant U .

In the following, an upper bound on the survival function $\mathbb{P}(\tau > t)$ will be derived. First, $\mathbb{P}(|\hat{p}_t - p| \geq D/2) = \mathbb{P}(|\hat{p}_t - \mathbb{E}(\hat{p}_t)| \geq D/2) \leq 2 \exp(-D^2 t/2)$ by Hoeffding's inequality, where

it was used that the maximum likelihood estimate \hat{p}_t can be expressed as an average of t Bernoulli(p) random variables. Second, using $g(t) \in o(t^\gamma)$ by Condition 2, there exists a $t_0 > 0$ such that the event $\{g(t) < D/2\}$ is implied by $\{D > t^\gamma\}$ for all $t > t_0$. Indeed, $g(t) \in o(t^\gamma)$ and $Dt^{-\gamma} > 1$ imply the existence of a $t_0 > 0$ such that $g(t) < g(t)Dt^{-\gamma} < D/2$ for all $t > t_0$.

The survival function $\mathbb{P}(\tau > t)$ can now be bounded above by conditioning on D . For $t > t_0$,

$$\begin{aligned}\mathbb{P}(\tau > t) &= \mathbb{P}(\tau > t | D \leq t^\gamma) \mathbb{P}(D \leq t^\gamma) + \mathbb{P}(\tau > t | D > t^\gamma) \mathbb{P}(D > t^\gamma) \\ &\leq (m+1)Ut^\gamma + \mathbb{P}(|\hat{p}_t - p| \geq D/2 \vee g(t) \geq D/2 | D > t^\gamma) \\ &\leq (m+1)Ut^\gamma + \mathbb{P}(|\hat{p}_t - p| \geq D/2 | D > t^\gamma) \\ &\leq (m+1)Ut^\gamma + 2 \exp\left(-\frac{1}{2}t^{2\gamma+1}\right),\end{aligned}$$

where it was used that $\mathbb{P}(\tau > t | D \leq t^\gamma) \leq 1$, $\mathbb{P}(D > t^\gamma) \leq 1$ and that $\tau > t$ if either $|\hat{p}_t - p| \geq D/2$ or $g(t) \geq D/2$ by the definition of τ in Section 2. The latter can be omitted as $\{D > t^\gamma\}$ implies $\{g(t) < D/2\}$ for $t > t_0$. Using $\gamma > -1/2$ by Condition 2, $t^{2\gamma+1} \rightarrow \infty$ in the argument of the exponential function as $t \rightarrow \infty$ and thus $\mathbb{P}(\tau > t) \in O(t^\gamma)$.

Using the fact that the cumulative distribution function (cdf) of the r th order statistic $X_{(r)}$ of $n \in \mathbb{N}$ independent and identically distributed random variables X_1, \dots, X_n with cdf F_X can be expressed as $F_{X_{(r)}}(x) = \sum_{i=r}^n \binom{n}{i} F_X^i(x) (1 - F_X(x))^{n-i}$ for $r \in \{1, \dots, n\}$ (David and Nagaraja, 2003), the expectation of $\tau_{(m-s)}$ can be bounded as

$$\begin{aligned}\mathbb{E}(\tau_{(m-s)}) &= \int_0^\infty 1 - F_{\tau_{(m-s)}}(t) dt \\ &= \int_0^\infty \sum_{i=0}^{m-s-1} \binom{m}{i} F_\tau^i(t) (1 - F_\tau(t))^{m-i} dt \leq \sum_{i=0}^{m-3} \binom{m}{i} \int_0^\infty \mathbb{P}(\tau > t)^{m-i} dt,\end{aligned}$$

where it was used that the stopping times are non-negative, that $F_\tau(t) \leq 1$ and that $2 \leq s < m$ implies $m-s-1 \leq m-3$. Using $\mathbb{P}(\tau > t) \in O(t^\gamma)$, the integrals $\int_0^\infty \mathbb{P}(\tau > t)^{m-i} dt$ behave like $\int_0^\infty t^{\gamma(m-i)} dt$ and converge as $\gamma < -1/3$ (cf. Condition 2) and $m \geq 3$ imply $\gamma(m-i) < -1$ for all $i \in \{0, \dots, m-3\}$. \square

References

- Andrews, D. and Buchinsky, M. (2000). A Threestep Method for Choosing the Number of Bootstrap Repetitions. *Econometrica*, 68(1):23–51.
- Andrews, D. and Buchinsky, M. (2003). Evaluation of a three-step method for choosing the number of bootstrap repetitions. *J Econometrics*, 103(1-2):345–386.
- Armitage, P. (1958). Numerical Studies in the Sequential Estimation of a Binomial Parameter. *Biometrika*, 45(1-2):1–15.
- Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J Roy Stat Soc B Met*, 57(1):289–300.
- Benjamini, Y. and Yekutieli, D. (2001). The control of the false discovery rate in multiple testing under dependency. *Ann Stat*, 29(4):1165–1188.
- Besag, J. and Clifford, P. (1991). Sequential Monte Carlo p-values. *Biometrika*, 78(2):301–304.
- Bonferroni, C. (1936). Teoria statistica delle classi e calcolo delle probabilità. *Pubblicazioni del R Istituto Superiore di Scienze Economiche e Commerciali di Firenze*, 8:3–62.

- Clopper, C. and Pearson, E. (1934). The Use of Confidence or Fiducial Limits Illustrated in the Case of the Binomial. *Biometrika*, 26(4):404–413.
- Darling, D. and Robbins, H. (1967a). Confidence sequences for mean, variance, and median. *P Natl Acad Sci USA*, 58(1):66–68.
- Darling, D. and Robbins, H. (1967b). Iterated logarithm inequalities. *P Natl Acad Sci USA*, 57(5):1188–92.
- David, N. and Nagaraja, H. (2003). *Order Statistics*. Wiley.
- Davidson, R. and MacKinnon, J. (2000). Bootstrap tests: How many bootstraps? *Economet Rev*, 19(1):55–68.
- Ding, D., Gandy, A., and Hahn, G. (2018). A simple method for implementing Monte Carlo tests. *arXiv:1611.01675*, pages 1–17.
- Fay, M. and Follmann, D. (2002). Designing Monte Carlo implementations of permutation or bootstrap hypothesis tests. *Am Stat*, 56(1):63–70.
- Fay, M., Kim, H.-J., and Hachey, M. (2007). On using truncated sequential probability ratio test boundaries for Monte Carlo implementation of hypothesis tests. *J Comput Graph Stat*, 16(4):946–967.
- Gandy, A. (2009). Sequential Implementation of Monte Carlo Tests With Uniformly Bounded Resampling Risk. *J Am Stat Assoc*, 104(488):1504–1511.
- Gandy, A. and Hahn, G. (2014). MMCTest – A Safe Algorithm for Implementing Multiple Monte Carlo Tests. *Scand J Stat*, 41(4):1083–1101.
- Gandy, A. and Hahn, G. (2016). A Framework for Monte Carlo based Multiple Testing. *Scand J Stat*, 43(4):1046–1063.
- Gandy, A. and Hahn, G. (2017). QuickMMCTest: quick multiple Monte Carlo testing. *Stat Comput*, 27(3):823–832.
- Guo, W. and Peddada, S. (2008). Adaptive Choice of the Number of Bootstrap Samples in Large Scale Multiple Testing. *Stat Appl Genet Mol Biol*, 7(1):1–16.
- Hahn, G. (2019). Optimal allocation of Monte Carlo simulations to multiple hypothesis tests. *Stat Comput*, pages 1–16.
- Hochberg, Y. (1988). A sharper Bonferroni procedure for multiple tests of significance. *Biometrika*, 75(4):800–802.
- Hoeffding, W. (1963). Probability inequalities for sums of bounded random variables. *J Am Stat Assoc*, 58(301):13–30.
- Holm, S. (1979). A Simple Sequentially Rejective Multiple Test Procedure. *Scand J Stat*, 6(2):65–70.
- Kim, H.-J. (2010). Bounding the resampling risk for sequential Monte Carlo implementation of hypothesis tests. *J Stat Plan Infer*, 140(7):1834–1843.
- Lai, T. (1976). On Confidence Sequences. *Ann Stat*, 4(2):265–280.
- Lin, D. (2005). An efficient Monte Carlo approach to assessing statistical significance in genomic studies. *Bioinformatics*, 21(6):781–787.

- Robbins, H. (1970). Statistical methods related to the law of the iterated logarithm. *Ann Math Statist*, 41(5):1397–1409.
- Rom, D. (1990). A sequentially rejective test procedure based on a modified Bonferroni inequality. *Biometrika*, 77(3):663–665.
- Sandve, G., Ferkingstad, E., and Nygård, S. (2011). Sequential Monte Carlo multiple testing. *Bioinformatics*, 27(23):3235–3241.
- Shaffer, J. (1986). Modified Sequentially Rejective Multiple Test Procedures. *J Am Stat Assoc*, 81(395):826–831.
- Sidak, Z. (1967). Rectangular confidence regions for the means of multivariate normal distributions. *J Am Stat Assoc*, 62(318):626–633.
- Silva, I. and Assunção, R. (2013). Optimal generalized truncated sequential Monte Carlo test. *J Multivariate Anal*, 121:33–49.
- Silva, I. and Assunção, R. (2018). Truncated sequential Monte Carlo test with exact power. *Braz J Probab Stat*, 32(2):215–238.
- Silva, I., Assunção, R., and Costa, M. (2009). Power of the sequential Monte Carlo test. *Sequential Anal*, 28(2):163–174.
- Simes, R. (1986). An improved Bonferroni procedure for multiple tests of significance. *Biometrika*, 73(3):751–754.
- van Wieringen, W., van de Wiel, M., and van der Vaart, A. (2008). A Test for Partial Differential Expression. *J Am Stat Assoc*, 103(483):1039–1049.
- Wald, A. (1945). Sequential Tests of Statistical Hypotheses. *Ann Math Statist*, 16(2):117–186.