

A global model for predicting the arrival of imported dengue infections

Jessica Liebig^{1*}, Cassie Jansen², Dean Paini³, Lauren Gardner^{1,4,5}, Raja Jurdak^{1,6,7}

¹Data61, Commonwealth Scientific and Industrial Research Organisation
Brisbane, Queensland, Australia

²Communicable Diseases Branch, Department of Health
Brisbane, Queensland, Australia

³Health & Biosecurity, Commonwealth Scientific and Industrial Research Organisation
Canberra, Australian Capital Territory, Australia

⁴Department of Civil Engineering, Johns Hopkins University
Baltimore, Maryland, USA

⁵School of Civil and Environmental Engineering, University of New South Wales
Sydney, New South Wales, Australia

⁶School of Electrical Engineering and Computer Science, Queensland University of Technology
Brisbane, Queensland, Australia

⁷School of Computer Science and Engineering, University of New South Wales
Sydney, New South Wales, Australia

Abstract

With approximately half of the world's population at risk of contracting dengue, this mosquito-borne disease is of global concern. International travellers significantly contribute to dengue's rapid and large-scale spread by importing the disease from endemic into non-endemic countries. To prevent future outbreaks and dengue from establishing in non-endemic countries, knowledge about the arrival time and location of infected travellers is crucial. We propose a network model that predicts the monthly number of dengue-infected air passengers arriving at any given airport. We consider international air travel volumes to construct weighted networks, representing passenger flows between airports. We further calculate the probability of passengers, who travel through the international air transport network, being infected with dengue. The probability of being infected depends on the destination, duration and timing of travel. Our findings shed light onto dengue importation routes and reveal country-specific reporting rates that have been until now largely unknown. This paper provides important new knowledge about the spreading dynamics of dengue that is highly beneficial for public health authorities to strategically allocate the often limited resources to more efficiently prevent the spread of dengue.

INTRODUCTION

The well connected structure of the global air transportation network and the steadily increasing volume of international travel has a vast impact on the rapid, large-scale spread of arboviral and other diseases [1, 2, 3, 4, 5, 6, 7]. A recent example of disease introduction to a novel region is the spread of the Zika virus from Brazil to Europe, the United States and other countries, which prompted the World Health Organisation (WHO) to announce a public health emergency of international concern in early 2016. Investigations confirmed that international viraemic travellers were a major contributing factor to the rapid spread [8].

With an estimated 50-100 million symptomatic infections each year [9, 10], dengue is ranked the most impor-

tant mosquito-borne disease [11, 12]. The rapid geographic spread is, to a great extent, driven by the increase in international air travel [13, 14]. In addition, dengue is severely under-reported, making it extremely challenging to monitor and prevent the spread of the disease. Presumably, 92% of symptomatic infections are not reported to health authorities [10]. Low reporting rates can have many reasons, including low awareness levels and misdiagnosis [9, 15].

Due to the rapid global spread of dengue as well as severe under-reporting, many countries are facing the threat of ongoing local transmission in the near future [11]. In non-endemic countries, local outbreaks are usually triggered by an imported case [16], a person who acquired the disease overseas and transmitted the virus to local mosquitoes. To prevent ongoing dengue transmission in non-endemic countries, it is critical to forecast the importation of disease cases into these areas and move from responsive contain-

*Corresponding author: jess.liebig@csiro.au

ment of dengue outbreaks to proactive outbreak mitigation measures.

The majority of existing models forecast relative rather than absolute risk of dengue importation and are unable to predict the total number of imported disease cases [13, 17, 18]. The few models that can predict absolute numbers are region-specific rather than global [19, 20, 21]. The most recently proposed model estimates the total number of imported dengue cases for 27 European countries [21], however, the model has several limitations: (i) Monthly incidence rates were based on dengue cases reported to the World Health Organisation (WHO) despite dengue being under-reported and the general consensus that the actual number of cases is much higher than the figures published by the WHO [10, 9]; (ii) Only 16 countries were considered as possible sources of importation. The authors reason that these 16 countries contribute 95% of all global dengue cases, referring to numbers published by the WHO. Since African countries do not report to the WHO, and dengue remains an under-reported disease in many other countries [22, 23, 24, 25], it is likely that the percentage contribution to the number of global dengue cases by the 16 selected countries is strongly biased; (iii) Seasonal distributions of dengue cases were inferred based on information from only two source countries (Latin American countries were assumed to have similar seasonalities to Brazil, while Thailand served as a proxy for countries in South-East Asia). The assertion that all countries within a given global region experience similar seasonal fluctuations in dengue infections is likely inaccurate. For example, dengue notifications peak between April and December in Thailand, while Indonesia reports the highest number of dengue cases from November to April [26].

The contribution of this paper is twofold. First, we develop a network model that overcomes the limitations of previous models by employing global air passenger volumes, country-specific dengue incidence rates and country-specific temporal infection patterns. We construct weighted directed networks, using data collected by the International Air Transportation Association (IATA) to capture the movement of air passengers. We calculate monthly, country-specific dengue incidence rates by combining data from the Global Health Data Exchange [27], the most comprehensive health database, and known seasonal patterns in reported dengue infections [26]. Further, we distinguish between two categories of travellers: returning residents and visitors. The number of days people from these two categories spend in an endemic country, and therefore the risk of being infectious on arrival, vary greatly. The model predicts the number of imported dengue cases per month for any given airport and can be applied with relative ease to other vector-borne diseases of global concern, such as malaria, Zika or chikungunya.

Second, we apply the model to infer time-varying, region-specific reporting rates, defined as the ratio of reported to actual infections. Dengue reporting rates vary greatly across space and time, often by several orders of magnitude,

and hence are difficult to determine [10]. The usual approach towards estimating country-specific reporting rates is to carry out cohort or capture-recapture studies that can be costly, are time consuming and may be biased [28]. Consequently, dengue reporting-rates remain unknown for most countries [10].

In this paper we focus on those countries that are most at risk of dengue introduction, i.e. non-endemic countries with vector presence. These countries will have the greatest benefit from our model as knowledge about the likely arrival times and places of infected people is crucial to prevent local outbreaks.

MATERIALS AND METHODS

CSIRO’s human research ethics committee CSSHREC has approved this study (approval number: Ethics Clearance 142/16). All data were analysed anonymously and individuals cannot be identified.

IATA Data

The International Air Transportation Association (IATA) has approximately 280 airline members who together contribute to approximately 83% of all air traffic. Data is collected in form of travel routes, detailing the origin, destination and stopover airports. It contains over 10,000 airports in 227 different countries and dependencies. For each route the total number of passengers per month is given. We do not have any information on stopover times and whether passengers are leaving the airport during their stopover and therefore assume that all passengers continue their journey to the final destination instantly. Table S1 lists the IATA 3-Letter Codes used to abbreviate airports in the main manuscript. As the recorded itineraries do not include any travel on chartered flights, we compare the IATA passenger volumes to official airport passenger statistics [29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47] to quantify the potential discrepancies between actual travel patterns and that reported by IATA. Table S2 lists the countries where the difference in passenger numbers is greater than 15% (at country level) and countries where airport statistics were not available and the tourist data suggests inaccuracies in the IATA data (i.e. the number of tourists arriving in a particular country is larger than the total number of passengers arriving). We also excluded Singapore as a source of importation for Australia for the following reason: The Department of Home Affairs publishes Arrival Card data [48] that can be used to validate the IATA data. A comparison of the monthly travel volume from Singapore to Australia revealed that the IATA data overestimates travel volumes by approximately 112% on average in 2011 and 2015. This may be due to individuals who travel from other countries to Singapore and then directly continue to Australia and do not book their entire trip in one itinerary (this would be recorded as two separate trips in the IATA data

that cannot be linked to each other). Due to this large discrepancy in the travel data we believe that our model will significantly overestimate the number of dengue infections imported from Singapore, and therefore exclude it as a source country for Australia.

The air transportation network

We begin by constructing twelve weighted, directed networks, using IATA data, to represent the monthly movement of air passengers during a given year. The networks are denoted $\mathcal{G}_m = (V, E)$, with $m = 1, \dots, 12$ indicating the month of the year. The node set V comprises more than 10,000 airports recorded by IATA. To distinguish the travellers by their country of embarkation, we represent the edges of the network as ordered triples, $(i, j, \omega_{i,j}(c, k)) \in E$, where $i, j \in V$ and $\omega_{i,j}(c, k)$ is a function that outputs the number of passengers who initially embarked in country c with final destination airport k and travel from airport i to airport j as part of their journey.

Incidence rates and seasonal distributions

Calculating the number of infected passengers requires daily infection probabilities. We derive these from country-level yearly estimates of symptomatic dengue incidence rates that are published together with their 95% confidence intervals by the Global Health Data Exchange [27]. The estimates are obtained using the model published in [10] and account for under-reporting.

We first deduce monthly incidence rates using information on dengue seasonality published by the International Association for Medical Assistance to Travellers [26]. To do so we associate a weight with each month that indicates the intensity of transmission. To assign the weights we use a modified cosine function with altered period that matches the length of the peak-transmission season. The function is shifted and its amplitude adjusted so that its maximum occurs midway through the peak-season with value equal to the length of the peak-season divided by 2π . The months outside the peak-season receive a weight of one if dengue transmission occurs year around and a weight of zero if dengue transmission ceases outside the peak-season. The weights are then normalised and multiplied by the yearly incidence rate for the corresponding country. Normalising the weights ensures that the sum of the monthly incidence rates is equal to the yearly incidence rate. To calculate the lower and upper bounds of the monthly incidence rates, we multiply the normalised weights by the lower and upper bounds of the 95% confidence interval given for the yearly incidence rates.

The average probability, $\beta_{c,m}$, of a person becoming infected on any given day during month m in country c is then given by

$$\beta_{c,m} = 1 - e^{-\gamma_{c,m}/d_m}, \quad (1)$$

where $\gamma_{c,m}$ is the monthly dengue incidence rate in country c during month m and d_m is number of days in month m . Note that Equation (1) converts the daily incidence rate into the probability of a single person becoming infected with dengue on any given day during month m .

Inferring the number of infected passengers

Next, we present a mathematical model that approximates the number of dengue-infected people for each edge in the network $\mathcal{G}_m(V, E)$. The time between being bitten by an infectious mosquito and the onset of symptoms is called the intrinsic incubation period (IIP). This period closely aligns with the latent period, after which dengue can be transmitted to mosquitoes [49]. The IIP lasts between 3 and 14 days (on average 5.5 days) and was shown to follow a gamma distribution of shape 53.8 and scale equal to 0.1 [50]. After completion of the IIP a person is infectious for approximately 2 to 10 days (on average 5 days) [51, 50]. The length of the infectious period was shown to follow a gamma distribution of shape 25 and scale equal to 0.2 [50]. We denote the sum of the IIP and the infectious period by n , which is rounded to the nearest integer after the summation. For travellers to import the infection from country c into a new location r they must have been infected with dengue within the last $n - 1$ days of their stay in country c . We now consider the following two cases: $t_c \geq n - 1$ and $t_c < n - 1$, where t_c is number of days spent in country c before arriving in region r . Since we do not know the exact date of arrival for travellers, we assume that arrival and departure dates fall within the same month and hence $\beta_{c,m}$ is the same for every day during the travel period.

If $t_c \geq n - 1$, that is the individual spent more time in country c than the sum of the lengths of the IIP and the infectious period, the probability of not being infected on return is equal to $(1 - \beta_{c,m})^{t_c} + [1 - (1 - \beta_{c,m})^{t_c - (n-1)}]$. The first term covers the possibility that the individual did not get infected whilst staying in country c and the second term covers the possibility that the individual got infected and recovered before arriving at a given airport (see Fig S1). Hence, the probability of a person, who arrives at a given airport from country c during month m , being infected with dengue is given by

$$\begin{aligned} p_{c,m} &= 1 - \left[(1 - \beta_{c,m})^{t_c} + 1 - (1 - \beta_{c,m})^{t_c - (n-1)} \right] \\ &= (1 - \beta_{c,m})^{t_c - (n-1)} - (1 - \beta_{c,m})^{t_c}. \end{aligned} \quad (2)$$

If $t_c < n - 1$, that is the individual spent less time in country c than the sum of the lengths of the IIP and the infectious period, the probability of not being infected on return is equal to $(1 - \beta_{c,m})^{t_c}$, which covers the possibility that the individual did not get infected whilst staying in country c . Since $t_c < n - 1$, the probability of recovery before arriving at a given airport is zero. Hence, the probability of a person, who arrives from country c at a given airport

during month m , being infected with dengue is given by

$$p_{c,m} = 1 - (1 - \beta_{c,m})^{t_c}. \quad (3)$$

We distinguish between two different types of travellers arriving at a given airport of region r : returning residents and visitors. We define a returning resident as a traveller who resides in region r and a visitor as a traveller who resides in country c and visits region r . Returning residents are expected to have stayed a couple of weeks in the endemic country, while visitors may have spent their whole life in the country.

Since we lack information on how long each individual spent in country c before arriving at an airport of region r , we substitute parameter t_c by $\langle t \rangle_c^{res}$ if the person is a returning resident, $\langle t \rangle_c^{res}$ being the average number of days a returning resident spends in country c before returning home. If the person is a visitor, parameter t_c is substituted by $\langle t \rangle_c^{vis}$, the average number of days a visitor spends in country c before arriving at an airport of region r . We distinguish between returning residents and visitors since $\langle t \rangle_c^{res} \ll \langle t \rangle_c^{vis}$.

We assume that the length of stay for returning residents follows a normal distribution with mean equal to 15 days and standard deviation of 2, i.e. $\langle t \rangle_c^{res} \sim \mathcal{N}(15, 2)$. A previous study has shown that employees around the world are on average entitled to approximately 15 days of annual leave [52]. On the other hand, visitors likely spent all their lives in the endemic country. We assume that $\langle t \rangle_c^{vis} \sim \mathcal{N}(\mu_{vis}, 0.1\mu_{vis})$, where μ_{vis} is equal to c 's me-

dian population age. Median population ages by country are published in the World Factbook by the Central Intelligence Agency [53].

For simplicity we do not take immunity to the different dengue strains into consideration.

Proportion of returning residents and visitors

Lastly, we need to infer the proportions of returning residents and visitors. As this information is not contained in the IATA itineraries, we use international tourism arrival data from the World Tourism Organisation [54]. The data contains the yearly number of international tourist arrivals by air for each destination country. From the IATA data we calculate the total number of arrivals per year for each country and hence can infer the ratio of visitors to returning residents. As we lack sufficient data, we assume that the ratio of visitors to residents is the same for each month.

Calculating the absolute number of infected passengers

Given the above, we can now determine the number of infected passengers $I_{k,m}$ arriving at airport k during month m as follows:

$$I_{k,m} = \sum_{i,j,c} \omega_{i,j}(c, k) [qp_{c,m}^{res} + (1 - q)p_{c,m}^{vis}], \quad (4)$$

where q is the proportion of residents inferred from the international tourism arrival data,

$$p_{c,m}^{res} = \begin{cases} (1 - \beta_{c,m})^{\langle t \rangle_c^{res} - (n-1)} - (1 - \beta_{c,m})^{\langle t \rangle_c^{res}} & \langle t \rangle_c^{res} \geq n - 1 \\ 1 - (1 - \beta_{c,m})^{\langle t \rangle_c^{res}} & \langle t \rangle_c^{res} < n - 1, \end{cases} \quad (5)$$

and

$$p_{c,m}^{vis} = \begin{cases} (1 - \beta_{c,m})^{\langle t \rangle_c^{vis} - (n-1)} - (1 - \beta_{c,m})^{\langle t \rangle_c^{vis}} & \langle t \rangle_c^{vis} \geq n - 1 \\ 1 - (1 - \beta_{c,m})^{\langle t \rangle_c^{vis}} & \langle t \rangle_c^{vis} < n - 1. \end{cases} \quad (6)$$

Evaluation of the models uncertainty

We performed a thousand runs of the model for each edge in the network, drawing the parameters from their respective distributions, to calculate the mean and standard deviation of dengue-infected passengers. In addition, we have conducted a global sensitivity analysis to identify the model parameters with the greatest influence. We used Sobol's method [55] with 100,000 samples to carry out the sensitivity analysis. The parameter ranges are shown in Table 1. The analysis was done with SALib [56], an open-source Python library.

Parameter	Range
$\beta_{c,m}$	[0.000001, 0.000445]
t_c (days)	[1, 29200]
n (days)	[5, 24]

Table 1: The model parameter ranges used in Sobol's method.

RESULTS

We run our model for two different years to explore the robustness of the proposed methodology. Specifically, the analysis is conducted for 2011 and 2015. The results for the year 2015 are presented in the main manuscript, while




Figure 1: Predicted dengue importations for August 2015. The map shows the output of our model for August 2015. The area of a node increases with the number of dengue cases imported through the corresponding airport. Airports that are predicted to not receive any infections are not shown on the map. Endemic countries are coloured dark grey. Countries that are non-endemic and where dengue vectors *Aedes aegypti* and/or *Aedes albopictus* are present are coloured in light grey. The blue circles correspond to the top ten airports identified in Fig 2. The map was created with the Python GeoPandas package and publicly available shapefiles from Natural Earth (<http://www.naturalearthdata.com/>).

the results for 2011 are presented in the supplementary material. Fig 1 shows the number of predicted imported dengue infections per airport for August 2015, where the area of a node increases with the number of dengue cases imported through the corresponding airport. The map clearly shows that many non-endemic regions where the dengue-transmitting vectors *Aedes aegypti* or *Aedes albopictus* are present (coloured in light grey) have airports that are predicted to receive a high number of dengue infections. For a list of dengue endemic and non-endemic countries see Table S3. As resources for the control and prevention of dengue are often limited [57], these countries face a high risk of future endemicity.

In Fig 2 and Fig S2 we plot the number of predicted dengue importations over time for the ten airports that receive the highest number of cases, lie in non-endemic regions with vector presence and where local cases have been reported in the past (more detailed plots with confidence intervals are shown in Fig S3). While the majority of airports listed in Fig 2 and Fig S2 are predicted to receive between 50 and 150 cases each month, Miami International Airport (MIA) is estimated to receive between 146 and 309 cases each month during both years. With Orlando International Airport (MCO) and Fort Lauderdale-Hollywood International Airport (FLL) also represented amongst the airports with the highest number of imported cases, Florida faces a high risk of local dengue outbreaks. Los Angeles International Airport (LAX) is predicted to receive the second highest number of imported cases. In 2011 its monthly predictions vary between 97 and 205 cases and in 2015 between 113 and 253 cases. The remaining airports listed in Fig 2 and Fig S2 are located in France, Germany, the Netherlands,

Texas, and Queensland, Australia. A full ranking of all airports located in non-endemic countries with vector presence can be found in Table S4 of the supplementary material.




Figure 2: Predicted monthly dengue importations by airport for 2015. The number of predicted imported dengue infections for the top ten airports in non-endemic countries/states with vector presence for each month in 2015. A break in a line indicates that the corresponding airport was not amongst the top ten during the respective month. Airports are abbreviated using the corresponding IATA code. A full list of abbreviations can be found in the supplementary material (see Table S1).

In addition to calculating the number of imported dengue infections per airport, the model further provides the number of infected passengers travelling between any two airports, thus revealing common importation routes. Table 2 and Table S5 list the routes that carry the highest number of infected passengers whose final destinations lie in non-endemic countries with vector presence. Table S6 lists the




Figure 3: Predicted dengue infections imported by returning residents and visitors in 2015. Here we show the results for non-endemic countries/states with vector presence with the highest number of predicted imported dengue cases in 2015. The bars are stacked to distinguish between returning residents (green) and visitors (blue). The blue solid line corresponds to the total number of imported cases. The error bars correspond to the model’s coefficient of variation (see Material and methods). The six countries were selected because they are predicted to receive the highest number of dengue importations, are non-endemic and dengue vectors are established.

routes that carry the highest number of infected passengers whose final destinations lie in non-endemic countries irrespective of whether vectors are present. For example, the route between Denpasar and Perth is ranked third in 2011 in Table S6, but it is not considered in the ranking shown in Table S5, as there are no vectors in Perth. Fig S4 shows a map of all importation routes into non-endemic countries with vector presence.

Orig.	Dest.	Pax	Month
SJU (Puerto Rico)	MCO (Florida)	51	Jul
PTP (Guadeloupe)	ORY (France)	37	Aug
FDF (Martinique)	ORY (France)	34	Aug
SJU (Puerto Rico)	FLL (Florida)	32	Jul
TPE (Taiwan)	LAX (California)	31	Aug
GRU (Brazil)	MIA (Florida)	29	Apr
DEL (India)	KBL (Afghanistan)	27	Aug
GDL (Mexico)	LAX (California)	24	Aug
CUN (Mexico)	MIA (Florida)	24	Aug
CUN (Mexico)	LAX (California)	22	Aug

Table 2: The ten routes with the highest predicted number of dengue-infected passengers with final destinations in non-endemic countries with vector presence. The table lists the direct routes with the highest predicted volume of dengue-infected passengers who continue to travel to non-endemic regions with vector presence and where local outbreaks have been reported in the past. The last column records the month during which the highest number of infected passengers are predicted.

In both years the highest predicted number of infected passengers are recorded during the northern hemisphere’s summer. The route between São Paulo International Airport (GRU) and Miami International Airport (MIA) is the exception, where the highest number of infected passen-

gers is predicted during April. The routes with the highest estimated number of dengue-infected passengers terminate at airports in countries that are non-endemic and where dengue-transmitting vectors are present.

Returning residents and visitors

Next, we aggregate airports by country/state to predict the number of imported dengue infections on a coarser level. For non-endemic countries that cover an area larger than 5,000,000 km² and where dengue vectors are present we aggregate airports by state. These countries are Russia, the United States of America and Australia. The comparison between passenger volumes recorded by IATA and official airport statistics indicated that the IATA data for Russia may be inaccurate, i.e. the difference in passenger numbers is larger than 15% (see Material and Methods). Hence, we did not perform a state-level analysis for this country. In Australia vectors are present only in Queensland [58]. While vectors have been observed in more than 40 different US states, autochthonous cases have been reported only in California, Florida, Hawaii and Texas [59].

Our model separately calculates the number of dengue-infected people amongst returning residents and visitors and hence we can identify which of these groups is more likely to import the disease into a given country or state. Fig 3 and Fig S5 show the results for six non-endemic countries/states with vector presence that are predicted to receive the highest number of dengue importations each month. Results for the remaining countries and states are shown in Figs S6 - S11. We observe that the contributions of returning residents and visitors to the total number of imported dengue




Figure 4: Predicted percentage contribution of dengue importations by country of acquisition in 2015. The predicted percentage contribution by source country and month in 2015. The size and colour of the circles indicate the percentage contribution of the corresponding country to the total number of imported cases. The y -labels indicate the yearly percentage contribution of the corresponding source country.

infections is predicted to vary greatly between the different countries and states. In Florida and Queensland returning residents are predicted to be the main source of dengue importation. In France and Italy approximately one third of all dengue infections are predicted to be imported by visitors while in Spain visitors import around 75% of all imported cases. For Switzerland we do not have any information about the ratio of returning residents to visitors. For the United States there is evidence in the form of surveillance reports that returning residents are indeed the main contributors to dengue importations [60]. For Queensland we predict that 95% and 94% of infections were imported by returning residents in 2011 and 2015, respectively. Our predictions are supported by Queensland’s dengue notification data (provided by Queensland Health), showing that 97% and 92% of all dengue importations in 2011 and 2015, respectively, were imported by returning residents.

Countries of acquisition

In addition to being able to distinguish between returning residents and visitors, the model also divides the imported cases according to their places of acquisition. Fig 4 and Fig S12 show the model’s estimated percentage contribution of dengue importations by source country.

Florida is predicted to import most infections from the

Caribbean and Latin America, with infections acquired in Puerto Rico (PRI) predicted to peak during June and July and infections acquired in Brazil predicted to peak between January and April. We hypothesise that Florida receives such a high number of imported dengue cases due to its close proximity to the Caribbean, which has been endemic since the 1970s [61]. France is predicted to receive many infections from the Caribbean, in particular from Martinique and Guadeloupe which are French overseas regions and hence a high volume of air traffic from these regions to metropolitan France is expected. These predictions align with the fact that outbreaks of dengue in France coincide with outbreaks in the French West Indies, where most reported cases are acquired [62, 63]. In Italy the model predicts that the most common countries of acquisition are India and Brazil. India and Brazil are also the most common countries of acquisition for Switzerland in 2011. In 2015 Switzerland is predicted to receive most of their dengue importations from India and Thailand. Spain is predicted to import the majority of infections from Latin America and the Caribbean. For Queensland the model predicts that imported cases are acquired mostly in South-East Asia with Indonesia being the largest source. This is in agreement with previous studies [64] and the dengue case data that was provided by Queensland Health. In addition, we performed a rank-based validation of these results.




Figure 5: Rank-based validation and correlation between reported and predicted imported cases for Queensland in 2015. (A) Countries are ranked by the total number of predicted and reported imported dengue cases. The reported ranking is then plotted against the predicted ranking. Countries that were ranked by the model, but did not appear in the dataset receive a rank of $i + 1$, where i is the number of unique importation sources according to the dengue case data. Similarly, countries that appeared in the data and were not ranked by the model receive a rank of $i + 1$. For circles that lie on the $x = y$ line (grey solid line) the predicted and reported rankings are equal. Circles that lie between the two dashed lines correspond to countries with a difference in ranking that is less than or equal to five. The circle areas are scaled proportionally to the number of reported cases that were imported from the corresponding country. Spearman’s rank correlation coefficient between the absolute numbers of reported and predicted importations is equal to 0.6. (B) The absolute number of reported dengue importations are plotted against the absolute number of predicted importations.

We obtained dengue case data from Queensland Health, which records the places of acquisition for each dengue case reported in Queensland. We rank the countries of acquisition by the total number of predicted and reported dengue-infected people who arrive in Queensland. We then plot the reported ranking against the predicted ranking. In addition, we plot the absolute number of reported importations against the absolute number of predicted importations and calculate Spearman’s rank correlation coefficient. Fig 5 and Fig S13 show the results.

The rank-based validation of our model demonstrates that overall, the model captures the different importation sources well. It does particularly well for the countries from which Queensland receives the most infections. Spearman’s rank correlation coefficient is equal to 0.6 for the year 2015 and equal to 0.58 for the year 2011. Below we explain some of the differences between the data and the model output.

For the rank-based validation the two largest outliers in both years are Fiji and Taiwan. The predicted ranking for Fiji in 2011 is 2, while the reported ranking is 10. In 2015 we estimate Fiji to be ranked fifth, however no cases were reported in 2015 and hence Fiji is ranked last amongst the reported cases. According to the Fijian government tourists are less likely to contract the disease than local residents as they tend to stay in areas that are not infested by *Aedes aegypti* mosquitoes [65] or where there is likely considerable control effort undertaken by tourism accommodation operators. Since the incidence rates incorporated into our model do not distinguish between different regions of a source country, the model is unable to account for such nuances. In 2011 and 2015 we estimate Taiwan to be ranked seventh and eighth, respectively, however no cases were reported in both years. This result is surprising as dengue occurs year-

round in Taiwan [26] and approximately 44,000 and 16,000 Queensland residents travelled to Taiwan in 2011 and 2015, respectively.

Some of the differences between the observed percentages and the predicted percentages can be explained by under-reporting. It is possible that dengue awareness among travellers to one country is greater than the awareness amongst travellers to another country. Travellers with higher awareness levels are more likely to report to a doctor if feeling unwell after their return.

Country-specific reporting rates

The reporting rate of a disease is defined as the ratio of reported infections to actual infections. Dengue reporting rates vary greatly across space and time and are difficult to determine [10]. The usual approach to estimating country-specific reporting rates is to carry out cohort or capture-recapture studies that can be costly, are time consuming and may be biased [28].

We utilised our model to infer country- and state-specific reporting rates of imported cases by performing a least squares linear regression without intercept.

Table 3 and Table S7 show the estimated yearly and seasonal reporting rates of imported cases for Queensland, Florida, France, Italy and Spain. To distinguish locally acquired and imported cases in Queensland, we use case-based data from Queensland Health where the country of acquisition is recorded. Travel-related dengue cases reported in Europe are published by the European Centre for Disease Prevention and Control (<http://ghdx.healthdata.org/gbd-results-tool>).

Data for Florida is available from the Florida De-

	Dec-Feb	Mar-May	Jun-Aug	Sep-Nov	Yearly
Queensland	32.4	48.9	18.6	22.6	28.6
Spain	14	14	31.7	26.3	23.5
Italy	4.5	6.8	9.2	13.1	9
France	3.8	6.9	9.7	7.1	7.2
Florida	0.9	0.7	1.2	2.7	1.4

Table 3: Yearly and seasonal reporting rates of imported cases in 2015. The table shows the estimated reporting rates of imported cases for Queensland, Spain, Italy, France and Florida. We estimate the reporting rates by using a least squares linear regression without intercept.

partment of Health (<http://www.floridahealth.gov/diseases-and-conditions/mosquito-borne-diseases/surveillance.html>).

The results show that estimated reporting rates of imported cases are highest in Queensland, in particular during autumn. This is expected as dengue awareness campaigns are intensified between November and April [66]. In contrast, Florida has the lowest dengue reporting rate (1.3% in 2011 and 1.4% in 2015). This finding is supported by a previous study which found that awareness levels in Florida are extremely low [67]. The estimated reporting rates for the European countries are also low; however, the model predicts a substantial increase from 2011 to 2015. The question why reporting rates in Queensland are higher is challenging to answer, as we do not have any information about the true number of imported cases. However, Queensland has one of the best dengue prevention programs in the world. According to Queensland Health, other states and countries frequently ask for training and advice regarding surveillance and awareness campaigns.

Model uncertainty

We found that the average coefficient of variation of our importation model is 19.5% across both years. That is, the model’s standard deviation is on average equal to 19.5% of its mean. Fig S14 shows the distribution of the coefficient of variation for several destinations.

The results from the global sensitivity analysis show that t_c is the most important of the three model parameters with a total-order index of 0.94 (see Fig S15). The different values of the first-order and total-order indices indicate interaction between the model parameters. The second-order indices show that there is significant interaction between parameters t_c and $\beta_{c,m}$ with a second-order index of 0.19, as well as between parameters t_c and n with a second-order index of 0.1.

Since the range of parameter t_c is large ([1, 29200] days), we performed the sensitivity analysis again for a shorter range of values ([1, 30] days) that is more realistic for returning residents who spend their holidays in an endemic country. In this case, parameter $\beta_{c,m}$, with a total-order index of 0.6, is more important than t_c , which has a total-order index of 0.35 (see Fig S15). The second-order indices show that there is still significant interaction between parameters t_c and $\beta_{c,m}$ with a second-order index of 0.06, and

between parameters t_c and n with a second-order index of 0.07.

DISCUSSION

To mitigate the risk of outbreaks from importation of dengue into non-endemic regions it is critical to predict the arrival time and location of infected individuals. We modelled the number of dengue infections arriving each month at any given airport, which enabled us to estimate the number of infections that are imported into different countries and states each month. In addition, the model determines the countries of acquisition and hence is able to uncover the routes along which dengue is most likely imported. Our results can also be used to estimate country- and state-specific reporting rates of imported cases.

Such knowledge can inform surveillance, education and risk mitigation campaigns to better target travellers along high risk importation routes at the most appropriate times. It will also help authorities to more efficiently surveil those airports with the highest risk of receiving dengue-infected passengers.

The model proposed here overcomes many of the shortcomings of previous models, however, it is not without limitations. Validation through comparison of reported cases to predicted cases is infeasible due to the high degree of under-reporting. However, we demonstrate that the coefficient of variation of the model with 19.5% on average is low (see Material and Methods). A rank-based validation for Queensland confirmed that the different importation sources are accurately predicted.

Incidence rates may vary considerably from region to region within the same country [65] and higher resolution data could improve the model’s predictions, as it would better reflect the export of dengue cases from the individual regions. Region-specific incidence rates can, for instance, be combined with spatial patterns of the visiting frequency of travellers to determine the likelihood of travellers to export dengue out of endemic countries. Additional data on individuals’ travel behaviour may also be beneficial, as it can be analysed to improve the estimation of the average time that a person has spent in a specific country before arriving at a given airport. Our assumption that returning residents and visitors are exposed to the same daily incidence rates is a simplification. Further details on the types of accom-

modation, for example, resorts vs local housing, could also be used to inform the daily incidence rates, due to variations in vector control. The global sensitivity analysis has revealed that t_c , the number of days a traveller has spent in country c , is the most important model parameter. Hence, additional data on individuals' travel behaviour may substantially improve the model. Knowledge about the exact age of visitors who reside in non-endemic countries would also improve the model. Currently, we assume that the age of a visitor is equal to the median age of the population of the country in which the visitor resides. In reality, the age of air passengers may differ from the median age, especially for developing countries.

In temperate regions local conditions may not allow for dengue to be transmitted during the winter months. Thus, even a large number of imported cases during those months would not trigger local outbreaks. Variable seasonality patterns due to El Niño Southern Oscillation can affect the spread of dengue in tropical and subtropical regions. An interesting direction for future research is to combine the here proposed model with knowledge of local conditions and weather phenomena like El Niño Southern Oscillation to evaluate the risk of local outbreaks. In this work we studied dengue importation via air travel. In future, we will also consider other modes of transportation to develop a more comprehensive model.

DATA AVAILABILITY STATEMENT

The air travel data used in this study are owned by a third party and were licensed for use under contract by International Air Travel Association (IATA)- Passenger Intelligence Services (PaxIS): <http://www.iata.org/services/statistics/intelligence/paxis/Pages/index.aspx>. The same data can be purchased for use by any other researcher by contacting: Phil GENNAOUI Regional Manager - Aviation Solutions (Asia Pacific) Tel: +65 6499 2314 — Mob: +65 9827 0414 gennaoui@iata.org — www.iata.org

Monthly dengue incidence rates for all countries for the years 2011 and 2015 are available as supplementary information files.

Median population age data is publicly available from the Central Intelligence Agency at <https://www.cia.gov/library/publications/resources/the-world-factbook/fields/343.html>.

International tourism arrival data used in this study are owned by a third party and cannot be shared publicly. The data is available for purchase from the World Tourism Organisation at <https://www.eunwto.org/action/doSearch?ConceptID=2445&target=topic>.

Dengue notification data from Australia is publicly available from the Australian Department of Health at http://www9.health.gov.au/cda/source/rpt_1_sel.cfm.

Dengue notification data from Europe is publicly available from the European Centre for Disease Prevention and Con-

trol at <http://atlas.ecdc.europa.eu/public/index.aspx>.

Case-based dengue notifications for Queensland cannot be shared as it contains confidential information. The authors gained access to this data in accordance with Section 284 of the Public Health Act 2005.

ACKNOWLEDGMENTS

We would like to thank Frank de Hoog and Simon Dunstall for their constructive feedback which helped us to improve the model. We would also like to thank Queensland Health for providing dengue outbreak data. This work is part of the DiNeMo project.

REFERENCES

- [1] David E. Bloom, Steven Black, and Rino Rappuoli. Emerging infectious diseases: A proactive approach. *Proc. Natl. Acad. Sci.*, 114(16):4055–4059, 2017.
- [2] Dirk Brockmann and Dirk Helbing. The hidden geometry of complex, network-driven contagion phenomena. *Science*, 342(6164):1337–1342, 2013.
- [3] Ilaria Dorigatti, Arran Hamlet, Ricardo Aguas, Lorenzo Cattarino, Anne Cori, Christl A Donnelly, Tini Garske, Natsuko Imai, and Neil M Ferguson. International risk of yellow fever spread from the ongoing outbreak in Brazil, December 2016 to May 2017. *Eurosurveillance*, 22(28):30572, 2017.
- [4] R. Guimera, S. Mossa, A. Turttschi, and L. A. N. Amaral. The worldwide air transportation network: Anomalous centrality, community structure, and cities' global roles. *Proc. Natl. Acad. Sci.*, 102(22):7794–7799, 2005.
- [5] Zhuojie Huang and Andrew J Tatem. Global malaria connectivity through air travel. *Malar. J.*, 12(1):269, 2013.
- [6] L. Hufnagel, D. Brockmann, and T. Geisel. Forecast and control of epidemics in a globalized world. *Proc. Natl. Acad. Sci.*, 101(42):15124–15129, 2004.
- [7] A. J. Tatem, Z. Huang, A. Das, Q. Qi, J. Roth, and Y. Qiu. Air travel and vector-borne disease movement. *Parasitology*, 139(14):1816–1830, 2012.
- [8] Isaac I Bogoch, Oliver J Brady, Moritz U G Kraemer, Matthew German, Marisa I Creatore, Manisha A Kulkarni, John S Brownstein, Sumiko R Mekaru, Simon I Hay, Emily Groot, Alexander Watts, and Kamran Khan. Anticipating the international spread of Zika virus from Brazil. *Lancet*, 387(10016):335–336, 2016.
- [9] Samir Bhatt, Peter W. Gething, Oliver J. Brady, Jane P. Messina, Andrew W. Farlow, Catherine L. Moyes, John M. Drake, John S. Brownstein, Anne G.

- Hoen, Osman Sankoh, Monica F. Myers, Dylan B. George, Thomas Jaenisch, G. R. William Wint, Cameron P. Simmons, Thomas W. Scott, Jeremy J. Farrar, and Simon I. Hay. The global distribution and burden of dengue. *Nature*, 496(7446):504–507, 2013.
- [10] Jeffrey D Stanaway, Donald S Shepard, Eduardo A Undurraga, Yara A Halasa, Luc E Coffeng, Oliver J Brady, Simon I Hay, Neeraj Bedi, Isabela M Bensenor, Carlos A Castañeda-Orjuela, Ting-Wu Chuang, Katherine B Gibney, Ziad A Memish, Anwar Rafay, Kingsley N Ukwaja, Naohiro Yonemoto, and Christopher J L Murray. The global burden of dengue: An analysis from the Global Burden of Disease Study 2013. *Lancet Infect. Dis.*, 16(6):712–723, 2016.
- [11] Natasha Evelyn Anne Murray, Mikkel Quam, and Annelies Wilder-Smith. Epidemiology of dengue: Past, present and future prospects. *Clin. Epidemiol.*, 5:299–309, 2013.
- [12] World Health Organization. WHO heralds new phase in the fight against neglected tropical diseases, 2013. Available at: http://www.who.int/mediacentre/news/releases/2013/ntds_report_20130116/en/ [Accessed 25/10/18].
- [13] Lauren Gardner and Sahotra Sarkar. A global airport-based risk model for the spread of dengue infection via the air transport network. *PLoS One*, 8(8):e72129, 2013.
- [14] John S Mackenzie, Duane J Gubler, and Lyle R Petersen. Emerging flaviviruses: The spread and resurgence of Japanese encephalitis, West Nile and dengue viruses. *Nat. Med.*, 10(12):S98–S109, 2004.
- [15] Cameron P. Simmons, Jeremy J. Farrar, Nguyen van Vinh Chau, and Bridget Wills. Dengue. *N. Engl. J. Med.*, 366(15):1423–1432, 2012.
- [16] Hsu P-S Chen C-D Lian I-B Chao D-Y Chang F-S, Tseng Y-T. Re-assess vector indices threshold as an early warning tool for predicting dengue epidemic in a dengue non-endemic country. *PLOS Negl Trop Dis.*, 9(9):e0004043, 2015.
- [17] Jan C. Semenza, Bertrand Sudre, Jennifer Miniot, Massimiliano Rossi, Wei Hu, David Kossowsky, Jonathan E. Suk, Wim Van Bortel, and Kamran Khan. International dispersal of dengue through air travel: Importation risk for Europe. *PLoS Negl. Trop. Dis.*, 8(12):e3278, 2014.
- [18] Lauren M. Gardner and Sahotra Sarkar. Risk of dengue spread from the Philippines through international air travel. *Transp. Res. Rec. J. Transp. Res. Board*, 2501:25–30, 2015.
- [19] A. Wilder-Smith, M. Quam, O. Sessions, J. Rocklöv, J. Liu-Helmersson, L. Franco, and K. Khan. The 2012 dengue outbreak in Madeira: Exploring the origins. *Eurosurveillance*, 19(8), 2014.
- [20] Mikkel B. Quam, Kamran Khan, Jennifer Sears, Wei Hu, Joacim Rocklöv, and Annelies Wilder-Smith. Estimating air travel-associated importations of dengue virus into Italy. *J. Travel Med.*, 22(3):186–193, 2015.
- [21] Eduardo Massad, Marcos Amaku, Francisco Antonio Bezerra Coutinho, Claudio José Struchiner, Marcelo Nascimento Burattini, Kamran Khan, Jing Liu-Helmersson, Joacim Rocklöv, Moritz U. G. Kraemer, and Annelies Wilder-Smith. Estimating the probability of dengue virus introduction and secondary autochthonous cases in Europe. *Sci. Rep.*, 8(1):4629, 2018.
- [22] Katherine Standish, Guillermina Kuan, William Avilés, Angel Balmaseda, and Eva Harris. High dengue case capture rate in four years of a cohort study in Nicaragua compared to national surveillance data. *PLoS Negl. Trop. Dis.*, 4(3):e633, 2010.
- [23] M. Kakkar. Dengue fever is massively under-reported in India, hampering our response. *BMJ*, 345(17):e8574, 2012.
- [24] S. Vong, S. Goyet, S. Ly, C. Ngan, R. Huy, V. Duong, O. Wichmann, G. W. Letson, H. S. Margolis, and P. Buchy. Under-recognition and reporting of dengue in Cambodia: A capture-recapture analysis of the National Dengue Surveillance System. *Epidemiol. Infect.*, 140(03):491–499, 2012.
- [25] T. Y. M. Wahyono, J. Nealon, S. Beucher, A. Prayitno, A. Moureau, S. Nawawi, H. Thabrany, and M. Nadjib. Indonesian dengue burden estimates: Review of evidence by an expert panel. *Epidemiol. Infect.*, 145(11):2324–2329, 2017.
- [26] International Association for Medical Assistance to Travellers (IAMAT). International Association for Medical Assistance to Travellers, 2018. Available at: <https://www.iamat.org/> [Accessed 6/11/18].
- [27] Global Burden of Disease Collaborative Network. Global Burden of Disease Study 2016 (GBD 2016) Results, 2017. Available at: <http://ghdx.healthdata.org/gbd-results-tool> [Accessed 9/11/18].
- [28] Nguyen T. Toan, Stefania Rossi, Gabriella Prisco, Nicola Nante, and Simonetta Viviani. Dengue epidemiology in selected endemic countries: Factors influencing expansion factors as estimates of underreporting. *Trop. Med. Int. Heal.*, 20(7):840–863, 2015.

- [29] Airline Network News and Analysis. Aviation database, 2018. Available at: <https://www.anna.aero/databases/> [Accessed 1/12/18].
- [30] Civil Aviation Administration of China. 2015 Civil Aviation Airport Throughput Ranking, 2016. Available at: http://www.caac.gov.cn/XXGK/XXGK/TJSJ/201603/t20160331_30105.html [Accessed 1/12/18].
- [31] Ministry of Land Infrastructure Transport and Tourism. Ministry of Land, Infrastructure, Transport and Tourism,, 2016. Available at: <https://web.archive.org/web/20161021205147/http://www.mlit.go.jp/common/001141840.pdf> [Accessed 1/12/18].
- [32] Ministry of Transportation Republic of Indonesia. Air transport traffic, 2018. Available at: <http://hubud.dephub.go.id/?en/llu/index/filter:bulan,0> [Accessed 1/12/18].
- [33] Airports Council International. North America airport traffic annual reports, 2015. Available at: <https://airportscouncil.org/intelligence/north-american-airport-traffic-reports/north-america-airport-traffic-annual-reports/> [Accessed 1/12/18].
- [34] Directorate General of Civil Aviation - Kuwait. Statistics, 2015. Available at: <https://www.dgca.gov.kw/en/civil-aviation/media-and-info/statistics> [Accessed 1/12/18].
- [35] EANA - Navegación Aérea Argentina. Statistical reports, 2016. Available at: <https://www.eana.com.ar/estadisticas#node-150> [Accessed 1/12/18].
- [36] Angkasa Pura II. Innovating beyond excellence - 2015 annual report, 2015. Available at: https://cms.angkasapura2.co.id/NUWEB.PUBLIC.FILES/angkasapura2/Annual_25_07_2016_09_04_23.pdf [Accessed 1/12/18].
- [37] Civil Aviation Authority of the Philippines. Passenger movement, 2015. Available at: <https://data.gov.ph/dataset/civil-aviation-authority-philippines-passenger-movement> [Accessed 1/12/18].
- [38] Instituto de Estadísticas de Puerto Rico. Carga y pasajeros aéreos y marítimos, 2015. Available at: http://www.estadisticas.gobierno.pr/iepr/Estadisticas/InventariodeEstad%C3%ADsticas/tabid/186/ctl/view_detail/mid/775/report_id/9485fbd8-efa5-4583-a8a4-a39c74eb846f/Default.aspx [Accessed 1/12/18].
- [39] Malaysia Airports. Connecting you seamlessly to the world - Annual report 2015, 2015. Available at: <http://mahb.listedcompany.com/misc/ar/ar2015.pdf> [Accessed 1/12/18].
- [40] Zambia: Transport Data Portal. Air transport statistics Zambia, 2016. Available at: <http://zambiamtc.opendataforafrica.org/ZMATS2015/air-transport-statistics-zambia?airports=1000000-kenneth-kaunda-international-airport> [Accessed 1/12/18].
- [41] California Department of Transportation. 2105:2014 air passenger and air cargo traffic activity report, 2016. Available at: <http://www.dot.ca.gov/hq/planning/aeronaut/documents/statistics/15trafficgrowthrpt.pdf> [Accessed 1/12/18].
- [42] Airports Authority of India. Passengers, 2016. Available at: <https://www.aai.aero/sites/default/files/traffic-news/Mar2k15annex3.pdf> [Accessed 1/12/18].
- [43] Associazione Italiana Gestori Aeroporti. Statistiche, 2016. Available at: <http://www.assaeroporti.com/statistiche/> [Accessed 1/12/18].
- [44] Office National des Aéroports. Passengers, 2017. Available at: [http://www.onda.ma/en/I-am-a-Professional/Companies/statistics/\(offset\)/30](http://www.onda.ma/en/I-am-a-Professional/Companies/statistics/(offset)/30) [Accessed 1/12/18].
- [45] Airports Company South Africa. Statistics, 2018. Available at: <http://www.airports.co.za/news/statistics> [Accessed 1/12/18].
- [46] Egyptian Holding Company for Airports and Air Navigation. Statistics, 2015. Available at: <http://www.ehcaan.com/statistics.aspx> [Accessed 1/12/18].
- [47] Airports Council International. Passenger Summary, 2018. Available at: <https://aci.aero/data-centre/annual-traffic-data/passengers/2015-final-summary/> [Accessed 1/12/18].
- [48] Department of Home Affairs. Overseas Arrivals and Departures, 2017. Available at: <https://data.gov.au/dataset/overseas-arrivals-and-departures> [Accessed 22/11/18].
- [49] Miranda Chan and Michael A. Johansson. The incubation periods of dengue viruses. *PLoS One*, 7(11):e50972, 2012.
- [50] G. Chowell, P. Diaz-Dueñas, J.C. Miller, A. Alcazar-Velazco, J.M. Hyman, P.W. Fenimore, and C. Castillo-Chavez. Estimation of the reproduction number of dengue fever from spatial epidemic data. *Math. Biosci.*, 208(2):571–589, 2007.
- [51] Duane J. Gubler and Gary G. Clark. Dengue/dengue hemorrhagic fever: The emergence of a global health problem. *Emerg. Infect. Dis.*, 1(2):55–57, 1995.

- [52] McCann D Messenger JC, Lee S. *Working time around the world: Trends in working hours, laws, and policies in a global comparative perspective*. 1st ed. London: Routledge, 2007.
- [53] Central Intelligence Agency. The World Factbook, 2017. Available at: <https://www.cia.gov/library/publications/the-world-factbook/fields/2177.html{#}aq> [Accessed 12/11/18].
- [54] World Tourism Organisation. All Countries: Inbound Tourism: Arrivals by mode of transport 1995 - 2017, 2018. Available at: <https://www.e-unwto.org/doi/abs/10.5555/unwtotfb0000271519952017201809> [Accessed 12/11/18].
- [55] Sobol IM. Global sensitivity indices for nonlinear mathematical models and their Monte Carlo estimates. *Math Comput Simulat.*, 55:271–280, 2001.
- [56] Usher W Herman J. SALib: An open-source Python library for sensitivity analysis. *The Journal of Open Source Software*, 2(9):97, 2017.
- [57] Amy C Morrison, Emily Zielinski-Gutierrez, Thomas W Scott, and Ronald Rosenberg. Defining challenges and proposing solutions for control of the virus vector *Aedes aegypti*. *PLoS Med.*, 5(3):e68, 2008.
- [58] Nigel W. Beebe, Robert D. Cooper, Pipi Mottram, and Anthony W. Sweeney. Australia’s dengue risk driven by human adaptation to climate change. *PLoS Negl. Trop. Dis.*, 3(5):e429, 2009.
- [59] Micah B. Hahn, Rebecca J. Eisen, Lars Eisen, Karen A. Boegler, Chester G. Moore, Janet McAllister, Harry M. Savage, and John-Paul Mutebi. Reported Distribution of *Aedes (Stegomyia) aegypti* and *Aedes (Stegomyia) albopictus* in the United States, 1995-2016 (Diptera: Culicidae). *J. Med. Entomol.*, 53(5):1169–1175, 2016.
- [60] Caitlin A.M. Van Dodewaard and Stephanie L. Richards. Trends in dengue cases imported into the United States from Pan America 2001–2012. *Environ. Health Insights*, 9:33–40, 2015.
- [61] Amarakoon, Dharmaratne, Anthony Chen, Sam Rawlins, Dave D. Chadee, Michael Taylor, and Roxann Stennett. Dengue epidemics in the Caribbean—temperature indices to gauge the potential for onset of dengue. *Mitig. Adapt. Strateg. Glob. Chang.*, 13(4):341–357, 2008.
- [62] Victor Vasquez, Elie Haddad, Alice Perignon, Stéphane Jaureguiberry, Ségolène Brichler, Isabelle Leparco-Goffart, and Eric Caumes. Dengue, chikungunya, and Zika virus infections imported to Paris between 2009 and 2016: Characteristics and correlation with outbreaks in the French overseas territories of Guadeloupe and Martinique. *Int. J. Infect. Dis.*, 72:34–39, 2018.
- [63] Guy La Ruche, Dominique Dejour-Salamanca, Pascale Bernillon, Isabelle Leparco-Goffart, Martine Ledrans, Alexis Armengaud, Monique Debruyne, Gérard-Antoine Denoyel, and Ségolène Brichler. Capturere-capture method for estimating annual incidence of imported dengue, France, 2007–2010. *Emerg. Infect. Dis.*, 19(11):1740–1748, 2013.
- [64] David Warrilow, Judith A. Northill, and Alyssa T. Pyke. Sources of dengue viruses imported into Queensland, Australia, 2002–2010. *Emerg. Infect. Dis.*, 18(11):1850–1857, 2012.
- [65] The Fijian Government. Minimal risk of dengue fever in major tourism areas of Fiji, 2014. Available at: <http://www.fiji.gov.fj/Media-Center/Press-Releases/MINIMAL-RISK-OF-DENGUE-FEVER-IN-MAJOR-TOURISM-AREA.aspx> [Accessed 8/6/18].
- [66] Queensland Health. Queensland dengue management plan 2015–2020, 2015. Available at: https://www.health.qld.gov.au/_data/assets/pdf_file/0022/444433/dengue-mgt-plan.pdf [Accessed 29/10/18].
- [67] Mary H. Hayden, Jamie L. Cavanaugh, Christopher Tittel, Melinda Butterworth, Steven Haenchen, Katherine Dickinson, Andrew J. Monaghan, and Kacey C. Ernst. Post outbreak review: Dengue preparedness and response in Key West, Florida. *Am. J. Trop. Med. Hyg.*, 93(2):397–400, 2015.
- [68] Centers for Disease Control and Prevention. Dengue map, 2019. Available at: <https://www.healthmap.org/dengue/en/> [Accessed 31/07/19].

SUPPORTING INFORMATION

Table S1: List of airport abbreviations

IATA 3-Letter Code	Name	City (Country/State)	IATA 3-Letter Code	Name	City (Country/State)
AEP	Jorge Newbery Airport	Buenos Aires (Argentina)	LAX	Los Angeles International Airport	Los Angeles (California)
BKK	Suvarnabhumi Airport	Bangkok (Thailand)	LHR	Heathrow Airport	London (UK)
BNE	Brisbane Airport	Brisbane (Queensland)	MAD	Adolfo Suárez Madrid-Barajas Airport	Madrid (Spain)
BOM	Chhatrapati Shivaji International Airport	Mumbai (India)	MCO	Orlando International Airport	Orlando (Florida)
CDG	Charles de Gaulle Airport	Paris (France)	MEX	Mexico City International Airport	Mexico City (Mexico)
COK	Cochin International Airport	Kochi (India)	MIA	Miami International Airport	Miami (Florida)
CUN	Cancún International Airport	Cancún (Mexico)	MNL	Ninoy Aquino International Airport	Manila (Philippines)
DEL	Indira Gandhi International Airport	New Delhi (India)	MTY	Monterrey International Airport	Apodaca (Mexico)
DFW	Dallas/Fort Worth International Airport	Dallas (Texas)	MLX	Milan Malpensa Airport	Milan (Italy)
DPS	Ngurah Rai International Airport	Denpasar (Indonesia)	NRT	Narita International Airport	Tokyo (Japan)
DXB	Dubai International Airport	Dubai (UAE)	ORY	Paris Orly Airport	Paris (France)
EZE	Ministro Pistarini International Airport	Buenos Aires (Argentina)	PER	Perth Airport	Perth (Western Australia)
FDF	Martinique Aimé Césaire International Airport	Fort-de-France (Martinique)	PTP	Pointe-à-Pitre International Airport	Pointe-à-Pitre (Guadeloupe)
FLL	Fort Lauderdale-Hollywood International Airport	Miami (Florida)	PUJ	Punta Cana International Airport	Punta Cana (Dominican Republic)
GDL	Miguel Hidalgo y Costilla Guadalajara International Airport	Guadalajara (Mexico)	SAL	Monseñor Óscar Arnulfo Romero International Airport	San Salvador (El Salvador)
GRU	São Paulo International Airport	São Paulo (Brazil)	SDQ	Las Américas International Airport	Punta Caucedo (Dominican Republic)
ICN	Incheon International Airport	Seoul (South Korea)	SFO	San Francisco International Airport	San Francisco (California)
IAH	George Bush Intercontinental Airport	Houston (Texas)	SJU	Luis Muñoz Marín International Airport	San Juan (Puerto Rico)
JFK	John F. Kennedy International Airport	New York City (New York)	STI	Cibao International Airport	Santiago de los Caballeros (Dominican Republic)
KBL	Hamid Karzai International Airport	Kabul (Afghanistan)	TPE	Taiwan Taoyuan International Airport	Taipei (Taiwan)

Table S2: List of non-endemic countries where IATA data is inaccurate

Algeria	Bahrain	Bonaire, Saint Eustatius & Saba	Bulgaria
Central African Republic	Croatia	Egypt	Federated States of Micronesia
Finland	Germany	Guinea-Bissau	Greece
Hungary	Iceland	Iran	Israel
Malawi	Morocco	Netherlands	Russian Federation
Serbia	Slovenia	South Africa	South Korea
Tanzania	Togo	The Gambia	Tunisia
Turkey	Uganda	Ukraine	Zambia

Table S3: List of countries indicating whether dengue vectors are present and whether the country is endemic. Information about endemicity was obtained from [68]. Information about vector presence was obtained from [26].

Country	Endemic	Vector presence	Country	Endemic	Vector presence	Country	Endemic	Vector presence
Afghanistan	no	yes	Ghana	no	yes	Pakistan	yes	yes
Albania	no	yes	Gibraltar	no	yes	Palau	yes	yes
Algeria	no	yes	Greece	no	yes	Palestine	no	yes
American Samoa	no	yes	Greenland	no	no	Panama	yes	yes
Andorra	no	no	Grenada	yes	yes	Papua New Guinea	yes	yes
Angola	yes	yes	Guadeloupe	yes	yes	Paraguay	yes	yes
Anguilla	yes	yes	Guam	no	no	Peru	yes	yes
Antarctica	no	no	Guatemala	yes	yes	Philippines	yes	yes
Antigua and Barbuda	yes	yes	Guinea	yes	yes	Poland	no	no
Argentina	yes	yes	Guinea-Bissau	no	yes	Portugal	no	yes
Armenia	no	yes	Guyana	yes	yes	Puerto Rico	yes	yes
Aruba	yes	yes	Haiti	yes	yes	Qatar	no	no
Australia	no	yes	Honduras	yes	yes	Reunion	yes	yes
Austria	no	yes	Hong Kong	yes	yes	Romania	no	yes
Azerbaijan	no	no	Hungary	no	yes	Russian Federation	no	yes
Bahrain	no	no	Iceland	no	no	Rwanda	no	yes
Bangladesh	yes	yes	India	yes	yes	Saint Helena	no	no
Barbados	yes	yes	Indonesia	yes	yes	Saint Kitts and Nevis	yes	yes
Belarus	no	no	Inner Hebrides	no	no	Saint Lucia	yes	yes
Belgium	no	yes	Iran	no	no	Saint Pierre and Miquelon	no	no
Belize	yes	yes	Iraq	no	no	Saint Vincent and the Grenadines	yes	yes
Benin	no	yes	Ireland	no	no	Samoa	no	yes
Bermuda	no	yes	Israel	no	yes	Sao Tome and Principe	no	no
Bhutan	yes	yes	Italy	no	yes	Saudi Arabia	yes	yes
Bolivia	yes	yes	Jamaica	yes	yes	Senegal	yes	yes
Bonaire, Saint Eustatius & Saba	no	yes	Japan	no	no	Serbia	no	yes
Bosnia and Herzegovina	no	yes	Jordan	no	yes	Seychelles	yes	yes
Botswana	no	no	Kazakhstan	no	no	Sierra Leone	yes	yes
Brazil	yes	yes	Kenya	yes	yes	Singapore	yes	yes
Brunei	yes	yes	Kiribati	no	yes	Sint Maarten	no	yes
Bulgaria	no	yes	Kuwait	no	no	Slovakia	no	yes
Burkina Faso	yes	yes	Kyrgyzstan	no	no	Slovenia	no	yes
Burundi	no	yes	Laos	yes	yes	Solomon Islands	yes	yes
Cambodia	yes	yes	Latvia	no	no	Somalia	yes	yes
Cameroon	yes	yes	Lebanon	no	yes	South Africa	no	no
Canada	no	no	Lesotho	no	no	South Korea	no	no
Cape Verde	yes	yes	Liberia	no	yes	South Sudan	yes	yes
Cayman Islands	yes	yes	Libya	no	no	Spain	no	yes
Central African Republic	no	yes	Liechtenstein	no	no	Sri Lanka	yes	yes
Chad	no	yes	Lithuania	no	no	Sudan	yes	yes
Channel Islands	no	no	Luxembourg	no	no	Suriname	yes	yes
Chile	no	no	Macau	yes	yes	Swaziland	no	no
China	yes	yes	Macedonia	no	yes	Sweden	no	no
Christmas Island	no	no	Madagascar	yes	yes	Switzerland	no	yes
Cocos (Keeling) Islands	no	no	Malawi	no	yes	Syria	no	yes
Colombia	yes	yes	Malaysia	yes	yes	Taiwan	yes	yes
Comoros	yes	yes	Maldives	no	yes	Tajikistan	no	no

Country	Endemic	Vector presence	Country	Endemic	Vector presence	Country	Endemic	Vector presence
Congo	no	yes	Mali	yes	yes	Tanzania	no	yes
Cook Islands	no	yes	Malta	no	yes	Thailand	yes	yes
Costa Rica	yes	yes	Marshall Islands	no	yes	The Bahamas	yes	yes
Cote d'Ivoire	yes	yes	Martinique	yes	yes	The Gambia	no	yes
Croatia	no	yes	Mauritania	no	yes	Timor-Leste	yes	yes
Cuba	yes	yes	Mauritius	yes	yes	Togo	no	yes
Curacao	no	yes	Mayotte	yes	yes	Tonga	no	yes
Cyprus	no	no	Mexico	yes	yes	Trinidad and Tobago	yes	yes
Czech Republic	no	yes	Moldova	no	no	Tunisia	no	no
Democratic Republic of the Congo	yes	yes	Monaco	no	yes	Turkey	no	yes
Denmark	no	no	Mongolia	no	no	Turkmenistan	no	no
Djibouti	yes	yes	Montenegro	no	yes	Turks and Caicos Islands	yes	yes
Dominica	yes	yes	Montserrat	yes	yes	Tuvalu	no	yes
Dominican Republic	yes	yes	Morocco	no	no	Uganda	no	yes
Ecuador	yes	yes	Mozambique	yes	yes	Ukraine	no	no
Egypt	no	yes	Myanmar	yes	yes	United Arab Emirates	yes	yes
El Salvador	yes	yes	Namibia	no	yes	United Kingdom	no	no
Equatorial Guinea	yes	yes	Nauru	no	yes	United States	no	yes
Eritrea	yes	yes	Nepal	yes	yes	United States Minor Outlying Islands	no	no
Estonia	no	no	Netherlands	no	yes	Uruguay	no	no
Ethiopia	yes	yes	New Caledonia	yes	yes	Uzbekistan	no	no
Falkland Islands	no	no	New Zealand	no	no	Vanuatu	yes	yes
Federated States of Micronesia	no	yes	Nicaragua	yes	yes	Venezuela	yes	yes
Fiji	no	yes	Niger	no	yes	Vietnam	yes	yes
Finland	no	no	Nigeria	yes	yes	Virgin Islands	yes	yes
France	no	yes	Niue	no	yes	Wallis and Futuna Islands	no	yes
French Guiana	yes	yes	Norfolk Island	no	no	Western Sahara	no	no
French Polynesia	no	yes	North Korea	no	no	Yemen	yes	yes
Gabon	yes	yes	Northern Mariana Islands	no	yes	Zambia	no	yes
Georgia	no	yes	Norway	no	no	Zimbabwe	no	yes
Germany	no	yes	Oman	yes	yes			

Table S4: Annual estimated imported dengue cases per airport

Code	Imported cases 2011	Imported cases 2015	Name	City	Country/State
MIA	2413	2547	Miami International	Miami	US/Florida
LAX	1518	1871	Los Angeles Intl	Los Angeles	US/California
CDG	941	1227	Charles De Gaulle	Paris-De Gaulle	France/Île-de-France
SFO	831	1166	San Francisco Intl	San Francisco	US/California
MCO	822	1036	Orlando Intl	Orlando	US/Florida
FLL	792	970	Ft Lauderdale Intl	Fort Lauderdale	US/Florida
ORY	652	788	Orly	Paris-Orly	France/Île-de-France
IAH	599	810	George Bush Intercontinental	Houston-Intercontinental	US/Texas
EZE	586	648	Ministro Pistarini	Buenos Aires	Argentina/Autonomous City of Buenos Aires
MAD	385	439	Adolfo Suarez-Barajas	Madrid	Spain/Community of Madrid
DFW	381	478	Dallas/Ft Worth Intl	Dallas/Fort Worth	US/Texas
BNE	379	529	Brisbane Intl	Brisbane	Australia/Queensland
MXP	356	408	Malpensa	Milan-Malpensa	Italy/Lombardy
FCO	343	424	Fiumicino	Rome-Da Vinci	Italy/Lazio
AEP	258	225	Jorge Newbery	Buenos Aires-Newbery	Argentina/Autonomous City of Buenos Aires
MLE	204	214	Ibrahim Nasir International	Male	Maldives/Malé
POS	198	150	Piarco International	Port of Spain	Trinidad and Tobago/Port of Spain
TPA	160	224	Tampa International	Tampa	US/Florida
SXM	159	184	Prinses Juliana International	St. Maarten	Sint Maarten
BCN	148	207	Barcelona	Barcelona	Spain/Catalonia
CUR	147	173	Hato International	Curacao	Curacao
SAN	142	181	San Diego International Airport	San Diego	US/California
HNL	123	153	Honolulu Intl	Honolulu/Oahu	US/Hawaii
BEY	122	162	Rafic Hariri International	Beirut	Lebanon/Beirut
KBL	109	155	Kabul International	Kabul	Afghanistan/Kabul
ACC	97	126	Kotoka International	Accra	Ghana/Greater Accra
SJC	88	99	San Jose Municipal	San Jose	US/California
SAT	80	127	San Antonio Intl	San Antonio	US/Texas
VCE	75	108	Marco Polo	Venice	Italy/Veneto
AUS	71	123	Austin-Bergstrom International Airport	Austin	US/Texas
SMF	70	94	Sacramento International	Sacramento	US/California
OAK	65	65	Metro Oakland Intl	Oakland	US/California
LYS	56	72	Satolas	Lyon	France/Auvergne-Rhône-Alpes
OOL	55	126	Coolangatta	Gold Coast	Australia/Queensland
JAX	53	74	Jacksonville Intl	Jacksonville	US/Florida
MRS	53	69	Marignane	Marseille	France/Provence-Alpes-Côte d'Azur
NCE	53	71	Cote D'Azur	Nice	France/Provence-Alpes-Côte d'Azur
BLQ	49	59	Guglielmo Marconi	Bologna	Italy/Emilia-Romagna
COR	46	78	Pajas Blancas	Córdoba	Argentina/Córdoba
TLS	42	53	Blagnac	Toulouse	France/Occitanie
COO	39	66	Cadjehoun	Cotonou	Benin/Littoral
ONT	39	43	Ontario Intl	Ontario	US/California
SAH	38	11	Sana'a International	Sana'a	Yemen/Sana'a
LIN	38	56	Linate	Milan-Linate	Italy/Lombardy
FAT	37	42	Fresno Yosemite International	Fresno	US/California
SNA	37	58	John Wayne Airport	Orange County	US/California
DLA	33	60	Douala International	Douala	Cameroon/Littoral
KGL	29	48	Kigali International Airport	Kigali	Rwanda/Kigali
PNS	27	31	Pensacola International	Pensacola	US/Florida
BOD	24	33	Merignac	Bordeaux	France/Nouvelle-Aquitaine
CAY	24	38	Felix Eboue	Cayenne	France/French Guiana
NAN	23	27	Nadi International	Nadi	Fiji/Ba
PBM	22	34	Johan A. Pengel Intl	Paramaribo	Suriname/Paramaribo
CNS	22	37	Cairns International	Cairns	Australia/Queensland
ELP	22	25	El Paso Intl	El Paso	US/Texas
NTE	21	23	Chateau Bougon	Nantes	France/Pays de la Loire
FLR	21	26	Peretola	Florence	Italy/Tuscany
BUR	18	9	Hollywood-Burbank	Burbank	US/California
HOU	18	38	William P Hobby	Houston-Hobby	US/Texas

Code	Imported cases 2011	Imported cases 2015	Name	City	Country/State
PPG	17	18	Pago Pago Intl	Pago Pago	American Samoa/Maoputasi County
TRN	17	20	Citta Di Torino	Turin	Italy/Piedmont
ROB	17	17	Roberts Intl	Monrovia-Roberts	Liberia
MFE	15	14	Miller International	McAllen	US/Texas
PBI	15	23	Palm Beach Intl	West Palm Beach	US/Florida
LGB	15	13	Long Beach Municipal	Long Beach	US/California
BZV	15	29	Maya Maya	Brazzaville	Congo/Brazzaville
NAP	14	16	Capodichino	Naples	Italy/Campania
TLH	14	21	Tallahassee International	Tallahassee	US/Florida
MDZ	13	20	El Plumerillo	Mendoza	Argentina/Mendoza
ROS	13	25	Islas Malvinas	Rosario	Argentina/Santa Fe
CRP	13	15	Corpus Christi Intl	Corpus Christi	US/Texas
RSW	13	14	Southwest Florida International	Fort Myers	US/Florida
PMI	13	15	Palma De Mallorca	Palma de Mallorca	Spain/Balearic Islands
MPL	13	18	Mediterranee	Montpellier	France/Occitanie
VPS	12	14	Destin-Ft Walton Beach Airport	Destin-Ft Walton Beach	US/Florida
VRN	12	8	Verona	Verona	Italy/Veneto
FGI	11	20	Fagali'I	Apia	Samoa/Tuamasaga
GNV	11	16	J R Alison Regional Municipal	Gainesville	US/Florida
OGG	11	13	Kahului	Kahului/Maui	US/Hawaii
VLC	11	16	Valencia Airport	Valencia	Spain/Valencian Community
CTA	11	11	Fontanarossa	Catania	Italy/Sicily
BJM	10	17	Bujumbura Intl	Bujumbura	Burundi/Bujumbura Mairie
TAB	10	8	ANR Robinson International	Tobago	Trinidad and Tobago
AGP	10	15	Malaga Airport	Malaga	Spain/Andalusia
ADE	10	3	Aden International	Aden	Yemen/Aden
HRE	10	22	Harare International	Harare	Zimbabwe/Harare
PNR	9	19	Pointe Noire	Pointe Noire	Congo
PUF	9	10	Uzein	Pau	France/Nouvelle-Aquitaine
BIO	9	15	Bilbao Airport	Bilbao	Spain/Basque Autonomous Community
GOA	9	10	Cristoforo Colombo	Genoa	Italy/Liguria
LPA	9	14	Gran Canaria	Gran Canaria	Spain/Canary Islands
GRK	9	11	Regional/R.Gray AAF	Killeen/Fort Hood	US/Texas
WDH	9	10	Windhoek Intl	Windhoek	Namibia/Khomas
MAF	9	13	Midland-Odessa Regl	Midland/Odessa	US/Texas
TRW	9	11	Bonriki International	Tarawa	Kiribati
TSV	9	12	Townsville International	Townsville	Australia/Queensland
PSP	9	11	Palm Springs Muni	Palm Springs	US/California
MLH	9	10	Euroairport	Mulhouse/Basel	France/Grand Est
BES	8	11	Bretagne	Brest	France/Brittany
NKC	8	13	Nouakchott	Nouakchott	Mauritania/Nouakchott
BRC	8	12	San Carlos Bariloche International	San Carlos Bariloche	Argentina/Río Negro
LBB	8	10	Preston Smith Intl	Lubbock	US/Texas
AMA	8	10	Rick Husband Intl	Amarillo	US/Texas
EYW	7	12	Key West Intl	Key West	US/Florida
NDJ	7	16	Ndjamena	N'Djamena	Chad/N'Djamena
DAL	7	22	Dallas Love Field	Dallas-Love	US/Texas
ECP	5	11	Northwest Florida Beaches International Airport	Panama City	US/Florida
NIM	5	12	Diori Hamani International Airport	Niamey	Niger/Niamey
SPN	3	11	Saipan International	Saipan	Northern Mariana Islands
SFB	0	10	Sanford International	Orlando-Sanford	US/Florida

Table S5: The ten routes with the highest predicted number of dengue-infected passengers with final destinations in non-endemic countries with vector presence.

Orig.	Dest.	Pax	Month
SJU (Puerto Rico)	MCO (Florida)	52	Jul
FDF (Martinique)	ORY (France)	34	Aug
CUN (Mexico)	MIA (Florida)	32	Aug
SDQ (Dominican Republic)	MIA (Florida)	30	Aug
CCS (Venezuela)	MIA (Florida)	28	Aug
GDL (Mexico)	LAX (California)	27	Aug
SJU (Puerto Rico)	FLL (Florida)	25	Jul
PUJ (Dominican Republic)	MIA (Florida)	24	Jul
MNL (Philippines)	LAX (California)	23	Jul
SJU (Puerto Rico)	MIA (Florida)	23	Jul

Table S6: The ten routes with the highest predicted number of dengue-infected passengers who continue to travel to non-endemic regions. The table lists the direct routes with the highest predicted volume of dengue-infected passengers who continue to travel to non-endemic regions irrespective of vector presence. The last column records the month during which the highest number of infected passengers are predicted.

2011				2015			
Origin	Destination	Pax	Month	Origin	Destination	Pax	Month
BOM	DXB	108	Jul	BOM	DXB	142	Aug
CUN	MEX	86	Aug	DEL	DXB	97	Aug
DPS	PER	77	Jan	CUN	MEX	75	Aug
SDQ	JFK	76	Aug	COK	DXB	72	Aug
STI	JFK	76	Aug	DPS	PER	65	Jan
DEL	DXB	72	Jul	MAA	DXB	59	Aug
MNL	ICN	71	Aug	MNL	ICN	95	Aug
DEL	LHR	65	Aug	HYD	DXB	55	Aug
MNL	NRT	65	Jul	SJU	JFK	57	Aug
MTY	MEX	62	Sep	DEL	LHR	59	Aug

Table S7: Yearly and seasonal reporting rates of imported cases in 2011.

	Dec-Feb	Mar-May	Jun-Aug	Sep-Nov	Yearly
Queensland	38.4	25.2	13.7	18.9	24.3
Italy	5.7	4	1.9	7.3	4.4
France	2.2	3.1	4.8	1.3	3
Florida	1.2	0.5	1	2.5	1.3




Figure S1: Illustration of the possibility of recovery before return. If an individual gets infected with dengue while overseas, but recovers before returning to region r , the individual cannot infect other people in region r .




Figure S2: Predicted monthly dengue importations by airport for 2011. The number of predicted imported dengue infections for the top ten airports in non-endemic countries/states with vector presence for each month in 2011. A break in a line indicates that the corresponding airport was not amongst the top ten during the respective month. Airports are abbreviated using the corresponding IATA code. A full list of abbreviations can be found in the supplementary material (see Table S1)


A**B**

Figure S3: Predicted monthly dengue importations by airport The number of predicted imported dengue infections for the top ten airports in non-endemic countries/states with vector presence for each month in (A) 2011 and (B) 2015. The error bars correspond to ± 1 standard deviation. Airports are abbreviated using the corresponding IATA code.




Figure S4: Dengue-infected passengers who continue to travel to non-endemic countries/states with vector presence for every route in the air transportation network This map corresponds to August 2015. The thickness as well as the colour of an edge represent the number of infected people travelling along the corresponding route. Blue represents relatively lower numbers of infected people, red represents relatively higher numbers of infected travellers and yellow represents the mid range.




Figure S5: Predicted dengue infections imported by returning residents and visitors in 2011 Here we show the results for non-endemic countries/states with vector presence with the highest number of predicted imported dengue cases in 2011. The bars are stacked to distinguish between returning residents (green) and visitors (blue). The blue solid line corresponds to the total number of imported cases. The error bars correspond to the model's coefficient of variation (see Material and methods).




Figure S6: Predicted imported dengue infections for returning residents and visitors for US states in 2011. The bars are stacked to distinguish between returning residents (green) and visitors (blue). The blue solid line corresponds to the total number of imported cases. The error bars correspond to the model's coefficient of variation (13.49%) that was inferred through Monte Carlo simulations.




Figure S7: Predicted imported dengue infections for returning residents and visitors for US states in 2015. The bars are stacked to distinguish between returning residents (green) and visitors (blue). The blue solid line corresponds to the total number of imported cases. The error bars correspond to the model's coefficient of variation (13.49%) that was inferred through Monte Carlo simulations.




Figure S8: Predicted imported dengue infections for returning residents and visitors for Australian states. ACT: Australian Capital Territory, NSW: New South Wales, NT: Northern Territory, SA: South Australia, TAS: Tasmania, VIC: Victoria, WA: Western Australia. The bars are stacked to distinguish between returning residents (green) and visitors (blue). The blue solid line corresponds to the total number of imported cases. The error bars correspond to the model's coefficient of variation (13.49%) that was inferred through Monte Carlo simulations.



Figure S9: Predicted imported dengue infections for returning residents and visitors for European countries in 2011. The bars are stacked to distinguish between returning residents (green) and visitors (blue). The blue solid line corresponds to the total number of imported cases. The error bars correspond to the model's coefficient of variation (13.49%) that was inferred through Monte Carlo simulations.




Figure S10: Predicted imported dengue infections for returning residents and visitors for European countries in 2015. The bars are stacked to distinguish between returning residents (green) and visitors (blue). The blue solid line corresponds to the total number of imported cases. The error bars correspond to the model's coefficient of variation (13.49%) that was inferred through Monte Carlo simulations.




Figure S11: Predicted imported dengue infections for returning residents and visitors for non-endemic African countries. The bars are stacked to distinguish between returning residents (green) and visitors (blue). The blue solid line corresponds to the total number of imported cases. The error bars correspond to the model's coefficient of variation (13.49%) that was inferred through Monte Carlo simulations.




Figure S12: Predicted percentage contribution of dengue importations by country of acquisition in 2011. The predicted percentage contribution by source country and month in 2011. The size and colour of the circles indicate the percentage contribution of the corresponding country to the total number of imported cases. The y-labels indicate the yearly percentage contribution of the corresponding source country.




Figure S13: Rank-based validation and correlation between reported and predicted imported cases for Queensland in 2011. (A) Countries are ranked by the total number of predicted and reported imported dengue cases. The reported ranking is then plotted against the predicted ranking. Countries that were ranked by the model, but did not appear in the dataset receive a rank of $i + 1$, where i is the number of unique importation sources according to the dengue case data. Similarly, countries that appeared in the data and were not ranked by the model receive a rank of $i + 1$. For circles that lie on the $x = y$ line (grey solid line) the predicted and reported rankings are equal. Circles that lie between the two dashed lines correspond to countries with a difference in ranking that is less than or equal to five. The circle areas are scaled proportionally to the number of reported cases that were imported from the corresponding country. Spearman's rank correlation coefficient between the absolute numbers of reported and predicted importations is equal to 0.58. (B) The absolute number of reported dengue importations are plotted against the absolute number of predicted importations.




Figure S14: The distribution of the coefficient of variation for several destinations. (A) Distributions for 2011. (B) Distributions for 2015.




Figure S15: Sobol's sensitivity analysis of the model's parameters. Parameter $\beta_{c,m}$ denotes the daily dengue incidence rate of country c during month m , parameter t_c denotes the number of days a traveller who arrives at a given airport has spent in country c and parameter n denotes the sum of the intrinsic incubation period and the infectious period in humans. **(A)** The first-order and total-order indices for the parameter ranges as shown in Table 1 of the main manuscript. The indices indicate that t_c is the most important model parameter. **(B)** The second-order indices for the parameter ranges as shown in Table 1 of the main manuscript. There is significant interactions between parameters t_c and $\beta_{c,m}$ and between parameters t_c and n . **(C)** The first-order and total-order indices for a shorter range of value ($[1, 30]$ days) for parameter t_c . In this case $\beta_{c,m}$ is the most important parameter. **(D)** The second-order indices for a shorter range of value ($[1, 30]$ days) for parameter t_c . There is still significant interaction between parameters t_c and $\beta_{c,m}$ and between parameters t_c and n .