Evidence of isotropy at large-scale from polarizations of radio sources

Prabhakar Tiwari¹ and Pankaj Jain²

National Astronomical Observatories, CAS, Beijing 100012, China e-mail: ptiwari@nao.cas.cn 2

Department of Physics, Indian Institute of Technology, Kanpur 208016, India

ABSTRACT

We test the isotropy of radio polarization angle orientations with a robust and reliable dual frequency polarimetric survey of active galactic nuclei (AGN). We find that the polarization orientations are consistent with the assumption of isotropy for scales larger than or equal to ~ 800 Mpc. This provides further evidence of isotropy at large distance scales and is likely to impose strong constraints on some of the physical mechanisms that may be invoked to explain past observations of alignment of radio and optical polarizations at large distance scales.

Key words. Cosmology: large-scale structure of Universe – general: polarization – galaxies: active

¹ National Astronomical Observatories, CAS, Beijing 1000 e-mail: ptiwari@nao.cas.cn
² Department of Physics, Indian Institute of Technology, F e-mail: pkjain@iitk.ac.in
April 7, 2024 **ABST**We test the isotropy of radio polarization angle orientatio survey of active galactic nuclei (AGN). We find that the p of isotropy for scales larger than or equal to ~ 800 Mpc. T scales and is likely to impose strong constraints on some o past observations of alignment of radio and optical polariza **Key words**. Cosmology: large-scale structure of Universe – **1. Introduction 1. Introduction**<l date. In particular the CMB itself shows several anomalies at large angular scales (de Oliveira-Costa et al. 2004; Ralston & Jain 2004; Schwarz et al. 2004, 2016; Aluri et al. 2017; Rath et al. 2018). The optical polarization from quasars shows polarization angle (PA) alignment over a very large scale of Gpc (Hutsemékers 1998), furthermore the radio polarizations also show similar alignment signal at Gpc scales (Hutsemékers et al. 2014).

Besides the large scale anisotropies described above, the JVAS/CLASS 8.4-GHz sample (Jackson et al. 2007) of flat-spectrum radio sources (FSRS) polarization angles also shows a significant evidence of alignment at distance scale of 150 Mpc (Tiwari & Jain 2013). However the significance of

these polarization angle alignment effects is not very large and requires confirmation with larger data samples. With same JVAS/CLASS data Joshi et al. (2007) report isotropy at relatively larger scales. Shurtleff (2014) reports less significant PA alignments in two circular regions of 24° radius on the sky. Jagannathan & Taylor (2014) also report the jet angle alignment across angular scales of up to 1.8 degrees (~ 53 Mpc at redshift one) from ELAIS N1 Deep radio survey. However, after so many different observations and PA alignment studies the situation is not very clear. These anisotropies if real and not some instrumental or observational artifacts potentially disrupt the modern standard cosmology and thus it is important to review and further investigate these claimed anisotropy signals with new and refined observations.

There exist many theoretical models which aim to explain the observed large scale optical alignment (Hutsemékers et al. 2014). These include mixing of electromagnetic radiation with hypothetical light pseudoscalar particles in correlated background magnetic fields (Jain et al. 2002; Agarwal et al. 2011; Piotrovich et al. 2009; Agarwal et al. 2012), vector perturbations (Morales & Saez 2007), large scale magnetic field coupled to dark energy (Urban & Zhitnitsky 2010), cosmic strings (Poltis & Stojkovic 2010; Hackmann et al. 2010), anisotropic expansion (Ciarcelluti 2012) and spontaneous violation of isotropy (Chakrabarty 2016). The radio alignment on small distance scales (Tiwari & Jain 2013; Jagannathan & Taylor 2014) may be explained by correlated supercluster magnetic field (Tiwari & Jain 2016). As we shall see our analysis is likely to impose significant constraints on these models.

In this work we employ the simultaneous dual frequency 86 GHz and 229 GHz polarimetric survey (Agudo et al. 2014) of radio flux and polarization for a large sample of 211

radio-loud active galactic nuclei (AGN) to test the hypothesis that the polarization vectors of AGNs are randomly oriented at large-scales in the sky. These are clear and reliable measure of radio polarizations from a dedicated polarimetric survey at two frequencies simultaneously and thus an excellent catalog to test isotropy at large-scales.

The outline of the paper is as follows. We provide the details about the survey and data in Section 2. In Section 3 we discuss the methods to measure the isotopy and quantify the anisotropy signal. We present the analysis and our detailed results in Section 4. In Section 5 we discuss and conclude this work.

2. Data sample

The dual frequency 86 GHz and 229 GHz catalog we use contains 221 radio-loud AGNs (Agudo et al. 2014). The observations were performed on the IRAM (Institute for Radio Astronomy in the Millimeter Range) 30 meter telescope with the XPOL polarimeter (Thum et al. 2008). The sample AGNs are flux limited and are above 1 Jy (total-flux) at 86 GHz, the redshift of 199 AGNs is known and ranges from z = 0.00068 to 3.408. The mean and median redshift of the sample is 0.937 and 0.859, respectively. The sample is dominated by blazars and contains 152 quasars, 32 BL Lacs and 21 radio galaxies and 6 unclassified sources. The linear polarization above 3σ median level of $\sim 1\%$ is detected for 183 sources. In the sample, for 22 sources the linear polarization angle is measured at both 86 and 229 GHz, and a good match between the PAs at these frequencies is seen (see Fig. 14 in Agudo et al. 2014). The sample is dominantly in the northern sky and covers the entire northern hemisphere nearly uniformly. We have shown the sky distribution of sources in Fig. 1. The short millimeter survey is an excellent probes of radio loud AGN jets and has several advantages over radio centimeter i.e. \sim GHz surveys. The millimeter radiation is predominately from the cores of synchrotron-emitting relativist jets and has negligible contribution from host AGN and its surroundings. It is also less affected by Faraday rotation and depolarization (Zavala & Taylor 2004; Agudo et al. 2010). The PAs in the sample are uniform between 0 to 180° , their distribution is shown in Fig. 2. The jet position angle with respect to polarization angle in sample are not preferably parallel or perpendicular (Agudo et al. 2014). In general the survey sample is robust and no suspicious systematics or unusual behaviour is observed. Further details about the observation and calibration can be found in Agudo et al. (2014).

3. Measure of anisotropy

The polarization angles in sample are on the hypothetical celestial sphere and directional measurement on this sphere corresponds to a particular coordinate system. To test our hypothesis of isotropy, we resort to Jain et al. (2004) coordinate independent statistics procedure and compare the polarization vector of a source after transporting it to the position of a reference source along the geodesic joining the two. We define the measure of isotropy as following (Hutsemékers 1998; Jain et al. 2004). We have redshift and angular position i.e. right ascension and declination given for each source. We define the separation between



Fig. 1: The sky distribution of sources and their respective polarization angles measured with respect to the local longitude.



Fig. 2: The polarization angle distribution in sample

any two given sources in terms of comoving distance. For calculating comoving distances we have assumed Λ CDM and cosmological parameters from the latest *Planck* results (Planck Collaboration et al. 2018). Next we consider the n_v nearest neighbours of a source situated at site k with its polarization angle ψ_k . Let ψ_i be the polarization angle of the source at i^{th} site within the n_v nearest neighbour set. A dispersion measure of polarization angles relative to k^{th} source ψ_k with its n_v nearest neighbour is written as,

$$d_k = \frac{1}{n_v} \sum_{i=1}^{n_v} \cos[2(\psi_i + \Delta_{i \to k}) - 2\psi_k],$$
(1)

where $\Delta_{i\to k}$ is a correction to angle ψ_i due to its parallel transport from site $i \to k$ (Jain et al. 2004). The polarization angles span a range of 0 to 180° and to make them behave like usual angles and take values over entire range 0 to 360° we multiply the polarization angles by two (Ralston & Jain 1999). The function d_k is the average of the cosine of the differences of the polarization vector at site k and those of its n_v nearest neighbour set and hence is a measure of the dispersion in angles. It takes on higher values for data with lower dispersions and vice versa. We take the average of d_k over all source sites and define this



Fig. 3: The average distance corresponding to n_v . The distance is computed assuming Λ CDM and the cosmological parameters are taken from the latest Planck results (Planck Collaboration et al. 2018). 'All sources (z)' sample contains 175 sources, all these have redshifts and good polarization measurements. 139 out of these 175 sources are quasars, 69 quasar are above redshift one.

as a measure of alignment in sample,

$$S_D = \frac{1}{N_t} \sum_{k=1}^{N_t} d_k,$$
 (2)

where N_t is the total number of sources in the sample. Similar alternate statistics can also be defined as a measure of alignment (Bietenholz 1986; Hutsemékers 1998; Jain et al. 2004; Tiwari & Jain 2013; Pelgrims & Cudell 2014). The nearest neighbour statistics uses number of nearest neighbour, n_v , as a proxy of distance. This is to fix *statistics* at each source location i.e. to have same n_v while calculating d_k . The average distance corresponding to n_v for different sub-samples studies in this work is given in Fig. 3.

The error in the statistic S_D is computed using jackknife method (Tiwari & Jain 2016) and the significance is computed by comparing the random sample S_D with the data S_D . The jackknife error on S_D for a given n_v is calculated by re-sampling the data. We eliminate the *i*th source from the full sample and calculate the correlation statistics $S_D(i)$. Given full sample statistics S_D the jackknife error δS_D in its estimation is given as,

$$(\delta S_D)^2 = \frac{(N_t - 1)}{N_t} \sum_{i=1}^{N_t} (S_D(i) - S_D)^2.$$
(3)

4. Analysis and results

Our full sample contains 211 sources; 175 out of these have both redshift and good polarization measurements and so a good sample for our analysis. There are 183 sources with jet angles measurements along with redshifts. The S_D measurements and its significance as compared to random sample are shown in Fig. 4. The σ significance i.e. how the sample behaves with respect to random isotropic and uniform polarization angles is defined as follow,

$$\sigma = \frac{S_D(\text{data}) - S_D(\text{random})}{\sqrt{(\delta S_D^{\text{data}})^2 + (\delta S_D^{\text{random}})^2}}$$
(4)



Fig. 4: The statistics S_D of all sources and its significance with respect to random isotropic polarization distribution. The error bars on data S_D are jackknife errors and the random polarization angle S_D is also shown with its RMS errors bars drawn from 1000 random samples. The σ significance is the difference between data S_D and random S_D as defined in Eq. 4.

where δS_D^{data} and $\delta S_D^{\text{random}}$ are jackknife errors on data S_D and RMS errors on random S_D^{random} , respectively. The results with full data in Fig. 4 shows no significant deviation from isotropy and are well within one sigma of the isotropic and uniform distribution of polarization angles. The jet angles also show good agreement with random jet angle distribution, the results are shown in Fig. 5. The average distance to first nearest neighbour in full sample is 719 Mpc, thus the observation of isotropy in this work applies for distance scales ≥ 719 Mpc. We are limited by source number density to probe smaller scales and so can't explore the polarization angle alignment signal claimed at scales of order 100 Mpc (Tiwari & Jain 2013).

There have been several observations of quasar polarization alignments at large distance scales (Hutsemékers et al. 2014; Pelgrims & Hutsemékers 2015, 2016) and so we also explore the anisotropy in quasars only sample. We have in total 139 quasars with redshift and measure of polarization angles in our full sample. These are evenly distributed over northern sky (Fig. 1) and centered around redshift one (see figure 2 and 3 in Agudo et al. 2014). Again, we do not see any alignment in this quasar only sample and the statistics agrees well with random distribution. The results are shown in Fig. 6. The least nearest neighbour distance in this sample is 849 Mpc and so the quasar polarization angle distribution is isotropic at least at this scale and above.

Next, we test if the alignment signal is redshift dependent, and if it is present with high redshift quasars (Pelgrims & Hutsemékers 2015). We consider quasars with redshift larger than one and calculate S_D , this sample contains 67 quasars and for this sample the average distance to first nearest neighbour is 1153 Mpc. We find that this sample also agrees well with random, no significant alignment is seen, the results are shown in Fig. 7. The quasar sample with redshift less than one is also consistent with isotropic polarization distribution.



Fig. 5: The statistics S_D of jet angles. Jet position angles are fairly isotropic and agree well within one σ with random sample. Other details same as in Fig. 4



Fig. 6: The statistics S_D exclusive to quasar sample. Other details same as in Fig. 4

5. Discussion and Conclusion

The polarization angle alignment of radio sources and quasars at large distance scales remains puzzling since very long (Birch 1982; Kendall & Young 1984; Hutsemékers 1998; Jain & Ralston 1999; Tiwari & Jain 2013; Hutsemékers et al. 2014; Pelgrims & Hutsemékers 2015, 2016). Surprisingly, most of these along with cosmic microwave background (CMB) dipole-quadrupoleoctopole (de Oliveira-Costa et al. 2004; Schwarz et al. 2004), radio galaxy distribution dipole (Singal 2011; Gibelyou & Huterer Rubart & Schwarz 2012;2013;Tiwari et al. 2015; Tiwari & Jain 2015; Tiwari & Nusser 2016; Colin et al. 2017), and polarizations at optical frequencies at cosmological scale indicate a preferred direction pointing roughly towards the Virgo cluster (Ralston & Jain 2004). Although there exist some explanations for the large-scale optical polarization alignment



Fig. 7: The quasar sample with redshift greater than one. Other details same as in Fig. 4

following axion-photon interaction (Agarwal et al. 2012), and radio polarization alignment at 150 Mpc scale in terms of galaxy supercluster magnetic field (Tiwari & Jain 2016); the alignment signal reported in Hutsemékers (1998); Hutsemékers et al. (2014); Pelgrims & Hutsemékers (2015, 2016) remains puzzling. In this work we explore on largescale millimeter radio polarization alignment and report isotropy.

The observed isotropy is likely to impose significant constrains on the proposed mechanisms for large scale polarization alignment, discussed in the Introduction (Jain et al. 2002; Agarwal et al. 2011; Piotrovich et al. 2009; Agarwal et al. 2012; Morales & Saez 2007; Urban & Zhitnitsky 2010; Poltis & Stojkovic 2010; Hackmann et al. 2010; Ciarcelluti 2012; Chakrabarty 2016). The constraints may be more stringent on mechanisms which predict an effect independent of frequency or a larger effect at smaller frequency. This is because such models will predict an alignment of same or higher strength in comparison to that seen at optical frequencies (Hutsemékers 1998). In contrast the constraints may not be very stringent on a mechanism, such as pseudoscalar-photon mixing (Jain et al. 2002), which increases with frequency. Many of these mechanisms rely on the presence of large scale correlated magnetic field. Hence the observed alignment may also be a probe of the correlations in the large scale magnetic field. However this relationship between alignment and magnetic field is not direct since it also relies on some other effect, such as the mixing of electromagnetic radiation with hypothetical pseudoscalar particles (Andriamonje et al. 2007; Tiwari 2012; Payez et al. 2012; Wouters & Brun 2013; Ayala et al. 2014; Tiwari 2017). Only on small distance scales of order 100 Mpc, it has been speculated that supercluster magnetic field may directly affect the alignment of galaxies (Tiwari & Jain 2016). However, in this case also the alignment effect will depend on the level of randomness introduced by local effects within individual galaxies.

The alignment of radio polarization reported earlier were usually in 2D due to unavailability of redshift, so to test if the alignment signal was a consequence of 2D projection, we calculate S_D for our sample at fixed redshift, i.e. projecting all sources at same redshift. Even with 2D statis-



Fig. 8: The statistics S_D calculated with 2D sample. Other details same as in Fig. 4



Fig. 9: Alignment at fixed distance. S_D is average dispersion over the spheres of radius 'Distance' on x-axis at each source site. The number of nearest neighbour n_v at each site varies here slightly. Other details same as in Fig. 4

tics we find the samples (all sources, quasar only sample) agreeing well with random distribution (Fig. 8).

We also test the data with alternate statistics. In particularly we test the data by averaging S_D at fixed distance i.e. averaging dispersion over the spheres of fixed radii at each sites. This also closely matches with random isotropic polarization distribution, results are shown in Fig. 9. Furthermore, the high redshift quasar distribution is also isotropic.

We have tested the large-scale radio polarization alignment signal with a robust and simultaneous dual frequency radio polarimetric survey (Agudo et al. 2014). We do not find any observation of alignment at large-scale and polarization angles of AGNs and jet angles are fairly uniform and isotropic at scale equal and above \sim Gpc. However, due to low number density we are unable to probe scales less than 719 Mpc and cannot examine the alignment claimed at relatively smaller scales (Tiwari & Jain 2013; Jagannathan & Taylor 2014). Furthermore with this sample we cannot explore the alignment with the axis of large quasar groups (LQGs) (Pelgrims & Hutsemékers 2016). This is because the LQGs are typically of smaller size in comparison to distance scale of the first neighbour in our sample.

We note that the jet position angle with respect to polarization angle in sample are not preferably parallel or perpendicular (Agudo et al. 2014) and this is somewhat unusual and unlike the predictions from axially symmetric jet models (Falle 1991; Martí et al. 1997; Komissarov 1999; Fendt et al. 2012). This probably indicate that the magnetic field or the particles in radio emission region are non axisymmetric and somehow due to local magnetic fields or other effects the polarization vector is misaligned with respect to jet position angle. However, this observation does not change our conclusion as we find isotropy with jet position angle as well.

Nevertheless, this work adds to the previous studies of large-scale radio polarization alignment anomalies. With this data we clearly see the isotropy above Gpc. These results further support the isotropy assumption in cosmology along with CMB and other supporting large-scale isotropy observations.

6. Acknowledgments

This work is supported by NSFC Grants 1171001024 and 11673025, the National Key Basic Research and Development Program of China (No. 2018YFA0404503) and the Science and Engineering Research Board (SERB), Government of India. The work is also supported by NAOC youth talent fund 110000JJ01.

References

- Abu-Zayyad, T., Aida, R., Allen, M., et al. 2012, The Astrophysical Journal, 757, 26
- Agarwal, N., Aluri, P. K., Jain, P., Khanna, U., & Tiwari, P. 2012, EPJC, 72, 15
- Agarwal, N., Kamal, A., & Jain, P. 2011, Phys. Rev., D83, 065014
- Agudo, I., Thum, C., Gómez, J. L., & Wiesemeyer, H. 2014, A&A, 566, A59
- Agudo, I., Thum, C., Wiesemeyer, H., & Krichbaum, T. P. 2010, ApJ. S. 189. 1
- Aluri, P. K., Ralston, J. P., & Weltman, A. 2017, Mon. Not. Roy. Astron. Soc., 472, 2410
- Andriamonje, S., Aune, S., Autiero, D., et al. 2007, JCAP, 4, 010
- Ayala, A., Domínguez, I., Giannotti, M., Mirizzi, A., & Straniero, O.
- 2014, Physical Review Letters, 113, 191302 Bennett, C. L., Larson, D., Weiland, J. L., et al. 2013, ApJ. S, 208, 20
- Bietenholz, M. F. 1986, AJ, 91, 1249
- Birch, P. 1982, Nature, 298, 451
- Chakrabarty, S. S. 2016, Physical Review D., 93, 123507
- Ciarcelluti, P. 2012, Mod. Phys. Lett., A27, 1250221
- Colin, J., Mohayaee, R., Rameez, M., & Sarkar, S. 2017, Mon. Not. Roy. Astron. Soc., 471, 1045
- de Oliveira-Costa, A., Tegmark, M., Zaldarriaga, M., & Hamilton, A. 2004, PhRvD, 69, 063516
- Falle, S. A. E. G. 1991, MNRAS, 250, 581
- Fendt, C., Porth, O., & Vaidya, B. 2012, in Journal of Physics Conference Series, Vol. 372, Journal of Physics Conference Series, 012011 Gibelyou, C. & Huterer, D. 2012, MNRAS, 427, 1994
- Hackmann, E., Hartmann, B., Lammerzahl, C., & Sirimachan, P. 2010, Phys. Rev., D82, 044024
- Hutsemékers, D. 1998, A&A, 332, 410
- Hutsemékers, D., Braibant, L., Pelgrims, V., & Sluse, D. 2014, A&A, 572, A18
- Jackson, N., Battye, R., Browne, I., et al. 2007, MNRAS, 376, 371
- Jagannathan, P. & Taylor, R. 2014, in American Astronomical Society Meeting Abstracts, Vol. 223, American Astronomical Society Meeting Abstracts #223, 150.34
- Jain, P., Narain, G., & Sarala, S. 2004, MNRAS, 347, 394

Article number, page 5 of 6

- Jain, P., Panda, S., & Sarala, S. 2002, PhRvD, 66, 085007
- Jain, P. & Ralston, J. P. 1999, Modern Physics Letters A, 14, 417
- Joshi, S., Battye, R., Browne, I., et al. 2007, MNRAS, 380, 162
- Kendall, D. G. & Young, G. A. 1984, MNRAS, 207, 637
 Komissarov, S. S. 1999, MNRAS, 308, 1069
- Martí, J. M., Müller, E., Font, J. A., Ibáñez, J. M. Z., & Marquina, A. 1997, ApJ, 479, 151
- Milne, E. A. 1933, Zeitschrift für Astrophysik, 6, 1
- Milne, E. A. 1935, Relativity, gravitation and world-structure
- Morales, J. A. & Saez, D. 2007, Phys. Rev., D75, 043011
- Payez, A., Cudell, J., & Hutsemekers, D. 2012, JCAP, 1207, 041 Pelgrims, V. & Cudell, J. R. 2014, MNRAS, 442, 1239
- Pelgrims, V. & Hutsemékers, D. 2015, MNRAS, 450, 4161
- Pelgrims, V. & Hutsemékers, D. 2016, A&A, 590, A53 Penzias, A. A. & Wilson, R. W. 1965, ApJ, 142, 419
- Piotrovich, M. Yu., Gnedin, Yu. N., & Natsvlishvili, T. M. 2009, Astrophysics, 52, 451, [Astrofiz.52, 451(2009)]
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, A&A, 594, A16
- Planck Collaboration, Aghanim, N., Akrami, Y., et al. 2018, ArXiv e-prints [arXiv:1807.06209]
- Poltis, R. & Stojkovic, D. 2010, Phys. Rev. Lett., 105, 161301
- Ralston, J. P. & Jain, P. 1999, IJMPD8 (1999) 537-547, 8, 537
- Ralston, J. P. & Jain, P. 2004, IJMPD, 13, 1857
- Rath, P. K., Samal, P. K., Panda, S., & Mishra, D. D. 2018, Mon. Not. Roy. Astron. Soc., 475, 4357
- Rubart, M. & Schwarz, D. J. 2013, A&A, 555 [arXiv:1301.5559]
- Schwarz, D. J., Copi, C. J., Huterer, D., & Starkman, G. D. 2016, Classical and Quantum Gravity, 33, 184001
- Schwarz, D. J., Starkman, G. D., Huterer, D., & Copi, C. J. 2004, PhRvL, 93, 221301
- Shurtleff, R. 2014 [arXiv:1408.2514]
- Singal, A. K. 2011, ApJL, 742, L23
- Thum, C., Wiesemeyer, H., Paubert, G., Navarro, S., & Morris, D. 2008, Publications of the Astronomical Society of the Pacific, 120, 777
- Tiwari, P. 2012, PhRvD, D86, 115025
- Tiwari, P. 2017, Phys. Rev., D95, 023005
- Tiwari, P. & Jain, P. 2013, Int.J.Mod.Phys., D22, 1350089 (26 pages)
- Tiwari, P. & Jain, P. 2015, MNRAS, 447, 2658
- Tiwari, P. & Jain, P. 2016, MNRAS, 460, 2698
- Tiwari, P., Kothari, R., Naskar, A., Nadkarni-Ghosh, S., & Jain, P. 2015, Astroparticle Physics, 61, 1
- Tiwari, P. & Nusser, A. 2016, Journal of Cosmology and Astroparticle Physics, 2016, 062
- Urban, F. R. & Zhitnitsky, A. R. 2010, Phys. Rev., D82, 043524
- Řípa, J. & Shafieloo, A. 2017, ApJ, 851, 15
- White, M., Scott, D., & Silk, J. 1994, ARAA, 32, 319
- Wouters, D. & Brun, P. 2013, Astrophys.J., 772, 44
- Zavala, R. T. & Taylor, G. B. 2004, ApJ, 612, 749