

Charged rotating black holes coupled with nonlinear electrodynamics Maxwell field in the mimetic gravity

G.G.L. Nashed^{1,2,3,*}, W. El Hanafy^{1,2,†} and Kazuharu Bamba^{4‡}

¹Centre for Theoretical Physics, The British University in Egypt, P.O. Box 43, El Sherouk City, Cairo 11837, Egypt

²Egyptian Relativity Group (ERG), Cairo University, Giza 12613, Egypt

³Mathematics Department, Faculty of Science, Ain Shams University, Cairo 11566, Egypt and

⁴Division of Human Support System, Faculty of Symbiotic Systems Science, Fukushima University, Fukushima 960-1296, Japan

In mimetic gravity, we derive D -dimension charged black hole solutions having flat or cylindrical horizons with zero curvature boundary. The asymptotic behaviours of these black holes behave as (A)dS. We study both linear and nonlinear forms of the Maxwell field equations in two separate contexts. For the nonlinear case, we derive a new solution having a metric with monopole, dipole and quadrupole terms. The most interesting feature of this black hole is that its dipole and quadrupole terms are related by a constant. However, the solution reduces to the linear case of the Maxwell field equations when this constant acquires a null value. Also, we apply a coordinate transformation and derive rotating black hole solutions (for both linear and nonlinear cases). We show that the nonlinear black hole has stronger curvature singularities than the corresponding known black hole solutions in general relativity. We show that the obtained solutions could have at most two horizons. We determine the critical mass of the degenerate horizon at which the two horizons coincide. We study the thermodynamical stability of the solutions. We note that the nonlinear electrodynamics contributes to process a second-order phase transition whereas the heat capacity has an infinite discontinuity.

I. INTRODUCTION

In the last few decades, several cosmological observations have confirmed that our universe is dominated by dark components: dark matter and dark energy [1–3]. The competition to criticize the origin and the nature of these ingredients either within observational or theoretical framework is going on equal footing. In the observational framework, it has been shown that about 26% of the energy content in our universe belongs to the sector of dark matter, however, about 69% specifies the dark energy. In the theoretical framework, modified gravity theories are the most appealing by invoking a geometrical origin to explain these ingredients [4–16]. Among modified gravity theories, we particularly mention two approaches that gain attention in literature, the curvature based $f(R)$ gravity, c.f. [17–22] and the torsion based $f(T)$ teleparallel gravity, c.f. [23–30] (for recent reviews on modified theories of gravity and the issue of dark energy, see, for example [15, 16, 22, 31–34]). Another modified scenario that has been proposed recently is the mimetic gravity one [35]. Mimetic theory has many applications in cosmology [36, 37, 39–41] as well as in the solar system [42–46], see also the recent review [47].

Although this theory provides geometrical foundations to explain dark components as in modified gravity, it presents a distinguishable framework to deal with that problem. The construction of the mimetic theory can be derived from the general relativity theory by separating the conformal degree of freedom of the gravitational field via re-parameterizing the physical metric by an auxiliary metric with a mimetic field ϕ . The equations of motion in the mimetic theory are characterized by an extra term sourced by the mimetic field. The mimetic scalar field is generated by a singular limit of the general informal transformation which is not invertible [48–50], and its kinetic term is provided to satisfy the kinematical constraint

$$g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi = -1. \quad (1)$$

Using the above constraint, the equations of motion have shown to be traceless, where the scalar field mimics pressureless cold dark matter (CDM) just as in general relativity even for vacuum solution [47]. While generalizing the model by invoking potential functions, one can produce a unified cosmic history nothing to speak of bouncing cosmology [51–54]. The mimetic gravity has shown to be invariant under Weyl transformation [55, 56]. On its perturbation level, the sound speed of the scalar fluctuations is exactly zero, $c_s = 0$, so to obtain a successful inflation one may impose a higher-derivative term $(\Box\phi)^2$ to grant the scalar fluctuation a nonzero sound speed [57, 58]. This feature may have important impacts on large structure and galaxy formation specially for very small but not vanishing values of the sound speed. In particular, the higher-derivative term departs

*Electronic address: nashed@bue.edu.eg

†Electronic address: waleed.elhanafy@bue.edu.eg

‡Electronic address: bamba@sss.fukushima-u.ac.jp

the properties of the mimetic dark matter from being a perfect dust-like fluid [59–61]. In these cases, imperfect dark matter could be relevant to the missing-satellites problem and the core-cusp problem [62].

One of the crucial issues in mimetic gravity is stability. In fact, there are many studies on the stability of mimetic gravity. Although, the theory does not suffer from gradient instabilities or those are associated to higher derivatives (Ostrogradski ghost), the theory might have ghost instability [63]. However, the later can be resolved by coupling higher derivatives of the mimetic field to curvature. Besides that it has been shown that the strong coupling scale can be raised to 10 TeV [64]. This cutoff scale on spatial momenta of ghosts makes the mimetic matter scenario is phenomenologically viable. At a lower scale, it reduces to GR supplemented by a fluid with small positive sound speed $c_s^2 \leq 10^{-20}$, which is compatible with the observations of the photon flux from vacuum decay $c_s^2 \geq 10^{-42}$. Also the authors of [59, 65], independently, found that the theory is stable if the density of the mimetic field is positive. This preliminary analysis favors the solutions with de Sitter backgrounds in presence of a cosmological constant, since in this case both the curvature and the trace of the matter energy-momentum tensor contribute to hold the energy density positive. However, the theory might still suffer from the gravitational instability associated with caustic surfaces of the geodesic flow [65]. Another work using the effective theory approach has been developed to overcome the problem of gradient instability [66]. Also, it has been shown that the mimetic constraint, namely Eq. (1), can be implemented in the action using a Lagrangian multiplier [67]. Using a suitable gravitational Lagrange multiplier could provide a healthy mimetic gravity free from ghosts [34]. Moreover, in Gauss-Bonnet theories ghost modes can be removed at the level of equations of motion [68].

Recently one can find several variants of mimetic gravity: The mimetic $f(R)$ gravity, where some quantum/string corrections can be taken into account by adding higher-order curvature invariants to the action of mimetic gravitation theory [51, 69]. Same technique has been applied to construct mimetic Gauss-Bonnet theories $f(G)$ [70], mimetic $f(R, T)$ [71], mimetic $f(R, \phi)$ [58], mimetic covariant Horava-like gravity [72, 73], mimetic Galileon gravity [74, 75], mimetic Horndeski gravity [76], unimodular-mimetic $f(R)$ gravity [52], mimetic Born-Infeld gravity [77, 78] and non-local mimetic $f(R)$ gravity [79]. Notably, the importance of higher-derivative invariants in the Lagrangian of the mimetic theory has been analyzed in [66, 80–85]. There are many other amendments of the mimetic theory have been considered, for example by including the vector-tensor mimetic gravity [86], bi-scalar mimetic models [87], the one in which the implementation of the limiting curvature hypothesis is considered to solve the issues of the cosmological singularity [36, 37] and the braneworld mimetic gravity [88]. Also, the models where the mimetic field couples to matter non-minimally [89] and the currents of baryon number have been also discussed in [90].

Modified gravity theories are usually tested via cosmological models and black hole physics. The aim of this work is to test mimetic gravity coupled to linear/nonlinear electrodynamics in the black hole physics domain. According to “no hair” theorem, the black hole is characterized by three conserved quantities: ADM mass, spin and charge. Static spherically symmetric spacetime is known as Schwarzschild black hole, rotating case is known as Kerr black hole, when static spherically symmetric black hole gains charges, it is called Reissner-Nordström (RN) and the rotating charged black hole is known as Kerr-Newmann (KN). The later type of black holes, interestingly, undergoes a phase transition associated with an infinite discontinuity of the black hole heat capacity [91]. In practice, mass and spin are tested and confirmed by observations. However, charged black holes are believed to be not existing in real scenarios. This is because that requires a rapid neutralization. On the other hand, it is interesting to investigate mechanisms of producing charged black holes. This has been examined even within classical framework, c.f [92], others are to study the charge production mechanism by involving dark matter [93–95]. Charged black hole solutions have been examined in several theoretical aspects c.f. [96–100]. However, the mimetic gravity has been proven to be a good candidate to describe dark matter. In this sense, we find studying charged black hole solutions in mimetic gravity as in the present paper is a step for more investigation in this regard. In astrophysics domain, the effect of charge on the merge rate of binary black holes has been studied recently [101, 102]. In [101], it has been shown that a fast radio burst (FRB) or a gamma-ray burst (GRB) can be explained, depending on the value of the black hole charge. It sets lower limits on the charge necessary to produce each phenomenon. It has been shown that, for a $10 M_\odot$ black hole, the merger can produce a FRB, if the charge of one members of the black hole charge is more than $\sim 10^{12}$ Coulombs. If its charge is more than $\sim 10^{16}$ Coulombs, it can generate a GRB. Future joint gravitational waves (GW)/GRB/FRB searches, specially after LIGO discoveries GW150914, GW151226 and LVT151012, may set some constraints on charged black holes.

Asymptotically anti-de Sitter (AdS) black holes have been studied extensively after Hawking-Page paper [103], in which they discussed a phase transition in the case of Schwarzschild-AdS black hole. Since then thermodynamics of more complicated AdS black holes have been investigated, a first order phase transition has been viewed in the case of Reissner-Nordström-anti de Sitter (RN-AdS) [104, 105]. Also, a second order phase transition has been realized in the case of KN-AdS [106]. In the mimetic $f(R)$ gravitational theory the action is adjusted by making use of a Lagrangian multiplier and mimetic potential, then a vacuum solution of RN-AdS black hole has been derived under some restrictions distinguishing this mimetic $f(R)$ variant from the $f(R)$ gravity [53]. It is the aim of the present study to derive a novel class of solutions of charged black holes coupled with the linear and the nonlinear electrodynamics Maxwell field in the context of mimetic gravitational theory. We derive analytical

charged black hole solutions in D -dimension, using the nonlinear electrodynamics. We show that the metric contains monopole, dipole and quadrupole terms. Interestingly, the dipole and quadrupole terms are strongly related to some constant so that these terms vanish when the constant has a nil value, and then the solution reduces to the linear Maxwell field solution.

The arrangement of this study is as follow: In Section II, a brief review of the mimetic gravitational theory is given. In Section III, a new black hole solution is derived. The solution behaves asymptotically as a flat spacetime. In Section IV, we apply a coordinate transformation to the obtained solution, then derive analytic D -dimension rotating charged solutions in the framework of mimetic theory. In Section V, we derive new D -dimension charged black hole solutions, using the nonlinear electrodynamics, which show that the metric has monopole, dipole and quadrupole terms. In Section VI, we study the properties of the black hole solutions derived in Sections III and V by calculating the curvature invariants. This show that the black hole solutions in the nonlinear electrodynamics case have singularities stronger than those derived from the linear case. In Section VII, we study the thermodynamic properties of the solutions derived in Sections III and V. Final section, is devoted to summarize the present study.

II. PRELIMINARIES OF MIMETIC GRAVITATIONAL THEORY

In this section, we discuss the case when mimetic gravity is coupled to electrodynamics in presence of a cosmological constant. The original mimetic gravity variant was first constructed in the frame of the general relativity to investigate dark matter in cosmology [35]. The construction of the theory depends on redefining of the physical metric $g_{\mu\nu}$ [35]:

$$g_{\alpha\beta} = -(\bar{g}^{\mu\nu} \partial_\mu \phi \partial_\nu \phi) \bar{g}_{\alpha\beta}, \quad (2)$$

where $\bar{g}_{\alpha\beta}$ is the conformal auxiliary metric, ϕ is the mimetic scalar field and $\bar{g}^{\alpha\beta}$ is the inverse of $\bar{g}_{\alpha\beta}$. Using Eq. (2) one can show that GR theory is invariant under the conformal transformation, i.e., $\bar{g}_{\alpha\beta} \rightarrow \omega(x_\mu) \bar{g}_{\alpha\beta}$ with $\omega(x_\mu)$ being an arbitrary function of the coordinates. Equation (2) shows that the mimetic field should satisfy Eq. (1).

Now we consider the mimetic Maxwell theory in the presence of a cosmological constant Λ . Thence, the action of this field is given by

$$S := \frac{1}{2\chi} \int d^D x \sqrt{-g(\bar{g}_{\mu\nu}, \phi)} \{R(\bar{g}_{\mu\nu}, \phi) - 2\Lambda\} - \int d^D x \sqrt{-g(\bar{g}_{\mu\nu}, \phi)} \mathcal{L}_{em}, \quad (3)$$

where χ is the D -dimensional gravitational constant $\chi = 2(D-3)\Omega_{D-1}G_D$ and G_D is the gravitation Newtonian constant in D -dimensions, the D -dimensional cosmological constant $\Lambda = -\frac{(D-1)(D-2)}{2\lambda^2}$ with a length scale λ of the dS spacetime, and $\mathcal{L}_{em} = F \wedge \star F$ is the Maxwell Lagrangian, with $F = dV$ and $V = V_\mu dx^\mu$ being the gauge potential 1-form [107, 108]. We denote the volume of $(D-1)$ -dimensional unit sphere by Ω_{D-1} , where

$$\Omega_{D-1} = \frac{2\pi^{(D-1)/2}}{\Gamma((D-1)/2)}, \quad (4)$$

with Γ being the gamma function that depends on the dimension of the space-time¹ and $g \equiv g(\bar{g}_{\mu\nu}, \phi)$ being the determinant of the physical metric defined by Eq. (2).

Varying the action of Eq. (3) with respect to the physical metric, one can derive the following equations of motion of the gravitational field

$$E_\mu{}^\nu \equiv G_\mu{}^\nu + \frac{1}{2}\delta_\mu^\nu \Lambda - \bar{\mathcal{T}}_\mu{}^\nu - \tilde{\mathcal{T}}_\mu{}^\nu = 0, \quad (5)$$

where $G_{\mu\nu}$ is Einstein tensor, $\tilde{\mathcal{T}}_\mu{}^\nu$ is the energy-momentum tensor of the mimetic field and $\bar{\mathcal{T}}_\mu{}^\nu$ is the energy-momentum tensor of the electromagnetic field

$$\bar{\mathcal{T}}_\mu{}^\nu = F_{\mu\alpha} F^{\nu\alpha} - \frac{1}{4}\delta_\mu^\nu F_{\alpha\beta} F^{\alpha\beta}. \quad (6)$$

Notably, the auxiliary metric does not appear explicitly in the field equations, but it implicitly does through the physical metric given in Eq. (1) and the mimetic field ϕ . The presence of the mimetic field in the field equations can be written as

$$\tilde{\mathcal{T}}_{\mu\nu} = -\left(G + \frac{D}{2}\Lambda - \bar{\mathcal{T}}\right)\partial_\mu \phi \partial_\nu \phi, \quad (7)$$

¹ For $D = 4$, one can recover $2(D-3)\Omega_{D-1} = 8\pi$.

where $G = -\left(\frac{D-2}{2}\right)R$ is the trace of Einstein tensor. Finally, the variation in term of the action (3) with respect to the vector potential V_μ yields [107]

$$\partial_\nu \left(\sqrt{-g} F^{\mu\nu} \right) = 0. \quad (8)$$

It is worth to mention that the energy-momentum tensors, $\mathcal{T}_{\mu\nu}^{\text{em}}$ and $\tilde{\mathcal{T}}_{\mu\nu}$, are conserved, i.e. satisfy the continuity equations $\nabla^\mu \mathcal{T}_{\mu\nu}^{\text{em}} = 0 = \nabla^\mu \tilde{\mathcal{T}}_{\mu\nu}$, where ∇ is the covariant derivative. Using the mimetic field constraint (1) and the energy-momentum tensor (7), the corresponding continuity reads

$$\nabla^\kappa \left(\left[G + \frac{D}{2} \Lambda - \mathcal{T}^{\text{em}} \right] \partial_\kappa \phi \right) = \frac{1}{\sqrt{-g}} \partial_\kappa \left(\sqrt{-g} \left[G + \frac{D}{2} \Lambda - \mathcal{T}^{\text{em}} \right] g^{\kappa\sigma} \partial_\sigma \phi \right) = 0. \quad (9)$$

Alternatively, one finds that (1) is satisfied identically, when (9) is used. It is straightforward to show that the trace of Eq. (5) has the form

$$\left[G + \frac{D}{2} \Lambda - \mathcal{T}^{\text{em}} \right] \left(1 + g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \right) \equiv 0, \quad (10)$$

which is satisfied identically due to the mimetic field constraint, namely Eq. (1). In conclusion, we note that the conformal degree of freedom provides a dynamical quantity, i.e. ($G \neq 0$), and therefore the mimetic theory has non-trivial solutions for the conformal mode even in the absence of matter [35].

III. D-DIMENSION CHARGED BLACK HOLES IN MIMETIC GRAVITY

In this section, we present D -dimension solution of a charged black hole in mimetic gravity. So we suppose the spacetime configuration is given by the metric

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2 \left(\sum_{i=1}^{\ell} d\theta_i^2 + \sum_{k=1}^{D-\ell-2} dz_k^2 \right), \quad (11)$$

where $0 \leq r < \infty$, $-\infty < t < \infty$, $0 \leq \theta_i < 2\pi$ and $-\infty < z_k < \infty$. Here $f(r)$ is unknown function of the radial coordinate r only. For the spacetime (11), we get the Ricci scalar

$$R = -\frac{r^2 f'' + 2(D-2)rf' + (D-2)(D-3)f}{r^2}. \quad (12)$$

Then, the non-vanishing components of Eqs. (5) and (6) read the following set of field equations

$$\begin{aligned} E_t^t &= \frac{(D-2)rf' + (D-2)(D-3)f + \Lambda r^2 + q'^2 r^2 + f n_{z_{(D-3)}}^2 - 2f n_{z_{(D-3)}} s' + f s'^2}{2r^2} \equiv 0, \\ E_t^{z_{(D-3)}} &= E_{z_{(D-3)}}^t = q' [n_{z_{(D-3)}} - s'] \equiv 0, \\ E_r^r &= \frac{1}{2r^2} \left\{ (D-2)rf' + (D-2)(D-3)f + \Lambda r^2 - \phi'^2 f [2(D-2)^2 f' r + (D-2)^2 (D-3)f + (D-2)f'' r^2 + 6\Lambda r^2] - q'^2 r^2 \right. \\ &\quad \left. - f n_{z_{(D-3)}}^2 + 2f n_{z_{(D-3)}} s' - f s'^2 + (D-4)f\phi'^2 [q'^2 r^2 + 2f s' n_{z_{(D-3)}} - f s'^2 - f n_{z_{(D-3)}}^2] \right\} \equiv 0, \\ E_{\theta_1}^{\theta_1} &= E_{\theta_2}^{\theta_2} = \dots = E_{\theta_\ell}^{\theta_\ell} = \frac{1}{2r^2} \left[2(D-3)rf' + (D-4)(D-3)f + r^2 f'' + \Lambda r^2 - r^2 q'^2 - 2f s' n_{z_{(D-3)}} + f s'^2 + f n_{z_{(D-3)}}^2 \right] \equiv 0, \\ E_{z_1}^{z_1} &= E_{z_2}^{z_2} = \dots = E_{z_{D-\ell-2}}^{z_{D-\ell-2}} = \frac{1}{2r^2} \left[2(D-3)rf' + (D-4)(D-3)f + r^2 f'' + \Lambda r^2 - r^2 q'^2 + 2f s' n_{z_{(D-3)}} - f s'^2 - f n_{z_{(D-3)}}^2 \right] \equiv 0, \end{aligned} \quad (13)$$

where $f' = \frac{df(r)}{dr}$, $s' = \frac{ds(r)}{dr}$, $q' = \frac{dq(r)}{dr}$ and $n_{z_{(D-3)}} = \frac{dn(z_{(D-3)})}{dz_{(D-3)}}$ with $q(r)$, $n(z_{(D-3)})$ and $s(r)$ being three unknown functions related to the electric and magnetic charges of the black hole. These are usually defined from the general form of the vector potential

$$V = q(r)dt + n(z_{(D-3)})dr + s(r)dz_{(D-3)}. \quad (14)$$

Notably, for vanishing $n(z_{(D-3)})$ and $s(r)$, one can generate D -dimension charged electric solutions in the mimetic gravitational theories. However, for non-vanishing $q(r)$, $n(z_{(D-3)})$ and $s(r)$, one expects rich physical properties to showup. We solve the field equations (13) as follows

$$\begin{aligned} q(r) &= \frac{c_1}{r^{D-3}}, \quad n(z_{(D-3)}) = c_2 z_{(D-3)}, \quad s(r) = c_3 r, \quad f(r) = -\frac{\Lambda r^2}{(D-1)(D-2)} + \frac{c_4}{r^{D-3}} + \frac{(D-3)c_1^2}{(D-2)r^{2(D-3)}}, \\ \phi(r) &= \pm \left[f_1(r) + c_4 \ln \sqrt{r - c_4/2 + f_1(r)} \right] \delta_4^4 + c_5, \quad \text{with} \quad f(r) = \frac{1}{f_1(r)} \left(2f_1(r) - (r - c_4) \tan^{-1} \left[\frac{f_2(r)}{2f_1(r)} \right] \right), \\ f_1(r) &= \sqrt{r^2 - c_4 r}, \quad f_2 = 2r - c_4. \end{aligned} \quad (15)$$

where c_i , $i = 1 \cdots 5$ are constants. It is worth to mention that Eq. (15) is an exact solution of Maxwell-mimetic gravitational theory given by Eqs. (5) and (6) in addition to the trace which given by Eq. (7). As clear from the obtained solution (15) that the mimetic field $\phi(r)$ is a function of the radial coordinate r in the 4-dimensional spacetime case, while in the case $D > 4$ the mimetic field becomes constant. Plugging the solution (15) into the spacetime metric (11), we write

$$\begin{aligned} ds^2 &= - \left\{ -\frac{\Lambda r^2}{(D-1)(D-2)} + \frac{c_4}{r^{D-3}} + \frac{(D-3)c_1^2}{(D-2)r^{2(D-3)}} \right\} dt^2 + \left\{ -\frac{\Lambda r^{D-1}}{(D-1)(D-2)} + \frac{c_4}{r^{D-3}} + \frac{(D-3)c_1^2}{(D-2)r^{2(D-3)}} \right\}^{-1} dr^2 \\ &+ r^2 \left(\sum_{i=2}^{\ell} d\theta_i^2 + \sum_{k=2}^{D-\ell-2} dz_k^2 \right). \end{aligned} \quad (16)$$

As clear the spacetime (16) is asymptotically (A)dS, also it is obvious that the magnetic fields related to $n(z_{(D-3)})$ and $s(r)$ do not contribute to the metric.

IV. ROTATING BLACK STRING SOLUTIONS

In this section, we derive rotating solutions satisfying the field equations (5) and (6) of Maxwell-mimetic theory. In order to do so, we apply the coordinate transformations

$$\theta'_i = -\Xi \theta_i + \frac{\omega_i}{\lambda^2} t, \quad t' = \Xi t - \sum_{i=1}^{\ell} \omega_i \theta_i, \quad (17)$$

where ω_i , $i \geq 1$ is the number of rotation parameters and Ξ is defined as

$$\Xi := \sqrt{1 + \sum_{i=1}^{\ell} \frac{\omega_i^2}{\lambda^2}}.$$

The rotation group in D -dimensions is so $(D-1)$ and the independent number of the rotation parameters for a localized object is equal to the number of Casimir operators, which is $[(D-1)/2]$, where $[y]$ is the integer part of y . Applying the transformation (17) to the metric (16), we get

$$ds^2 = -f(r) \left[\Xi dt' - \sum_{i=1}^{\ell} \omega_i d\theta'_i \right]^2 + \frac{dr^2}{f(r)} + \frac{r^2}{\lambda^4} \sum_{i=1}^{\ell} [\omega_i dt' - \Xi \lambda^2 d\theta'_i]^2 + r^2 \sum_{k=1}^{D-\ell-2} dz_k^2 - \frac{r^2}{\lambda^2} \sum_{i < j} (\omega_i d\theta'_j - \omega_j d\theta'_i)^2, \quad (18)$$

where $f(r)$ is given by Eq. (15). We note that the static configuration (16) can be recovered as a special case when the rotation parameters ω_ℓ are made to vanish. Also, it is important to mention that the vanishing of the quantities c_1 and c_4 leads to an odd AdS spacetime. On the other hand, it is easy task to show that the limiting metric is a Minkowski spacetime, since all curvature components vanish identically.

In general the coordinate transformation (17) is admitted only locally [109, 110], since it relates time to periodic coordinate θ_ℓ . On other words, the spacetimes (18) and (11) can be locally mapped into each other but not globally and thus they are different. This has been discussed in more detail in [111], for similar coordinate transformation, it has been shown that if the first Betti number of a manifold has a non-vanishing value, then there are no global diffeomorphisms can connect the two spacetimes. Therefore, the manifold parameterized globally by the rotation parameters ω_ℓ is different from the static spacetime. The solution (15) shows that the first Betti number is one, which characterizes the cylindrical or toroidal horizons.

V. NEW BLACK HOLES WITH NONLINEAR ELECTRODYNAMICS IN MIMETIC GRAVITY

In this section, we consider the mimetic theory with nonlinear electrodynamics in the presence of a cosmological constant. Therefore, we take the action

$$\mathcal{S}_{NL} := \frac{1}{2\chi} \int d^D x \sqrt{-g(\bar{g}_{\mu\nu}, \phi)} \{R(\bar{g}_{\mu\nu}, \phi) - 2\Lambda\} - \int d^D x \sqrt{-g(\bar{g}_{\mu\nu}, \phi)} \mathcal{L}(F), \quad (19)$$

where $\mathcal{L}(F)$ is the Lagrangian of the nonlinear electrodynamics. Alternatively, we could reexpress $\mathcal{L}(F)$ in terms of Legendre transformation

$$Q = 2F \mathcal{L}_F - \mathcal{L}, \quad \text{where} \quad \mathcal{L}_F = \frac{\partial \mathcal{L}}{\partial F}. \quad (20)$$

It is useful to define the second-order tensor

$$P_{\mu\nu} = \mathcal{L}_F F_{\mu\nu}, \quad (21)$$

where the linear Maxwell field is recovered by setting $\mathcal{L}_F = 1$. As clear from the above that Q is a function of P , where [112, 113]

$$P = \frac{1}{4} P_{\mu\nu} P^{\mu\nu} = [\mathcal{L}_F]^2 F. \quad (22)$$

Thus, we have

$$dQ = \frac{1}{\mathcal{L}_F} d\{[\mathcal{L}_F]^2 F\} = Q_P dP, \quad \text{where} \quad Q_P = \frac{\partial Q}{\partial P}.$$

Varying the action of Eq. (19) with respect to the physical metric, one can write the gravitational field equations

$$E_\mu{}^\nu = G_\mu{}^\nu + \frac{1}{2} \delta_\mu^\nu \Lambda - \overset{\text{NL}}{\mathfrak{T}}_\mu{}^\nu - \widetilde{\mathfrak{T}}_\mu{}^\nu \equiv 0, \quad (23)$$

and Maxwell field equations of the nonlinear electrodynamics [113]

$$\partial_\nu (\sqrt{-g} P^{\mu\nu}) = 0. \quad (24)$$

In the above, the energy-momentum tensor,

$$\overset{\text{NL}}{\mathfrak{T}}_\mu{}^\nu = 2(Q_P P_{\mu\alpha} P^{\nu\alpha} - \delta_\mu^\nu [2P Q_P - Q]), \quad (25)$$

whose a non-vanishing trace. In addition, the mimetic field contributes in the field equations as

$$\widetilde{\mathfrak{T}}_{\mu\nu} = -\left(G + \frac{D}{2} \Lambda - \overset{\text{NL}}{\mathfrak{T}}\right) \partial_\mu \phi \partial_\nu \phi. \quad (26)$$

Similar to the linear case, the energy-momentum tensors, $\overset{\text{NL}}{\mathfrak{T}}_{\mu\nu}$ and $\widetilde{\mathfrak{T}}_{\mu\nu}$ are conserved, i.e. $\nabla^\mu \overset{\text{NL}}{\mathfrak{T}}_{\mu\nu} = 0 = \nabla^\mu \widetilde{\mathfrak{T}}_{\mu\nu}$. Using Eqs. (1) and (26), we write the continuity equation of the mimetic field

$$\nabla^\kappa \left(\left[G + \frac{D}{2} \Lambda - \overset{\text{NL}}{\mathfrak{T}} \right] \partial_\kappa \phi \right) \equiv \frac{1}{\sqrt{-g}} \partial_\kappa \left(\sqrt{-g} \left[G + \frac{D}{2} \Lambda - \overset{\text{NL}}{\mathfrak{T}} \right] g^{\kappa\sigma} \partial_\sigma \phi \right) = 0. \quad (27)$$

It is useful to give the trace of Eq. (23)

$$\left(G + \frac{D}{2} \Lambda - \overset{\text{NL}}{\mathfrak{T}} \right) (1 + g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi) = 0. \quad (28)$$

As clear, Eq. (28) is satisfied identically, if Eq. (1) is used. Since $G \neq 0$ or $\overset{\text{NL}}{\mathfrak{T}} \neq 0$, the mimetic theory at hand has non-trivial solutions and the conformal degree of freedom, remarkably, provides a dynamical quantity [35].

We apply the field equations (23) and (24) to the spacetime metric (11), where the non-vanishing components are

$$\begin{aligned}
E_t^t &\equiv \frac{(D-2)rf' + (D-2)(D-3)f + \Lambda r^2 - 2r^2Q(r)}{2r^2} = 0, \\
E_r^r &\equiv \frac{1}{2fr^2} \left\{ (D-2)rf' + (D-2)(D-3)f + \Lambda r^2 - \phi'^2 f [2(D-2)^2 f' r + (D-2)^2 (D-3)f + (D-2)f'' r^2 + D\Lambda r^2 - 2Dr^2Q(r)] \right. \\
&\quad \left. - 2r^2Q(r) - 2(D-2)\phi'^2 f \phi' \varphi''^{-1} r^2 Q(r)' \right\} = 0, \\
E_{\theta_1}^{\theta_1} &= E_{\theta_2}^{\theta_2} = \dots = E_{\theta_\ell}^{\theta_\ell} = E_{z_1}^{z_1} = E_{z_2}^{z_2} = \dots = E_{z_{D-\ell-2}}^{z_{D-\ell-2}} \\
&\equiv \frac{1}{2r^2\varphi''} \left[2(D-3)rf' \varphi'' + \varphi'' \{ (D-4)(D-3)f + r^2 f'' + \Lambda r^2 - 2r^2Q(r) \} + 2r^2 \varphi' Q' \right] = 0.
\end{aligned} \tag{29}$$

In the above, $Q(r)$ is an arbitrary function representing nonlinear electrodynamics, and $\varphi(r)$ is an unknown function reproducing D -dimension electric charge in the vector potential where $P = d\mathcal{V}$ and $\mathcal{V} = \mathcal{V}_\mu dx^\mu$ being the gauge potential 1-form which defined in the non-linear case as

$$\mathcal{V} = \varphi(r)dr. \tag{30}$$

The general solution of the system of differential equations (29) have the form

$$\begin{aligned}
Q(r) &= \frac{c_6}{r^{2(D-2)}} + \frac{c_7}{r^{2(D-3)}} + \frac{c_7^2}{r^{3(D-2)}}, \quad \phi(r) = \left[f_1(r) + c_9 \ln \sqrt{r - c_9/2 + f_1(r)} \right], \quad \text{with } f_1(r) = \sqrt{r^2 - c_9 r}, \\
\varphi(r) &= \frac{c_8}{r^{D-3}}, \quad f(r) = \frac{\Lambda r^2}{(D-1)(D-2)} + \frac{c_9}{r^{D-3}} - \frac{2c_6}{(D-2)(D-3)r^{2(D-3)}} - \frac{2c_7}{(D-2)^2 r^{2D-5}} - \frac{4c_7^2}{(D-2)(2D-5)r^{(3D-8)}},
\end{aligned} \tag{31}$$

where c_i , $i = 6 \dots 9$ are constants. From the above solutions, we can conclude: In the solution (31) as clear from the $Q(r)$ function that the 4-dimension nonlinear electrodynamics consists of monopole, dipole and quadrupole terms. Interestingly, the constant associated with the monopole is different from that appears with the dipole and quadrupole terms. In this case, one can get the form of the nonlinear electrodynamics by setting the constant $c_7 = 0$. On the other hand, the mimetic field $\phi(r)$ is constant in this case.

It is worth to mention that Eq. (31) is an exact solution to Maxwell-mimetic gravitational theory that is given by Eqs. (23) and (24). Then, we write explicitly the line elements corresponding to the above solution as

$$\begin{aligned}
ds^2 &= - \left\{ \frac{\Lambda r^2}{(D-1)(D-2)} + \frac{c_9}{r^{D-3}} - \frac{2c_6}{(D-2)(D-3)r^{2(D-3)}} - \frac{2c_7}{(D-2)^2 r^{2D-5}} - \frac{4c_7^2}{(D-2)(2D-5)r^{(3D-8)}} \right\} dt^2 \\
&\quad + \left\{ \frac{\Lambda r^2}{(D-1)(D-2)} + \frac{c_9}{r^{D-3}} - \frac{2c_6}{(D-2)(D-3)r^{2(D-3)}} - \frac{2c_7}{(D-2)^2 r^{2D-5}} - \frac{4c_7^2}{(D-2)(2D-5)r^{(3D-8)}} \right\}^{-1} dr^2 \\
&\quad + r^2 \left(\sum_{i=2}^{\ell} d\theta_i^2 + \sum_{k=2}^{D-\ell-2} dz_k^2 \right),
\end{aligned} \tag{32}$$

One can easily recognize that the spacetime (32) is asymptotically (A)dS.

Similar to what has been done in Sec. IV, we add angular momentum to the spacetime (32) by applying the transformation (17). Therefore, the metric becomes

$$ds^2 = -f(r) \left[\Xi dt' - \sum_{i=2}^{\ell} \omega_i d\theta'_i \right]^2 + \frac{dr^2}{f(r)} + \frac{r^2}{\lambda^4} \sum_{i=1}^{\ell} [\omega_i dt' - \Xi \lambda^2 d\theta'_i]^2 + r^2 \sum_{k=1}^{D-\ell-2} dz_k^2 - \frac{r^2}{\lambda^2} \sum_{i < j}^{\ell} (a_i d\phi'_j - a_j d\phi'_i)^2, \tag{33}$$

where $f(r)$ is given by Eq. (31). We note that coupling gravity and nonlinear electrodynamics has been studied in gravitational theories other than mimetic gravity, c.f [114, 115]. In [115], GR gravity is coupled to nonlinear electrodynamics to study thermodynamics of magnetically charged black holes. Also, in [114], quadratic curvature gravity is coupled with nonlinear electrodynamics to obtain regular black hole solutions, in that work the perturbative solution has been derived up to first order, where the regularity of the obtained solution depends on two free parameters of the model. In both models, nonlinear electrodynamic contribution has been assumed to have a particular form to reduce to RN asymptotically, then the solutions have been adjusted to satisfy the field equations. This is in contrast to the model at hand, where the solutions, namely (31), have been obtained directly from the equations of motion without pre-assumptions of the nonlinear electrodynamics contribution.

VI. FEATURES OF THE BLACK HOLE SOLUTIONS

Now we are going to discuss some relevant features of the charged black hole solutions presented in previous sections for the linear and nonlinear cases.

Firstly, in the linear Maxwell field case, the metric of the charged black hole solution (16) takes the form

$$ds^2 = - \left[r^2 \Lambda_{eff} - \frac{m}{r^{D-3}} + \frac{(D-3)q^2}{(D-2)r^{2(D-3)}} \right] dt^2 + \frac{dr^2}{\left[r^2 \Lambda_{eff} - \frac{m}{r^{D-3}} + \frac{(D-3)q^2}{(D-2)r^{2(D-3)}} \right]^{-1}} + r^2 \left(\sum_{i=2}^{\ell} d\theta_i^2 + \sum_{k=2}^{D-\ell-2} dz_k^2 \right), \quad (34)$$

where $\Lambda_{eff} = -\frac{\Lambda}{(D-1)(D-2)}$, $c_1 = q$ and $c_4 = m$. The above equation shows clearly that the metric in the linear case is the Reissner-Nordström solution which behaves asymptotically as dS background. By taking the limit $q \rightarrow 0$, we get the dS non-charged black holes. However, the horizons of the metric (34) are given by the real positive roots of $\Gamma(r) = 0$, where $\Gamma(r) = r^{4(D-3)}\Lambda - mr^{D-3} + \frac{(D-3)q^2}{(D-2)}$, see [116]. For the model at hand, namely the spacetime metric (34), taking $x = r^2$, we find that the constraint $\Gamma(x) = 0$ gives two positive real roots in the 4-dimension case, one of the roots represents the black hole event (inner) horizon, r_b and the other one represents a cosmological (outer) horizon, r_c . Similarly, in the 5-dimension case, the constraint $\Gamma(x) = 0$ gives three positive roots, i.e. the solution has three horizons. In general, in the D -dimension case, the constraint $\Gamma(x) = 0$ gives $(D-2)$ horizons.

Secondly, in the the nonlinear Maxwell field case, the spacetime metrics of the charged black hole solutions (32) takes the form

$$ds^2 = - \left\{ \Lambda_{eff} r^2 - \frac{m}{r^{D-3}} + \frac{q^2}{(D-2)(D-3)r^{2(D-3)}} + \frac{q_1}{(D-2)^2 r^{2D-5}} + \frac{q_1^2}{4(D-2)(2D-5)r^{3D-8}} \right\} dt^2 \\ + \left\{ \Lambda_{eff} r^2 - \frac{m}{r^{D-3}} - \frac{q^2}{(D-2)(D-3)r^{2(D-3)}} + \frac{q_1}{(D-2)^2 r^{2D-5}} + \frac{q_1^2}{4(D-2)(2D-5)r^{3D-8}} \right\}^{-1} dr^2 \\ + r^2 \left(\sum_{i=2}^{\ell} d\theta_i^2 + \sum_{k=2}^{D-\ell-2} dz_k^2 \right), \quad (35)$$

where $c_9 = -m$, $c_6 = -q^2/2$ and $c_7 = -q_1/2$. Equation (35) shows clearly that the metric of the charged black hole in the nonlinear Maxwell case is different from RN black hole. This difference is due to the existence of the dipole and quadrupole terms that are related by the constant q_1 . However, in the case that $q_1 = 0$, the solution reduces to the RN case. It is of interest to note that all the charged terms appear in Eq. (35) are reproduced from the arbitrary function, $Q(r)$, which characterizes the nonlinear electrodynamics. By taking the limit $q_1 \rightarrow 0$, the solution goes to the (A)dS charged black hole of the linear case. Additionally, we investigate the number of horizons of the spacetime metric (35). In the 4-dimension case, by taking $x = r^3$, similar to the linear electrodynamics case, we find that the constraint $\Gamma(x) = 0$ has two positive real roots, one for the black hole even horizon, r_b , and the other is for a cosmological horizon, r_c . In the 5-dimension case, the constraint $\Gamma(x) = 0$ gives three positive roots and in the general D -dimension case, the constraint $\Gamma(x) = 0$ gives $(D-2)$ horizons.

Next we discuss some physical properties of the solutions (15) and (31) by investigating the singularity behaviours and the their stabilities.

A. Visualization of black holes' singularities

We investigate the physical singularities by calculating some the curvature invariants. In both cases, linear and nonlinear electrodynamics, see (15) and (31), it is clear that the function $f(r)$ is irregular when $f_1(r) = 0$. Therefore, it is worth to investigate the regularity of the solutions at $f_1(r) = 0$. For the solutions (15), we evaluate the scalar invariants

$$R^{\mu\nu\lambda\rho} R_{\mu\nu\lambda\rho} = \frac{L_1(r)}{r^{4(D-2)}}, \quad R^{\mu\nu} R_{\mu\nu} = \frac{L_2(r)}{r^{4(D-2)}}, \quad R = \frac{L_3(r)}{r^{2(D-2)}}, \quad (36)$$

where $R^{\mu\nu\lambda\rho} R_{\mu\nu\lambda\rho}$, $R^{\mu\nu} R_{\mu\nu}$, R are the Kretschmann scalars, the Ricci tensors square, the Ricci scalars, respectively, also $L_i(r)$ are some polynomial functions in the radial coordinate r . Equation (36) shows that the solutions, at $r = 0$, have true singularities.

For the solutions (31), we evaluate the invariants

$$R^{\mu\nu\lambda\rho} R_{\mu\nu\lambda\rho} = \left(\frac{L_4(r)}{r^{6(D-2)}} \right), \quad R^{\mu\nu} R_{\mu\nu} = \left(\frac{L_5(r)}{r^{6(D-2)}} \right), \quad R = \left(\frac{L_6(r)}{r^{3(D-2)}} \right). \quad (37)$$

As clear all invariants have true singularities at $r = 0$. Remarkably, at the limit $r \rightarrow 0$, the behaviours of the Kretschmann scalars, the Ricci tensor square and the Ricci scalars of the nonlinear charged black hole solutions are given by $K = R_{\mu\nu}R^{\mu\nu} \sim r^{-6(D-2)}$, $R \sim r^{-3(D-2)}$ in contrast with the solutions of the linear Maxwell mimetic theory which have $K = R_{\mu\nu}R^{\mu\nu} \sim r^{-4(D-2)}$ and $R \sim r^{-2(D-2)}$. This shows clearly that the singularities in the nonlinear electrodynamics case is stronger than that has been obtained in the linear Maxwell mimetic gravity case. Notably, one may check if geodesics are extendible beyond these regions. According to Tipler and Królak [117, 118], this indicates the strength of the singularity. This topic will be discussed in forthcoming studies.

B. Energy conditions

Energy conditions provide important tools to examine and better understand cosmological models and/or strong gravitational fields. We are interested in the study of energy conditions in the nonlinear electrodynamics case, since the linear case is well known in the general relativity theory. As mentioned before, we focus on the nonlinear electrodynamics black hole solutions (31). The Energy conditions are classified into four categories: The strong energy (SEC), the weak energy (WEC), the null energy (NEC) and the dominant energy conditions [119, 120]. To fulfill those, the inequalities below must be satisfied

$$\begin{aligned} SEC : \rho + p_r &\geq 0, \quad \rho + p_r + 2p_t \geq 0, \\ WEC : \rho &\geq 0, \quad \rho + p_r \geq 0, \quad \rho + p_t \geq 0, \\ NEC : \rho + p_r &\geq 0, \quad \rho + p_t \geq 0, \\ DEC : \rho &\geq |p_r|, \quad \rho \geq |p_t|, \end{aligned} \quad (38)$$

where $\mathfrak{T}_0^{\text{NL}} = \rho$, $\mathfrak{T}_1^{\text{NL}} = p_r$ and $\mathfrak{T}_2^{\text{NL}} = \mathfrak{T}_3^{\text{NL}} = p_t$ are the density, radial and tangential pressures, respectively. Straightforward calculations of black hole solutions (31), in 4-dimensions, give

$$\begin{aligned} SEC : \rho + p_r &= \frac{4q^2r^2 - 4q_1r + q_1^2}{8r^6} > 0, \quad \rho + p_r + 2p_t = -\frac{q_1(2r + q_1)}{8r^6} < 0, \\ WEC : \rho &= \frac{4q^2r^2 - 4q_1r + q_1^2}{16r^6} > 0, \quad \rho + p_r = \frac{4q^2r^2 - 4q_1r + q_1^2}{8r^6} > 0, \quad \rho + p_t = \frac{8q^2r^2 + 2q_1r - q_1^2}{16r^6} > 0, \\ NEC : \rho + p_r &= \frac{4q^2r^2 - 4q_1r + q_1^2}{8r^6} > 0, \quad \rho + p_t = \frac{8q^2r^2 + 2q_1r - q_1^2}{8r^6} > 0, \\ DEC : \rho &\geq |p_r| \quad (\text{satisfied}), \quad \rho \geq |p_t| \quad (\text{satisfied}). \end{aligned} \quad (39)$$

This shows that the SEC is the only condition that is not satisfied. Remarkably, the SEC breaking is due to the contribution of the charge q_1 which characterizes the nonlinear function $Q(r)$. This shows clearly how the nonlinear contribution of the charge q_1 strengthens the singularity as discussed in the previous subsection. The SEC has a relatively clear geometrical interpretation which is the convergence of timelike geodesics which is manifestly related to the construction of singularities. We remind that the mimetic theory is stable if the density of the mimetic field is positive, which favors the solutions with de Sitter backgrounds in presence of a positive cosmological constant as in the work at hand [65] (see also [59]). In conclusion, the validity of the WEC in-return guarantees the positivity of the energy density in agreement with stability condition we just mentioned.

VII. THERMODYNAMICAL STABILITY AND PHASE TRANSITION

Black hole thermodynamics is one of the most exciting topics in physics, since it subjugates the laws of thermodynamics to understand gravitational/quantum physics of black holes. Two main approaches have been proposed to extract thermodynamical quantities of black holes: The first approach was given by Gibbons and Hawking [121, 122] to study thermal properties of the Schwarzschild black hole solution using the Euclidean continuation. The second approach is to identify gravitational surface and then define temperature then study the stability of the black hole [123–125]. Here in this study, we follow the second approach to study the thermodynamics of the (A)dS black holes derived in Eq. (34) and (35) then discussed their stability. These black holes are portrayed by the mass, m , the charges (monopole, q , dipole and quadrupole, q_1) and also by the cosmological constant Λ .

One can calculate the horizons of the linear, Eq. (34), and the nonlinear, Eq. (35), electrodynamics cases by finding the roots of $f(r) = 0$. In 4-dimensional spacetime, these can be seen in Figs. 1(a) and 1(b) for particular values of the model parameters. The plots show the two roots of $f(r)$ which determine the black hole r_b and the cosmological r_c horizons of the solutions at hand which in agreement with the results of Sec. VI. We note that, in the linear case, for $m > 0$, $q > 0$ and $\Lambda > 0$, we find that these




FIG. 1: Schematic plots of the function $f(r)$ which characterizes the event horizons by setting $f(r) = 0$: (a) For the linear Maxwell mimetic gravity theory, the function $f(r)$ is given by (15); (b) For the nonlinear electrodynamics mimetic gravity theory, the function $f(r)$ is given by (31). Also the plots indicate stronger singularities, as $r \rightarrow 0$, in the nonlinear electrodynamics case.

two roots are possible when $m > m_{min} = \frac{2}{3}6^{1/4}q^{3/2}\Lambda^{1/4}$. Interestingly, when $m = m_{min}$, we determine the degenerate horizons $r_{dg} = (q/\sqrt{6\Lambda})^{1/2}$ at which $r_b = r_c$, that is Nariai black hole. The thermodynamics of Nariai black hole has been studied in several works, c.f. [115, 126, 127]. Otherwise, when $m < m_{min}$, there is no black hole. This can be shown in Fig. 2(a), while Fig. 2(b) is showing the dependence of the degenerate horizon on the charge in the linear electrodynamics case. Similarly, the degenerate horizon can be determined in the nonlinear electrodynamics case. In 4-dimensional spacetime, the solutions which have two horizons similar to our model can be obtained for Schwarzschild-dS and Kerr-Schild class [128–130], RN black holes surrounded with quintessence [131], minimal model of a regular black hole [132], and also in the case of Bardeen black holes which are spherically symmetric solution of the noncommutative geometry [115, 127, 133, 134]. In our calculations, we use positive values of the cosmological constant, since we are interested in the double horizons solutions. However, it is worth to mention that negative cosmological constant produces a pattern similar to the case of two vacuum scales spacetime which connects two de Sitter vacua at $r \rightarrow 0$ and $r \rightarrow \infty$. The later is characterized by at most three horizons [135, 136]. However, in our case we have exactly one horizon.

The black hole thermodynamical stability is related to the sign of its heat capacity C_h . In the following, we analyze the thermal stability of the black hole solutions via the behaviour of their heat capacities [104, 137, 138]

$$C_h = \frac{dE_h}{dT_h} = \frac{\partial m}{\partial r_h} \left(\frac{\partial T}{\partial r_h} \right)^{-1}, \quad (40)$$

where E_h is the energy. If the heat capacity $C_h > 0$ ($C_h < 0$), the black hole is thermodynamically stable (unstable), respectively. To a better understanding of this process, we assume that at some point and due to thermal fluctuations, the black hole absorbs more radiation than it emits, which means that its heat capacity is positive. This means that the mass of the black hole indefinitely increases. On the contrary, the black hole emits more radiation than it absorbs, which means the heat capacity is negative. This means that the black hole mass indefinitely decreases until it disappears completely. Thus, black holes with negative heat capacities are thermally unstable.

In order to evaluate Eq. (40), it requires us to derive analytical formulae of $m_h \equiv m(r_h)$ and $T_h \equiv T(r_h)$. Firstly, we calculate the black hole mass within an even horizon r_h . We set $f(r_h) = 0$, then we get

$$\begin{aligned} m_{h_{Eq.(34)}} &= r_h^{(D-3)} \left(r_h^2 \Lambda_{eff} + \frac{(D-3)q^2}{(D-2)r_h^{2(D-3)}} \right), \\ m_{h_{Eq.(35)}} &= r_h^{(D-3)} \left(r_h^2 \Lambda_{eff} + \frac{q^2}{(D-2)(D-3)r_h^{2(D-3)}} + \frac{q_1}{(D-2)^2 r_h^{2D-5}} + \frac{q_1^2}{4(D-2)(2D-5)r_h^{(3D-8)}} \right). \end{aligned} \quad (41)$$

The above equations show that the black hole total mass is given as a function of the horizon radius and the charge. One can also find the degenerate horizon by setting $\partial m_h / \partial r_h = 0$, in 4 dimensions; this gives $r_{dg} = (q/\sqrt{6\Lambda})^{1/2}$ as previously obtained.




FIG. 2: Schematic plots of the degenerate horizons in the linear Maxwell mimetic gravity: (a) The plot of $f(r)$ shows the black hole event horizon, r_b , and the cosmological horizon, r_c , where $m > m_{min}$. At $m = m_{min}$ ($r_b = r_c$), the black hole has a degenerate horizon at r_{dg} . Otherwise, $m < m_{min}$ the black hole is naked; (b) The plot shows the dependence of the degenerate horizon on the charge q at fixed mass $m = m_{min}$. Here we take $D = 4$ and $\Lambda = 0.058$.

For fixed charge values in both linear and nonlinear charged black holes, we plot the horizon mass-radius relation as is depicted in Fig. 3(a). As seen from this figure, the horizon mass-radius relation is characterized by

$$m(r_h \rightarrow 0) \rightarrow \infty, \quad m(r_h \rightarrow \infty) \rightarrow \infty. \quad (42)$$

Also, one can find that there is a minimal mass at the degenerate horizon whereas the double horizons (event and cosmological) coincide. For larger masses, the double horizons are separated, while smaller masses do not show horizons. This confirms the results of Fig. 2(a). In this sense, we find that the model at hand shares some features with the minimal model of a regular black hole [132]. However, as shown by Eq. (42), there is no minimal length of the black hole event horizon as in the minimal model scenario.

The Hawking temperature of the black holes can be obtained by requiring the absence of singularity at the horizon in the Euclidean sector of the black hole solutions. Secondly, we obtain the associated temperature with the outer event horizon $r = r_h$ as [139]

$$T = \frac{\kappa}{2\pi}, \quad \text{where } \kappa \text{ is the surface gravity defined as } \kappa = \frac{f'(r_h)}{2}. \quad (43)$$

The Hawking temperatures associated with the black hole solutions (34) and (35) are

$$\begin{aligned} T_{h_{Eq.(34)}} &= \frac{1}{4\pi} \left(\frac{(D-1)(D-2)r_h^{2(D-2)}\Lambda_{eff} - q^2}{(D-2)r_h^{2D-5}} \right), \\ T_{h_{Eq.(35)}} &= \frac{1}{4\pi} \left\{ \frac{(D-1)(D-2)r_h^{2(D-2)}\Lambda_{eff} - q^2}{(D-2)r_h^{2D-5}} + \frac{(D-3)q_1}{(D-2)^2 r_h^{2(D-2)}} + \frac{(D-3)q_1^2}{4(D-2)(2D-5)r_h^{(3D-7)}} \right\} \end{aligned} \quad (44)$$

where T_h is the Hawking temperature at the event horizon. For linear and nonlinear electrodynamics, in the 4-dimensional spacetime, we plot the horizon temperatures-radius relation in Fig. 3(b) for particular values the black hole parameters. The figure shows that the horizon temperature T_h vanishes at the degenerate horizon $r_h = r_{dg}$. At $r_h < r_{dg}$, the horizon temperature goes below absolute zero forming an ultracold black hole. As noted earlier by Davies [91] that there is no obvious reasons from thermodynamics prevent a black hole temperature to go below absolute zero and turn it to naked singularity. In fact this is the case presented in Fig. 3(b) at $r_h < r_{min}$ region. However, this case of ultracold black hole is justified in the presence of phantom energy field [140], and explains the decreasing mass pattern in Fig. 3(a). Indeed, this case suits well with mimetic gravity theories whereas the SEC breaking indicate a phantom mimetic field. Also, it would be useful to examine some potential patterns in that case. At $r_h > r_{dg}$, the horizon temperature becomes positive. For values r_h larger enough, the horizon temperatures of both linear




FIG. 3: Schematic plots of some thermodynamical quantities of the black hole solutions: (a) The horizon mass-radius relation determines the minimal mass required to identify the degenerate horizon; (b) The temperature vanishes on the horizon radius r_h ; (c) The heat capacity shows the locally unstable event horizon which is characterized by the negative $C_h < 0$. Also, the nonlinear electrodynamics case shows a second-order phase transition as C_h diverges.

and nonlinear charged black holes behave similarly. Including the gravitational effect of thermal radiation, one can show that at some very high temperature T_{max} the radiation would become unstable and collapse to a black hole [103]. Hence, the pure AdS solution is only stable at temperatures $T < T_{max}$. Above T_{max} , only the heavy black holes would have stable configurations [103].

Next we evaluate the horizon heat capacity C_h , therefore we substitute Eqs. (41) and (44) into Eq. (40). Hence, we write

$$C_h = \frac{2\pi r_h^2 [(D-1)(D-2)r_h^{3D-4}\Lambda_{eff} - q^2 r_h^D]}{(D-1)(D-2)r_h^{2D}\Lambda + (D-5)r_h^4 q^2},$$

$$C_h = \frac{(2D-5)\pi r_h^{D-2} \left[[4(D-1)(D-2)r_h^{3(D-2)}\Lambda_{eff} - q_1^2] r_h^{2D-5} - 4q_1 r_h^{3D-8} \right] r_h^{2(D-3)} - 4q^2 r_h^{5D-13}}{r_h^{7D-19} [(D-5)[4(D-1)(D-2)r_h^2\Lambda + 4(2D-5)q^2 r_h^{-2(D-3)} - 8(D-3)r_h^{-2(D-3)}q_1] - (3D-7)(D-3)r_h^{-(3D-8)}q_1^2]}.$$
(45)

It is not easy to extract information directly from Eqs. (45), therefore we plot them in Fig. 3(c) for particular values of the black hole parameters. As we show on this figure, in both linear and nonlinear charged black holes, the horizon heat capacities vanish at the degenerate horizon r_{dg} as well as their horizon temperatures. In the linear electrodynamics, the heat capacity has negative values for $r_h < r_{dg}$, but it has positive values for $r_h > r_{dg}$. In the nonlinear electrodynamics, the heat capacity is positive for all values of the horizon radius r_h except for an intermediate region $r_c < r_h < r_{dg}$, it is negative and characterized by a second-order phase transition at r_c whereas the heat capacity has an infinite discontinuity.

In conclusion, in the linear charged black hole case, a typical pattern of the heat capacity has been obtained in literature c.f [141]. However, we find that the negative heat capacity region is associated with a positive temperature on the contrary to our case. In mimetic gravity with linear electrodynamics, the negative heat capacity region, i.e $r_h < r_{dg}$, is associated with a temperature below absolute zero. This can be justified in the presence of a phantom field as in our case whereas the SEC is not fulfilled, see Eqs. (39). At $r_h = r_{dg}$, the temperature on the event horizon is exactly zero as well as the heat capacity as shown on Figs 3(b) and 3(c). For the case $r_h > r_{dg}$, both temperature and heat capacity are positive and the solution is in a thermal equilibrium. Indeed, thermodynamics stability of the charged black hole in AdS spacetime has been widely studied in many theories, thermodynamics of Bardeen (regular) black holes c.f. [126], Schwarzschild-AdS in two vacuum scales case [142], and also similar work in the noncommutative geometry [114, 143–145]. Indeed, all these solutions are characterized by a second-order phase transition where the heat capacity has an infinite discontinuity similar to our case, while the heat capacity remains negative at $r_h > r_{dg}$. This is in contrast to our case, as clear from Fig. 3(c), whereas the heat capacity crosses to positive regions as r_h goes to larger values. This qualitative difference is due to the nonlinear electrodynamics contribution. As we have mentioned before, the nonlinear contribution of the model at hand is derived from the equations of motions in contrast to other models which pre-assume the form of the nonlinear terms to produce RN asymptotically.

VIII. SUMMARY AND PROSPECTIVES

Day after day dark matter is being confirmed by astrophysical and cosmological observations. Two main streams have been proposed to explain dark matter, modifications of Einstein's general relativity and modifications of the standard model by introducing new particle species. It has been shown that these two classes are not different after all [146]. In fact, every modified gravity model has new degrees of freedom besides the usual massless graviton. As we mentioned in the introduction that the mimetic gravity is a good candidate to explain CDM presence. This motivates us to explore the theory in astrophysics domain, by investigating possible new solutions of charged rotating black holes.

For this aim, we derive the field equations of Maxwell mimetic gravitational theory. We apply these field equations to D -manifold with ℓ angular coordinates, k -dimension Euclidean metric and one unknown function $f(r)$ of the radial coordinate. In addition, we use a generalized vector potential which includes three unknown functions, one of them is related to the electric charge and the other two functions are related to the magnetic field. In this context, we derive charged D -dimension black hole solutions that possess the mass and the electric charge of the black hole. These behave asymptotically as (A)dS. Then, we apply a coordinate transformation relating the angular coordinates and the temporal coordinate, which allows to derive D -dimension rotating charged black hole solutions.

Similarly, in the nonlinear electrodynamics coupled to mimetic gravity, we derive D -dimension charged black hole solutions. We apply the corresponding field equations which allow to obtain a new D -dimension charged black hole solutions. Interestingly, the new black holes show interesting physical properties, besides the monopole term the solutions contain nonlinear effects related to the dipole and quadrupole charges. The later terms are characterized by a common constant, so that its vanishing derives the solution to the linear case. We show that the asymptotic behaviours of this class of the black hole solutions behaves as (A)dS spacetime. Similar to the Maxwell electrodynamics case, we apply a coordinate transformation which allows to obtain new D -dimension rotating charged black hole solutions analytically.

Also, we calculate the invariants constructed from the curvature, namely Kretschmann invariants K , the Ricci tensors squared $R_{\mu\nu}R^{\mu\nu}$ and the Ricci scalars R , to study possible singularities of the obtained solutions. This shows that they have a true singularities at $r = 0$. However, in the Maxwell mimetic gravity, the asymptotic behaviour of these invariants are $K = R_{\mu\nu}R^{\mu\nu} \sim r^{-4(D-2)}$ and $R \sim r^{-2(D-2)}$. On the other hand, their asymptotic behaviours are $K = R_{\mu\nu}R^{\mu\nu} \sim r^{-6(D-2)}$ and $R \sim r^{-3(D-2)}$. This indicates that the nonlinear electrodynamics mimetic gravity produces singularities stronger than the Maxwell electrodynamics case. In addition, we calculate the number of horizons, in the linear (nonlinear) case the solutions have $D - 2$ horizons. This again proves that the singularities of the black hole solutions in the nonlinear case are stronger than the linear case. Then, we investigate the fulfilment of the energy conditions in the context of the obtained solutions. We verified the fulfilment of the WEC which guarantees the positivity of the energy density. Supported by the preliminary studies [59, 65], the obtained results are in favor of the absence of ghosts. However, we remind that the perturbation analysis of the present theory is a mandatory to fully investigate the ghost problem. Since here we focus on finding new solutions of charged black holes, we leave the perturbation analysis to be carried out separately elsewhere in the future.

Finally, for the linear and the nonlinear electrodynamics cases, we study the horizons showing that the solutions could have at most two horizons, a black hole event horizon r_b and a cosmological one r_c . We determine a minimum value of the black hole mass at which the two horizons coincide forming the degenerate horizon, above this minimum mass the black hole would have two horizons, below the minimum mass it there is no black hole. Also, we study thermodynamics and thermal phase transitions of the obtained black holes. In the linear electrodynamics case, the temperature drops below absolute zero when $r_h < r_{min}$ forming an ultracold black hole, while the heat capacity is being negative at these regions and so it is unstable. Similar conclusions can be derived for the nonlinear electrodynamics case, however it is characterized by a second-order phase transition whereas the heat capacity has an infinite discontinuity.

Acknowledgments

The work of GGLN is partially supported by the Egyptian Ministry of Scientific Research under project No. 24-2-12. Moreover, the work of KB was supported in part by the JSPS KAKENHI Grant Number JP 25800136 and Competitive Research Funds for Fukushima University Faculty (18RI009).

-
- [1] Adam G. Riess *et al.* (Supernova Search Team), “Observational evidence from supernovae for an accelerating universe and a cosmological constant,” *Astron. J.* **116**, 1009–1038 (1998), [arXiv:astro-ph/9805201 \[astro-ph\]](#).
 - [2] S. Perlmutter *et al.* (Supernova Cosmology Project), “Measurements of Omega and Lambda from 42 high redshift supernovae,” *Astrophys. J.* **517**, 565–586 (1999), [arXiv:astro-ph/9812133 \[astro-ph\]](#).

- [3] P. A. R. Ade *et al.* (Planck), “Planck 2015 results. XIII. Cosmological parameters,” *Astron. Astrophys.* **594** (2016), 10.1051/0004-6361/201525830, arXiv:1502.01589 [astro-ph.CO] .
- [4] Salvatore Capozziello, “Curvature quintessence,” *Int. J. Mod. Phys. ,* 483–492 (2002), arXiv:gr-qc/0201033 [gr-qc] .
- [5] Sean M. Carroll, Vikram Duvvuri, Mark Trodden, and Michael S. Turner, “Is cosmic speed - up due to new gravitational physics?” *Phys. Rev. ,* 043528 (2004), arXiv:astro-ph/0306438 [astro-ph] .
- [6] Shin’ichi Nojiri and Sergei D. Odintsov, “Where new gravitational physics comes from: M Theory?” *Phys. Lett. ,* 5–11 (2003), arXiv:hep-th/0307071 [hep-th] .
- [7] A. D. Dolgov and Masahiro Kawasaki, “Can modified gravity explain accelerated cosmic expansion?” *Phys. Lett. ,* 1–4 (2003), arXiv:astro-ph/0307285 [astro-ph] .
- [8] Takeshi Chiba, “ $1/R$ gravity and scalar - tensor gravity,” *Phys. Lett. ,* 1–3 (2003), arXiv:astro-ph/0307338 [astro-ph] .
- [9] Sean M. Carroll, Antonio De Felice, Vikram Duvvuri, Damien A. Easson, Mark Trodden, and Michael S. Turner, “The Cosmology of generalized modified gravity models,” *Phys. Rev. ,* 063513 (2005), arXiv:astro-ph/0410031 [astro-ph] .
- [10] Shin’ichi Nojiri and Sergei D. Odintsov, “Introduction to modified gravity and gravitational alternative for dark energy,” *Theoretical physics: Current mathematical topics in gravitation and cosmology. Proceedings, 42nd Karpacz Winter School, Ladek, Poland, February 6-11, 2006, eConf ,* 06 (2006), [Int. J. Geom. Meth. Mod. Phys.4,115(2007)], arXiv:hep-th/0601213 [hep-th] .
- [11] Richard P. Woodard, “Avoiding dark energy with $1/r$ modifications of gravity,” *The invisible universe: Dark matter and dark energy. Proceedings, 3rd Aegean School, Karfas, Greece, September 26-October 1, 2005, Lect. Notes Phys.* **720**, 403–433 (2007), arXiv:astro-ph/0601672 [astro-ph] .
- [12] Shin’ichi Nojiri and Sergei D. Odintsov, “Modified gravity and its reconstruction from the universe expansion history,” *Einstein’s legacy: From the theoretical paradise to astrophysical observations, J. Phys. Conf. Ser.* **66**, 012005 (2007), arXiv:hep-th/0611071 [hep-th] .
- [13] M. I. Wanas, “The other side of gravity and geometry: Antigravity and anticurvature,” *Adv. High Energy Phys.* **2012**, 752613 (2012).
- [14] Timothy Clifton, Pedro G. Ferreira, Antonio Padilla, and Constantinos Skordis, “Modified Gravity and Cosmology,” *Phys. Rept.* **513**, 1–189 (2012), arXiv:1106.2476 [astro-ph.CO] .
- [15] Salvatore Capozziello and Mariafelicia De Laurentis, “Extended Theories of Gravity,” *Phys. Rept.* **509**, 167–321 (2011), arXiv:1108.6266 [gr-qc] .
- [16] Kazuharu Bamba, Salvatore Capozziello, Shin’ichi Nojiri, and Sergei D. Odintsov, “Dark energy cosmology: the equivalent description via different theoretical models and cosmography tests,” *Astrophys. Space Sci.* **342**, 155–228 (2012), arXiv:1205.3421 [gr-qc] .
- [17] Shin’ichi Nojiri and Sergei D. Odintsov, “Modified gravity with negative and positive powers of the curvature: Unification of the inflation and of the cosmic acceleration,” *Phys. Rev. ,* 123512 (2003), arXiv:hep-th/0307288 [hep-th] .
- [18] S. Capozziello, Vincenzo F. Cardone, and A. Troisi, “Reconciling dark energy models with $f(R)$ theories,” *Phys. Rev. ,* 043503 (2005), arXiv:astro-ph/0501426 [astro-ph] .
- [19] Luca Amendola, David Polarski, and Shinji Tsujikawa, “Are $f(R)$ dark energy models cosmologically viable ?” *Phys. Rev. Lett.* **98**, 131302 (2007), arXiv:astro-ph/0603703 [astro-ph] .
- [20] Salvatore Capozziello, S. Nojiri, S. D. Odintsov, and A. Troisi, “Cosmological viability of $f(R)$ -gravity as an ideal fluid and its compatibility with a matter dominated phase,” *Phys. Lett. ,* 135–143 (2006), arXiv:astro-ph/0604431 [astro-ph] .
- [21] Jose A. R. Cembranos, “Dark Matter from R^2 -gravity,” *Phys. Rev. Lett.* **102**, 141301 (2009), arXiv:0809.1653 [hep-ph] .
- [22] Shin’ichi Nojiri and Sergei D. Odintsov, “Unified cosmic history in modified gravity: from $f(R)$ theory to Lorentz non-invariant models,” *Phys. Rept.* **505**, 59–144 (2011), arXiv:1011.0544 [gr-qc] .
- [23] Christian G. Boehmer, Atifah Mussa, and Nicola Tamanini, “Existence of relativistic stars in $f(T)$ gravity,” *Class. Quant. Grav.* **28**, 245020 (2011), arXiv:1107.4455 [gr-qc] .
- [24] Han Dong, Ying-bin Wang, and Xin-he Meng, “Extended Birkhoff’s Theorem in the $f(T)$ Gravity,” *Eur. Phys. J. ,* 2002 (2012), arXiv:1203.5890 [gr-qc] .
- [25] G. L. Nashed, “FRW in quadratic form of $f(T)$ gravitational theories,” *Gen. Rel. Grav.* **47**, 75 (2015), arXiv:1506.08695 [gr-qc] .
- [26] Kazuharu Bamba, G. G. L. Nashed, W. El Hanafy, and Sh. K. Ibraheem, “Bounce inflation in $f(T)$ Cosmology: A unified inflaton-quintessence field,” *Phys. Rev. ,* 083513 (2016), arXiv:1604.07604 [gr-qc] .
- [27] A. Paliathanasis, S. Basilakos, E. N. Saridakis, S. Capozziello, K. Atazadeh, F. Darabi, and M. Tsamparlis, “New Schwarzschild-like solutions in $f(T)$ gravity through Noether symmetries,” *Phys. Rev. ,* 104042 (2014), arXiv:1402.5935 [gr-qc] .
- [28] G. G. L. Nashed and W. El Hanafy, “Analytic rotating black hole solutions in N -dimensional $f(T)$ gravity,” *Eur. Phys. J. ,* 90 (2017), arXiv:1612.05106 [gr-qc] .
- [29] A. Awad, W. El Hanafy, G. G. L. Nashed, S. D. Odintsov, and V. K. Oikonomou, “Constant-roll Inflation in $f(T)$ Teleparallel Gravity,” *JCAP* **1807**, 026 (2018), arXiv:1710.00682 [gr-qc] .
- [30] A. Awad, W. El Hanafy, G. G. L. Nashed, and Emmanuel N. Saridakis, “Phase Portraits of general $f(T)$ Cosmology,” *JCAP* **1802**, 052 (2018), arXiv:1710.10194 [gr-qc] .
- [31] Valerio Faraoni and Salvatore Capozziello, *Beyond Einstein Gravity*, Vol. 170 (Springer, Dordrecht, 2011).
- [32] Kazuharu Bamba and Sergei D. Odintsov, “Inflationary cosmology in modified gravity theories,” *Symmetry* **7**, 220–240 (2015), arXiv:1503.00442 [hep-th] .
- [33] Yi-Fu Cai, Salvatore Capozziello, Mariafelicia De Laurentis, and Emmanuel N. Saridakis, “ $f(T)$ teleparallel gravity and cosmology,” *Rept. Prog. Phys.* **79**, 106901 (2016), arXiv:1511.07586 [gr-qc] .
- [34] S. Nojiri, S. D. Odintsov, and V. K. Oikonomou, “Modified Gravity Theories on a Nutshell: Inflation, Bounce and Late-time Evolution,” *Phys. Rept.* **692**, 1–104 (2017), arXiv:1705.11098 [gr-qc] .
- [35] Ali H. Chamseddine and Viatcheslav Mukhanov, “Mimetic Dark Matter,” *JHEP* **11**, 135 (2013), arXiv:1308.5410 [astro-ph.CO] .
- [36] Ali H. Chamseddine and Viatcheslav Mukhanov, “Nonsingular Black Hole,” *Eur. Phys. J. ,* 183 (2017), arXiv:1612.05861 [gr-qc] .
- [37] Ali H. Chamseddine and Viatcheslav Mukhanov, “Resolving Cosmological Singularities,” *JCAP* **1703**, 009 (2017), arXiv:1612.05860 [gr-qc] .

- [38] Nashed G. G. L. “Spherically symmetric black hole solution in mimetic gravity and anti-evaporation,” *International Journal of Geometric Methods in Modern Physics* **15**, 1850154-1056 (2018) .
- [39] Hassan Saadi, “A Cosmological Solution to Mimetic Dark Matter,” *Eur. Phys. J. ,* **14** (2016), [arXiv:1411.4531 \[gr-qc\]](#) .
- [40] Zahra Haghani, Tiberiu Harko, Hamid Reza Sepangi, and Shahab Shahidi, “Cosmology of a Lorentz violating Galileon theory,” *JCAP* **1505**, 022 (2015), [arXiv:1501.00819 \[gr-qc\]](#) .
- [41] E. H. Baffou, M. J. S. Houndjo, M. Hamani-Daouda, and F. G. Alvarenga, “Late time cosmological approach in mimetic $f(R, T)$ gravity,” *Eur. Phys. J. ,* **708** (2017), [arXiv:1706.08842 \[gr-qc\]](#) .
- [42] Ratbay Myrzakulov and Lorenzo Sebastiani, “Spherically symmetric static vacuum solutions in Mimetic gravity,” *Gen. Rel. Grav.* **47**, 89 (2015), [arXiv:1503.04293 \[gr-qc\]](#) .
- [43] Andrea Addazi and Antonino Marciano, “Evaporation and Antievaporation instabilities,” *Symmetry* **9**, 249 (2017), [arXiv:1710.07962 \[gr-qc\]](#) .
- [44] Artyom V. Astashenok and Sergei D. Odintsov, “From neutron stars to quark stars in mimetic gravity,” *Phys. Rev. ,* **063008** (2016), [arXiv:1512.07279 \[gr-qc\]](#) .
- [45] Ratbay Myrzakulov, Lorenzo Sebastiani, Sunny Vagnozzi, and Sergio Zerbini, “Static spherically symmetric solutions in mimetic gravity: rotation curves and wormholes,” *Class. Quant. Grav.* **33**, 125005 (2016), [arXiv:1510.02284 \[gr-qc\]](#) .
- [46] Eugeny Babichev and Sabir Ramazanov, “Gravitational focusing of Imperfect Dark Matter,” *Phys. Rev. ,* **024025** (2017), [arXiv:1609.08580 \[gr-qc\]](#) .
- [47] L. Sebastiani, S. Vagnozzi, and R. Myrzakulov, “Mimetic gravity: a review of recent developments and applications to cosmology and astrophysics,” *Adv. High Energy Phys.* **2017**, 3156915 (2017), [arXiv:1612.08661 \[gr-qc\]](#) .
- [48] Nathalie Deruelle and Josephine Rua, “Disformal Transformations, Veiled General Relativity and Mimetic Gravity,” *JCAP* **1409**, 002 (2014), [arXiv:1407.0825 \[gr-qc\]](#) .
- [49] Fang-Fang Yuan and Peng Huang, “Induced geometry from disformal transformation,” *Phys. Lett. ,* **120–124** (2015), [arXiv:1501.06135 \[gr-qc\]](#) .
- [50] Guillem Domenech, Shinji Mukohyama, Ryo Namba, Atsushi Naruko, Rio Saitou, and Yota Watanabe, “Derivative-dependent metric transformation and physical degrees of freedom,” *Phys. Rev. ,* **084027** (2015), [arXiv:1507.05390 \[hep-th\]](#) .
- [51] Shin’ichi Nojiri and Sergei D. Odintsov, “Mimetic $f(R)$ gravity: inflation, dark energy and bounce,” *Mod. Phys. Lett. ,* **1450211** (2014), [arXiv:1408.3561 \[hep-th\]](#) .
- [52] S. D. Odintsov and V. K. Oikonomou, “Unimodular Mimetic $f(R)$ Inflation,” *Astrophys. Space Sci.* **361**, 236 (2016), [arXiv:1602.05645 \[gr-qc\]](#) .
- [53] V. K. Oikonomou, “A note on Schwarzschildde Sitter black holes in mimetic $f(R)$ gravity,” *Int. J. Mod. Phys. ,* **1650078** (2016), [arXiv:1605.00583 \[gr-qc\]](#) .
- [54] Jibitesh Dutta, Wompherdeiki Khyllep, Emmanuel N. Saridakis, Nicola Tamanini, and Sunny Vagnozzi, “Cosmological dynamics of mimetic gravity,” *JCAP* **1802**, 041 (2018), [arXiv:1711.07290 \[gr-qc\]](#) .
- [55] Hassan Firouzjahi, Mohammad Ali Gorji, and Seyed Ali Hosseini Mansoori, “Instabilities in Mimetic Matter Perturbations,” *JCAP* **1707**, 031 (2017), [arXiv:1703.02923 \[hep-th\]](#) .
- [56] Katrin Hammer and Alexander Vikman, “Many Faces of Mimetic Gravity,” (2015), [arXiv:1512.09118 \[gr-qc\]](#) .
- [57] Ali H. Chamseddine, Viatcheslav Mukhanov, and Alexander Vikman, “Cosmology with Mimetic Matter,” *JCAP* **1406**, 017 (2014), [arXiv:1403.3961 \[astro-ph.CO\]](#) .
- [58] Leila Mirzaghali and Alexander Vikman, “Imperfect Dark Matter,” *JCAP* **1506**, 028 (2015), [arXiv:1412.7136 \[gr-qc\]](#) .
- [59] Masud Chaichian, Josef Kluson, Markku Oksanen, and Anca Tureanu, “Mimetic dark matter, ghost instability and a mimetic tensor-vector-scalar gravity,” *JHEP* **12**, 102 (2014), [arXiv:1404.4008 \[hep-th\]](#) .
- [60] O. Malaeb, “Hamiltonian Formulation of Mimetic Gravity,” *Phys. Rev. ,* **103526** (2015), [arXiv:1404.4195 \[gr-qc\]](#) .
- [61] Masooma Ali, Viqar Husain, Shohreh Rahmati, and Jonathan Ziprick, “Linearized gravity with matter time,” *Class. Quant. Grav.* **33**, 105012 (2016), [arXiv:1512.07854 \[gr-qc\]](#) .
- [62] Fabio Capela and Sabir Ramazanov, “Modified Dust and the Small Scale Crisis in CDM,” *JCAP* **1504**, 051 (2015), [arXiv:1412.2051 \[astro-ph.CO\]](#) .
- [63] Jibril Ben Achour, David Langlois, and Karim Noui, “Degenerate higher order scalar-tensor theories beyond Horndeski and disformal transformations,” *Phys. Rev. D* **93**, 124005 (2016), [arXiv:1602.08398 \[gr-qc\]](#) .
- [64] S. Ramazanov, F. Arroja, M. Celería, S. Matarrese, and L. Pilo, “Living with ghosts in Horava-Lifshitz gravity,” *JHEP* **06**, 020 (2016), [arXiv:1601.05405 \[hep-th\]](#) .
- [65] A. O. Barvinsky, “Dark matter as a ghost free conformal extension of Einstein theory,” *JCAP* **1401**, 014 (2014), [arXiv:1311.3111 \[hep-th\]](#) .
- [66] Shin’ichi Hirano, Sakine Nishi, and Tsutomu Kobayashi, “Healthy imperfect dark matter from effective theory of mimetic cosmological perturbations,” *JCAP* **1707**, 009 (2017), [arXiv:1704.06031 \[gr-qc\]](#) .
- [67] S. Nojiri, S. D. Odintsov, and V. K. Oikonomou, “Ghost-Free $f(R)$ Gravity with Lagrange Multiplier Constraint,” *Phys. Lett. ,* **44–49** (2017), [arXiv:1710.07838 \[gr-qc\]](#) .
- [68] S. Nojiri, S. D. Odintsov, and V. K. Oikonomou, “Ghost-free Gauss-Bonnet Theories of Gravity,” (2018), [arXiv:1811.07790 \[gr-qc\]](#) .
- [69] Genly Leon and Emmanuel N. Saridakis, “Dynamical behavior in mimetic $f(R)$ gravity,” *JCAP* **1504**, 031 (2015), [arXiv:1501.00488 \[gr-qc\]](#) .
- [70] Artyom V. Astashenok, Sergei D. Odintsov, and V. K. Oikonomou, “Modified GaussBonnet gravity with the Lagrange multiplier constraint as mimetic theory,” *Class. Quant. Grav.* **32**, 185007 (2015), [arXiv:1504.04861 \[gr-qc\]](#) .
- [71] D. Momeni, R. Myrzakulov, and E. Gdeikli, “Cosmological viable mimetic $f(R)$ and $f(R, T)$ theories via Noether symmetry,” *Int. J. Geom. Meth. Mod. Phys.* **12**, 1550101 (2015), [arXiv:1502.00977 \[gr-qc\]](#) .
- [72] Ratbay Myrzakulov, Lorenzo Sebastiani, Sunny Vagnozzi, and Sergio Zerbini, “Mimetic covariant renormalizable gravity,” *Fund. J.*

- Mod. Phys. **8**, 119–124 (2015), [arXiv:1505.03115 \[gr-qc\]](#) .
- [73] Guido Cognola, Ratbay Myrzakulov, Lorenzo Sebastiani, Sunny Vagnozzi, and Sergio Zerbini, “Covariant Horava-like and mimetic Horndeski gravity: cosmological solutions and perturbations,” *Class. Quant. Grav.* **33**, 225014 (2016), [arXiv:1601.00102 \[gr-qc\]](#) .
- [74] Zahra Haghani, Tiberiu Harko, Hamid Reza Sepangi, and Shahab Shahidi, “The scalar Einstein-aether theory,” (2014), [arXiv:1404.7689 \[gr-qc\]](#) .
- [75] Yevgeniya Rabochaya and Sergio Zerbini, “A note on a mimetic scalartensor cosmological model,” *Eur. Phys. J.* , **85** (2016), [arXiv:1509.03720 \[gr-qc\]](#) .
- [76] Frederico Arroja, Nicola Bartolo, Purnendu Karmakar, and Sabino Matarrese, “The two faces of mimetic Horndeski gravity: disformal transformations and Lagrange multiplier,” *JCAP* **1509**, 051 (2015), [arXiv:1506.08575 \[gr-qc\]](#) .
- [77] Mariam Bouhmadi-Lpez, Che-Yu Chen, and Pisin Chen, “Primordial Cosmology in Mimetic Born-Infeld Gravity,” *JCAP* **1711**, 053 (2017), [arXiv:1709.09192 \[gr-qc\]](#) .
- [78] Che-Yu Chen, Mariam Bouhmadi-Lpez, and Pisin Chen, “Black hole solutions in mimetic Born-Infeld gravity,” *Eur. Phys. J.* , **59** (2018), [arXiv:1710.10638 \[gr-qc\]](#) .
- [79] Ratbay Myrzakulov and Lorenzo Sebastiani, “Non-local $f(R)$ -mimetic gravity,” *Astrophys. Space Sci.* **361**, 188 (2016), [arXiv:1601.04994 \[gr-qc\]](#) .
- [80] Sabir Ramazanov, “Initial Conditions for Imperfect Dark Matter,” *JCAP* **1512**, 007 (2015), [arXiv:1507.00291 \[gr-qc\]](#) .
- [81] S. A. Paston, “Forms of action for perfect fluid in General Relativity and mimetic gravity,” *Phys. Rev.* , 084059 (2017), [arXiv:1708.03944 \[gr-qc\]](#) .
- [82] Yunlong Zheng, Liuyuan Shen, Yicen Mou, and Mingzhe Li, “On (in)stabilities of perturbations in mimetic models with higher derivatives,” *JCAP* **1708**, 040 (2017), [arXiv:1704.06834 \[gr-qc\]](#) .
- [83] Yong Cai and Yun-Song Piao, “Higher order derivative coupling to gravity and its cosmological implications,” *Phys. Rev.* , 124028 (2017), [arXiv:1707.01017 \[gr-qc\]](#) .
- [84] Kazufumi Takahashi and Tsutomu Kobayashi, “Extended mimetic gravity: Hamiltonian analysis and gradient instabilities,” *JCAP* **1711**, 038 (2017), [arXiv:1708.02951 \[gr-qc\]](#) .
- [85] Mohammad Ali Gorji, Seyed Ali Hosseini Mansoori, and Hassan Firouzjahi, “Higher Derivative Mimetic Gravity,” *JCAP* **1801**, 020 (2018), [arXiv:1709.09988 \[astro-ph.CO\]](#) .
- [86] Rampei Kimura, Atsushi Naruko, and Daisuke Yoshida, “Extended vector-tensor theories,” *JCAP* **1701**, 002 (2017), [arXiv:1608.07066 \[gr-qc\]](#) .
- [87] Emmanuel N. Saridakis and Minas Tsoukalas, “Bi-scalar modified gravity and cosmology with conformal invariance,” *JCAP* **1604**, 017 (2016), [arXiv:1602.06890 \[gr-qc\]](#) .
- [88] Naser Sadeghnezhad and Kourosh Nozari, “Braneworld Mimetic Cosmology,” *Phys. Lett.* , 134–140 (2017), [arXiv:1703.06269 \[gr-qc\]](#) .
- [89] Sunny Vagnozzi, “Recovering a MOND-like acceleration law in mimetic gravity,” *Class. Quant. Grav.* **34**, 185006 (2017), [arXiv:1708.00603 \[gr-qc\]](#) .
- [90] Liuyuan Shen, Yicen Mou, Yunlong Zheng, and Mingzhe Li, “Direct couplings of mimetic dark matter and their cosmological effects,” *Chin. Phys.* , 015101 (2018), [arXiv:1710.03945 \[gr-qc\]](#) .
- [91] P. C. W. Davies, “Thermodynamics of Black Holes,” *Proc. Roy. Soc. Lond.* , 499–521 (1977).
- [92] Robert M. Wald, “Black hole in a uniform magnetic field,” *Phys. Rev.* , 1680–1685 (1974).
- [93] A. De Rujula, S. L. Glashow, and U. Sarid, “CHARGED DARK MATTER,” *Nucl. Phys.* , 173–194 (1990).
- [94] Kris Sigurdson, Michael Doran, Andriy Kurylov, Robert R. Caldwell, and Marc Kamionkowski, “Dark-matter electric and magnetic dipole moments,” *Phys. Rev.* , 083501 (2004), [Erratum: *Phys. Rev.D*73,089903(2006)], [arXiv:astro-ph/0406355 \[astro-ph\]](#) .
- [95] Vitor Cardoso, Caio F. B. Macedo, Paolo Pani, and Valeria Ferrari, “Black holes and gravitational waves in models of minicharged dark matter,” *JCAP* **1605**, 054 (2016), [arXiv:1604.07845 \[hep-ph\]](#) .
- [96] Gamal G. L. Nashed, “Charged axially symmetric solution, energy and angular momentum in tetrad theory of gravitation,” *Int. J. Mod. Phys.* , 3181–3197 (2006), [arXiv:gr-qc/0501002 \[gr-qc\]](#) .
- [97] Gamal Gergess Lamee Nashed, “Charged axially symmetric solution and energy in teleparallel theory equivalent to general relativity,” *Eur. Phys. J.* , 851–857 (2007), [arXiv:0706.0260 \[gr-qc\]](#) .
- [98] Gamal Gergess Lamee Nashed, “Charged Dilaton, Energy, Momentum and Angular-Momentum in Teleparallel Theory Equivalent to General Relativity,” *Eur. Phys. J.* , 291–302 (2008), [arXiv:0804.3285 \[gr-qc\]](#) .
- [99] Gamal G. L. Nashed, “Brane World black holes in Teleparallel Theory Equivalent to General Relativity and their Killing vectors, Energy, Momentum and Angular-Momentum,” *Chin. Phys.* **B19**, 020401 (2010), [arXiv:0910.5124 \[gr-qc\]](#) .
- [100] Gamal G. L. Nashed, “Reissner Nordstrom solutions and Energy in teleparallel theory,” *Mod. Phys. Lett.* , 2241–2250 (2006), [arXiv:gr-qc/0401041 \[gr-qc\]](#) .
- [101] Bing Zhang, “Mergers of Charged Black Holes: Gravitational Wave Events, Short Gamma-Ray Bursts, and Fast Radio Bursts,” *Astrophys. J.* **827** (2016), 10.3847/2041-8205/827/2/L31, [arXiv:1602.04542 \[astro-ph.HE\]](#) .
- [102] Ruomin Zhu and Thomas Osburn, “Inspiral into a charged black hole,” *Phys. Rev.* , 104058 (2018), [arXiv:1802.00836 \[gr-qc\]](#) .
- [103] S. W. Hawking and Don N. Page, “Thermodynamics of Black Holes in anti-De Sitter Space,” *Commun. Math. Phys.* **87**, 577 (1983).
- [104] Andrew Chamblin, Roberto Emparan, Clifford V. Johnson, and Robert C. Myers, “Charged AdS black holes and catastrophic holography,” *Phys. Rev.* , 064018 (1999), [arXiv:hep-th/9902170 \[hep-th\]](#) .
- [105] Andrew Chamblin, Roberto Emparan, Clifford V. Johnson, and Robert C. Myers, “Holography, thermodynamics and fluctuations of charged AdS black holes,” *Phys. Rev.* , 104026 (1999), [arXiv:hep-th/9904197 \[hep-th\]](#) .
- [106] P. C. W. Davies, “Thermodynamic Phase Transitions of Kerr-Newman Black Holes in De Sitter Space,” *Class. Quant. Grav.* **6**, 1909 (1989).
- [107] A. M. Awad, S. Capozziello, and G. G. L. Nashed, “ D -dimensional charged Anti-de-Sitter black holes in $f(T)$ gravity,”

- JHEP **07**, 136 (2017), [arXiv:1706.01773 \[gr-qc\]](#) .
- [108] Salvatore Capozziello, P. A. Gonzalez, Emmanuel N. Saridakis, and Yerko Vasquez, “Exact charged black-hole solutions in D -dimensional $f(T)$ gravity: torsion vs curvature analysis,” JHEP **02**, 039 (2013), [arXiv:1210.1098 \[hep-th\]](#) .
- [109] J. P. S. Lemos, “Cylindrical black hole in general relativity,” Phys. Lett. , 46–51 (1995), [arXiv:gr-qc/9404041 \[gr-qc\]](#) .
- [110] Adel M. Awad, “Higher dimensional charged rotating solutions in (A)dS space-times,” Class. Quant. Grav. **20**, 2827–2834 (2003), [arXiv:hep-th/0209238 \[hep-th\]](#) .
- [111] John Stachel, “Globally stationary but locally static space-times: A gravitational analog of the Aharonov-Bohm effect,” Phys. Rev. , 1281–1290 (1982).
- [112] I. H. Salazar, A. Garcia, and J. Plebanski, “Duality Rotations and Type D Solutions to Einstein Equations With Nonlinear Electromagnetic Sources,” J. Math. Phys. **28**, 2171–2181 (1987).
- [113] Eloy Ayon-Beato and Alberto Garcia, “New regular black hole solution from nonlinear electrodynamics,” Phys. Lett. , 25 (1999), [arXiv:hep-th/9911174 \[hep-th\]](#) .
- [114] Waldemar Berej, Jerzy Matyjasek, Dariusz Tryniecki, and Mariusz Woronowicz, “Regular black holes in quadratic gravity,” Gen. Rel. Grav. **38**, 885–906 (2006), [arXiv:hep-th/0606185 \[hep-th\]](#) .
- [115] Yun Soo Myung, Yong-Wan Kim, and Young-Jai Park, “Thermodynamics of regular black hole,” Gen. Rel. Grav. **41**, 1051–1067 (2009), [arXiv:0708.3145 \[gr-qc\]](#) .
- [116] Dominic Brecher, Jianyang He, and Moshe Rozali, “On charged black holes in anti-de Sitter space,” JHEP **04**, 004 (2005), [arXiv:hep-th/0410214 \[hep-th\]](#) .
- [117] Frank J. Tipler, “Singularities in conformally flat spacetimes,” Physics Letters A **64**, 8 – 10 (1977).
- [118] C.J.S. Clarke and A. Królak, “Conditions for the occurrence of strong curvature singularities,” Journal of Geometry and Physics **2**, 127 – 143 (1985).
- [119] S. W. Hawking and G. E. R. Ellis, *The Large Scale Structure of Spacetime* (Cambridge University Press, Cambridge, 1973).
- [120] Gamal G. L. Nashed, “Isotropic stars in higher-order torsion scalar theories,” Adv. High Energy Phys. **2016**, 7020162 (2016), [arXiv:1602.07193 \[physics.gen-ph\]](#) .
- [121] C. J. Hunter, “The Action of instantons with nut charge,” Phys. Rev. , 024009 (1999), [arXiv:gr-qc/9807010 \[gr-qc\]](#) .
- [122] S. W. Hawking, C. J. Hunter, and Don N. Page, “Nut charge, anti-de Sitter space and entropy,” Phys. Rev. , 044033 (1999), [arXiv:hep-th/9809035 \[hep-th\]](#) .
- [123] J. D. Bekenstein, “Black holes and the second law,” Lett. Nuovo Cim. **4**, 737–740 (1972).
- [124] Jacob D. Bekenstein, “Black holes and entropy,” Phys. Rev. , 2333–2346 (1973).
- [125] G. W. Gibbons and S. W. Hawking, “Cosmological Event Horizons, Thermodynamics, and Particle Creation,” Phys. Rev. , 2738–2751 (1977).
- [126] Yun Soo Myung, Yong-Wan Kim, and Young-Jai Park, “Quantum Cooling Evaporation Process in Regular Black Holes,” Phys. Lett. , 221–225 (2007), [arXiv:gr-qc/0702145 \[GR-QC\]](#) .
- [127] Wontae Kim, Hyeonjoon Shin, and Myungseok Yoon, “Anomaly and Hawking radiation from regular black holes,” J. Korean Phys. Soc. **53**, 1791–1796 (2008), [arXiv:0803.3849 \[gr-qc\]](#) .
- [128] I. G. Dymnikova, “De Sitter-Schwarzschild Black Hole: its Particlelike Core and Thermodynamical Properties,” International Journal of Modern Physics D **5**, 529–540 (1996).
- [129] Irina Dymnikova, “Cosmological term as a source of mass,” Class. Quant. Grav. **19**, 725–740 (2002), [arXiv:gr-qc/0112052 \[gr-qc\]](#) .
- [130] Irina Dymnikova, “Generic Features of Thermodynamics of Horizons in Regular Spherical Space-Times of the Kerr-Schild Class,” Universe **4**, 63 (2018).
- [131] K. Ghaderi and B. Malakolkalami, “Thermodynamics of the Schwarzschild and the ReissnerNordström black holes with quintessence,” Nucl. Phys. , 10–18 (2016).
- [132] Sean A. Hayward, “Formation and evaporation of regular black holes,” Phys. Rev. Lett. **96**, 031103 (2006), [arXiv:gr-qc/0506126 \[gr-qc\]](#) .
- [133] Piero Nicolini, Anais Smailagic, and Euro Spallucci, “Noncommutative geometry inspired Schwarzschild black hole,” Phys. Lett. , 547–551 (2006), [arXiv:gr-qc/0510112 \[gr-qc\]](#) .
- [134] M. Sharif and Wajiha Javed, “Thermodynamics of a Bardeen black hole in noncommutative space,” Can. J. Phys. **89**, 1027–1033 (2011), [arXiv:1109.6627 \[gr-qc\]](#) .
- [135] K. A. Bronnikov, A. Dobosz, and I. G. Dymnikova, “Nonsingular vacuum cosmologies with a variable cosmological term,” Class. Quant. Grav. **20**, 3797–3814 (2003), [arXiv:gr-qc/0302029 \[gr-qc\]](#) .
- [136] Kirill Bronnikov, Irina Dymnikova, and Evgeny Galaktionov, “Multi-horizon spherically symmetric spacetimes with several scales of vacuum energy,” Class. Quant. Grav. **29**, 095025 (2012), [arXiv:1204.0534 \[gr-qc\]](#) .
- [137] Khireddine Nouicer, “Black holes thermodynamics to all order in the Planck length in extra dimensions,” Class. Quant. Grav. **24**, 5917–5934 (2007), [Erratum: Class. Quant. Grav. **24**, 6435 (2007)], [arXiv:0706.2749 \[gr-qc\]](#) .
- [138] Irina Dymnikova and Micha Korpuk, “Thermodynamics of regular cosmological black holes with the de sitter interior,” Entropy **13**, 1967–1991 (2011).
- [139] S. W. Hawking, “Particle Creation by Black Holes,” Euclidean quantum gravity, Commun. Math. Phys. **43**, 199–220 (1975), [167(1975)].
- [140] E. O. Babichev, V. I. Dokuchaev, and Yu N. Eroshenko, “Black holes in the presence of dark energy,” Phys. Usp. **56**, 1155–1175 (2013), [Usp. Fiz. Nauk **189**, no.12, 1257 (2013)], [arXiv:1406.0841 \[gr-qc\]](#) .
- [141] Meng-Sen Ma and Ren Zhao, “Stability of black holes based on horizon thermodynamics,” Phys. Lett. , 278–283 (2015), [arXiv:1511.03508 \[gr-qc\]](#) .
- [142] Irina Dymnikova and Michal Korpuk, “Regular black hole remnants in de Sitter space,” Phys. Lett. , 12–18 (2010).
- [143] Jingyun Man and Hongbo Cheng, “The calculation of the thermodynamic quantities of the Bardeen black hole,”

- [Gen. Rel. Grav. **46**, 1660 \(2014\), arXiv:1304.5686 \[hep-th\]](#) .
- [144] R. Tharanath, Jishnu Suresh, and V. C. Kuriakose, “Phase transitions and Geometrothermodynamics of Regular black holes,” [Gen. Rel. Grav. **47**, 46 \(2015\), arXiv:1406.3916 \[gr-qc\]](#) .
- [145] R. V. Maluf and Juliano C. S. Neves, “Thermodynamics of a class of regular black holes with a generalized uncertainty principle,” [Phys. Rev. , 104015 \(2018\), arXiv:1801.02661 \[gr-qc\]](#) .
- [146] Xavier Calmet and Iber Kuntz, “What is modified gravity and how to differentiate it from particle dark matter?” [Eur. Phys. J. , 132 \(2017\), arXiv:1702.03832 \[gr-qc\]](#) .