

A lattice formulation of $\mathcal{N} = 2$ supersymmetric SYK model

Mitsuhiro KATO^a, Makoto SAKAMOTO^b and Hiroto SO^c

*^aInstitute of Physics
University of Tokyo, Komaba
Meguro-ku, Tokyo 153-8902, Japan*

*^bDepartment of Physics
Kobe University
Nada-ku, Hyogo 657-8501, Japan*

*^cDepartment of Physics
Ehime University
Bunkyo-chou 2-5, Matsuyama 790-8577, Japan*

ABSTRACT

We construct $\mathcal{N} = 2$ supersymmetric SYK model on one-dimensional (euclidean time) lattice. One nilpotent supersymmetry is exactly realized on the lattice in use of the cyclic Leibniz rule (CLR).

Sachdev-Ye-Kitaev (SYK) model[1, 2] and its generalizations have been attracting much attentions in several contexts.¹ In these analysis, large N limit is fully utilized to obtain effective theory in the IR region. Finite N analysis, however, is also interesting to see, for example, some sort of stringy corrections for AdS/CFT-type correspondence.

For finite N analysis beyond perturbation, one approach is the exact diagonalization of the hamiltonian[5] based on the realization of fermion operators by gamma matrices. Another approach may be the lattice formulation on the euclidean time. The latter seems convenient for the calculation of multi-point correlation function of the operators with distinct times, especially for the comparison with results from the effective field theory of bi-local collective modes. Also, if higher-dimensional extension becomes possible, Monte Carlo simulation will be numerically less expensive than exact diagonalization of the hamiltonian. So we concentrate lattice formulation in the present paper. Among others we will focus on lattice formulation of the supersymmetric generalization[6] of SYK model. It is actually highly non-trivial, since realizing supersymmetry on lattice is a very difficult task[7].

The present authors have been studying a way for realizing nilpotent subalgebra of the supersymmetry on lattice in use of the cyclic Leibniz rule (CLR)[8, 9]. In the present letter, as an application of the CLR, we will construct $\mathcal{N} = 2$ supersymmetric SYK model[6] on lattice. As will be seen, one of the two supersymmetries is exactly realized on lattice thanks to the CLR.

Let us consider $\mathcal{N} = 2$ supersymmetric SYK model[6] whose hamiltonian H is given by the anti-commutator of two nilpotent supercharges Q and \bar{Q}

$$H = \{Q, \bar{Q}\}, \quad Q^2 = 0, \quad \bar{Q}^2 = 0, \quad (1)$$

where supercharges are defined by N complex fermions $\psi^i, \bar{\psi}^i$ ($i = 1, \dots, N$) and complex random couplings C_{ijk}, \bar{C}_{ijk} with totally anti-symmetric indices:

$$Q = \frac{i}{3!} C_{ijk} \psi^i \psi^j \psi^k, \quad \bar{Q} = \frac{i}{3!} \bar{C}_{ijk} \bar{\psi}^i \bar{\psi}^j \bar{\psi}^k. \quad (2)$$

These fermions satisfy the following anti-commutation relations

$$\{\psi^i, \psi^j\} = 0, \quad \{\bar{\psi}^i, \bar{\psi}^j\} = 0, \quad \{\psi^i, \bar{\psi}^j\} = \delta^{ij}. \quad (3)$$

The random couplings have non-zero second moment under quenched average

$$\overline{C_{ijk} \bar{C}_{ijk}} = \frac{2J}{N^2} \quad (4)$$

¹For recent review see, for example, [3, 4] and references therein.

with a characteristic constant J which controls the strength of the interaction.

The above supercharges are cubic in fermion, thereby hamiltonian has quartic interactions. This can be generalized by taking any odd number of fermions in supercharges. Hereafter we use generalized supercharges with q fermions. Corresponding action in euclidean time is defined by

$$S = \int dt \left\{ \bar{\psi}^i \partial_t \psi^i - \bar{b}^i b^i + \frac{i^{\frac{q-1}{2}}}{(q-1)!} C_{j_1 j_2 \dots j_q} \bar{b}^{j_1} \psi^{j_2} \dots \psi^{j_q} + \frac{i^{\frac{q-1}{2}}}{(q-1)!} \bar{C}_{j_1 j_2 \dots j_q} b^{j_1} \bar{\psi}^{j_2} \dots \bar{\psi}^{j_q} \right\}. \quad (5)$$

Here we have introduced complex auxiliary variables b^i, \bar{b}^i in order to realize supersymmetry linearly and make the action off-shell invariant. C and \bar{C} are q indices generalization of the random couplings. Supersymmetry transformation for each variable is

$$\begin{aligned} \delta_Q \psi^i &= 0, & \delta_{\bar{Q}} \psi^i &= \bar{b}^i, \\ \delta_Q b^i &= 0, & \delta_{\bar{Q}} b^i &= \partial_t \bar{\psi}^i, \\ \delta_Q \bar{\psi}^i &= b^i, & \delta_{\bar{Q}} \bar{\psi}^i &= 0, \\ \delta_Q \bar{b}^i &= \partial_t \psi^i, & \delta_{\bar{Q}} \bar{b}^i &= 0, \end{aligned} \quad (6)$$

where we omit transformation parameters, so that δ_Q and $\delta_{\bar{Q}}$ should be treated as Grassmann odd quantities.

Now let us make a lattice version of (5) and (6). First, we replace the variables $\psi^i(t), \bar{\psi}^i(t), b^i(t)$ and $\bar{b}^i(t)$ by the lattice variables $\psi_n^i, \bar{\psi}_n^i, b_n^i$ and \bar{b}_n^i where n stands for a lattice site. Then supersymmetry transformation should become

$$\begin{aligned} \delta_Q \psi_n^i &= 0, & \delta_{\bar{Q}} \psi_n^i &= \bar{b}_n^i, \\ \delta_Q b_n^i &= 0, & \delta_{\bar{Q}} b_n^i &= (\nabla^{(T)} \bar{\psi}^i)_n, \\ \delta_Q \bar{\psi}_n^i &= b_n^i, & \delta_{\bar{Q}} \bar{\psi}_n^i &= 0, \\ \delta_Q \bar{b}_n^i &= (\nabla^{(T)} \psi^i)_n, & \delta_{\bar{Q}} \bar{b}_n^i &= 0, \end{aligned} \quad (7)$$

where $\nabla^{(T)}$ is an appropriate difference operator. We use superscript (T) in order to distinguish it from the difference operator in the action for which we use $\nabla^{(A)}$.

Next, we construct a lattice action in the following form

$$S = S_{\text{kin}} + S_M + S_{\bar{M}}, \quad (8)$$

$$S_{\text{kin}} = \bar{\psi}_n^i \nabla_{nm}^{(A)} \psi_m^i - \bar{b}_n^i b_n^i, \quad (9)$$

$$S_M = \frac{i^{\frac{q-1}{2}}}{(q-1)!} C_{j_1 j_2 \dots j_q} M_{n_1 n_2 \dots n_q} \bar{b}_{n_1}^{j_1} \psi_{n_2}^{j_2} \dots \psi_{n_q}^{j_q}, \quad (10)$$

$$S_{\bar{M}} = \frac{i^{\frac{q-1}{2}}}{(q-1)!} \bar{C}_{j_1 j_2 \dots j_q} \bar{M}_{n_1 n_2 \dots n_q} b_{n_1}^{j_1} \bar{\psi}_{n_2}^{j_2} \dots \bar{\psi}_{n_q}^{j_q}, \quad (11)$$

where repeated lattice site indices are summed. $M_{n_1 n_2 \dots n_q}$ and $\bar{M}_{n_1 n_2 \dots n_q}$ are complex coefficients which define multiple product of variables. Note that the last $q - 1$ site indices of M and \bar{M} are totally symmetric. This action gives (5) in the naive continuum limit as far as M and \bar{M} go to 1 and ∇ goes to ∂_t . Requiring the invariance of the action under either transformation δ_Q or $\delta_{\bar{Q}}$ in (7), we obtain the conditions which should be satisfied by $\nabla^{(A)}$, $\nabla^{(T)}$, M and \bar{M} . For example if we require $\delta_{\bar{Q}}$ -invariance, then we have

$$\delta_{\bar{Q}} S_{\text{kin}} = -\bar{\psi}_n^i (\nabla_{nm}^{(A)} + \nabla_{mn}^{(T)}) \bar{\psi}_m^i = 0, \quad (12)$$

$$\delta_{\bar{Q}} S_M = \frac{i^{\frac{q-1}{2}}}{(q-1)!} C_{j_1 j_2 \dots j_q} M_{n_1 n_2 \dots n_q} \sum_{k=2}^q (-1)^{k-2} \bar{b}_{n_1}^{j_1} \psi_{n_2}^{j_2} \dots \psi_{n_{k-1}}^{j_{k-1}} \bar{b}_{n_k}^{j_k} \psi_{n_{k+1}}^{j_{k+1}} \dots \psi_{n_q}^{j_q} = 0, \quad (13)$$

$$\delta_{\bar{Q}} S_{\bar{M}} = \frac{i^{\frac{q-1}{2}}}{(q-1)!} \bar{C}_{j_1 j_2 \dots j_q} \bar{M}_{mn_2 \dots n_q} \nabla_{mn_1}^{(T)} \bar{\psi}_{n_1}^{j_1} \bar{\psi}_{n_2}^{j_2} \dots \bar{\psi}_{n_q}^{j_q} = 0. \quad (14)$$

The first condition (12) is satisfied if difference operator ∇ meets

$$\nabla_{nm}^{(A)} + \nabla_{mn}^{(T)} = 0. \quad (15)$$

The second condition (13) is satisfied if we make

$$M \text{ to be totally symmetric for all } q \text{ indices.} \quad (16)$$

And the third condition (14) is satisfied if \bar{M} meets

$$\sum_{\text{permutation of } \{n_1 \dots n_q\}} \nabla_{mn_1}^{(T)} \bar{M}_{mn_2 \dots n_q} = 0. \quad (17)$$

The summation in (17) can be reduced to the one in only cyclic permutation due to the totally symmetric nature of the last $q - 1$ indices. Thus this relation is nothing but the cyclic Leibniz rule (CLR)[8, 9].

It should be stressed that the CLR (17) is not an abstract relation but has many concrete solutions. For example, if we simply take $\nabla^{(A)} = \nabla^{(T)}$, then

$$M_{lmn} = \delta_{l,m} \delta_{l,n}, \quad (18)$$

$$\bar{M}_{lmn} = \frac{1}{6} (2\delta_{l,m-1} \delta_{l,n-1} + \delta_{l,m+1} \delta_{l,n-1} + \delta_{l,m-1} \delta_{l,n+1} + 2\delta_{l,m+1} \delta_{l,n+1}), \quad (19)$$

$$\nabla_{mn} = \frac{1}{2} (\delta_{m+1,n} - \delta_{m-1,n}) \quad (20)$$

is one of the ultralocal² solutions of (15), (16) and (17) for $q = 3$. Systematic construction of the solutions for the CLR with symmetric difference operator can be found in [10].

²Here ultralocal means the operator defined in a region with finite extent on a lattice. Local operators on the lattice include not only ultralocal ones but also ones with infinite extent which decay exponentially.

Although this might be the simplest solution, we would have species doubler with it. So we propose alternative solution

$$M_{lmn} = \delta_{l,m} \delta_{l,n}, \quad (21)$$

$$\begin{aligned} \bar{M}_{lmn} = & \frac{1}{24} \left[2(1+r)^2 \delta_{l+1,m} \delta_{l+1,n} + 2(1-r)^2 \delta_{l-1,m} \delta_{l-1,n} \right. \\ & + (1-r^2)(\delta_{l-1,m} \delta_{l+1,n} + \delta_{l+1,m} \delta_{l-1,n}) \\ & + (3-r)(1+r)(\delta_{l+1,m} \delta_{l,n} + \delta_{l,m} \delta_{l+1,n}) \\ & + (3+r)(1-r)(\delta_{l-1,m} \delta_{l,n} + \delta_{l,m} \delta_{l-1,n}) \\ & \left. + 2(3+r^2) \delta_{l,m} \delta_{l,n} \right], \end{aligned} \quad (22)$$

$$\nabla_{mn}^{(T)} = \frac{1}{2} [\delta_{m+1,n} - \delta_{m-1,n} + r(\delta_{m+1,n} + \delta_{m-1,n} - 2\delta_{m,n})], \quad (23)$$

$$\nabla_{mn}^{(A)} = \frac{1}{2} [\delta_{m+1,n} - \delta_{m-1,n} - r(\delta_{m+1,n} + \delta_{m-1,n} - 2\delta_{m,n})]. \quad (24)$$

Here r is a real parameter and the kinetic action with this $\nabla^{(A)}$ contains so-called Wilson term (r corresponds to the Wilson term coefficient) which lifts doublers up with cutoff-scale mass.

For $r = 1$ case, which corresponds to the forward difference operator, we can write down solutions with generic q :

$$\begin{aligned} M_{l n_1 \dots n_{q-1}} &= \frac{1}{q!} \sum_P \left(\delta_{l+1,n_1} \dots \delta_{l+1,n_{q-1}} + \delta_{l+1,n_1} \dots \delta_{l+1,n_{q-2}} \delta_{l,n_{q-1}} + \delta_{l+1,n_1} \dots \delta_{l+1,n_{q-3}} \delta_{l,n_{q-2}} \delta_{l,n_{q-1}} \right. \\ &\quad \left. + \dots + \delta_{l+1,n_1} \delta_{l,n_2} \dots \delta_{l,n_{q-1}} + \delta_{l,n_1} \dots \delta_{l,n_{q-1}} \right) \end{aligned} \quad (25)$$

$$= \frac{1}{q!} \sum_P \sum_{k=0}^{q-1} \left[\prod_{a=1}^k \delta_{l+1,n_a} \right] \left[\prod_{b=k+1}^{q-1} \delta_{l,n_b} \right] \quad (26)$$

where P stands for the summation over all permutations of $(n_1, n_2, \dots, n_{q-1})$, and $\prod_{a=1}^0 \delta_{l+1,n_a} =$

$1 = \prod_{b=q}^{q-1} \delta_{l,n_b}$ is understood.

Thus we have a concrete way how to construct $\delta_{\bar{Q}}$ -invariant lattice action for $\mathcal{N} = 2$ supersymmetric SYK model. If you need δ_Q -invariant action, just exchange the roles of M and \bar{M} .

A few remarks are in order:

We cannot require both δ_Q and $\delta_{\bar{Q}}$ invariances. Because these impose simultaneously the CLR relation and totally symmetric indices to M and \bar{M} , but there is no local solution of

CLR with symmetric M or \bar{M} [8].

For $\delta_{\bar{Q}}$ -invariance M is totally symmetric, but \bar{M} is not because of CLR and locality. Therefore \bar{M} is not complex conjugate of M , so that the resulting action is not hermitian. This “could-be sign problem” is of order $O(a)$ where a is lattice constant, and disappears at least in the naive continuum limit.

It seems the CLR approach is a unique way to realize supersymmetry for the models of this type. The other approaches, like a method through the Nicolai map or a method of finding a nilpotent transformation without difference operator[11, 12, 13], may not work for the model. In particular, the last term of the action (5) is $\delta_{\bar{Q}}$ -invariant but not $\delta_{\bar{Q}}$ -exact, therefor topological field theoretic approach cannot be applied. This supports the uniqueness of the CLR approach.

Acknowledgements

This work is supported in part by the Grant-in-Aid for Scientific Research No.25287049 (M.K.), No.15K05055 (M.S.) and No.25400260 (H.S.) by the Japanese Ministry of Education, Science, Sports and Culture.

References

- [1] S. Sachdev and J. Ye, “Gapless spin fluid ground state in a random, quantum Heisenberg magnet,” Phys. Rev. Lett. **70** (1993) 3339 doi:10.1103/PhysRevLett.70.3339 [cond-mat/9212030].
- [2] A. Kitaev, “ A simple model of quantum holography,” KITP strings seminar and Entanglement 2015 program (Feb. 12, April 7, and May 27, 2015). <http://online.kitp.ucsb.edu/online/entangled15/>.
- [3] V. Rosenhaus, “An introduction to the SYK model,” arXiv:1807.03334 [hep-th].
- [4] G. Sárosi, “AdS₂ holography and the SYK model,” PoS Modave **2017** (2018) 001 doi:10.22323/1.323.0001 [arXiv:1711.08482 [hep-th]].
- [5] T. Li, J. Liu, Y. Xin and Y. Zhou, JHEP **1706** (2017) 111 doi:10.1007/JHEP06(2017)111 [arXiv:1702.01738 [hep-th]].

- [6] W. Fu, D. Gaiotto, J. Maldacena and S. Sachdev, “Supersymmetric Sachdev-Ye-Kitaev models,” *Phys. Rev. D* **95** (2017) no.2, 026009 Addendum: [*Phys. Rev. D* **95** (2017) no.6, 069904] doi:10.1103/PhysRevD.95.069904, 10.1103/PhysRevD.95.026009 [arXiv:1610.08917 [hep-th]].
- [7] M. Kato, M. Sakamoto and H. So, “Taming the Leibniz Rule on the Lattice,” *JHEP* **0805** (2008) 057 doi:10.1088/1126-6708/2008/05/057 [arXiv:0803.3121 [hep-lat]].
- [8] M. Kato, M. Sakamoto and H. So, “A criterion for lattice supersymmetry: cyclic Leibniz rule,” *JHEP* **1305** (2013) 089 doi:10.1007/JHEP05(2013)089 [arXiv:1303.4472 [hep-lat]].
- [9] M. Kato, M. Sakamoto and H. So, “Non-renormalization theorem in a lattice supersymmetric theory and the cyclic Leibniz rule,” *PTEP* **2017** (2017) no.4, 043B09 doi:10.1093/ptep/ptx045 [arXiv:1609.08793 [hep-lat]].
- [10] D. Kadoh and N. Ukita, “General solution of the cyclic Leibniz rule,” *PTEP* **2015** (2015) no.10, 103B04 doi:10.1093/ptep/ptv140 [arXiv:1503.06922 [hep-lat]].
- [11] S. Catterall and S. Karamov, “Exact lattice supersymmetry: The Two-dimensional N=2 Wess-Zumino model,” *Phys. Rev. D* **65** (2002) 094501 doi:10.1103/PhysRevD.65.094501 [hep-lat/0108024].
- [12] S. Catterall, “Lattice supersymmetry and topological field theory,” *JHEP* **0305** (2003) 038 doi:10.1088/1126-6708/2003/05/038 [hep-lat/0301028].
- [13] J. Giedt and E. Poppitz, “Lattice supersymmetry, superfields and renormalization,” *JHEP* **0409** (2004) 029 doi:10.1088/1126-6708/2004/09/029 [hep-th/0407135].