

# Universal Aspects of $U(1)$ Gauge Field Localization on Branes in $D$ -dimensions

L. F. F. Freitas,<sup>a</sup> G. Alencar<sup>a</sup> and R. R. Landim<sup>a</sup>

<sup>a</sup>Universidade Federal do Ceará - UFC,

Departamento de Física, Universidade Federal do Ceará, 60451-970 Fortaleza, Ceará, Brazil

E-mail: [luizfreitas@fisica.ufc.br](mailto:luizfreitas@fisica.ufc.br), [geovamaciel@gmail.com](mailto:geovamaciel@gmail.com)

**ABSTRACT:** In this work, we study the general properties of the  $D$ -vector field localization on  $(D - d - 1)$ -brane with co-dimension  $d$  in asymptotically AdS spacetime. We consider a conformally flat metric with the warp factor depending only on the transverse extra dimensions. It is known that the free gauge field is not confined in co-dimension one braneworlds. In addition, in most of the higher co-dimension models a general mechanism to solve this is lacking. For co-dimension one models, this problem was solved by using non-minimal couplings with gravity. Here we employ this mechanism stressing the universal aspects that lead to localization of the  $D$ -vector field. With this, an analytical solution for  $U(1)$  gauge field (zero-mode of transverse  $(D - d)$ -vector field) on the brane can always be obtained, and a localization condition can be found by setting up the bulk geometry as asymptotically AdS. Beyond that, the existence of this massless mode has as a consequence to exclude any possible tachyonic modes of the theory. Therefore, our solution has an universal validity for any warp factor and is independent of the particular model considered. We also obtain some conditions in which the confinement of gravity ensures the localization of the gauge field. Finally, it is studied the scalar components of  $D$ -vector field. A drawback in co-dimension one models is that the scalar components of  $D$ -vector field are not localized simultaneously with the gauge field. As a general result, we show that if we consider the coupling with the tensor and the Ricci scalar in a higher co-dimension, there is an indication that both sectors will be localized. As a concrete example, the above techniques are applied for the intersecting brane model. For this model, it is obtained that the branes introduce boundary conditions that fix all parameters of the model in such a way that both sectors, gauge and scalar fields, are confined.

**KEYWORDS:** Fields Localization, Geometrical Coupling,  $U(1)$  Gauge Field

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Review About Abelian Vector Field Localization</b>	<b>3</b>
<b>3</b>	<b>General Aspects of Vector Field Localization</b>	<b>5</b>
3.1	Localization of Transversal Sector of $U(1)$ Vector Field – $\hat{\mathcal{A}}^\mu$	7
3.1.1	Localization of Zero-Mode – $\hat{\mathcal{A}}_0^\mu$	8
3.2	Localization of the Scalar Components – $\mathcal{B}^j$	11
3.2.1	Localization of Zero-Mode – $\mathcal{B}_0^j$	14
<b>4</b>	<b>Application: Intersecting Brane Model</b>	<b>16</b>
<b>5</b>	<b>Final Remarks</b>	<b>21</b>
<b>A</b>	<b>Appendix</b>	<b>24</b>
A.1	Splitting of the Action (3.1) in the Sectors $\mathcal{S}_\perp[\hat{\mathcal{A}}_\mu]$ and $\mathcal{S}[\phi, B_k]$	24

---

## 1 Introduction

The formulation of theories in spacetimes with more than 4-dimensions as a tool to solve problems in physics is not new [1, 2]. However, only after the development of string theory and the compactification mechanisms of extra dimensions, at the end of the last century, this tool began to be regarded a possible real description of nature [3]. A feature of these higher dimensional theories was the need of the extra dimensions to be compactified into a very small spatial volume inaccessible in the available energy range. This because the Newton’s gravitational law depends explicitly on the number of spatial dimensions, and it indicates the presence of only three large spatial dimensions. The first to speculate about the possibility of these extra dimensions being non-compact were Rubakov and Shaposhnikov [4]. The authors showed that a large extra dimension is allowed as long as the fields of the Standard Model (SM), as well as the gravity, are confined to a 4-dimensional hypersurface. In such a way that our energy scale (TeV) does not allow us to access them.

In this direction, L. Randall and R. Sundrum (RS) proposed two models with warped geometry in an  $AdS_5$  spacetime with delta-like **3**-branes [5, 6]. The RS-I model, proposed to solve the Higgs hierarchy problem, considers a **5**-dimensional universe  $(x^\mu, \phi)$  with the spatial dimension  $\phi$  compactified under a circle with an orbifold symmetry  $S_1/Z_2$ . At the fixed points  $(\phi = 0, \pi)$  are placed two delta-like **3**-branes, and the **3**-brane at  $\phi = \pi$  would correspond to our universe with all SM fields confined. The RS-II model considers only one

delta-like **3**-brane with a non-compact and infinitely large extra dimension  $(x^\mu, y)$ , and it was proposed as an alternative to the compactification. On both models, the gravity and the scalar fields are localized on the **3**-brane. Despite this, the other fields of SM are not confined as expected by RS [7–9]. Soon after the success of RS models, other proposals of braneworld with localized gravity were presented in five dimensions (**5D**): smooth versions of RS-II (thick branes) [10, 11]; thick branes with inner structure [12, 13]; cosmological models, where the **3**-brane has a *Robertson-Walker* metric [14, 15]. Also, other solutions in spacetimes with more than **5D** as **3**-brane generated by string-like or vortex topological defects in **6D** [16, 17]; braneworld models generated by the intersection of delta-like branes [18, 19], and other proposals [20–24, 58].

In all the above models, the issue of the SM fields localization is always an important point to be verified [25–30], mainly the confinement of  $U(1)$  gauge field. It is a well known fact that the free abelian gauge field is not confined in **5D** braneworld models [9, 32, 33]. In **6D** or higher dimensional models, the confinement seems to be possible since there is only one infinitely large extra dimension [8, 34, 35], however, a more detailed analysis, by exploring the Hodge duality symmetry [58] for example, shows the opposite. Some attempts to solve this problem were performed. In most cases, introducing new degrees of freedom such as interaction terms with fermionic or scalar (*dilatonic*) fields [11, 36, 38–43]. Although these mechanisms allow to confine the  $U(1)$  gauge field, other questions arise: what is the meaning of these new fields for the theory? or, do such mechanisms work for other braneworld models?

Recently, Ghoroku *et al* developed a localization mechanism in RS-II model without the need of introducing new degrees of freedom [44]. They introduced a mass term and a non-covariant interaction between the vector field and the **3**-brane. This mechanism works, however, there is no solid motivation for the introduction of a coupling with the **3**-brane. Furthermore, they introduce a free parameter in the theory. Based on this method, a purely geometric localization mechanism was proposed in Refs. [18, 45, 46], where an interaction term of the vector field with the Ricci scalar is added. This *geometric coupling* allows us to localize the massless mode of abelian vector field and has the advantage that it is covariant and does not introduce new degrees of freedom neither free parameters in the theory. Beyond that, the interaction with the **3**-brane arises as a consequence of the coupling of the vector field with the gravity. Afterward, this mechanism was applied to the massive modes which the resonant modes of vector and  $p$ -form fields can be studied [47–49]; also, by looking for phenomenological consequences, as a residual non-zero mass for the photon due to the existence of extra dimensions [50]; beyond this, the application of this non-minimal coupling with gravity to analyze the localization of other fields [51, 52]. All this points were developed in warped models with only *one* infinitely large extra dimension.

Despite the above results, a generalization of the geometrical coupling mechanism to more than one transverse extra dimensions is lacking. As presented above many other scenarios of braneworld with more extra dimensions were proposed allowing a more rich gravitational configuration. Furthermore, the vector field will have more scalar components which can play an important role on the brane. In view of all this, in this manuscript we study the vector field localization in a generic spacetime with an arbitrary number of large

extra dimensions. First, we look for the universal aspects of localization for the two sectors of the  $\mathbf{D}$ -vector field, the gauge  $(\mathbf{D} - \mathbf{d})$ -vector field and the scalar components on the brane. By universal it means aspects that do not depend on any specific braneworld model, but only on the fact that they are asymptotically AdS. Also, due to the localization of such fields be valid for a wide variety of braneworld models. In this way, we look for the possibility of both components can be simultaneously localized for some range of the parameters of the model. We want to make it clear that the localization of the scalar components does not necessarily imply corrections to the Coulomb law, instead such components could be interpreted as Higgs fields, for example. As a concrete application of our results, we consider the intersecting brane model cited above.

This work is organized as follows: In section (2), a review of the main results about abelian vector field localization through the geometric coupling is made. In section (3), the general aspects of the confinement of the  $\mathbf{U}(1)$  gauge field on a generic braneworld scenario are discussed. It is also done a study about the localization of scalar components. In section (4), we carry out an application of our general results for the specific case of intersecting branes. The conclusions are left for section (5).

## 2 Review About Abelian Vector Field Localization

In a general way, when we talk about localization of fields in RS-like braneworlds, it means that we want to factor out the action

$$S = \int d^4x d^{D-4}z \sqrt{-g^{(D)}} \mathcal{L}_{(\text{matter})}^{(D)}, \quad (2.1)$$

into a sector containing an effective action on the  $\mathbf{3}$ -brane and an integral in the coordinates of extra dimensions, *i.e.*,

$$S = \int d^{D-4}z f(z) \int d^4x \sqrt{-g^{(4)}} \mathcal{L}_{(\text{matter})}^{(4)} = K \int d^4x \sqrt{-g^{(4)}} \mathcal{L}_{(\text{matter})}^{(4)}. \quad (2.2)$$

Thus, we say that the theory is well-defined, *i.e.*, the field is localized on the brane, when the integral  $K$  is finite. As mentioned above, we will restrict ourselves to the localization of the vector field in a generic  $\mathbf{D}$ -dimensional RS-like braneworld scenario. In doing so, the gravitational field will only be used as a background and we are not interested in its dynamics. Before considering the  $\mathbf{D}$ -dimensional case, it is instructive to briefly review some results found in the literature on this subject.

Let us start with the free gauge field in  $\mathbf{5D}$  RS-II model, where the action can be written as [9]

$$S = -\frac{1}{4} \int d^4x dy \sqrt{-g} g^{MN} g^{PQ} \mathcal{F}_{MP} \mathcal{F}_{NQ}. \quad (2.3)$$

In the above equation,  $\mathcal{F}_{MP} = \partial_M \mathcal{A}_P - \partial_P \mathcal{A}_M$  is the field strength,  $\mathcal{A}_M = (\mathcal{A}_\mu, \mathcal{A}_5)$  is the five dimensional gauge field and  $g_{MN}$  is the background metric of RS-II model given by

$$ds^2 = g_{MN} dX^M dX^N = \frac{1}{(1 + k|y|)^2} (\eta_{\mu\nu} dx^\mu dx^\nu + dy^2). \quad (2.4)$$

The gauge-invariance of action (2.3) can be used to choose  $\partial_\mu \mathcal{A}^\mu = 0$ , and also  $\mathcal{A}_5 = 0$ . With these assumptions, the variation of (2.3), followed by the standard separation of variables  $\mathcal{A}_\mu = A_\mu(x)\chi(y)$ , gives the equations

$$\partial_\mu F^{\mu\nu}(x) = m^2 A^\nu(x) \quad (2.5)$$

and

$$-e^{-\sigma(y)} \partial_5 \left( e^{\sigma(y)} \partial^5 \chi(y) \right) = m^2 \chi(y). \quad (2.6)$$

In Eq. (2.5), the contraction of the greek indexes  $(\mu, \nu \dots)$  is done with the Minkowski metric. Besides all this, the action (2.3) can be factored as

$$S = - \int dy e^{\sigma(y)} \chi^2 \int d^4x \left[ \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} m^2 A^\mu A_\mu \right]. \quad (2.7)$$

Now we turn to the analysis of the zero mode ( $m^2 = 0$ ) of Eqs. (2.5) and (2.6). In this case the equation (2.6) gives a constant solution  $\chi_0 = c_0$ . With this, it is easy to check that, from Eq. (2.7), the integral of localization for the gauge field  $A_{(0)}^\mu$  (massless mode of vector field  $A^\mu$ ) is not finite, viz,

$$\int dy e^{\sigma(y)} \chi_0^2 = c_0^2 \int dy e^{\sigma(y)} = c_0^2 \int_{-\infty}^{\infty} \frac{dy}{1 + k|y|} \rightarrow \infty, \quad (2.8)$$

and therefore the gauge field is not confined for this **5D** RS-II model.

Another possibility is to consider the **6D** string-like braneworld model with background metric given by [8, 37]

$$ds^2 = e^{2\sigma(r)} (\eta_{\mu\nu} dx^\mu dx^\nu + R_0^2 d\theta^2 + dr^2), \quad (2.9)$$

which has one compactified and one large extra dimension. It is possible to write an expression analogous to Eq. (2.7) for the zero-mode of gauge field as

$$S = - \int_0^{2\pi} R_0 d\theta \int_0^\infty dr e^{2\sigma(r)} \chi_0^2(r) \int d^4x \frac{1}{4} F_{\mu\nu} F^{\mu\nu}. \quad (2.10)$$

In order to obtain the above action, we used  $A_r = A_\theta = 0$  and the gauge condition  $\partial_\mu A^\mu = 0$ . Similarly to the case in **5** dimensions, we can get an equation which drives the massive modes given by [8, 37]

$$-e^{-2\sigma(r)} \partial_r \left( e^{2\sigma(r)} \partial_r \chi(r) \right) = m^2 \chi(r). \quad (2.11)$$

Again, there is a constant solution for the zero-mode, but now the integral in the extra dimensions is finite. However it has been pointed by Duff *et al* that the constant solution is not consistent with the Hodge Duality symmetry present in the model [58]. By considering this symmetry these authors show that the only consistent solution cannot be localized for any co-dimension one braneworld model [58]. Therefore, it is necessary to obtain a mechanism to confine this field consistently. Initially, interaction terms with other matter

fields were introduced. However, a mechanism that does not introduce new degrees of freedom was lacking. Only recently, a powerful mechanism, by using non-minimal couplings with gravity, was proposed to solve that problem.

The origin of the non-minimal coupling mechanism can be traced back to Ref. [44], where K. Ghoroku and A. Nakamura developed a localization mechanism for the vector field without introducing new degrees of freedom. The authors proposed an action in **5**-dimensions given by

$$S = - \int d^4x dz \sqrt{-g} \left[ \frac{1}{4} g^{MN} g^{PQ} \mathcal{F}_{MP} \mathcal{F}_{NQ} + \frac{1}{2} (M^2 + \tau \delta(z)) g^{MN} \mathcal{A}_M \mathcal{A}_N \right] \quad (2.12)$$

where  $g_{MN}$  is given in Eq. (2.4),  $M^2$  is the mass of vector field and  $\tau$  is a coupling constant between the vector field and the **3**-brane. Despite the fact that the above action is not gauge invariant, the authors show that a gauge field with all desired properties is trapped over the **3**-dimensional brane. For this, they proposed  $\mathcal{A}_N = (\hat{A}_\mu^T + \mathcal{A}_\mu^L, \mathcal{A}_5)$ , with  $\partial_\mu \hat{A}_T^\mu = 0$ , and showed that it is possible to confine the massless mode of the transverse field  $\hat{A}_T^\mu$  when  $M > 0$  and  $\tau = -2k \left( \sqrt{1 + M^2/k^2} - 1 \right)$ . However, the authors do not give a solid motivation to add an interaction term with the **3**-brane.

Based on this mechanism, Refs. [18, 45, 46] recently proposed a localization mechanism by addition of an interaction term between the vector field and the Ricci scalar  $R$ . They start of an action given by

$$S = - \int d^4x dz \sqrt{-g} \left[ \frac{1}{4} g^{MN} g^{PQ} \mathcal{F}_{MP} \mathcal{F}_{NQ} + \frac{\gamma_1}{2} R g^{MN} \mathcal{A}_M \mathcal{A}_N \right], \quad (2.13)$$

where  $\gamma_1$  is a parameter fixed by the boundary conditions and  $R$  is given by  $R = 16k\delta(z) - 20k^2$ . This model, called *geometric coupling*, has no additional degrees of freedom or free parameters and it provides an ‘origin’ for the interaction terms proposed in (2.12). As in the Ghoroku model, here the action is not gauge invariant. Nevertheless, the authors also find a gauge field confined on the brane. For this, again it is used  $\mathcal{A}_N = (\hat{A}_\mu + \partial_\mu \Phi, \mathcal{A}_5)$ , where  $\partial_\mu \hat{A}^\mu = 0$  defines the transverse gauge field  $\hat{A}^\mu$ . The localization of the massless mode of  $\hat{A}^\mu$  occurs for  $\gamma_1 = -1/16$ , a valid result not only for RS-II model, but also for its smooth versions. It makes this mechanism extremely interesting and powerful. Although the localization of gauge field is the important point in this study, the scalar components of vector field can play an important role on the **3**-brane. In Ref. [45], the authors also studied the scalar component  $\mathcal{A}_5$  and they showed that it can be performed, but for a different value of  $\gamma_1$ . Therefore, this kind of coupling only with the Ricci scalar does not allow the confinement of both sectors simultaneously in this co-dimension 1 model.

### 3 General Aspects of Vector Field Localization

In this section we will generalize the results of the previous section by considering non-minimal couplings of vector field with the scalar and the Ricci tensor in a  $D$ -dimensional

spacetime. To do this, let us start from the action

$$S_2 = - \int d^{(D-d)} x d^d y \sqrt{-g} \left[ \frac{1}{4} \mathcal{F}_{MN} \mathcal{F}^{MN} + \frac{\lambda_1}{2} R \mathcal{A}_M \mathcal{A}^M + \frac{\lambda_2}{2} R_{MN} \mathcal{A}^M \mathcal{A}^N \right]. \quad (3.1)$$

In the above equation  $\mathcal{A}_M(x, y)$  is the vector field in  $D$ -dimensions,  $\mathcal{F}_{MP} = \partial_M \mathcal{A}_P - \partial_P \mathcal{A}_M$  is the field strength tensor and  $R$  and  $R_{NM}$  are the scalar and Ricci tensor respectively. Since we want to study general aspects that do not depend on any specific braneworld model we will consider a generic background metric given by

$$ds^2 = g_{MN} dX^M dX^N = e^{2\sigma(y)} \left( \eta_{\mu\nu} dx^\mu dx^\nu + \eta_{jk} dy^j dy^k \right), \quad (3.2)$$

where  $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1, \dots)$ ,  $\eta_{jk} = \delta_{jk}$  (Kronecker Delta) and the warp factor  $\sigma(y)$  depends only on the transverse extra dimensions  $y^j$ . Throughout the manuscript, capital indexes  $M, N$  assume value on all  $D$  dimensions; greek indexes are related to brane coordinates and run from  $\mu, \nu = (1, 2, \dots, D-d)$  and latin indexes are related to the  $d$  extra dimensions and run from  $j, k = (D-d+1, D-d+2, \dots, D)$ .

At this point is important to note that the action (3.1) is invariant by general coordinate transformation. Thus, when we perform a Lorentz transformation at the brane,

$$A^{M'} = L^{M'}_M A^M \quad \rightarrow \quad \begin{pmatrix} A^{\mu'} \\ B^{j'} \end{pmatrix} = \begin{pmatrix} \Lambda^{\mu'}_\mu & 0 \\ 0 & \delta^{j'}_j \end{pmatrix} \begin{pmatrix} A^\mu \\ B^j \end{pmatrix}, \quad (3.3)$$

where  $\Lambda^{\mu'}_\mu$  is an usual Lorentz transformation in Minkowski spacetime. It makes clear that the components  $A^\mu$  will be a Lorentz vector at the brane. Also, the components  $B^j$  will be Lorentz scalars at the brane. Therefore, it is convenient to split the analysis of localization for this two fields (sectors).

Another important point is that we do not treat the backreaction of vector field, which is considered as a small perturbation. Despite the fact that in the above action gravity is non-minimally coupled to the vector field, this will not change the gravity dynamics and it can be considered just as a background. In this way, the vector field localization can be treated separately. As in the models of section (2), the above action is also not gauge invariant. As we have seen, this is not a problem since an effective gauge field can still be obtained on the brane. Here, we will show the same: despite the fact that in  $D$  dimensions the action is not gauge invariant, an effective action on the brane can be obtained and it contains a gauge field with all desired properties: massless, gauge and Lorentz invariant.

Finally, in this and the next sections we will need of explicit expressions for the scalar and the Ricci tensor obtained from the metric (3.2). The Ricci tensor is given by

$$\begin{aligned} R_{MN} = & - \left[ (D-2) \delta_M^k \delta_N^j + \eta_{MN} \eta^{kj} \right] \partial_k \partial_j \sigma(y) \\ & + (D-2) \left[ \delta_M^k \delta_N^j - \eta_{MN} \eta^{kj} \right] \partial_k \sigma(y) \partial_j \sigma(y); \end{aligned} \quad (3.4)$$

and the Ricci scalar by

$$R = - (D-1) e^{-2\sigma(y)} \eta^{jk} [2 \partial_k \partial_j \sigma(y) + (D-2) \partial_k \sigma(y) \partial_j \sigma(y)]. \quad (3.5)$$

In the above equations and from now on we will use the definition  $\partial_k \equiv \frac{\partial}{\partial y^k}$ .

### 3.1 Localization of Transversal Sector of $U(1)$ Vector Field – $\hat{\mathcal{A}}^\mu$

Due to the above discussion, and without loss of generality, it is convenient to split the  $D$ -dimensional vector field as  $\mathcal{A}_N = (\hat{\mathcal{A}}_\mu + \partial_\mu \phi, \mathcal{B}_k)$ , where  $\hat{\mathcal{A}}_\mu$  is the transverse sector of the vector field on the brane defined such that  $\eta^{\mu\nu} \partial_\mu \hat{\mathcal{A}}_\nu = 0$ . Note that the splitting of  $\mathcal{A}_N$  field does not specify any gauge for the vector field, in fact the action (3.1) is not gauge-invariant, therefore such symmetry can not be used to exclude degrees of freedom.

With the above definitions the action (3.1) can be split in two independent parts (see appendix A)

$$S = S_\perp[\hat{\mathcal{A}}_\mu] + S[\phi, \mathcal{B}_k], \quad (3.6)$$

where  $S_\perp[\hat{\mathcal{A}}_\mu]$  contains only the transverse sector

$$S_\perp = - \int d^{(D-d)} x d^d y \sqrt{-g} \left\{ \frac{1}{4} g^{\mu\nu} g^{\rho\lambda} \hat{\mathcal{F}}_{\mu\rho} \hat{\mathcal{F}}_{\nu\lambda} + \frac{1}{2} g^{\mu\nu} g^{jk} \partial_j \hat{\mathcal{A}}_\mu \partial_k \hat{\mathcal{A}}_\nu \right. \\ \left. + \frac{\lambda_1}{2} R g^{\mu\nu} \hat{\mathcal{A}}_\mu \hat{\mathcal{A}}_\nu + \frac{\lambda_2}{2} g^{\mu\nu} g^{\rho\lambda} R_{\mu\rho} \hat{\mathcal{A}}_\nu \hat{\mathcal{A}}_\lambda \right\}, \quad (3.7)$$

and another part contains only the longitudinal and scalar components of the vector field,  $\partial_\mu \phi$  and  $\mathcal{B}_k$  respectively. The explicit form of  $S[\phi, \mathcal{B}_k]$  was not written because it is not necessary for the discussion of this section. In next section, we will study the localization of  $\mathcal{B}_k$  components directly from (3.1). From the action (3.7) and using the metric (3.2), we can obtain the following equation of motion for the transverse sector  $\hat{\mathcal{A}}_\mu$ ,

$$-e^{-(D-4)\sigma} \partial_k \left( e^{(D-4)\sigma} \partial^k \hat{\mathcal{A}}^\lambda \right) + \lambda_1 R e^{2\sigma} \hat{\mathcal{A}}^\lambda + \lambda_2 R^\lambda{}_\mu \hat{\mathcal{A}}^\mu = \partial_\nu \hat{\mathcal{F}}^{\nu\lambda}, \quad (3.8)$$

where the Minkowski metric was used to lower/raise indexes.

In order to solve the equation (3.8) we first need of explicit form of  $R_{\mu\rho}$ . From Eq. (3.4) we can get that it is diagonal and given by

$$R_{\mu\rho} = - \left[ \partial_k \partial^k \sigma(y) + (D-2) \partial_k \sigma(y) \partial^k \sigma(y) \right] \eta_{\mu\rho} \equiv -h(y) \eta_{\mu\rho}. \quad (3.9)$$

With this and by performing the standard decomposition  $\hat{\mathcal{A}}_\nu = A_\mu(x) \chi(y) e^{-\frac{(D-4)}{2}\sigma(y)}$ , we find that Eq. (3.8) reduces to the following equations. One for the transverse vector field  $A_\mu(x)$  given by

$$\partial_\nu F^{\nu\mu}(x) = m^2 A^\mu(x) \quad (3.10)$$

and another which drives the mass modes of the theory, given by

$$-\partial_k \partial^k \chi + \left[ \frac{(D-4)^2}{4} \partial_k \sigma(y) \partial^k \sigma(y) + \frac{(D-4)}{2} \partial_k \partial^k \sigma(y) \right. \\ \left. + \lambda_1 R e^{2\sigma} - \lambda_2 h(y) \right] \chi = m^2 \chi. \quad (3.11)$$

With this, the action (3.7) can be written as

$$S_\perp = - \int d^d y \chi^2 \int d^{D-d} x \left\{ \frac{1}{4} F_{\mu\rho} F^{\mu\rho} + \frac{1}{2} m^2 A_\nu A^\nu \right\} \quad (3.12)$$

and a consistent effective theory localized over the brane requires that the integral

$$K = \int d^d y \chi^2 \quad (3.13)$$

be finite. The expression (3.11) is a Schrödinger-like equation with potential given by

$$U(y^j) = c_1 \partial_k \partial^k \sigma(y) + c_2 \partial_k \sigma(y) \partial^k \sigma(y), \quad (3.14)$$

where we have used the relations (3.5) and (3.9) to obtain the above result. Equation (3.11) is very convenient because it allows us to obtain qualitative informations about the system just by the analysis of potential  $U$ . As known from quantum mechanics, the asymptotic behavior of  $U$  determines whether the states  $\chi(y)$  can be normalized or not. If  $\lim_{|y| \rightarrow \infty} U(y) = \infty$ , then we have a confining system and all states can be normalized; if otherwise  $\lim_{|y| \rightarrow \infty} U(y) = -\infty$ , then the states cannot be normalized; and finally, when  $\lim_{|y| \rightarrow \infty} U(y) = u_0$  (constant) we have two possibilities: a) if  $m^2 < u_0$  the modes are normalized and b) if  $m^2 = u_0$  the analysis must be performed case by case. As we do not know anything about the warp factor  $\sigma(y)$ , these questions can not be solved yet. Thus, let us obtain a solution for the zero-mode and carry out the discussion of localization.

### 3.1.1 Localization of Zero-Mode – $\hat{\mathcal{A}}_0^\mu$

Let us propose a zero-mode solution for equation (3.11) as  $\chi_0 = e^{a\sigma(y)}$ . It is easy to see that it will satisfy (3.11) when  $a = c_1$  and  $a^2 = c_2$ . These conditions can be reduced to the condition  $c_1^2 = c_2$  and can always be satisfied by adjusting the coupling constants  $\lambda_1$  and  $\lambda_2$  in the action (3.7). This possible values of  $\lambda_1$  and  $\lambda_2$  will be analyzed latter. An important consequence of the condition  $c_1^2 = c_2$  is that it allows us to factor Eq. (3.11) as

$$[-\partial_k + c_1 \partial_k \sigma(y)] [\partial^k + c_1 \partial^k \sigma(y)] \chi(y^j) = m^2 \chi(y^j). \quad (3.15)$$

This is analogous to a supersymmetric quantum mechanical problem, such that we can conclude that there are not tachyonic modes in the spectrum, *i.e.*,  $m^2 \geq 0$ . About the normalization of zero-mode  $\chi_0(y) = e^{a\sigma(y)}$ , we can infer from Eq. (3.13) that  $\chi_0^2$  should go to zero faster than  $|y^j|^{-d}$ , when  $|y^j| \rightarrow \infty$ . This is the most we can obtain by considering an arbitrary  $\sigma(y)$ .

In order to reach some conclusion about the localization of zero mode we will need of more information about our system. As said before, we can obtain information about normalization by analyzing the asymptotic behavior of  $U$ . Note that our effective potential (3.14) depends only on the background warp factor  $\sigma(y)$ , so we must specify, at least, the behavior of  $\sigma(y)$  when  $|y^j| \rightarrow \infty$ . Since we look for RS-like models we will required an asymptotically *AdS* background. This means that we must consider  $\lim_{|y^j| \rightarrow \infty} R = -\kappa$ , with  $\kappa > 0$  and constant. This leads to the following asymptotic behavior for  $\sigma(y)$ :

$$\lim_{|y^j| \rightarrow \infty} \sigma(y^j) = -\ln \left( \sum_j \beta_j |y^j| \right), \quad (3.16)$$

with  $\beta_j$ 's constants. With (3.16) we finally obtain from (3.14) that  $\lim_{|y| \rightarrow \infty} U(y^j) = 0$ . Therefore the analysis of the potential is not conclusive about the localization of massless mode ( $\mathbf{m}^2 = 0$ ) and we must analyze the solution of the equations of motion. By plugging (3.16) in our solution to the zero-mode, we can get the asymptotic behavior of our solution which is given by

$$\lim_{|y^j| \rightarrow \infty} \chi_0^2(y^j) = \left( \sum_j \beta_j |y^j| \right)^{-2a}. \quad (3.17)$$

If now we substitute this in the integral (3.13), we can conclude that it will be finite if we have  $2a = 2c_1 > d$ , where  $d$  is the number of large transverse extra dimensions. Therefore we have two conditions given by

$$c_1^2 = c_2 \quad \text{and} \quad 2c_1 > d \quad (3.18)$$

which must be imposed to our system.

With the conditions (3.18) we can analyze for which range of parameters  $\lambda_1$  and  $\lambda_2$  the massless mode ( $U(1)$  gauge field) can be confined on the brane. For this we use the relations (3.5) and (3.9) to obtain explicit forms of  $c_1$  and  $c_2$ , which are given by

$$c_1 = \frac{D-4}{2} - 2\lambda_1(D-1) - \lambda_2 \quad \text{and} \quad c_2 = \frac{(D-4)^2}{4} - (\lambda_1(D-1) + \lambda_2)(D-2). \quad (3.19)$$

Let us list some particular cases:

- (i) *Free gauge field* ( $\lambda_1 = \lambda_2 = 0$ ) - From Eq. (3.19) we obtain  $2c_1 = (D-4)$ . Thus, as we must have  $2c_1 > d$  in order to confine  $\hat{\mathcal{A}}^\mu$ , we get  $(D-4) > d$ . For example, if we consider a **3**-brane ( $d = D-4$ ), we can conclude that the free gauge field cannot be localized for any number of extra dimensions.
- (ii) *Ricci scalar coupling* ( $\lambda_2 = 0$ ) - When we use that  $c_1^2 = c_2$ , we obtain that  $\lambda_1 = \frac{D-6}{4(D-1)}$  and by using the localization condition  $2c_1 > d$ , we obtain  $2 > d$ . Therefore, the gauge field is localized only for models with co-dimension one. It is important to stress that for this and the next cases, *Hodge duality* symmetry does not apply due to the presence of the interaction term.
- (iii) *Ricci tensor coupling* ( $\lambda_1 = 0$ ) - this allows us to write  $2c_1 = (D-4) - 2\lambda_2 > d$ . From  $c_1^2 = c_2$  we get  $\lambda_2 = -2$  and the localization of the transverse sector  $\hat{\mathcal{A}}^\mu$  occurs for any number of extra dimensions.
- (iv) Finally, we have the case where  $\lambda_1$  and  $\lambda_2$  are non-zero. For this situation, there is a sub case where we can impose  $\lambda_1 = -\frac{1}{2}\lambda_2$ . This constraint allows us to combine the interaction terms with the scalar and the Ricci tensor in action (3.7) to generate the Einstein tensor. This particular coupling gives us the same conclusions of the case (i). For the general situation, we obtain from  $c_1^2 = c_2$  the relation

$$\lambda_2^{(\pm)} = -2\lambda_1(D-1) - 1 \pm \sqrt{\lambda_1(D-1)(D-2) + 1} \quad (3.20)$$

and consequently

$$c_1^{(\pm)} = \frac{D-2}{2} \pm \sqrt{\lambda_1(D-1)(D-2)+1}. \quad (3.21)$$

By imposing  $2c_1 > d$  we get

$$\lambda_1 \geq -\frac{1}{(D-1)(D-2)}, \quad \text{for } c_1^{(+)} \quad (3.22)$$


and

$$-\frac{1}{(D-1)(D-2)} \leq \lambda_1 < -\frac{4-(D-d-2)^2}{4(D-1)(D-2)}, \quad \text{for } c_1^{(-)}. \quad (3.23)$$


Therefore the localization can always be accomplished by just imposing that  $\lambda_1$  satisfy the above conditions. For example, if we choose  $d = D - 4$  we obtain that  $\lambda_1$  must belongs to the range  $[-\frac{1}{(D-1)(D-2)}, \infty)$  for  $c_1^{(+)}$  or  $[-\frac{1}{(D-1)(D-2)}, 0)$  for  $c_1^{(-)}$ . We should point to the fact that in this case the parameters  $\lambda_1, \lambda_2$  are not completely fixed. This will be important when we consider the localization of scalar components of the vector field in the next section.

The above discussion reveals some important points about the gauge field localization. Among the main results, we stress that the background is completely generic. We did not need to specify whether the model consists of a delta or a thick-like brane. Thus, this localization mechanism does not suffer from the ‘problem’ of being sensitive to the thickness of the brane. Another important point is that the existence of a zero-mode solution for the gauge field imply that the model has no tachyon. These are the reasons why the geometric coupling has a general validity, allowing the confinement of  $U(1)$  gauge field for an wide variety of braneworld models.

To conclude the discussion about transverse gauge field, we will carry out a comparative analysis between our zero-mode  $\chi^{(\text{gauge})}(y) = \exp[c_1\sigma(y)]$  and the one found for the gravity field in Ref. [53]. Since the consistence of gravity is the starting point of braneworld models, it is interesting to analyze if its localization imply the localization of the gauge field. In ref. [53], Csaba Csáki *et.al.* performed a study about the universal aspects of gravity localization in braneworld models. From the standard Einstein-Hilbert action, the authors studied the metric fluctuations on the background like that of Eq. (3.2). An interesting point is that the zero-mode found for the gravitational field is given by  $\psi^{(\text{gravity})}(y) = \exp\left[\left(\frac{D-2}{2}\right)\sigma(y)\right]$ . By imposing that this solution is square integrable they reach a general condition for gravity localization given by  $D-2 > d$ . In the case of the gauge field, in Eq. (3.18) we also found a similar condition given by  $2c_1 > d$ . In this way, if  $2c_1 \geq (D-2)$ , the localization of gravitational field is enough to ensure the localization of gauge field. Looking at the cases (i-iv) above discussed, only the configurations (iii) and (iv) can satisfy  $2c_1 \geq (D-2)$ . For the case (iii)  $2c_1 = D \geq (D-2)$ , therefore we get that the localization of gravity implies the localization of the gauge field. For the case (iv) we see that only the solution  $c_1^{(+)}$  can satisfy the relation  $2c_1 \geq (D-2)$ . In Figs. (1) and (2) we give a plot of values of  $\lambda_1$  and  $\lambda_2$ , as a function of the dimension  $D$ , which allow the localization of gauge field for some cases discussed above.



**Figure 1:** Case (ii) of the sections (3.1.1) and (3.2.1) - The blue solid line gives the values of  $\lambda_1(D)$  which allow the localization of the  $\hat{\mathcal{A}}^\mu$  field. The region below the dashed line shows the values which allow the localization of the  $\mathcal{B}^j$  fields. Also, the region below the dotted line are the values of  $\lambda_1(D)$  for which the localization of gravity ensures the localization of  $\hat{\mathcal{A}}^\mu$ .



**Figure 2:** Case (iii) of the sections (3.1.1) and (3.2.1) - The blue solid line gives the values of  $\lambda_2(D)$  which allow the localization of the  $\hat{\mathcal{A}}^\mu$  field. The region below the dashed line shows the values which allow the localization of the  $\mathcal{B}^j$  fields. Also, the region below the dotted line are the values of  $\lambda_2(D)$  for which the localization of gravity ensures the localization of  $\hat{\mathcal{A}}^\mu$ .

### 3.2 Localization of the Scalar Components – $\mathcal{B}^j$

It is common in the literature to consider the trivial solution for the fields  $\mathcal{B}_j$  [see section (2)]. The reason is that if these fields are not localized, its backreaction could modify the background. Despite of this, in this section we must consider the more general case and study the possibility of localizing the components  $\mathcal{B}_j$ . We will emphasize on the possibility that these fields can be confined simultaneously with the transverse gauge field  $\hat{\mathcal{A}}^\mu$ . We must point that the localization of these components have no consequence over the Coulomb law. Actually, by Eq. (3.3), an observer at the brane sees these components as *scalar fields* and these have no relation to Coulomb's scalar potential. However, the effective theory for this fields over the brane could be interpreted as Higgs fields or maybe as dark energy, for example.

In order to study the localization of the scalar components  $\mathcal{B}_j$ , we will carry out the variation of action (3.1) with respect to  $\mathcal{A}^M$ . By doing this, we get the following equation of motion

$$\frac{1}{\sqrt{-g}} \partial_N \left( \sqrt{-g} g^{MN} g^{PQ} \mathcal{F}_{MP} \right) = \lambda_1 R g^{MQ} \mathcal{A}_M + \lambda_2 g^{MQ} g^{NP} R_{MN} \mathcal{A}_P, \quad (3.24)$$

and due to the anti-symmetry of  $\mathcal{F}_{MP}$ , we also obtain the constraint

$$\partial_Q \left( \lambda_1 R \sqrt{-g} g^{MQ} \mathcal{A}_M + \lambda_2 \sqrt{-g} g^{MQ} g^{NP} R_{MN} \mathcal{A}_P \right) = 0. \quad (3.25)$$

From these expressions, the equation of motion for the scalar fields  $\mathcal{B}^j$  can be obtained. For this we choose  $Q = k$  in the Eq. (3.24) to obtain

$$\begin{aligned} \partial_\mu \partial^\mu \mathcal{B}^j + e^{-(D-4)\sigma} \partial_k \left( e^{(D-4)\sigma} \mathcal{B}^{kj} \right) \\ - \left( \lambda_1 R \delta_k^j + \lambda_2 e^{-2\sigma} R_k^j \right) e^{2\sigma} \mathcal{B}^k = \partial^j \partial_\mu \mathcal{A}^\mu. \end{aligned} \quad (3.26)$$

We also write the constraint (3.25) as

$$\begin{aligned} e^{-(D-2)\sigma} \partial_k \left[ e^{(D-2)\sigma} \left( \lambda_1 R \delta_k^j + \lambda_2 e^{-2\sigma} R_k^j \right) \mathcal{B}^k \right] \\ + \left[ \lambda_1 R - \lambda_2 e^{-2\sigma} h(y) \right] \partial_\mu \mathcal{A}^\mu = 0. \end{aligned} \quad (3.27)$$

In the above equations  $\mathcal{B}^{kj} = \partial^k \mathcal{B}^j - \partial^j \mathcal{B}^k$  and the indexes are lowered/raised using the Minkowski metric (remember that  $\eta_{jk} = \delta_{jk}$ ). Looking at the equations (3.26) and (3.27), we realize that the solution for the fields  $\mathcal{B}^j$  cannot be obtained as directly as for the field  $\hat{\mathcal{A}}^\mu$ . Actually, since the equations (3.26) are coupled, its complete solution is model-dependent, *i.e.*, the warp factor must be known. Despite of this, some general properties of the system can be obtained by studying the asymptotic behavior of Eqs. (3.26) and (3.27).

From now on we will look for a solution of Eqs. (3.26) and (3.27) when  $|\mathbf{y}^j| \rightarrow \infty$ . In this way, we can study under what conditions they are convergent and square integrable. In this limit, the asymptotic warp factor (3.16) can be used and thus we get some simplifications on our system. First we see from Eq. (3.4) that  $R_{jk}$  can be written as

$$\begin{aligned} R_{jk} &= (D-2) [-\partial_j \partial_k \sigma(y) + \partial_j \sigma(y) \partial_k \sigma(y)] - h(y) \eta_{jk} \\ &\equiv (D-2) \Omega_{jk} - h(y) \eta_{jk}, \end{aligned} \quad (3.28)$$

with  $h(y)$  defined in Eq. (3.9). By using the asymptotic warp factor (3.16) we can find

$$\Omega_{jm} = -\partial_m \partial_j \sigma(y) + \partial_m \sigma(y) \partial_j \sigma(y) \rightarrow 0, \quad (3.29)$$

and thus (3.28) gets the form

$$R_{jk} \rightarrow h(y) \eta_{jk}. \quad (3.30)$$

Another simplification is that  $\lambda_1 R - \lambda_2 h(y) e^{-2\sigma} \equiv g(y)$  goes to a constant value  $g(y) \rightarrow C_1$ . With these simplifications the constraint (3.27) simplifies to

$$e^{-(D-2)\sigma} \partial_k \left[ e^{(D-2)\sigma} \mathcal{B}^k \right] + \partial_\mu \mathcal{A}^\mu = 0. \quad (3.31)$$

Substituting this in Eq. (3.26) we finally get

$$e^{-(D-4)\sigma} \partial_k \left[ e^{(D-4)\sigma} \partial^k \mathcal{B}^j \right] + \partial_\rho \partial^\rho \mathcal{B}^j - g(y) e^{2\sigma} \mathcal{B}^j \\ + 2 \partial^j \mathcal{B}^k \partial_k \sigma + (D-2) \mathcal{B}^k \partial^j \partial_k \sigma = 0. \quad (3.32)$$

The above equation is the main one which we will try to solve. In this way, we intend, at least, to get the some conditions for which the scalar components can be localized on the brane.

Let us propose the transformation  $\mathcal{B}^k(x, y) = B^k(x, y) e^{-\frac{(D-2)}{2}\sigma}$  in Eq. (3.32). With this, the last two terms can be eliminated and we obtain

$$\partial_k \partial^k B^j(x, y) - \left[ \frac{(D-2)}{2} \partial_k \partial^k \sigma + \frac{(D-2)(D-6)}{4} \partial_k \sigma \partial^k \sigma + g(y) e^{2\sigma} \right] B^j(x, y) \\ + \partial_\rho \partial^\rho B^j(x, y) + 2 \left[ \partial^j B^k(x, y) - \partial^k B^j(x, y) \right] \partial_k \sigma = 0. \quad (3.33)$$

Note that the above equation has a very convenient form. We can carry out a contraction of Eq. (3.33) with  $\partial_j \sigma$  to obtain an equation that can be easily solved. By doing this and using the asymptotic warp factor (3.16) we obtain

$$\left[ \frac{(D-2)}{2} \left( \partial_k \partial^k \sigma + \frac{(D-6)}{2} \partial_k \sigma \partial^k \sigma \right) + g(y) e^{2\sigma} \right] \Phi(x, y) \\ - \partial_k \partial^k \Phi(x, y) = \partial_\rho \partial^\rho \Phi(x, y), \quad (3.34)$$

where we have defined

$$\sum_j \text{sgn}(y^j) \beta_j B^j(x, y) \equiv \Phi(x, y) \quad (3.35)$$

with  $\text{sgn}(y^j)$  the sign function defined in standard way. This definition can be used in Eq. (3.33) in order to remove the coupled terms. By doing this, we get a non-homogeneous source-like term as follows

$$\partial_k \partial^k B^j(x, y) - \left[ \frac{(D-2)}{2} \partial_k \partial^k \sigma + \frac{(D-2)(D-6)}{4} \partial_k \sigma \partial^k \sigma + g(y) \right] B^j(x, y) \\ + \partial_\rho \partial^\rho B^j(x, y) - 2 \partial^k B^j(x, y) \partial_k \sigma = 2 e^\sigma \partial^j \Phi(x, y). \quad (3.36)$$

Note that the source of Eq. (3.36) is given by the solution of Eq. (3.34). Therefore, in order to solve (3.36) we first need to solve Eq. (3.34). Before this we can use the known fact of partial differential equation that the most general solution is given by the sum  $B^j(x, y) = \tilde{B}^j(x, y) + \tilde{B}_p^j(x, y)$ , where  $\tilde{B}^j(x, y)$  is a solution to the homogeneous equation

$$\partial_k \partial^k \tilde{B}^j(x, y) - \left[ \frac{(D-2)}{2} \partial_k \partial^k \sigma + \frac{(D-2)(D-6)}{4} \partial_k \sigma \partial^k \sigma + g(y) \right] \tilde{B}^j(x, y) \\ + \partial_\rho \partial^\rho \tilde{B}^j(x, y) - 2 \partial^k \tilde{B}^j(x, y) \partial_k \sigma = 0; \quad (3.37)$$

and  $\tilde{B}_p^j(x, y)$  is a particular solution of the complete equation

$$\begin{aligned} \partial_k \partial^k \tilde{B}_p^j(x, y) - \left[ \frac{(D-2)}{2} \partial_k \partial^k \sigma + \frac{(D-2)(D-6)}{4} \partial_k \sigma \partial^k \sigma + g(y) \right] \tilde{B}_p^j(x, y) \\ + \square \tilde{B}_p^j(x, y) - 2 \partial^k \tilde{B}_p^j(x, y) \partial_k \sigma = 2e^\sigma \partial^j \Phi(x, y); \end{aligned} \quad (3.38)$$

In this way, Eq. (3.33) is reduced to the above system of equations (3.34), (3.37) and (3.38). The next step is to perform a separation of variables of these equations. First let us propose  $\Phi(x, y) = \theta(x)\zeta(y)$ , thus we get the two equations

$$\square_x \theta(x) = M^2 \theta(x); \quad (3.39)$$

and

$$-\partial_k \partial^k \zeta(y) + U(y) \zeta(y) = M^2 \zeta(y), \quad (3.40)$$

where

$$U(y) = \frac{(D-2)}{2} \left( \partial_k \partial^k \sigma + \frac{(D-6)}{2} \partial_k \sigma \partial^k \sigma \right) + g(y) e^{2\sigma}. \quad (3.41)$$

If now we substitute  $\Phi(x, y) = \theta(x)\zeta(y)$  in Eq. (3.38) we see that the only way to separate the variables is by choosing  $\tilde{B}_p^j(x, y) = \theta(x)Z^j(y)$ . Therefore, the final form of the non-homogeneous equation is given by

$$\begin{aligned} \partial_k \partial^k Z^j(y) - \left[ \frac{(D-2)}{2} \partial_k \partial^k \sigma + \frac{(D-2)(D-6)}{4} \partial_k \sigma \partial^k \sigma + g(y) \right] Z^j(y) \\ + M^2 Z^j(y) - 2 \partial^k Z^j(y) \partial_k \sigma = 2e^\sigma \partial^j \zeta(y). \end{aligned} \quad (3.42)$$

Thus our final solution is in the form

$$B^j(x, y) = \tilde{B}^j(x, y) + \theta(x)Z^j(y) \quad (3.43)$$

where, as said before,  $\tilde{B}^j(x, y)$  is a solution of Eq. (3.37),  $\theta(x)$  the solution of Eq. (3.39) and  $Z^j(y)$  the solution of Eq. (3.42). At this moment, the issue of zero-mode localization for the fields  $B^j(x, y)$  can be attained.

### 3.2.1 Localization of Zero-Mode – $\mathcal{B}_0^j$

Before to study the zero-mode localization for the scalar components, we should point that from definition (3.35) and by using the Eq. (3.39), we see that

$$\square_x \Phi(x, y) = M^2 \Phi(x, y) = \sum_j \text{sgn}(y^j) \beta_j \square_x B^j(x, y). \quad (3.44)$$

If we are studying the zero-mode ( $M^2 = 0$ ) of the field  $\Phi(x, y)$ , then we should have  $\sum_j \text{sgn}(y^j) \beta_j \square_x B^j(x, y) = 0$ . With this, we conclude that  $\square_x B^j(x, y) = 0$  for each field  $B^j(x, y)$  independently. Therefore, the massless mode analysis of  $\Phi(x, y)$  corresponds to the massless mode analysis of  $B^j(x, y)$ .

Now we can turn to the zero-mode solution of the fields  $B^j(x, y)$ . First, in view of the Eq. (3.42), we need to solve Eq. (3.40) for the zero-mode  $\zeta_0(y)$ . By proposing the ansatz  $\zeta_0(y) = e^{\tilde{b}\sigma(y)}$  for Eq. (3.40) with  $M^2 = 0$ , we see that this is a solution when

$$\tilde{b}_{\pm} = -\frac{1}{2} \pm \frac{1}{2} [(D-3)^2 - 4(D-1)(\lambda_2 + \lambda_1 D)]^{\frac{1}{2}} \quad (3.45)$$

Note that Eq. (3.40) is a Schrödinger-like equation with potential  $U$  given in Eq. (3.41). The potential  $U$  is even by spatial inversion ( $y^j \rightarrow -y^j$ ), and this generates a solutions for the field  $\Phi(x, y)$  with well-defined parity (even or odd). We saw earlier that the warp factor (3.16) is an even function of  $y^j$ . Therefore, since our solution  $\zeta_0(y)$  is a power of the warp factor, we find that the field  $\Phi_0(x, y)$  is even by the exchange  $y^j \rightarrow -y^j$ .

With  $\zeta_0(y)$  obtained above, we can now look for a solution of the non-homogeneous equation (3.42), which becomes

$$\begin{aligned} \partial_k \partial^k Z_0^j(y) - \left[ \frac{(D-2)}{2} \partial_k \partial^k \sigma + \frac{(D-2)(D-6)}{4} \partial_k \sigma \partial^k \sigma + g(y) \right] Z_0^j(y) \\ - 2\partial^k Z_0^j(y) \partial_k \sigma = 2\tilde{b} e^{(\tilde{b}+1)\sigma(y)} \partial^j \sigma(y). \end{aligned} \quad (3.46)$$

To solve the above equation first note that, as discussed previously, the field  $\Phi_0(x, y)$  has even parity. Since we also have that  $\Phi(x, y) = \sum_j \text{sgn}(y^j) \beta_j B^j(x, y)$ ,  $\text{sgn}(y^j) B_0^j(x, y)$  must also be even. This implies that  $Z_0^j(y)$  must be odd. With this it is easy to check that the solution for (3.46) is obtained from the ansatz  $Z_0^j(y) = \text{sgn}(y^j) |y^j| e^{(1+\tilde{b})\sigma(y)}$  (without sum in  $j$ ).

About the solution for the homogeneous Eq. (3.37), this can be found by performing a very similar treatment to that used in section (3.1.1). By doing this, we find

$$\tilde{B}^j(x, y) = \text{sgn}(y^j) e^{(\tilde{b}+1)\sigma} \bar{B}_0^j(x). \quad (3.47)$$

Finally, we arrive at the final solution for our system, given by

$$\begin{aligned} \mathcal{B}_0^j(x, y) &= e^{-\frac{(D-2)}{2}\sigma} B_0^j(x, y) \\ &= \text{sgn}(y^j) e^{\left[\tilde{b} - \frac{(D-4)}{2}\right]\sigma} \left[ \bar{B}_0^j(x) + \theta_0(x) |y^j| \right] \equiv \text{sgn}(y^j) f^j(x, |y|). \end{aligned} \quad (3.48)$$

Beyond that, by definition (3.35) and due to the form of the warp factor (3.16), the fields  $B_0^j(x, y)$  must satisfy

$$\sum_j \beta_j \bar{B}_0^j(x) = 0. \quad (3.49)$$

Eq. (3.49) gives us a constraint on the fields  $\bar{B}_0^j(x)$ , which at the end leads to the correct number of degrees of freedom. After all this, we can analyze under what conditions the zero-mode of the scalar components can be confined on the brane.

In the appendix (A) we split the action (3.1) in two sectors given by Eq. (A.4) for the gauge field  $\hat{\mathcal{A}}^\mu$  and Eq. (A.5) that contain the scalar fields  $\mathcal{B}^j$ . In this last expression, the kinetic term that we must use to deal with the localization of massless mode is given by

$$S_0[\mathcal{B}^j] = -\frac{1}{2} \int d^{(D-d)} x d^d y \sqrt{-g} g^{\mu\nu} g^{jk} \partial_\mu \mathcal{B}_j \partial_\nu \mathcal{B}_k. \quad (3.50)$$

We are interested only in the convergence of the integral when  $|\mathbf{y}^j| \rightarrow \infty$ . Thus, for large  $|\mathbf{y}^j|$ ,  $\theta_0(\mathbf{x})|\mathbf{y}^j|$  is the dominant term of our solution (3.48). Therefore, if it is square integrable, all the solution will be. In our case, the  $\mathbf{y}^j$  integral in the action (3.50) will be finite when  $2\tilde{\mathbf{b}}_{\pm} > d + 2$ . With this condition, we can analyze each of the cases bellow:

- (i) *Free scalar field* ( $\lambda_1 = \lambda_2 = 0$ ) - from the condition  $2\tilde{\mathbf{b}}_{\pm} > d + 2$ , we get  $\pm(D - 3) > d + 3$ . For the case  $D = 4 + d$ , for example, it is easy to see that it is not possible to localize the scalar components.
- (ii) *Ricci scalar coupling* ( $\lambda_2 = 0$ ) - In this case, the convergence condition  $2\tilde{\mathbf{b}}_{\pm} > d + 2$  leads to a constraint for  $\lambda_1$  given by  $\lambda_1 < -\frac{(d+3)^2 - (D-3)^2}{4D(D-1)}$ . With this, only the solution  $\tilde{\mathbf{b}}_+$  allows a zero-mode localized for the scalar components. The Fig. (1) shows the values of  $\lambda_1(D)$  that allow the localization of scalar components over a **3**-brane ( $D = d + 4$ ). Remember that for the gauge field  $\hat{\mathcal{A}}_{\mu}$ , we obtained  $\lambda_1 = \frac{D-6}{4(D-1)}$ . From the Fig. (1), we see that this value of  $\lambda_1$  is not inside the region that allows the localization of two sectors simultaneously on the **3**-brane.
- (iii) *Ricci tensor coupling* ( $\lambda_1 = 0$ ) - For this, we get the constrain  $\lambda_2 < -\frac{(d+3)^2 - (D-3)^2}{4(D-1)}$  and, again, only the solution  $\tilde{\mathbf{b}}_+$  allows a zero-mode localized for the scalar components. For this case, the value  $\lambda_2 = -2$ , defined in item (iii) of section (3.1.1), obeys this relation for any number of transverse extra dimensions. In this way, there is the possibility of 'trapping' simultaneously both sectors on a **3**-brane with this kind of coupling. This can also be seen in Figure (2).
- (iv) Finally, with  $\lambda_1$  and  $\lambda_2$  nonzero, we get  $\lambda_1 D + \lambda_2 < -\frac{(d+3)^2 - (D-3)^2}{4(D-1)}$ . Thus, we also can have both sectors localized simultaneously on a **3**-brane.

Of course, we only can ensure that the scalar components  $\mathcal{B}^j$  are localized if we obtain the complete solution for them. Anyway, we get at least one general condition that must be imposed on our system such that we get confined massless scalar fields. In the next section, we will apply this general results for a specific braneworld model.


## 4 Application: Intersecting Brane Model

In section (3), the general aspects of the  $D$ -vector field localization in a generic braneworld model was studied. Among the main general results, we obtained that the confinement of this higher dimensional vector field generates an effective theory for one  $U(1)$  gauge field  $\hat{\mathcal{A}}^{\mu}$ . Beyond this, there is an indicative that an effective scalar theory related to components  $\mathcal{B}^j$  can be also obtained. For the gauge field, it was found a zero-mode solution which is valid for any warp factor. However for the scalar components, it was only possible to find an asymptotic zero-mode solution. Here, we will apply the general results obtained in last section to a specific braneworld model and a more detailed discussion on the scalar components localization will be carried out. We will use the intersecting brane model developed by Arkani-Hamed *et. al.* [19]. In this model, the background metric is

given by Eq. (3.2) with the warp factor

$$\sigma(y^j) = -\ln\left(1 + k \sum_j |y^j|\right). \quad (4.1)$$

The authors show that gravity is localized in a warped model generated by the intersection of  $d$  delta-like  $(D-2)$ -branes. In the figure (3) it is shown an example with two transverse extra dimensions. For this case we have two delta-like 4-branes intersecting one each other generating one 3 dimensional brane.



**Figure 3:** Intersecting branes model with two transverse extra dimensions.

We will focus our attention on the discussion of the massless scalar components  $\mathcal{B}_j$  of section (3.2.1). This is because the solution for these components were obtained only in an asymptotic approximation, and it would be interesting to find a complete solution for these fields. In this way, we will be able to verify whether the results for scalar components can lead to a reasonable effective theory on the brane. Beyond this, from now on we will fix our brane with 3 spatial dimensions, *i.e.*,  $D = d + 4$ . As the branes in (4.1) are delta-like, the zero-mode solutions  $\mathcal{B}_0^k = B^k(x, y)e^{-\frac{(D-2)}{2}\sigma}$ , with  $B^k(x, y)$  given by Eq. (3.48), must be valid for all  $y^j \neq 0$ . Thus, we only have to obtain the boundary conditions imposed by the branes on such solutions. An important point of this braneworld model is that each  $(D-2)$ -brane impose one boundary condition on the fields  $\mathcal{B}_j$ . Beyond this, at the intersection points we also must impose another boundary condition. Looking at the figure (3), it means that each 4-brane will introduce one boundary condition and the 3-brane will impose another one.

As we saw, all the analysis for scalar fields was performed from equations (3.26) and (3.27). For the present case, the warp factor (4.1) allows us to write the following common factor of these equations

$$\lambda_1 R e^{2\sigma} \delta_j^k + \lambda_2 R_j^k = C_0 e^{\sigma(y)} \delta_j^k \sum_m \delta(y^m) - C_1 e^{2\sigma} \delta_j^k + C_2 e^{\sigma} \delta(y^j) \delta_j^k. \quad (4.2)$$

In the above expression the constants  $C_0$ ,  $C_1$  and  $C_2$  are given by

$$\begin{aligned} C_0 &= 2k [\lambda_2 + 2\lambda_1 (D-1)]; \\ C_1 &= k^2 (D-1) (D-4) (\lambda_2 + \lambda_1 D); \\ C_2 &= 2k\lambda_2 (D-2). \end{aligned} \quad (4.3)$$

From this we can carry out some considerations about the constraint (3.27). It can be written in the form

$$e^{-(D-2)\sigma} \partial_k \left[ e^{(D-2)\sigma} \left( C_0 e^{-\sigma} \sum_m \delta(y^m) - C_1 + C_2 e^{-\sigma} \delta(y^k) \right) \mathcal{B}^k \right] + \left[ C_0 e^{-\sigma} \sum_m \delta(y^m) - C_1 \right] \partial_\mu \mathcal{A}^\mu = 0. \quad (4.4)$$

Due to the presence of Dirac's delta, the following procedures can be performed. We will consider the above constraint for  $y^j \neq 0$

$$\partial_\mu \mathcal{A}^\mu + e^{-(D-2)\sigma} \text{sgn}(y^k) \partial_k \left[ e^{(D-2)\sigma} f^k \right] = 0, \quad (4.5)$$

where  $f^k$  was defined in Eq. (3.48). Next, we assume that it is valid for all  $y^j$ . In order to this to be valid we must substitute Eq. (4.5) in Eq. (4.4) such that it becomes

$$2C_0 \sum_{m \neq k} \delta(y^m) e^{-\sigma} \sum_k f^k \delta(y^k) - C_0 k \sum_{m \neq k} \delta(y^m) \sum_k f^k - 2C_1 \delta(y^k) f^k = 0. \quad (4.6)$$

With this we can consider that we have solved our constraint and the above equation will impose boundary conditions on the fields  $\mathcal{B}^k$ . Beyond this boundary conditions, the fields  $\mathcal{B}^k$  must also obey a set of boundary conditions obtained from equations of motion (3.26). We saw that for this intersecting branes model each brane must impose a boundary condition on the fields  $\mathcal{B}^k$ . By using the figure (3) as example, we will carry out one integration of Eqs. (3.26) and (4.6) into the range  $[-\epsilon, \epsilon]$  for each extra coordinate keeping the others fixed, this will give us a set of boundary conditions. Next, we perform an integration in small volume containing the intersection point. This will give us another boundary conditions. For an arbitrary number of extra dimensions we must only extend this procedure.

By using the above procedure to Eq. (4.6) we get the following boundary conditions

$$C_0(D-5) \sum_k f^k(0) = 0; \quad (4.7)$$

$$2C_1 f^n(y^n = 0) - C_0 k \sum_{k \neq n} f^k(y^n = 0) = 0. \quad (4.8)$$

Applying the same to Eq. (3.26) we get

$$(C_0 + C_2) \mathcal{B}_0^k(y^k = 0, y^j) \Big|_{y^j=c; j \neq k} = 0; \quad (4.9)$$

$$C_0 e^{\sigma(y)} \mathcal{B}_0^k(y^m = 0, y^j) \Big|_{y^j=c; m \neq j} = \left[ \frac{d}{dy^m} \mathcal{B}_0^k(y^m, y^j) \right]_{y^m=-\frac{\epsilon}{2}, y^j=c; m \neq j}^{y^m=\frac{\epsilon}{2}}; \quad (4.10)$$

$$\frac{(dC_0 + C_2)}{2} \mathcal{B}_0^k(0) = k c_1 (d-1) \mathcal{B}_0^k(0) - k c_1 \text{sgn}(y^k = 0) \sum_{\substack{j \\ j \neq k}} \mathcal{B}_0^j(0^+). \quad (4.11)$$

The above equations are all the boundary conditions which must be imposed on the scalar fields. First, the equations (4.9) and (4.11) will be trivially satisfied if we define  $\text{sgn}(\mathbf{0}) = \mathbf{0}$ . Second, the condition (4.10) imposes the constraint  $\mathbf{C}_0 = -2\mathbf{k}(\tilde{\mathbf{b}} + \mathbf{1}) + \mathbf{k}(D - 2)$  on the parameters of the theory. With this and by using Eqs. (3.45) and (4.3) we get

$$\lambda_2^{(\pm)} = -2\lambda_1(D - 1) - 1 \pm \sqrt{\lambda_1(D - 1)(D - 2) + 1}. \quad (4.12)$$

The expression (4.12) is exactly the relation obtained in Eq. (3.20) for the localization of gauge field  $\hat{\mathbf{A}}^\mu$ . This is an excellent result, since we wish to confine both sectors (gauge and scalar fields) simultaneously. Finally, the equation (4.7) gives us  $\sum_{\mathbf{k}} \bar{\mathbf{B}}^{\mathbf{k}}(\mathbf{x}) = \mathbf{0}$  and the Eq. (4.8) leads to  $\boldsymbol{\theta}(\mathbf{x}) = \mathbf{0}$  and  $2\mathbf{C}_1 + \mathbf{C}_0\mathbf{k} = \mathbf{0}$ . This last relation allows us to fix the value of the parameter  $\lambda_1$  in equation (4.12). By doing this we get

$$\lambda_1 = \frac{(D^2 - 5D + 5)(D^2 - 7D + 13)}{(D - 4)^2(D - 2)(D - 1)}. \quad (4.13)$$

With this, all the parameters of the theory were fixed and, fortunately, with both sectors localized. Looking at this value of  $\lambda_1$  we see that is not possible to localize both sectors in the case (iii) discussed in sections (3.1.1) and (3.2.1). The reason is that in case (iii) we have  $\lambda_1 = \mathbf{0}$  and this is possible in Eq. (4.13) only for non-integer values of  $D$ .

Now, the localization of the gauge and the scalar fields for the case (iv) of sections (3.1.1) and (3.2.1) can be performed. Since both solutions are power of the warp factor as follows  $e^{a\sigma(\mathbf{y})}$ . We will carry out the analysis of the values of  $\mathbf{a}$  obtained previously for both sectors.

- (1) *Gauge field* - Case (iv) of section (3.1.1): for this case, the value of  $\lambda_1$  showed in Eq. (4.13) gives us

$$\mathbf{a} = \mathbf{c}_1^{(\pm)} = \frac{D - 2}{2} \pm \sqrt{\lambda_1(D - 1)(D - 2) + 1} = \frac{D - 2}{2} \pm \frac{(D - 3)^2}{(D - 4)}. \quad (4.14)$$


Thus, as discussed in section (3.1.1), we have two solutions for the gauge field. In Fig. (4), we have a plot (solid red line) of the localization condition  $2\mathbf{c}_1 > d = D - 4$ . In addition, we also have the two solutions (4.14):  $\mathbf{c}_1^{(+)}$  (dashed blue line) and  $\mathbf{c}_1^{(-)}$  (dotted black line). From this graphic, it is clear that the only solution that satisfies the localization condition is  $\mathbf{c}_1^{(+)}$ . This conclusion agree with the general results obtained in (3.1.1).

- (2) *Scalar fields* - Case (iv) of section (3.2.1): for this case, the value of  $\lambda_1$  given by Eq. (4.13) allows us to write


$$\mathbf{a} = \tilde{\mathbf{b}}^{(\pm, \pm)} = -\frac{1}{2} \pm \frac{1}{2} \left[ (D - 3)^2 - 4(D - 1)(\lambda_2^{(\pm)} + \lambda_1 D) \right]^{\frac{1}{2}}. \quad (4.15)$$

Where  $\tilde{\mathbf{b}}^{(\pm, +)} = -\frac{1}{2} \pm \sqrt{\dots \lambda_2^{(+)}}$  and  $\tilde{\mathbf{b}}^{(\pm, -)} = -\frac{1}{2} \pm \sqrt{\dots \lambda_2^{(-)}}$ . In this way, we have four solutions for each scalar field. We should point that the localization condition for the scalar components found after the equation (3.50) was obtained considering

$\theta_0(x) \neq 0$ . However the boundary conditions (4.9)-(4.8) led us to deduce that  $\theta_0(x)$  is zero, so we have to consider a change in the localization condition, instead of  $2\tilde{b} > d + 2$ , we should consider  $2\tilde{b} > d$ . It is easily verified by put the solution (3.48), with  $\theta_0(x) = 0$ , in Eq. (3.50). The Fig. (5) shows a plot (solid red line) of the above new localization condition and the four solutions (4.15):  $\tilde{b}^{(+,+)}$  (dashed blue line),  $\tilde{b}^{(+,-)}$  (dashed purple line) and  $\tilde{b}^{(-,\pm)}$  (dotted black lines). Again, as showed graphically, only the solutions  $\tilde{b}^{(+,\pm)}$  can satisfy the localization condition. Therefore only in these cases the scalar field is localized on the **3**-brane.



**Figure 4:** Comparative plot of solutions  $c_1^{(+)}$  (dashed blue line),  $c_1^{(-)}$  (dotted black line) and the localization condition - L.C. (red line).



**Figure 5:** Comparative plot of solutions  $\tilde{b}^{(+,+)}$  (dashed blue line),  $\tilde{b}^{(+,-)}$  (dashed purple line),  $\tilde{b}^{(-,\pm)}$  (dotted black line) and the localization condition - L.C. (red line).

It is important to stress that the localization of both sectors was only possible in this model because the interaction terms with the tensor and the Ricci scalar were present. Beyond this, a complete solution for the scalar components is model-dependent, therefore it is not possible ensure that the scalar components will be really confined for others warp factors.

To conclude, let us obtain the effective theory of these scalar components on the **3**-brane. In the discussion of last paragraph, we obtained that  $\theta(x)$  is zero, so the solution (3.48) is given by

$$\mathcal{B}_0^j(x, y) = e^{-\frac{(D-2)}{2}\sigma(y^j)} B^j(x, y) = \text{sgn}(y^j) e^{\left(\tilde{b}^{(+)} - \frac{(D-4)}{2}\right)\sigma(y^j)} \bar{B}^j(x). \quad (4.16)$$

With this, we obtain the following effective action

$$\begin{aligned} S_0[\mathcal{B}^j] &= -\frac{1}{2} \int d^4x d^d y e^{(D-4)\sigma(y^j)} \partial_\mu \mathcal{B}_{0j} \partial^\mu \mathcal{B}_0^j \\ &= -\frac{1}{2} \int d^d y e^{2\tilde{b}^{(+)}\sigma(y^j)} \int d^4x \partial_\mu \bar{B}^j(x) \partial^\mu \bar{B}_j(x) \end{aligned} \quad (4.17)$$

with the integrals over the extra dimensions finite. In addition, we must impose the constraint

$$\sum_{j=1}^d \bar{B}^j(x) = 0. \quad (4.18)$$

In the particular case of  $D = 7$  and  $d = 3$ , Eq. (4.17) can be written as

$$S_0 \sim -\frac{1}{2} \int d^4x [\partial^\mu \bar{B}_0^1 \partial_\mu \bar{B}_0^1 + \partial^\mu \bar{B}_0^2 \partial_\mu \bar{B}_0^2 + \partial^\mu \bar{B}_0^3 \partial_\mu \bar{B}_0^3]. \quad (4.19)$$

Now, we can use (4.18) and redefine the fields such that the effective action for the zero mode of the scalar field is given by

$$S_0 \sim -\int d^4x \left[ \frac{1}{2} \partial^\mu \bar{\bar{B}}_0^1(x) \partial_\mu \bar{\bar{B}}_0^1(x) + \frac{1}{2} \partial^\mu \bar{\bar{B}}_0^2(x) \partial_\mu \bar{\bar{B}}_0^2(x) \right]. \quad (4.20)$$

That is the action for free scalar fields.

## 5 Final Remarks

In this work the general properties of the vector field localization on braneworlds with co-dimension  $d$  in asymptotically AdS spacetime were studied. It was considered a  $D$  dimensional bulk with a generic conformally flat metric  $e^{2\sigma}(\eta_{\mu\nu} dx^\mu dx^\nu + \delta_{ij} dy^i dy^j)$ , which the warp factor depends only on the transverse extra dimensions  $\sigma(y)$ . In this context, we use a non-minimal coupling between gravity and the  $D$ -vector field  $\mathcal{A}_N = (\mathcal{A}_\mu, \mathcal{B}_k)$  as a localization mechanism. The study of zero-mode localization for the fields  $\mathcal{A}_\mu$  (gauge field) and  $\mathcal{B}_k$  (scalar fields) was separated in some particular cases: (a) non-minimal coupling only with *Ricci scalar*; (b) only with *Ricci tensor*; and (c) the case with both interaction terms.

In section (3.1), we analyzed the abelian vector field problem, where the features of the background geometry allowed us to obtain a Schrödinger-type equation with the potential given by (3.14). Such equation has a general analytic solution for the massless mode given by a function of the warp factor,  $\chi_0^{(gauge)} = e^{a\sigma(y)}$ , which is a solution when  $a^2 = c_1^2 =$

$\mathbf{c}_2$ , with  $\mathbf{c}_1$  and  $\mathbf{c}_2$  presented in Eq. (3.19). Due to its shape, this solution becomes valid for a wide variety of warp factors, either for delta-like or smooth branes. Furthermore, the mere existence of this zero-mode solution excludes any possible tachyonic mode of the theory. This is a powerful result, because, unlike other localization methods, the geometric coupling does not depend of the specific braneworld model, but only of the properties of spacetime itself. Among the main aspects, the localization condition ( $2\mathbf{c}_1 > \mathbf{d}$ ) can be obtained by imposing only that the background is an asymptotically AdS spacetime. In section (3.1.1) was made a detailed analysis of the cases (a)-(c) above mentioned, where we obtained: For case (a), the analytic solution is given by  $\chi_0^{(gauge)} = e^{\sigma(y)}$ , with  $\mathbf{a} = \mathbf{c}_1 = \mathbf{1}$  and the coupling constant given by  $\lambda_1 = \frac{D-6}{4(D-1)}$ . Therefore, from the localization condition, such zero-mode solution is confined when  $\mathbf{d} < 2$ , *i.e.*, in co-dimension one models. For case (b), we get the analytic solution  $\chi_0^{(gauge)} = e^{\frac{D}{2}\sigma(y)}$  and the coupling constant  $\lambda_2 = -2$ . This solution is localized for any number of transverse extra dimensions, since the localization condition becomes  $\mathbf{d} < D$  and can always be satisfied. Finally for case (c), the analytic solution is obtained when  $\mathbf{a} = \mathbf{c}_1^{(\pm)} = \frac{D-2}{2} \pm \sqrt{\lambda_1(D-1)(D-2)+1}$ , with  $\lambda_1$  a 'free' parameter. This solution is also localized on a **3**-brane for any number of extra dimensions, provided that the parameter  $\lambda_1$  belongs to the range  $(-(D-1)^{-1}(D-2)^{-1}, 0)$  for  $\mathbf{c}_1^{(-)}$ , or to the range  $(-(D-1)^{-1}(D-2)^{-1}, \infty)$  for  $\mathbf{c}_1^{(+)}$ . To conclude this discussion about the effective gauge field, we obtained some conditions for which the localization of gravity ensures the localization of gauge field sector in this scenarios. As discussed at the end of the section (3.1.1), the zero-mode solution obtained in Ref. [53] for the gravitational field is given by  $\psi_0^{(gravity)} = e^{\frac{(D-2)}{2}\sigma}$ . When we compare this solution to the above analysis for the gauge field, we can see that if the coupling of vector field with the Ricci tensor is present in the theory, then the localization of gravity is enough to ensure the localization of gauge field. This is an interesting result, since that all the consistency of the model will depend only on the fact that gravity is consistent.

We also considered the localization of the scalar components  $\mathcal{B}^j$  of the  $D$ -vector field  $\mathcal{A}_M$ . As said in the introduction, in co-dimension one models the scalar component is never localized simultaneously with the gauge field sector. This is a drawback since the backreaction of this field could alter the AdS vacuum. In section (3.2), we studied this problem and showed that when more co-dimensions are considered there is an indication that such scalar components can be localized simultaneously with the gauge field sector  $\mathcal{A}_\mu$ . Differently of this sector, a general analytical treatment of the  $\mathcal{B}^j$  was not found. This is due to the fact that the equation of motion (3.26) can not be diagonalized and therefore are always coupled. However, as we are interested in convergence conditions of the solution, an asymptotic treatment was performed for the cases (a)-(c) above. With this in mind the asymptotic solutions was found in Eq. (3.48). This solution indicates that the localization of the gauge and scalar components of the vector field can be simultaneously obtained only for the cases (b) and (c). Therefore only when the interaction with the Ricci tensor is switched on, as showed in fig. (1) and (2). We should stress that this is another important result of this work, since the localization of both components ensures that the backreaction of  $\mathcal{B}^j$  will not jeopardise the AdS feature of the vacuum. Moreover,

the localization of these components does not imply any modification on the Coulomb law, instead they could be interpreted as Higgs fields, or even dark energy. However, we could not ensure that these components are really confined because there is no guarantee that the solutions will be regular in all range of integration. In order to fully solve the scalar components problem, a specific background must be considered.

As an application, the intersecting brane model, developed by Arkani-Hamed *et al*, was used to discuss in a more detailed way these general results presented by us. Beyond that, it was possible to obtain the complete solutions for scalar components given by (4.16). As said before, the branes introduce a set of boundary conditions that fix all the parameters of the model. With this, we verified that the localization of the scalar and gauge components is not possible for the case (b) above. Therefore a fully consistent model is possible only for the case (c), when couplings with the *scalar* and the *Ricci tensor* are considered. Furthermore, in Eq. (4.20) we found the effective field theory for the scalar sector at the **3**-brane, which are free massless scalar fields. With this we get that the our final effective action has one free gauge field plus  $(d - 1)$  free scalar fields. The scalar fields can play an important role in cosmology and particle physics and in principle can provides phenomenological consequences of the geometrical localization mechanism. However, in the moment, this is beyond the scope of this paper and can be treated in a future work.

## Acknowledgement

We acknowledge the financial support provided by Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico (FUNCAP), the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and FUNCAP/CNPq/PRONEX.

## A Appendix

### A.1 Splitting of the Action (3.1) in the Sectors $S_{\perp}[\hat{\mathcal{A}}_{\mu}]$ and $S[\phi, B_k]$

In section (3.1), we proposed the separation  $\mathcal{A}_N = (\hat{\mathcal{A}}_{\mu} + \partial_{\mu}\phi, B_k)$ , where  $\partial_{\mu}\hat{\mathcal{A}}^{\mu} = 0$  is the transverse sector of the effective abelian vector field on the brane. This proposal allowed carry out the separation of action (3.1) in one part containing only the transverse sector and another part with longitudinal and scalar components  $B_k$  sectors. Here, we will only clarify this procedure.

From the action (3.1) we have the kinetic term in  $D$ -dimensions,

$$\frac{1}{4}\mathcal{F}_{MN}\mathcal{F}^{MN} = \frac{1}{4}\hat{\mathcal{F}}_{\mu\nu}\hat{\mathcal{F}}^{\mu\nu} + \frac{1}{4}\mathcal{B}_{jk}\mathcal{B}^{jk} + \frac{1}{2}\mathcal{F}_{\mu j}\mathcal{F}^{\mu j}, \quad (\text{A.1})$$

where  $(\mu, \nu, \dots)$  are related to the brane coordinates and  $(j, k, \dots)$  are related to extra dimensions. The first term in expression (A.1) is already written as a function only of the transverse sector  $\hat{\mathcal{A}}^{\mu}$ . The last term, which still has coupled terms of transverse sector with the other sectors, can be written as follows

$$\begin{aligned} \frac{1}{2}\mathcal{F}_{\mu j}\mathcal{F}^{\mu j} &= \frac{1}{2}g^{\mu\nu}g^{jk}\left(\partial_{\mu}\mathcal{B}_j - \partial_j\hat{\mathcal{A}}_{\mu} - \partial_j\partial_{\mu}\phi\right)\left(\partial_{\nu}\mathcal{B}_k - \partial_k\hat{\mathcal{A}}_{\nu} - \partial_k\partial_{\nu}\phi\right) \\ &= \frac{1}{2}g^{\mu\nu}g^{jk}\partial_{\mu}\mathcal{B}_j\partial_{\nu}\mathcal{B}_k - g^{\mu\nu}g^{jk}\partial_{\mu}\mathcal{B}_j\partial_k\partial_{\nu}\phi + \frac{1}{2}g^{\mu\nu}g^{jk}\partial_j\partial_{\mu}\phi\partial_k\partial_{\nu}\phi \\ &\quad + \frac{1}{2}g^{\mu\nu}g^{jk}\partial_j\hat{\mathcal{A}}_{\mu}\partial_k\hat{\mathcal{A}}_{\nu} + g^{\mu\nu}g^{jk}\partial_j\hat{\mathcal{A}}_{\mu}\partial_k\partial_{\nu}\phi - g^{\mu\nu}g^{jk}\partial_{\mu}\mathcal{B}_j\partial_k\hat{\mathcal{A}}_{\nu}. \end{aligned} \quad (\text{A.2})$$

The last two terms in this relation can be converted in boundary terms (in the coordinates of the brane) due to the condition  $\eta^{\mu\nu}\partial_{\nu}\hat{\mathcal{A}}_{\mu} = 0$ . Thus, if we consider that the boundary terms are zero, then the action (3.1) can be written as follows

$$S = S_{\perp}[\hat{\mathcal{A}}_{\mu}] + S[\phi, \mathcal{B}_k], \quad (\text{A.3})$$

where

$$\begin{aligned} S_{\perp} &= -\int d^{(D-d)}x d^d y \sqrt{-g} \left\{ \frac{1}{4}g^{\mu\nu}g^{\rho\lambda}\hat{\mathcal{F}}_{\mu\rho}\hat{\mathcal{F}}_{\nu\lambda} + \frac{1}{2}g^{\mu\nu}g^{jk}\partial_j\hat{\mathcal{A}}_{\mu}\partial_k\hat{\mathcal{A}}_{\nu} \right. \\ &\quad \left. + \frac{\lambda_1}{2}Rg^{\mu\nu}\hat{\mathcal{A}}_{\mu}\hat{\mathcal{A}}_{\nu} + \frac{\lambda_2}{2}g^{\mu\nu}g^{\rho\lambda}R_{\mu\rho}\hat{\mathcal{A}}_{\nu}\hat{\mathcal{A}}_{\lambda} \right\} \end{aligned} \quad (\text{A.4})$$

and

$$\begin{aligned} S[\phi, \mathcal{B}_k] &= -\int d^{(D-d)}x d^d y \sqrt{-g} \left\{ \frac{1}{4}g^{jm}g^{kl}\mathcal{B}_{jk}\mathcal{B}_{ml} + \frac{1}{2}g^{\mu\nu}g^{jk}\partial_{\mu}\mathcal{B}_j\partial_{\nu}\mathcal{B}_k \right. \\ &\quad - g^{\mu\nu}g^{jk}\partial_{\mu}\mathcal{B}_j\partial_k\partial_{\nu}\phi + \frac{1}{2}g^{\mu\nu}g^{jk}\partial_j\partial_{\mu}\phi\partial_k\partial_{\nu}\phi \\ &\quad \left. + \frac{\lambda_1}{2}Rg^{jk}\mathcal{B}_k\mathcal{B}_j + \frac{\lambda_2}{2}g^{lm}g^{jk}R_{jl}\mathcal{B}_m\mathcal{B}_k \right\}. \end{aligned} \quad (\text{A.5})$$

## References

- [1] G. Nordström, *On the possibility of unifying the electromagnetic and the gravitational fields*, Phys. Z. 15 (1914) 504–506.
- [2] T. Kaluza, *On the Problem of Unity in Physics*, Sitzungsber. Preuss. Akad. Wiss. Berlin. (Math. Phys.) 96 (1921) 966–972. O. Klein, *Quantentheorie und fünfdimensionale Relativitätstheorie*, Zeitschrift für Physik A. 37 (1926) 895–906. O. Klein, *The Atomicity of Electricity as a Quantum Theory Law*, Nature, 118 (1926) 51.
- [3] K. Akama, *Pregeometry*, Lect. Notes Phys. 176 (1982) 267–271; K. Akama, *An Early Proposal of “Brane World”*, [[hep-th/0001113](#)]
- [4] V. A. Rubakov and M. E. Shaposhnikov, *Do We Live Inside a Domain Wall?*, Phys. Lett. B. 125 (1983) 136.
- [5] L. Randall and R. Sundrum, *A Large mass hierarchy from a small extra dimension*, Phys. Rev. Lett. 83 (1999) 3370. [[hep-ph/9905221](#)]
- [6] L. Randall and R. Sundrum, *An Alternative to compactification*, Phys. Rev. Lett. 83 (1999) 4690. [[hep-th/9906064](#)]
- [7] S. Chang, J. Hisano, H. Nakano, N. Okada, M. Yamaguchi, *Bulk Standard Model in the Randall-Sundrum Background*, Phys. Rev. D 62 (2000) 084025. [[hep-ph/9912498](#)]
- [8] I. Oda, *Localization of Various Bulk Fields on a Brane*, (2000). [[hep-th/0009074](#)]
- [9] B. Bajc and G. Gabadadze, *Localization of matter and cosmological constant on a brane in anti-de Sitter space*, Phys. Lett. B. 474 (2000) 282. [[hep-th/9912232](#)]
- [10] M. Gremm, *Four-dimensional gravity on a thick domain wall*, Phys. Lett. B. 478 (2000) 434. [[hep-th/9912060](#)]
- [11] A. Kehagias and K. Tamvakis, *Localized gravitons, gauge bosons and chiral fermions in smooth spaces generated by a bounce*, Phys. Lett. B. 504 (2001) 38. [[hep-th/0010112](#)]
- [12] D. Bazeia, J. Menezes and R. Menezes, *New global defect structures*, Phys. Rev. Lett. 91 (2003) 241601. [[hep-th/0305234](#)]
- [13] D. Bazeia and A. R. Gomes, *Bloch brane*, JHEP. 0405 (2004) 012. [[hep-th/0403141](#)]
- [14] Y. X. Liu, Z. H. Zhao, S. W. Wei and Y. S. Duan, *Bulk Matters on Symmetric and Asymmetric de Sitter Thick Branes*, JCAP. 0902 (2009) 003. [[hep-th/0901.0782](#)]
- [15] H. Guo, A. Herrera-Aguilar, Y. X. Liu, D. Malagon-Morejon, and R. R. Mora-Luna, *Localization of bulk matter fields, the hierarchy problem and corrections to Coulomb’s law on a pure de Sitter thick braneworld*, Phys. Rev. D. 87 (2013) 095011. [[hep-th/1103.2430](#)]
- [16] M. Gogberashvili, and D. Singleton, *Brane in 6D with increasing gravitational trapping potential*, Phys. Rev. D. 69 (2003). [[hep-th/0305241](#)]
- [17] J. E. G. Silva, V. Santos and C. A. S. Almeida, *Gravity localization in a string-cigar braneworld*, Class. Quant. Grav. 30 (2013) 025005. [[hep-th/1208.2364](#)]
- [18] A. Flachi and M. Minamitsuji, *Field localization on a brane intersection in anti-de Sitter spacetime*, Phys. Rev. D. 79 (2009) 104021. [[hep-th/0903.0133](#)]
- [19] N. Arkani-Hamed, S. Dimopoulos, G. R. Dvali and N. Kaloper, *Infinitely large new dimensions*, Phys. Rev. Lett. 84 (2000) 586. [[hep-th/9907209](#)]

- [20] A. G. Cohen and D. B. Kaplan, *Solving the hierarchy problem with noncompact extra dimensions*, Phys. Lett. B. 470 (1999) 52-58. [[hep-th/9910132](#)]
- [21] J. E. Kim, B. Kyae and H. M. Lee, *Localized gravity and mass hierarchy in  $D = 6$  with a Gauss-Bonnet term*, Phys. Rev. D. 64 (2001).
- [22] O. Corradini, A. Iglesias, Z. Kakushadze and P. Langfelder, *Gravity on a 3-brane in 6D bulk*, Phys. Lett. B. 521 (2001) 96-104. [[hep-th/0108055](#)]
- [23] N. B. Cendejas and A. H. Aguilar, *Localization of 4D gravity on pure geometrical Thick branes*, Phys. Rev. D. 73 (2006). [[hep-th/0603184](#)]
- [24] Y. X. Liu, C.-E. Fu, H. Guo and H.-T. Li, *Deformed brane with finite extra dimension*, Phys. Rev. D. 73 (2006). [[hep-th/1102.4500](#)]
- [25] S. Randjbar-Daemi and M. Shaposhnikov, *Fermion zero-modes on brane-worlds*, Phys. Lett. B 492 (2000) 361-364. [[hep-th/0008079](#)]
- [26] R. Koley and S. Kar, *Scalar kinks and fermion localisation in warped spacetimes*, Class. Quant. Grav. 22 (2005) 753-768. [[hep-th/0407158](#)]
- [27] C. Ringeval, P. Peter and J. P. Uzan, *Localization of massive fermions on the brane*, Phys. Rev. D. 65 (2002) 044016. [[hep-th/0109194](#)]
- [28] R. R. Landim, G. Alencar, M. O. Tahim and R. N. Costa Filho, *New analytical solutions for bosonic field trapping in thick branes*, Phys. Lett. B. 731 (2014). [[hep-th/1310.2147v2](#)]
- [29] A. Melfo, N. Pantoja and J. D. Tempo, *Fermion localization on thick branes*, Phys. Rev. D. 73 (2006) 044033. [[hep-th/0601161](#)]
- [30] C.-E. Fu, Y. Zhong and Y.-X. Liu,  *$U(1)$  gauge vector field on a codimension-2 brane*, (2018). [[hep-th/1810.02081](#)]
- [31] G. Alencar, R. R. Landim, M. O. Tahim and R. N. Costa Filho, *Gauge field emergence from Kalb-Ramond localization*, Phys. Lett. B. 742 (2015) 256-260. [[hep-th/1409.5042](#)]
- [32] A. Pomarol, *Gauge boson in a five-dimensional theory with localized gravity*, Phys. Lett. B. 486 (2000) 153. [[hep-ph/9911294](#)]
- [33] H. Davoudiasl, J. L. Hewett and T. G. Rizzo, *Bulk gauge fields in the Randall-Sundrum model*, Phys. Lett. B. 473 (2000) 43. [[hep-ph/9911262](#)]
- [34] P. Midodashvili, *Brane in 6D and localization of matter fields*, (2003). [[hep-th/0308003](#)]
- [35] M. Giovannini, *Gauge field localization on Abelian vortices in six dimensions*, Phys. Rev. D. 66 (2002). [[hep-th/0205139](#)]
- [36] I. Oda, *A new mechanism for trapping of photon*, (2001). [[hep-th/0103052](#)]
- [37] I. Oda, *Localization of Matters on a String-like Defect*, (2000). [[hep-th/0006203](#)]
- [38] G. Dvali, G. Gabadadze and M. Shifman, *(Quasi)Localized Gauge Field on a Brane: Dissipating Cosmic Radiation to Extra Dimensions?*, Phys. Lett. B. 497 (2001) 271-280. [[hep-th/0010071](#)]
- [39] R. Guerrero, A. Melfo, N. Pantoja and R. Rodriguez, *Gauge field localization on brane worlds*, Phys. Rev. D. 81 (2010) 086004. [[hep-th/0912.0463](#)]
- [40] B. Batell and T. Gherghetta, *Yang-Mills localization in warped space*, Phys. Rev. D. 75 (2006). [[hep-th/0611305](#)]

- [41] C. A. Vaquera-Araujo and O. Corradini, *Localization of abelian gauge fields on thick branes*, O. Eur. Phys. J. C (2015) 75: 48. [[hep-th/1406.2892](#)]
- [42] W. T. Cruz, M. O. Tahim and C. A. S. Almeida, *Gauge field localization on a dilatonic deformed brane*, Phys. Lett. B. 686 (2010) 259-263.
- [43] W. T. Cruz, A. R. P. Lima and C. A. S. Almeida, *Gauge field localization on the Bloch Brane*, Phys. Rev. D. 87 (2012) 045018. [[hep-th/1211.7355](#)]
- [44] K. Ghoroku and A. Nakamura, *Massive vector trapping as a gauge boson on a brane*, Phys. Rev. D. 65 (2002) 084017. [[hep-th/0106145](#)]
- [45] G. Alencar, R. R. Landim, M. O. Tahim and R. N. Costa Filho, *Gauge Field Localization on the Brane Through Geometrical Coupling*, Phys. Lett. B. 739 (2014) 125. [[hep-th/1409.4396](#)]
- [46] Z. H. Zhao, Q. Y. Xie and Y. Zhong, *New localization method of  $U(1)$  gauge vector field on flat branes in (asymptotic)  $AdS_5$  spacetime*, Class. Quant. Grav. 32 (2015) 035020. [[hep-th/1406.3098](#)]
- [47] R. R. Landim, G. Alencar, M. O. Tahim and R. N. Costa Filho, *A Transfer Matrix Method for Resonances in Randall-Sundrum Models*, JHEP. 1108 (2011) 071. [[hep-th/1105.5573](#)]
- [48] I. C. Jardim, G. Alencar, R. R. Landim and R. N. Costa Filho, *Does Geometric Coupling Generates Resonances?*, EPL. 115 (2016) 51001. [[hep-th/1505.00689](#)]
- [49] I. C. Jardim, G. Alencar, R. R. Landim and R. N. Costa Filho, *Massive  $p$ -form trapping as a  $p$ -form on a brane*, JHEP. 1504 (2015) 003. [[hep-th/1410.6756](#)]
- [50] G. Alencar, C. R. Muniz, R. R. Landim, I. C. Jardim and R. N. Costa Filho, *Photon mass as a probe to extra dimensions*, Phys. Lett. B 759 (2016) 138-1406. [[hep-th/1511.03608](#)]
- [51] G. Alencar, I. C. Jardim, R. R. Landim, C. R. Muniz and R. N. Costa Filho, *Generalized nonminimal couplings in Randall-Sundrum scenarios*, Phys. Rev. D 93 (2016) no. 12, 124064. [[hep-th/1506.00622](#)]
- [52] G. Alencar, R. R. Landim, C. R. Muniz and R. N. Costa Filho, *Nonminimal couplings in Randall-Sundrum scenarios*, Phys. Rev. D 92 (2015) 066006. [[hep-th/1502.02998](#)]
- [53] C. Csaki, J. Erlich, T. J. Hollowood and Y. Shirman, *Universal aspects of gravity localized on thick branes*, Nucl. Phys. B 581 (2000) 309. [[hep-th/0001033](#)]
- [54] P. Dimopoulos, K., Farakos, A. Kehagias and G. Koutsoumbas, *Lattice Evidence for Gauge Field Localization on a Brane*, Nucl. Phys. B 617 (2001) 237-252. [[hep-th/0007079](#)]
- [55] M. Giovannini, *Vector field localization and negative tension branes*, Phys. Rev. D 65 (2002) 124019. [[hep-th/0204235](#)]
- [56] Y. X. Liu, Z. G. Xu, F. W. Chen and S. W. Wei, *New localization mechanism of fermions on braneworlds*, Phys. Rev. D 89 (2014) 086001. [[hep-th/1312.4145](#)]
- [57] W. D. Goldberger and M. B. Wise, *Bulk fields in the Randall-Sundrum compactification scenario*, Phys. Rev. D. 60 (1999) 107505. [[hep-ph/9907218](#)]
- [58] M. J. Duff and J. T. Liu, *Hodge duality on the brane*, Phys. Lett. B 508 (2001) 381. [[hep-th/0010171](#)]