Finite- and infinite-volume thermodynamics around the zero of the pressure in deconfining SU(2) Quantum Yang-Mills theory

Ralf Hofmann

Institut für Theoretische Physik, Universität Heidelberg, Philosophenweg 16, D-69120 Heidelberg, Germany; r.hofmann@thphys.uni-heidelberg.de

Abstract

We re-address the self-intersection region in a figure-eight shaped center-vortex loop containing a frequently perturbed *BPS monopole* subject to a core-oscillation frequency ω_0 , rectifying a numerical error in estimating the region's radius r_0 in comparison to the spatial coarse-graining scale of infinite-volume thermodynamics. Implications are discussed. We also compute the lowest frequency Ω_0 of a spherically symmetric plasma oscillation within a *neutral* and spatially homogeneous ball-like region of deconfining phase in dependence of its radius R_0 . For $r_0 = R_0$ we compare ω_0 with Ω_0 . We point out how the idealisations, which are assumed in this work, will have to be relaxed in order to address the emission of electromagnetic radiation and of non-intersecting as well as self-intersecting center-vortex loops away from the surface region of macroscopically sized plasma balls.

1 Introduction

The phase structure and the deconfining thermal ground state of (electric-magnetic dually interpreted) SU(2) Yang-Mills thermodynamics suggests the existence of a finite-extent particle with a magnetic moment provided by the electric flux along a figure-eight-like configuration of a self-intersecting center-vortex loop, immersed into the confining phase. This soliton is stable, possesses a mass m_0 mainly arising from the deconfining energy density of the ball-like, Bohr-radius sized self-intersection region whose radius r_0 is about fifty times larger than the Compton wave length l_C . This object exhibits one unit of electric charge, immersed into this deconfining region as a BPS monopole of mass m_m whose core localises to within a radius matching l_C , which frequently and indeterministically is relocated and forced to oscillate at frequency ω_0 [1, 2] by the action of a trivial-holonomy caloron/anticaloron center [3] in the deconfining thermal ground state [4]. Such a model of the electron (and other charged leptons) concretely represents Louis de Broglie's ideas on the (quantum) thermodynamics of the isolated particle [5, 6], based on the important observation that its mass m_0 must be related to a time-periodic phenomenon of (circular) frequency ω_0 as $m_0 c^2 = \hbar \omega_0 - c$ the speed of light in vacuum, \hbar Planck's reduced unit of action – in the rest frame of this particle. Note that a Lorentz boost generates out of a spatially flat oscillation a propagating wave, whose (de Broglie) wavelength λ relates to the spatial momentum p of the particle as $\lambda = \frac{2\pi\hbar}{p}$, and the increase of rest energy m_0c^2 by virtue of the boost can be decomposed into a reduction of internal heat plus the mechanical work invested. The reason why this model of the electron complies with the particle's apparent structurelessness of charge distribution, as inferred from scattering experiments with energy-momentum transfers up to several TeV, is the thermal nature of the self-intersection region (maximum entropy), the locus of electric charge essentially being undetermined within the large volume $\sim \frac{4}{3}\pi r_0^3$. On the scale of the electron mass m_0 the center-vortex loop without a region of self-intersection is massless and prone to curve shrinking [7]. Emergent in the confining phase of the same SU(2) Yang-Mills theory, such a soliton is a natural candidate for the (electron) neutrino.

There are implications of this model for high-temperature plasma physics (in considering transport properties in view of the turbulent behaviour induced by unstable centervortex loops with higher self-intersection numbers [8, 9]) and for the physics of strongly correlated electrons in a spatial plane (parts of the correlations intrinsically arising from the non-local magnetic interactions of vortex sectors [7] and the thermal, indeterministic nature of the electric charge density [4]) underlying, e.g., the phenomenon of high- T_c superconductivity. Both fields of research require long-term efforts, however.

One concern of the present paper is to discuss a few amendments to [4] where the above-sketched model of the electron was discussed. In particular, we re-state results on r_0 and m_m and the ratio m_m/m_0 under the use of the proper (and not erroneous as in [4]) monopole mass formula $m_m = \frac{4\pi}{e(T_0)}H_{\infty}(T_0)$ with $e(T_0) = 12.96$ the value of the effective gauge coupling at the temperature $T_0 = 1.32 T_c$ (T_c the critical temperature for the deconfining-preconfining phase transition), where the deconfining pressure P vanishes. Here $H_{\infty}(T_0) = \pi T_0$ denotes the asymptotic modulus of the adjoint Higgs field in the BPS monopole configuration, linking it to (anti)caloron dissociation at maximum non-trivial holonomy [10, 11, 12, 13, 14] (and therefore electron-positron pair creation).

There is no essential change in numerical values up to this point. However, in comput-

ing the ratio $\frac{r_0}{|\phi|^{-1}(T_0)}$ a substantial error was committed in [4]: Instead of $\frac{r_0}{|\phi|^{-1}(T_0)} \sim 160$ the actual value turns out to be $\frac{r_0}{|\phi|^{-1}(T_0)} \sim 0.104$ which corresponds to $\frac{r_0}{T_0^{-1}} \sim 1.29$. (The radius r_0 is denoted as R_0 in [4].) Therefore, it is incorrect to state, as was done in [4], that the self-intersection region of radius r_0 represents infinite-volume thermodynamics. Rather, this region is deeply contained within the center of the accomodating caloron or anticaloron, and, judged by the approximate attainment of the asymptotic harmonic dependence of ϕ on Euclidean time when integrating the according field-strength correlator on a caloron or anticaloron in singular gauge over the instanton-scale parameter [14], the use of infinite-volume thermodynamics can thus be considered a coarse resemblence only.

The other objective of our present work is a discussion of an extended, neutral plasma ball at T_0 , immersed into the confining phase, where infinite-volume thermodynamics is well saturated in the bulk (disregarding the non-thermal situation within a thin shell of preconfining/turbulent confining phase leading to surface-tension effects) for a sufficiently large ball radius R_0 . In particular, we ask what the lowest frequency Ω_0 is, associated with a spherically symmetric breathing mode describing a homogeneous plasma oscillation about T_0 . This requires the determination of the longitudinal sound speed c_s at T_0 which turns out to be $c_s = 0.479 c$. Note that, in spite of the vanishing pressure at T_0 , exhibiting a cancellation of the contributions from the thermal ground state and its (partially massive) excitations, this is not much below the ultrarelativistic-gas limit $c_s = \frac{1}{\sqrt{3}} c \sim 0.577 c$. Subsequently, we then compare Ω_0 and $\omega_0 = m_0 c^2/\hbar$ at $R_0 = r_0$.

This paper is organised as follows. In Sec. 2 we re-address the computation of r_0 using the proper expression for the monopole mass m_m . The value of r_0 is not affected much compared to the one obtained in [4]. We also compute the ratio of m_m to m_0 . However, due to a calculational error the ratio of r_0 to the coarse-graining scale $|\phi|^{-1}$ turns to out be much smaller than what was found in [4] and what was taken as a proof that infinitevolume thermodynamics is valid at face value. We argue, however, that the ratio r_0/T_0^{-1} permits an approximately thermodynamical treatment because the asymptotic harmonic time dependence of the integral over the field-strength correlator, rendering the Euclidean time dependence of field ϕ a mere gauge choice [14], is reasonably well approached when cutting off the instanton-scale-parameter integral at r_0 . In Sec. 3 we address a situation where the deconfining, homogeneous plasma, considered at temperatures around T_0 , is void of an explicit monopole but undergoes spherically symmetric, oscillatory breathing by virtue of a finite longitudinal sound velocity c_s . We reliably compute the square of c_s by appealing to the derivatives w.r.t. temperature of the pressure and the energy density, employing the evolution equation for the mass of the off-Cartan modes. Both situations, monopole driven oscillation and spherically symmetric breathing, are compared in terms of their frequencies, the former exhibiting a much more rapid oscillation than the latter. Finally, in Sec. 4 we summarise and discuss our results and provide an outlook on future work.

2 Self-intersection region of a figure-eight shaped centervortex loop

In [4] we have proposed a model of the free electron, based on the phase structure of SU(2) Yang-Mills thermodynamics and the work in [1, 2], investigating the response of a

BPS monopole to a spherically symmetric initial perturbation and the spectrum of normal modes. Here we would like to correct some numerical statements arising from an incorrect monopole mass formula (Eq. (18) of [4]). Also, we point out an error in Eq. (21) of [4], entailing a conceptual re-interpretation of the physics associated with the self-intersection region. From now on we work in super-natural units: $c = \hbar = k_B = 1$ where k_B denotes Boltzmann's constant.

The mass m_m of a BPS monopole is given as [15]

$$m_m = \frac{4\pi}{e} H_\infty \,, \tag{1}$$

where e denotes the defining gauge coupling of the adjoint Higgs model and H_{∞} the spatially asymptotic modulus of the Higgs field. Assuming that the monopole was liberated by the dissociation of a maximum-holonomy caloron at $T_0 = 1.32 T_c$, the temperature where the thermodynamical (one-loop) pressure vanishes, we have [14, 13]

$$e(T_0) = 12.96, \qquad H_{\infty}(T_0) = \pi T_0.$$
 (2)

The electron's rest mass m_0 is, on one hand, given by the circular frequency ω_0 of monopole-core oscillation, found to be equal to the mass of the two off-Cartan modes in [1, 2]

$$\omega_0 = eH_\infty \,. \tag{3}$$

On the other hand, m_0 decomposes into m_m and the energy contained within an approximately ball-like region of radius r_0 (the contribution of the two vortex loops in negligible [14], r_0 is denoted as R_0 in [4]) due to the deconfining plasma of energy density $\rho(T_0)$

$$m_{0} = 12.96 H_{\infty}(T_{0}) = m_{m} + \frac{4\pi}{3} r_{0}^{3} \rho(T_{0})$$

$$= H_{\infty}(T_{0}) \left(\frac{4\pi}{12.96} + 8.31 \times \frac{128\pi}{3} \left(\frac{r_{0}}{18.31} \right)^{3} H_{\infty}^{3}(T_{0}) \right).$$
(4)

For later use, we introduce the dimensionless temperature $\lambda \equiv \frac{2\pi T}{\Lambda}$ where Λ denotes the Yang-Mills scale, related to m_0 as [4]

$$\Lambda = \frac{1}{118.6} \, m_0 \,. \tag{5}$$

Solving Eq. (4) for r_0 yields

$$r_0 = 4.043 \, H_\infty^{-1} \tag{6}$$

instead of $r_0 = 4.10 H_{\infty}^{-1}$ as obtained in [4]. It is instructive to compute the relative contribution of m_m to m_0 and the ratio of Compton wave length $l_C = m_0^{-1} = \frac{1}{12.96 H_{\infty}(T_0)}$ [1, 2] to r_0

$$\frac{m_m}{m_0} = \frac{4\pi}{(12.96)^2} = 0.0748, \qquad \frac{l_C}{r_0} = \frac{1}{52.40}.$$
(7)

Moreover, radius r_0 compares to the spatial coarse-graining scale $|\phi|^{-1}(T_0)$, turning the Euclidean time dependence of the field strength correlation within a caloron or anticaloron center into a mere choice of gauge [14], as follows

$$\frac{r_0}{|\phi|^{-1}(T_0)} = \frac{4.043}{\sqrt{2\left(\frac{118.6}{12.96}\right)^3}} = 0.1033.$$
(8)




Figure 1: Saturation towards a harmonic Euclidean time dependence of the contribution of a Harrington-Shepard caloron to the field-strength correlation defining the phase of the field ϕ as a function of the scaled cutoff $\xi \equiv \frac{\rho}{\beta}$ for the instanton-scale-parameter integration. Cutting off at $\rho \sim r_0 = 1.29\beta_0$ suggests that there are (mild) deviations from a harmonic dependence. Figure taken from [16].

Due to a simple calculational error in Eq. (22) of [4] this result is qualitatively different and implies that the "thermodynamics" we discussed so far actually occurs deep within the center of a caloron or anticaloron of a scale parameter $\rho \sim |\phi|^{-1}(T_0)$ (the coarse-graining scale of infinite-volume thermodynamics in the deconfining phase [14]). Since

$$\frac{r_0}{\beta_0} = 1.29 \qquad (\beta_0 \equiv \frac{1}{T_0})$$
 (9)

Fig. 1 suggests that the scale parameter integral, defining the phase of ϕ [14], does not quite saturate a harmonic dependence on Euclidean time when cut off at $\rho \sim r_0$. Therefore, our above discussion, which assumes that such a saturation occurs within the volume $\frac{4\pi}{3}r_0^3$, is only an approximate account of the quantum physics within the self-intersection region: the monopole is always close to the locus of action at the inmost point of the caloron or anticaloron, rendering this region a highly jittery object, not only concerning the spherical shell of preconfining/confining phase that represents the boundary of the ball but also its deep bulk. Still, the thermodynamical approximation employed here should yield reasonable estimates of all spatial scales and of the critical temperature $T_c =$ $13.87/(2\pi \times 118.6) m_0 = 9.49 \text{ keV}$ [4].

3 Lowest spherically symmetric breathing mode

Let us now address a situation, where thermodynamics can be applied without restrictions. Namely, we would like to compute the frequency Ω_0 of the lowest spherically symmetric breathing mode of a deconfining SU(2) Yang-Mills plasma ball of homogeneous energy density ρ and pressure P, whose temperature oscillates about T_0 . Surface effects, arising from the transition between the deconfining (bulk) and the confining (exterior of ball) phases can be neglected for a sufficiently large ball mass $M \equiv \frac{4}{3}\pi R_0^3 \rho(R_0)$, and the according expression for Ω_0 reads [18, 17]

$$\Omega_0 = \frac{\pi c_s \Lambda}{\bar{R}_0} \,, \tag{10}$$

where the dimensionless quantities c_s (longitudinal sound velocity) and \bar{R}_0 (R_0 in units of the inverse Yang-Mills scale Λ^{-1}) are defined as

$$c_s^2 \equiv \left. \frac{\frac{d\bar{P}}{d\lambda}}{\frac{d\bar{\rho}}{d\lambda}} \right|_{\lambda = \lambda_0} \tag{11}$$

and $\bar{R}_0 \equiv R_0 \Lambda$. In Eq. (11) the use of the one-loop pressure $P \equiv \Lambda^4 \bar{P}$ and of the one-loop energy density $\rho \equiv \Lambda^4 \bar{\rho}$ are excellent approximations (modified by higher-loop corrections on the 1%-level [14]). Amusingly, an estimate of Ω_0 by virtue of a linearisation of the force-balance equation $\frac{4}{3}\pi R^3 \rho(R)\ddot{R} = 4\pi R^2 P(R)$ about R_0 and in employing energy conservation,

$$\bar{R} \equiv R\Lambda = \left(\frac{3\bar{M}}{4\pi\bar{\rho}(\bar{R})}\right)^{1/3} \qquad (\bar{M} \equiv \Lambda M), \tag{12}$$

replaces the factor of π in Eq. (10) by a factor of three. For \bar{P} and $\bar{\rho}$ we have [14]

$$\bar{P}(2a,\lambda) \equiv -\frac{2\lambda^4}{(2\pi)^6} \left[2\tilde{P}(0) + 6\tilde{P}(2a) \right] - 2\lambda,$$

$$\bar{\rho}(2a,\lambda) \equiv \frac{2\lambda^4}{(2\pi)^6} \left[2\tilde{\rho}(0) + 6\tilde{\rho}(2a) \right] + 2\lambda,$$

$$a = a(\lambda) \equiv 2\pi e(\lambda) \lambda^{-3/2} \qquad (e(\lambda_0) = 12.96),$$
(13)

where

$$\tilde{P}(y) \equiv \int_0^\infty dx \, x^2 \log\left[1 - \exp\left(-\sqrt{x^2 + y^2}\right)\right],$$

$$\tilde{\rho}(y) \equiv \int_0^\infty dx \, x^2 \frac{\sqrt{x^2 + y^2}}{\exp\left(\sqrt{x^2 + y^2}\right) - 1}.$$
(14)

Taking into account implicit (via $a(\lambda)$) and explicit dependences of \bar{P} and $\bar{\rho}$ on λ and employing the evolution equation [14]

$$1 = -\frac{24\lambda^3}{(2\pi)^6} \left(\lambda \frac{da}{d\lambda} + a\right) a D(2a) , \qquad (15)$$

one derives

$$\frac{d\bar{P}}{d\lambda} = -\frac{\lambda^3}{(2\pi)^6} \left(16\tilde{P}(0) + 48(\tilde{P}(2a) - a^2D(2a)) \right),
\frac{d\bar{\rho}}{d\lambda} = \frac{\lambda^3}{(2\pi)^6} \left(16\tilde{\rho}(0) + 48\left(\tilde{\rho}(2a) - a^2(D(2a) - F(2a))\right) \right) + 2\left(1 - \frac{D(2a) - F(2a)}{D(2a)} \right),$$
(16)

where

$$D(y) \equiv \int_{0}^{\infty} dx \, \frac{x^{2}}{\sqrt{x^{2} + y^{2}}} \frac{1}{\exp\left(\sqrt{x^{2} + y^{2}}\right) - 1},$$

$$F(y) \equiv \int_{0}^{\infty} dx \, x^{2} \frac{\exp\left(\sqrt{x^{2} + y^{2}}\right)}{\left(\exp\left(\sqrt{x^{2} + y^{2}}\right) - 1\right)^{2}}.$$
(17)

Substituting Eqs.(16) into Eq. (11) at $\lambda_0 = 18.31$, we numerically obtain

$$c_s(\lambda_0) = 0.479$$
. (18)

For Eq. (10) this yields

$$\Omega_0 = 1.506 \,\frac{\Lambda}{\bar{R}_0} \,. \tag{19}$$

Let us now compare the monopole-core induced frequency ω_0 of the self-intersection region of the figure-eight shaped center-vortex loop (model of the electron) with Ω_0 at one and the same radius

$$r_0 = R_0 = 4.043 \, H_\infty^{-1}(T_0) \,, \tag{20}$$

see Eq. (6). For Ω_0 this yields

$$\Omega_0 = 0.372 \, H_\infty(T_0) \tag{21}$$

such that $\frac{\omega_0}{\Omega_0} = \frac{12.96}{0.372} = 34.84$. This large ratio is suggestive since the oscillation in the self-intersection region – quantum initiated by caloron or anticaloron action – is induced by the classical dynamics of a monopole core [1, 2] whose size matches the Compton wave length l_C while the lowest symmetric breathing mode of the neutral deconfining ball is a consequence of sound propagation in bulk thermodynamics, spatially supported by a much larger system of Bohr-radius dimensions.

Eq. (19) is the more reliable the larger R_0 is. Isotropy breaking effects, which associate with the neglected surface dynamics of the ball and/or the excitation of spherically nonsymmetric oscillation states, cause this surface to (electromagnetically) radiate with a spectrum that is dominated by frequencies around $\nu_0 \sim \frac{\Omega_0}{2\pi}$, corresponding to a wave length $l_0 = \frac{1}{\nu_0} \sim \frac{2\pi R_0}{1.506}$.

4 Summary and Discussion

This paper's purpose was to compare two situations in which a ball-like region of deconfining phase in SU(2) Yang-Mills thermodynamics oscillates about the zero of the pressure at temperature T_0 : the self-intersection region of a figure-eight shaped, solitonic centervortex loop (a model of the electron) containing a frequently perturbed BPS monopole, whose classical core dynamics drives this oscillation of (circular) frequency ω_0 (up to a factor \hbar coincident with the rest energy m_0c^2 of the soliton [5, 6]). Furthermore, we have considered a homogeneous, electrically neutral region whose lowest spherically symmetric oscillatory excitation of (circular) frequency Ω_0 is supported by a finite speed of sound c_s , in turn based on the thermal ground state's quantum excitations which are microscopically mediated by the unit of action \hbar localised within the center of a caloron/anticaloron [14]. At the same radius, $r_0 = R_0 = \frac{4.043}{\pi T_0}$, we obtain $\frac{\omega_0}{\Omega_0} = 34.84$ which makes explicit the very different cause of oscillation in either case.

We have noticed a numerical error in [4] concerning an estimate of the system size r_0 in terms of the spatial coarse-graining scale $|\phi|^{-1}$. The correct result states that finite-size effects cannot be excluded at face value since the region of self-intersection actually is contained deeply within the ball of spatial coarse-graining corresponding to the infinitevolume situation. However, the asymptotic harmonic time dependence of the integrated field strength correlation [14], required for the introduction of the field ϕ , is approached to some extent when cutting the instanton-scale-parameter integration off at r_0 such that, approximately, one can still rely on infinite-volume thermodynamics.

The present work only represents a first step in studying the plasma dynamics of a ball-like region of deconfining phase at T_0 . More realistically, the physics of a certain boundary shell should be taken into account. The preconfining-phase part of this shell is superconducting (condensate of electrically charged, massless monopoles) [14] with implications for the stabilisation of the highly turbulent gas of self-intersecting vortex loops (electrically charged particles) within a region externally adjacent to this shell. Also, we did not address the evaporation physics (emission of non-intersecting and self-intersecting center-vortex loops from the surface of this shell) in case of macroscopically sized balls, see [19], and how this process affects the oscillation dynamics and electromagnetic emission. Our results on the spherically symmetric oscillations of the homogeneous and macroscopic plasma could be relevant for models of Cepheid variables beyond the usual kappa mechanism of star dynamics and the description of certain, quasi-stabilised, compact and radiating objects created within atmospheric discharges. It is not unlikely to provide for a conceptual impact on plasma stabilisation in terrestial fusion experiments.

Acknowledgments: We would like to acknowledge a useful conversation with Anton Plech.

References

- Fodor, G. and Racz, I. What Does a Strongly Excited 't Hooft-Polyakov Magnetic Monopole Do? *Phys. Rev. Lett.* 2004, *92*, 151801.
- [2] Forgacs, P. and Volkov, M. S. Resonant Excitations of the 't Hooft-Polyakov Monopole. Phys. Rev. Lett. 2004, 92, 151802.
- [3] Harrington, B. J. and Shepard, H. K., Periodic Euclidean solutions and the finitetemperature Yang-Mills gas. *Phys. Rev. D* 1978, 17, 2122.
- [4] Hofmann, R. The isolated electron: De Broglie's "hidden" thermodynamics, SU(2) Quantum Yang-Mills theory, and a strongly perturbed BPS monopole. *Entropy* 2017, 19, 575.
- [5] de Broglie, L. Recherches sur la theorie des quanta. Ann. de Phys. 1925, 10th series,1.
- [6] de Broglie, L. The thermodynamics of the isolated particle. *Gauthier-Villars Editor* 1964, Paris, 1.
- [7] Moosmann, J. and Hofmann, R. Evolving center-vortex loops. ISRN Math. Phys. 2012, 2012, 236783.
- [8] Hofmann, R. Yang-Mills thermodynamics at low temperature. Mod. Phys. Lett. A 2007, 22, 657.
- [9] Bender, C. M. and Wu, T. T. Anharmonic oscillator. Phys. Rev 1969, 184, 1231.

- [10] Nahm, W. A simple formalism for the BPS monopole. *Phys. Lett. B* 1980, 90, 413. Nahm, W. All self-dual multimonopoles for arbitrary gauge groups. *CERN preprint TH-3172* 1981.
 Nahm, W. The Construction of all Self-dual Multimonopoles by the ADHM Method. In *Monopoles in Quantum Field Theory*, ed. N. Craigie *et al.* World Scientific, Singapore, 1982, p. 87.
 Nahm, W. Self-dual monopoles and calorons. In *Trieste Group Theor. Method*, 1983, p. 189.
- [11] Kraan, T. C. and Van Baal, P. Exact T-duality between calorons and Taub-NUT spaces. *Phys. Lett. B* 1998, 428, 268.
 Kraan, T. C. and Van Baal, P. Periodic instantons with non-trivial holonomy. *Nucl. Phys. B* 1998, 533, 627.
- [12] Lee, K. and Lu, C. SU(2) calorons and magnetic monopoles. Phys. Rev. D 1998, 58, 025011.
- [13] Diakonov, D. et al. Quantum weights of dyons and of instantons. Phys. Rev. D 2004, 70, 036003.
- [14] Hofmann, R. The thermodynamics of quantum Yang–Mills theory: theory and application. World Scientific, Singapore, 2012 (1st ed.) and 2016 (2nd ed.).
- [15] Prasad, M. K. and Sommerfield, C. M. Exact Classical Solution for the 't Hooft Monopole and the Julia-Zee Dyon. *Phys. Rev. Lett.*, **1975**, *35*, 760.
- [16] Herbst, U. and Hofmann, R. Emergent Inert Adjoint Scalar Field in SU(2) Yang-Mills Thermodynamics due to Coarse-Grained Topological Fluctuations. *ISRN High Energy Physics* 2012, Article ID 373121.
- [17] Hartland, G. V., Hu, M., and Sader, J. E. Softening of the symmetry breaking mode in gold particles by laser-induced heating. J. Phys. Chem. B 2003, 107, 7472.
- [18] Lamb, H. On the vibrations of an elastic sphere. Proc. London Math. Soc. 1882, 13, 189.
- [19] Giacosa, F. and Hofmann, R. Bursts of low-energy electron-positron pairs in TeVrange collider physics. *Mod. Phys. Lett. A* 2009, 24, 1937.