

Maxwell's equal-area law with several pairs of conjugate variables for RN-AdS black holes

Xiong-Ying Guo, Huai-Fan Li*, Ren Zhao

*Department of Physics,
Shanxi Datong University, Datong 037009, China
Institute of Theoretical Physics,
Shanxi Datong University, Datong 037009, China*

Abstract

In this paper, using Maxwell's equal-area law we study the phase transition of charged AdS black holes by choosing different independent conjugate variables. As is well known, the phase transition can be characterized by the state function of the system, the determination of the phase transition point has nothing to do with the choice of independent conjugate variables. To studying the thermodynamic properties of AdS black holes we give the conditions under which the independent conjugate variables are chosen. When the charge of the black hole is invariable, according to the conditions we find that the phase transition is related to the electric potential and the horizon radius of the charged black hole. Keeping the cosmological constant as a fixed parameter, the phase transition of a charged AdS black hole is related to the ratio of the event horizon to cosmological constant of the black hole. This conclusion is of great importance for us to study the critical phenomenon of black holes and to improve the self-consistent geometry theory of black hole thermodynamic.

February 12, 2022

*Email: huaifan.li@stu.xjtu.edu.cn; huaifan999@sxdt dx.edu.cn(H.-F. Li)

Contents

1	Introduction	2
2	Thermodynamics of charged AdS black hole	3
3	The equal-area law of charged AdS black hole in extended phase space	4
3.1	The construction of equal-area law in $P - V$ diagram	5
3.2	The construction of equal-area law in $T - S$ diagram	6
3.3	The construction of equal-area law in $P - v$ diagram	7
3.4	The construction of equal-area law in $Q^2 - \Psi$ and $Q - \Phi$ diagram . . .	9
4	Discussions and conclusions	11

1 Introduction

The study of the thermodynamic properties of black holes has always been one of the topics of interest to theoretical physicists. In recent years, the study of black hole thermodynamics in AdS and dS space has attracted wide attentions [1–29]. Moreover, the cosmological constant Λ is used as the thermodynamic pressure for the AdS black hole in the extended phase space, and its conjugate variable has dimension of volume. Calculating the critical components and finding the phase transition of AdS black hole, this phase transition is analogous to the Van der Waals liquid-gas phase transition [1–4, 28–40]. Some researches are done by using the conjugate variables (P, v) [16, 18, 31], some researches are done by the conjugate variation (T, S) [1, 16, 20, 39]. In the previous works, we had were studied the critical behaviors of black holes according to (Q, Φ) , through the study we find that the second-order phase transition is independent of the choose of conjugate variable, while the critical exponents are dependent on the choice of conjugate variables [41].

For a normal canonical thermodynamic system, the phase transition point is the state function of the system. The determination of the phase transition point is independent of the choice of independent conjugate variables. Using the Maxwell’s equal area law we discuss that the phase transition points of AdS black hole should be the same with different independent conjugate variables [42–48]. When the phase transition points obtained by the independent conjugate variable are different from those

obtained by other independent conjugate variable, we believe that this set of independent conjugate variable cannot be regarded as the conjugate variable of AdS black hole. The conditions for selecting independent dual state parameters for studying the thermodynamic properties of black holes are given. It is not only helpful for further understanding the properties of entropy, temperature and heat capacity of black holes, but also is of great importance to the improvement of self-consistent geometry theory of black hole thermodynamic.

This paper is organized as follows. In the next section we review the thermodynamics of charged AdS black hole. In section 3 we study the critical behavior of charged AdS black hole with Maxwell's equal area law according to different conjugate variables (P, V) , (T, S) and (P, v) , respectively. We also study the critical behavior by the conjugate variables (Q^2, Ψ) and (Q, Φ) to find the condition that the phase transition occurs. In section 4, we give conclusion and discussion for the choice of the conjugate variable in the study of thermodynamics of black hole.

2 Thermodynamics of charged AdS black hole

The action of Einstein-Maxwell theory in the background of AdS spacetime is [1]

$$I = \frac{1}{16\pi} \int d^4x \sqrt{-g} (R - 2\Lambda - F_{\mu\nu} F^{\mu\nu}). \quad (2.1)$$

To start with we review some basic thermodynamic properties of the spherical charged AdS black hole. In Schwarzschild-like coordinates the metric reads

$$ds^2 = -f(r)dt^2 + f^{-1}dr^2 + r^2 d\Omega_2^2, \quad (2.2)$$

Here, $d\Omega_2^2$ stands for the standard element on S^2 and the function f is given by

$$f(r) = 1 - \frac{2M}{r} + \frac{Q^2}{r^2} + \frac{r^2}{l^2}, \quad (2.3)$$

The position of the black hole event horizon is determined as a larger root of $f(r_+) = 0$. The parameter M represents the ADM mass of the black hole and in our set up it is associated with the enthalpy of the system. Q represents the electric charge. Using the Euclidean trick, one can identify the black hole temperature

$$T = \frac{1}{4\pi r_+} \left(1 + \frac{3r_+^2}{l^2} - \frac{Q^2}{r_+^2} \right) = \frac{1}{4\pi r_+} \left(1 + 8\pi P r_+^2 - \frac{Q^2}{r_+^2} \right), \quad (2.4)$$

and the corresponding entropy

$$S = \frac{A}{4}, \quad A = 4\pi r_+^2. \quad (2.5)$$

Thus, the ADM mass of black hole is enthalpy $H = M$ [49] and is obtained as [19]

$$M(S, Q, P) = \frac{1}{6\sqrt{\pi S}}(3S + 8PS^2 + 3\pi Q^2). \quad (2.6)$$

The intensive parameters conjugate to S , Q and P are defined by $T = \left(\frac{\partial M}{\partial S}\right)_{P,Q}$, $\Phi = \left(\frac{\partial M}{\partial Q}\right)_{S,P}$, $V = \left(\frac{\partial M}{\partial P}\right)_{S,Q}$, where T is the temperature, $\Phi = \frac{Q}{r_+}$ is the electric potential and the thermodynamics volume V is obtained via $V = \frac{4\pi}{3}r_+^3$. The new variable $\Psi = \left(\frac{\partial M}{\partial Q^2}\right)_{S,P}$, which conjugate to Q^2 , is the inverse of the specific volume $v = 2r_+$ in the natural unit where $l_p = 1$ [1]. Therefore, the above thermodynamic relation satisfies the first law as:

$$dM = TdS + \Psi dQ^2 + VdP, \quad dM = TdS + \Phi dQ + VdP. \quad (2.7)$$

Comparing Eq.(2.7) with the first law equations of the general thermodynamics system, we find that $\Psi(\Phi)$ is conjugated to the $Q^2(Q)$, respectively. From scaling argument, we arrive at the Smarr formula as :

$$M = 2TS + 2\Psi Q^2 - VP, \quad M = 2TS + \Phi Q - VP. \quad (2.8)$$

In the previous works, the critical phenomena of charged AdS black hole with different choices of the conjugated variables have been studied. [1, 19–22]. Taking Q as a constant, we can obtain the value of critical point as

$$T_c = \frac{\sqrt{6}}{18\pi Q}, \quad r_c = \sqrt{6}Q, \quad P_c = \frac{1}{96\pi Q^2}. \quad (2.9)$$

While taking the l as constant, the value of critical point as

$$T_c = \frac{2}{\pi l \sqrt{6}}, \quad r_c = \frac{l}{\sqrt{6}}, \quad Q_c = \frac{l}{6}. \quad (2.10)$$

3 The equal-area law of charged AdS black hole in extended phase space

When the charge of the AdS black hole is invariant, the state equation of charged AdS black hole is given by Eq.(2.4). The state equation can be write as $f(T, P, V) = 0$, which means the particle number of system is invariant. First, we investigate the condition that the phase transition occur with the different conjugate variables (P, V) and (T, S) , respectively, by Maxwell's equal area law.

3.1 The construction of equal-area law in $P - V$ diagram

When the charge Q of AdS black hole is invariant, the horizontal axis of two-phase coexistence region in charged AdS black hole are S_2 and S_1 , respectively, the pressure on the vertical axis as P_0 , which depends on the horizon radius of black hole r_+ . From the Maxwell's equal area law,

$$P_0(V_2 - V_1) = \int_{V_1}^{V_2} P dV, \quad (3.1)$$

we can obtain

$$P_0 = \frac{T_0}{2r_1} - \frac{1}{8\pi r_1^2} + \frac{Q^2}{8\pi r_1^4}, \quad P_0 = \frac{T_0}{2r_2} - \frac{1}{8\pi r_2^2} + \frac{Q^2}{8\pi r_2^4}, \quad (3.2)$$

$$2P_0 = \frac{3T_0(1+x)}{2r_2(1+x+x^2)} - \frac{3}{4\pi r_2^2(1+x+x^2)} + \frac{3Q^2}{4\pi r_2^4 x(1+x+x^2)}. \quad (3.3)$$

here $x = r_1/r_2$, T_0 is the black hole temperature. From Eq. (3.2), we obtain

$$0 = T_0 - \frac{1}{4\pi r_2 x}(1+x) + \frac{Q^2}{4\pi r_2^3 x^3}(1+x^2)(1+x), \quad (3.4)$$

$$2P_0 = \frac{T_0}{2r_2 x}(1+x) - \frac{1}{8\pi r_2^2 x^2}(1+x^2) + \frac{Q^2}{8\pi r_2^4 x^4}(1+x^4) \quad (3.5)$$

From Eqs.(3.3) and (3.5), we find that

$$\frac{1}{4\pi r_2 x} = \frac{T_0(1+x)(1-2x+x^2)}{(1+x-4x^2+x^3+x^4)} + \frac{Q^2[(1+x^4)(1+x+x^2)-6x^3]}{4\pi r_2^3 x^3(1+x-4x^2+x^3+x^4)}. \quad (3.6)$$

From Eq. (3.4),

$$T_0 = \frac{(1+x)}{4\pi r_2 x} - \frac{Q^2}{4\pi r_2^3 x^3}(1+x)(1+x^2), \quad (3.7)$$

substituting Eq.(3.7) into Eq. (3.6), we obtain

$$r_2^2 = \frac{Q^2(1+2x-6x^2+2x^3+x^4)}{x^2(1-x)^2} = \frac{Q^2}{x^2}(1+4x+x^2) = Q^2 f_Q(x). \quad (3.8)$$

Substituting Eq.(3.8) into Eq. (3.7), we find that

$$T_0 = \frac{(1+x)}{4\pi x Q f^{1/2}(x)} - \frac{(1+x)(1+x^2)}{4\pi Q f^{3/2}(x) x^3} \quad (3.9)$$

Taking $T_0 = \chi T_c$, $T_c = \frac{\sqrt{6}}{18\pi Q}$ is critical temperature. When $0 < \chi < 1$, so Eq.(3.9) can be rewrite as

$$\chi x^2 f_Q^{3/2}(x) \frac{1}{3\sqrt{6}} = (1+x). \quad (3.10)$$

We can derive the radius of black hole horizon r_2 or r_1 with Eqs. (3.8) and (3.10). We find that the temperature T_0 and the pressure P_0 satisfy the conduction of the first-order phase transition according to the Ehrenfest's classification. From Eqs.(2.4),(3.8) and (3.10), we can plot the phase diagram $P - V$ with different χ in Fig.1.




Figure 1: (Color online) The simulated phase transition and the boundary of a two-phase coexistence on the base of isotherms in the $P - V$ diagram for charged anti-de Sitter black holes. The temperature of isotherms from top to bottom. The green line is the critical isotherm diagram. In the calculation setting $Q = 1$.

3.2 The construction of equal-area law in $T - S$ diagram

When the cosmological constant l is invariant, the horizontal axis of two-phase coexistence region in charged AdS black hole are S_2 and S_1 , respectively, the temperature on the vertical axis as $T_0 (T_0 \leq T_c)$, which depend on the horizon radius of black hole r_+ . From the Maxwell's equal area law,

$$T_0(S_2 - S_1) = \int_{S_1}^{S_2} T dS = \int_{r_1}^{r_2} \frac{1}{2} \left(1 + \frac{3r_+^2}{l^2} - \frac{Q^2}{r_+^2} \right) dr_+, \quad (3.11)$$

$$T_0 = \frac{1}{4\pi r_2} \left(1 + \frac{3r_2^2}{l^2} - \frac{Q_0^2}{r_2^2} \right), \quad T_0 = \frac{1}{4\pi r_1} \left(1 + \frac{3r_1^2}{l^2} - \frac{Q_0^2}{r_1^2} \right), \quad (3.12)$$

$$2\pi T_0 r_2^2 (1 - x^2) = r_2 (1 - x) + \frac{r_2^3}{l^2} (1 - x^3) - Q_0^2 \frac{1 - x}{r_2 x}. \quad (3.13)$$

From Eq.(3.12), we can obtain

$$0 = -\frac{r_2 - r_1}{r_2 r_1} + \frac{3}{l^2} (r_2 - r_1) + Q_0^2 \frac{r_2^3 - r_1^3}{r_2^3 r_1^3} = -\frac{1 - x}{r_2 x} + \frac{3r_2}{l^2} (1 - x) + \frac{Q_0^2}{r_2^3 x^3} (1 - x^3), \quad (3.14)$$

$$8\pi T_0 = \frac{r_2 + r_1}{r_2 r_1} + \frac{3}{l^2}(r_2 + r_1) - Q_0^2 \frac{r_2^3 + r_1^3}{r_2^3 r_1^3} = \frac{1+x}{r_2 x} + \frac{3}{l^2} r_2 (1+x) - \frac{Q_0^2}{r_2^3 x^3} (1+x^3). \quad (3.15)$$

Combining Eqs.(3.14) and (3.15), we can obtain

$$\frac{r_2^2}{l^2} x = 1 - \frac{Q_0^2(1+x-4x^2+x^3+x^4)}{r_2^2 x^2 (1-x)^2} = 1 - \frac{Q_0^2(1+3x+x^2)}{r_2^2 x^2}. \quad (3.16)$$

Substituting Eq.(3.16) into Eq.(3.14), we have

$$r_2^2 = \frac{Q_0^2(1+4x+x^2)}{x^2} = Q^2 f_Q(x). \quad (3.17)$$

We find that Eq.(3.17) is the same as Eq.(3.8). Substituting Eqs.(3.16) and (3.17) into Eq.(3.15), we can obtain

$$2\pi T_0 x Q f_Q^{3/2}(x) = (1+x) f_Q(x) - \frac{(1+3x+3x^2+x^3)}{x^2}. \quad (3.18)$$

Taking $T_0 = \chi T_c$, where $T_c = \frac{\sqrt{6}}{18\pi Q}$ is the critical temperature. When $0 < \chi < 1$, the Eq.(3.18) can be rewrite as

$$\chi x^2 f_Q^{3/2}(x) \frac{1}{3\sqrt{6}} = (1+x). \quad (3.19)$$

From Eqs.(2.4), (3.8) and (3.10), we can plot the $T - S$ phase diagram with different χ in Fig. 2.




Figure 2: (Color online) The simulated phase transition and the boundary of a two-phase coexistence on the base of isotherms in the $T - S$ diagram for charged anti-de Sitter black holes. The temperature of isotherms from top to bottom. The green line is the critical isotherm diagram.

3.3 The construction of equal-area law in $P - v$ diagram

The horizontal axis of two-phase coexistence region in charged AdS black hole are $v_2 = 2r_2$ and $v_1 = 2r_1$, respectively, the pressure on the vertical axis as P_0 . From the

Maxwell's equal area law,

$$P_0(v_2 - v_1) = \int_{v_1}^{v_2} P dv \quad (3.20)$$

So similarly, we can obtain

$$r_2^2 = Q^2 \frac{4(1-x^3) + 3(1+x^2)(1+x) \ln x}{x^2 3(2(1-x) + (1+x) \ln x)} = Q^2 F_Q(x). \quad (3.21)$$

Taking $T_0 = \chi T_c$, here $T_c = \frac{\sqrt{6}}{18\pi Q}$ is critical temperature. When $0 < \chi < 1$, we have

$$\chi x^3 F_Q^{3/2}(x) \frac{4}{3\sqrt{6}} = F_Q(x) x^2 (1+x) - (1+x^2)(1+x). \quad (3.22)$$

From Eqs. (3.21) and (3.22), we can plot phase diagram $P - v$ with different χ in Figs. 3. From Eqs. (3.19) and (3.22), we can plot $\chi - x$ with differential duality conjugate




Figure 3: (Color online) The simulated phase transition and the boundary of a two-phase coexistence on the base of isotherms in the $P - v$ diagram for charged anti-de Sitter black holes. The temperature of isotherms from top to bottom. The green line is the critical isotherm diagram. we have set $Q = 1$.

variables in Fig. 4.

From Eqs.(3.8), (3.11), (3.17) and (3.19), we know that the first-order phase transition point of charged AdS black hole is same with differential conjugate variables (P, V) and (T, S) in same temperature and the charge of AdS black hole is invariable. When we take the conjugate variable (P, v) , the first-order phase transition point of charged AdS black hole is different from that of charged AdS black hole with conjugate variable (P, V) and (T, S) , however, the second-order phase transition is the same. We know that the first-order phase transition point is independent of the the choice of conjugate variable with the same temperature for the particle numbers constant, so the conjugate variables (P, v) can not be used as the states parameters of the first-order phase transition of charged AdS black hole.




Figure 4: (Color online) The $\chi - x$ diagram for charged anti-de Sitter black holes with differential duality state parameters. The line corresponding with conjugate variables $P - V$, $P - v$ and $Q - \Phi$ from top to bottom.

From Eqs.(3.8),(3.17) and (3.21), the ratio of the phase transition point r_2 or r_1 to the charge of black hole is constant with a fixed temperature. So the phase transition of charged AdS black hole depends on the ratio of the phase transition point r_2 or r_1 to the charge Q of black hole, while it is independent of the radius r_2 or r_1 of the event horizon of the black hole. When the charge Q of charge AdS black hole is constant and the temperature is fixed, the radius of the horizon of black hole $r_+ < r_1$, the charged AdS black hole corresponds to the liquid phase of van der Waals system, the radius of the horizon of black hole $r_+ > r_2$, the charged AdS black hole corresponds to the gas phase of van der Waals system, the radius of the horizon of black hole $r_2 \geq r_+ \geq r_1$ corresponds to the gas-liquid coexistence of van der Waals system. So when the temperature $T < T_c$ of the charged AdS black hole, there are three different states in the same pressure P_0 . From the different ratio of the charge to the radius of black hole horizon of black hole, We define these different states as high potential state, medium potential state, low potential state.

3.4 The construction of equal-area law in $Q^2 - \Psi$ and $Q - \Phi$ diagram

When the charge of the AdS black hole of space is variant, taking $Q^2 - \Psi$ as independent duality parameter. The horizontal axis of two-phase coexistence region in charged AdS black hole are Ψ_2 and Ψ_1 , respectively, the temperature on the vertical axis as Q_0^2 , which is dependent on the horizon radius of black hole r_+ . From the Maxwell's equal area law,

$$Q_0^2(\Psi_1 - \Psi_2) = \int_{\Psi_2}^{\Psi_1} Q^2 d\Psi. \quad (3.23)$$

$$Q_0^2 = r_2^2 + \frac{3r_2^4}{l^2} - 4\pi r_2^3 T_0, \quad Q_0^2 = r_1^2 + \frac{3r_1^4}{l^2} - 4\pi r_1^3 T_0. \quad (3.24)$$

we can obtain

$$Q_0^2 = r_2^2 x + \frac{r_2^4}{l^2} x(1+x+x^2) - 2\pi T_0 r_2^3 x(1+x), \quad (3.25)$$

and

$$0 = (1+x) + \frac{3r_2^2}{l^2} (1+x^2)(1+x) - 4\pi T_0 r_2 (1+x+x^2), \quad (3.26)$$

$$2Q_0^2 = r_2^2(1+x^2) + \frac{3r_2^4}{l^2} (1+x^4) - 4\pi T_0 r_2^3 (1+x^3). \quad (3.27)$$

Combining Eqs. (3.25), (3.26) and (3.27), we can get

$$r_2 = \frac{(1+x)}{\pi T_0 (1+4x+x^2)} = \frac{f_l(x)}{\pi T_0}. \quad (3.28)$$

Taking $T_0 = \frac{T_c}{\chi} = \frac{1}{\chi\pi l} \sqrt{\frac{2}{3}}$, we can obtain [19]

$$3\chi^2(1+x)^2 = 2(1+4x+x^2). \quad (3.29)$$

Taking (Q, Φ) as independent conjugate variables, The horizontal axis of two-phase coexistence region in charged AdS black hole are Φ_2 and Φ_1 , respectively, the pressure on the vertical axis as Q_0 . From the Maxwell's equal area law,

$$Q_0(\Phi_1 - \Phi_2) = \int_{\Phi_2}^{\Phi_1} Q d\Phi. \quad (3.30)$$

So similarly, we can obtain

$$r_2 = \frac{(1+x)}{\pi T_0 (1+4x+x^2)} = \frac{f_l(x)}{\pi T_0}. \quad (3.31)$$

Setting $T_0 = \frac{T_c}{\chi} = \frac{1}{\chi\pi l} \sqrt{\frac{2}{3}}$, we can obtain

$$r_2 = \frac{(1+x)}{\pi T_0 (1+4x+x^2)} = \chi l \sqrt{\frac{3}{2}} f_l(x) \quad (3.32)$$

$$3\chi^2(1+x)^2 = 2(1+4x+x^2). \quad (3.33)$$

From Eqs. (2.4), (3.31) and (3.33), we can plot the phase diagram $Q^2 - \Psi$ and $Q - \Phi$ with different χ in Fig. 5. We can plot the $\chi - x$ from (3.32) in Fig.4. From Eqs.(3.28), (3.29), (3.31) and (3.33), we find that the phase transition point is same for the two




Figure 5: (Color online)(left) The behavior of isothermal $Q^2 - \Psi$ diagram of charged anti-de Sitter black holes. (right) The behavior of isothermal $Q - \Phi$ diagram of charged anti-de Sitter black holes. The temperature of isotherms increases from top to bottom. The green line is the critical isotherm diagram. we have set $l = 1$.

conjugate variables (Q^2, Ψ) and (Q, Φ) with the fixed temperature. So, the first laws of thermodynamics represented by the two conjugate variables (Q^2, Ψ) and (Q, Φ) are equivalent.

From Eq.(3.32), the ratio of the phase transition point r_2 or r_1 to the cosmological constant l of space is constant with a fixed temperature for the particle number variance. So the phase transition of charged AdS black hole dependent with the ratio of the phase transition point r_2 or r_1 to the cosmological constant l , is independent of the radius r_2 or r_1 of the event horizon of the black hole. When the cosmological constant l is constant and the temperature $T > T_c$ is fixed, the radius of the horizon of black hole $r_+ < r_1$, the charged AdS black hole corresponds to the liquid phase of van der Waals system, the radius of the horizon of black hole $r_+ > r_2$, the charged AdS black hole corresponds to the gas phase of van der Waals system, the radius of the horizon of black hole $r_2 \geq r_+ \geq r_1$ corresponds to the gas-liquid coexistence of van der Waals system. So when the temperature $T > T_c$ of the charged AdS black hole, there are three different states in the same pressure Q . From the different ratio of the radius of black hole horizon of black hole to the cosmological constant l of space, We define these different states as larger phase, medium phase, small phase.

4 Discussions and conclusions

The study of black hole thermodynamics have gained attention as the black hole, especially the thermodynamics of black hole directly involve the gravitation, statistics, particle physics and field theory [50–55]. Even so it is unclear in the precise statistical

description of a black hole's thermodynamic state the thermodynamic properties of black holes and the critical phenomenon in ordinary thermodynamic systems still is a research subject widely concerned.

In this paper, we choose different conjugate variables and use Maxwell's equal-area law to study the phase transition of charged AdS black hole. We find that the phase transition point which obtained by choosing the conjugate variable (P, V) and (T, S) , respectively, is equal at the same temperature. The phase transition point obtained by the conjugate variable (P, v) is different with that obtained by choosing the conjugate variable (P, V) and (T, S) . For the phase transition point of AdS black hole is the states function of system, and is independent of the process, so we think that the conjugate variable (P, v) cannot be independent conjugate variable to study the phase transition of AdS black hole.

From the study of these section 3.1, 3.2 and 3.3, we obtain that when the temperature $T < T_c$ of the charged AdS black hole, there are three differential states in the same pressure P_0 . From the differential ratio of the charge to the radius of black hole horizon of black hole, We define these different states as high potential state, medium potential state, low potential state. From the study of this section 3.4, when the temperature $T > T_c$ of the charged AdS black hole, there are three differential states in the same pressure Q . From the differential ratio of the radius of black hole horizon of black hole to the cosmological constant l , We define these different states as larger phase, medium phase, small phase. As we have seen, the phase transition of charged AdS black hole, not only depend on the event horizon radius of black hole, but also depend on the ratio the charge potential to the radius of black hole horizon. It would be worthwhile to better understand the critical behaviour of black hole, and perfect the self-consistent geometry theory of black hole thermodynamic.

Acknowledgements

We would like to thank Prof. Zong-Hong Zhu and Meng-Sen Ma for their indispensable discussions and comments. This work was supported by the Young Scientists Fund of the National Natural Science Foundation of China (Grant No.11205097), in part by the National Natural Science Foundation of China (Grant No.11475108), Supported by Program for the Innovative Talents of Higher Learning Institutions of Shanxi, the Natural Science Foundation of Shanxi Province, China (Grant No.201601D102004) and

the Natural Science Foundation for Young Scientists of Shanxi Province,China (Grant No.2012021003-4).

References

- [1] D. Kubiznak, R. B. Mann, *P-V criticality of charged AdS black holes*, JHEP 1207, 033 (2012), arXiv:1205.0559
- [2] B. P. Dolan, D. Kastor, D. Kubiznak, R. B. Mann, J. Traschen, *Thermodynamic Volumes and Isoperimetric Inequalities for de Sitter Black Holes*, Phys.Rev. D. 87. 104017, arXiv:1301.5926
- [3] S. Gunasekaran, D. Kubiznak, R. B. Mann, *Extended phase space thermodynamics for charged and rotating black holes and Born-Infeld vacuum polarization*, JHEP 1211, 110 (2012), arXiv:1208.6251
- [4] Rong-Gen Cai, Li-Min Cao, Li Li, and Run-Qiu Yang, *P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space*, JHEP 1309, 1-22 (2013), arXiv:1306.6233
- [5] Y. Sekiwa, *Thermodynamics of de Sitter black holes: Thermal cosmological constant*, Phys. Rev. D 73,084009 (2006), hep-th/0602269
- [6] Zi-xu Zhao, Ji-liang Jing, *Ehrenfest scheme for complex thermodynamic systems in full phase space*. JHEP 11, 037 (2014), arXiv:1405.2640
- [7] Meng-Sen Ma, Ren Zhao, Yan-Song Liu, *Phase transition and thermodynamic stability of topological black holes in Horava-Lifshitz gravity*, Classical and Quantum Gravity, 34, 165009 (2017), arXiv:1604.06998
- [8] Meng-Sen Ma, Rui-Hong Wang, *Peculiar P-V criticality of topological Hova-Lifshitz black holes*, Phys. Rev. D 96, 024052 (2017), arXiv:1707.09156
- [9] S. H. Hendi, B. Eslam Panah, S. Panahiyan, M. S. Talezadeh, *Geometrical thermodynamics and P-V criticality of the black holes with power-law Maxwell field*, Eur. Phys. J. C 77 (2017) 133, arXiv:1612.00721

- [10] Z. Dayyani, A. Sheykhi, M. H. Dehghani, *Counterterm method in Einstein dilaton gravity and the critical behavior of dilaton black hole with a power-Maxwell field*, Phys. Rev. D 95, 84004 (2017)
- [11] De-Cheng Zou, Yun-qi Liu, Rui-hong Yue, *Behavior of quasinormal modes and Van der Waals-like phase transition of charged AdS black holes in massive gravity*, Eur. Phys. J. C 77, 365 (2017), arXiv:1702.08118
- [12] Peng Cheng, Shao-Wen Wei, Yu-Xiao Liu, *Critical phenomena in the extended phase space of Kerr-Newman-AdS black holes*, Phys. Rev. D 94, 024025 (2016), arXiv:1603.08694
- [13] R. Banerjee, B. R. Majhi, S. Samanta, *Thermogeometric phase transition in a unified framework*, Phys. Lett. B 767 25-28 (2017), arXiv:1611.06701
- [14] R. Banerjee, D. Roychowdhury, *Thermodynamics of phase transition in higher dimensional AdS black holes*, JHEP 11,004(2011),arXiv:1109.2433
- [15] R. Banerjee, D. Roychowdhury, *Critical behavior of Born Infeld AdS black holes in higher dimensions*, Phys. Rev. D 85, 104043 (2012) arXiv:1203.0118
- [16] A. Dey, S. Mahapatra and T. Sarkar, *Thermodynamics and entanglement entropy with Weyl corrections*, Phys. Rev. D 94, 026006(2016), arXiv:1512.07117
- [17] K. Bhattacharya, B. R. Majhi, S. Samanta, *van der Waals criticality AdS black holes: A phenomenological study*, Phys. Rev. D 96, 084037 (2017)
- [18] Xiao-Xiong Zeng, Li-Fang Li, *van der Waals phase transition in the framework of holography*, Phys. Lett. B 764, 100 (2017)
- [19] A. Dehyadegari, A. Sheykhi, A. Montakhap, *Critical behavior and microscopic structure of charged AdS black holes*, Phys. Lett. B 768, 235(2017), arXiv:1607.05333
- [20] Rong-Gen Cai, Shan-Ming Ruan, Shao-Jiang Wang, Run-Qiu Yang, Rong-Hui Peng, *Complexity Growth for AdS Black Holes*, JHEP 1609, 161(2016), arXiv:1606.08307

- [21] Jia-Lin Zhang, Rong-Gen Cai, Hongwei Yu, *Phase transition and Thermodynamical geometry of Reissner-Nordström-AdS Black Holes in Extended Phase Space*, Phys. Rev. D 91, 044028(2015), arXiv:1502.01428
- [22] Jia-Lin Zhang, Rong-Gen Cai, Hongwei Yu, *Phase transition and thermodynamical geometry for Schwarzschild AdS black hole in AdS⁵ spacetime*, JHEP 02, 143(2015), arXiv:1409.5305
- [23] Shao-Wen Wei, Yu-Xiao Liu, *Photon orbits and thermodynamic phase transition of d -dimensional charged AdS black holes*, Phys. Rev. D 97, 104027(2018), arXiv:1711.01522
- [24] Shou-Long Li, Hao. Wei, *Holographic Entanglement Entropy and Van der Waals transitions in Einstein-Maxwell-Dilaton theory*, arXiv:1809.03810
- [25] Shou-Long, Hong-Da Lyu, Hua-Kai Deng and Hao. Wei, *$P - v$ Criticality in Gauged Supergravities*, arXiv:1809.03471
- [26] Yan-Gang Miao and Zhen-Ming Xu, *Thermodynamics of Noncommutative High-Dimensional AdS Black Holes with Non-Gaussian Smeared Matter Distributions*, Eur. Phys. J. C, 76, 217(2016), arXiv:1511.00853.
- [27] Yan-Gang Miao and Zhen-Ming Xu, *Validity of Maxwell Equal Area Law for Black Holes Conformally Coupled to Scalar Fields in AdS₅ Spacetime*, Eur. Phys. J. C, 77, 403(2017), arXiv:1610.01769
- [28] Yi-Fei Wang, Ming Zhang, Wen-Biao Liu, *Coexistence curve and molecule number density of AdS topological charged black hole in massive gravity*, arXiv: 1711.04403
- [29] A. Dehyadegari, A. Sheykhi, *Reentrant phase transition of Born-Infeld-AdS black holes*, Phys. Rev. D 98, 024011(2018), arXiv:1711.01151
- [30] K. Jafarzade, J. Sadeghi, *Phase Transition of charged Rotational Black Hole and Quintessence*, arXiv:1710.08642
- [31] Ren Zhao, Li-Chun Zhang *Entropy of higher-dimensional charged de Sitter black holes and phase transition*, Commun. Theor. Phys. 70, 578-584(2018), arXiv:1710.07225

- [32] A. Övgün, *P-v criticality of a specific black hole in $f(R)$ gravity coupled with Yang-Mills field*, Advances in High Energy Physics, 2018, 8153721(2018), arXiv:1710.06795
- [33] Z. Dayyani, A. Sheykhi, M. H. Dehghani, S. Hajkhalili, *Critical behavior and phase transition of dilaton black holes with nonlinear electrodynamics*, Eur. Phys. J. C 78, 152 (2018), arXiv:1709.06875
- [34] M. K. Zangeneh, A. Dehyadegari, A. Sheykhi, R. B. Mann, *Microscopic Origin of Black Hole Reentrant Phase Transitions*, Phys. Rev. D 97, 084054 (2018), arXiv:1709.04432
- [35] Ming Zhang, De-Cheng Zou, Rui-Hong Yue, *Reentrant phase transitions and triple points of topological AdS black holes in Born-Infeld-massive gravity*, Advances in High Energy Physics 2017, 3819246 (2017), arXiv:1707.04101
- [36] A. Sahay, R. Jha, *Geometry of criticality, supercriticality and Hawking-Page transitions in Gauss-Bonnet-AdS black holes*, Phys. Rev. D 96, 126017 (2017), arXiv:1707.03629
- [37] Xiao-Xiong Zeng, Yi-Wen Han, *Holographic Van der Waals phase transition for a hairy black hole*, Advances in High Energy Physics, 2017, 2356174(2017), arXiv:1706.02024
- [38] Shao-Wen Wei, Bin Liang, Yu-Xiao Liu, *Critical phenomena and chemical potential of charged AdS black hole*, Phys. Rev. D 96, 124018 (2017), arXiv:1705.08596
- [39] Shao-Wen Wei, Yu-Xiao Liu, *Critical phenomena and thermodynamic geometry of charged Gauss-Bonnet AdS black holes*, Phys. Rev. D 87, 044014 (2013), arXiv:1209.1707
- [40] Ren Zhao, Hui-Hua Zhao, Meng-Sen Ma, Li-Chun Zhang *On the critical phenomena and thermodynamics of charged topological dilaton AdS black holes*, Eur. Phys. J. C 73, 2645(2013)
- [41] Huai-fan Li, Meng-Sen Ma, Li-Chun Zhang, and Ren Zhao, *Entropy of Kerr-de Sitter black hole*. Nuclear Physics B 920, 211-220 (2017)

- [42] Huai-fan Li Hui-Hua Zhao, Li-Chun Zhang, and Ren Zhao, *Clapeyron equation and phase equilibrium properties in higher dimensional charged topological dilaton AdS black holes with a nonlinear source*, Eur. Phys. J. C (2017) 77:295
- [43] Huai-Fan Li, Xiong-ying Guo, Hui-Hua Zhao, Ren Zhao, *Maxwell's equal area law for black holes in power Maxwell invariant*, General Relativity and Gravitation. 2017, 49(8): 111 [1610.05428]
- [44] Hui-Hua Zhao, Li-Chun Zhang, and Ren Zhao, *Two-Phase Equilibrium Properties in Charged Topological Dilaton AdS Black Holes*, Advances in High Energy Physics, 2016, 2021748(2016)
- [45] Hui-Hua Zhao , Li-Chun Zhang, Meng-Sen. Ma, and Ren. Zhao, *Phase transition and clapeyron equation of black holes in higher dimensional AdS spacetime*, Class. Quantum Grav. 32 (2015) 145007
- [46] Li-Chun Zhang, Hui-Hua Zhao, Ren Zhao, Meng-Sen Ma, *Equal Area Laws and Latent Heat for d-Dimensional RN-AdS Black Hole*, Advances in High Energy Physics, 816728 (2014)
- [47] A. Belhaj, M. Chabab, H. Ei Moumni, K. Masmar, and M. B. Sedra, *Maxwell's equal-area law of Gauss-bonnet-Anti-de Sitter black hole*, Eur. Phys. J. C (2015) 75:71
- [48] E. Spallucci, A. Smailagic, *Maxwells equal-area law for charged Anti-de Sitter black holes*, Phys. Lett. B, 723, 436(2013), arXiv:1305.3379
- [49] D, Kastor, S, Ray, and J, Traschen, *Enthalpy and the Mechanics of AdS Black Holes*, Class. Quant. Grav. 26, 195011 (2009).
- [50] J. D. Bekenstein, *Black holes and the second law*, Lett. Nuovo Cimento, 4, 737 (1972).
- [51] J. D. Bekenstein, *Generalized second law of thermodynamics in black hole physics*, Phys. Rev. D 9, 3292 (1974).
- [52] J. D. Bekenstein, *Extraction of energy and charge from a black hole*, Phys. Rev. D 7, 949 (1973).

- [53] J. M. Bardeen, B. Carter, S. W. Hawking, *The Four laws of black hole mechanics*, Commun. Math. Phys. 31, 161 (1973).
- [54] S. W. Hawking, *Black Hole Explosions*, Nature, 248, 30 (1974).
- [55] S. W. Hawking, *Particle Creation by Black Holes*, Commun. Math. Phys. 43, 199 (1975).