

The $T\bar{T}$ perturbation and its geometric interpretation

Riccardo Conti,¹ Stefano Negro,² Roberto Tateo¹

¹*Dipartimento di Fisica and Arnold-Regge Center, Università di Torino, and INFN, Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy.*

²*C.N. Yang Institute for Theoretical Physics, New York Stony Brook, NY 11794-3840. U.S.A.*

E-mail: riccardo.conti@to.infn.it, stefano.negro@stonybrook.edu,
tateo@to.infn.it

ABSTRACT: Starting from the recently-discovered $T\bar{T}$ -perturbed Lagrangians, we prove that the deformed solutions to the classical EoMs for bosonic field theories are equivalent to the unperturbed ones but for a specific field-dependent local change of coordinates. This surprising geometric outcome is fully consistent with the identification of $T\bar{T}$ -deformed 2D quantum field theories as topological JT gravity coupled to generic matter fields. Although our conclusion is valid for generic interacting potentials, it first emerged from a detailed study of the sine-Gordon model and in particular from the fact that solitonic pseudo-spherical surfaces embedded in $\mathbb{R}^3$ are left invariant by the deformation. Analytic and numerical results concerning the perturbation of specific sine-Gordon soliton solutions are presented.

Contents

1	Introduction	1
2	Classical integrable equations and embedded surfaces	3
2.1	Construction of the solitonic surfaces	4
2.2	The case of sine-Gordon	5
3	The $T\bar{T}$-deformed sine-Gordon model and its associated surfaces	6
3.1	From the deformed to the undeformed model through a local change of coordinates	9
4	A geometric map for N-boson fields and arbitrary potential	10
5	$T\bar{T}$-deformed soliton solutions in the sine-Gordon model	13
5.1	The one-kink solution	13
5.2	The two-kink solution	15
5.3	The breather	16
6	The shock-wave phenomenon and the Hagedorn-type transition	18
7	Conclusions	21
A	Short review on surfaces embedded in $\mathbb{R}^3$	22
B	Computation of the fundamental quadratic forms from sine-Gordon ZCR	26

1 Introduction

The deformation of 2D quantum field theories [1, 2] by the Zamolodchikov’s $T\bar{T}$ operator [3], has recently attracted the attention of theoretical physicists due to the many important links with string theory [4–7] and AdS/CFT [8–17].

A remarkable property of this perturbation, discovered in [1, 2], concerns the evolution of the quantum spectrum at finite volume R , with periodic boundary conditions, in terms of the $T\bar{T}$ coupling constant τ . The spectrum is governed by the inhomogeneous Burgers equation

$$\partial_\tau E_n(R, \tau) = \frac{1}{2} \partial_R (E_n^2(R, \tau) - P_n^2(R)) , \quad (1.1)$$

where $E_n(R, \tau)$ and $P_n(R)$ are the total energy and momentum of a generic energy eigenstate $|n\rangle$, respectively. Equation (1.1) is valid also for non-integrable models.

Notice that (1.1) reveals an important feature of $T\bar{T}$ -deformed QFTs: the interaction between the perturbing operator and the geometry, through the coupling τ . The latter property is a basic requirement for any sensible theory of gravity but in the current case it naturally emerges, non perturbatively and at full quantum level, from a specific irrelevant perturbation of Lorentz-invariant Quantum Field Theories (QFTs). An important link with JT topological gravity was noticed and studied in [18], where it was shown that JT gravity coupled to matter leads to a scattering phase matching that associated to the $T\bar{T}$ perturbation [1, 2, 4–6, 19–21].

Studies of partition functions [2, 22–24] have led to a proof of the uniqueness of this perturbation [25] under the assumption that the theory on the torus is invariant under modular transformations and that the energy of a given eigenstate is a function only of τ and of the energy and momentum of the corresponding state at $\tau = 0$. Furthermore, starting from the JT-gravity setup, in [23] the hydrodynamic-type equation (1.1) was recovered. The latter result together with [18] confirms, beyond any reasonable doubt, the equivalence between the $T\bar{T}$ deformation and JT topological gravity coupled to generic matter field.

The aim of this paper is to address the problem concerning the classical interpretation of the $T\bar{T}$ perturbation following the more direct approach proposed in [2] and further developed in [26, 27]. The current analysis is based on the observation [1, 2] that (1.1) directly implies a self-consistent flow equation for the deformed Lagrangian $\mathcal{L}^{(\tau)}$

$$\partial_\tau \mathcal{L}^{(\tau)} = \text{Det} \left(T_{\mu\nu}^{(\tau)} \right) , \quad T_{\mu\nu}^{(\tau)} = -\frac{2}{\sqrt{|g|}} \frac{\delta(\mathcal{L}^{(\tau)} \sqrt{|g|})}{\delta g^{\mu\nu}} , \quad (1.2)$$

where $g = \text{Det} (g_{\mu\nu})$ and $T\bar{T} = -\pi^2 \text{Det} (T_{\mu\nu})$ is the classical counterpart of Zamolodchikov’s operator.

Starting from the unperturbed Lagrangian $\mathcal{L}^{(0)}$ equation (1.2) can be solved giving the $T\bar{T}$ -deformed exact result $\mathcal{L}^{(\tau)}$. Adopting this strategy, the Nambu-Goto classical Lagrangian in the static gauge was recovered [2] along with the deformation of bosonic models with generic interacting potential [2, 26, 27], WZW and σ -models [16, 26, 28], and the Thirring model [26].

There are many reasons to study these newly-discovered set of classical Lagrangians. First of all, according to [18, 23], these systems should correspond to JT gravity coupled to non-topological matter, a fact that is by no mean evident from the Lagrangian point of view.

Secondly, when the starting model is integrable, there should be a general way to deform the whole integrable model machinery. For example, a generalisation of the ODE/IM correspondence [29–31] should lead to an alternative method to obtain the quantum spectrum at finite volume [32, 33] and it may open the way to the inclusion of the $T\bar{T}$ inside the Wilson Loops/Scattering Amplitudes setup, in $\text{AdS}_5/\text{CFT}_4$ [34, 35] and perhaps also to consistently deform the Argyres-Douglas theory [36–38].

The main purpose of this article is to prove that, for bosonic theories with arbitrary interacting potentials, the $T\bar{T}$ perturbation has indeed the alternative interpretation as a

space-time deformation. In Euclidean coordinates the change of variables is

$$dx^\mu = \left(\delta^\mu_\nu + \tau \tilde{T}^\mu_\nu(\mathbf{y}) \right) dy^\nu, \quad \mathbf{y} = (y^1, y^2), \quad (1.3)$$

$$dy^\mu = \left(\delta^\mu_\nu + \tau (\tilde{T}^{(\tau)})^\mu_\nu(\mathbf{x}) \right) dx^\nu, \quad \mathbf{x} = (x^1, x^2), \quad (1.4)$$

with $\tilde{T}^\mu_\nu = -\epsilon^\mu_\rho \epsilon^\sigma_\nu T^\rho_\sigma$ and $(\tilde{T}^{(\tau)})^\mu_\nu = -\epsilon^\mu_\rho \epsilon^\sigma_\nu (T^{(\tau)})^\rho_\sigma$, where $T = T^{(0)}$ and $T^{(\tau)}$ are the unperturbed and perturbed stress-energy tensor in the set of coordinates $\mathbf{y}$ and $\mathbf{x}$, respectively. Then, any solution of the perturbed EoMs can be mapped onto the $\tau = 0$ corresponding solution, *i.e.*

$$\phi^{(\tau)}(\mathbf{x}) = \phi^{(0)}(\mathbf{y}(\mathbf{x})), \quad (1.5)$$

where the r.h.s. of (1.5)¹ is defined on a deformed space-time with metric

$$\mathbf{g}'_{\mu\nu} = \delta_{\mu\nu} - \tau \epsilon_{\mu\rho} \epsilon^\sigma_\nu (2T + \tau T^2)^\rho_\sigma. \quad (1.7)$$

In fact (1.4) corresponds to a natural generalization of the Virasoro conditions used in the GGRT treatment of the NG string [39],² and it matches precisely the generalisation corresponding to classical JT gravity [18, 23].

2 Classical integrable equations and embedded surfaces

It is an established fact that integrable equations in two dimensions admit an interpretation in terms of surfaces embedded inside an N -dimensional space. The two oldest examples of this connection, dating back to the works of 19th century geometers [40, 41], are the sine-Gordon and Liouville equations. They appear as the Gauss-Mainardi-Codazzi (GMC) system of equations (A.14) for, respectively, pseudo-spherical and minimal surfaces embedded in the Euclidean space $\mathbb{R}^3$. As proved by Bonnet [42], any surface embedded in $\mathbb{R}^3$ is uniquely determined (up to its position in the ambient space) by two rank 2 symmetric tensors: the metric $g_{\mu\nu}$ (A.4) and the second fundamental tensor $d_{\mu\nu}$ (A.6). Their intuitive role is to measure, respectively, the length of an infinitesimal curve and the displacement of its endpoint from the tangent plane at the starting point. One can then use $g_{\mu\nu}$ and $d_{\mu\nu}$ to study the motion of a frame anchored to the surface. The result is a system of linear differential equations, known as *Gauss-Weingarten* equations (A.9, A.10). The GMC system appears then as the consistency condition for this linear system, effectively constraining the “moduli space” consisting of the two tensors $g_{\mu\nu}$ and $d_{\mu\nu}$.

The search for a general correspondence originated in the works of Lund, Regge, Pohlmeyer and Getmanov [43–45] and was subsequently formalised by Sym [46–50] who showed that any

¹Notice that from (1.5) it follows that $\phi^{(\tau)}(\mathbf{x})$ fulfills the Burgers-type equation

$$\partial_\tau \phi^{(\tau)}(\mathbf{x}) + (\partial_\tau x^\mu) \partial_\mu \phi^{(\tau)}(\mathbf{x}) = 0, \quad (1.6)$$

which may justify the wave-breaking phenomena observed in section 5. In our results x^μ is always linear in τ , however we could not find an explicit expression for $\partial_\tau x^\mu$ valid in general.

²See [8] for a clarifying discussion related to the current topic.

integrable system whose associated linear problem is based on a semi-simple Lie algebra $\mathfrak{g}$ can be put in the form of a GMC system for a surface embedded in a $\dim(\mathfrak{g})$ -dimensional surface.³ In this section we will shortly review Sym's results for the general setting and concentrate on the case of sine-Gordon model. We will use the following conventions

$$\mathbf{z} = (z^1, z^2) , \quad \partial_\mu f(\mathbf{z}) \equiv \frac{\partial}{\partial z^\mu} f(\mathbf{z}) , \quad \forall f : \mathbb{R}^2 \rightarrow \mathbb{R} , \quad \mu = 1, 2 .$$

2.1 Construction of the solitonic surfaces

Let us consider a generic 2-dimensional system of non-linear partial differential equations for a set of real fields $\{\phi_i(\mathbf{z})\}$ admitting a *Zero Curvature Representation* (ZCR) for a pair of functions L_1 and L_2 taking values in a d -dimensional representation of a semi-simple Lie algebra⁴ $\mathfrak{g}$ ($\dim(\mathfrak{g}) = N$):

$$\partial_2 L_1 - \partial_1 L_2 + [L_1, L_2] = 0 . \quad (2.1)$$

The functions L_μ depend on $\mathbf{z}$ through the fields $\phi_i(\mathbf{z})$ and their derivatives and on a real spectral parameter λ :

$$L_\mu \equiv L_\mu(\mathbf{z}|\lambda) \equiv L_\mu(\{\phi_i(\mathbf{z})\}, \{\partial_\nu \phi_i(\mathbf{z})\}, \dots | \lambda) . \quad (2.2)$$

The Zero Curvature Representation can be interpreted as the compatibility condition for a system of first-order linear partial differential equations involving an auxiliary $d \times d$ matrix-valued function $\Phi \equiv \Phi(\mathbf{z}|\lambda)$

$$\partial_\mu \Phi = L_\mu \Phi , \quad \mu = 1, 2 , \quad (2.3)$$

commonly known as *associated linear problem*. Assuming $\Phi(\mathbf{z}_0|\lambda) \in G$ as initial condition, with G being the Lie group associated to $\mathfrak{g}$, equation (2.3) allows, in principle, to recover a single-valued function $\Phi \in G$ in the whole $\mathbb{R}^2$. This function can then be used to construct the following object

$$r(\mathbf{z}|\lambda) = \Phi^{-1}(\mathbf{z}|\lambda) \frac{\partial}{\partial \lambda} \Phi(\mathbf{z}|\lambda) , \quad (2.4)$$

which is interpreted as the coordinate description of a λ -family of surfaces embedded into the N -dimensional affine space $\mathfrak{g}$. Moreover, equipping the affine space $\mathfrak{g}$ with a non-degenerate scalar product (i.e. the Killing form of the semi-simple Lie algebra), we can convert $\mathfrak{g}$ into an N -dimensional flat space. In other words, we can find an orthonormal basis $\{\mathbf{e}^i\}$ of $\mathfrak{g}$ with respect to the Killing form and then extract the quantities r_i from the identity

$$r = \sum_{i=1}^N r_i \mathbf{e}^i = \Phi^{-1}(\mathbf{z}|\lambda) \frac{\partial}{\partial \lambda} \Phi(\mathbf{z}|\lambda) . \quad (2.5)$$

³An interesting additional result of Sym concerns the existence of the same kind of connection for spin systems and σ -models.

⁴Here we abuse notations by denoting with $\mathfrak{g}$ both the algebra and its d -dimensional representation. The same applies for the associated Lie Group G .

The vector $\mathbf{r} = (r_1, r_2, \dots, r_N)^T$ is then the position vector of a family of surfaces embedded in N -dimensional flat space,⁵ parametrised by λ . These are called *solitonic surfaces* and satisfy the following properties:

1. their GMC system reduces to the ZCR (2.1). This means that any integrable system whose EoMs can be represented as a ZCR depending on a spectral parameter λ , can be associated to a particular class of surfaces;
2. they are invariant with respect to λ -independent gauge transformation of the pair L_μ . This fact provides a way to prove the equivalence of distinct soliton systems up to gauge transformations and independent coordinate redefinitions, see [49];
3. their metric tensor (induced by the flat space $\mathfrak{g}$) is explicitly computed from the pair L_μ as

$$g_{\mu\nu} = \text{Tr} \left(\mathbf{Ad} \left(\frac{\partial L_\mu}{\partial \lambda} \right) \mathbf{Ad} \left(\frac{\partial L_\nu}{\partial \lambda} \right) \right), \quad (2.6)$$

where $\mathbf{Ad}$ denotes the adjoint representation of the algebra $\mathfrak{g}$. Consequently, any intrinsic property of the soliton surface is determined uniquely by the ZCR.

2.2 The case of sine-Gordon

Let us now consider the specific case of the sine-Gordon equation

$$\partial \bar{\partial} \phi = \frac{m^2}{\beta} \sin(\beta \phi), \quad (2.7)$$

where we set $\mathbf{z} = (z^1, z^2) = (z, \bar{z})$. The ZCR for this model is well known

$$L_1^{\text{sG}} = \mathcal{Z} = \frac{\beta}{2} \partial \phi \mathcal{S}^3 + \mathfrak{i} m \lambda \left[\cos \left(\frac{\beta}{2} \phi \right) \mathcal{S}^1 - \sin \left(\frac{\beta}{2} \phi \right) \mathcal{S}^2 \right], \quad (2.8)$$

$$L_2^{\text{sG}} = \bar{\mathcal{Z}} = -\frac{\beta}{2} \bar{\partial} \phi \mathcal{S}^3 + \mathfrak{i} \frac{m}{\lambda} \left[\cos \left(\frac{\beta}{2} \phi \right) \mathcal{S}^1 + \sin \left(\frac{\beta}{2} \phi \right) \mathcal{S}^2 \right], \quad (2.9)$$

where $\mathcal{S}^j$ are the generators of $\mathfrak{su}(2)$

$$[\mathcal{S}^i, \mathcal{S}^j] = \varepsilon^{ij}_k \mathcal{S}^k. \quad (2.10)$$

Since $\dim(\mathfrak{su}(2)) = 3$, we know that we are dealing with a surface embedded in the Euclidean plane $\mathbb{R}^3$ ($\mathfrak{su}(2)$ is compact). As mentioned in section 2, Bonnet theorem [42] tells us that any surface in $\mathbb{R}^3$ is completely specified (modulo its position) by its first and second fundamental

⁵The signature of this space depends on the real form chosen for the algebra; for example $\mathfrak{sl}(2) \simeq \mathfrak{so}(2, 1)$ give rise to surfaces in Minkowski space $\mathbb{R}^{2,1}$.

quadratic forms, which can be computed easily:⁶

$$\text{I}^{\text{sG}} = g_{\mu\nu}^{\text{sG}} dz^\mu dz^\nu = 2m^2 \left[(dz)^2 - \frac{2}{\lambda^2} \cos(\beta\phi) dz d\bar{z} + \frac{1}{\lambda^4} (d\bar{z})^2 \right], \quad (2.11)$$

$$\text{II}^{\text{sG}} = d_{\mu\nu}^{\text{sG}} dz^\mu dz^\nu = 2m^2 \frac{\sqrt{2}}{\lambda} \sin(\beta\phi) dz d\bar{z}. \quad (2.12)$$

From (2.11) and (2.12) one can then extract the Gaussian and the mean curvatures using (A.8):

$$K^{\text{sG}} = \text{Det} \left(d_{\mu\rho}^{\text{sG}} (g^{\text{sG}})^{\rho\nu} \right) = -\frac{\lambda^2}{2}, \quad H^{\text{sG}} = d_{\mu\nu}^{\text{sG}} (g^{\text{sG}})^{\nu\mu} = \frac{\lambda}{\sqrt{2}} \cot(\beta\phi), \quad (2.13)$$

with $g_{\mu\nu}^{\text{sG}} (g^{\text{sG}})^{\nu\rho} = \delta_\mu^\rho$. The fact that K^{sG} is constant negative tells us that we are dealing with a pseudo-spherical surface, which we were expecting from the old results of Bour [40]. Thus, for this specific case, the solitonic surfaces correspond to pseudo-spherical ones, with the spectral parameter λ playing the role of Gaussian curvature.

3 The $\text{T}\bar{\text{T}}$ -deformed sine-Gordon model and its associated surfaces

Let us now apply the Sym formalism sketched above to the $\text{T}\bar{\text{T}}$ -deformed sine-Gordon model [27]

$$\partial \left(\frac{\bar{\partial}\phi}{S} \right) + \bar{\partial} \left(\frac{\partial\phi}{S} \right) = \frac{V'}{4S} \left(\frac{S+1}{1-\tau V} \right)^2, \quad (3.1)$$

$$S = \sqrt{1 + 4\tau(1-\tau V) \partial\phi \bar{\partial}\phi}, \quad (3.2)$$

$$V = 2 \frac{m^2}{\beta^2} (1 - \cos(\beta\phi)), \quad V' = 2 \frac{m^2}{\beta} \sin(\beta\phi), \quad (3.3)$$

and derive the geometric properties of the associated surfaces. We start with the ZCR, which was found in [27]

$$L_1^{\text{T}\bar{\text{T}}} \equiv \mathcal{Z} = \beta \frac{\partial\phi}{2S} \mathcal{S}^3 + 2im \left[F_+ \cos\left(\frac{\beta}{2}\phi\right) \mathcal{S}^1 - F_- \sin\left(\frac{\beta}{2}\phi\right) \mathcal{S}^2 \right], \quad (3.4)$$

$$L_2^{\text{T}\bar{\text{T}}} \equiv \bar{\mathcal{Z}} = -\beta \frac{\bar{\partial}\phi}{2S} \mathcal{S}^3 + 2im \left[\bar{F}_+ \cos\left(\frac{\beta}{2}\phi\right) \mathcal{S}^1 + \bar{F}_- \sin\left(\frac{\beta}{2}\phi\right) \mathcal{S}^2 \right], \quad (3.5)$$

⁶These can be recovered by plugging (2.5) in the classical geometry formulae

$$g_{\mu\nu} = \partial_\mu \mathbf{r} \cdot \partial_\nu \mathbf{r}, \quad d_{\mu\nu} = -\partial_\mu \partial_\nu \mathbf{r} \cdot \mathbf{n},$$

where $\mathbf{n}$ is the normal unit vector to the plane spanned by $\partial_1 \mathbf{r}$ and $\partial_2 \mathbf{r}$:

$$\mathbf{n} = \frac{\partial_1 \mathbf{r} \times \partial_2 \mathbf{r}}{|\partial_1 \mathbf{r} \times \partial_2 \mathbf{r}|}.$$

where

$$F_+ = \left(\lambda B_+ + \frac{1}{\lambda} (\partial\phi)^2 B_- \right), \quad F_- = \left(\lambda B_+ - \frac{1}{\lambda} (\partial\phi)^2 B_- \right), \quad (3.6)$$

$$\bar{F}_+ = \left(\frac{1}{\lambda} B_+ + \lambda (\bar{\partial}\phi)^2 B_- \right), \quad \bar{F}_- = \left(\frac{1}{\lambda} B_+ - \lambda (\bar{\partial}\phi)^2 B_- \right), \quad (3.7)$$

with

$$B_+ = \frac{(S+1)^2}{8S(1-\tau V)}, \quad B_- = \frac{\tau}{2S}. \quad (3.8)$$

Again we have a ZCR based on the algebra $\mathfrak{su}(2)$ and thus a surface embedded in $\mathbb{R}^3$. We need then to recover the fundamental forms I and II, whose computation, although straightforward as in the case of sine-Gordon, is lengthy and cumbersome. Sparing the uninteresting details, we present directly the results

$$\text{I}^{\text{T}\bar{\text{T}}} = g_{\mu\nu}^{\text{T}\bar{\text{T}}} dz^\mu dz^\nu = \frac{m^2}{2S^2} \left(\frac{S+1}{1-\tau V} \right)^2 \hat{g}_{\mu\nu} dz^\mu dz^\nu, \quad (3.9)$$

$$\text{II}^{\text{T}\bar{\text{T}}} = d_{\mu\nu}^{\text{T}\bar{\text{T}}} dz^\mu dz^\nu = \frac{m^2 \sin(\beta\phi)}{\sqrt{2}\lambda(1-\tau V)} \left(\frac{S+1}{S} \right)^2 \hat{d}_{\mu\nu} dz^\mu dz^\nu, \quad (3.10)$$

where the matrices $\hat{g}_{\mu\nu}$ and $\hat{d}_{\mu\nu}$ are

$$\hat{g}_{\mu\nu} = \begin{pmatrix} \left(\frac{S+1}{2} - \frac{S-1}{2\lambda^2} \frac{\partial\phi}{\partial\phi} \right)^2 + \frac{S^2-1}{4\lambda^2} \frac{\beta^2 V}{m^2} \frac{\partial\phi}{\partial\phi} & \frac{S^2-1}{4} \left(\frac{\bar{\partial}\phi}{\partial\phi} + \frac{1}{\lambda^4} \frac{\partial\phi}{\partial\phi} \right) - \frac{S^2+1}{2\lambda^2} \cos(\beta\phi) \\ \frac{S^2-1}{4} \left(\frac{\bar{\partial}\phi}{\partial\phi} + \frac{1}{\lambda^4} \frac{\partial\phi}{\partial\phi} \right) - \frac{S^2+1}{2\lambda^2} \cos(\beta\phi) & \left(\frac{S+1}{2\lambda^2} - \frac{S-1}{2} \frac{\bar{\partial}\phi}{\partial\phi} \right)^2 + \frac{S^2-1}{4\lambda^2} \frac{\beta^2 V}{m^2} \frac{\bar{\partial}\phi}{\partial\phi} \end{pmatrix}_{\mu\nu},$$

$$\hat{d}_{\mu\nu} = \begin{pmatrix} \tau (\partial\phi)^2 \frac{S^2+1}{4(1-\tau V)} \\ \frac{S^2+1}{4(1-\tau V)} \tau (\bar{\partial}\phi)^2 \end{pmatrix}_{\mu\nu}. \quad (3.11)$$

One easily verifies that in the $\tau \rightarrow 0$ limit, which implies $S \rightarrow 1$, one recovers the fundamental forms of sine-Gordon

$$\text{I}^{\text{T}\bar{\text{T}}} \xrightarrow{\tau \rightarrow 0} 2m^2 \begin{pmatrix} 1 & -\frac{1}{\lambda^2} \cos(\beta\phi) \\ -\frac{1}{\lambda^2} \cos(\beta\phi) & \frac{1}{\lambda^4} \end{pmatrix}_{\mu\nu} dz^\mu dz^\nu = \text{I}^{\text{sG}}, \quad (3.12)$$

$$\text{II}^{\text{T}\bar{\text{T}}} \xrightarrow{\tau \rightarrow 0} m^2 \frac{\sqrt{2}}{\lambda} \sin(\beta\phi) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}_{\mu\nu} dz^\mu dz^\nu = \text{II}^{\text{sG}}. \quad (3.13)$$

What is striking about the matrices (3.11) is that, although their dependence on τ is complicated, they recombine in such a way that the Gaussian and mean curvature do not depend explicitly on it! In fact these two geometric invariants are exactly the same as the unperturbed sine-Gordon model:

$$K^{\text{T}\bar{\text{T}}} = -\frac{\lambda^2}{2} = K^{\text{sG}}, \quad H^{\text{T}\bar{\text{T}}} = \frac{\lambda}{\sqrt{2}} \cot(\beta\phi) = H^{\text{sG}}. \quad (3.14)$$

This suggests that the solitonic surface corresponding to a particular solution of the $T\bar{T}$ -deformed sine-Gordon equation is the same as the one associated to the undeformed model, what changes should be the coordinate system used to describe it. For the sake of completeness, we have reported in Figure 1 examples of embedded pseudo-spherical surfaces related to one-kink solutions, a stationary breather and a two-kink solution. The plots were obtained implementing the method described in [51]. The embedded surfaces in $\mathbb{R}^3$, as we have just




Figure 1. Pseudo-spherical solitonic surfaces associated to kink and breather solutions. Figure 1a represents the Dini surface, corresponding to a moving kink, while in Figure 1b the famous Beltrami pseudo-sphere is represented. The latter surface is obtained from Dini’s surface by taking the stationary limit of the kink solution. Figures 1c and 1d correspond to the pseudo-spherical surfaces associated to a stationary breather and to a two-kink solution, respectively.

argued and will be explicitly shown in the next section, are independent of the deformation parameter τ , being it re absorbable through a local change of coordinates.

The corresponding soliton solutions, described in section 5, are instead affected by the $T\bar{T}$ in a highly non-trivial way. For instance, they generally possess critical values in τ corresponding to shock-wave phenomena, *i.e.* branching of the solutions. Examples of shock-wave phenomena and square root-type transitions in the classical energy – similar to the Hagedorn transition at quantum level – will be discussed in sections 5 and 6 for specific solutions of the deformed sine-Gordon model.

3.1 From the deformed to the undeformed model through a local change of coordinates

Thus we have inferred that there must exist a coordinate system $\mathbf{w} = (w^1(\mathbf{z}), w^2(\mathbf{z})) = (w(\mathbf{z}), \bar{w}(\mathbf{z}))$ in which the matrices $g_{\mu\nu}^{\text{T}\bar{\text{T}}}$ and $d_{\mu\nu}^{\text{T}\bar{\text{T}}}$ assume the same form as $g_{\mu\nu}^{\text{sG}}$ and $d_{\mu\nu}^{\text{sG}}$, respectively. In formulae

$$g_{\mu\nu}^{\text{sG}} dw^\mu dw^\nu = g_{\mu\nu}^{\text{T}\bar{\text{T}}} dz^\mu dz^\nu \implies g_{\mu\nu}^{\text{sG}} \frac{dw^\mu}{dz^\rho} \frac{dw^\nu}{dz^\sigma} = g_{\rho\sigma}^{\text{T}\bar{\text{T}}} , \quad (3.15)$$

$$d_{\mu\nu}^{\text{sG}} dw^\mu dw^\nu = d_{\mu\nu}^{\text{T}\bar{\text{T}}} dz^\mu dz^\nu \implies d_{\mu\nu}^{\text{sG}} \frac{dw^\mu}{dz^\rho} \frac{dw^\nu}{dz^\sigma} = d_{\rho\sigma}^{\text{T}\bar{\text{T}}} . \quad (3.16)$$

It is now a matter of simple algebraic manipulations to obtain the following equations for the new coordinates

$$\partial w = \frac{(S+1)^2}{4S(1-\tau V)} , \quad \bar{\partial} \bar{w} = \frac{(S+1)^2}{4S(1-\tau V)} , \quad (3.17)$$

$$\bar{\partial} w = \frac{\tau}{S} (\bar{\partial} \phi)^2 , \quad \partial \bar{w} = \frac{\tau}{S} (\partial \phi)^2 . \quad (3.18)$$

Let us now use the latter relations to find the partial derivatives of the field ϕ in the coordinates $\mathbf{w}$:

$$\begin{pmatrix} \partial \phi \\ \bar{\partial} \phi \end{pmatrix} = \mathcal{J} \begin{pmatrix} \partial \phi / \partial w \\ \partial \phi / \partial \bar{w} \end{pmatrix} , \quad \mathcal{J} = \begin{pmatrix} \partial w & \partial \bar{w} \\ \bar{\partial} w & \bar{\partial} \bar{w} \end{pmatrix} . \quad (3.19)$$

The result is

$$\partial \phi = \frac{1}{1-\tau(\mathcal{K}+V)} \frac{\partial \phi}{\partial w} , \quad \bar{\partial} \phi = \frac{1}{1-\tau(\mathcal{K}+V)} \frac{\partial \phi}{\partial \bar{w}} , \quad (3.20)$$

where we have defined the following function

$$\mathcal{K} = \frac{\partial \phi(\mathbf{w})}{\partial w} \frac{\partial \phi(\mathbf{w})}{\partial \bar{w}} . \quad (3.21)$$

With the help of (3.20), we can now find the expression for S in the coordinates $\mathbf{w}$

$$S = \sqrt{1 + 4\tau(1-\tau V) \partial \phi \bar{\partial} \phi} = \frac{1 + \tau(\mathcal{K} - V)}{1 - \tau(\mathcal{K} + V)} . \quad (3.22)$$

We can then write the Jacobian matrix $\mathcal{J}$ and its inverse $\mathcal{J}^{-1}$ in terms of $\mathbf{w}$ as

$$\begin{aligned} \mathcal{J} &= \begin{pmatrix} \partial w & \partial \bar{w} \\ \bar{\partial} w & \bar{\partial} \bar{w} \end{pmatrix} = \frac{1}{(1-\tau V)^2 - \tau^2 \mathcal{K}^2} \begin{pmatrix} 1-\tau V & \tau \left(\frac{\partial \phi}{\partial w} \right)^2 \\ \tau \left(\frac{\partial \phi}{\partial \bar{w}} \right)^2 & 1-\tau V \end{pmatrix} , \\ \mathcal{J}^{-1} &= \begin{pmatrix} \partial_w z & \partial_w \bar{z} \\ \partial_{\bar{w}} z & \partial_{\bar{w}} \bar{z} \end{pmatrix} = \begin{pmatrix} 1-\tau V & -\tau \left(\frac{\partial \phi}{\partial w} \right)^2 \\ -\tau \left(\frac{\partial \phi}{\partial \bar{w}} \right)^2 & 1-\tau V \end{pmatrix} . \end{aligned} \quad (3.23)$$

This results allows us to express the partial derivatives of any function $f(\mathbf{z})$ as partial derivatives with respect to the new coordinates

$$\begin{pmatrix} \partial f \\ \bar{\partial} f \end{pmatrix} = \mathcal{J} \begin{pmatrix} \partial f / \partial w \\ \partial f / \partial \bar{w} \end{pmatrix}, \quad (3.24)$$

and we can then apply all the above formulae to the equation (3.1), obtaining

$$\partial \left(\frac{\bar{\partial} \phi}{S} \right) + \bar{\partial} \left(\frac{\partial \phi}{S} \right) = 2 \frac{\frac{\partial}{\partial w} \frac{\partial}{\partial \bar{w}} \phi}{(1 + \tau(\mathcal{K} - V))^2} - 2\tau \frac{V'}{(1 - \tau V)^2 - \tau^2 \mathcal{K}^2} \frac{\mathcal{K}}{(1 + \tau(\mathcal{K} - V))}, \quad (3.25)$$

$$\frac{V'}{4S} \left(\frac{S+1}{1 - \tau V} \right)^2 = \frac{V'}{(1 - \tau V)^2 - \tau^2 \mathcal{K}^2}. \quad (3.26)$$

The equality of (3.25) and (3.26) yields then

$$\frac{2 \frac{\partial}{\partial w} \frac{\partial}{\partial \bar{w}} \phi - V'}{(1 + \tau(\mathcal{K} - V))^2} = 0. \quad (3.27)$$

4 A geometric map for N-boson fields and arbitrary potential

We have seen, in the preceding section, how the $\text{T}\bar{\text{T}}$ deformation of the sine-Gordon model can be interpreted as a field-dependent coordinate transformation. We arrived at this interesting conclusion by exploiting the relation existing amongst ZCR of soliton equations and the classical geometry of surfaces embedded in flat space. Although this connection was pivotal in guiding us to the map (3.17, 3.18), from that point on we did not make explicit mention to the form of the potential. In other words, we can consider all formulae from (3.17) to (3.27) to be valid for any 2-dimensional single scalar system.

More generally, the results (3.23, 3.27) admit a straightforward generalisation to the case of N bosonic fields ϕ_i , ($i = 1, \dots, N$) interacting with a generic derivative-independent potential $V(\phi_i)$

$$\mathcal{L}_N^{(\tau)}(\mathbf{z}) = \frac{V}{1 - \tau V} + \frac{-1 + \sqrt{1 + 4\bar{\tau} \left(\mathcal{L}_{\text{free}}^{(0)} - \bar{\tau} \mathcal{B} \right)}}{2\bar{\tau}}, \quad (4.1)$$

$$\mathcal{L}_{\text{free}}^{(0)} = \sum_{i=1}^N \partial \phi_i \bar{\partial} \phi_i, \quad \mathcal{B} = |\partial \vec{\phi} \times \bar{\partial} \vec{\phi}|^2, \quad (4.2)$$

with $\bar{\tau} = \tau(1 - \tau V)$, arising as a $\text{T}\bar{\text{T}}$ -deformation of [26, 27]

$$\mathcal{L}_N^{(0)} = \sum_{i=1}^N \partial \phi_i \bar{\partial} \phi_i + V(\phi_i). \quad (4.3)$$

The generalization of (3.23) to the N -boson case is

$$\begin{aligned}\mathcal{J}_N &= \begin{pmatrix} \frac{\partial w}{\partial w} & \frac{\partial \bar{w}}{\partial \bar{w}} \end{pmatrix} = \frac{1}{(1 - \tau V)^2 - \tau^2 (\mathcal{K}_N)^2} \begin{pmatrix} 1 - \tau V & \tau \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial w} \right)^2 \\ \tau \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial \bar{w}} \right)^2 & 1 - \tau V \end{pmatrix}, \\ \mathcal{J}_N^{-1} &= \begin{pmatrix} \frac{\partial w z}{\partial \bar{w} z} & \frac{\partial w \bar{z}}{\partial \bar{w} \bar{z}} \end{pmatrix} = \begin{pmatrix} 1 - \tau V & -\tau \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial w} \right)^2 \\ -\tau \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial \bar{w}} \right)^2 & 1 - \tau V \end{pmatrix},\end{aligned}\quad (4.4)$$

with $(\mathcal{K}_N)^2 = \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial w} \right)^2 \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial \bar{w}} \right)^2$. In fact we have verified that the deformed EoMs resulting from (4.1) are mapped by (4.4) into the undeformed EoMs associated to $\mathcal{L}_N^{(0)}$.

It is instructive to translate (4.4) in Euclidean coordinates. Considering

$$\begin{aligned}\left(\frac{\partial}{\partial w} + \frac{\partial}{\partial \bar{w}} \right) (z + \bar{z}) &= 2 + \tau \left(-\sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial \bar{w}} \right)^2 - \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial w} \right)^2 - 2V \right), \\ \left(\frac{\partial}{\partial w} - \frac{\partial}{\partial \bar{w}} \right) (z - \bar{z}) &= 2 - \tau \left(\sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial \bar{w}} \right)^2 - \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial w} \right)^2 + 2V \right), \\ \left(\frac{\partial}{\partial w} + \frac{\partial}{\partial \bar{w}} \right) (z - \bar{z}) &= \tau \left(\sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial \bar{w}} \right)^2 - \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial w} \right)^2 \right), \\ \left(\frac{\partial}{\partial w} - \frac{\partial}{\partial \bar{w}} \right) (z + \bar{z}) &= -\tau \left(\sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial \bar{w}} \right)^2 - \sum_{i=1}^N \left(\frac{\partial \phi_i}{\partial w} \right)^2 \right),\end{aligned}\quad (4.5)$$

and moving to Euclidean coordinates both in the $\mathbf{z}$ and in the $\mathbf{w}$ frames

$$\begin{cases} z = x^1 + \mathbf{i} x^2 \\ \bar{z} = x^1 - \mathbf{i} x^2 \end{cases}, \quad \begin{cases} w = y^1 + \mathbf{i} y^2 \\ \bar{w} = y^1 - \mathbf{i} y^2 \end{cases} \rightarrow \begin{cases} \frac{\partial}{\partial w} + \frac{\partial}{\partial \bar{w}} = \frac{\partial}{\partial y^1} \\ \frac{\partial}{\partial w} - \frac{\partial}{\partial \bar{w}} = -\mathbf{i} \frac{\partial}{\partial y^2} \end{cases}\quad (4.6)$$

we find

$$\frac{\partial x^1}{\partial y^1} = 1 + \tau T_2^2(\mathbf{y}), \quad \frac{\partial x^2}{\partial y^2} = 1 + \tau T_1^1(\mathbf{y}), \quad \frac{\partial x^1}{\partial y^2} = \frac{\partial x^2}{\partial y^1} = -\tau T_2^1(\mathbf{y}), \quad (4.7)$$

where $T_\nu^\mu(\mathbf{y})$ is the stress energy tensor of the undeformed theory, $T = T^{(0)}$. Expressions (4.7) can be more compactly rewritten as

$$\frac{\partial x^\mu}{\partial y^\nu} = \delta^\mu_\nu + \tau \tilde{T}_\nu^\mu(\mathbf{y}), \quad \tilde{T}_\nu^\mu(\mathbf{y}) = -\epsilon^\mu_\rho \epsilon^\sigma_\nu T^\rho_\sigma(\mathbf{y}). \quad (4.8)$$

From (4.8) the inverse Jacobian in Euclidean coordinates reads

$$\mathcal{J}_N^{-1} = \begin{pmatrix} \frac{\partial x^1}{\partial y^1} & \frac{\partial x^2}{\partial y^1} \\ \frac{\partial x^1}{\partial y^2} & \frac{\partial x^2}{\partial y^2} \end{pmatrix} = \begin{pmatrix} 1 + \tau T_2^2 & -\tau T_2^1 \\ -\tau T_1^2 & 1 + \tau T_1^1 \end{pmatrix}, \quad (4.9)$$

and thus the metric, in the set of coordinates $\mathbf{y}$, is

$$\mathbf{g}'_{\mu\nu} = \frac{\partial x^\rho}{\partial y^\mu} \frac{\partial x^\sigma}{\partial y^\nu} \mathbf{g}_{\rho\sigma} = \delta_{\mu\nu} - \tau \epsilon_{\mu\rho} \epsilon^\sigma{}_\nu (2T + \tau T^2)^\rho{}_\sigma , \quad (4.10)$$

where we used the fact that $\mathbf{g}_{\rho\sigma} = \delta_{\rho\sigma}$. Translating the first expression of (4.4) in $\mathbf{z}$ coordinates and then moving to Euclidean coordinates, one obtains the inverse relation of (4.8)

$$\frac{\partial y^\mu}{\partial x^\nu} = \delta^\mu{}_\nu + \tau (\tilde{T}^{(\tau)})^\mu{}_\nu(\mathbf{x}) , \quad (\tilde{T}^{(\tau)})^\mu{}_\nu(\mathbf{x}) = -\epsilon^\mu{}_\rho \epsilon^\sigma{}_\nu (T^{(\tau)})^\rho{}_\sigma(\mathbf{x}) , \quad (4.11)$$

where $(T^{(\tau)})^\mu{}_\nu(\mathbf{x})$ is the stress energy tensor of the deformed theory.

Finally let us conclude this section with a couple of remarks:

- Consider the transformation of the Lagrangian⁷ (4.1) under the on-shell map (4.4)

$$\mathcal{L}_N^{(\tau)}(\mathbf{z}(\mathbf{w})) = \frac{\mathcal{L}_N^{(0)}(\mathbf{w}) + \tau \left((\mathcal{K}_N)^2 - V^2 \right)}{1 - 2\tau V - \tau^2 \left((\mathcal{K}_N)^2 - V^2 \right)} . \quad (4.13)$$

Using the latter expression together with

$$\text{Det}(\mathcal{J}_N^{-1}) = \text{Det}(\mathcal{J}_N)^{-1} = 1 - 2\tau V - \tau^2 \left((\mathcal{K}_N)^2 - V^2 \right) , \quad (4.14)$$

we find that the action transforms as

$$\begin{aligned} \mathcal{A}[\phi] &= \int dz d\bar{z} \mathcal{L}_N^{(\tau)}(\mathbf{z}) = \int dw d\bar{w} |\text{Det}(\mathcal{J}_N^{-1})| \mathcal{L}_N^{(\tau)}(\mathbf{z}(\mathbf{w})) \\ &= \int dw d\bar{w} \left(\mathcal{L}_N^{(0)}(\mathbf{w}) + \tau \text{T}\bar{\text{T}}^{(0)}(\mathbf{w}) \right) , \end{aligned} \quad (4.15)$$

where $\text{T}\bar{\text{T}}^{(0)}(\mathbf{w}) = (\mathcal{K}_N)^2 - V^2$. Thus, we conclude that the action is not invariant under the change of variables. This is not totally surprising since the map (4.4) is on-shell, however it is remarkable that the (bare) perturbing field can be so easily identified once the change of variables is performed. Again, our result matches with [18], where the $\text{T}\bar{\text{T}}^{(0)}$ term emerges as a JT gravity contribution to the action.

- Notice that the EoMs associated to (4.1) for a generic potential V are invariant under the transformation⁸

$$\mathbf{z} \rightarrow \gamma \mathbf{z} , \quad \tau \rightarrow \gamma \tau , \quad V \rightarrow V - c , \quad (4.16)$$

⁷In the $N = 1$ case, the transformed Lagrangian takes an even simpler expression

$$\mathcal{L}_1^{(\tau)}(\mathbf{z}(\mathbf{w})) = \frac{\mathcal{L}_1^{(0)}(\mathbf{w})}{1 - \tau \mathcal{L}_1^{(0)}(\mathbf{w})} . \quad (4.12)$$

⁸We thank Sergei Dubovsky for questioning us about the possible existence of such symmetry of the EoMs.

with c constant and $\gamma = 1/(1 - \tau c)$, which corresponds to the following change of variables at the level of the solutions

$$\phi_i^{(\tau)}(\mathbf{z})|_V = \phi_i^{(\gamma\tau)}(\gamma\mathbf{z})|_{V-c}, \quad i = 1, \dots, N, \quad (4.17)$$

where the notation $\phi_i^{(\tau)}|_V$ means that $\phi_i^{(\tau)}$ is solution to the deformed EoMs with potential V .

5 T $\bar{T}$ -deformed soliton solutions in the sine-Gordon model

In this section we show how to compute T $\bar{T}$ -deformed solutions of the sG model by explicitly evaluating the change of variables on specific solutions $\phi(\mathbf{w})$ of the undeformed theory. The idea is to solve the following sets of differential equations derived from the inverse Jacobian (3.23)

$$\begin{cases} \frac{\partial z(\mathbf{w})}{\partial w} = 1 - \tau V(\phi(\mathbf{w})) \\ \frac{\partial z(\mathbf{w})}{\partial \bar{w}} = -\tau \left(\frac{\partial \phi(\mathbf{w})}{\partial \bar{w}} \right)^2 \end{cases}, \quad \begin{cases} \frac{\partial \bar{z}(\mathbf{w})}{\partial w} = -\tau \left(\frac{\partial \phi(\mathbf{w})}{\partial w} \right)^2 \\ \frac{\partial \bar{z}(\mathbf{w})}{\partial \bar{w}} = 1 - \tau V(\phi(\mathbf{w})) \end{cases}, \quad (5.1)$$

for $z(\mathbf{w})$ and $\bar{z}(\mathbf{w})$. Then from the inverse map, *i.e.* $\mathbf{w}(\mathbf{z})$, we evaluate the expression of the deformed solution as

$$\phi^{(\tau)}(\mathbf{z}) = \phi^{(0)}(\mathbf{w}(\mathbf{z})). \quad (5.2)$$

In the following we will deal only with some of the simplest solutions of the sG model. In principle our approach applies for all the solutions, although we could not find an explicit result for the integrated map in the cases involving more than two solitons.

For sake of clarity, the computations shown in the following sections will be carried on in light cone coordinates, *i.e.* $(z, \bar{z})$ and $(w, \bar{w})$, however the plots will be displayed using space and time coordinates $(x, t) = (x^1, x^2)$.

5.1 The one-kink solution

Let us start with the one-kink solution moving with velocity v

$$\phi_{1\text{-kink}}^{(0)}(\mathbf{w}) = 4 \arctan \left(e^{\frac{m}{\beta} (aw + \frac{1}{a}\bar{w})} \right), \quad a = \sqrt{\frac{1-v}{1+v}}. \quad (5.3)$$

With the identification $\phi(\mathbf{w}) = \phi_{1\text{-kink}}^{(0)}(\mathbf{w})$, equations (5.1) can be easily integrated yielding

$$\begin{aligned} z(\mathbf{w}) &= w - 4\tau \frac{m}{a\beta} \tanh \left[\frac{m}{\beta} \left(aw + \frac{1}{a}\bar{w} \right) \right], \\ \bar{z}(\mathbf{w}) &= \bar{w} - 4\tau \frac{am}{\beta} \tanh \left[\frac{m}{\beta} \left(aw + \frac{1}{a}\bar{w} \right) \right], \end{aligned} \quad (5.4)$$

where the constants of integration are fixed consistently with the $\tau = 0$ case. Notice that from (5.3) we have

$$\frac{m}{\beta} \left(aw + \frac{1}{a}\bar{w} \right) = \ln \left(\tan \left(\frac{\phi_{1\text{-kink}}^{(0)}(\mathbf{w})}{4} \right) \right), \quad (5.5)$$




Figure 2. The $T\bar{T}$ -deformed moving one-kink solution ($m = \beta = 1$, $a = 2$), for different values of the perturbation parameter τ . Figure 2b represents the undeformed solution. Figure 2a corresponds to $\tau = -1/4$, while Figures 2c and 2d correspond to $\tau = 1/8$ and $\tau = 1/3$, respectively. Notice that at $\tau = 1/3$ a shock-wave singularity occurs.

and thus expressions (5.4) become

$$z(\mathbf{w}) = w + 4\tau \frac{m}{a\beta} \cos\left(\frac{\phi_{1\text{-kink}}^{(0)}(\mathbf{w})}{2}\right), \quad \bar{z}(\mathbf{w}) = \bar{w} + 4\tau \frac{a m}{\beta} \cos\left(\frac{\phi_{1\text{-kink}}^{(0)}(\mathbf{w})}{2}\right), \quad (5.6)$$

which are easily inverted as

$$\begin{aligned} w(\mathbf{z}) &= z - 4\tau \frac{m}{a\beta} \cos\left(\frac{\phi_{1\text{-kink}}^{(0)}(\mathbf{w}(\mathbf{z}))}{2}\right) = z - 4\tau \frac{m}{a\beta} \cos\left(\frac{\phi_{1\text{-kink}}^{(\tau)}(\mathbf{z})}{2}\right), \\ \bar{w}(\mathbf{z}) &= \bar{z} - 4\tau \frac{am}{\beta} \cos\left(\frac{\phi_{1\text{-kink}}^{(0)}(\mathbf{w}(\mathbf{z}))}{2}\right) = \bar{z} - 4\tau \frac{am}{\beta} \cos\left(\frac{\phi_{1\text{-kink}}^{(\tau)}(\mathbf{z})}{2}\right). \end{aligned} \quad (5.7)$$

Finally, plugging (5.7) into (5.3) we find

$$\frac{m}{\beta} \left(az + \frac{1}{a} \bar{z} \right) = 8\tau \frac{m^2}{\beta^2} \cos\left(\frac{\phi_{1\text{-kink}}^{(\tau)}(\mathbf{z})}{2}\right) + \ln\left(\tan\left(\frac{\phi_{1\text{-kink}}^{(\tau)}(\mathbf{z})}{4}\right)\right), \quad (5.8)$$

which is exactly the deformed one-kink solution found in [27]. In Figure 2 the solution is represented for different values of τ . Notice that for negative values of τ (Figure 2a) the solution stretches w.r.t the undeformed one (Figure 2b), while for positive values of τ (Figures 2c and 2d) it bends and becomes multi-valued. In particular $\tau = 1/8$ (Figure 2c) is the delimiting value corresponding to a shock wave singularity.

5.2 The two-kink solution

Consider now the solution which describes the scattering between two kinks with velocities v_1 and v_2

$$\phi_{2\text{-kink}}^{(0)}(\mathbf{w}) = 4 \arctan \left(\frac{a_1 + a_2}{a_2 - a_1} \frac{e^{\frac{m}{\beta}(a_1 w + \frac{1}{a_1} \bar{w} + k_1)} - e^{\frac{m}{\beta}(a_2 w + \frac{1}{a_2} \bar{w} + k_2)}}{1 + e^{\frac{m}{\beta}(a_1 w + \frac{1}{a_1} \bar{w} + k_1)} e^{\frac{m}{\beta}(a_2 w + \frac{1}{a_2} \bar{w} + k_2)}} \right), \quad (5.9)$$

where again $a_i = \sqrt{\frac{1-v_i}{1+v_i}}$, $i = 1, 2$, and k_i , $i = 1, 2$, are constant phases. Compared to the one-kink case, this time the sets of differential equations (5.1) are more complicated to integrate. It is useful to parametrize the solutions $z(\mathbf{w})$ and $\bar{z}(\mathbf{w})$ of (5.1) in terms of the combinations

$$u_i(\mathbf{w}) = \frac{m}{\beta} \left(a_i w + \frac{1}{a_i} \bar{w} + k_i \right), \quad i = 1, 2. \quad (5.10)$$

Performing the change of variables $\mathbf{u} = (u_1(\mathbf{w}), u_2(\mathbf{w}))$

$$\begin{cases} \frac{\partial z}{\partial u_1} = \frac{\beta}{m} \frac{a_1 \left(\frac{\partial z}{\partial w} - a_2^2 \frac{\partial z}{\partial \bar{w}} \right)}{a_1^2 - a_2^2} \\ \frac{\partial z}{\partial u_2} = -\frac{\beta}{m} \frac{a_2 \left(\frac{\partial z}{\partial w} - a_1^2 \frac{\partial z}{\partial \bar{w}} \right)}{a_1^2 - a_2^2} \end{cases}, \quad \begin{cases} \frac{\partial \bar{z}}{\partial u_1} = \frac{\beta}{m} \frac{a_1 \left(\frac{\partial \bar{z}}{\partial w} - a_2^2 \frac{\partial \bar{z}}{\partial \bar{w}} \right)}{a_1^2 - a_2^2} \\ \frac{\partial \bar{z}}{\partial u_2} = -\frac{\beta}{m} \frac{a_2 \left(\frac{\partial \bar{z}}{\partial w} - a_1^2 \frac{\partial \bar{z}}{\partial \bar{w}} \right)}{a_1^2 - a_2^2} \end{cases}, \quad (5.11)$$

and plugging (5.1) into (5.11) with the identification $\phi(\mathbf{w}) \equiv \phi_{2\text{-kink}}^{(0)}(\mathbf{w})$, we obtain two sets of differential equations which can be solved for $\mathbf{z}(\mathbf{u})$, giving

$$\begin{aligned} z(\mathbf{u}) &= \frac{\beta}{m} \frac{a_1 u_1 - a_2 u_2}{a_1^2 - a_2^2} - 4\tau \frac{m}{\beta} \frac{(a_1^2 - a_2^2)(a_1 \tanh u_2 - a_2 \tanh u_1)}{a_1 a_2 (a_1^2 + a_2^2 - 2a_1 a_2 (\text{sech } u_1 \text{sech } u_2 + \tanh u_1 \tanh u_2))}, \\ \bar{z}(\mathbf{u}) &= \frac{\beta}{m} \frac{a_1 a_2 (a_1 u_2 - a_2 u_1)}{a_1^2 - a_2^2} - 4\tau \frac{m}{\beta} \frac{(a_1^2 - a_2^2)(a_1 \tanh u_1 - a_2 \tanh u_2)}{a_1^2 + a_2^2 - 2a_1 a_2 (\text{sech } u_1 \text{sech } u_2 + \tanh u_1 \tanh u_2)}. \end{aligned} \quad (5.12)$$

As in the previous section, the constants of integration in (5.12) are fixed by imposing the consistency with the $\tau = 0$ case. In order to find the deformed two-kink solution $\phi_{2\text{-kink}}^{(\tau)}(\mathbf{z}) = \phi_{2\text{-kink}}^{(0)}(\mathbf{u}(\mathbf{z}))$, we should solve (5.12) for $\mathbf{u}(\mathbf{z})$. Since this is analytically very complicated, we resort to numerical inversion. In Figure 3 the deformed solution $\phi_{2\text{-kink}}^{(\tau)}(\mathbf{z})$ is reported for different values of τ . The picture is quite similar to the one-kink case. In fact, for negative values of τ (Figure 3a) the solution stretches w.r.t. the undeformed one (Figure 3b), while for positive values of τ (Figures 3c and 3d) it bends and again it becomes multi-valued. Unlike the one-kink case, here it is not possible to find analytically the delimiting value of τ corresponding to the shock singularity.




Figure 3. The $T\bar{T}$ -deformed two-kink solution ($m = \beta = 1$, $a_1 = 2$, $a_2 = 3$), for different values of the perturbation parameter τ . Figure 3b represents the undeformed solution. Figure 3a corresponds to $\tau = -1/4$, while Figures 3c and 3d correspond to τ , *i.e.* $\tau = 1/10$ and $\tau = 1/6$, respectively.

5.3 The breather

Another interesting solution is the breather with envelope speed $v = 0$

$$\phi_{\text{breather}}^{(0)}(\mathbf{w}) = 4 \arctan \left(\tan \psi \frac{\sin \left(-\frac{m}{\beta}(w - \bar{w}) \cos \psi + \bar{k} \right)}{\cosh \left(\frac{m}{\beta}(w + \bar{w}) \sin \psi + k \right)} \right), \quad (5.13)$$

where ψ is a parameter related to the period T of one full oscillation via $T = \frac{2\pi}{\cos \psi}$ and $k, \bar{k}$ are constant phases. In analogy with the two-kink case, it is useful to use the same strategy and parametrize the solutions $\mathbf{z}(\mathbf{w})$ of (5.1) in terms of

$$u(\mathbf{w}) = \frac{m}{\beta}(w + \bar{w}) \sin \psi + k, \quad \bar{u}(\mathbf{w}) = -\frac{m}{\beta}(w - \bar{w}) \cos \psi + \bar{k}. \quad (5.14)$$




Figure 4. The $T\bar{T}$ -deformed stationary breather solution with envelope speed $v = 0$ ($m = \beta = 1$, $\psi = \frac{2}{5}\pi$), for different values of the perturbation parameter τ . Figure 4b represents the undeformed solution, Figure 4a corresponds to $\tau = -1/2$, while Figures 4c and 4d correspond to $\tau = 1/10$ and $\tau = 1/5$, respectively.

Performing the change of variables $\mathbf{u}(\mathbf{w}) = (u(\mathbf{w}), \bar{u}(\mathbf{w}))$, one finds

$$\begin{cases} \frac{\partial z}{\partial u} = \frac{\beta}{m} \frac{1}{2 \sin \psi} \left(\frac{\partial z}{\partial w} + \frac{\partial z}{\partial \bar{w}} \right) \\ \frac{\partial z}{\partial \bar{u}} = \frac{\beta}{m} \frac{1}{2 \cos \psi} \left(-\frac{\partial z}{\partial w} + \frac{\partial z}{\partial \bar{w}} \right) \end{cases}, \quad \begin{cases} \frac{\partial \bar{z}}{\partial u} = \frac{\beta}{m} \frac{1}{2 \sin \psi} \left(\frac{\partial \bar{z}}{\partial w} + \frac{\partial \bar{z}}{\partial \bar{w}} \right) \\ \frac{\partial \bar{z}}{\partial \bar{u}} = \frac{\beta}{m} \frac{1}{2 \cos \psi} \left(-\frac{\partial \bar{z}}{\partial w} + \frac{\partial \bar{z}}{\partial \bar{w}} \right) \end{cases}, \quad (5.15)$$

and again plugging (5.1) into (5.15) with the identification $\phi(\mathbf{w}) \equiv \phi_{\text{breather}}^{(0)}(\mathbf{w})$, one gets two sets of differential equations which can be solved for $\mathbf{z}(\mathbf{u})$ giving

$$\begin{aligned} z(\mathbf{u}) &= \frac{\beta}{m} \left(\frac{u}{2 \sin \psi} - \frac{\bar{u}}{2 \cos \psi} \right) - 8\tau \frac{m}{\beta} \sin \psi \frac{\cos \bar{u}}{\cosh u} \frac{\sec \bar{u} \sinh u + \operatorname{sech} u \sin \bar{u} \tan \psi}{1 + (\tan \psi \sin \bar{u} \operatorname{sech} u)^2}, \\ \bar{z}(\mathbf{u}) &= \frac{\beta}{m} \left(\frac{u}{2 \sin \psi} + \frac{\bar{u}}{2 \cos \psi} \right) - 8\tau \frac{m}{\beta} \sin \psi \frac{\cos \bar{u}}{\cosh u} \frac{\sec \bar{u} \sinh u - \operatorname{sech} u \sin \bar{u} \tan \psi}{1 + (\tan \psi \sin \bar{u} \operatorname{sech} u)^2}. \end{aligned} \quad (5.16)$$

As for the two-kink example, the constants of integration in (5.16) are fixed according to the $\tau = 0$ case, and again the solution $\mathbf{u}(\mathbf{z})$ to (5.16) is computed numerically. The deformed solution $\phi_{\text{breather}}^{(\tau)}(\mathbf{z})$ is displayed in Figure 4 for different values of τ . The result is similar to

the previous cases: the solution stretches for negative values of τ (Figure 4a) and it bends for positive values of τ (Figure 4c and 4d) w.r.t. the undeformed one (Figure 4b). However, notice that in this case the shock phenomenon occurs in both positive and negative directions of τ , and consequently the solution becomes multi-valued (Figures 4a and 4d) for $|\tau|$ sufficiently large.

6 The shock-wave phenomenon and the Hagedorn-type transition

In this section we will discuss the emergence of critical phenomena in the classical solutions, *i.e.* the shock-wave singularity and the square root-type transition, and comment on the relations among them. We will use as a guide example the stationary T $\bar{\text{T}}$ -deformed elliptic solution of the sG model derived in [27], where we set $\rho = 1/\kappa > 0$ and $m = \beta = 1$,

$$x = \frac{1}{2\sqrt{\rho}} \left[(\rho + 4\tau) \mathbf{F} \left(\frac{\phi(x)}{2} \middle| -\rho \right) - 8\tau \mathbf{E} \left(\frac{\phi(x)}{2} \middle| -\rho \right) \right], \quad (6.1)$$

defined on a cylinder of radius R fixed. Due to the following properties of the elliptic functions

$$\begin{aligned} \mathbf{F}(z + n\pi | \gamma) &= \mathbf{F}(z | \gamma) + 2n \mathbf{K}(\gamma), \\ \mathbf{E}(z + n\pi | \gamma) &= \mathbf{E}(z | \gamma) + 2n \mathbf{E}(\gamma), \quad z, \gamma \in \mathbb{C}, \quad n \in \mathbb{Z}, \end{aligned} \quad (6.2)$$

the solution $\phi(x)$ can be interpreted as a stationary 1-kink with twisted boundary conditions

$$\phi(x + R) = \phi(x) + 2\pi, \quad (6.3)$$

where the radius R is

$$R = \frac{1}{\sqrt{\rho}} \left((\rho + 4\tau) \mathbf{K}(-\rho) - 8\tau \mathbf{E}(-\rho) \right). \quad (6.4)$$

We stress that R is kept fixed while $\rho = \rho(\tau, R)$ is considered as a function of τ and R , defined implicitly through (6.4). Differentiating both sides of (6.4) w.r.t. τ and R and solving for $\partial_\tau \rho$ and $\partial_R \rho$ one finds

$$\partial_\tau \rho = -\frac{8\rho(1+\rho)(2\mathbf{E}(-\rho) - \mathbf{K}(-\rho))}{(\rho + 4\tau)\mathbf{E}(-\rho)}, \quad \partial_R \rho = \frac{2\rho^{3/2}(1+\rho)}{(\rho + 4\tau)\mathbf{E}(-\rho)}. \quad (6.5)$$

We shall now compute the energy on the cylinder. The components of the Hilbert stress-energy tensor $T_{\mu\nu}^{(\tau)}$ are

$$T_{22}^{(\tau)} \equiv \mathcal{H}^{(\tau)} = \frac{V}{1 - \tau V} + \frac{1 + \tau(1 - \tau V)\phi_x^2 - S}{2S\tau(1 - \tau V)} = \frac{2(2 + \rho V)}{\rho(1 - 2\tau V) - 4\tau}, \quad (6.6)$$

$$T_{12}^{(\tau)} = T_{21}^{(\tau)} \equiv \mathcal{P}^{(\tau)} = -\frac{\phi_t \phi_x}{2S} = 0, \quad (6.7)$$

$$T_{11}^{(\tau)} = -\frac{V}{1 - \tau V} - \frac{1 - \tau(1 - \tau V)\phi_t^2 - S}{2S\tau(1 - \tau V)} = \frac{4}{\rho + 4\tau}, \quad (6.8)$$

where we used the following expressions for ϕ_t and ϕ_x derived from (6.1)

$$\phi_t = 0, \quad \phi_x = \frac{2\sqrt{\rho}\sqrt{4+\rho V}}{\rho(1-2\tau V)-4\tau}, \quad (6.9)$$

and

$$S = \sqrt{1 + \tau(1 - \tau V)(\phi_x^2 - \phi_t^2)} = \frac{\rho + 4\tau}{\rho(1 - 2\tau V) - 4\tau}. \quad (6.10)$$

Notice that the apparent pole singularity at $\tau = 1/V$ in $T_{11}^{(\tau)}$ and $T_{22}^{(\tau)}$ disappears once (6.9) is used in (6.6) and (6.8). Finally the energy and momentum at finite volume R are

$$E^{(\tau)} = \int_{x_0}^{x_0+R} \mathcal{H}^{(\tau)}(x) dx = \int_{\phi(x_0)=0}^{\phi(x_0+R)=2\pi} \frac{\mathcal{H}^{(\tau)}(\phi)}{\phi_x} d\phi = \frac{4}{\sqrt{\rho}} (2\mathbf{E}(-\rho) - \mathbf{K}(-\rho)), \quad (6.11)$$

$$P^{(\tau)} = \int_{x_0}^{x_0+R} \mathcal{P}^{(\tau)}(x) dx = \int_{\phi(x_0)=0}^{\phi(x_0+R)=2\pi} \frac{\mathcal{P}^{(\tau)}(\phi)}{\phi_x} d\phi = 0, \quad (6.12)$$

$$K^{(\tau)} = \int_{x_0}^{x_0+R} T_{11}^{(\tau)}(x) dx = \int_{\phi(x_0)=0}^{\phi(x_0+R)=2\pi} \frac{T_{11}^{(\tau)}(\phi)}{\phi_x} d\phi = \frac{4R}{\rho + 4\tau}, \quad (6.13)$$

where $x_0 = 0 \pmod{R}$. From (6.5), (6.11) and (6.13) one can prove that the energy fulfils the Burgers equation (1.1) with $P_n = 0$

$$\partial_\tau E^{(\tau)} = \frac{1}{2} E^{(\tau)} \partial_R E^{(\tau)} = -\frac{1}{R} \det \left(\int_{x_0}^{x_0+R} T_{\mu\nu}^{(\tau)}(x) dx \right) = - \int_{x_0}^{x_0+R} \det \left(T_{\mu\nu}^{(\tau)}(x) \right) dx, \quad (6.14)$$

where the last equality in (6.14) shows the factorization property of the $T\bar{T}$ operator at the classical level. Since the energy $E^{(\tau)}$ fulfils a Burgers equation, it is expected to have a square root-type singularity.⁹ The critical radius $R_c(\tau)$ corresponds to a value of R such that the first derivative of $E^{(\tau)}(R)$ w.r.t. R diverges. One easily checks that

$$\partial_R E^{(\tau)} = -\frac{4}{\rho + 4\tau}, \quad (6.16)$$

thus the first derivative is divergent at the radius $R_c(\tau)$ defined through the equation

$$\rho(\tau, R_c(\tau)) = -4\tau. \quad (6.17)$$

According to (6.4) and (6.11), the critical radius and the corresponding energy turn out to be

$$R_c(\tau) = 4\sqrt{-\tau} \mathbf{E}(4\tau), \quad E_c^{(\tau)} \equiv E^{(\tau)}(R_c) = \frac{2}{\sqrt{-\tau}} (\mathbf{K}(4\tau) - 2\mathbf{E}(4\tau)). \quad (6.18)$$

⁹It is worth to notice that the unperturbed energy $E^{(0)}$ displays the following divergent behavior for small R

$$E^{(0)} = \frac{\pi^2}{R} + 2R - \frac{R^3}{2\pi^2} + \mathcal{O}(R^7), \quad (6.15)$$

which resembles that of a CFT.




Figure 5. The kink solution to the $T\bar{T}$ -deformed sG model on a cylinder of radius R (a) and the corresponding energies as functions of R (b).

To find the behavior of $E^{(\tau)}$ as a function of R close to the branch singularity R_c , we first expand R and $E^{(\tau)}$ in powers of the small quantity $\varepsilon = \rho + 4\tau$

$$\begin{aligned} R - R_c &= \frac{R_c}{128\tau^2(1-4\tau)} \varepsilon^2 + \mathcal{O}(\varepsilon^3), \\ E^{(\tau)} - E_c^{(\tau)} &= \frac{R_c}{16\tau^2(1-4\tau)} \varepsilon + \mathcal{O}(\varepsilon^2), \end{aligned} \quad (6.19)$$

then, removing ε , one finds

$$E^{(\tau)} - E_c^{(\tau)} = \pm \frac{\sqrt{R_c}}{\tau\sqrt{2-8\tau}} \sqrt{R - R_c} + \mathcal{O}(R - R_c), \quad (6.20)$$

which gives a square root branch point at R_c for the energy.

Now we would like to briefly discuss the effect of the shock-wave singularities of the deformed solution on the Hamiltonian density. To compute the range of values of τ where the solution becomes multi-valued, we first identify the zeros of $\text{Det}(\mathcal{J}^{-1})$:

$$\text{Det}(\mathcal{J}^{-1}) = 0 \quad \Longleftrightarrow \quad x = \frac{\sqrt{\rho}}{2} \text{dn}^{-1} \left(\pm \sqrt{\frac{\rho + 4\tau}{8\tau}} \middle| -\rho \right), \quad (6.21)$$

where $\text{dn}^{-1}(z|\gamma)$ is the inverse of the Jacobi elliptic function $\text{dn}(z|\gamma)$. From the reality properties of $\text{dn}^{-1}(z|\gamma)$ it follows that x is real for

$$\rho > 0 \quad \wedge \quad \tau_1^* = \frac{\rho}{4+8\rho} < \tau < \frac{\rho}{4} = \tau_2^*, \quad (6.22)$$

where the critical values¹⁰ τ_1^* and τ_2^* corresponds to shock-wave singularities of the solution at $\phi = \pi$ and $\phi = 0, 2\pi$, respectively. The Hamiltonian density (6.6) is indeed singular when

$$\tau = \frac{\rho}{4 + 2\rho V} = \frac{\rho}{4 + 8\rho \sin^2(\phi/2)}, \quad (6.23)$$

which corresponds to the range of singular values of τ (6.22) as ϕ interpolates from 0 to 2π . However, it is important to stress that these branching singularities do not affect the total energy (6.11), which remains smooth in τ , since the singularities cancel out when dividing by ϕ_x in (6.11). In Figure 5 we displayed the behaviour of $\phi(x)$ (Figure 5a) and $E^{(\tau)}(R)$ (Figure 5b) for various values of τ . We see that the shock-wave phenomenon and the square root-type singularity occur at positive and negative values of τ , respectively.

7 Conclusions

Starting from the $T\bar{T}$ -deformed Lagrangians proposed in [2, 26, 27], the main result of this article is the direct derivation of the exact one-to-one map between solutions of the unperturbed and deformed equations of motion, which takes the general form (1.4,1.5). The result matches the topological gravity predictions of [18, 23] but it should be possible to obtain the fundamental equations (1.4,1.5,1.7) also by working within the framework introduced by Cardy in [22].

We initially arrived to this conclusion by studying the well known classical relation between sine-Gordon, the associated Lax operators and pseudo-spherical surfaces embedded in $\mathbb{R}^3$. We think that this alternative and more explicit approach to the problem may provide a complementary point of view compared to [18, 23] and open the way to the implementation of further integrable model tools, such as the Inverse Scattering Method and the ODE/IM correspondence within the $T\bar{T}$ /JT framework.

There are many theoretical aspects that deserve to be further explored. First of all, it would be conceptually very important to study fermionic theories and supersymmetric sigma models. In [26], it was argued that for the $T\bar{T}$ -perturbed Thirring model the Lagrangian truncates at second order in τ , such a truncation is not totally surprising, however the sine-Gordon Lagrangian is instead deformed in an highly non trivial way and it would be nice to identify the mechanism which allows to preserve the quantum equivalence between the two systems. Secondly, it would be important to continue the investigation of deformed 2D Yang-Mills [27], along the lines started in the interesting recent work [52]. These studies might also serve as a guide for the inclusion of the $T\bar{T}$ inside the Wilson Loop/Scattering Amplitude setup [34, 35] (see also the remarks in the outlook section of [52]).

Finally, it would also be interesting to study the generalisation of our results to the JT case described in [53–57] and to check whether for any of the higher-dimensional models discussed in [22, 26, 27, 58] there could exist a map, between deformed and undeformed solutions, similar to equations (1.4,1.5).

¹⁰Notice that, in the $\rho \rightarrow \infty$ limit, one recovers the 1-kink solution, and the critical range reduce to $\tau > \tau_1^* = \frac{1}{8}$, since $\tau_2^* \rightarrow \infty$.

Note: We have recently been informed that the coordinate map between deformed and undeformed classical Lagrangian systems was also independently introduced by Chih-Kai Chang and studied in an on-going research project involving also Christian Ferko and Savdeep Sethi.

Acknowledgments – We are especially grateful to Sasha Zamolodchikov, Sylvain Lacroix for inspiring discussions and help, and to Sergei Dubovsky for kindly guiding us through the recent literature connecting the $T\bar{T}$ perturbation to JT gravity. We also thank Andrea Cavaglià, Riccardo Borsato, Chih-Kai Chang, Yunfeng Jiang, Zohar Komargodski, Marc Magro, Marc Mezei, Leonardo Santilli, Alessandro Sfondrini, István Szécsényi and Miguel Tierz for useful discussions on related topics. This project was partially supported by the INFN project SFT, the EU network GATIS+, NSF Award PHY-1620628, and by the FCT Project PTDC/MAT-PUR/30234/2017 “Irregular connections on algebraic curves and Quantum Field Theory”.

A Short review on surfaces embedded in $\mathbb{R}^3$

The purpose of this appendix is to briefly review the basic concepts related to the classical theory of surfaces embedded in the Euclidean space $\mathbb{R}^3$. We will follow the standard constructive approach which can be found, for example, in [51]. Let us start by considering a surface Σ together with the vector-valued function $\mathbf{r}(\mathbf{z}) \in \mathbb{R}^3$, describing its embedding into 3-dimensional flat space. It is clear that the two vectors

$$\mathbf{r}_\mu = \frac{\partial}{\partial z^\mu} \mathbf{r}, \quad \mu = 1, 2, \quad (\text{A.1})$$

span the tangent plane $T_P \Sigma$ to the surface at any non-critical point $P \in \Sigma$.¹¹ We will disregard the subtleties arising with the presence of critical points and suppose that $\mathbf{r}_1(\mathbf{z}) \neq \mathbf{r}_2(\mathbf{z})$ for all points $P \in \Sigma$. This basis of $T_P \Sigma$ can be improved to a basis σ of $\mathbb{R}^3$ by adding the unit normal vector $\mathbf{n}$

$$\sigma = \{\mathbf{r}_1, \mathbf{r}_2, \mathbf{n}\}, \quad \mathbf{n} = \frac{\mathbf{r}_1 \times \mathbf{r}_2}{|\mathbf{r}_1 \times \mathbf{r}_2|}. \quad (\text{A.2})$$

The surface Σ inherits a metric structure from the ambient space $\mathbb{R}^3$ and its line element, also known as *first fundamental quadratic form*, is

$$I \equiv ds^2 = d\mathbf{r} \cdot d\mathbf{r} = \mathbf{r}_\mu \cdot \mathbf{r}_\nu dz^\mu dz^\nu. \quad (\text{A.3})$$

The tensor

$$g_{\mu\nu} = \mathbf{r}_\mu \cdot \mathbf{r}_\nu, \quad (\text{A.4})$$

is called *first fundamental tensor* or *metric tensor* of the surface Σ . According to the classical theorem by Bonnet [42] any surface embedded in flat 3-space is uniquely determined, up to isometries, by the first and the *second fundamental quadratic form*, defined as

$$II = -d\mathbf{r} \cdot d\mathbf{n} = -\mathbf{r}_\nu \cdot \left(\frac{\partial}{\partial z^\mu} \mathbf{n} \right) dz^\mu dz^\nu = \left(\frac{\partial}{\partial z^\mu} \mathbf{r}_\nu \right) \cdot \mathbf{n} dz^\mu dz^\nu. \quad (\text{A.5})$$

¹¹A critical point of a surface is, in this context, defined as a point P_c such that $\mathbf{r}_1(\mathbf{z}_c) = \mathbf{r}_2(\mathbf{z}_c)$.

The tensor

$$d_{\mu\nu} = \left(\frac{\partial}{\partial z^\mu} \mathbf{r}_\nu \right) \cdot \mathbf{n} , \quad (\text{A.6})$$

describes the projection of the vectors $\frac{\partial}{\partial z^\mu} \mathbf{r}_\nu(P)$ on the normal direction and tells us how much the surface curves away from the tangent space in an infinitesimal interval around the point P . These two tensors can be combined into the object

$$s_\mu^\nu = d_{\mu\rho} g^{\rho\nu} , \quad g_{\mu\rho} g^{\rho\nu} = \delta_\mu^\nu , \quad (\text{A.7})$$

known as *shape* or *Weingarten operator*, whose eigenvalues κ_1, κ_2 are the *principal curvatures* of the surface Σ . The latter quantities are geometric invariants, meaning that they do not change under coordinate transformations. Usually they are combined into the *Gauss* and *mean* curvatures

$$K = \kappa_1 \kappa_2 = \text{Det}(s_\mu^\nu) , \quad H = \frac{\kappa_1 + \kappa_2}{2} = \frac{1}{2} s_\mu^\mu . \quad (\text{A.8})$$

The tensors $g_{\mu\nu}$ and $d_{\mu\nu}$ determine the structural equations for embedded surfaces, comprising the *Gauss equations*

$$\frac{\partial}{\partial z^\mu} \mathbf{r}_\nu = \Gamma_{\mu\nu}^\rho \mathbf{r}_\rho + d_{\mu\nu} \mathbf{n} , \quad (\text{A.9})$$

and the *Weingarten equations*

$$\frac{\partial}{\partial z^\mu} \mathbf{n} = s_\mu^\nu \mathbf{r}_\nu , \quad (\text{A.10})$$

where we introduced the *Christoffel symbols* for the metric

$$\Gamma_{\mu\nu}^\rho = \frac{1}{2} g^{\rho\sigma} \left(\frac{\partial}{\partial z^\nu} g_{\mu\sigma} + \frac{\partial}{\partial z^\mu} g_{\nu\sigma} - \frac{\partial}{\partial z^\sigma} g_{\mu\nu} \right) . \quad (\text{A.11})$$

These equations describe how the frame σ moves on the surface and can be collected into the following linear system

$$\frac{\partial}{\partial z^\mu} \sigma = U_\mu \sigma , \quad (\text{A.12})$$

with¹²

$$U_1 = \begin{pmatrix} \Gamma_{11}^1 & \Gamma_{11}^2 & d_{11} \\ \Gamma_{12}^1 & \Gamma_{12}^2 & d_{12} \\ -s_1^1 & -s_1^2 & 0 \end{pmatrix} , \quad U_2 = \begin{pmatrix} \Gamma_{12}^1 & \Gamma_{12}^2 & d_{12} \\ \Gamma_{22}^1 & \Gamma_{22}^2 & d_{22} \\ -s_2^1 & -s_2^2 & 0 \end{pmatrix} . \quad (\text{A.13})$$

These structural equations are subject to a set of compatibility conditions called *Gauss-Mainardi-Codazzi* (GMC) system, which takes the form of a zero curvature condition on the matrices U_μ

$$\partial_2 U_1 - \partial_1 U_2 + [U_1, U_2] = 0 . \quad (\text{A.14})$$

Note that the matrices U_μ do not form a Lax pair in the usual sense, since no spectral parameter is present. Moreover, these matrices do not belong to any particular semi-simple Lie algebra. Specialising this general construction to the sine-Gordon case, we will show how

¹²Note that $\Gamma_{\mu\nu}^\rho = \Gamma_{\nu\mu}^\rho$ and $d_{\mu\nu} = d_{\nu\mu}$.

to build a proper Lax pair out of the matrices U_μ .

As a first example, consider a pseudo-spherical surface. In this case the Gauss curvature is $K = -\mu^2 < 0$, with constant μ , and one can choose as parametric curves the asymptotic lines, for which $d_{11} = d_{22} = 0$. Setting $\Delta^2 = \text{Det}(g_{\mu\nu})$, we see that

$$K = -\frac{d_{12}^2}{\Delta^2}. \quad (\text{A.15})$$

After some manipulations [51], it can be shown that in this case the Mainardi-Codazzi equations imply

$$\Gamma_{12}^1 = \Gamma_{12}^2 = 0 \implies \frac{\partial}{\partial z^2}(g_{11}) = \frac{\partial}{\partial z^1}(g_{22}) = 0. \quad (\text{A.16})$$

Defining the angle ω between the parametric lines as

$$\cos \omega = \frac{g_{12}}{\sqrt{g_{11}g_{22}}}, \quad \sin \omega = \frac{\Delta}{\sqrt{g_{11}g_{22}}}, \quad (\text{A.17})$$

we have the following expression for the fundamental forms

$$\text{I} = g_{11} (dz^1)^2 + 2\sqrt{g_{11}g_{22}} \cos \omega dz^1 dz^2 + g_{22} (dz^2)^2, \quad (\text{A.18})$$

$$\text{II} = 2\mu\sqrt{g_{11}g_{22}} \sin \omega dz^1 dz^2. \quad (\text{A.19})$$

Now, given the (anti-)holomorphicity of g_{11} and g_{22} we can rescale the variables z^μ to $z'^\mu = \sqrt{g_{\mu\mu}} z^\mu$ (no summation on repeated indices here) in terms of which one has¹³

$$\text{I} = (dz'^1)^2 - 2\cos \omega dz'^1 dz'^2 + (dz'^2)^2, \quad (\text{A.20})$$

$$\text{II} = 2\mu \sin \omega dz'^1 dz'^2. \quad (\text{A.21})$$

It is possible to show that the GMC system (A.14) reduces to the sine-Gordon equation

$$\frac{\partial}{\partial z'^1} \frac{\partial}{\partial z'^2} \omega = \mu^2 \sin \omega. \quad (\text{A.22})$$

Let us now consider the matrices U_μ

$$U_1 = \begin{pmatrix} \omega_1 \cot \omega & -\omega_1 \csc \omega & 0 \\ 0 & 0 & \mu \sin \omega \\ \mu \cot \omega & -\mu \csc \omega & 0 \end{pmatrix},$$

$$U_2 = \begin{pmatrix} 0 & 0 & \mu \sin \omega \\ -\omega_2 \csc \omega & \omega_2 \cot \omega & 0 \\ -\mu \csc \omega & \mu \cot \omega & 0 \end{pmatrix}, \quad (\text{A.23})$$

where $\omega_\mu = \frac{\partial}{\partial z^\mu} \omega$. The matrices (A.23) do not belong to $\mathfrak{su}(2)$, as we would expect, and contain no trace of the spectral parameter λ . We can fix these apparent problems by the

¹³This corresponds to a parametrization of the surface by arc-length along the asymptotic lines.

following considerations. First we notice that the triple $\sigma = \{\mathbf{r}_1, \mathbf{r}_2, \mathbf{n}\}$ is not orthonormal. However, the rotation

$$\sigma \longrightarrow \tilde{\sigma} = M\sigma, \quad M = \begin{pmatrix} 1 & 0 & 0 \\ -\cot \omega & \csc \omega & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (\text{A.24})$$

which corresponds to a gauge transformation on the matrices U_μ

$$U_\mu \longrightarrow \tilde{U} = (\partial_\mu M) M^{-1} + M U_\mu M^{-1}, \quad (\text{A.25})$$

leaves the compatibility equation – the sine-Gordon equation – invariant and maps (A.23) into

$$\tilde{U}_1 = \begin{pmatrix} 0 & -\omega_1 & 0 \\ \omega_1 & 0 & \mu \\ 0 & -\mu & 0 \end{pmatrix}, \quad \tilde{U}_2 = \begin{pmatrix} 0 & 0 & \mu \sin \omega \\ 0 & 0 & -\mu \cos \omega \\ -\mu \sin \omega & \mu \cos \omega & 0 \end{pmatrix}, \quad (\text{A.26})$$

which now belong to the $\mathfrak{su}(2)$ algebra. Finally, the spectral parameter can be recovered by noticing that the sine-Gordon equation is invariant under the following transformation

$$(z'^1, z'^2, \mu) = \left(\alpha \tilde{z}^1, \beta \tilde{z}^2, \frac{1}{\sqrt{\alpha\beta}} m \right), \quad (\text{A.27})$$

for any constant α and β . Choosing $\alpha = \sqrt{2}m$ and $\beta = \sqrt{2}\frac{m}{\lambda^2}$ and writing $\omega = \beta\phi$, we obtain

$$\text{I} = 2m^2 \left((d\tilde{z}^1)^2 - \frac{2}{\lambda^2} \cos \beta\phi d\tilde{z}^1 d\tilde{z}^2 + \frac{1}{\lambda^4} (d\tilde{z}^2)^2 \right), \quad (\text{A.28})$$

$$\text{II} = 2\sqrt{2}\frac{m^2}{\lambda} \sin \beta\phi d\tilde{z}^1 d\tilde{z}^2, \quad (\text{A.29})$$

which coincides with the quadratic forms (2.11, 2.12).

Finally, as another interesting example of integrable model associated to embedded surfaces, let us briefly discuss a constant mean curvature surface, *i.e.* a surface such that $H = \text{const.}$. In this case one can choose conformal coordinates, in which the fundamental forms simplify to

$$\text{I} = \frac{2}{H^2} e^\omega dz^1 dz^2, \quad (\text{A.30})$$

$$\text{II} = \frac{1}{H} \left[A_1 (dz^1)^2 + 2e^\omega dz^1 dz^2 + A_2 (dz^2)^2 \right]. \quad (\text{A.31})$$

Some simple computation shows that the GCM equations are equivalent to the system

$$\frac{\partial}{\partial z^1} \frac{\partial}{\partial z^2} \omega = e^\omega - A_1 A_2 e^{-\omega}, \quad (\text{A.32})$$

$$\frac{\partial}{\partial z^2} A_1 = \frac{\partial}{\partial z^1} A_2 = 0, \quad (\text{A.33})$$

which is known as *modified sinh-Gordon equation*. Its Gauss curvature is

$$K = H^2 (1 - A_1 A_2 e^{-2\omega}) . \quad (\text{A.34})$$

Rescaling the field as $\omega \rightarrow \omega + 2 \ln H$, the functions A_i as $A_i \rightarrow H A_i$ and sending $H \rightarrow 0$ yields a minimal surface and reduces the GMC system to Liouville equation

$$\frac{\partial}{\partial z^1} \frac{\partial}{\partial z^2} \omega = K e^\omega , \quad K = -A_1 A_2 e^{-2\omega} . \quad (\text{A.35})$$

B Computation of the fundamental quadratic forms from sine-Gordon ZCR

While in the preceding appendix we presented the derivation of soliton equations starting from the basic geometric data of some particular surface, here we wish to follow the reverse path and explicitly show how to obtain the forms (2.11, 2.12) starting from sine-Gordon ZCR (2.8, 2.9). First of all we need to find a basis of $\mathfrak{su}(2)$ with respect to the Killing form

$$(a, b)_K = \text{Tr}(\mathbf{Ad}(a) \mathbf{Ad}(b)) , \quad a, b \in \mathfrak{su}(2) . \quad (\text{B.1})$$

In the adjoint representation one has $T^i = \mathbf{Ad}(\mathcal{S}^i)$, with

$$T^1 = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} , \quad T^2 = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} , \quad T^3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix} , \quad (\text{B.2})$$

and

$$(T^i, T^j)_K = -2\delta^{ij} . \quad (\text{B.3})$$

The orthonormal basis is easily found to be

$$\mathbf{e}^i = \frac{\mathbf{i}}{\sqrt{2}} \mathcal{S}^i , \quad (\text{B.4})$$

and we see that for a pair of matrices A and B belonging to the 2-dimensional representation of $\mathfrak{su}(2)$, one has

$$(A, B)_K = 4 \text{Tr}(AB) . \quad (\text{B.5})$$

Now we need the partial derivatives of r (2.4)

$$r = \Phi^{-1} \frac{\partial}{\partial \lambda} \Phi \implies \frac{\partial}{\partial z^\mu} r = \Phi^{-1} \frac{\partial}{\partial \lambda} (L_\mu \Phi) - \Phi^{-1} L_\mu \Phi \Phi^{-1} \frac{\partial}{\partial \lambda} \Phi , \quad (\text{B.6})$$

where we have used the linear system $\partial_\mu \Phi = L_\mu \Phi$. We have then

$$\frac{\partial}{\partial z^\mu} r = \Phi^{-1} \frac{\partial L_\mu}{\partial \lambda} \Phi . \quad (\text{B.7})$$

We can immediately compute the metric tensor $g_{\mu\nu}$

$$g_{\mu\nu} = \left(\frac{\partial r}{\partial z^\mu}, \frac{\partial r}{\partial z^\nu} \right)_K = 4 \operatorname{Tr} \left(\frac{\partial r}{\partial z^\mu} \frac{\partial r}{\partial z^\nu} \right) = 4 \operatorname{Tr} \left(\frac{\partial L_\mu}{\partial \lambda} \frac{\partial L_\nu}{\partial \lambda} \right). \quad (\text{B.8})$$

Inserting the expressions (2.8, 2.9) we obtain

$$g_{\mu\nu} = 2m^2 \begin{pmatrix} 1 & -\frac{1}{\lambda^2} \cos(\beta\phi) \\ -\frac{1}{\lambda^2} \cos(\beta\phi) & \frac{1}{\lambda^4} \end{pmatrix}_{\mu\nu}. \quad (\text{B.9})$$

The second derivatives of r follow from simple computations

$$\frac{\partial}{\partial z^\mu} \frac{\partial}{\partial z^\nu} r = \Phi^{-1} \left(\frac{\partial}{\partial z^\nu} \frac{\partial L_\mu}{\partial \lambda} + \left[\frac{\partial L_\mu}{\partial \lambda}, L_\nu \right] \right) \Phi. \quad (\text{B.10})$$

The matrix version of the unit normal is

$$n = \sum_{i=1}^3 n_i \mathcal{S}^i = \frac{1}{2\sqrt{2}} \frac{\left[\frac{\partial r}{\partial z^1}, \frac{\partial r}{\partial z^2} \right]}{\sqrt{\operatorname{Det} \left(\left[\frac{\partial r}{\partial z^1}, \frac{\partial r}{\partial z^2} \right] \right)}}. \quad (\text{B.11})$$

We obtain that

$$\operatorname{Det} \left(\left[\frac{\partial r}{\partial z^1}, \frac{\partial r}{\partial z^2} \right] \right) = \left(\frac{m^2}{2\lambda^2} \sin(\beta\phi) \right)^2. \quad (\text{B.12})$$

We can finally compute the second fundamental tensor

$$\begin{aligned} d_{\mu\nu} &= \left(\frac{\partial}{\partial z^\mu} \frac{\partial}{\partial z^\nu} r, n \right)_K \\ &= \frac{1}{\sqrt{2}} \frac{\lambda^2}{m^2 \sin(\beta\phi)} \operatorname{Tr} \left(\left[\frac{\partial L_1}{\partial \lambda}, \frac{\partial L_2}{\partial \lambda} \right] \left(\frac{\partial}{\partial z^\nu} \frac{\partial L_\mu}{\partial \lambda} + \left[\frac{\partial L_\mu}{\partial \lambda}, L_\nu \right] \right) \right). \end{aligned} \quad (\text{B.13})$$

The explicit expression is

$$d_{\mu\nu} = \frac{\sqrt{2}m^2}{\lambda} \sin(\beta\phi) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (\text{B.14})$$

References

- [1] F. A. Smirnov and A. B. Zamolodchikov, *On space of integrable quantum field theories*, *Nucl. Phys.* **B915** (2017) 363–383 [[arXiv:1608.05499](#)].
- [2] A. Cavaglià, S. Negro, I. M. Szécsényi and R. Tateo, *TT -deformed 2D Quantum Field Theories*, *JHEP* **10** (2016) 112 [[arXiv:1608.05534](#)].
- [3] A. B. Zamolodchikov, *Expectation value of composite field TT in two-dimensional quantum field theory*, [hep-th/0401146](#).
- [4] S. Dubovsky, R. Flauger and V. Gorbenko, *Solving the Simplest Theory of Quantum Gravity*, *JHEP* **09** (2012) 133 [[arXiv:1205.6805](#)].

- [5] S. Dubovsky, R. Flauger and V. Gorbenko, *Effective String Theory Revisited*, *JHEP* **09** (2012) 044 [[arXiv:1203.1054](#)].
- [6] M. Caselle, D. Fioravanti, F. Gliozzi and R. Tateo, *Quantisation of the effective string with TBA*, *JHEP* **07** (2013) 071 [[arXiv:1305.1278](#)].
- [7] C. Chen, P. Conkey, S. Dubovsky and G. Hernandez-Chifflet, *Undressing Confining Flux Tubes with $T\bar{T}$* , [arXiv:1808.01339](#).
- [8] L. McGough, M. Mezei and H. Verlinde, *Moving the CFT into the bulk with $T\bar{T}$* , *JHEP* **04** (2018) 010 [[arXiv:1611.03470](#)].
- [9] G. Turiaci and H. Verlinde, *Towards a 2d QFT Analog of the SYK Model*, *JHEP* **10** (2017) 167 [[arXiv:1701.00528](#)].
- [10] A. Giveon, N. Itzhaki and D. Kutasov, *$T\bar{T}$ and LST*, *JHEP* **07** (2017) 122 [[arXiv:1701.05576](#)].
- [11] A. Giveon, N. Itzhaki and D. Kutasov, *A solvable irrelevant deformation of AdS_3/CFT_2* , *JHEP* **12** (2017) 155 [[arXiv:1707.05800](#)].
- [12] M. Asrat, A. Giveon, N. Itzhaki and D. Kutasov, *Holography Beyond AdS*, *Nucl. Phys.* **B932** (2018) 241–253 [[arXiv:1711.02690](#)].
- [13] G. Giribet, *$T\bar{T}$ -deformations, AdS/CFT and correlation functions*, *JHEP* **02** (2018) 114 [[arXiv:1711.02716](#)].
- [14] P. Kraus, J. Liu and D. Marolf, *Cutoff AdS_3 versus the $T\bar{T}$ deformation*, *JHEP* **07** (2018) 027 [[arXiv:1801.02714](#)].
- [15] W. Cottrell and A. Hashimoto, *Comments on $T\bar{T}$ double trace deformations and boundary conditions*, [arXiv:1801.09708](#).
- [16] M. Baggio and A. Sfondrini, *Strings on NS-NS Backgrounds as Integrable Deformations*, *Phys. Rev.* **D98** (2018), no. 2 021902 [[arXiv:1804.01998](#)].
- [17] J. P. Babaro, V. F. Foit, G. Giribet and M. Leoni, *$T\bar{T}$ type deformation in the presence of a boundary*, *JHEP* **08** (2018) 096 [[arXiv:1806.10713](#)].
- [18] S. Dubovsky, V. Gorbenko and M. Mirbabayi, *Asymptotic fragility, near AdS_2 holography and $T\bar{T}$* , *JHEP* **09** (2017) 136 [[arXiv:1706.06604](#)].
- [19] G. Mussardo and P. Simon, *Bosonic type S matrix, vacuum instability and CDD ambiguities*, *Nucl. Phys.* **B578** (2000) 527–551 [[hep-th/9903072](#)].
- [20] A.I. B. Zamolodchikov, *From tricritical Ising to critical Ising by thermodynamic Bethe ansatz*, *Nucl. Phys.* **B358** (1991) 524–546.
- [21] S. Dubovsky, V. Gorbenko and M. Mirbabayi, *Natural Tuning: Towards A Proof of Concept*, *JHEP* **09** (2013) 045 [[arXiv:1305.6939](#)].
- [22] J. Cardy, *The $T\bar{T}$ deformation of quantum field theory as a stochastic process*, [arXiv:1801.06895](#).
- [23] S. Dubovsky, V. Gorbenko and G. Hernandez-Chifflet, *$T\bar{T}$ partition function from topological gravity*, *JHEP* **09** (2018) 158 [[arXiv:1805.07386](#)].
- [24] S. Datta and Y. Jiang, *$T\bar{T}$ deformed partition functions*, *JHEP* **08** (2018) 106 [[arXiv:1806.07426](#)].

- [25] O. Aharony, S. Datta, A. Giveon, Y. Jiang and D. Kutasov, *Modular invariance and uniqueness of $T\bar{T}$ deformed CFT*, [arXiv:1808.02492](#).
- [26] G. Bonelli, N. Doroud and M. Zhu, *$T\bar{T}$ -deformations in closed form*, *JHEP* **06** (2018) 149 [[arXiv:1804.10967](#)].
- [27] R. Conti, L. Iannella, S. Negro and R. Tateo, *Generalised Born-Infeld models, Lax operators and the $T\bar{T}$ perturbation*, *JHEP* **11** (2018) 007 [[arXiv:1806.11515](#)].
- [28] A. Dei and A. Sfondrini, *Integrable spin chain for stringy Wess-Zumino-Witten models*, *JHEP* **07** (2018) 109 [[arXiv:1806.00422](#)].
- [29] P. Dorey and R. Tateo, *Anharmonic oscillators, the thermodynamic Bethe ansatz, and nonlinear integral equations*, *J. Phys.* **A32** (1999) L419–L425 [[hep-th/9812211](#)].
- [30] V. V. Bazhanov, S. L. Lukyanov and A. B. Zamolodchikov, *Spectral determinants for Schrodinger equation and Q operators of conformal field theory*, *J. Statist. Phys.* **102** (2001) 567–576 [[hep-th/9812247](#)].
- [31] P. Dorey, C. Dunning and R. Tateo, *The ODE/IM Correspondence*, *J. Phys.* **A40** (2007) R205 [[hep-th/0703066](#)].
- [32] S. L. Lukyanov and A. B. Zamolodchikov, *Quantum Sine(h)-Gordon Model and Classical Integrable Equations*, *JHEP* **07** (2010) 008 [[arXiv:1003.5333](#)].
- [33] P. Dorey, S. Faldella, S. Negro and R. Tateo, *The Bethe Ansatz and the Tzitzeica-Bullough-Dodd equation*, *Phil. Trans. Roy. Soc. Lond.* **A371** (2013) 20120052 [[arXiv:1209.5517](#)].
- [34] L. F. Alday, D. Gaiotto and J. Maldacena, *Thermodynamic Bubble Ansatz*, *JHEP* **09** (2011) 032 [[arXiv:0911.4708](#)].
- [35] L. F. Alday, J. Maldacena, A. Sever and P. Vieira, *Y-system for Scattering Amplitudes*, *J. Phys.* **A43** (2010) 485401 [[arXiv:1002.2459](#)].
- [36] D. Gaiotto, *Opers and TBA*, [arXiv:1403.6137](#).
- [37] K. Ito and H. Shu, *ODE/IM correspondence and the Argyres-Douglas theory*, *JHEP* **08** (2017) 071 [[arXiv:1707.03596](#)].
- [38] A. Grassi and M. Mariño, *A Solvable Deformation Of Quantum Mechanics*, [arXiv:1806.01407](#).
- [39] P. Goddard, J. Goldstone, C. Rebbi and C. B. Thorn, *Quantum dynamics of a massless relativistic string*, *Nucl. Phys.* **B56** (1973) 109–135.
- [40] E. Bour, *Théorie de la déformation des surfaces*, *J. de l'École Imperiale Polytech.* **19** (1862) 1–48.
- [41] J. Liouville, *Sur l'équation aux différences partielles $\frac{d^2 \ln \lambda}{dudv} \pm \frac{\lambda}{2a^2} = 0$* , *J. de mathématiques pures et appliquées* **18** (1853) 71–72.
- [42] P. O. Bonnet, *Mémoire sur la théorie des surfaces applicables sur une surface donnée*, *J. de l'École Polytech.* **42** (1867) 1–151.
- [43] K. Pohlmeyer, *Integrable Hamiltonian Systems and Interactions through Quadratic Constraints*, *Commun. math. Phys.* **46** (1976) 207–221.
- [44] F. Lund and T. Regge, *Unified approach to strings and vortices with soliton solutions*, *Phys. Rev.* **D 14** (1976) 1524–1535.

- [45] B. S. Getmanov, *New Lorentz-invariant system with exact multisoliton solutions*, *JETP Lett.* **25** (1977) 119–122.
- [46] A. Sym, *Soliton Surfaces*, *Lett. al Nuovo Cimento* **33** (1982) 394–400.
- [47] A. Sym, *Soliton Surfaces. II. Geometric Unification of Solvable Nonlinearities*, *Lett. al Nuovo Cimento* **36** (1983) 307–312.
- [48] A. Sym, *Soliton Surfaces. III. Solvable nonlinearities with trivial geometry.*, *Lett. al Nuovo Cimento* **39** (1984) 193–196.
- [49] A. Sym, *Soliton Surfaces. VI. Gauge Invariance and Final Formulation of the Approach.*, *Lett. al Nuovo Cimento* **41** (1984) 353–360.
- [50] A. Sym, *Soliton Surfaces. V. Geometric Theory of Loop Solitons.*, *Lett. al Nuovo Cimento* **41** (1984) 33–40.
- [51] C. Rogers and W. Schief, *Bäcklund and Darboux transformations: geometry and modern applications in soliton theory*, vol. 30. Cambridge University Press, 2002.
- [52] L. Santilli and M. Tierz, *Large N phase transition in $T\bar{T}$ -deformed 2d Yang-Mills theory on the sphere*, [arXiv:1810.05404](#).
- [53] A. Bzowski and M. Guica, *The holographic interpretation of $J\bar{T}$ -deformed CFTs*, [arXiv:1803.09753](#).
- [54] M. Guica, *An integrable Lorentz-breaking deformation of two-dimensional CFTs*, [arXiv:1710.08415](#).
- [55] S. Chakraborty, A. Giveon and D. Kutasov, *$J\bar{T}$ deformed CFT_2 and string theory*, *JHEP* **10** (2018) 057 [[arXiv:1806.09667](#)].
- [56] L. Apolo and W. Song, *Strings on warped AdS_3 via $T\bar{J}$ deformations*, [arXiv:1806.10127](#).
- [57] O. Aharony, S. Datta, A. Giveon, Y. Jiang and D. Kutasov, *Modular covariance and uniqueness of $J\bar{T}$ deformed CFTs*, [arXiv:1808.08978](#).
- [58] M. Taylor, *TT deformations in general dimensions*, [arXiv:1805.10287](#).