

Proxy Controls and Panel Data

Ben Deaner*

July 28, 2020

Abstract

We present a flexible approach to the identification and estimation of causal objects in nonparametric, non-separable models with confounding. Key to our analysis is the use of ‘proxy controls’: covariates that do not satisfy a standard ‘unconfoundedness’ assumption but are informative proxies for variables that do. Our analysis applies to both cross-sectional and panel models. Our identification results motivate a simple and ‘well-posed’ nonparametric estimator and we analyze its asymptotic properties. In panel settings, our methods provide a novel approach to the difficult problem of identification with non-separable general heterogeneity and fixed T . In panels, observations from different periods serve as proxies for unobserved heterogeneity and our key identifying assumptions follow from restrictions on the serial dependence structure. We apply our methodology to two empirical settings. We estimate causal effects of grade retention on cognitive performance using cross-sectional variation and we estimate a structural Engel curve for food using panel data.

The threat of confounding is the defining challenge of empirical economics. Confounding is familiar to quantitative researchers in all fields, but it is of particular concern to economists, who are interested primarily in causal inference and whose data are usually drawn from observational studies. Ideally, a researcher can combat confounding by conditioning on ‘perfect controls’: variables that explain all association between assigned treatments and potential outcomes. Unfortunately, perfect controls are often unavailable, but the researcher may have access to covariates that can proxy for an ideal set of unobserved perfect controls. These ‘proxy controls’ could be a set of test scores that proxy for innate ability, or they could be self-reported wages and years in education in place of socio-economic status.

*PhD candidate, Massachusetts Institute of Technology. Email at bdeaner@mit.edu. Previous editions of this work can be found at <https://arxiv.org/abs/1810.00283> (first version September 30th 2018). Many thanks to my advisers Whitney Newey, Anna Mikusheva, and Jerry Hausman for numerous helpful comments on this work and thanks to those who attended the Econometric society session C1 at the ASSA 2020 Annual Meeting in San Diego for useful feedback and likewise to those who attended via Zoom the Treatment Effects 2 session of the 2020 Econometric Society World Congress. Thanks as well to those who attended presentations on this project at MIT, in particular Florian Gunsilius, Alberto Abadie, Claire Lazar, and others.

A naive approach would treat the proxies as if they were perfect controls. For example, one could treat test scores as if they did in fact perfectly measure ability. However, if the proxies mis-measure the perfect controls, then controlling for the proxies in the conventional manner need not remove all confounding and the resulting estimates would likely be asymptotically biased.

We develop new results on non-parametric identification and estimation of causal objects when only proxy controls are available. We suggest a non-parametric estimation procedure based on our identification results that is straightforward to implement and is, to the best of our knowledge, novel. We establish consistency and a convergence rate for our estimator under our identifying assumptions and some other primitive conditions.

While our analysis applies in cross-sectional settings, our results are well-suited to the context of panel data. In panels, observations from other time periods can be informative proxies for factors that explain the confounding (i.e., perfect controls). To illustrate, suppose that innate ability is a confounding factor. Then an individual's innate ability is associated with both the individual's treatment assignments and potential outcomes. It follows that the history of treatment assignments is informative about innate ability. We provide conditions on the serial dependence structure of the data and latent variables so that one can form proxy controls from past observations that satisfy our conditional independence restrictions. Thus we add to the panel literature by suggesting a novel non-parametric approach to identification and estimation of causal objects in fixed- T panel models with general, non-separable heterogeneity.

Our analysis treats identification and estimation with proxy controls as a measurement error problem. Proxy controls mis-measure a set of latent perfect controls. To account for the measurement error, the researcher divides the available proxy controls into two groups and, in effect, uses one group of proxy controls to instrument for the other. The validity of this approach does not require that the proxy controls be valid instruments in the standard sense. Instead, the proxy controls must satisfy a conditional independence restriction which, loosely speaking, states that the two sets of proxy controls are only related through their mutual association with the unobserved perfect controls. In addition, we require that the proxy controls are sufficiently informative (defined in terms of statistical completeness) about the latent perfect controls.

We specify conditions under which the estimation problem suggested by our identification result is 'well-posed'. The well-posedness of our estimation problem allows us to derive simple and fast rates of convergence under primitive conditions. We present an easy-to-implement non-parametric estimator that builds on our identification results and analyze its properties.

To demonstrate the usefulness of our methodology we apply it to two very different real world data problems. We revisit the empirical setting of [Fruehwirth *et al.* \(2016\)](#) who use data from the Early Childhood Longitudinal Study of Kindergartners (ECLS-K) to estimate the causal impact of grade retention on the performance of US students in cognitive tests. We use their data to estimate some of the same counterfactual objects using our methods. While the ECLS-K provides four periods of panel data, we use this setting to demonstrate

the application of our methods to cross-sectional data.

In addition, we use data from the Panel Survey of Income Dynamics (PSID) to estimate a structural Engel curve for food. In this case our analysis is premised upon a Markov-type serial dependence restriction.

To summarize, our contribution is fourfold. We provide new results for identification with proxy controls that apply to both the cross-sectional and panel settings. We show that in dynamic panel settings our conditional independence restrictions follow from conditions on the serial dependence structure. We propose a novel estimation method based on our identification results and establish convergence rates under primitive conditions. Finally, we use our methodology to estimate causal objects in two separate empirical settings.

Related Literature

In the biometrics literature, [Miao *et al.* \(2018\)](#) identify an average structural function when controls are mis-measured and develop a related statistical test, they do not provide an estimator. Their work in turn builds on the work of [Kuroki & Pearl \(2014\)](#) for discrete variables.

Our identification results are closely related to those in [Miao *et al.* \(2018\)](#). However, our identification results differ in key ways. Firstly, our analysis allows us to identify the conditional average structural function (CASF) or the conditional distribution of potential outcomes (conditional on assigned treatments). By contrast, [Miao *et al.* \(2018\)](#) only identify the unconditional average structural function/marginal distribution. Identification of the CASF allows us to identify important counterfactual objects like the average effect of treatment on the treated (for further examples see our empirical applications). Secondly, we introduce a condition, Assumption 4.i, which implies the well-posedness of estimators based on our characterization of the CASF. Well-posedness is crucial for deriving fast and simple convergence rates for our estimator. Thirdly, we consider panel models and present panel-type assumptions that are sufficient for our key identifying assumptions. Further comparison of our identification results and those of [Miao *et al.* \(2018\)](#) can be found in Appendix A.2.

[Hu & Schennach \(2008\)](#) provide identification results for nonparametric and nonseparable models with measurement error and present a related estimator. [Rokkanen \(2015\)](#) employs results from [Hu & Schennach \(2008\)](#) to achieve identification using proxy controls in a regression discontinuity setting. We treat the proxy controls problem as one of measurement error and our assumptions resemble some of those in [Hu & Schennach \(2008\)](#). In particular, both of our analyses require completeness assumptions involving latent factors and conditional independence of some observables conditional on the unobserved factors. Unlike [Hu & Schennach \(2008\)](#), we do not require a normalization like mean- or median-unbiasedness of the mis-measured variables and we are able to provide a simpler, constructive identification of causal objects and an uncomplicated, well-posed estimation method. We are able to do this because in our problem the measurement error is only in control variables and not in treatment variables. By contrast, the results of [Hu & Schennach \(2008\)](#) apply for measurement

error in treatments rather than just controls. Identification with proxy controls is somewhat more tractable than the problem of measurement error in treatment variables. We are not interested in the causal effect of the latent perfect controls themselves and we do not recover any distributions involving the latent perfect controls, even as an intermediate step in estimation. See Appendix A.2 for further discussion.

Our panel analysis follows a long line of work in which observations from other periods are used to account for unobserved heterogeneity. This approach is the basis of classic methods like those of Hausman & Taylor (1981), Holtz-Eakin *et al.* (1988), and Arellano & Bond (1991). A recent literature takes a similar approach to control for unobserved heterogeneity that is non-additive and appears in non-linear models. For example, the work of Arellano & Bonhomme (2016), Freyberger (2018) and Evdokimov (2009).

A key step in our estimation method is a Penalized Sieve Minimum Distance (PSMD) procedure. PSMD estimators are analyzed extensively in Chen & Pouzo (2012) and Chen & Pouzo (2015). PSMD estimation generalizes earlier methods for estimating conditional moment restriction models, for example those presented in Newey & Powell (2003) and Ai & Chen (2003).

There is a sizable literature examining the effects of grade retention on cognitive and social success. For a meta-analysis of some earlier literature on this topic see Jimerson (2001). Our analysis builds on the work of Fruehwirth *et al.* (2016). We use the data available in the supplement of their paper and their code to clean that data and we estimate some of the same causal effects as they do in their work.

Recent work to estimate structural (i.e., causal) Engel curves for food in a nonparametric or semiparametric fashion includes the instrumental variables approach of Blundell *et al.* (2007) and the panel approach of Chernozhukov *et al.* (2015). For a short survey see Lewbel (2008).

1 General Model and Identification

Consider the following structural model:

$$Y = y_0(X, U) \tag{1.1}$$

Y is an observed dependent variable, X is a column vector of observables that represents the levels of assigned treatments, and U is a (potentially infinite-dimensional) vector that represents unobserved heterogeneity. The ‘structural function’ y_0 is not assumed to be of any particular parametric form.

The model above incorporates both cross-sectional and panel settings. In the panel case the model applies for a particular period t , that is, for a particular cross-sectional slice of the panel data. We could make the time-dependence explicit and rewrite the model above as $Y_t = y_{0,t}(X_t, U_t)$.

Throughout the discussion it is assumed that the structural function y_0 in (1.1) captures the causal effect of X on Y . For clarity, we situate our analysis in the potential outcomes framework. If a unit has realization of the heterogeneity

U of u , then $y_0(x, u)$ is the unit's 'potential outcome' from treatment level x . That is, the outcome that would have been observed had the treatment of that unit been set to level x . Thus U captures all heterogeneity in the potential outcomes. We assume throughout (without loss of generality) that the mapping between the potential outcomes $y_0(\cdot, u)$ and heterogeneity in potential outcomes u is one-to-one. This means that independence of some variable with the heterogeneity U is equivalent to independence of that variable with the potential outcomes $y_0(\cdot, U)$.

The focus of this paper is on the identification and estimation of conditional average potential outcomes, where we condition on the assigned treatments X . We refer to the function that returns the conditional average potential outcomes as the 'conditional average structural function' (CASF).

The CASF \bar{y} is defined formally as follows:

$$\bar{y}(x_1|x_2) = E[y_0(x_1, U)|X = x_2]$$

In words, suppose we draw a unit at random from the sub-population who were assigned treatment $X = x_2$. Then the expected counterfactual outcome had the unit instead received treatment level x_1 is $\bar{y}(x_1|x_2)$.¹

One may also be interested in identifying average potential outcomes conditional on the treatments as well as some additional variables S which could represent say, membership of a demographic sub-group. That is, one may wish to identify $E[y_0(x_1, U)|X = x_2, S = s]$. Our results extend straight-forwardly to this case. For instance, if S is discrete one can simply apply our analysis to the sub-population with S at some fixed s .²

By transforming the model, one can define an even richer set of counterfactual objects in terms of the CASF of the transformed model. For example, let y be some fixed scalar and consider the transformation $w \mapsto 1\{w \leq y\}$. Let \tilde{Y} be the transformed outcome variable, that is $\tilde{Y} = 1\{Y \leq y\}$, and let \tilde{y}_0 be the transformed structural function, that is $\tilde{y}_0(x, u) = 1\{y_0(x, u) \leq y\}$. The transformed model is $\tilde{Y} = \tilde{y}_0(X, U)$.

The conditional cumulative distribution function of the potential outcomes in the original model can be written as:

$$\begin{aligned} P(y_0(x_1, U) \leq y|X = x_2) &= E[1\{y_0(x_1, U) \leq y\}|X = x_2] \\ &= E[\tilde{y}_0(x_1, U)|X = x_2] \end{aligned}$$

The right-hand side of the final equality above is the conditional average structural function for the transformed model. Our identifying assumptions do not refer to Y directly but instead to the latent variable U , thus if our assumptions apply for the original model they also apply for the transformed model. Therefore, if we can identify the CASF, then we can also identify the conditional

¹Note that if X is continuously distributed then $\bar{y}(x_1|x_2)$ is only uniquely defined for x_2 up to a set of F_X -measure zero, where F_X is the law of X .

²A previous working version of this paper explicitly incorporated additional conditioning variables S which could be continuous, discrete or a mixture of both. For ease of exposition we have dropped this feature from the current draft.

cumulative distribution function of the potential outcomes. This in turn implies identification of the conditional quantiles.

1.1 Proxy Controls

A common approach to identification in the presence of confounding relies on the presence of what we term ‘perfect controls’. A vector of perfect controls is an observable random vector W^* , so that conditioning on W^* , the treatments X and the heterogeneity in potential outcomes are independent. Formally, we write $U \perp\!\!\!\perp X|W^*$. We use this notation to denote conditional independence throughout the paper.

To achieve identification with observed perfect controls, one generally requires additional conditions, in particular something like a full support assumption for the distribution of W^* . We give sufficient conditions in Assumption 1 below.

Assumption 1 (Perfect Controls). i. $U \perp\!\!\!\perp X|W^*$ ii. The joint distribution $F_{(W^*, X)}$ is absolutely continuous with the product of the distributions F_{W^*} and F_X . iii. $E[|Y|] < \infty$ and for F_X -almost all x $E[|y_0(x, U)|] < \infty$.

Under Assumption 1 the CASF satisfies:

$$\bar{y}(x_1|x_2) = E[E[Y|W^*, X = x_1]|X = x_2]$$

Where the equality holds for F_X -almost all x_1 and x_2 . If W^* is observed, then the RHS of the final equality depends only on the distribution of observables, thus the equation above identifies $\bar{y}(x_1|x_2)$. The characterization above is well-known but we formally state and prove it in Proposition B.1 in Appendix B.

When perfect controls W^* are unavailable the researcher may have access to proxy controls. In particular, we suppose that the researcher has access to two vectors of proxies V and Z . We present assumptions that imply identification when only proxy controls V and Z are available. The assumptions refer to the vector of perfect controls W^* for which V and Z act as proxies. Since W^* is unobserved, the assumptions can be understood to state that a vector of latent variables W^* exists that simultaneously satisfies all the conditions in our assumptions. To argue persuasively that the assumptions are plausible in a given setting, a researcher will generally have to choose a particular set of unobserved perfect controls W^* and argue that the assumptions hold for those controls.

The proxy controls V and Z can be understood as measurements of W^* that are subject to non-classical (i.e., non-zero mean and non-additive) noise. To account for the measurement error, the researcher in effect uses the proxy controls in Z as instruments for the proxy controls in V . Neither V nor Z is required to be independent of the latent factors W^* .

As we discuss in Section 2, it may be useful to allow for the possibility that the vectors V and Z have components in common. Our assumptions do not

preclude this so long as the shared components are non-random conditional on W^* (e.g., if the shared components are also components of W^*). We denote the vector of shared components by \bar{W} and refer to them in Assumption 3 below. If Z and V do not share components then one can ignore \bar{W} in that assumption.

Assumption 2 (Conditional Independence). i. $U \perp\!\!\!\perp Z|(X, W^*)$ ii. $V \perp\!\!\!\perp (X, Z)|W^*$

Assumption 3 (Informativeness). i. For $F_{(X, \bar{W})}$ -almost all (x, \bar{w}) , for any function $\delta \in L_2(F_{W^*|\bar{W}=\bar{w}, X=x})$:

$$E[\delta(W^*)|X = x, Z = z] = 0 \iff \delta(w^*) = 0$$

Where the first equality is $F_{Z|X=x, \bar{W}=\bar{w}}$ -almost sure and the second $F_{W^*|X=x, \bar{W}=\bar{w}}$ -almost sure. ii. For $F_{(X, \bar{W})}$ -almost all (x, \bar{w}) , for any function $\delta \in L_2(F_{W^*|X=x, \bar{W}=\bar{w}})$:

$$E[\delta(W^*)|X = x, V = v] = 0 \iff \delta(w^*) = 0$$

Where the first equality is $F_{V|X=x, \bar{W}=\bar{w}}$ -almost sure and the second $F_{W^*|X=x, \bar{W}=\bar{w}}$ -almost sure.

Assumption 2.i states that after conditioning on the treatments X and perfect controls W^* , any remaining variation in the proxy controls Z is independent of the heterogeneity in potential outcomes U . If the remaining variation in Z is interpreted as measurement error, then the measurement error must be independent of potential outcomes. Assumption 2.ii states that any dependence between V and (X, Z) is explained by their mutual association with the perfect controls W^* . We emphasize that the independence between V and (X, Z) in Assumption 2.ii is conditional. Without conditioning on the perfect controls W^* , V could be strongly associated with both X and Z . Again, note that neither Assumption 2.i nor 2.ii requires either V or Z be independent of W^* .

Assumptions 3.i and 3.ii state, loosely speaking, that both V and Z are sufficiently informative about the unobserved perfect controls W^* . The informativeness conditions are in terms of ‘completeness’, or more precisely, L_2 -completeness (Andrews (2017)). Completeness is used to achieve identification in the non-parametric instrumental variables (NPIV) models of Newey & Powell (2003) and Ai & Chen (2003). In the NPIV context, completeness is an instrumental relevance condition analogous to the rank condition for identification in linear IV (see Newey & Powell (2003) for discussion). With this interpretation, 3.i states that conditional on any given value of assigned treatments X , Z is a relevant instrument for W^* , and 3.ii states that conditioning on X , V is a relevant instrument for W^* . Some sufficient conditions for statistical completeness can be found in D’Haultfoeulle (2011) and Hu & Shiu (2018). In some settings L_2 -completeness is generic in a certain sense (Andrews (2017), Chen *et al.* (2014)).

In the linear IV case, the rank condition can only hold if the number of instruments exceeds the number of endogenous regressors, this is known as the

‘order condition’. Such a condition is not, strictly speaking, necessary for L_2 -completeness in nonparametric models. However, an order condition is necessary in certain special cases, for example the conditional Gaussian case discussed in [Newey & Powell \(2003\)](#). Therefore, in practice, it seems prudent to require that the order condition hold: that each of the vectors V and Z be of a weakly larger dimension than W^* .

In addition to Assumptions 1, 2 and 3 we require that either one of two regularity conditions hold. The conditions refer to sequences of functions $\{\mu_k\}_{k=1}^\infty$, $\{u_k\}_{k=1}^\infty$ and $\{v_k\}_{k=1}^\infty$ that depend on the joint distribution of Z , V and X . The definition of these functions is somewhat technical and so we relegate it to Appendix A.

Assumption 4.i refers to $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}$, the Radon-Nikodym derivative of $F_{V|X=x_2}$ with respect to $F_{V|X=x_1}$. This must exist under Assumptions 1.ii and 2.ii (see Proposition B.2 in Appendix B). If the distribution of V given X admits a conditional probability density function then this equals $f(v|x_1)/f(v|x_2)$ where $f(v|x)$ is the probability density of V at v conditional on $X = x$.

Assumption 4 (Regularity). For both 4.i and 4.ii below assume the following: Assumptions A.1, 1.ii, and 2.ii hold, and for $F_{\bar{W}}$ -almost all \bar{w} and F_X -almost all x_1 and x_2 , $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V)$ and $E[Y|X, Z]$ have finite mean squares conditional on $(X, \bar{W}) = (x_1, \bar{w})$.

i. For some function \tilde{C} with $E[\tilde{C}(x_1, x_2, \bar{W})|X = x_1] = C(x_1, x_2) < \infty$:

$$\sum_{k=1}^{\infty} \frac{1}{\mu_k(x_1, \bar{w})^2} E\left[\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V) u_k(X, V) | X = x_1, \bar{W} = \bar{w}\right]^2 \leq \tilde{C}(x_1, x_2, \bar{w})$$

ii. For some function \tilde{D} with $E[\tilde{D}(X, \bar{w})|X = x] = D(x) < \infty$:

$$\sum_{k=1}^{\infty} \frac{1}{\mu_k(x, \bar{w})^2} E\left[E[Y|X, Z] v_k(X, Z) | X = x, \bar{W} = \bar{w}\right]^2 \leq \tilde{D}(x, \bar{w})$$

Assumption 4 first imposes some previously stated assumptions and states that $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V)$ and $E[Y|X, Z]$ have finite conditional mean squares. This ensures the terms of the sums in 4.i and 4.ii are well-defined. 4.i and 4.ii each require that generalized Fourier coefficients go to zero sufficiently quickly. These restrictions can be understood as smoothness conditions (see [Hall & Horowitz \(2005\)](#)). Conditions of the same form are used elsewhere in the literature, for example to establish existence of an NPIV regression function in [Darolles *et al.* \(2011\)](#). [Miao *et al.* \(2018\)](#) employs a condition that closely resembles Assumption 4.ii for a similar purpose to us.

Theorem 1.1 (Identification). *Suppose Assumptions 1, 2, and 3 hold. Then:*

a. If 4.ii (and not necessarily 4.i) holds, there exists a function γ with $\gamma(x_1, \cdot) \in L_2(F_{V|X=x_1})$ for F_X -almost all x_1 , so that:

$$E[Y - \gamma(X, V) | X, Z] = 0 \tag{1.2}$$

And for any such γ , for F_X -almost all x_1 and x_2 :

$$\bar{y}(x_1|x_2) = E[\gamma(x_1, V)|X = x_2]$$

b. If 4.i (and not necessarily 4.ii) holds, there exists a function φ with $\varphi(x_1, x_2, \cdot) \in L_2(F_{Z|X=x_1})$ for F_X -almost all x_1 , so that:

$$E[\varphi(x_1, x_2, Z)|X = x_1, V] = \frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V) \quad (1.3)$$

And for any such φ , for F_X -almost all x_1 and x_2 :

$$\bar{y}(x_1|x_2) = E[Y\varphi(x_1, x_2, Z)|X = x_1]$$

Theorem 1.1 characterizes the CASF in terms of observables and thus establishes identification. Theorem 1.2 below establishes that the equations that characterize the CASF in 1.1.a and 1.1.b are well-posed.

Theorem 1.2 (Well-Posedness). *Suppose Assumptions 1, 2, and 3 hold. Then:*

a. *If 4.i (and not necessarily 4.ii) holds, for any $\tilde{\gamma}(x_1, \cdot) \in L_2(F_{V|X=x_1})$, and for F_X -almost all x_1 and x_2 :*

$$\begin{aligned} & (\bar{y}(x_1|x_2) - E[\tilde{\gamma}(x_1, V)|X = x_2])^2 \\ & \leq C(x_1, x_2) E \left[(E[Y - \tilde{\gamma}(X, V)|X, Z])^2 \middle| X = x_1 \right] \end{aligned} \quad (1.4)$$

b. *If 4.ii holds (and not necessarily 4.i) hold, for any function $\tilde{\varphi}(x_1, x_2, \cdot) \in L_2(F_{Z|X=x_1})$, and for F_X -almost all x_1 and x_2 :*

$$\begin{aligned} & (\bar{y}(x_1|x_2) - E[Y\tilde{\varphi}(x_1, x_2, Z)|X = x_1])^2 \\ & \leq D(x_1) E \left[\left(E[\tilde{\varphi}(X, x_2, Z)|X, V] - \frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V) \right)^2 \middle| X = x_1 \right] \end{aligned} \quad (1.5)$$

Theorems 1.1 and 1.2 suggest a two-step approach to estimation. Recall the moment condition in part a. of Theorem 1:

$$E[\gamma(X, V) - Y|X, Z] = 0 \quad (1.6)$$

The equation above is equivalent to a non-parametric instrumental variables (NPIV) moment condition in which V is the vector of endogenous regressors, X is a vector of exogenous regressors, and Z is a vector of instruments. Suppose $\hat{\gamma}$ solves an empirical analogue of the moment condition (1.6). In a second step Theorems 1.1 and 1.2 suggests we estimate the CASF by:

$$\bar{y}(x_1|x_2) \approx \hat{E}[\hat{\gamma}(x_1, V)|X = x_2]$$

where ' \hat{E} ' denotes some empirical analogue of the conditional expectation and the first-stage estimate $\hat{\gamma}$ is treated as non-random in the expectation. The

inequality in Theorem 1.2.a implies that if $\hat{\gamma}$ satisfies the population moment condition (1.6) with small error (in a mean-squared sense), then $E[\hat{\gamma}(x_1, V)|X = x_2]$ is close to the CASF $\bar{y}(x_1|x_2)$. If, in addition, $E[\hat{\gamma}(x_1, V)|X = x_2]$ is close to the sample analogue $\hat{E}[\hat{\gamma}(x_1, V)|X = x_2]$, then the latter provides a good estimate of the CASF. This motivates our estimator in Section 3.

The inequality in Theorem 1.2.a suggests that the estimation problem based on Theorem 1.1.a is well-posed in a certain sense. In short, if $\hat{\gamma}$ is close to satisfying the moment condition (1.6) then its conditional mean is close to the true CASF. Well-posedness allows us to derive simple convergence rates for our estimation method that are comparable to those in standard non-parametric regression and do not depend on any ‘sieve-measure of ill-posedness’ (Chen & Pouzo (2012)).

The well-posedness of our problem may be surprising because it is well-known that estimation of an NPIV regression function is generally ill-posed. However, we characterize the CASF as a linear functional (specifically a conditional mean) of an NPIV regression function. Estimation of a sufficiently smooth linear functional of an NPIV regression function is well-posed. In particular, the existence of a solution to (1.3) guarantees sufficient smoothness of the relevant linear functional. Because Assumption 4 ensures this existence, this assumption is crucial to the well-posedness. Existence conditions of a similar kind are shown to be closely related to root- n estimability of linear functionals of NPIV regression functions by Severini & Tripathi (2012) and Ichimura & Newey (2017).

One could also motivate an estimator based on part b. of Theorem 1.1. In particular one would first estimate $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(\cdot)$ and then solve for φ in an empirical analogue of equation (1.3). One would then plug the empirical solution $\hat{\varphi}$ into an empirical analogue of the conditional expectation $E[Y\hat{\varphi}(x_1, x_2, Z)|X = x_1]$, where as with $\hat{\gamma}$ above, $\hat{\varphi}$ is treated as non-random in the expectation.

2 Panel Models

The analysis in the previous section applies to the model (1.1) which may apply in both cross-sectional and panel settings. In panel settings, observations from previous periods are a natural source of proxy controls. Loosely speaking, if the same factors explain the confounding in each period (there are time-invariant perfect controls), then treatment assignments in other periods are informative about the confounding. Thus we can form vectors V and Z using treatments (and possibly outcomes) from different periods. Then the conditional independence restrictions in Assumptions 1.i and 2 can be understood in terms of the serial dependence structure. In this section we specify some forms of serial dependence so that our conditional independence restrictions apply for corresponding choices of V and Z . In panels the order condition places restrictions on the number of time periods.

In the panel setting, the data have a ‘time’ dimension and a ‘unit’ dimension. To apply our analysis in the panel case we rewrite the model (1.1) with time

subscripts as $Y_t = y_{0,t}(X_t, U_t)$. Then for each group there is an associated draw of the random variables (X_1, \dots, X_T) , (U_1, \dots, U_T) and a resulting sequence of outcomes (Y_1, \dots, Y_T) . This specification above allows for dynamic models with feedback, for example if X_t includes lags of Y_t .

In the panel setting our goal is to identify and estimate for a particular value of t , $E[y_{0,t}(x_1, U_t)|X_t = x_2]$. This is the conditional average potential outcome at period t from treatment x_1 conditional on assignment of treatment x_2 at t . In this context Assumptions 1.i, 2.i and 2.ii state that $U_t \perp\!\!\!\perp (X_t, Z)|W^*$ and $V \perp\!\!\!\perp (X_t, Z)|W^*$ ³

Below we present two cases in which the conditional independence restrictions above follow from more primitive conditions on the panel structure for appropriate choices of V and Z .

2.1 Markov Treatment Assignments and Predetermination

Suppose we are interested in the CASF at some fixed period t . Suppose that conditional on some (possibly period t -specific) latent variables \tilde{W}^* , the following conditional independence restriction holds:

$$U_t \perp\!\!\!\perp (X_1, \dots, X_t)|\tilde{W}^* \quad (2.1)$$

In words, the condition above states that the history of treatments up to and including period t is only related to potential outcomes through some factors \tilde{W}^* . If \tilde{W}^* is taken to represent some persistent latent factors, then the restriction is a non-parametric analogue of the ‘predetermination’ condition often employed in linear panel models. One justification for the assumption is as follows. Suppose we interpret \tilde{W}^* to contain all persistent factors in the potential outcomes. Then any remaining variation in U_t represents shocks to potential outcomes. In this case the assumption states that the history of treatments up to and including time t is uninformative about these shocks. However, the assumption allows for the possibility that shocks to potential outcomes impact (or are otherwise associated with) future treatment assignments.

Let $\lfloor t/2 \rfloor$ denote the largest natural number weakly less than half of t . Suppose that conditional on the latent variables \tilde{W}^* , the regressors satisfy a first-order Markov dependence structure at $\lfloor t/2 \rfloor$. Formally:

$$(X_1, \dots, X_{\lfloor t/2 \rfloor - 1}) \perp\!\!\!\perp (X_{\lfloor t/2 \rfloor + 1}, \dots, X_T)|(\tilde{W}^*, X_{\lfloor t/2 \rfloor}) \quad (2.2)$$

That is, conditional on the latent variables \tilde{W}^* , the treatment assignments for periods strictly prior to the given period $\lfloor t/2 \rfloor$ are only related to treatments after $\lfloor t/2 \rfloor$ through the treatment at $\lfloor t/2 \rfloor$.

Proposition 2.1 below shows that in this setting Assumptions 1.i and 2 hold for V and Z composed of particular lagged treatments.

³We have used that, $U_t \perp\!\!\!\perp (X_t, Z)|W^*$ is equivalent to the combination of $U_t \perp\!\!\!\perp X_t|W^*$ and $U_t \perp\!\!\!\perp Z|(X_t, W^*)$.

Proposition 2.1. *Suppose that (2.1) and (2.2) hold. Set $V = (X_1, \dots, X_{\lfloor t/2 \rfloor})$, $Z = (X_{\lfloor t/2 \rfloor}, \dots, X_{t-1})$, and $W^* = (\tilde{W}^*, X_{\lfloor t/2 \rfloor})$. Then Assumptions 1.i and 2 hold: $U_t \perp\!\!\!\perp (X_t, Z) | W^*$ and $V \perp\!\!\!\perp (X_t, Z) | W^*$.*

Note that we treat $X_{\lfloor t/2 \rfloor}$ as an observable perfect control, we therefore include it in both V and Z . Given the Markov structure, conditioning on the treatment at period $\lfloor t/2 \rfloor$ removes the dependence between V and Z .

2.2 Markov Treatment Assignments and Heterogeneity

We now give conditions under which Z and V may be composed not only of treatment assignments from periods other than t , but also the outcomes from other periods. We strengthen the conditional independence restriction from the previous subsection:

$$U_t \perp\!\!\!\perp (X_1, \dots, X_t, U_1, \dots, U_{t-1}) | \tilde{W}^* \quad (2.3)$$

The above states that any dependence between treatment assignments in all periods up to and including t and heterogeneity in potential outcomes in all periods up to and including t , is explained by the (possibly period- t specific) factors \tilde{W}^* .

We suppose that conditional on the latent variables \tilde{W}^* , both the treatment assignments and heterogeneity follow a first-order Markov dependence structure. Formally, conditional on $(\tilde{W}^*, X_{\lfloor t/2 \rfloor}, U_{\lfloor t/2 \rfloor})$:

$$(X_1, \dots, X_{\lfloor t/2 \rfloor - 1}, U_1, \dots, U_{\lfloor t/2 \rfloor - 1}) \perp\!\!\!\perp (X_{\lfloor t/2 \rfloor + 1}, \dots, X_t, U_{\lfloor t/2 \rfloor + 1}, \dots, U_t) \quad (2.4)$$

The following proposition shows that if we set V , Z and W^* much as in the previous subsection but now with lagged outcomes, then Assumptions 1.i and 2 hold.

Proposition 2.2. *Suppose that (2.3) and (2.4) hold. Suppose we set $W^* = (\tilde{W}^*, X_{\lfloor t/2 \rfloor}, Y_{\lfloor t/2 \rfloor})$ and set V and Z as follows:*

$$\begin{aligned} V &= (X_1, \dots, X_{\lfloor t/2 \rfloor}, Y_1, \dots, Y_{\lfloor t/2 \rfloor}) \\ Z &= (X_{\lfloor t/2 \rfloor}, \dots, X_{t-1}, Y_{\lfloor t/2 \rfloor}, \dots, Y_{t-1}) \end{aligned}$$

Then Assumptions 1.i and 2 hold: $U_t \perp\!\!\!\perp (X_t, Z) | W^$ and $V \perp\!\!\!\perp (X_t, Z) | W^*$.*

2.3 Assumption 3 and the Order Condition in Panels

Assumption 3 consists of two L_2 -completeness conditions. As discussed in Section 1, these conditions can be understood as instrumental relevance conditions: conditioning on X_t , both V and Z must be relevant instruments for the perfect controls W^* .

In the Markov treatment assignment case discussed above, both V and Z are likely to be strongly associated with the perfect controls $W^* = (\tilde{W}^*, X_{\lfloor t/2 \rfloor})$. Both Z and V contain $X_{\lfloor t/2 \rfloor}$ and in addition some treatment assignments for

periods other than t . Recall that by the predetermination condition \tilde{W}^* are latent variables that explain the confounding between (X_1, \dots, X_t) and U_t . If there is confounding in each period that is explained by the presence of those same variables \tilde{W}^* then each component of Z and V ought to be informative about \tilde{W}^* .

Note that if T is large then there are more observations from different periods from which to form V and Z . V and Z are then more likely to satisfy Assumption 3. As we discuss in Section 1, it is prudent to require that an order condition hold, i.e., that V and Z each be of a weakly larger dimension than W^* . In the first-order Markov treatment assignment example when treatments are scalar, Z is of length $t - \lfloor t/2 \rfloor$ and V is of length $\lfloor t/2 \rfloor$. Therefore, the order condition requires that \tilde{W}^* be of length at most $\lfloor t/2 \rfloor - 1$. If we are interested in the CASF at the final period T then \tilde{W}^* must be of length no greater than $\lfloor T/2 \rfloor - 1$.

3 Estimation

In this section we describe our estimation procedure and analyze its asymptotic properties. The key step in estimation corresponds to penalized sieve minimum distance (PSMD) estimation (see [Chen & Pouzo \(2012\)](#), [Chen & Pouzo \(2015\)](#)). Our estimator can be applied in panel settings or to cross-sectional data. To emphasize this generality we return to the notation in Section 1 in which we suppress time subscripts.

Let $\{(Y_i, X_i, Z_i, V_i)\}_{i=1}^n$ be a sample of n observations of the variables Y , X , Z , and V . In the panel case, Y_i and X_i should be understood to come from one fixed period t . For each n let ϕ_n be a column vector of basis functions defined on the support of (X, V) . The first stage of the procedure consists of non-parametric regression. The practitioner estimates regression functions g , π_n and α_n which are defined by:

$$\begin{aligned} g(x, z) &= E[Y|X = x, Z = z] \\ \pi_n(x, z) &= E[\phi_n(X, V)|X = x, Z = z] \\ \alpha_n(x_1, x_2) &= E[\phi_n(x_1, V)|X = x_2] \end{aligned}$$

The estimation of each of the functions above can be carried out using a standard non-parametric regression method like local-linear regression or series least-squares. Denote estimates of the fitted values $g(X_i, Z_i)$, $\pi_n(X_i, Z_i)$, and $\alpha_n(x_1, x_2)$ by \hat{g}_i , $\hat{\pi}_{n,i}$, and $\hat{\alpha}_n(x_1, x_2)$ respectively.

Let P be some penalty function. Let $\lambda_{0,n}$ be a positive scalar penalty parameter. In the second stage, the researcher evaluates a vector of coefficients $\hat{\theta}$ that minimize the penalized least-squares objective:

$$\frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i} \theta)^2 + \lambda_{0,n} P(\theta) \quad (3.1)$$

The estimate of the CASF is then given by:

$$\bar{y}(x_1|x_2) \approx \hat{\alpha}_n(x_1, x_2)' \hat{\theta} \quad (3.2)$$

Our consistency and convergence rate results pertain to a particular version of the method described above. This is also the procedure we implement in our empirical applications. We take P to be a ridge penalty (i.e., $P(\theta) = \|\theta\|_2^2$, where $\|\cdot\|_2$ is the Euclidean norm). Letting $\hat{\Sigma}_{\lambda_{0,n}} = \frac{1}{n} \sum_{i=1}^n \hat{\pi}_{n,i} \hat{\pi}'_{n,i} + \lambda_{0,n} I_n$, the minimizer of (3.1) then has a closed-form solution:

$$\hat{\theta} = \hat{\Sigma}_{\lambda_{0,n}}^{-1} \frac{1}{n} \sum_{i=1}^n \hat{\pi}_{n,i} \hat{g}_i \quad (3.3)$$

We assume the basis functions $\phi_n(x, v)$ are multiplicatively separable in x and v . In particular there are length $k(n)$ and $l(n)$ vectors of functions ρ_n and χ_n so that $\phi_n(x, v) = \rho_n(v) \otimes \chi_n(x)$ where ‘ \otimes ’ is the Kronecker product. The first-stage regressions are then carried out using series ridge regression with penalty parameters $\lambda_{1,n}$, $\lambda_{2,n}$, and $\lambda_{3,n}$. Let ψ_n be a length- $m(n)$ vector of basis functions defined on the support of (X, Z) and let $\psi_{n,i} = \psi_n(X_i, Z_i)$. Let $\rho_{n,i} = \rho_n(V_i)$ and $\chi_{n,i} = \chi_n(X_i)$. For any λ define $\hat{\Omega}_\lambda = \frac{1}{n} \sum_{i=1}^n \psi_{n,i} \psi'_{n,i} + \lambda I_{m(n)}$ and $\hat{G}_\lambda = \frac{1}{n} \sum_{i=1}^n \chi_{n,i} \chi'_{n,i} + \lambda I_{l(n)}$ (where I_k is the k -by- k identity matrix). We obtain estimates \hat{g}_i , $\hat{\pi}_{n,i}$, and $\hat{\alpha}_n(x_1, x_2)$ as follows:

$$\begin{aligned} \hat{g}_i &= \psi'_{n,i} \hat{\Omega}_{\lambda_{1,n}}^{-1} \frac{1}{n} \sum_{j=1}^n \psi_{n,j} Y_j \quad (3.4) \\ \hat{\pi}_{n,i} &= (\psi'_{n,i} \hat{\Omega}_{\lambda_{2,n}}^{-1} \frac{1}{n} \sum_{j=1}^n \psi_{n,j} \rho_{n,j}) \otimes \chi_{n,i} \\ \hat{\alpha}_n(x_1, x_2) &= (\chi_n(x_2)' \hat{G}_{\lambda_{3,n}}^{-1} \frac{1}{n} \sum_{j=1}^n \chi_{n,j} \rho_{n,j}) \otimes \chi_n(x_1) \end{aligned}$$

3.1 Consistency and Convergence Rate

We prove the consistency of our estimator and derive convergence rates under primitive conditions similar to those common in the literature on standard nonparametric regression. The well-posedness of our estimation problem plays a crucial role in our results. Well-posedness allows us to derive convergence rates that do not depend on the ‘sieve-measure of ill-posedness’ that is characteristic of PSMD estimation of in nonparametric instrumental variables (NPIV) models. Moreover, our smoothness assumptions pertain only to reduced-form objects, i.e., conditional probability densities and expectations involving only observables. This contrasts with the NPIV case, which typically requires smoothness of the structural function. We discuss intuition for the difference between the behavior of our PSMD estimator and NPIV estimators later in this section.

In order to state Assumptions 5.1, 5.2, 5.3, and 5.4 below, we introduce some additional notation. $\|\cdot\|_2$ is the Euclidean norm of a vector and $\|\cdot\|_{op}$ the corresponding operator norm of a matrix. For sequences of scalars a_n and b_n , $a_n \lesssim b_n$ if and only if the sequence $|a_n/b_n|$ is bounded, and $a_n \prec b_n$ if and only if $a_n \lesssim b_n$ but not $b_n \lesssim a_n$. For sequences of random scalars $a_n \lesssim_p b_n$

if and only if a_n/b_n is stochastically bounded and $a_n \prec_p b_n$ if and only if $a_n \lesssim_p b_n$ but not $b_n \lesssim_p a_n$.

For any $s, c > 0$ let $\Lambda_s^d(c)$ be the space of smooth functions defined as follows. For any vector $q \in \mathbb{N}_0^d$, let D_q be the partial derivative operator. That is, for any scalar function δ on \mathbb{R}^d , $D_q[\delta](r) = \frac{\partial^{\|q\|_1}}{\partial^{q_1} r_1 \partial^{q_2} r_2 \dots \partial^{q_{\dim(R)}} r_{\dim(R)}} \delta(r)$. Then $\delta \in \Lambda_s^d(c)$ if and only if, for any $q \in \mathbb{N}_0^d$ with $\|q\|_1 \leq \lfloor s \rfloor$, $D_q[\delta]$ exists and has magnitude bounded uniformly by c , and for all $\|q\|_1 = \lfloor s \rfloor$ and $r_1, r_2 \in \mathbb{R}^d$:

$$|D_q[\delta](r_1) - D_q[\delta](r_2)| \leq c \|r_1 - r_2\|_2^{s - \lfloor s \rfloor}$$

‘esssup’ denotes the essential supremum: $\text{esssup } R$ is the infimum of the set of scalars greater than random variable R with probability 1. Finally, we let $Q_n = E[\rho_{n,i} \rho'_{n,i}]$, $G_n = E[\chi_{n,i} \chi'_{n,i}]$, and $\Omega_n = E[\psi_{n,i} \psi'_{n,i}]$.

Assumption 5.1 (Bases). i. Q_n is non-singular with $\xi_{\rho,n} = \text{esssup } \|Q_n^{-1/2} \rho_{n,i}\|_2$ finite. For any $s > 0$ uniformly over any $c > 0$ and $\delta \in \Lambda_s^{\dim(V)}(c)$:

$$\inf_{\beta \in \mathbb{R}^{k(n)}} E[(\delta(V) - \rho_n(V)' \beta)^2]^{1/2} \lesssim c \ell_{\rho,n}(s)$$

ii. **Either** X has finite discrete support and for sufficiently large n any scalar function on \mathcal{X} is a linear transformation of χ_n , **or** G_n is non-singular with $\xi_{\chi,n} = \text{esssup } \|G_n^{-1/2} \chi_{n,i}\|_2$ finite, and for any $s > 0$, uniformly over all $c > 0$ and $\delta \in \Lambda_s^{\dim(X)}(c)$, if $\beta^*[\delta]$ minimizes $E[(\delta(X) - \chi_n(X)' \beta)^2]$ then:

$$\text{esssup } |\delta(X) - \chi_n(X)' \beta^*[\delta]| \lesssim c \ell_{\chi,n}(s)$$

iii. Ω_n is non-singular with $\xi_{\psi,n} = \text{esssup } \|\Omega_n^{-1/2} \psi_{n,i}\|_2$ finite, and for any $s > 0$, uniformly over all $c > 0$ and $\delta \in \Lambda_s^{\dim(X,Z)}(c)$, if $\beta^*[\delta]$ minimizes $E[(\delta(X, Z) - \psi_n(X, Z)' \beta)^2]$ then:

$$\text{esssup } |\delta(X, Z) - \psi_n(X, Z)' \beta^*[\delta]| \lesssim c \ell_{\psi,n}(s)$$

Assumption 5.2 (Densities, Conditional Variance). i. The joint distribution $F_{(X,Z,V)}$ is absolutely continuous with the product of the marginals $F_{(X,Z)} \otimes F_V$. ii. The Radon-Nikodym derivative $\frac{dF_{(X,Z,V)}}{dF_X \otimes F_V}$ is bounded above and away from zero. iii. X has finite discrete support **or** X is continuously distributed on support $\mathcal{X} \in \mathbb{R}^{\dim(X)}$, X admits a probability density f_X that is bounded above and away from zero, and there exist $b > 0$ and $\underline{r} > 0$, so that for any $x \in \mathcal{X}$ and $0 < b' \leq b$, $\text{vol}(B_{x,b'} \cap \mathcal{X}) \geq \underline{r} \text{vol}(B_{0,b'})$, where $B_{x,b'}$ is the Euclidean ball of radius b' centered at x and $\text{vol}(\cdot)$ returns the volume. iv. there exists $\bar{\sigma}_Y < \infty$ so that with probability 1, $E[Y^2 | X, Z] \leq \bar{\sigma}_Y^2$.

Assumption 5.3 (Smoothness). In each case, for $F_{(X,Z,V)}$ -almost all (x, z, v) , i. $\frac{dF_{(X,Z,V)}}{F_{(X,Z)} \otimes dF_V}(x, z, \cdot) \in \Lambda_{s_1}^{\dim(V)}(c_1)$ and $\frac{dF_{(X,Z,V)}}{F_{(X,Z)} \otimes dF_V}(\cdot, z, v) \in \Lambda_{s_2}^{\dim(X)}(c_2)$ ii. $g(\cdot, z) \in \Lambda_{s_3}^{\dim(X)}(c_3)$, iii. $\frac{dF_{(X,Z,V)}}{F_{(X,Z)} \otimes dF_V}(\cdot, \cdot, v) \in \Lambda_{s_4}^{\dim(X,Z)}(c_4)$, iv. Either X has finite discrete support or $\frac{dF_{(X,Z,V)}}{dF_X \otimes F_V}(\cdot, v) \in \Lambda_{s_5}^{\dim(X,Z)}(c_5)$.

Assumption 5.4 (Sieve Space Growth). i. $\frac{\xi_{\rho,n}^2 \log(k(n))}{n} \prec 1$, ii. $\frac{\xi_{\chi,n}^2 \log(l(n))}{n} \prec 1$, iii. $\frac{\xi_{\psi,n}^2 \log(m(n))}{n} \prec 1$

Assumption 5.1 specifies conditions on the basis functions. In particular, the assumption specifies the rate at which the basis functions can approximate smooth functions. Similar assumptions are ubiquitous in the literature on sieve regression and precise bounds for particular basis functions can be found in the approximation literature (see DeVore & Lorentz (1993)). For example, suppose X is continuous and the basis functions ρ_n , χ_n , and ψ_n are either spline series, local polynomial partition series, or Cohen–Daubechies–Vial wavelets, each of order $s_0 \geq s$. Then $\xi_{\psi,n}^2 \lesssim m(n)$, $\xi_{\rho,n}^2 \lesssim k(n)$, $\xi_{\chi,n}^2 \lesssim l(n)$, $\ell_{\psi,n}(s) \lesssim m(n)^{-s/\dim(X,Z)}$, $\ell_{\rho,n}(s) \lesssim k(n)^{-s/\dim(V)}$, and $\ell_{\chi,n}(s) \lesssim l(n)^{-s/\dim(X)}$ (see for example Chernozhukov *et al.* (2014)). In the case of X discrete, if ψ_n is a product basis of χ_n and order $s_0 \geq s$ splines, local polynomials, or wavelets for Z , then we achieve $\ell_{\psi,n}(s) \lesssim m(n)^{-s/\dim(X)}$.

Assumption 5.2.i is a weak condition on the joint distribution of V and (X, Z) , it is satisfied if, for example, V , X , and Z have non-zero joint probability density on a rectangular support. Assumption 5.2.ii holds if, for example, X and V have joint probability density that is bounded above and away from zero on a rectangular support. The condition on the support of X in the continuous case in 5.2.iii is a very weak regularity condition on the boundary. It holds, for example, if \mathcal{X} is rectangular. 5.2.iv is standard in the non-parametric regression literature and is trivially true if Y is bounded above and below.

Assumption 5.3 lists smoothness conditions on some reduced-form objects, primarily Radon-Nikodym derivatives of probability distributions of observables. If the relevant joint probability densities exist, then a Radon-Nikodym derivative is equal to a ratio of conditional probability densities. For example, if V admits a conditional probability density function, conditional on (X, Z) , then $\frac{dF_{(X,Z,V)}}{F_{(X,Z)} \otimes dF_V}(v, x, z) = \frac{f(v|x,z)}{f(v)}$ where $f(v|x, z)$ is the density of V at v conditional on (X, Z) equal to (x, z) and $f(v)$ is the marginal density of V at v . Thus, if the relevant densities exist the smoothness conditions in 5.3 are, apart from in the case of 5.3.ii, smoothness assumptions regarding ratios of probability densities. Recall that $g(x, z) = E[Y|X = x, Z = z]$ is a reduced-form regression function, smoothness restrictions on such an object are standard in the non-parametric regression literature.

Assumption 5.4 restricts the rate at which the numbers of basis functions can grow. This assumption allows us to apply Rudelson’s matrix law of large numbers (Rudelson (1999)).

Theorem 3.1 below is an immediate corollary to the more general result in Proposition 3.4 in Appendix B. The more general result allows for the possibility that the smoothness coefficients s_1, s_2, \dots, s_5 vary and that the bases χ_n and ψ_n differ. Moreover, the more general proposition does not assume the penalty parameters are set rate-optimally. Instead, Proposition 3.4 provides rates of convergence in terms of the penalty parameters. This greater generality is achieved at the expense of simplicity, which is why we relegate Proposition

3.4 to the appendix.

Theorem 3.1 (Convergence). Let Assumptions 1, 2, 3, and 5 hold and Assumption 4 with $D(x_1)$ and $C(x_1, x_2)$ bounded uniformly over F_X -almost all x_1 and x_2 . Suppose that the penalty parameters $\lambda_{0,n}$, $\lambda_{1,n}$, $\lambda_{2,n}$, and $\lambda_{3,n}$ are chosen rate-optimally. In addition, suppose $\ell_{\psi,n}(s) \lesssim 1$, $l(n) \gtrsim 1$, that the smoothness parameters s_1, s_2, \dots, s_5 are all equal to s and that the smallest and largest eigenvalues of Q_n and G_n are bounded above and away from zero uniformly over n . Then if X has finite and discrete support:

$$\begin{aligned} & |\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| \\ & \lesssim_p \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ & + \ell_{\psi,n}(s) + \ell_{\rho,n}(s) \end{aligned}$$

And otherwise, uniformly over F_X -almost all x_1 and x_2 :

$$\begin{aligned} & |\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| \\ & \lesssim_p \xi_{\chi,n}^2 \min\{\sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n}\} \\ & + \xi_{\chi,n} \xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ & + \xi_{\chi,n}^2 \ell_{\chi,n}(s) + \xi_{\Omega,n} \ell_{\psi,n}(s) + \xi_{\Omega,n} \ell_{\rho,n}(s) + \xi_{\Omega,n} (\xi_{\rho,n} \ell_{\chi,n}(1))^{\bar{s}} \end{aligned}$$

Where $\bar{s} = \min\{\frac{1}{2}, \frac{s}{s+1}\}$ and $\xi_{\Omega,n} = \text{ess sup} \|\tilde{\Omega}_n(X)\|_{op} \leq \xi_{\psi,n}$, where $\tilde{\Omega}_n(x) = E[\psi_{n,i} \psi'_{n,i} | X_i = x]^{1/2} \Omega_n^{-1/2}$.

For specific choices of basis functions and smoothness s , one can optimize the rates in Theorem 3.1. For example, consider the case discussed above in which all the basis functions other than those in ψ_n are splines, local polynomial partitions, or Cohen–Daubechies–Vial wavelets of order $s_0 \geq s$. $\psi_n(x, z)$ is a Kronecker product of those basis functions for the z argument and χ_n for the x argument. Suppose X has finite discrete support, then choosing $k(n)$, $m(n)$, and $l(n)$ to optimize the rate in Theorem 3.1 we get:

$$|\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| \lesssim_p n^{-\frac{s}{2s + \dim(Z) + \dim(V)}}$$

In both the discrete and the continuous cases, so long as $\ell_{\rho,n}(s)$, $\ell_{\chi,n}(s)$, $\ell_{\psi,n}(s)$ go to zero at a rate that is exponential in the smoothness s , then as $s \rightarrow \infty$ the optimal rate approaches the parametric rate of $n^{-1/2}$.

Discussion

The penalty parameters $\lambda_{0,n}$, $\lambda_{1,n}$, $\lambda_{2,n}$ and $\lambda_{3,n}$ that achieve the rates in Theorem 3.1 are selected as follows. The penalties $\lambda_{1,n}$, $\lambda_{2,n}$, and $\lambda_{3,n}$, which appear in the first-stage regressions, simply need to converge to zero sufficiently quickly so that the terms involving them are dominated. For instance, setting

$\lambda_{1,n} = \lambda_{2,n} = \lambda_{3,n} = 0$ optimizes the rate in Proposition 3.4. Note that we do not prove optimality of the rates we derive: it may be possible to derive a faster rate than that in Proposition 3.4 under the same assumptions. Therefore we cannot rule out that the rate in Proposition 3.4 is sub-optimal and that the optimal rate is not optimized by setting these penalties to zero. Furthermore, even if setting $\lambda_{1,n}$, $\lambda_{2,n}$, and $\lambda_{3,n}$ non-zero does not lead to any asymptotic improvements it may be helpful in finite samples. By contrast, the rate we provide in Proposition 3.4 requires that $\lambda_{0,n}$ not go to zero too quickly. The constraint on the rate at which $\lambda_{0,n}$ can go to zero is stated in full in Proposition 3.4 and the rate in Theorem 3.1 is achieved by letting the constraint bind.

While our estimator is of the PSMD type, our convergence rate results differ markedly from those of PSMD estimators of the structural function in NPIV models. In particular, our results do not depend on a sieve-measure of ill-posedness. The key difference between our estimation problem and NPIV estimation is that the latter is typically ill-posed, whereas Theorem 1.2 establishes that under our identifying assumptions, our estimation problem is well-posed. For some intuition let us first explain why an NPIV problem related to our procedure is ill-posed. Let γ satisfy the conclusion of Theorem 1.1.a. γ then plays the role of the structural function in an NPIV model in which Z is the instrument, V is an endogenous regressor, and X is an exogenous regressor. For a PSMD estimate of γ , one evaluates $\hat{\theta}$ that minimizes (3.1), and an estimate of $\gamma(x, v)$ is then $\phi_n(x, v)' \hat{\theta}$. This problem is ‘ill-posed’ in the sense that the difference between $\gamma(x, v)$ and $\phi_n(x, v)' \hat{\theta}$ may be very large even if $\hat{\theta}$ sets a population analogue of the objective in (3.1) close to zero. However, we do not wish to estimate γ we wish to estimate the conditional mean of γ , which equals $\bar{y}(x_1|x_2)$. The conditional mean of $\phi_n(x, v)' \hat{\theta}$ is $\alpha_n(x_1, x_2)' \hat{\theta}$. Theorem 1.2.a gives conditions that imply $\alpha_n(x_1, x_2)' \hat{\theta}$ is close to $\bar{y}(x_1|x_2)$ so long as $\hat{\theta}$ makes a population analogue of the objective in (3.1) small.

Note as well that, because γ is not our object of interest, we are able to avoid placing any smoothness restrictions on this function. This is fortunate because γ does not have a clear structural interpretation in our model, in contrast with NPIV. In fact, we only impose smoothness on reduced-form objects (e.g., conditional probability densities and conditional expectations involving only observables).

The well-posedness mentioned above relies on Theorem 1.2.a which requires Assumption 4.i but not 4.ii. However, Assumption 4.ii also plays a crucial role in our analysis. Assumption 4.ii implies that there is a γ that solves Theorem 1.1.a with a bounded norm. This allows us to show that if the penalty $\lambda_{n,0}$ goes to zero sufficiently slowly, then (under some additional assumptions) the mean square of $\phi_n(X, V)' \hat{\theta}$ grows slowly with n or is bounded. This enables us to control the difference between $\alpha(x_1, x_2)' \hat{\theta}$ and $\hat{\alpha}_n(x_1, x_2)' \hat{\theta}$, which is equal to the estimation error from regressing dependent variable $\phi_n(X, V)' \hat{\theta}$ on $\chi_n(X)$. The error from regression is (roughly speaking) large when the variance of the dependent variable is large, hence the need to keep the mean square of $\phi_n(X, V)' \hat{\theta}$ small.

4 Empirical Applications

We apply our methodology to real data. In order to emphasize the applicability of our approach to both cross-sectional and panel models we present two separate empirical settings. In our first application we use cross-sectional variation to estimate causal effects, and in the second application we exploit the panel structure of the data.

Details for the implementation of our estimation method (choice of basis functions, first-stage regression methods) can be found in Appendix A.3. In short, we use separable high-order polynomial bases and carry out the first-stage regressions using ridge and similarly use a ridge penalty in the minimum distance step.

4.1 Causal Impact of Grade Retention

Fruehwirth *et al.* (2016) examine the causal effect of being made to repeat a particular grade level on the cognitive development of US students. They use data from the ECLS-K panel study which contains panel data on the early cognitive development of US children. We use our methods to examine the effect of grade retention on the cognitive outcomes of children in the 1998-1999 kindergarten school year using cleaned data available with their paper. Following Fruehwirth *et al.* (2016), we take our outcome variables to be the tests scores in reading and math when aged approximately eleven. Also in line with Fruehwirth *et al.* (2016) our treatments are indicators for retention in kindergarten, ‘early’ (in first or second grade) and ‘late’ (in third or fourth grade). The cleaned data from Fruehwirth *et al.* (2016) contains only students who are retained at most once in the sample period and no students who skip a grade.

Estimation of the causal effect of grade retention is challenging because students that repeat a grade typically do so due to poor academic performance. Poor academic performance in the past may be associated with poor cognitive ability in the future, with retention status held fixed. Thus a difference of mean outcomes is likely overly negative compared to the true causal effect of retention.

The ECLS-K dataset contains scores that measure a student’s behavioral and social skills in kindergarten as well as the student’s performance in a range of cognitive tests at different ages. To account for the confounding Fruehwirth *et al.* (2016) estimate a latent factor model with a particular structure. They assume that all confounding between grade retention and potential future cognitive test scores is due entirely to the presence of three latent factors: cognitive ability, behavioral ability and general ability. Fruehwirth *et al.* (2016) then use test scores to recover the distribution of the latent factors and their loadings. They assume a particular multiplicative structure between the factors (which are time-invariant) and time-specific factor loadings in both their outcome and selection equations.

In our approach, W^* plays a role analogous to the latent factors in the analysis of Fruehwirth *et al.* (2016), and our approach is valid under similar

assumptions about independence in the noise of our test scores. However, we do not assume any particular factor structure.

We let the set of proxies V contain the student’s scores on the behavioral and social skills tests in kindergarten and Z contain the kindergarten cognitive test scores. [Fruehwirth *et al.* \(2016\)](#) note that these scores are from tests taken prior to any retention. We take the unobserved factors W^* to measure the student’s classroom success early in kindergarten.

Recall that Assumption 3 essentially requires that both the test scores Z and the behavioral scores V would each be relevant instruments for W^* . Z directly (but likely with bias and noise) measures academic success and so Z and W^* are likely closely associated. V measures behavioral and social skills which likely reflect brain development and impact learning, thus V and W^* are also plausibly closely associated. Each of Z and V are three dimensional, so an order condition in this context would require W^* comprise at most three latent factors.

In this context Assumptions 1.i requires that retention status and potential age-eleven test scores are only related due to mutual dependence on underlying kindergarten academic success. Assumption 2.i requires that a student’s kindergarten test scores are only related to the student’s potential age-eleven test scores through the mutual dependence on classroom success and retention status. This is plausible if individual-specific noise in Z results from day of the test factors that are unlikely to be associated with potential outcomes.

Assumption 2.ii requires that the student’s behavioral scores are only related to cognitive test scores and retention status through mutual association with the student’s underlying academic success in kindergarten. This is plausible if noise in Z results from exogenous day-of-test factors and retention decisions depend only on teachers’ and parents’ assessments of the child’s classroom success W^* as well as factors like attitudes toward retention which are (other than through W^*) independent of the behavioral scores V . Note that our assumptions allow for the possibility that retention decisions X depend directly on Z .

We apply the estimation method set out in Section 3 to estimate average effects of retention at different grades. Table 1 below presents our results, and corresponds roughly to Table 4 in [Fruehwirth *et al.* \(2016\)](#).

We estimate that retention in kindergarten, in first or second grade, and in third or fourth grade raises the average scores for reading and math for those students who were in fact retained at these ages. In other words, the average effect of treatment on the treated (ATT) is estimated to be positive in all cases. This contrasts with [Fruehwirth *et al.* \(2016\)](#) who estimate mostly negative ATTs.

At present, we lack theoretical results that establish valid inference based on the standard errors given in Table 1. Nonetheless, it is worth noting that if one were to perform inference in the usual way with these standard errors, none of our estimates effects would be statistically significant at the 90% level.

The first column in each table gives the counterfactual effects of retention at different ages for those students who were not retained at any of the ages covered in our data. In all cases we find that those students who were not

Table 1: Effects of Grade Retention on Cognitive Performance

(a) Reading Ability				
$n = 1998$				
Difference from non-retention:	Observed retention status:			
	Not retained	Retained kindergarten	Retained early	Retained late
Retained kindergarten	-0.07 (0.12)	0.07 (0.14)	-0.05 (0.30)	0.14 (0.27)
Retained early	-0.07 (0.13)	0.11 (0.26)	0.01 (0.15)	-0.05 (0.24)
Retained late	-0.33 (0.14)	-0.25 (0.43)	-0.32 (0.43)	0.15 (0.16)

(b) Math Ability				
$n = 1999$				
Difference from non-retention:	Observed retention status:			
	Not retained	Retained kindergarten	Retained early	Retained late
Retained kindergarten	-0.12 (0.14)	0.13 (0.19)	0.06 (0.29)	0.15 (0.24)
Retained early	-0.17 (0.16)	0.12 (0.34)	0.11 (0.17)	-0.02 (0.30)
Retained late	-0.43 (0.22)	-1.30 (0.67)	-1.38 (0.75)	0.02 (0.15)

Proxy control estimates of the treatment effects for groups with different treatment statuses. The sample size n differs because for some individuals not all three outcomes are available. Numbers in parentheses are standard errors calculated as the standard deviation of the estimates over 1000 nonparametric bootstrap draws

retained would not have performed better had they been retained. Again, the contrasts with [Fruehwirth *et al.* \(2016\)](#) who, perhaps surprisingly, find positive effects in all cases.

In all, our results paint a more positive picture of schools' retention policies than those of [Fruehwirth *et al.* \(2016\)](#). In nearly all cases they find those who are retained on average suffered as a result and those who were not retained would have benefited from retention. We find the opposite: the retained students benefited from their retention and those not retained would not have benefited. However, the large bootstrapped standard errors suggest that our results are imprecise and (although presently we do not have results for valid inference) do not appear statistically significant.

4.2 Structural Engel Curve for Food

A household's Engel curve for a particular class of good captures the relationship between the share of the household's budget spent on that class and the total expenditure of the household. An Engel curve is 'structural' if it captures the effect of an exogenous change in total expenditure. Imagine an ideal experiment in which the household's total expenditure is chosen by a researcher using

a random number generator and the household then chooses how to allocate that total expenditure between different classes of goods. Then the resulting relationship between the total expenditure and budget share is a structural Engel curve.

A nonparametric regression of the budget share spent on food and the total expenditure spent on certain classes of goods is unlikely to represent the average structural Engel curve for food. This is because total expenditure is chosen by the household and thus depends upon the household’s underlying consumption preferences, but these same preferences partially determine the household’s expenditure on food.

We estimate average structural Engel curves for food eaten at home using data from the Panel Study of Income Dynamics (PSID). The PSID study follows US households over a number years and record expenditure on various classes of goods. We use ten periods of data from the surveys carried out every two years between 1999 and 2017 inclusive. We drop all households whose household heads are not married or cohabiting and drop all households for which we lack the full ten periods of data for all variables included in our analysis, leaving us with a sample of 840 households. We take as the total expenditure the sums of expenditures on food (both at home and away from home), housing, utilities, transportation, education, childcare and healthcare.

We apply the approach to identification with fixed- T panels described in Section 2, in particular the Markov treatment assignment and predetermination case. Let X_t denote the total expenditure in period t , which is our treatment of interest in this setting. We aim to estimate the average and conditional average structural Engel curve for period T . Let \tilde{W}^* consist of factors that capture heterogeneity in household preferences.

In this setting, the Markov treatment assumption requires that total expenditure in the past and in the future are only related through the expenditure today and the household’s consumption preferences. Conditioning on today’s expenditure X_t is important because conditioning on preferences household, expenditure may still be serially correlated due to persistence in household assets. Furthermore, total expenditure may reflect wages and labor supply which may be serially correlated even controlling for preferences.


The assumption of predetermination at time T is can be stated formally as $U_T \perp\!\!\!\perp (X_1, \dots, X_T) | \tilde{W}^*$. Here U_T represents period- T -specific variation in the household’s tendency to allocate a percentage of total expenditure onto consumption of food at home. Thus the above states that time-varying preferences for spending money on food are only related to the history of total expenditure through the mutual association with the underlying time-invariant preferences of the household \tilde{W}^* .

Given these assumptions we set V and Z in line with the suggestions in Section 2. That is, $V = (X_1, \dots, X_5)$ and $Z = (X_5, \dots, X_9)$. To justify Assumption 3 in this setting consider that underlying household preferences \tilde{W}^* presumably impact expenditure in every period (not just period T) and thus are associated with V and Z which are composed of past treatments. The order condition here requires that the dimension of preferences \tilde{W}^* be no greater than four.

Sub-Figure 1.a plots our nonparametric estimate of the average over our sample of the structural Engel curve for food. The sub-figure shows a downward-sloping Engel curve that (with a log scale for total expenditure) is subtly concave. The downward slope of the curve suggests that food is a normal good, at least in aggregate.

At present, we lack results establishing the validity of the confidence bands in Figure 1. We include them nonetheless because they may be informative about the variability of our estimates.

Figure 1: Demand for Food



Estimates are plotted at 100 points evenly spaced (in levels not logs) between the 10% and 90% quantiles of total expenditure. For Sub-Figure 1.a we estimate $\frac{1}{n} \sum_{i=1}^n \bar{y}(x, X_i)$ at each point x on the grid. For Sub-Figure 1.b we estimate $\bar{y}(1.1x, x) - \bar{y}(x, x)$ for each x . The confidence bands are uniform and evaluated using 1000 bootstrap draws and with pointwise standard errors equal to pointwise standard deviations of the estimated over the bootstrap draws.

Sub-Figure 1.b presents estimates of the average change in the budget share of food from an exogenous 10% increase in total expenditure broken down by the observed total expenditure. This is the difference of two conditional average structural Engel curves. In all cases the estimated change in expenditure share is negative, which again would be true of a normal good.

Conclusion

We present new results on identification and estimation with proxy controls. Our analysis applies in cross-sections and can be specialized to panel models. We apply our methodology to an empirical setting that based on cross-sectional variation and panel variation.

The present work raises questions for future research. In particular, the properties of our estimation method under misspecification have yet to be analyzed. The well-posedness of our problem suggests the sensitivity to a failure of our

conditional independence assumptions may be limited, but the well-posedness depends crucially on Assumption 4 and the implications of a failure of this assumption are less clear. A related topic is the behavior of our approach when the number of proxy controls is allowed to grow with the sample size, and $T \rightarrow \infty$ in the panel case. We plan to explore these avenues in future research.

References

- Ai, Chunrong, & Chen, Xiaohong. 2003. Efficient estimation of models with conditional moment restrictions containing unknown functions. *Econometrica*, **71**(6), 1795–1843.
- Andrews, Donald WK. 2017. Examples of L2-complete and boundedly-complete distributions. *Journal of Econometrics*, **199**(2), 213–220.
- Arellano, Manuel, & Bond, Stephen. 1991. Some tests of specification for panel data: Monte Carlo evidence and an application to employment equations. *The review of economic studies*, **58**(2), 277–297.
- Arellano, Manuel, & Bonhomme, Stéphane. 2016. Nonlinear panel data estimation via quantile regressions. *The Econometrics Journal*, **19**, C61–C94.
- Blundell, Richard, Chen, Xiaohong, & Kristensen, Dennis. 2007. Semiparametric IV estimation of shape-invariant Engel curves. *Econometrica. Journal of the Econometric Society*, **75**(6), 1613–1669.
- Chen, Xiaohong, & Pouzo, Demian. 2012. Estimation of Nonparametric Conditional Moment Models with Possibly Nonsmooth Generalized Residuals. *Econometrica*.
- Chen, Xiaohong, & Pouzo, Demian. 2015. Sieve Wald and QLR Inferences on Semi/Nonparametric Conditional Moment Models. *Econometrica*, **83**, 1013–1079.
- Chen, Xiaohong, Chernozhukov, Victor, Lee, Sokbae, & Newey, Whitney K. 2014. Local Identification of Nonparametric and Semiparametric Models. *Econometrica*.
- Chernozhukov, Victor, Chetverikov, Denis, & Kato, Kengo. 2014. Anti-concentration and honest, adaptive confidence bands. *The Annals of Statistics*, **42**, 1787–1818.
- Chernozhukov, Victor, Fernández-Val, Iván, Hoderlein, Stefan, Holzmann, Hajo, & Newey, Whitney. 2015. Nonparametric identification in panels using quantiles. *Journal of Econometrics*, **188**, 378–392.
- Darolles, Serge, Fan, Yanqin, Florens, Jean-Pierre, & Renault, Eric. 2011. Nonparametric Instrumental Regression. *Econometrica*.

- DeVore, Ronald A., & Lorentz, George G. 1993. *Constructive Approximation*. Springer-Verlag.
- D'Haultfoeuille, Xavier. 2011. On the completeness condition in nonparametric instrumental problems. *Econometric Theory*, **27**(3), 460–471.
- Evdokimov, Kirill. 2009. *Identification and Estimation of a Nonparametric Panel Data Model with Unobserved Heterogeneity*.
- Freyberger, Joachim. 2018. Non-parametric panel data models with interactive fixed effects. *Review of Economic Studies*, **85**(3), 1824–1851.
- Fruehwirth, Jane Cooley, Navarro, Salvador, & Takahashi, Yuya. 2016. How the Timing of Grade Retention Affects Outcomes: Identification and Estimation of Time-Varying Treatment Effects. *Journal of Labor Economics*, **34**, 979–1021.
- Hall, Peter, & Horowitz, Joel L. 2005. Nonparametric methods for inference in the presence of instrumental variables. *The Annals of Statistics*, **33**, 2904–2929.
- Hausman, Jerry A., & Taylor, William E. 1981. Panel Data and Unobservable Individual Effects. *Econometrica*, **49**, 1377.
- Holtz-Eakin, Douglas, Newey, Whitney, & Rosen, Harvey S. 1988. Estimating Vector Autoregressions with Panel Data. *Econometrica*, **56**, 1371.
- Hu, Yingyao, & Schennach, Susanne M. 2008. Instrumental Variable Treatment of Nonclassical Measurement Error Models. *E*, **76**, 195–216.
- Hu, Yingyao, & Shiu, Ji-Liang. 2018. Nonparametric identification using instrumental variables: sufficient conditions for completeness. *Econometric Theory*, **34**(3), 659–693.
- Ichimura, Hidehiko, & Newey, Whitney K. 2017. *The influence function of semiparametric estimators*.
- Jimerson, Shane R. 2001. Meta-analysis of grade retention research: Implications for practice in the 21st century. *School psychology review*, **30**(3), 420–437.
- Kress, Rainer. 1999. *Linear Integral Equations*. Springer.
- Kuroki, Manabu, & Pearl, Judea. 2014. Measurement Bias and Effect Restoration in Causal Inference. *Biometrika*.
- Lewbel, Arthur. 2008. Engel Curves. In: *The New Palgrave Dictionary of Economics, 2nd Edition*. Palgrave.
- Miao, Wang, Geng, Zhi, & Tchetgen, Eric J. Tchetgen. 2018. Identifying causal effects with proxy variables of an unmeasured confounder. *Biometrika*, **105**, 987–993.

- Newey, Whitney K., & Powell, James L. 2003. Instrumental Variable Estimation of Nonparametric Models. *Econometrica*, **71**, 1565–1578.
- Rokkanen, Miikka AT. 2015. *Exam schools, ability, and the effects of affirmative action: Latent factor extrapolation in the regression discontinuity design*. Working paper.
- Rudelson, M. 1999. Random Vectors in the Isotropic Position. *Journal of Functional Analysis*, **164**, 60–72.
- Severini, Thomas A., & Tripathi, Gautam. 2012. Efficiency bounds for estimating linear functionals of nonparametric regression models with endogenous regressors. *Journal of Econometrics*, **170**, 491–498.
- Sunder, Viakalathur Shankar. 2016. *Operators on Hilbert space*. Vol. 71. Springer.

Appendix A.1: Results Relating to Assumption 4

Assumption 4 refers to sequences of functions $\{\mu_k\}_{k=1}^\infty$, $\{u_k\}_{k=1}^\infty$ and $\{v_k\}_{k=1}^\infty$. In this Appendix we define these functions and their properties under Assumption A.1. We also state and prove Lemma A.1 which is used in our proofs of Theorems 1.1 and 1.2.

First, let us introduce some additional notation. The vectors Z and V may share some common components. Denote the shared components by \bar{W} and let \tilde{V} and \tilde{Z} contain the remaining components of V and Z respectively. Thus we can decompose $V = (\tilde{V}, \bar{W})$, $Z = (\tilde{Z}, \bar{W})$.

For each (x, \bar{w}) in the support of (X, \bar{W}) define a linear operator $A_{x, \bar{w}} : L_2(F_{\tilde{V}|X=x, \bar{W}=\bar{w}}) \rightarrow L_2(F_{\tilde{Z}|X=x, \bar{W}=\bar{w}})$ by:

$$A_{x, \bar{w}}[\delta](\tilde{z}) = E[\delta(\tilde{V})|\tilde{Z} = \tilde{z}, X = x, \bar{W} = \bar{w}]$$

The adjoint of this linear operator $A_{x, \bar{w}}^* : L_2(F_{\tilde{Z}|X=x, \bar{W}=\bar{w}}) \rightarrow L_2(F_{\tilde{V}|X=x, \bar{W}=\bar{w}})$ is given by:

$$A_{x, \bar{w}}^*[\delta](\tilde{v}) = E[\delta(\tilde{Z})|\tilde{V} = \tilde{v}, X = x, \bar{W} = \bar{w}]$$

Assumption A.1 below implies that the operators $A_{x, \bar{w}}$ and $A_{x, \bar{w}}^*$ are ‘compact’ (map bounded sets of functions into relatively compact sets). Compactness of a linear operator implies the existence of a singular system for that operator. Thus the assumption allows us to define singular systems indexed by (x, \bar{w}) for each operator $A_{x, \bar{w}}$ and $A_{x, \bar{w}}^*$. Singular systems of linear operators in functional analysis are analogous to the singular value decompositions of matrices in standard linear algebra. For background on singular value decompositions of linear operators between function spaces see [Kress \(1999\)](#).

Assumption A.1 (Compact Operator). The following holds for $F_{(X, \bar{W})}$ -almost all (x, \bar{w}) . Let ‘ F_{prod} ’ denote the product measure of \tilde{V} and \tilde{Z} conditional on $(X, \bar{W}) = (x, \bar{w})$.⁴ The conditional joint measure $F_{(\tilde{V}, \tilde{Z})|X=x, \bar{W}=\bar{w}}$ is absolutely continuous with respect to F_{prod} and the corresponding Radon-Nikodym derivative is square integrable with respect to F_{prod} :

$$\int \left(\frac{dF_{(\tilde{V}, \tilde{Z})|X=x, \bar{W}=\bar{w}}}{dF_{prod}}(\tilde{v}, \tilde{z}) \right)^2 F_{prod}(d\tilde{v}, d\tilde{z}) < \infty$$

Under Assumption A.1 for $F_{(X, \bar{W})}$ -almost all (x, \bar{w}) there exists a unique singular system (indexed by (x, \bar{w})) $\{(u_k^{(x, \bar{w})}, v_k^{(x, \bar{w})}, \mu_k^{(x, \bar{w})})\}_{k=1}^{\infty}$ for $A_{x, \bar{w}}$. $\mu_k^{(x, \bar{w})}$ is the k^{th} singular value of $A_{x, \bar{w}}$. $u_k^{(x, \bar{w})}$ is a real-valued function that maps from the support of \tilde{V} conditional on $(X, \bar{W}) = (x, \bar{w})$. $v_k^{(x, \bar{w})}$ is a scalar valued function that maps from the support of \tilde{Z} conditional on $(X, \bar{W}) = (x, \bar{w})$. $u_k^{(x, \bar{w})}$ and $v_k^{(x, \bar{w})}$ are known as the k^{th} singular functions of the operator $A_{x, \bar{w}}$.⁵

The functions $\{(u_k, v_k, \mu_k)\}_{k=1}^{\infty}$, referred to in Assumption 4, are then defined as follows. For each k , and (x, \bar{w}) , $u_k(x, v) = u_k^{(x, \bar{w})}(\tilde{v})$ where $v = (v, \bar{w})$ (recall we can decompose $V = (\tilde{V}, \bar{W})$). Similarly $v_k(x, z) = v_k^{(x, \bar{w})}(\tilde{z})$ where $z = (z, \bar{w})$ and $\mu_k(x, \bar{w}) = \mu_k^{(x, \bar{w})}$.

The following lemma is an application of Theorems 15.16 and 15.18 (Picard) in [Kress \(1999\)](#).

Lemma A.1. *Suppose Assumptions 1, 2 and 3 and Assumption A.1 hold. Then:*

1., *Assumption 4.i implies that there exists a function φ so that for F_X -almost all x_1 and x_2 , $E[\varphi(x_1, x_2, Z)^2|X = x_1] \leq C(x_1, x_2)$ with $C(x_1, x_2) < \infty$ and:*

$$E[\varphi(x_1, x_2, Z)|X = x_1, V] = \frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V)$$

2., *4.ii implies that there exists a function γ with $E[\gamma(X, V)^2|X = x] \leq D(x)$ with $D(x) < \infty$ for F_X -almost all x and:*

$$E[\gamma(X, V)|X, Z] = E[Y|X, Z]$$

Proof. First, we restate the conclusions of the lemma in terms of the existence of solutions to operator equations involving $A_{x_1, \bar{w}}^*$ and $A_{x, \bar{w}}$.

Suppose that for $F_{\bar{W}}$ -almost all \bar{w} and F_X -almost all x_1 and x_2 the function $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w}))$ is in the range of $A_{x_1, \bar{w}}^*$. That is, there exists a function $\tilde{\varphi}_{x_1, x_2, \bar{w}} \in L_2(F_{\tilde{Z}|X=x_1, \bar{W}=\bar{w}})$ so that:

$$E[\tilde{\varphi}_{x_1, x_2, \bar{w}}(\tilde{Z})|X = x_1, \bar{W} = \bar{w}, \tilde{V} = \tilde{v}] = \frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\tilde{v}, \bar{w}))$$

⁴In more conventional notation F_{prod} is equal to $F_{\tilde{V}|X=x, \bar{W}=\bar{w}} \otimes F_{\tilde{Z}|X=x, \bar{W}=\bar{w}}$.

⁵See, e.g., [Kress \(1999\)](#) Theorem 15.16 and associated discussion.

For $F_{\tilde{V}|X=x_1, \bar{W}=\bar{w}}$ -almost all \tilde{v} . Further suppose that for $F_{\bar{W}|X=x_1}$ -almost all \bar{w} , has $\tilde{\varphi}_{x_1, x_2, \bar{w}}$ has $L_2(F_{\tilde{Z}|X=x_1, \bar{W}=\bar{w}})$ -norm bounded by $\sqrt{\tilde{C}(x_1, x_2, \bar{w})}$, that is:

$$E[\tilde{\varphi}_{x_1, x_2, \bar{w}}(\tilde{Z})^2 | X = x_1, \bar{W} = \bar{w}] \leq \tilde{C}(x_1, x_2, \bar{w})$$

and $E[\tilde{C}(x_1, x_2, \bar{W}) | X = x_1] \leq C(x_1, x_2)$. Then setting $\varphi(x_1, x_2, (\tilde{z}, \bar{w})) = \tilde{\varphi}_{x_1, x_2, \bar{w}}(\tilde{z})$ we see conclusion 1. of the lemma holds.

Similarly, the following implies conclusion 2. For $F_{\bar{W}}$ -almost all \bar{w} and F_X -almost all x the following conditions hold. The function $E[Y|Z = (\cdot, \bar{w}), X = x]$ is in the range of $A_{x, \bar{w}}$ and that the solution $\tilde{\gamma}_{x, \bar{w}}$ has $L_2(F_{\tilde{V}|X=x, \bar{W}=\bar{w}})$ -norm bounded by $\sqrt{\tilde{D}(x, \bar{w})}$ and $E[\tilde{D}(x, \bar{w}) | X = x] \leq D(x)$.

To establish the relevant functions are in the ranges of the desired operators we apply Theorem 15.18 (Picard) in [Kress \(1999\)](#). We restate the theorem below and then show its premises hold under our assumptions.

Theorem 15.18 (Picard) in [Kress \(1999\)](#) states the following. Let $T : H_1 \rightarrow H_2$ be a compact linear operator from a Hilbert space H_1 to a Hilbert space H_2 with singular system $\{(u_k^{(T)}, v_k^{(T)}, \mu_k^{(T)})\}_{k=1}^\infty$. Then a function $\delta \in H_2$ is in the range of T (i.e., there exists a function $f \in H_1$ with $T[f] = \delta$) if and only if δ is in the orthogonal complement of the null space of the adjoint T^* and for some finite scalar c :

$$\sum_{k=1}^{\infty} \frac{1}{(\mu_k^{(T)})^2} |\langle \delta, v_k^{(T)} \rangle|^2 \leq c \quad (1)$$

In which case the solution f with smallest norm has norm \sqrt{c} . The bilinear form $\langle \cdot, \cdot \rangle$ is the inner product associated with the Hilbert space H_2 .

The operator $A_{x, \bar{w}}$ maps from the Hilbert space $L_2(F_{\tilde{V}|X=x, \bar{W}=\bar{w}})$ to the space $L_2(F_{\tilde{Z}|X=x, \bar{W}=\bar{w}})$, the latter of which has inner product given by:

$$\langle f_1, f_2 \rangle = E[f_1(\tilde{Z})f_2(\tilde{Z}) | X = x, \bar{W} = \bar{w}]$$

The adjoint $A_{x, \bar{w}}^*$ maps from $L_2(F_{\tilde{Z}|X=x, \bar{W}=\bar{w}})$ to $L_2(F_{\tilde{V}|X=x, \bar{W}=\bar{w}})$, the latter of which has inner product given by:

$$\langle f_1, f_2 \rangle = E[f_1(\tilde{V})f_2(\tilde{V}) | X = x, \bar{W} = \bar{w}]$$

Thus to apply Picard's theorem and show $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w}))$ is in the range of $A_{x_1, \bar{w}}^*$ so that there is a solution with norm weakly less than $\sqrt{\tilde{C}(x_1, x_2, \bar{w})}$ we need to show the following. a. $A_{x_1, \bar{w}}^*$ is compact and has a singular system, b. that $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w}))$ is an element of $L_2(F_{\tilde{V}|X=x_1, \bar{W}=\bar{w}})$, and c. that $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w}))$ is in the orthogonal complement of $A_{x_1, \bar{w}}$. Finally, we must show that d. (1) holds for $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w}))$ in place of δ and for the relevant Hilbert space and linear operator $A_{x_1, \bar{w}}^*$ and with c bounded by $\tilde{C}(x_1, x_2, \bar{w})$.

Similarly, to apply Picard's theorem and show $E[Y|Z = (\cdot, \bar{w}), X = x]$ is in the range of $A_{x, \bar{w}}$ so that there is a solution with norm bounded by $\sqrt{\tilde{D}(x, \bar{w})}$, we need to show that e. $A_{x, \bar{w}}$ is compact and has a singular system, f. that $E[Y|Z = (\cdot, \bar{w}), X = x]$ is an element of $L_2(F_{\tilde{Z}|X=x, \bar{W}=\bar{w}})$ and g. that $E[Y|Z = (\cdot, \bar{w}), X = x]$ is in the orthogonal complement of $A_{x, \bar{w}}^*$. And further, h. that (1) holds for $E[Y|Z = (\cdot, \bar{w}), X = x]$ in place of δ and for the relevant Hilbert space and linear operator $A_{x, \bar{w}}$ and with c bounded by $\tilde{D}(x, \bar{w})$

First we show that Assumptions A.1 implies that the operator $A_{x, \bar{w}}$ and its adjoint $A_{x, \bar{w}}^*$ are compact and therefore have unique singular systems (points a. and e. above). To see this first note that:

$$\begin{aligned} A_{x, \bar{w}}[\delta](\tilde{z}) &= E[\delta(\tilde{V})|\tilde{Z} = \tilde{z}, X = x, \bar{W} = \bar{w}] \\ &= \int \frac{dF_{(\tilde{V}, \tilde{Z})|X=x, \bar{W}=\bar{w}}}{dF_{prod}}(\tilde{v}, \tilde{z})\delta(\tilde{w}^*)F_{\tilde{V}|X=x, \bar{W}=\bar{w}}(d\tilde{v}) \end{aligned}$$

Thus $A_{x, \bar{w}} : L_2(F_{\tilde{V}|X=x, \bar{W}=\bar{w}}) \rightarrow L_2(F_{\tilde{Z}|X=x, \bar{W}=\bar{w}})$ is an integral operator with kernel $\frac{dF_{(\tilde{V}, \tilde{Z})|X=x, \bar{W}=\bar{w}}}{dF_{prod}}$ and Assumption A.1 states that the kernel is square integral with respect to the product measure of $F_{\tilde{V}|X=x, \bar{W}=\bar{w}}$ and $F_{\tilde{Z}|X=x, \bar{W}=\bar{w}}$. This implies that the operator $A_{x, \bar{w}}$ is Hilbert-Schmidt and therefore compact (see for example Section 3.3.1 of [Sunder \(2016\)](#)). Compactness of an operator implies compactness of its adjoint (alternatively we could simply repeat the steps above for $A_{x, \bar{w}}^*$). If $A_{x, \bar{w}}$ is compact then by Theorem 15.16 of [Kress \(1999\)](#) it admits a singular system. Note that the singular system of the adjoint $A_{x, \bar{w}}^*$ is the same as for $A_{x, \bar{w}}$ but with the roles of the singular functions $u_k^{(x, \bar{w})}$ and $v_k^{(x, \bar{w})}$ switched for each k . Thus we have shown a. and e. hold.

Next note that the first part of Assumptions 4.i states that

$$E \left[\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\tilde{V}, \bar{W}))^2 \middle| X = x_1 \right] < \infty$$

or in other words:

$$\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w})) \in L_2(F_{\tilde{V}|X=x_1, \bar{W}=\bar{w}})$$

Similarly the first part of 4.ii is equivalent to $E[Y|Z = (\cdot, \bar{w}), X = x] \in L_2(F_{\tilde{Z}|X=x_1, \bar{W}=\bar{w}})$. Thus 4.i and 4.ii imply point b. and f. respectively.

Now let us show that under Assumptions 1 and 2, c. and g. hold, that is the functions $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w}))$ and $E[Y|Z = (\cdot, \bar{w}), X = x]$ are in the orthogonal complements of the null spaces of operators $A_{x_1, \bar{w}}$ and $A_{x, \bar{w}}^*$ respectively.

Under Assumption 2.ii, for F_X -almost all x_1 and x_2 , and F_V -almost all v :

$$E \left[\frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}(W^*) \middle| X = x_1, V = v \right] = \frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(v)$$

For intermediate steps that show the above see (.2) in the proof of Theorem 1.1 in Appendix B. Now, let a function δ be in the null space of $A_{x_1, \bar{w}}$, that is $F_{\tilde{Z}|X=x_1, \bar{W}=\bar{w}}$ -almost surely:

$$E[\delta(\tilde{V})|X = x_1, \bar{W} = \bar{w}, \tilde{Z} = z] = 0$$

Then by iterated expectations and Assumption 2.ii:

$$\begin{aligned} & E[\delta(\tilde{V})|X = x_1, \bar{W} = \bar{w}, \tilde{Z} = z] \\ &= E[E[\delta(\tilde{V})|X, W^*]|X = x_1, \bar{W} = \bar{w}, \tilde{Z} = z] \\ &= 0 \end{aligned}$$

And so by Assumption 3.i $F_{W^*|X=x_1, \bar{W}=\bar{w}}$ -almost surely:

$$E[\delta(\tilde{V})|X = x_1, W^* = w^*] = 0$$

But then we see that the $L_2(F_{\tilde{V}|X=x_1, \bar{W}=\bar{w}})$ -inner product of δ and $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w}))$ is zero:

$$\begin{aligned} & E\left[\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V)\delta(\tilde{V})\middle|X = x_1, \bar{W} = \bar{w}\right] \\ &= E\left[E\left[\frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}(W^*)\middle|X, V\right]\delta(\tilde{V})\middle|X = x_1, \bar{W} = \bar{w}\right] \\ &= E\left[\frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}(W^*)E[\delta(\tilde{V})|X = x_1, W^*]\middle|X = x_1, \bar{W} = \bar{w}\right] \\ &= 0 \end{aligned}$$

Where the first equality follows by substituting for $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}$, the second by iterated expectations and the third because $E[\delta(\tilde{V})|X = x_1, W^*]$ is $F_{W^*|X=x_1, \bar{W}=\bar{w}}$ -almost surely zero. Since the inner product is zero for any δ in the null-space of $A_{x_1, \bar{w}}$, by definition $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w}))$ is in the orthogonal complement of the null-space. Thus c. holds.

Next note that Assumption 2.i and iterated expectations implies:

$$E[E[Y|W^*, X]|X = x, Z = z] = E[Y|X = x, Z = z]$$

Let a function δ be in the null space of $A_{x, \bar{w}}$, that is, for $F_{\tilde{V}|X=x, \bar{W}=\bar{w}}$ -almost all \tilde{v} :

$$E[\delta(\tilde{Z})|X = x, \bar{W} = \bar{w}, \tilde{V} = \tilde{v}] = 0$$

Then by iterated expectations and Assumption 2.ii:

$$\begin{aligned} & E[\delta(\tilde{Z})|X = x, \bar{W} = \bar{w}, \tilde{V} = \tilde{v}] \\ &= E[E[\delta(\tilde{Z})|X, W^*]|X = x, \bar{W} = \bar{w}, \tilde{V} = \tilde{v}] \\ &= 0 \end{aligned}$$

And so by Assumption 3.ii $F_{W^*|X=x, \bar{W}=\bar{w}}$ -almost surely:

$$E[\delta(\tilde{V})|X = x, W^* = w^*] = 0$$

But then we see that the $L_2(F_{\tilde{Z}|X=x, \bar{W}=\bar{w}})$ -inner product of δ and $E[Y|X = x, Z = (\cdot, \bar{w})]$ is zero:

$$\begin{aligned} & E \left[E[Y|X, Z] \delta(\tilde{Z}) \middle| X = x, \bar{W} = \bar{w} \right] \\ &= E \left[E[E[Y|X, W^*]|X, Z] \delta(\tilde{Z}) \middle| X = x, \bar{W} = \bar{w} \right] \\ &= E \left[E[Y|X, W^*] E[\delta(\tilde{Z})|X = x, W^*] \middle| X = x, \bar{W} = \bar{w} \right] \\ &= 0 \end{aligned}$$

And so $E[Y|X = x, Z = (\cdot, \bar{w})]$ is in the null space of $A_{x, \bar{w}}^*$. Thus g. holds.

Finally, points d. and h. One can then see that for each given (x_1, \bar{w}) Assumption 4.i. is precisely the condition (.1) where $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}((\cdot, \bar{w}))$ is the function δ , the inner-product is that of the space $L_2(F_{\tilde{V}|X=x, \bar{W}=\bar{w}})$, and the singular values, $\mu_k^{(T)}$ and functions $v_k^{(T)}$ are those of $A_{x, \bar{w}}^*$. In particular, $\mu_k^{(T)}$ and $v_k^{(T)}$ are given by $\mu_k^{(x, \bar{w})} = \mu_k(x, \bar{w})$ and $u_k^{(x, \bar{w})} = u_k(x, (\cdot, \bar{w}))$. Moreover, c is replaced by $\tilde{C}(x_1, x_2, \bar{w})$. For each given (x, \bar{w}) Assumption 4.ii is precisely the condition (.1) with δ given by $E[Y|Z = (\cdot, \bar{w}), X = x]$, the inner-product that of $L_2(F_{\tilde{Z}|X=x, \bar{W}=\bar{w}})$, and the singular system that of $A_{x, \bar{w}}$ and c equal to $\tilde{D}(x, \bar{w})$. □

Appendix A.2: Additional Comparison with Miao et. al. and Rokkanen

As we discuss in the literature review, some of our identification results in Section 1 resemble those of [Miao et al. \(2018\)](#). In particular, we identify the CASF (equivalently, the conditional distribution of potential outcomes) under very similar assumptions to those under which they identify the unconditional average structural function (equivalently, the marginal/unconditional distribution of potential outcomes). We first became aware of their work in between our posting of the first and second versions of our paper on Arxiv. Apart from identifying different objects, our results also differ from those of [Miao et al. \(2018\)](#) in that they do not consider panel settings nor provide an estimator based on their results. They also do not provide an identification result of the form similar our Theorem 1.1.b nor do they provide conditions for well-posedness. Below we provide a more detailed comparison of our identifying assumptions with theirs.

Our Assumption 2 is equivalent to their restriction (f). Our Assumptions 3.i and 3.ii play a similar role to the completeness assumptions in Conditions 2 and 3 of [Miao *et al.* \(2018\)](#). If there are no shared components \bar{W} and either our Assumption 2 holds or their conditional independence restriction (f) holds, then their Conditions 2 and 3 are stronger than our Assumption 3. In particular, under Assumption 2, our Assumption 3.i is weaker than the assumption $E[\delta(V)|X = x, Z] = 0 \iff \delta(V) = 0$ which is analogous to the corresponding assumption in their paper. Our Assumptions A.1 and 4.ii are of a very similar form to Assumptions A1, A2 and A3 in [Miao *et al.* \(2018\)](#) and play a similar role in part a. of our Theorem 1.1 to the role of A1-A3 in their paper.

Our characterization of the CASF in Theorem 1.1.a somewhat resembles the characterization of $p(y|do(x))$ in [Miao *et al.* \(2018\)](#). Their analysis does not use the potential outcomes framework, however $p(y|do(x))$ roughly represents the marginal distribution of potential outcomes. To see why this correspondence is not exact, note that in the discrete outcome case they equate $p(y|do(x))$ with (in our notation) $E[E[1\{Y = y\}|X = x, W^*]]$. To ensure that this equals the marginal distribution of potential outcomes one requires Assumption 1.i in our paper hold. [Miao *et al.* \(2018\)](#) do not make an assumption analogous to our Assumption 1.i but our reading of their work is that such an assumption is implicit in their analysis (alternatively, one could understand their work to equate W^* and U).

[Rokkanen \(2015\)](#) gives conditions for identification in a regression discontinuity design. His work applies the measurement error results of [Hu & Schennach \(2008\)](#) to the proxy control setting. We focus on comparison with [Rokkanen \(2015\)](#) rather than directly with [Hu & Schennach \(2008\)](#) because [Rokkanen \(2015\)](#) is an application of their work to a setting that is relatively close to our own. Here we provide more detail on the differences and similarities between our assumptions and discuss the reason for those differences. While there are similarities between our assumptions, our characterization of the CASF does not resemble that of [Rokkanen \(2015\)](#).

Section 2.3 of [Rokkanen \(2015\)](#) refers to variables M_1 , M_2 and M_3 . To compare assumptions we take these to be V , Z and X respectively, however his analysis allows M_3 to be some additional set of proxy controls rather than the treatments.

Our Assumptions 1.i, 2.i, and 2.ii resemble but are weaker than the analogous Assumptions C.2 and D.1 in [Rokkanen \(2015\)](#). More precisely, we require that (X, Z) be independent of V conditional on W^* , whereas an application of C.2 requires that X , Z , and V be jointly independent of W^* . Further, our Assumptions 1.i and 2.i are equivalent to independence of U with (X, Z) conditional on W^* , whereas D.1 would require that U be independent of (X, Z, V) conditional on W^* . C.1 would require that (X, V, Z, W^*) have joint distribution bounded by the product of their marginals (it also requires that V , Z and W^* have probability densities on their supports), by contrast we only requires Assumption 1.ii which states that the joint distribution of (X, W^*) be dominated by the product of their marginals. We require L_2 -completeness in Assumption 3, [Rokkanen \(2015\)](#) requires bounded completeness (which is weaker) in As-

sumptions C.5 and D.2. Since we take M_3 to be X , D.2 is redundant. In C.5 the completeness does not involve conditioning on a fixed X but under his Assumption C.1 this is equivalent to completeness conditional on X . [Rokkanen \(2015\)](#) does not require a condition like Assumption 4 which we use to establish a simple well-posed characterization of the CASF.

Crucially, we do not require anything like Condition C.4 in [Rokkanen \(2015\)](#) which requires that the mis-measured perfect controls satisfy a normalization like mean- or median-unbiasedness. The reason that we do not require a normalization of this kind (which is taken from [Hu & Schennach \(2008\)](#)) is that the results in [Hu & Schennach \(2008\)](#) apply for measurement error in treatments rather than just controls. Suppose that treatment variables are mis-measured in the following way: rather than observe the true treatments we instead observe the treatments multiplied by an unknown constant. In this case there is little hope of recovering a treatment effect without some normalization. Now suppose that the treatments are measured correctly and instead the controls are multiplied by an unknown constant. If the original controls and treatments satisfy an unconfoundedness assumption then so do the mis-measured controls and treatments. Therefore the mis-measurement is of no consequence and there is no need for a normalization. Our results show that this reasoning applies for more general and stochastic forms of measurement error

Appendix A.3: Implementation Details

Recall the definition of the estimator from Section 3 in which ridge regression is used for all non-parametric regressions, a ridge penalty is used in the PSMD step, and ϕ_n is multiplicatively separable (so that $\phi_n(x, v) = \rho_n(v) \otimes \chi_n(x)$). In both applications we choose basis functions ρ_n , χ_n , and ψ_n as follows. We let ρ_n and χ_n consist of the first 5-order power series of the components of v and x respectively. So letting v_j denote the j^{th} components of v which is of length $\dim(V)$, $\rho_n(v)$ contains all distinct functions of the form $v_1^{a_1} \times v_2^{a_2} \times \dots \times v_{\dim(V)}^{a_{\dim(V)}}$ where a_1, \dots, a_L are positive integers with $\sum_j^{\dim(V)} a_j \leq 5$. We define $\chi_n(x)$ analogously for x , likewise we define $\psi_n(x, z)$ analogously with the vector v replaced by the vector (x, z) . When performing ridge regressions and in the PSMD step we first normalize the basis functions to have sample mean zero and standard deviations equal to unity and as such we do not penalize the intercept.

Appendix B: Proofs

Proofs for Section 1

Proposition B.1. *Under Assumption 1:*

$$\bar{y}(x_1|x_2) = E[E[Y|W^*, X = x_1]|X = x_2]$$

Proof. By iterated expectations:

$$\bar{y}(x_1|x_2) = E[E[y_0(x_2, U)|W^*, X = x_2]|X = x_2]$$

By Assumption 1.i and the definition of y_0 :

$$\begin{aligned} E[y_0(x_1, U)|W^* = w^*, X = x_2] &= E[y_0(x_1, U)|W^* = w^*, X = x_1] \\ &= E[Y|W^* = w^*, X = x_1] \end{aligned}$$

Under Assumption 1.ii $E[Y|W^* = w^*, X = x_1]$ is well-defined for $F_{W^*|X=x_2}$ -almost all w^* (rather than just $F_{W^*|X=x_1}$ -almost all w^*). So we can substitute to get:

$$\bar{y}(x_1|x_2) = E[E[Y|W^*, X = x_1]|X = x_2]$$

□

To understand the use of Assumption 1.ii in the proof above suppose that for each x in the support of X , W^* admits a conditional probability density function $f_{W^*|X}(\cdot|x)$, then we can write:

$$\bar{y}(x_1|x_2) = \int E[Y|W^* = w^*, X = x_1] f_{W^*|X}(w^*|x_2) dw^*$$

Thus the conditional expectation $E[Y|W^* = w^*, X = x_1]$ must be identified for almost all w^* for which $f_{W^*|X}(w^*|x_2) > 0$. But $E[Y|W^* = w^*, X = x_1]$ is only uniquely defined for almost all w^* for which $f_{W^*|X}(w^*|x_1) > 0$. Thus we need the two conditional pdfs to be non-zero on the same support. Assumption 1.ii strengthens this full-support requirement to allow for W^* without a conditional probability density function (say, because F_{W^*} has mass points).

Proposition B.2. *Under Assumption 1.ii and 2.ii the Radon-Nikodym derivative $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}$ exists.*

Proof. Let $F_{(X,V)}[\mathcal{A}]$ be the probability that (X, V) is in the set \mathcal{A} . Note that:

$$\begin{aligned} F_{(X,V)}[\mathcal{A}] &= E\left[P[(X, V) \in \mathcal{A}|X, W^*]\right] \\ &= \int P[(x, V) \in \mathcal{A}|W^* = w^*] F_{(X,W^*)}(d(x, w^*)) \end{aligned}$$

Where the first equality follows by iterated expectations and the second by Assumption 2.ii. Let $F_X \otimes F_V[\mathcal{A}]$ be the product measure of \mathcal{A} . We have:

$$\begin{aligned} F_X \otimes F_V[\mathcal{A}] &= \int P[(x, V) \in \mathcal{A}] F_X(dx) \\ &= \int \int P[(x, V) \in \mathcal{A}|W^* = w^*] F_{W^*}(dw^*) F_X(dx) \\ &= \int P[(x, V) \in \mathcal{A}|W^* = w^*] F_X \otimes F_{W^*}(d(x, w^*)) \end{aligned}$$

Where $F_X \otimes F_{W^*}$ is the product measure of F_X and F_{W^*} . The first equality and second above follow by iterated expectations. But by Assumption 1.ii, the measure $F_{(X, W^*)}$ is non-zero on precisely the sets for which $F_X \otimes F_{W^*}$ is non-zero. Since $P[(x, V) \in \mathcal{A} | W^* = w^*]$ is weakly positive it follows that $F_{(X, V)}[\mathcal{A}]$ is strictly positive if and only if $F_X \otimes F_V[\mathcal{A}]$ is strictly positive. Since this holds for any \mathcal{A} , $F_X \otimes F_V$ and $F_{(X, V)}$ are absolutely continuous. Existence of $\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}$ then follows by the Radon-Nikodym theorem. \square

Proof Theorem 1.1. Throughout the proof, statements involving x_1 and x_2 should be understood to hold for F_X -almost all x_1 and x_2 .

First Theorem 1.1.a. By Lemma A.1, under Assumption 4.ii there exists a function γ with $E[\gamma(X, V)^2 | X] < \infty$ so that:

$$E[Y - \gamma(X, V) | X, Z] = 0$$

Fix such a γ , by iterated expectations and Assumption 2.ii:

$$\begin{aligned} E[\gamma(X, V) | X, Z] &= E[E[\gamma(X, V) | X, W^*, Z] | X, Z] \\ &= E[E[\gamma(X, V) | X, W^*] | X, Z] \end{aligned}$$

And by iterated expectations and Assumption 2.i:

$$\begin{aligned} E[Y | X, Z] &= [y_0(X, U) | X, Z] \\ &= E[E[y_0(X, U) | X, W^*, Z] | X, Z] \\ &= E[E[y_0(X, U) | X, W^*] | X, Z] \end{aligned}$$

And so:

$$E\left[E[y_0(X, U) | X, W^*] - E[\gamma(X, V) | X, W^*] \middle| X, Z\right] = 0$$

But then by Assumption 3.i:

$$E[y_0(X, U) | X, W^*] = E[\gamma(X, V) | X, W^*]$$

By Assumption 1.i and 1.ii:

$$E[y_0(X, U) | X = x_1, W^*] = E[y_0(x_1, U) | X = x_2, W^*]$$

And by Assumption 2.ii:

$$E[\gamma(X, V) | X = x_1, W^*] = E[\gamma(x_1, V) | X = x_2, W^*]$$

And so:

$$E[y_0(x_1, U) | X = x_2, W^*] = E[\gamma(x_1, V) | X = x_2, W^*]$$

The LHS has finite expectation conditional on $X = x_2$ by Assumption 1.iii, and thus so does the RHS above. By iterated expectations:

$$E[y_0(x_1, U) | X = x_2] = E[\gamma(x_1, V) | X = x_2]$$

And by definition the LHS equals $\bar{y}(x_1|x_2)$. To see that the mean on the RHS above is well-defined note that:

$$\begin{aligned} & E[\gamma(x_1, V)|X = x_2] \\ &= E[\gamma(x_1, V) \frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V)|X = x_1] \\ &\leq E[\gamma(x_1, V)^2|X = x_1] E[\frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V)^2|X = x_1] \end{aligned}$$

Where the inequality follows by Cauchy-Schwartz. The RHS in the last line is finite by Assumption 4.ii.

Now for Theorem 1.1.b. By Lemma A.1, under Assumption 4.i there exists a function φ with $E[\varphi(X, x_2, Z)^2|X] < \infty$ so that:

$$E[\varphi(x_1, x_2, Z)|X = x_1, V] = \frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V)$$

Fix such a φ . The next step refers to $\frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}$, which is the Radon-Nikodym derivative of $F_{W^*|X=x_2}$ with respect to $F_{W^*|X=x_1}$. By the Radon-Nikodym theorem this exists if the two distributions are absolutely continuous which in turn is implied by Assumption 1.ii.

Note that under Assumption 2.ii:

$$E\left[\frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}(W^*) \middle| X = x_1, V\right] = \frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(V)$$

To see this note that:

$$\begin{aligned} & E\left[\frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}(W^*) \middle| X = x_1, V = v\right] \tag{.2} \\ &= E\left[\frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}(W^*) \frac{dF_{W^*|X=x_1, V=v}}{dF_{W^*|X=x_2}}(W^*) \middle| X = x_2\right] \\ &= E\left[\frac{dF_{W^*|X=x_1, V=v}}{dF_{W^*|X=x_1}}(W^*) \middle| X = x_2\right] \\ &= E\left[\frac{dF_{V|W^*, X=x_1}}{dF_{V|X=x_1}}(v, W^*) \middle| X = x_2\right] \end{aligned}$$

Where we let $\frac{dF_{V|W^*, X=x_1}}{dF_{V|X=x_1}}(v, w^*)$ denote the Radon-Nikodym derivative of $F_{V|W^*=w^*, X=x_2}$ with respect to $F_{V|X=x_1}$. The first equality follows by the Radon-Nikodym theorem, the second by the chain rule, the third by properties of Radon-Nikodym derivatives. Now note that by Assumption 2.ii:

$$\frac{dF_{V|W^*, X=x_1}}{dF_{V|X=x_1}}(v, W^*) = \frac{dF_{V|W^*, X=x_2}}{dF_{V|X=x_1}}(v, W^*)$$

and so:

$$\begin{aligned}
& E \left[\frac{dF_{V|W^*, X=x_1}}{dF_{V|X=x_1}}(v, W^*) \middle| X = x_2 \right] \\
&= E \left[\frac{dF_{V|W^*, X=x_2}}{dF_{V|X=x_1}}(v, W^*) \middle| X = x_2 \right] \\
&= \frac{dF_{V|X=x_2}}{dF_{V|X=x_1}}(v)
\end{aligned}$$

Where the final equality follows by the properties of the Radon-Nikodym derivative. Further, by iterated expectations and Assumption 2.ii:

$$\begin{aligned}
E[\varphi(x_1, x_2, Z)|X = x_1, V] &= E[E[\varphi(x_1, x_2, Z)|W^*, X, V]|X = x_1, V] \\
&= E[E[\varphi(x_1, x_2, Z)|W^*, X]|X = x_1, V]
\end{aligned}$$

So we have:

$$E \left[\frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}(W^*) - E[\varphi(x_1, x_2, Z)|W^*, X] \middle| X = x_1, V = v \right] = 0$$

By Assumption 3.ii this implies:

$$E[\varphi(x_1, x_2, Z)|W^*, X = x_1] = \frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}(W^*) \quad (.3)$$

Next note that by iterated expectations and Assumptions 1.i and 1.ii:

$$\begin{aligned}
E[y_0(x_1, U)|X = x_2] &= E[E[y_0(x_1, U)|X = x_2, W^*]|X = x_2] \\
&= E[E[y_0(X, U)|X = x_1, W^*]|X = x_2] \quad (.4)
\end{aligned}$$

And by the Radon-Nikodym theorem:

$$\begin{aligned}
& E[E[Y|X = x_1, W^*]|X = x_2] \\
&= E \left[\frac{dF_{W^*|X=x_2}}{dF_{W^*|X=x_1}}(W^*) E[y_0(X, U)|X, W^*] \middle| X = x_1 \right]
\end{aligned}$$

Substituting for the LHS by (.4) and for the Radon-Nikodym derivative on the RHS by (.3) we get:

$$\begin{aligned}
& E[y_0(x_1, U)|X = x_2] \\
&= E \left[E[\varphi(x_1, x_2, Z)|W^*, X] E[y_0(X, U)|X, W^*] \middle| X = x_1 \right]
\end{aligned}$$

Next note that:

$$\begin{aligned}
& E \left[E[\varphi(x_1, x_2, Z)|W^*, X] E[y_0(X, U)|X, W^*] \Big| X = x_1 \right] \\
&= E \left[\varphi(x_1, x_2, Z) E[y_0(X, U)|X, W^*] \Big| X = x_1 \right] \\
&= E \left[\varphi(x_1, x_2, Z) E[y_0(X, U)|X, W^*, Z] \Big| X = x_1 \right] \\
&= E \left[\varphi(x_1, x_2, Z) y_0(X, U) \Big| X = x_1 \right] \\
&= E \left[\varphi(x_1, x_2, Z) Y \Big| X = x_1 \right]
\end{aligned}$$

Where the first equality follows by iterated expectations, the second by Assumption 2.i, the third by iterated expectations and the final by the definition of Y . Combining we get:

$$E[y_0(x_1, U)|X = x_2] = E \left[\varphi(x_1, x_2, Z) Y \Big| X = x_1 \right]$$

□

Proof Theorem 1.2. First we prove 1.2.a. Note that by The Radon-Nikodym theorem, for any $\tilde{\gamma}$ with $E[\tilde{\gamma}(X, V)^2|X = x]$ finite for F_X -almost all x , we have:

$$\begin{aligned}
& E[\tilde{\gamma}(x_1, V)|X = x_2] \\
&= E \left[\tilde{\gamma}(X, V) \frac{dF_V|X=x_2}{dF_V|X=x_1}(V) \Big| X = x_1 \right] \\
&= E[\tilde{\gamma}(X, V) E[\varphi(x_1, x_2, Z)|X, V] \Big| X = x_1]
\end{aligned}$$

Where φ satisfies the equations for φ in conclusion 1. of Lemma A.1. By iterated expectations:

$$\begin{aligned}
& E[\tilde{\gamma}(X, V) E[\varphi(x_1, x_2, Z)|X, V] \Big| X = x_1] \\
&= E[\tilde{\gamma}(X, V) \varphi(x_1, x_2, Z) \Big| X = x_1] \\
&= E[E[\tilde{\gamma}(X, V)|X, Z] \varphi(x_1, x_2, Z) \Big| X = x_1]
\end{aligned}$$

Under the conditions of 1.2.a, Theorem 1.1.b holds, and so combining the characterization of $\bar{y}(x_1|x_2)$ in Theorem 1.1.b and the equation above we get:

$$\begin{aligned}
& \bar{y}(x_1|x_2) - E[\tilde{\gamma}(x_1, V)|X = x_2] \\
&= E \left[(Y - E[\tilde{\gamma}(X, V)|X, Z]) \varphi(x_1, x_2, Z) \Big| X = x_1 \right]
\end{aligned}$$

And so applying Cauchy-Schwartz:

$$\begin{aligned} & (\bar{y}(x_1|x_2) - E[\tilde{\gamma}(x_1, V)|X = x_2])^2 \\ & \leq E[\varphi(x_1, x_2, Z)^2|X = x_1] \\ & \quad \times E\left[\left(Y - E[\tilde{\gamma}(X, V)|X, Z]\right)^2\middle|X = x_1\right] \end{aligned}$$

But Lemma A.1 states that we can find a solution φ for which $E[\varphi(x_1, x_2, Z)^2|X = x_1]$ is less than $C(x_1, x_2)$. Thus from the above we get:

$$\begin{aligned} & (\bar{y}(x_1|x_2) - E[\tilde{\gamma}(x_1, V)|X = x_2])^2 \\ & \leq C(x_1, x_2)E\left[\left(Y - E[\tilde{\gamma}(X, V)|X, Z]\right)^2\middle|X = x_1\right] \end{aligned}$$

Now part b. By iterated expectations for any $\tilde{\varphi}$ with $E[\tilde{\varphi}(x_1, x_2, Z)^2|X = x_1] \leq \infty$ for F_X -almost all x_1 and x_2 , we have:

$$E[\tilde{\varphi}(x_1, x_2, Z)Y|X = x_1] = E[\tilde{\varphi}(x_1, x_2, Z)E[Y|X, Z]|X = x_1]$$

Letting γ satisfy the equations in conclusion 2. of Lemma A.1 we can substitute out $E[Y|X, Z]$ in the above to get:

$$\begin{aligned} & E[\tilde{\varphi}(x_1, x_2, Z)Y|X = x_1] \\ & = E[\tilde{\varphi}(x_1, x_2, Z)E[\gamma(X, V)|X, Z]|X = x_1] \\ & = E[E[\tilde{\varphi}(x_1, x_2, Z)|X, V]\gamma(X, V)|X = x_1] \end{aligned} \tag{.5}$$

Where the second equality follows by iterated expectations. Recall that under the conditions of part a. of the Theorem 1.1:

$$\bar{y}(x_1|x_2) = E[\gamma(x_1, V)|X = x_2] \tag{.6}$$

By the Radon-Nikodym theorem:

$$E[\gamma(x_1, V)|X = x_2] = E\left[\gamma(X, V)\frac{dF_{V|X=x_2}(V)}{dF_{V|X=x_1}(V)}\middle|X = x_1\right]$$

Combining with (.5) and (.6) we get:

$$\begin{aligned} & \bar{y}(x_1|x_2) - E[\tilde{\varphi}(x_1, x_2, Z)Y|X = x_1] \\ & = E\left[\gamma(X, V)\left(\frac{dF_{V|X=x_2}(V)}{dF_{V|X=x_1}(V)} - E[\tilde{\varphi}(x_1, x_2, Z)|X, V]\right)\middle|X = x_1\right] \end{aligned}$$

By Cauchy-Schwartz:

$$\begin{aligned} & (\bar{y}(x_1|x_2) - E[\tilde{\varphi}(x_1, x_2, Z)Y|X = x_1])^2 \\ & = E[\gamma(X, V)^2|X = x_1] \\ & \quad \times E\left[\left(\frac{dF_{V|X=x_2}(v)}{dF_{V|X=x_1}(v)} - E[\tilde{\varphi}(x_1, x_2, Z)|X, V]\right)^2\middle|X = x_1\right] \end{aligned}$$

Recall that Lemma A.1 states that we can find such a γ with $E[\gamma(x_1, V)^2 | X = x_1] \leq D(x_1)$. So from the above we have:

$$\begin{aligned} & (\bar{y}(x_1|x_2) - E[\tilde{\varphi}(x_1, x_2, Z)Y | X = x_1])^2 \\ &= D(x_1) E \left[\left(\frac{dF_V | X=x_2}{dF_V | X=x_1}(v) - E[\tilde{\varphi}(x_1, x_2, Z) | X, V] \right)^2 \middle| X = x_1 \right] \end{aligned}$$

□

Proofs for Section 2

The proofs of Propositions 2.1 and 2.2 below use the following three facts about conditional independence. Let W_1, W_2, W_3 , and W_4 be random variables. We have:

$$W_1 \perp\!\!\!\perp W_2 | (W_3, W_4) \implies W_1 \perp\!\!\!\perp (W_2, W_3) | (W_3, W_4) \quad (.7)$$

$$W_1 \perp\!\!\!\perp (W_2, W_3) | W_4 \implies W_1 \perp\!\!\!\perp (W_2, W_3) | (W_3, W_4) \quad (.8)$$

$$W_1 \perp\!\!\!\perp (W_2, W_3) | W_4 \implies W_1 \perp\!\!\!\perp W_2 | W_4 \quad (.9)$$

Proof of Proposition 2.1. Let us restate the conditions of the proposition:

$$(X_1, \dots, X_{\lfloor t/2 \rfloor - 1}) \perp\!\!\!\perp (X_{\lfloor t/2 \rfloor + 1}, \dots, X_T) | (\tilde{W}^*, X_{\lfloor t/2 \rfloor}) \quad (.10)$$

$$U_t \perp\!\!\!\perp (X_1, \dots, X_t) | \tilde{W}^* \quad (.11)$$

Using (.8) we see (.11) implies:

$$U_t \perp\!\!\!\perp (X_1, \dots, X_t) | (\tilde{W}^*, X_{\lfloor t/2 \rfloor})$$

And applying (.9):

$$U_t \perp\!\!\!\perp (X_{\lfloor t/2 \rfloor}, \dots, X_t) | (\tilde{W}^*, X_{\lfloor t/2 \rfloor})$$

Substituting the definitions of Z and W^* gives $U_t \perp\!\!\!\perp (X_t, Z) | W^*$.

Twice applying (.7), (.10) implies that:

$$(X_1, \dots, X_{\lfloor t/2 \rfloor}) \perp\!\!\!\perp (X_{\lfloor t/2 \rfloor}, \dots, X_t) | (\tilde{W}^*, X_{\lfloor t/2 \rfloor})$$

Substituting the definitions of V, Z and W^* the above is equivalent to $V \perp\!\!\!\perp (X_t, Z) | W^*$. □

Proof of Proposition 2.2. Let us restate the conditions of the proposition. Firstly,

$$U_t \perp\!\!\!\perp (X_1, \dots, X_t, U_1, \dots, U_{t-1}) | \tilde{W}^* \quad (.12)$$

and conditional on $(\tilde{W}^*, X_{\lfloor t/2 \rfloor}, U_{\lfloor t/2 \rfloor})$:

$$(X_1, \dots, X_{\lfloor t/2 \rfloor - 1}, U_1, \dots, U_{\lfloor t/2 \rfloor - 1}) \perp\!\!\!\perp (X_{\lfloor t/2 \rfloor + 1}, \dots, X_t, U_{\lfloor t/2 \rfloor + 1}, \dots, U_t) \quad (.13)$$

Recall from Section 1 that for any x and any $u_1 \neq u_2$ we have $y_{0,t}(x, u_1) \neq y_{0,t}(x, u_2)$. Therefore (.12) implies:

$$U_t \perp\!\!\!\perp (X_1, \dots, X_t, Y_1, \dots, Y_{t-1}) | \tilde{W}^*$$

Applying (.8) we get:

$$U_t \perp\!\!\!\perp (X_1, \dots, X_t, Y_1, \dots, Y_{t-1}) | (\tilde{W}^*, X_{\lfloor t/2 \rfloor}, Y_{\lfloor t/2 \rfloor})$$

Applying (.9):

$$U_t \perp\!\!\!\perp (X_{\lfloor t/2 \rfloor}, \dots, X_t, Y_{\lfloor t/2 \rfloor}, \dots, Y_{t-1}) | (\tilde{W}^*, X_{\lfloor t/2 \rfloor}, Y_{\lfloor t/2 \rfloor})$$

Substituting the definitions of Z and W^* gives $U_t \perp\!\!\!\perp (X_t, Z) | W^*$.

Again, using that for any x and any $u_1 \neq u_2$ we have $y_{0,t}(x, u_1) \neq y_{0,t}(x, u_2)$, (.13) implies that conditional on $(\tilde{W}^*, X_{\lfloor t/2 \rfloor}, Y_{\lfloor t/2 \rfloor})$:

$$(X_1, \dots, X_{\lfloor t/2 \rfloor - 1}, Y_1, \dots, Y_{\lfloor t/2 \rfloor - 1}) \perp\!\!\!\perp (X_{\lfloor t/2 \rfloor + 1}, \dots, X_t, Y_{\lfloor t/2 \rfloor + 1}, \dots, Y_{t-1})$$

Applying (.7) we get that conditional on $(\tilde{W}^*, X_{\lfloor t/2 \rfloor}, Y_{\lfloor t/2 \rfloor})$:

$$(X_1, \dots, X_{\lfloor t/2 \rfloor}, Y_1, \dots, Y_{\lfloor t/2 \rfloor}) \perp\!\!\!\perp (X_{\lfloor t/2 \rfloor}, \dots, X_t, Y_{\lfloor t/2 \rfloor}, \dots, Y_{t-1})$$

Substituting the definitions of V , Z and W^* the above is equivalent to $V \perp\!\!\!\perp (X_t, Z) | W^*$. \square

Proofs for Section 3

We now give proofs for results in Section 3. Throughout this section expectations with a subscript only integrate over the random variables in that subscript, in other words the expectation treats all other random variables as fixed. For example, $E_Z[Z|X] = \int z X F_Z(dz) = E[Z]X$ and $E_V[V|Z|X = x] = \int v Z F_{V|X=x}(dv) = E[V|X = x]Z$, and $E_Z[(\pi(x, Z)' \hat{\theta})^2 | X = x] = \int (\pi(x, z)' \hat{\theta})^2 F_{Z|X=x}(dz)$.

The following proposition gives rates at which the reduced-form function $(x, z) \mapsto g(x, z)$ can be approximated uniformly by linear combinations of the components of $(x, z) \mapsto \pi_n(x, z)$, which are conditional expectations of the basis functions $\phi_n(X, V)$

Proposition 3.1. *Suppose Assumptions 1, 2, and 3 hold, Assumption 4.ii holds with $D(X)$ bounded above by a constant with probability 1, and Assumptions 5.1.i, 5.1.ii, 5.2, 5.3.i, and 5.3.ii hold. Then there is a sequence $\{\theta_n\}_{n=1}^{\infty}$ with $E[|\phi_n(X, V)' \theta_n|^2]$ bounded above uniformly over n so that, uniformly over $F_{(X,Z)}$ -almost all (x, z) :*

$$|g(x, z) - \pi_n(x, z)' \theta_n| \lesssim \ell_{\rho, n}(s_1)$$

if X has finite discrete support, and otherwise:

$$|g(x, z) - \pi_n(x, z)' \theta_n| \lesssim \ell_{\rho, n}(s_1) + (\xi_{\rho, n} \ell_{\chi, n}(1))^{\bar{s}}$$

Where $\tilde{s} = \frac{\min\{s_2, s_3, 1\}}{\min\{s_2, s_3, 1\} + 1}$ and the above is uniform over $F_{(X,Z)}$ -almost all (x, z) .

Proof. First note that for any length- $k(n)l(n)$ column vector θ_n , there is a $l(n)$ -by- $k(n)$ matrix $\tilde{\theta}_n$ so that $(\rho_n(V) \otimes \chi_n(X))' \theta_n = \chi_n(X)' \tilde{\theta}_n \rho_n(V)$ and vice-versa. In particular, we can let the entry of $\tilde{\theta}_n$ in the j^{th} row and k^{th} column be the $(j-1)l(n) + k$ -th entry of θ_n . For convenience we will find a $l(n)$ -by- $k(n)$ matrix rather than a length- $k(n)l(n)$ vector.

We will now show that for each n , there exists a vector-valued function β_n so that $E[(\beta_n(X)' \rho_n(V))^2]^{1/2} \lesssim 1$ and:

$$\text{ess sup } |g(X, Z) - \beta_n(X)' E[\rho_n(V) | X, Z]| \lesssim \ell_{\rho, n}(s_1)$$

Combining Assumptions 5.1.i and 5.3.i, uniformly over $F_{(X,Z)}$ -almost all (x, z) :

$$\inf_B E \left[\left(\frac{dF_{(X,Z,V)}}{d(F_{(X,Z)} \otimes F_V)}(x, z, V) - \rho_n(V)' B \right)^2 \right]^{1/2} \lesssim \ell_{\rho, n}(s_1)$$

This implies that there exist sequences of functions B_n and r_n so that for $F_{(X,Z,V)}$ -almost all (x, z, v) :

$$\frac{dF_{(X,Z,V)}}{d(F_{(X,Z)} \otimes F_V)}(x, z, v) = \rho_n(v)' B_n(x, z) + r_n(x, z, v) \quad (.14)$$

Where:

$$E[r_n(x, z, V)^2]^{1/2} \lesssim \ell_{\rho, n}(s_1)$$

By Assumptions 1, 2, 3 and 4.ii there is a γ with $E[\gamma(x, V)^2 | X = x]^{1/2} \leq D(x)$ so that $g(x, z) = E[\gamma(x, V) | X = x, Z = z]$ (see Lemma A.1). Thus, for any vector-valued function of appropriate dimension $\beta_n(x)$ we get for $F_{(X,Z)}$ -almost all (x, z) :

$$\begin{aligned} & g(x, z) - \beta_n(x)' E[\rho_n(V) | X = x, Z = z] \\ &= E[\gamma(x, V) - \beta_n(x)' \rho_n(V) | X = x, Z = z] \\ &= E \left[(\gamma(x, V) - \beta_n(x)' \rho_n(V)) \frac{dF_{(X,Z,V)}}{d(F_{(X,Z)} \otimes F_V)}(x, z, V) \right] \end{aligned}$$

Where the second equality follows by the definition of the Radon-Nikodym derivative. Substituting (.14), the RHS above becomes:

$$\begin{aligned} & \left(E[\gamma(x, V) \rho_n(V)'] - \beta_n(x)' E[\rho_n(V) \rho_n(V)'] \right) B_n(x, z) \quad (.15) \\ & + E \left[(\gamma(x, V) - \beta_n(x)' \rho_n(V)) r_n(x, z, V) \right] \end{aligned}$$

Assuming $E[\rho_n(V)\rho_n(V)']$ is non-singular, we can set $\beta_n(x)$ so that:

$$\beta_n(x) = E[\rho_n(V)\rho_n(V)']^{-1}E[\rho_n(V)\gamma(x, V)]$$

Substituting into (.15) the first term disappears and we get:

$$\begin{aligned} & g(x, z) - \beta_n(x)'E[\rho_n(V)|X = x, Z = z] \\ &= E\left[(\gamma(x, V) - \beta_n(x)'\rho_n(V))r_n(x, z, V)\right] \end{aligned}$$

By Cauchy-Schwartz:

$$\begin{aligned} & |E[(\gamma(x, V) - \beta_n(x)'\rho_n(V))r_n(x, z, V)]| \\ & \leq E\left[(\gamma(x, V) - \beta_n(x)'\rho_n(V))^2\right]^{1/2} E[r_n(x, z, V)^2]^{1/2} \\ & \lesssim \ell_{\rho, n}(s_1)E\left[(\gamma(x, V) - \beta_n(x)'\rho_n(V))^2\right]^{1/2} \end{aligned}$$

Note that $\beta_n(x)'\rho_n(v)$ is a least $L_2(F_V)$ -norm projection of $\gamma(x, V)$ onto $\rho_n(V)$ and so:

$$\begin{aligned} E\left[(\gamma(x, V) - \beta_n(x)'\rho_n(V))^2\right] & \leq E[\gamma(x, V)^2] \\ & = E\left[\gamma(x, V)^2 \frac{dF_X \otimes F_V}{dF_{(X, V)}}(x, V) | X = x\right] \\ & \leq \frac{\bar{D}^2}{\underline{c}} \end{aligned}$$

Where $\bar{D} < \infty$ is an almost-sure upper bound on $D(X)$ which exists by supposition, and $\underline{c} > 0$ is a lower bound on $\frac{dF_{(X, V)}}{dF_X \otimes F_V}$ which exists by Assumption 5.2.ii. Note that by properties of the least-squares projection we also have:

$$E[(\beta_n(x)'\rho_n(V))^2]^{1/2} \leq \bar{D} \quad (.16)$$

This in turn implies:

$$E[(\beta_n(x)'\rho_n(V))^2 | X = x] \leq \frac{\bar{D}^2}{\underline{c}}$$

And so:

$$E[(\beta_n(X)'\rho_n(V))^2]^{1/2} \leq \frac{\bar{D}}{\sqrt{\underline{c}}} \lesssim 1 \quad (.17)$$

In all, we get that uniformly over $F_{(X, Z)}$ -almost all (x, z) :

$$|g(x, z) - \beta_n(x)'E[\rho_n(V)|X = x, Z = z]| \lesssim \ell_{\rho, n}(s_1) \quad (.18)$$

Now consider the case of X with discrete finite support. Both $\beta_n(X)$ and $\chi_n(X)$ then have discrete finite support and by Assumption 5.1.ii for n sufficiently large any function defined on \mathcal{X} is a linear transformation of χ_n . Therefore, there exists a matrix θ_n so that for F_X -almost all x :

$$\chi_n(x)' \theta_n = \beta_n(x)$$

And so from (.18) we immediately get:

$$E \left[\left(g(X, Z) - \chi_n(x)' \theta_n E[\rho_n(V) | X, Z] \right)^2 \middle| X = x \right]^{1/2} \lesssim \ell_{\rho, n}(s_1)$$

Moreover, from (.17) we get $E[(\chi_n(X)' \theta_n \rho_n(V))^2]^{1/2} \lesssim 1$, and we are done.

The case of continuously distributed X requires more work. The function β_n defined above may not be smooth, to address this we first show that we can smooth-out β_n without incurring too much additional approximation error and then show we can approximate the smoothed out function by a linear combination of the basis functions that compose χ_n . Let $\{b_n\}_{n=1}^\infty$ be a sequence of strictly positive scalars with $b_n \rightarrow 0$ and for each n , define a linear operator M_n by:

$$M_n[\delta](x) = \frac{\int_{\mathcal{X} \cap B_{x, b_n}} \delta(x') dx'}{\int_{\mathcal{X} \cap B_{x, b_n}} dx'}$$

Where B_{x, b_n} denotes the Euclidean ball in $\mathbb{R}^{\dim(X)}$ of radius b_n centered at x . Note that under the Assumption 5.2.ii, for sufficiently large n there exists $\underline{r} > 0$ so that for all $x \in \mathcal{X}$:

$$1 \geq \frac{\int_{\mathcal{X} \cap B_{x, b_n}} dx'}{\int_{B_{0, b_n}} dx'} \geq \underline{r}$$

We will use M_n to smooth out β , in particular let $\tilde{\beta}_n$ be the smoothed analogue of β given by:

$$\tilde{\beta}_n(x) = M_n[\beta(X)](x)$$

It is not difficult to see that:

$$\sup_{x \in \mathcal{X}} \|E[\rho_n(V) \rho_n(V)']^{1/2} \tilde{\beta}_n(x)\|_2 \leq \sup_{x \in \mathcal{X}} \|E[\rho_n(V) \rho_n(V)']^{1/2} \beta_n(x)\|_2^2$$

We will show $E[|\tilde{\beta}_n(X)' \rho_n(V)|^2]$ is bounded uniformly over n . Note that:

$$\begin{aligned}
\int_{\mathcal{X}} |\tilde{\beta}_n(x)' \rho_n(v)|^2 dx &= \int_{\mathcal{X}} |M_n[\beta_n(X)' \rho_n(v)](x)|^2 dx \\
&\leq \int_{\mathcal{X}} \int_{\mathcal{X}} \frac{1\{|x' - x|_2 \leq b_n\}}{\int_{\mathcal{X} \cap B_{x, b_n}} dx'} |\beta_n(x')' \rho_n(v)|^2 dx' dx \\
&\leq \frac{1}{\underline{r}} \int_{\mathcal{X}} \int_{\mathcal{X}} \frac{1\{|x' - x|_2 \leq b_n\}}{\int_{B_{0, b_n}} dx'} dx |\beta_n(x')' \rho_n(v)|^2 dx' \\
&\leq \frac{1}{\underline{r}} \int_{\mathcal{X}} |\beta_n(x)' \rho_n(v)|^2 dx
\end{aligned}$$

Where the first equality uses the definition of $\tilde{\beta}_n$, the subsequent inequality follows by Jensen's inequality, and the next inequality by swapping the order of integration (valid by Tonelli's theorem as the integrand is positive) and using $\frac{\int_{\mathcal{X} \cap B_{x, b_n}} dx'}{\int_{B_{0, b_n}} dx'} \geq \underline{r}$. The final inequality follows from $\int_{\mathcal{X}} \frac{1\{|x' - x|_2 \leq b_n\}}{\int_{B_{0, b_n}} dx'} dx \leq 1$. Now applying the upper and lower bounds $\bar{f} < \infty$ and $\underline{f} > 0$ on the density of X , which exist by Assumption 5.2.iii, the above implies:

$$E[|\tilde{\beta}_n(X)' \rho_n(v)|^2] \leq \frac{\bar{f}}{\underline{r}\underline{f}} E[|\beta_n(X)' \rho_n(v)|^2]$$

Finally, integrating both sides above over v against the measure F_V and applying the upper and lower bounds $\bar{c} < \infty$ and $\underline{c} > 0$ on $\frac{dF_{(X, V)}}{dF_X \otimes F_V}$, which exist by Assumption 5.2.ii, we get:

$$\begin{aligned}
E[|\tilde{\beta}_n(X)' \rho_n(V)|^2] &\leq \frac{\bar{c}\bar{f}}{\underline{c}\underline{r}\underline{f}} E[|\beta_n(X)' \rho_n(V)|^2] \\
&\leq \frac{\bar{c}\bar{f}}{\underline{c}^2 \underline{r}\underline{f}} \bar{D}^2
\end{aligned} \tag{.19}$$

Where the final inequality follows by (.17). Now we will show that the function $\tilde{\beta}_n(\cdot)' E[\rho_n(V)|X = x, Z = z]$ is Lipschitz continuous (i.e., an element of $\Lambda_1^{\dim(X)}(c)$ for some $c > 0$). As an intermediate step we establish that for any function δ with $|\delta(x)| \leq c$ for all $x \in \mathcal{X}$:

$$|M_n[\delta](x_1) - M_n[\delta](x_2)| \leq \frac{2}{\underline{r}} c \frac{\dim(X)}{b_n} \|x_1 - x_2\|_2 \tag{.20}$$

To see this first note that:

$$\begin{aligned}
& M_n[\delta](x_1) - M_n[\delta](x_2) \\
&= \frac{\int_{\mathcal{X} \cap B_{x_1, b_n}} \delta(x) dx}{\int_{\mathcal{X} \cap B_{x_1, b_n}} dx} - \frac{\int_{\mathcal{X} \cap B_{x_2, b_n}} \delta(x) dx}{\int_{\mathcal{X} \cap B_{x_2, b_n}} dx} \\
&= \frac{\int_{\mathcal{X} \cap B_{x_1, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} \delta(x) dx - \int_{\mathcal{X} \cap B_{x_2, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} \delta(x) dx}{\int_{\mathcal{X} \cap B_{x_2, b_n}} dx} \\
&+ \left(\frac{1}{\int_{\mathcal{X} \cap B_{x_1, b_n}} dx} - \frac{1}{\int_{\mathcal{X} \cap B_{x_2, b_n}} dx} \right) \int_{\mathcal{X} \cap B_{x_1, b_n}} \delta(x) dx \\
&= \frac{\int_{\mathcal{X} \cap B_{x_1, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} \delta(x) dx - \int_{\mathcal{X} \cap B_{x_2, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} \delta(x) dx}{\int_{\mathcal{X} \cap B_{x_2, b_n}} dx} \\
&+ \frac{\int_{\mathcal{X} \cap B_{x_2, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} dx - \int_{\mathcal{X} \cap B_{x_1, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} dx}{\int_{\mathcal{X} \cap B_{x_2, b_n}} dx} \frac{\int_{\mathcal{X} \cap B_{x_1, b_n}} \delta(x) dx}{\int_{\mathcal{X} \cap B_{x_1, b_n}} dx}
\end{aligned}$$

Using that $|\delta(x)| \leq c$ for all $x \in \mathcal{X}$ and applying the triangle inequality we get:

$$\begin{aligned}
& |M_n[\delta](x_1) - M_n[\delta](x_2)| \\
&\leq 2c \frac{\int_{\mathcal{X} \cap B_{x_1, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} dx + \int_{\mathcal{X} \cap B_{x_2, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} dx}{\int_{\mathcal{X} \cap B_{x_2, b_n}} dx} \\
&\leq \frac{\int_{B_{0, b_n}} dx}{\int_{\mathcal{X} \cap B_{x_2, b_n}} dx} 2c \frac{\int_{B_{x_1, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} dx + \int_{B_{x_2, b_n} \setminus [B_{x_1, b_n} \cap B_{x_2, b_n}]} dx}{\int_{B_{0, b_n}} dx} \\
&\leq \frac{2}{\underline{r}} c \frac{2 \int_{B_{0, b_n}} dx - 2 \int_{B_{x_1, b_n} \cap B_{x_2, b_n}} dx}{\int_{B_{0, b_n}} dx}
\end{aligned}$$

Where the final inequality uses $\frac{\int_{\mathcal{X} \cap B_{x, b_n}} dx'}{\int_{B_{0, b_n}} dx'} \geq \underline{r}$. Suppose $\|x_1 - x_2\|_2 \leq 2b_n$, then we have:

$$B_{\frac{x_1 + x_2}{2}, b_n - \frac{1}{2}\|x_2 - x_1\|_2} \subseteq (B_{x_1, b_n} \cap B_{x_2, b_n})$$

Therefore:

$$\int_{B_{\frac{x_1 + x_2}{2}, b_n - \frac{1}{2}\|x_2 - x_1\|_2}} dx \leq \int_{B_{x_1, b_n} \cap B_{x_2, b_n}} dx$$

And so by the volumes of Euclidean balls we get:

$$\frac{2 \int_{B_{0, b_n}} dx - 2 \int_{B_{x_1, b_n} \cap B_{x_2, b_n}} dx}{\int_{B_{0, b_n}} dx} \leq 2 - 2 \left(1 - \frac{\|x_2 - x_1\|_2}{2b_n}\right)^{\dim(X)}$$

Thus, (still assuming $\|x_1 - x_2\|_2 \leq 2b_n$) we get:

$$\begin{aligned} |M_n[\delta](x_1) - M_n[\delta](x_2)| &\leq \frac{2}{r} c \left[2 - 2\left(1 - \frac{\|x_1 - x_2\|_2}{2b_n}\right)^{\dim(X)} \right] \\ &\leq \frac{2}{r} c \frac{\dim(X)}{b_n} \|x_1 - x_2\|_2 \end{aligned}$$

For the second inequality we have used that for any $0 \leq y \leq 2b_n$ we have $2 - 2\left(1 - \frac{y}{2b_n}\right)^{\dim(X)} \leq \frac{\dim(X)}{b_n} y$ (they are equal at $y = 0$ and the latter has larger derivative in y over $[0, 2b_n]$). Note that if $\|x_1 - x_2\|_2 \geq 2b_n$ then the expression above is greater than $2c$ which clearly bounds $|M_n[\delta](x_1) - M_n[\delta](x_2)|$ and so we have that for any $\|x_1 - x_2\|_2$ the inequality holds.

Now, we will upper bound the function $\beta_n(\cdot)' E[\rho_n(V) | X = x, Z = z]$. Note that (.16) is equivalent to:

$$\|E[\rho_n(V)\rho_n(V)]^{1/2} \beta_n(x)\|_2 \leq \bar{D}$$

Using the above and the definition of $\xi_{\rho,n}$ we get:

$$\begin{aligned} &|\beta_n(x_1)' E[\rho_n(V) | X = x_2, Z = z]| \\ &\leq \|E[\rho_n(V)\rho_n(V)]^{1/2} \beta_n(x_1)\|_2 \|E[\rho_n(V)\rho_n(V)]^{-1/2} E[\rho_n(V) | X = x_2, Z = z]\|_2 \\ &\leq \bar{D} \text{ess sup} \|E[\rho_n(V)\rho_n(V)]^{-1/2} \rho_n(V)\|_2 \\ &= \bar{D} \xi_{\rho,n} \end{aligned}$$

Where ‘ess sup’ is the essential supremum over the distribution of V . Using the upper bound above, and applying (.20) we get:

$$\begin{aligned} &|\tilde{\beta}_n(x_1)' E[\rho_n(V) | X = x, Z = z] - \tilde{\beta}_n(x_2)' E[\rho_n(V) | X = x, Z = z]| \\ &= |M_n[\beta_n(\cdot)' E[\rho_n(V) | X = x, Z = z]](x_1) \\ &\quad - M_n[\beta_n(\cdot)' E[\rho_n(V) | X = x, Z = z]](x_2)| \\ &\leq \tilde{c} \frac{\xi_{\rho,n}}{b_n} \|x_1 - x_2\|_2 \end{aligned}$$

And thus $\tilde{\beta}_n(\cdot)' E[\rho_n(V) | X = x, Z = z] \in \Lambda_1^{\dim(X)}(\tilde{c} \frac{\xi_{\rho,n}}{b_n})$, where \tilde{c} is some constant that is independent of n .

Next we show that replacing β_n with $\tilde{\beta}_n$ does not lose us much in terms of approximation error. Adding and subtracting terms and applying the triangle inequality we get:

$$\begin{aligned} &|g(x, z) - \tilde{\beta}_n(x)' E[\rho_n(V) | X = x, Z = z]| \\ &\leq \left| \frac{\int_{\mathcal{X} \cup B_{x,b_n}} (g(x', z) - \beta_n(x')' E[\rho_n(V) | X = x', Z = z]) dx'}{\int_{\mathcal{X} \cup B_{x,b_n}} dx'} \right| \\ &+ |g(x, z) - M_n[g(\cdot, z)]| \\ &+ \left| \frac{\int_{\mathcal{X} \cup B_{x,b_n}} \beta_n(x')' (E[\rho_n(V) | X = x, Z = z] - E[\rho_n(V) | X = x', Z = z]) dx'}{\int_{\mathcal{X} \cup B_{x,b_n}} dx'} \right| \end{aligned}$$

For the first term on the RHS of the inequality note that:

$$\begin{aligned}
& \left| \left(\int_{\mathcal{X} \cup B_{x, b_n}} dx' \right)^{-1} \int_{\mathcal{X} \cup B_{x, b_n}} (g(x', z) - \beta_n(x')' E[\rho_n(V) | X = x', Z = z]) dx' \right| \\
& \leq \sup_{x' \in \mathcal{X}} |g(x', z) - \beta_n(x')' E[\rho_n(V) | X = x', Z = z]| \\
& \lesssim \ell_{\rho, n}(s_1)
\end{aligned}$$

Where, for the last step we have used (.18). Next, it is easy to see that for any $\delta \in \Lambda_s^{\dim(X)}(c)$ we have:

$$|\delta(x) - M_n[\delta](x)| \leq c b_n^{\min\{s, 1\}}$$

And so, using Assumption 5.3.ii, the second term satisfies:

$$|g(x, z) - M_n[g(\cdot, z)]| \leq c_3 b_n^{\min\{s_3, 1\}}$$

For notational convenience define:

$$q(v, x, x', z) = \frac{dF_{(X, Z, V)}}{dF_{(X, Z)} \otimes F_V}(x, z, v) - \frac{dF_{(X, Z, V)}}{dF_{(X, Z)} \otimes F_V}(x', z, v)$$

For the third term, note that:

$$\begin{aligned}
& \left| \int_{\mathcal{X} \cap B_{x, b_n}} \beta_n(x')' (E[\rho_n(V) | X = x, Z = z] - E[\rho_n(V) | X = x', Z = z]) dx' \right| \\
& = \left| \int_{\mathcal{X} \cap B_{x, b_n}} E[\beta_n(x')' \rho_n(V) q(V, x, x', z)] dx' \right| \\
& \leq \int_{\mathcal{X} \cap B_{x, b_n}} |E[\beta_n(x')' \rho_n(V) q(V, x, x', z)]| dx' \\
& \leq \int_{\mathcal{X} \cap B_{x, b_n}} E[(\beta_n(x')' \rho_n(V))^2]^{1/2} E[q(V, x, x', z)^2]^{1/2} dx' \\
& \leq \bar{D} \int_{\mathcal{X} \cap B_{x, b_n}} E[q(V, x, x', z)^2]^{1/2} dx'
\end{aligned}$$

Where the first equality follows by the Radon-Nikodym theorem, the subsequent inequality by Jensen's inequality, the second inequality by Cauchy-Schwartz, and the final inequality by (.16). Note that by the reverse triangle inequality:

$$\begin{aligned}
& |E[q(V, x, x_1, z)^2]^{1/2} - E[q(V, x, x_2, z)^2]^{1/2}| \\
& \leq E[|q(V, x, x_1, z) - q(V, x, x_2, z)|^2]^{1/2} \\
& = E \left[\left(\frac{dF_{(X, Z, V)}}{dF_{(X, Z)} \otimes F_V}(x_1, z, V) - \frac{dF_{(X, Z, V)}}{dF_{(X, Z)} \otimes F_V}(x_2, z, V) \right)^2 \right]^{1/2}
\end{aligned}$$

And by Assumption 5.3.i the final term above is bounded by $c_2 \|x_1 - x_2\|_2^{\min\{s_2, 1\}}$. So we get:

$$\left| \frac{\int_{\mathcal{X} \cap B_{x, b_n}} \beta_n(x')' (E[\rho_n(V)|X = x, Z = z] - E[\rho_n(V)|X = x', Z = z]) dx'}{\int_{\mathcal{X} \cap B_{x, b_n}} dx'} \right| \leq c_2 b_n^{\min\{s_2, 1\}}$$

And so in all:

$$\begin{aligned} & |g(x, z) - \tilde{\beta}_n(x)' E[\rho_n(V)|X = x, Z = z]| \\ & \leq \frac{\tilde{c}_1}{\sqrt{\underline{c}}} \bar{D} \ell_{\rho, n}(s_1) + c_1 b_n^{\min\{s_2, 1\}} + c_2 b_n^{\min\{s_3, 1\}} \end{aligned}$$

Now, let $G_n = E[\chi_n(X)\chi_n(X)']$ and define the matrix θ_n by:

$$\theta_n = G_n^{-1} E[\chi_n(X)' \tilde{\beta}_n(X)']$$

Recall that $\tilde{\beta}_n(\cdot)' E[\rho_n(V)|X = x, Z = z] \in \Lambda_1^{\dim(X)}(\tilde{c} \frac{\xi_{\rho, n}}{b_n})$, by Assumption 5.1.ii we get:

$$\begin{aligned} & |\chi_n(x)' \theta_n E[\rho_n(V)|X = x, Z = z] - \tilde{\beta}_n(x)' E[\rho_n(V)|X = x, Z = z]| \\ & = |\chi_n(x)' G_n^{-1} E[\chi_n(X)' \tilde{\beta}_n(X)' E[\rho_n(V)|X = x, Z = z]] \\ & \quad - \tilde{\beta}_n(x)' E[\rho_n(V)|X = x, Z = z]| \\ & \lesssim \frac{\xi_{\rho, n}}{b_n} \ell_{\chi, n}(1) \end{aligned}$$

And so, by the triangle inequality:

$$\begin{aligned} & |g(x, z) - \chi_n(x)' \theta_n E[\rho_n(V)|X = x, Z = z]| \\ & \leq |g(x, z) - \tilde{\beta}_n(x)' E[\rho_n(V)|X = x, Z = z]| \\ & \quad + |\chi_n(x)' \theta_n E[\rho_n(V)|X = x, Z = z] - \tilde{\beta}_n(x)' E[\rho_n(V)|X = x, Z = z]| \\ & \lesssim \ell_{\rho, n}(s_1) + b_n^{\min\{s_2, 1\}} + b_n^{\min\{s_3, 1\}} + \frac{\xi_{\rho, n}}{b_n} \ell_{\chi, n}(1) \end{aligned}$$

Choosing b_n rate-optimally the above gives:

$$\begin{aligned} & |g(x, z) - \chi_n(x)' \theta_n E[\rho_n(V)|X = x, Z = z]| \\ & \leq O\left(\ell_{\rho, n}(s_1) + (\xi_{\rho, n} \ell_{\chi, n}(1))^{\tilde{s}}\right) \end{aligned}$$

Where $\tilde{s} = \frac{\min\{s_2, s_3, 1\}}{\min\{s_2, s_3, 1\} + 1}$. Finally, By properties of least squares projection:

$$\begin{aligned} E[|\chi_n(X)' \theta_n \rho_n(v)|^2] &= E\left[|\chi_n(X)' R_n^{-1} E[\chi_n(X)' \tilde{\beta}_n(X)' \rho_n(v)]|^2\right] \\ &\leq E[|\tilde{\beta}_n(X)' \rho_n(v)|^2] \end{aligned}$$

And so:

$$\int |\chi_n(x)' \theta_n \rho_n(v)|^2 F_X \otimes F_V(d(x, v)) \leq \int |\tilde{\beta}_n(x)' \rho_n(v)|^2 F_X \otimes F_V(d(x, v))$$

Using the upper and lower bounds $\bar{c} < \infty$ and $\underline{c} > 0$ on $\frac{dF_{(X,V)}}{dF_X \otimes F_V}$, which exist by Assumption 5.2.ii, this implies:

$$\begin{aligned} E[|\chi_n(X)' \theta_n \rho_n(V)|^2] &\leq \frac{\bar{c}}{\underline{c}} E[|\tilde{\beta}_n(X)' \rho_n(V)|^2] \\ &\leq \frac{\bar{c}^2 \bar{f}}{\underline{c}^3 r \underline{f}} \bar{D}^2 \\ &\lesssim 1 \end{aligned}$$

Where the final inequality follows by (19). So $E[|\chi_n(X)' \theta_n \rho_n(V)|^2]$ is bounded above uniformly over n , and we are done. \square

Lemma 3.1. *Suppose Assumptions 1, 2, 3, and 4.i hold. Then, for any sequence of appropriately sized vectors $\{\theta_n\}_{n=1}^\infty$:*

$$\begin{aligned} |\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| &\lesssim_p E[|g(x_1, Z) - \pi_n(x_1, Z)' \theta_n|^2 | X = x_1]^{1/2} \\ &\quad + E_Z[|\pi_n(x_1, Z)'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2} \\ &\quad + |(\alpha_n(x_1, x_2) - \hat{\alpha}_n(x_1, x_2))' \hat{\theta}| \end{aligned}$$

Where the expectation operator E_Z integrates over Z (and not over $\hat{\theta}$). Suppose the constant $C(x_1, x_2)$ in Assumption 4.i is bounded uniformly over some set of values for (x_1, x_2) . Then the rate above is uniform over those values of (x_1, x_2) .

Proof. Applying the well-posedness result of Theorem 2.a with $\gamma(X, V) = \Phi(X, V)' \hat{\theta}$ we get:

$$\begin{aligned} &|\bar{y}(x_1|x_2) - E_Z[\phi_n(x_1, V)' \hat{\theta} | X = x_2]| \\ &\leq C(x_1, x_2)^{1/2} E_Z \left[\left(E_{(Y,X,V)} [Y - \phi_n(X, V)' \hat{\theta} | X, Z] \right)^2 \Big| X = x_1 \right]^{1/2} \end{aligned}$$

Substituting the definitions of α_n , π_n , and g_n we see the above is equivalent to:

$$\begin{aligned} & |\bar{y}(x_1|x_2) - \alpha_n(x_1, x_2)' \hat{\theta}| \\ & \leq C(x_1, x_2)^{1/2} E_Z [|g(x_1, Z) - \pi_n(x_1, Z)' \hat{\theta}|^2 | X = x_1]^{1/2} \end{aligned}$$

Applying the triangle inequality we get:

$$\begin{aligned} & |\bar{y}(x_1|x_2) - \alpha_n(x_1, x_2)' \hat{\theta}| \\ & \leq C(x_1, x_2)^{1/2} E [|g(x_1, Z) - \pi_n(x_1, Z)' \theta_n|^2 | X = x_1]^{1/2} \\ & \quad + C(x_1, x_2)^{1/2} E_Z [|\pi_n(x_1, Z)'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2} \end{aligned}$$

Also by the triangle inequality:

$$\begin{aligned} |\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| & \leq |\bar{y}(x_1|x_2) - \alpha_n(x_1, x_2)' \hat{\theta}| \\ & \quad + |(\alpha_n(x_1, x_2) - \hat{\alpha}_n(x_1, x_2))' \hat{\theta}| \end{aligned}$$

Combining gives the result. \square

Lemma 3.2 below simply lists some consequences of Rudelson's matrix law of large numbers [Rudelson \(1999\)](#) which are used throughout subsequent proofs.

Lemma 3.2. *Suppose that for each n , $\{\zeta_{n,i}\}_{i=1}^n$ is a sequence of iid length- $q(n)$ random vectors so that $\Xi_n = E[\zeta_{n,i} \zeta_{n,i}']$ is nonsingular. If $\frac{\text{ess sup } \|\Xi_n^{-1/2} \zeta_{n,i}\|_2^2 \log(q(n))}{n} \prec 1$ then letting $\hat{\Xi}_n = \frac{1}{n} \sum_{i=1}^n \zeta_{n,i} \zeta_{n,i}'$:*

$$\begin{aligned} \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} & \lesssim_p \frac{\text{ess sup } \|\Xi_n^{-1/2} \zeta_{n,i}\|_2^2 \log(q(n))}{n} \\ & \prec_p 1 \end{aligned}$$

Where I is the identity matrix of dimension $q(n)$. Moreover, $\|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2}\|_{op} \lesssim_p 1$, $\|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op} \lesssim_p 1$, $\|\Xi_n^{1/2} \hat{\Xi}_n^{-1/2}\|_{op} \lesssim_p 1$ and uniformly over all $\lambda \geq 0$, $\|\Xi_n^{1/2} (\hat{\Xi}_n + \lambda I)^{-1} \Xi_n^{1/2}\|_{op} \lesssim_p 1$.

Proof. The first result, $\|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} \prec_p 1$, follows immediately from the matrix LLN ([Rudelson \(1999\)](#)). By the triangle inequality:

$$\begin{aligned} \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2}\|_{op} & \leq \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} + \|I\|_{op} \\ & \lesssim_p 1 \end{aligned}$$

Next note that using the triangle inequality and elementary properties of the operator norm:

$$\begin{aligned} \|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op} & \leq \|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2} (\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I)\|_{op} + 1 \\ & \leq \|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op} \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} + 1 \end{aligned}$$

And so, if $\|I - \Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2}\|_{op} < 1$:

$$\|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op} \leq (1 - \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op})^{-1}$$

We have already established $\|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} \prec_p 1$ and so:

$$\|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op} \prec_p 1$$

Next note that $\|\hat{\Xi}_n^{-1/2} \Xi_n^{1/2}\|_{op} \leq \|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op}^{1/2}$ and $\|\Xi_n^{1/2} \hat{\Xi}_n^{-1/2}\|_{op} \leq \|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2}\|_{op}^{1/2}$. Using these two inequalities, the triangle inequality, and elementary properties of the operator norm:

$$\begin{aligned} & \|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2}\|_{op} \\ & \leq \|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2} - I\|_{op} + 1 \\ & = \|\hat{\Xi}_n^{-1/2} \Xi_n^{1/2} (\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I) \Xi_n^{1/2} \hat{\Xi}_n^{-1/2}\|_{op} + 1 \\ & \leq \|\hat{\Xi}_n^{-1/2} \Xi_n^{1/2}\|_{op} \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} \|\Xi_n^{1/2} \hat{\Xi}_n^{-1/2}\|_{op} + 1 \\ & \leq \|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op}^{1/2} \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} \|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2}\|_{op}^{1/2} + 1 \end{aligned}$$

Suppose that $\|\Xi_n^{1/2} \hat{\Xi}_n^{-1/2}\|_{op} \geq 1$. Then $\|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2}\|_{op} \geq 1$ and thus $\|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2}\|_{op} \geq \|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2}\|_{op}^{1/2}$, so we get:

$$\begin{aligned} & \|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2}\|_{op} \\ & \leq \|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op}^{1/2} \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} \|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2}\|_{op} \quad (.21) \\ & + 1 \quad (.22) \end{aligned}$$

By our earlier results, with probability approaching 1, $\|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op}^{1/2} \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} < 1$, in which case from (.22) we get:

$$\|\hat{\Xi}_n^{-1/2} \Xi_n \hat{\Xi}_n^{-1/2}\|_{op} \leq (1 - \|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op}^{1/2} \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op})^{-1} \quad (.23)$$

We showed earlier that if $\|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op} < 1$ then:

$$\|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op} \leq (1 - \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op})^{-1} \quad (.24)$$

In all, combining (.23) and (.24) we get that if $\|\Xi_n^{1/2} \hat{\Xi}_n^{-1/2}\|_{op} \geq 1$ then with probability approaching 1:

$$\|\Xi_n^{1/2} \hat{\Xi}_n^{-1} \Xi_n^{1/2}\|_{op} \leq \frac{1 - \|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op}}{1 - 2\|\Xi_n^{-1/2} \hat{\Xi}_n \Xi_n^{-1/2} - I\|_{op}}$$

Note that $\|\Xi_n^{1/2}\hat{\Xi}_n^{-1/2}\|_{op} \leq \|\Xi_n^{1/2}\hat{\Xi}_n^{-1}\Xi_n^{1/2}\|_{op}^{1/2}$. Therefore, without assuming $\|\Xi_n^{1/2}\hat{\Xi}_n^{-1/2}\|_{op} \geq 1$ we have that, with probability approaching 1:

$$\|\Xi_n^{1/2}\hat{\Xi}_n^{-1/2}\|_{op} \leq \min \left\{ 1, \sqrt{\frac{1 - \|I - \Xi_n^{-1/2}\hat{\Xi}_n\Xi_n^{-1/2}\|_{op}}{1 - 2\|I - \Xi_n^{-1/2}\hat{\Xi}_n\Xi_n^{-1/2}\|_{op}}} \right\} \\ \lesssim_p 1$$

Finally note that:

$$\|\Xi_n^{1/2}(\hat{\Xi}_n + \lambda I)^{-1}\Xi_n^{1/2}\|_{op} = \|(\Xi_n^{-1/2}\hat{\Xi}_n\Xi_n^{-1/2} + \lambda\Xi_n^{-1})^{-1}\|_{op} \\ \leq \|(\Xi_n^{-1/2}\hat{\Xi}_n\Xi_n^{-1/2})^{-1}\|_{op} \\ = \|\Xi_n^{1/2}\hat{\Xi}_n^{-1}\Xi_n^{1/2}\|_{op} \\ \lesssim_p 1$$

Where the first inequality holds because $\lambda\Xi_n^{-1}$ is positive definite and for any positive definite matrices A and B we have $\|(A+B)^{-1}\|_{op} \leq \|A^{-1}\|_{op}$. \square

Lemma 3.3. *Let Assumptions 5.1.i, 5.1.ii, and 5.2.ii hold. Then there exist constants $\underline{c} > 0$ and $\bar{c} < \infty$ so that for all n and $\theta \in \mathbb{R}^{k(n)l(n)}$:*

$$\mu_{\min}(Q_n)\mu_{\min}(Q_n)\underline{c}\|\theta\|_2^2 \\ \leq E[|\phi_n(X, V)' \theta|^2] \\ \leq \mu_{\max}(Q_n)\mu_{\max}(Q_n)\bar{c}\|\theta\|_2^2$$

Where $\mu_{\min}(Q_n)$ and $\mu_{\max}(Q_n)$ respectively denote smallest and largest eigenvalues of Q_n and likewise for G_n . Furthermore, for all n , all $\theta \in \mathbb{R}^{k(n)l(n)}$ and F_X -almost all x :

$$E[|\phi_n(x, V)' \theta|^2] \leq \frac{\xi_{X,n}^2}{\underline{c}} E[|\phi_n(X, V)' \theta|^2]$$

Proof. Applying the separability of ϕ_n and properties of the Kronecker product we have that for any $\theta \in \mathbb{R}^{k(n)l(n)}$:

$$E[|\phi_n(X, V)' \theta|^2] = E[\theta' \phi_n(X, V) \phi_n(X, V)' \theta] \\ = \theta' E[(\rho_n(V) \otimes \chi_n(X)) (\rho_n(V) \otimes \chi_n(X))'] \theta \\ = E \left[\theta' [(\rho_n(V) \rho_n(V)') \otimes (\chi_n(X) \chi_n(X)')] \theta \right] \quad (.25)$$

By Assumption 5.2.ii there exist constants $\underline{c} > 0$ and $\bar{c} < \infty$ so that $\underline{c} \leq \frac{dF_{(X,V)}}{dF_X \otimes F_V}(x, v) \leq \bar{c}$ for $F_{(X,V)}$ -almost all (x, v) . Therefore, for any $\theta \in \mathbb{R}^{k(n)l(n)}$:

$$\begin{aligned}
& \underline{c}\theta' \left(E[\rho_n(V)\rho_n(V)'] \otimes E[\chi_n(X)\chi_n(X)'] \right) \theta \\
& \leq E \left[\theta' [(\rho_n(V)\rho_n(V)') \otimes (\chi_n(X)\chi_n(X)')] \theta \right] \\
& \leq \bar{c}\theta' \left(E[\rho_n(V)\rho_n(V)'] \otimes E[\chi_n(X)\chi_n(X)'] \right) \theta
\end{aligned}$$

By elementary properties of the Kronecker product, $A \otimes B$ is non-singular if and only if A and B are each non-singular. By Assumptions 5.1.i and 5.1.ii, $E[\rho_n(V)\rho_n(V)']$ and $E[\chi_n(X)\chi_n(X)']$ are non-singular and thus so is their Kronecker product. Substituting (.25) the above implies:

$$\begin{aligned}
& \underline{c} \| (E[\rho_n(V)\rho_n(V)'] \otimes E[\chi_n(X)\chi_n(X)'])^{-1} \|_{op}^{-1} \|\theta\|_2^2 \\
& \leq E[|\phi_n(X, V)\theta|^2] \\
& \leq \bar{c} \| E[\rho_n(V)\rho_n(V)'] \otimes E[\chi_n(X)\chi_n(X)'] \|_{op} \|\theta\|_2^2
\end{aligned}$$

By elementary properties of the Kronecker product, the smallest eigenvalue of $A \otimes B$ (equal to $\|(A \otimes B)^{-1}\|_{op}^{-1}$ if A and B are symmetric and non-singular) is the product of the smallest eigenvalues of A and B and the largest eigenvalue of $A \otimes B$ (equal to $\|A \otimes B\|_{op}$ if A and B are symmetric and non-singular) is the product of the largest eigenvalues of the two matrices. Therefore:

$$\mu_{min}(Q_n)\mu_{min}(Q_n)\underline{c}\|\theta\|_2^2 \leq E[|\phi_n(X, V)\theta|^2] \leq \mu_{max}(Q_n)\mu_{max}(Q_n)\bar{c}\|\theta\|_2^2$$

Now for the second statement of the lemma. Let ι be the function that maps a length- $k(n)l(n)$ column vector θ to a $k(n)$ -by- $l(n)$ matrix $\tilde{\theta}$ so that the (j, k) entry of $\tilde{\theta}$ is the $(j-1)l(n)+k$ -th entry of θ . Then for any length- $k(n)l(n)$ column vector θ , $(\rho_n(v) \otimes \chi_n(x))' \theta = \rho_n(v)' \iota(\theta) \chi_n(x)$. Note that for any $\theta \in \mathbb{R}^{k(n)l(n)}$:

$$\begin{aligned}
E[(\rho_n(V)' \iota(\theta) \chi_n(x))^2] &= E[\|(\chi_n(x)\chi_n(x)')^{1/2} \iota(\theta)' \rho_n(V)\|_2^2] \\
&\leq \|(\chi_n(x)\chi_n(x)')^{1/2} G_n^{-1/2}\|_{op}^2 E[\|G_n^{1/2} \iota(\theta)' \rho_n(V)\|_2^2]
\end{aligned}$$

Again, by Assumption 5.2.ii there exists $\bar{c} < \infty$ and $\underline{c} > 0$ so that $\underline{c} \leq \frac{dF_{X \otimes V}(x, v)}{dF_X \otimes F_V}(x, v) \leq \bar{c}$ for $F_{(X, V)}$ -almost all (x, v) , and so:

$$\begin{aligned}
E[\|G_n^{1/2} \iota(\theta)' \rho_n(V)\|_2^2] &= \int |\chi_n(x)' \iota(\theta)' \rho_n(v)|^2 F_X \otimes F_V(d(x, v)) \\
&\leq \frac{1}{\underline{c}} E[|\chi_n(X)' \iota(\theta)' \rho_n(V)|^2]
\end{aligned}$$

And further, note that:

$$\begin{aligned} \|(\chi_n(x)\chi_n(x)')^{1/2}G_n^{-1/2}\|_{op}^2 &= \|G_n^{-1/2}\chi_n(x)\|_2^2 \\ &\leq \xi_{\chi,n}^2 \end{aligned}$$

Combining we get:

$$E[(\rho_n(V)'\iota(\theta)\chi_n(x))^2] \leq \frac{1}{\underline{c}}\xi_{\chi,n}^2 E[(\rho_n(V)'\iota(\theta)\chi_n(X))^2]$$

Or equivalently:

$$E[|\phi_n(x, V)'\theta|^2] \leq \frac{1}{\underline{c}}\xi_{\chi,n}^2 E[|\phi_n(X, V)'\theta|^2]$$

□

Lemma 3.4. *Let Assumptions 5.1.i-iii, 5.2.i-ii, 5.3.i, 5.3.iii, and 5.4.i-iii hold and let $\ell_{\psi,n}(s_4) \prec 1$. Then uniformly over all $\theta \in \mathbb{R}^{k(n)l(n)}$ and F_X -almost all x :*

$$\begin{aligned} &E_Z \left[\left(\pi_n(x, Z)'\theta - \hat{\pi}_n(x, Z)'\theta \right)^2 \middle| X = x \right]^{1/2} \\ &\lesssim_p E[|\phi_n(X, V)'\theta|^2]^{1/2} \xi_{\chi,n} \xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ &+ E[|\phi_n(X, V)'\theta|^2]^{1/2} \xi_{\chi,n} (\xi_{\Omega,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \ell_{\psi,n}(s_4)) \end{aligned}$$

Furthermore, uniformly over all $\theta \in \mathbb{R}^{k(n)l(n)}$

$$\begin{aligned} &\left(\frac{1}{n} \sum_{i=1}^n |(\pi_{n,i} - \hat{\pi}_{n,i})'\theta|^2 \right)^{1/2} \\ &\lesssim_p E[|\phi_n(X, V)'\theta|^2]^{1/2} \xi_{\chi,n} (\min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ &+ E[|\phi_n(X, V)'\theta|^2]^{1/2} \xi_{\chi,n} (\lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \ell_{\psi,n}(s_4)) \end{aligned}$$

Proof. For each n and i define the length- $k(n)l(n)$ column vector $\epsilon_{n,i} = \rho_{n,i} - E[\rho_n(V)|X_i, Z_i]$. By construction:

$$E[Q_n^{-1/2}\epsilon_{n,i}|X_i, Z_i] = 0$$

Further, note that:

$$\begin{aligned} E[||Q_n^{-1/2}\epsilon_{n,i}||_2^2] &\leq 2E[||Q_n^{-1/2}\rho_{n,i}||_2^2] + 2E[||E[Q_n^{-1/2}\rho_n(V)|X_i, Z_i]||_2^2] \\ &\leq 4E[||Q_n^{-1/2}\rho_{n,i}||_2^2] \\ &= 4k(n) \end{aligned} \tag{.26}$$

The first inequality above follows by the definition of $\epsilon_{n,i}$ and Young's inequality and the second follows by positivity of the variance. The equality then follows by definition of Q_n . Next, note that with probability 1:

$$\begin{aligned}
\|Q_n^{-1/2}\epsilon_{n,i}\|_2 &\leq \|Q_n^{-1/2}\rho_{n,i}\|_2 + \|E[Q_n^{-1/2}\rho_n(V)|X_i, Z_i]\|_2 \\
&\leq \text{ess sup } \|Q_n^{-1/2}\rho_n(V)\|_2 + \text{ess sup } \|E[Q_n^{-1/2}\rho_n(V)|X, Z]\|_2 \\
&\leq 2\text{ess sup } \|Q_n^{-1/2}\rho_n(V)\|_2 \\
&\leq 2\xi_{\rho,n}
\end{aligned} \tag{.27}$$

Where the first inequality follows by the triangle inequality and definition of $\epsilon_{n,i}$, and the third inequality holds because Jensen's inequality gives:

$$\|E[Q_n^{-1/2}\rho_n(V)|X, Z]\|_2^2 \leq E[\|Q_n^{-1/2}\rho_n(V)\|_2^2|X, Z]$$

and the RHS above is clearly bounded by $\text{ess sup } \|Q_n^{-1/2}\rho_n(V)\|_2$. For each n define the $m(n)$ -by- $k(n)$ matrix β_n by:

$$\beta_n = \Omega_n^{-1}E[\psi_n(X, Z)E[\rho_n(V)|X, Z]']$$

Then define r_n by:

$$r_n(x, z) = E[\rho_n(V)|X = x, Z = z] - \beta_n'\psi_n(x, z)$$

Note that $E[\psi_{n,i}r_n(X_i, Z_i)'] = 0$. By the Radon-Nikodym Theorem, Assumption 5.2.i implies that for any $\theta \in \mathbb{R}^{k(n)}$:

$$E[\rho_n(V)'\theta|X = x, Z = z] = E[\rho_n(V)'\theta \frac{dF_{(X,Z,V)}}{d(F_{(X,Z)} \otimes F_V)}(x, z, V)]$$

By Assumption 5.3.iii, for F_V -almost all v , $\frac{dF_{(X,Z,V)}}{d(F_{(X,Z)} \otimes F_V)}(\cdot, \cdot, v) \in \Lambda_{s_4}^{\dim(X,Z)}(c_4)$. So for any vector $q \in \mathbb{N}_0^{\dim(X,Z)}$ with $\|q\|_1 \leq [s_4]$, the partial derivative $D_q \left[\frac{dF_{(X,Z,V)}}{d(F_{(X,Z)} \otimes F_V)}(\cdot, \cdot, v) \right](x, z)$ exist and has magnitude less than c_4 uniformly over $F_{(X,Z,V)}$ -almost all (x, z, v) . By the dominated convergence theorem we can differentiate under the integral to get:

$$\begin{aligned}
&|D_q \left[E[\rho_n(V)'\theta|X = \cdot, Z = \cdot] \right](x, z)| \\
&= |E \left[\rho_n(V)'\theta D_q \left[\frac{dF_{(X,Z,V)}}{d(F_{(X,Z)} \otimes F_V)}(\cdot, \cdot, V) \right](x, z) \right]| \\
&\leq E[|\rho_n(V)'\theta|^2]^{1/2} \text{ess sup } |D_q \left[\frac{dF_{(X,Z,V)}}{d(F_{(X,Z)} \otimes F_V)}(\cdot, \cdot, V) \right](x, z)| \\
&\leq c_4 \|Q_n^{1/2}\theta\|_2
\end{aligned} \tag{.28}$$

Moreover, for any $\|q\|_1 = \lfloor s_4 \rfloor$, we have:

$$\begin{aligned} & \left| D_q \left[\frac{dF_{(X,Z,V)}}{dF_{(X,Z)} \otimes F_V}(\cdot, \cdot, v) \right] (x_1, z_1) - D_q \left[\frac{dF_{(X,Z,V)}}{dF_{(X,Z)} \otimes F_V}(\cdot, \cdot, v) \right] (x_2, z_2) \right| \\ & \leq c_4 \left(\|x_1 - x_2\|_2^2 + \|z_1 - z_2\|_2^2 \right)^{\frac{s_4 - \lfloor s_4 \rfloor}{2}} \end{aligned}$$

Again, differentiating under the integral:

$$\begin{aligned} & \left| D_q \left[E[\rho_n(V)' \theta | X = x_1, Z = z_1] \right] - D_q \left[E[\rho_n(V)' \theta | X = x_2, Z = z_2] \right] \right| \\ & = \left| E \left[\rho_n(V)' \theta \left(D_q \left[\frac{dF_{(X,Z,V)}}{dF_{(X,Z)} \otimes F_V}(\cdot, \cdot, V) \right] (x_1, z_1) - D_q \left[\frac{dF_{(X,Z,V)}}{dF_{(X,Z)} \otimes F_V}(\cdot, \cdot, V) \right] (x_2, z_2) \right) \right] \right| \\ & \leq E[|\rho_n(V)' \theta|^2]^{1/2} \\ & \times \text{ess sup} \left| D_q \left[\frac{dF_{(X,Z,V)}}{dF_{(X,Z)} \otimes F_V}(\cdot, \cdot, V) \right] (x_1, z_1) - D_q \left[\frac{dF_{(X,Z,V)}}{dF_{(X,Z)} \otimes F_V}(\cdot, \cdot, V) \right] (x_2, z_2) \right| \\ & \leq c_4 \|Q_n^{1/2} \theta\|_2 \left(\|x_1 - x_2\|_2^2 + \|z_1 - z_2\|_2^2 \right)^{\frac{s_4 - \lfloor s_4 \rfloor}{2}} \quad (.29) \end{aligned}$$

(.28) and (.29) together imply:

$$(x, z) \mapsto E[\rho_n(V)' \theta | X = x, Z = z] \in \Lambda_{s_4}^{\dim(X,Z)}(c_4 \|Q_n^{1/2} \theta\|_2)$$

Using Assumption 5.1.iii, the above implies that uniformly over all $\theta \in \mathbb{R}^{k(n)}$ and $F_{(X,Z)}$ -almost all (x, z) :

$$\frac{r_n(x, z)' \theta}{\|Q_n^{1/2} \theta\|_2} \lesssim \ell_{\psi, n}(s_4)$$

Which in turn implies that (uniformly):

$$\|Q_n^{-1/2} r_n(x, z)\|_2 \lesssim \ell_{\psi, n}(s_4) \quad (.30)$$

Now decompose:

$$\pi_n(x, z)' \theta = \psi_n(x, z)' \beta_n \iota(\theta) \chi(x) + r_n(x, z)' \iota(\theta) \chi(x)$$

Recall that for $\theta \in \mathbb{R}^{k(n)\iota(n)}$, $\phi_n(x, v)' \theta = \rho_n(v)' \iota(\theta) \chi_n(x)$, where ι is defined as in Lemma 3.3. Substituting the above and using the formulas for π_n and $\hat{\pi}_n$ we get:

$$\begin{aligned} & (\pi_n(x, z) - \hat{\pi}_n(x, z))' \theta \\ & = r_n(x, z)' \iota(\theta) \chi_n(x) \\ & - \psi_n(x, z)' \hat{\Omega}_{\lambda_{2,n}}^{-1} \frac{1}{n} \sum_{i=1}^n \psi_{n,i}(\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x) \\ & - \psi_n(x, z)' \hat{\Omega}_{\lambda_{2,n}}^{-1} \lambda_{2,n} \beta_n \iota(\theta) \chi_n(x) \end{aligned}$$

Where $r_{n,i} = r_n(X_i, Z_i)$. By the triangle inequality:

$$\begin{aligned}
& E_Z \left[\left(\pi_n(x, Z)' \theta - \hat{\pi}_n(x, Z)' \theta \right)^2 \middle| X = x \right]^{1/2} \\
& \leq E[|r_n(x, Z)' \iota(\theta) \chi_n(x)|^2 | X = x]^{1/2} \\
& + \|\tilde{\Omega}_n(x)^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \frac{1}{n} \sum_{i=1}^n \psi_{n,i}(\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x)\|_2 \\
& + \|\tilde{\Omega}_n(x)^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \lambda_{2,n} \beta_n \iota(\theta) \chi_n(x)\|_2
\end{aligned} \tag{.31}$$

Taking (.30) and applying Cauchy-Schwartz, uniformly over $\theta \in \mathbb{R}^{k(n), l(n)}$ and $F_{(X,Z)}$ -almost all (x, z) and F_X -almost all x_1 :

$$\begin{aligned}
|r_n(x, z)' \iota(\theta) \chi_n(x_1)| & \leq \|r_n(x, z) Q_n^{-1/2}\|_2 \|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2 \\
& \lesssim E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{X,n} \ell_{\psi,n}(s_4)
\end{aligned} \tag{.32}$$

Where the final equality uses Lemma 3.3 which states that $\|Q_n^{-1/2} \iota(\theta) \chi_n(x)\|_2 \leq O(E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{X,n})$. This implies (again uniformly):

$$E[|r_n(x, Z)' \iota(\theta) \chi_n(x)|^2 | X = x]^{1/2} \lesssim E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{X,n} \ell_{\psi,n}(s_4)$$

Using the definition of the operator norm and recalling that $\xi_{\Omega,n}$ is defined as the smallest almost sure bound on $\|\tilde{\Omega}_n(X)^{1/2} \Omega_n^{1/2}\|_{op}$ we get:

$$\begin{aligned}
& \|\tilde{\Omega}_n(x)^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \frac{1}{n} \sum_{i=1}^n \psi_{n,i}(\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x)\|_2 \\
& \leq \xi_{\Omega,n} \|\Omega_n^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \Omega_n^{1/2}\|_{op} \left\| \frac{1}{n} \sum_{i=1}^n (\Omega_n^{-1/2} \psi_{n,i})(\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x) \right\|_2 \\
& = \xi_{\Omega,n} \|\Omega_n^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \Omega_n^{1/2}\|_{op} \\
& \times \left(\sum_{l=1}^{m(n)} \left| \frac{1}{n} \sum_{i=1}^n [\Omega_n^{-1/2} \psi_{n,i}]_l (\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x) \right|^2 \right)^{1/2} \\
& \leq \xi_{\Omega,n} \|\Omega_n^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \Omega_n^{1/2}\|_{op} \|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2 \\
& \times \left(\sum_{l=1}^{m(n)} \left\| \frac{1}{n} \sum_{i=1}^n [\Omega_n^{-1/2} \psi_{n,i}]_l Q_n^{-1/2} (\epsilon_{n,i} + r_{n,i}) \right\|_2^2 \right)^{1/2}
\end{aligned}$$

Where $[\Omega_n^{-1/2} \psi_{n,i}]_l$ is the l^{th} component of the vector $\Omega_n^{-1/2} \psi_{n,i}$. Note that:

$$\begin{aligned}
& E \sum_{l=1}^{m(n)} \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n [\Omega_n^{-1/2} \psi_{n,i}]_l Q_n^{-1/2} (\epsilon_{n,i} + r_{n,i}) \right\|_2^2 \\
&= E \left[\left\| \Omega_n^{-1/2} \psi_{n,i} \right\|_2^2 \left\| \epsilon_{n,i} + r_{n,i} \right\|_2^2 \right] \\
&\leq 2E \left[\left\| \Omega_n^{-1/2} \psi_{n,i} \right\|_2^2 \left\| Q_n^{-1/2} \epsilon_{n,i} \right\|_2^2 \right] + 2E \left[\left\| \Omega_n^{-1/2} \psi_{n,i} \right\|_2^2 \left\| Q_n^{-1/2} r_{n,i} \right\|_2^2 \right] \\
&\leq 2 \min \left\{ \xi_{\psi,n}^2 E \left[\left\| Q_n^{-1/2} \epsilon_{n,i} \right\|_2^2 \right], \xi_{\rho,n}^2 E \left[\left\| \Omega_n^{-1/2} \psi_{n,i} \right\|_2^2 \right] \right\} \\
&\quad + 2 \text{ess sup} \left\| Q_n^{-1/2} r_n(X, Z) \right\|_2^2 E \left[\left\| \Omega_n^{-1/2} \psi_{n,i} \right\|_2^2 \right] \\
&\leq 4 \min \{ \xi_{\psi,n}^2 k(n), \xi_{\rho,n}^2 m(n) \} + 2O(m(n)^2 \ell_{\psi,n}(s_4)) \\
&= O \left(\min \{ \xi_{\psi,n}^2 k(n), \xi_{\rho,n}^2 m(n) \} \right) \tag{.33}
\end{aligned}$$

Where the first equality follows because $E[\psi_{n,i}(\epsilon_{n,i} + r_{n,i})'] = 0$ for all n and the data are iid. The first inequality follows from Young's inequality, the second by the Holder inequality, the third by (.26), (.27), and (.30). The final equality follows because $\xi_{\rho,n}$ is increasing and $\ell_{\psi,n}(s_4)$ is decreasing.

In all:

$$\begin{aligned}
& E \left[\left\| \tilde{\Omega}_n(x)^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \frac{1}{n} \sum_{i=1}^n \psi_{n,i}(\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x) \right\|_2 \right] \\
&\lesssim_p \frac{2\xi_{\Omega,n} \left\| Q_n^{1/2} \iota(\theta) \chi_n(x) \right\|_2 \left\| \Omega_n^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \Omega_n^{1/2} \right\|_{op}}{\sqrt{n}} \min \left\{ \sqrt{\xi_{\psi,n}^2 k(n)}, \sqrt{\xi_{\rho,n}^2 m(n)} \right\}
\end{aligned}$$

By Lemma 3.2, we have $\left\| \Omega_n^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \Omega_n^{1/2} \right\|_{op} \lesssim_p 1$, and again by Lemma 3.3, $\left\| Q_n^{-1/2} \iota(\theta) \chi_n(x) \right\|_2 \lesssim E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{\chi,n}$ and so, uniformly over $\theta \in \mathbb{R}^{k(n)l(n)}$ and F_X -almost all x :

$$\begin{aligned}
& E \left[\left\| \tilde{\Omega}_n(x)^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \frac{1}{n} \sum_{i=1}^n \psi_{n,i}(\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x) \right\|_2 \right] \\
&= \lesssim_p E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{\chi,n} \xi_{\Omega,n} \min \left\{ \sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n} \right\}
\end{aligned}$$

Finally, consider the term $\left\| \tilde{\Omega}_n(x)^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \lambda_{2,n} \beta_n \iota(\theta) \chi_n(x) \right\|_2$. Note that:

$$\begin{aligned}
& \left\| \tilde{\Omega}_n(x)^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \lambda_{2,n} \beta_n \iota(\theta) \chi_n(x) \right\|_2 \\
&\leq \lambda_{2,n} \xi_{\Omega,n} \left\| \Omega_n^{1/2} \hat{\Omega}_{\lambda_{2,n}}^{-1} \Omega_n^{1/2} \right\|_{op} \left\| \Omega_n^{-1} \right\|_{op} \left\| \Omega_n^{1/2} \beta_n \iota(\theta) \chi_n(x) \right\|_2
\end{aligned}$$

By Lemma 3.2 $\|\Omega_n^{1/2}\hat{\Omega}_{\lambda_{2,n}}^{-1}\Omega_n^{1/2}\|_{op} \lesssim_p 1$. Furthermore:

$$\begin{aligned} \|\Omega_n^{1/2}\beta_n\iota(\theta)\chi_n(x)\|_2^2 &= E[|\psi_n(X, Z)\beta_n\iota(\theta)\chi_n(x)|^2] \\ &\leq E[E[\rho_n(V)|X, Z]'\iota(\theta)\chi_n(x)]^2 \\ &\leq E[\rho_n(V)'\iota(\theta)\chi_n(x)]^2 \\ &= \|Q^{1/2}\iota(\theta)\chi_n(x)\|_2^2 \\ &\lesssim \xi_{\chi,n}^2 E[|\phi_n(X, V)'\theta|^2] \end{aligned}$$

Where the first inequality holds by the properties of least-squares projection, the second inequality by positivity of the variance, and the final inequality by Lemma 3.3. Thus we have uniformly over $\theta \in \mathbb{R}^{k(n)\iota(n)}$ and F_X -almost all x :

$$\|\tilde{\Omega}_n(x)^{1/2}\hat{\Omega}_{\lambda_{2,n}}^{-1}\lambda_{2,n}\beta_n\iota(\theta)\chi_n(x)\|_2 \leq O(E[|\phi_n(X, V)'\theta|^2]^{1/2}\|\Omega_n^{-1}\|_{op}\lambda_{2,n}\xi_{\Omega,n}\xi_{\chi,n})$$

Substituting the results above into (.31) we get that uniformly over $\theta \in \mathbb{R}^{k(n)\iota(n)}$ and F_X -almost all x :

$$\begin{aligned} &E_Z \left[(\pi_n(x, z)'\theta - \hat{\pi}_n(x, z)'\theta)^2 \middle| X = x \right]^{1/2} \\ &\lesssim_p E[|\phi_n(X, V)'\theta|^2]^{1/2} \xi_{\chi,n}\xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ &+ E[|\phi_n(X, V)'\theta|^2]^{1/2} \xi_{\chi,n}(\lambda_{2,n}\xi_{\Omega,n}\|\Omega_n^{-1}\|_{op} + \ell_{\psi,n}(s_4)) \end{aligned}$$

Next we consider $\frac{1}{n} \sum_{i=1}^n |(\pi_{n,i} - \hat{\pi}_{n,i})'\theta|^2$. By the triangle inequality, Cauchy-Schwartz and the definition of the operator norm:

$$\begin{aligned} &\left(\frac{1}{n} \sum_{i=1}^n |(\pi_{n,i} - \hat{\pi}_{n,i})'\theta|^2 \right)^{1/2} \\ &\leq \left(\frac{1}{n} \sum_{i=1}^n |r'_{n,i}\iota(\theta)\chi_{n,i}|^2 \right)^{1/2} \\ &+ \left(\|\hat{\Omega}_n^{1/2}\Omega_n^{-1/2}\|_{op} \|\Omega_n^{1/2}\hat{\Omega}_{\lambda_{2,n}}^{-1}\Omega_n^{1/2}\|_{op} \right. \\ &\times \left. \|\Omega_n^{-1/2} \frac{1}{n} \sum_{j=1}^n \psi_{n,j}(\epsilon_{n,j} + r_{n,j})'Q_n^{1/2}\|_2 \|Q_n^{-1/2}\iota(\theta)\chi_{n,i}\|_2 \right) \\ &+ \left(\|\hat{\Omega}_n^{1/2}\Omega_n^{-1/2}\|_{op} \|\Omega_n^{1/2}\hat{\Omega}_{\lambda_{2,n}}^{-1}\Omega_n^{1/2}\|_{op} \|\Omega_n^{-1}\|_{op} \right. \\ &\times \left. \|\lambda_{2,n}\Omega_n^{1/2}\beta_n Q_n^{1/2}\|_2 \|Q_n^{-1/2}\iota(\theta)\chi_{n,i}\|_2 \right) \end{aligned}$$

By (.32):

$$\left(\frac{1}{n} \sum_{i=1}^n |r'_{n,i}\iota(\theta)\chi_{n,i}|^2 \right)^{1/2} \lesssim_p \|Q_n^{1/2}\iota(\theta)\chi_n(x)\|_2 \xi_{\chi,n} \ell_{\psi,n}(s_4)$$

By Lemma 3.2 $\|\hat{\Omega}_n^{1/2}\Omega_n^{-1/2}\|_{op} \leq \|\Omega_n^{-1/2}\hat{\Omega}_n\Omega_n^{-1/2}\|_{op}^{1/2} \lesssim_p 1$ and $\|\Omega_n^{1/2}\hat{\Omega}_{\lambda_{2,n}}^{-1}\Omega_n^{1/2}\|_{op} \lesssim_p 1$, combining this, the above, and (3.3) we get:

$$\begin{aligned} & \left(\frac{1}{n} \sum_{i=1}^n |(\pi_{n,i} - \hat{\pi}_{n,i})'\theta|^2 \right)^{1/2} \\ & \lesssim_p E[|\phi_n(X, V)'\theta|^2]^{1/2} \xi_{\chi,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ & + E[|\phi_n(X, V)'\theta|^2]^{1/2} \xi_{\chi,n} (\ell_{\psi,n}(s_4) + \lambda_{2,n}\|\Omega_n^{-1}\|_{op}) \end{aligned}$$

□

Lemma 3.5. *Let Assumptions 5.1.i-iii, 5.2.i-ii, 5.3.i, 5.3.iii, and 5.4.i-iii hold. Suppose $E[|\phi_n(X, V)'\theta_n|^2]$ is bounded above uniformly over n , and uniformly over $F_{(X,Z)}$ -almost all (x, z) :*

$$|g(x, z) - \pi_n(x, z)'\theta_n| \lesssim \kappa_n$$

Suppose in addition that:

$$\lambda_{0,n} \lesssim_p \mu_{\min}(Q_n)\mu_{\min}(G_n) \frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i}\theta_n)^2$$

Then:

$$E_{(X,V)}[|\phi_n(X, V)'(\hat{\theta} - \theta_n)|^2]^{1/2} \lesssim_p \sqrt{\frac{\mu_{\max}(Q_n)\mu_{\max}(G_n)}{\mu_{\min}(Q_n)\mu_{\min}(G_n)}}$$

Proof. By supposition $E[(\phi_n(X, V)'\theta_n)^2]^{1/2} \lesssim 1$. Now recall that $\hat{\theta}$ is defined by:

$$\hat{\theta} = \arg \min_{\theta \in \mathbb{R}^{K(n)}} \frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i}\theta)^2 + \lambda_{0,n}\|\theta\|_2^2$$

and so we have:

$$\begin{aligned} \lambda_{0,n}\|\hat{\theta}\|_2^2 & \leq \frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i}\hat{\theta})^2 + \lambda_{0,n}\|\hat{\theta}\|_2^2 \\ & \leq \frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i}\theta_n)^2 + \lambda_{0,n}\|\theta_n\|_2^2 \end{aligned}$$

And so:

$$\|\hat{\theta}\|_2^2 \leq \|\theta_n\|_2^2 + \lambda_{0,n}^{-1} \frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i}\theta_n)^2$$

Using Lemma 3.3 the above implies:

$$E_{(X,Z)}[|\phi_n(X,V)'\hat{\theta}|^2] \leq \frac{\mu_{max}(Q_n)\mu_{max}(G_n)\bar{c}}{\mu_{min}(Q_n)\mu_{min}(G_n)\underline{c}} E[|\phi_n(X,V)'\theta_n|^2] \\ + \mu_{max}(Q_n)\mu_{max}(G_n)\lambda_{0,n}^{-1} \frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i}\theta_n)^2$$

By supposition:

$$\lambda_{0,n} \gtrsim \mu_{min}(Q_n)\mu_{min}(G_n) \frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i}\theta_n)^2$$

And $E[|\phi_n(X,V)'\theta_n|^2] \lesssim 1$, and so:

$$\mu_{max}(Q_n)\mu_{max}(G_n)\lambda_{0,n}^{-1} \frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i}\theta_n)^2 \\ \lesssim_p \frac{\mu_{max}(Q_n)\mu_{max}(G_n)\bar{c}}{\mu_{min}(Q_n)\mu_{min}(G_n)\underline{c}} E[|\phi_n(X,V)'\theta_n|^2]$$

Therefore:

$$E_{(X,Z)}[|\phi_n(X,V)'\hat{\theta}|^2] \lesssim_p \frac{\mu_{max}(Q_n)\mu_{max}(G_n)}{\mu_{min}(Q_n)\mu_{min}(G_n)} E[|\phi_n(X,V)'\theta_n|^2]$$

By the triangle inequality:

$$E_{(X,V)}[(\phi_n(X,V)'(\hat{\theta} - \theta_n))^2]^{1/2} \\ \leq E[(\phi_n(X,V)'\theta_n)^2]^{1/2} + E_{(X,V)}[(\phi_n(X,V)'\hat{\theta})^2]^{1/2}$$

Using $E[|\phi_n(X,V)'\theta_n|^2] \lesssim 1$ and noting that $\frac{\mu_{max}(Q_n)\mu_{max}(G_n)}{\mu_{min}(Q_n)\mu_{min}(G_n)} \leq 1$ we combine to get:

$$E_{(X,V)}[(\phi_n(X,V)'(\hat{\theta} - \theta_n))^2]^{1/2} \lesssim_p \sqrt{\frac{\mu_{max}(Q_n)\mu_{max}(G_n)}{\mu_{min}(Q_n)\mu_{min}(G_n)}}$$

□

Proposition 3.2. *Let Assumptions 5.1.i, 5.1.ii, 5.1.iii, 5.2.i, 5.2.ii, 5.3.i, 5.3.iii, and 5.4.i-iii hold. Suppose $E[|\phi_n(X,V)'\theta_n|^2]$ is bounded above uniformly over n , and uniformly over $F_{(X,Z)}$ -almost all (x, z) :*

$$|g(x, z) - \pi_n(x, z)'\theta_n| \lesssim \kappa_n$$

Suppose in addition that $\lambda_{0,n}$ goes to zero sufficiently slowly, precisely:

$$\begin{aligned}
& \lambda_{0,n}^{1/2} (\mu_{\min}(Q_n) \mu_{\min}(G_n))^{-1/2} \\
& \lesssim_p (1 + \ell_{\psi,n}(s_3)) \sqrt{m(n)/n} + \ell_{\psi,n}(s_3) + \lambda_{1,n} \|\Omega_n^{-1}\|_{op} \\
& + \xi_{\chi,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\
& + \xi_{\chi,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \xi_{\chi,n} \ell_{\psi,n}(s_4) \\
& + \kappa_n
\end{aligned}$$

Then:

$$\begin{aligned}
& E_Z [|\pi_n(x_1, Z)'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2} \\
& \lesssim_p \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} \xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} (\xi_{\Omega,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \ell_{\psi,n}(s_4)) \\
& + \xi_{\Omega,n} \lambda_{0,n}^{1/2} \underline{\mu}_n^{-1/2}
\end{aligned}$$

Where $\bar{\mu}_n = \sqrt{\mu_{\max}(Q_n) \mu_{\max}(G_n)}$ and $\underline{\mu}_n = \sqrt{\mu_{\min}(Q_n) \mu_{\min}(G_n)}$.

Proof. Step 1: Prove the conditions of Lemma 3.5 are satisfied
By the triangle inequality:

$$\begin{aligned}
\left(\frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)^2\right)^{1/2} & \leq \left(\frac{1}{n} \sum_{i=1}^n (\hat{g}_i - g_i)^2\right)^{1/2} \\
& + \left(\frac{1}{n} \sum_{i=1}^n ((\hat{\pi}_{n,i} - \pi_{n,i})' \theta_n)^2\right)^{1/2} \\
& + \left(\frac{1}{n} \sum_{i=1}^n (g_i - \pi'_{n,i} \theta_n)^2\right)^{1/2} \tag{.34}
\end{aligned}$$

For the first term, define $\epsilon_i = Y_i - g_i$ and $r_{n,i} = g_i - \psi'_{n,i} \beta_n$ where $\beta_n = \Omega_n^{-1} E[\psi_{n,i} g_i]$. Then using the formula for \hat{g}_i :

$$\begin{aligned}
\hat{g}_i - g_i & = \psi'_{n,i} \hat{\Omega}_{\lambda_{1,n}}^{-1} \frac{1}{n} \sum_{j=1}^n \psi_{n,j} (\epsilon_j + r_{n,j}) \\
& + r_{n,i} - \lambda_{1,n} \psi'_{n,i} \hat{\Omega}_{\lambda_{1,n}}^{-1} \beta_n
\end{aligned}$$

Then applying the triangle inequality and the definition of the operator norm we get:

$$\begin{aligned}
& \left(\frac{1}{n} \sum_{i=1}^n (\hat{g}_i - g_i)^2 \right)^{1/2} \\
& \leq \|\hat{\Omega}_n^{1/2} \Omega_n^{-1/2}\|_{op} \|\Omega_n^{1/2} \hat{\Omega}_{\lambda_{1,n}}^{-1} \Omega_n^{1/2}\|_{op} \left\| \frac{1}{n} \sum_{j=1}^n \Omega_n^{-1/2} \psi_{n,j} (\epsilon_j + r_{n,j}) \right\|_2 \\
& \quad + \left(\frac{1}{n} \sum_{i=1}^n r_{n,i}^2 \right)^{1/2} + \lambda_{1,n} \|\hat{\Omega}_n^{1/2} \hat{\Omega}_{\lambda_{1,n}}^{-1} \beta_n\|_2
\end{aligned}$$

By Lemma 3.2, $\|\Omega_n^{1/2} \hat{\Omega}_{\lambda_{1,n}}^{-1} \Omega_n^{1/2}\|_{op} \lesssim_p 1$ and $\|\hat{\Omega}_n^{1/2} \Omega_n^{-1/2}\|_{op} \leq \|\Omega_n^{-1/2} \hat{\Omega}_n \Omega_n^{-1/2}\|_{op}^{1/2} \lesssim_p 1$. By Assumption 5.3.ii $g \in \Lambda_{s_3}^{\dim(X,Z)}(c_3)$ and so by Assumption 5.1.iii, for some constant $\tilde{c} > 0$, $|r_{n,i}| \leq \tilde{c} \ell_{\psi,n}(s_3)$ almost surely, which implies $(\frac{1}{n} \sum_{i=1}^n r_{n,i}^2)^{1/2} \lesssim_p \ell_{\psi,n}(s_3)$ and that $E[r_{n,i}^2]^{1/2} \leq \tilde{c} \ell_{\psi,n}(s_3)$. By Assumption 5.2.iv, $E[\epsilon_{n,j}^2 | X_i, Z_i] \leq \bar{\sigma}_Y^2$ almost surely. Therefore:

$$\begin{aligned}
& E\left[\left| \Omega_n^{-1/2} \psi_{n,j} (\epsilon_{n,j} + r_{n,j}) \right|^2 \right]^{1/2} \\
& \leq E\left[\left| \Omega_n^{-1/2} \psi_{n,j} \epsilon_{n,j} \right|^2 \right]^{1/2} + E\left[\left| \Omega_n^{-1/2} \psi_{n,j} r_{n,j} \right|^2 \right]^{1/2} \\
& = E\left[\left| \Omega_n^{-1/2} \psi_{n,j} \right|^2 E[\epsilon_{n,j}^2 | X_i, Z_i] \right]^{1/2} + E\left[\left| \Omega_n^{-1/2} \psi_{n,j} r_{n,j} \right|^2 \right]^{1/2} \\
& \leq (\bar{\sigma}_Y + \tilde{c} \ell_{\psi,n}(s_3)) \sqrt{m(n)}
\end{aligned}$$

Note that $E[\Omega_n^{-1/2} \psi_{n,j} (\epsilon_{n,j} + r_{n,j})] = 0$ and observations are iid, so by the above and a Markov inequality argument:

$$\left\| \frac{1}{n} \sum_{j=1}^n \Omega_n^{-1/2} \psi_{n,j} (\epsilon_j + r_{n,j}) \right\|_2 \lesssim_p (1 + \ell_{\psi,n}(s_3)) \sqrt{\frac{m(n)}{n}}$$

Where we have used that $\ell_{\psi,n}(s_3) < 1$. Note that:

$$\begin{aligned}
& \lambda_{1,n} \|\hat{\Omega}_n^{1/2} \hat{\Omega}_{\lambda_{1,n}}^{-1} \beta_n\|_2 \\
& \leq \lambda_{1,n} \|\hat{\Omega}_n^{1/2} \Omega_n^{-1/2}\|_{op} \\
& \quad \times \|\Omega_n^{1/2} \hat{\Omega}_{\lambda_{1,n}}^{-1} \Omega_n^{1/2}\|_{op} \|\Omega_n^{-1}\|_{op} \|\Omega_n^{1/2} \beta_n\|_2
\end{aligned}$$

Again, by Lemma 3.2, $\|\Omega_n^{1/2} \hat{\Omega}_{\lambda_{1,n}}^{-1} \Omega_n^{1/2}\|_{op} \lesssim_p 1$ and $\|\hat{\Omega}_n^{1/2} \Omega_n^{-1/2}\|_{op} \lesssim_p 1$. Finally, by the properties of least squares projection

$$\begin{aligned}
\|\Omega_n^{1/2} \beta_n\|_2 & = E[|\psi'_{n,i} \beta_n|^2] \leq E[g_i^2] \\
& \leq E[Y_i^2] \\
& \lesssim 1
\end{aligned}$$

And so $\lambda_{1,n} \|\hat{\Omega}_n^{1/2} \hat{\Omega}_{\lambda_{1,n}}^{-1} \beta_n\|_2 \lesssim_p \lambda_{1,n}$. In all:

$$\left(\frac{1}{n} \sum_{i=1}^n (\hat{g}_i - g_i)^2\right)^{1/2} \lesssim_p (1 + \ell_{\psi,n}(s_3)) \sqrt{\frac{m(n)}{n}} + \ell_{\psi,n}(s_3) + \lambda_{1,n} \|\Omega_n^{-1}\|_{op}$$

By Lemma 3.4 and $E[|\phi_n(X, V)' \theta_n|^2]^{1/2} \prec 1$:

$$\begin{aligned} & \left(\frac{1}{n} \sum_{i=1}^n |(\pi_{n,i} - \hat{\pi}_{n,i})' \theta_n|^2\right)^{1/2} \\ & \lesssim_p \xi_{\chi,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ & \quad + \xi_{\chi,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \xi_{\chi,n} \ell_{\psi,n}(s_4) \end{aligned}$$

And note that because the data are iid:

$$\begin{aligned} \left(\frac{1}{n} \sum_{i=1}^n (g_i - \pi'_{n,i} \theta_n)^2\right)^{1/2} & \lesssim_p E[|g_i - \pi'_{n,i} \theta_n|^2]^{1/2} \\ & \lesssim_p \kappa_n \end{aligned}$$

Substituting the previous three rates into (.34) we get:

$$\begin{aligned} & \left(\frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)^2\right)^{1/2} \\ & \lesssim_p (1 + \ell_{\psi,n}(s_3)) \sqrt{\frac{m(n)}{n}} + \ell_{\psi,n}(s_3) + \lambda_{1,n} \|\Omega_n^{-1}\|_{op} \\ & \quad + \xi_{\chi,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ & \quad + \xi_{\chi,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \xi_{\chi,n} \ell_{\psi,n}(s_4) \\ & \quad + \kappa_n \end{aligned}$$

By supposition, $\lambda_{0,n}^{1/2} (\mu_{\min}(Q_n) \mu_{\min}(G_n))^{-1/2}$ goes to zero more slowly in probability than the RHS above, therefore:

$$\lambda_{1,n} \lesssim_p \mu_{\min}(Q_n) \mu_{\min}(G_n) \frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)^2$$

And so by Lemma 3.5:

$$E_{(X,V)} [|\phi_n(X, V)'(\hat{\theta} - \theta_n)|^2]^{1/2} \lesssim_p \sqrt{\frac{\mu_{\max}(Q_n) \mu_{\max}(G_n)}{\mu_{\min}(Q_n) \mu_{\min}(G_n)}}$$

Step 2: Derive a rate for $E_Z [|\pi_n(x_1, Z)'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2}$

By the triangle inequality:

$$\begin{aligned}
& E_Z [|\pi_n(x_1, Z)'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2} \\
& \leq E_Z [|(\pi_n(x_1, Z) - \hat{\pi}_n(x_1, Z))'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2} \\
& + E_Z [|\hat{\pi}_n(x_1, Z)'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2}
\end{aligned}$$

From Lemma 3.4 and Step 1 we immediately get that:

$$\begin{aligned}
& E_Z [|(\pi_n(x_1, Z) - \hat{\pi}_n(x_1, Z))'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2} \\
& \lesssim_p \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi, n} \xi_{\Omega, n} \min\{\sqrt{\xi_{\psi, n}^2 k(n)/n}, \sqrt{\xi_{\rho, n}^2 m(n)/n}\} \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi, n} (\xi_{\Omega, n} \lambda_{2, n} \|\Omega_n^{-1}\|_{op} + \ell_{\psi, n}(s_4))
\end{aligned}$$

Where $\bar{\mu}_n = \sqrt{\mu_{max}(Q_n)\mu_{max}(G_n)}$ and $\underline{\mu}_n = \sqrt{\mu_{min}(Q_n)\mu_{min}(G_n)}$. Now consider the term $E_Z [|\hat{\pi}_n(x_1, Z)'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2}$. Let $\hat{\beta}_n = \hat{\Omega}_{\lambda_{2, n}}^{-1} \frac{1}{n} \sum_{i=1}^n \psi_{n, i} \rho_{n, i}$ so that $\hat{\pi}_n(x, z) = (\hat{\beta}_n' \psi_n(x, z)) \otimes \chi_n(x)$. Let ι be defined as in Lemma 3.3. Then note that for any length- $k(n)l(n)$ column vector θ , $\hat{\pi}_n(x, z)' \theta = \psi_n(x, z)' \hat{\beta}_n \iota(\theta) \chi_n(x)$. Substituting we get:

$$\begin{aligned}
& E_Z [(\hat{\pi}_n(x_1, Z)'(\hat{\theta} - \theta_n))^2 | X = x_1] \\
& = E_Z [(\psi_n(x_1, Z)' \hat{\beta}_n \iota(\hat{\theta} - \theta_n) \chi_n(x_1))^2 | X = x_1] \\
& = \|E[\psi_n(X, Z) \psi_n(X, Z)' | X = x_1]^{1/2} \hat{\beta}_n \iota(\hat{\theta} - \theta_n) \chi_n(x_1)\|_2^2 \\
& = \|\hat{\Omega}_n(x_1)^{1/2} \Omega_n^{-1/2} \Omega_n^{1/2} \hat{\beta}_n \iota(\hat{\theta} - \theta_n) \chi_n(x_1)\|_2^2 \\
& \leq \|\hat{\Omega}_n(x_1)^{1/2} \Omega_n^{-1/2}\|_{op}^2 \|\Omega_n^{1/2} \hat{\beta}_n \iota(\hat{\theta} - \theta_n) \chi_n(x_1)\|_2^2 \\
& \leq \xi_{\Omega, n}^2 \|\Omega_n^{1/2} \hat{\Omega}_n^{-1/2}\|_{op}^2 \|\hat{\Omega}_n^{1/2} \hat{\beta}_n \iota(\hat{\theta} - \theta_n) \chi_n(x_1)\|_2^2
\end{aligned}$$

By Lemma 3.2 $\|\Omega_n^{1/2} \hat{\Omega}_n^{-1/2}\|_{op} \lesssim_p 1$. Now for the final term on the RHS above, first note that:

$$\begin{aligned}
& \|\hat{\Omega}_n^{1/2} \hat{\beta}_n \iota(\hat{\theta} - \theta_n) \chi_{n, i}\|_2^2 \\
& = \frac{1}{n} \sum_{i=1}^n \|\psi'_{n, i} \hat{\beta}_n \iota(\hat{\theta} - \theta_n) \chi_{n, i}\|_2^2 \\
& = \frac{1}{n} \sum_{i=1}^n \|\hat{\pi}'_{n, i}(\hat{\theta} - \theta_n)\|_2^2 \\
& = \|\hat{\Sigma}_n^{1/2}(\hat{\theta} - \theta_n)\|_2^2
\end{aligned}$$

Where $\hat{\Sigma}_n = \frac{1}{n} \sum_{i=1}^n \hat{\pi}_{n, i} \hat{\pi}'_{n, i}$. Next we'll decompose $\hat{\theta} - \theta_n$. Recall $\hat{\theta} = \hat{\Sigma}_{\lambda_{0, n}}^{-1} \frac{1}{n} \sum_{i=1}^n \hat{\pi}_{n, i} \hat{g}_i$ and so we have:

$$\begin{aligned}\hat{\theta} - \theta_n &= \hat{\Sigma}_{\lambda_{0,n}}^{-1} \frac{1}{n} \sum_{i=1}^n \hat{\pi}_{n,i} (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n) \\ &\quad - \lambda_{0,n} \hat{\Sigma}_{\lambda_{0,n}}^{-1} \theta_n\end{aligned}$$

Note that:

$$\|\hat{\Sigma}_{\lambda_{0,n}}^{1/2} (\hat{\theta} - \theta_n)\|_2^2 - \|\hat{\Sigma}_n^{1/2} (\hat{\theta} - \theta_n)\|_2^2 = \lambda_{0,n} \|\hat{\theta} - \theta_n\|_2^2$$

The RHS is positive and so:

$$\|\hat{\Sigma}_n^{1/2} (\hat{\theta} - \theta_n)\|_2 \leq \|\hat{\Sigma}_{\lambda_{0,n}}^{1/2} (\hat{\theta} - \theta_n)\|_2$$

By the triangle inequality:

$$\begin{aligned}\|\hat{\Sigma}_{\lambda_{0,n}}^{1/2} (\hat{\theta} - \theta_n)\|_2 &\leq \|\hat{\Sigma}_{\lambda_{0,n}}^{-1/2} \frac{1}{n} \sum_{i=1}^n \hat{\pi}_{n,i} (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)\|_2 \\ &\quad + \lambda_{0,n} \|\hat{\Sigma}_{\lambda_{0,n}}^{-1/2} \theta_n\|_2\end{aligned}$$

Let's bound the final term above. First note that:

$$\begin{aligned}\|\hat{\Sigma}_{\lambda_{0,n}}^{-1/2}\|_{op} &= \|(\hat{\Sigma}_n + \lambda_{0,n} I)^{-1}\|_{op}^{1/2} \\ &\leq \lambda_{0,n}^{-1/2}\end{aligned}$$

And so:

$$\lambda_{0,n} \|\hat{\Sigma}_{\lambda_{0,n}}^{-1/2} \theta_n\|_2 \leq \lambda_{0,n}^{1/2} \|\hat{\Sigma}_{\lambda_{0,n}}^{-1/2}\|_{op} \|\theta_n\|_2$$

Next note that (by definition of the operator norm):

$$\begin{aligned}&\|\hat{\Sigma}_{\lambda_{0,n}}^{-1/2} \frac{1}{n} \sum_{i=1}^n \hat{\pi}_{n,i} (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)\|_2 \\ &\leq \|\hat{\Sigma}_{\lambda_{0,n}}^{-1/2} \hat{\Sigma}_n^{1/2}\|_{op} \left\| \frac{1}{n} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \hat{\pi}_{n,i} (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n) \right\|_2\end{aligned}$$

And that:

$$\begin{aligned}\|\hat{\Sigma}_{\lambda_{0,n}}^{-1/2} \hat{\Sigma}_n^{1/2}\|_{op} &= \|\hat{\Sigma}_n^{1/2} \hat{\Sigma}_{\lambda_{0,n}}^{-1} \hat{\Sigma}_n^{1/2}\|_{op}^{1/2} \\ &= \|(I + \lambda_{0,n} \hat{\Sigma}_n^{-1})^{-1}\|_{op}^{1/2} \\ &\leq 1\end{aligned}$$

Note that $\|\frac{1}{n} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \hat{\pi}_{n,i} (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)\|_2$ is the square root of the sum of squares residuals from least squares projection of $(\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)$ onto $\hat{\pi}_{n,i}$ and so:

$$\|\frac{1}{n} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \hat{\pi}_{n,i} (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)\|_2 \leq \left(\frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)^2\right)^{1/2}$$

Combining we get:

$$\|\hat{\Sigma}_n^{1/2}(\hat{\theta} - \theta_n)\|_2 \leq \lambda_{0,n}^{1/2} \|\theta_n\|_2 + \left(\frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)^2\right)^{1/2}$$

From Lemma 3.3 we get:

$$\begin{aligned} \|\theta_n\|_2 &\leq \underline{\mu}_n^{-1/2} E[|\phi_n(X, V)' \theta_n|^2]^{1/2} \\ &\lesssim \underline{\mu}_n^{-1/2} \end{aligned}$$

And so:

$$\|\hat{\Sigma}_n^{1/2}(\hat{\theta} - \theta_n)\|_2 \lesssim_p \lambda_{0,n}^{1/2} \underline{\mu}_n^{-1/2} + \left(\frac{1}{n} \sum_{i=1}^n (\hat{g}_i - \hat{\pi}'_{n,i} \theta_n)^2\right)^{1/2}$$

In Step 1 we showed that the second term on the RHS goes to zero in probability faster than the first term, and so:

$$\|\hat{\Sigma}_n^{1/2}(\hat{\theta} - \theta_n)\|_2 \lesssim_p \lambda_{0,n}^{1/2} (\mu_{\min}(Q_n) \mu_{\min}(Q_n))^{-1/2}$$

And so:

$$E_Z [(\hat{\pi}_n(x_1, Z)'(\hat{\theta} - \theta_n))^2 | X = x_1]^{1/2} \lesssim_p \xi_{\Omega,n} \lambda_{0,n}^{1/2} \underline{\mu}_n^{-1/2}$$

And in all:

$$\begin{aligned} &E_Z [|\pi_n(x_1, Z)'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2} \\ &\lesssim_p \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} \xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ &\quad + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} (\xi_{\Omega,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \ell_{\psi,n}(s_4)) \\ &\quad + \xi_{\Omega,n} \lambda_{0,n}^{1/2} \underline{\mu}_n^{-1/2} \end{aligned}$$

□

Lemma 3.6. *Let Assumptions 5.1.ii, 5.2.i-ii, 5.3.iv, and 5.4.ii hold. Uniformly over $\theta \in \mathbb{R}^{k(n)l(n)}$ and F_X -almost all x_1 and x_2 :*

$$\begin{aligned} &(\alpha_n(x_1, x_2) - \hat{\alpha}_n(x_1, x_2))' \theta \\ &\lesssim_p E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{\chi,n} (1 + \sqrt{\xi_{\chi,n}^2 l(n)/n}) \zeta_n \\ &\quad + E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{\chi,n}^2 \min\{\sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n}\} \\ &\quad + \lambda_{3,n} \xi_{\chi,n}^2 E[|\phi_n(X, V)' \theta|^2]^{1/2} \|G_n^{-1}\|_{op} \end{aligned}$$

Where $\zeta_n = 0$ if X has finite discrete support and $\zeta_n = \ell_{\chi,n}(s_5)$ otherwise.

Proof. We follow similar steps to Lemma 3.4. For each n and i define the length- $k(n)$ column vector $\epsilon_{n,i} = \rho_{n,i} - E[\rho_n(V)|X_i]$. By construction:

$$E[\epsilon_{n,i}|X_i] = 0$$

By a similar argument to that in the first part of Lemma 3.4:

$$\begin{aligned} E[\|Q_n^{-1/2}\epsilon_{n,i}\|_2^2] &\leq 2E[\|Q_n^{-1/2}\rho_{n,i}\|_2^2] + 2E[\|E[Q_n^{-1/2}\rho_n(V)|X_i]\|_2^2] \\ &\leq 4E[\|Q_n^{-1/2}\rho_{n,i}\|_2^2] \\ &= 4k(n) \end{aligned} \tag{.35}$$

By a similar argument to Lemma 3.4, with probability 1:

$$\begin{aligned} \|Q_n^{-1/2}\epsilon_{n,i}\|_2 &\leq \|Q_n^{-1/2}\rho_{n,i}\|_2 + \|E[Q_n^{-1/2}\rho_n(V)|X_i]\|_2 \\ &\leq \text{ess sup}\|Q_n^{-1/2}\rho_n(V)\|_2 + \text{ess sup}\|E[Q_n^{-1/2}\rho_n(V)|X]\|_2 \\ &\leq 2\xi_{\rho,n} \end{aligned} \tag{.36}$$

For each n define the $l(n)$ -by- $k(n)$ matrix μ_n by:

$$\mu_n = G_n^{-1}E[\chi_n(X)E[\rho_n(V)|X]']$$

Then define r_n by:

$$r_n(x) = E[\rho_n(V)|X = x] - \mu_n' \psi_n(x)$$

Note that $E[\psi_{n,i}r_n(X_i)'] = 0$. By the Radon-Nikodym Theorem, Assumption 5.2.i implies that for any $\theta \in \mathbb{R}^{k(n)}$:

$$E[\rho_n(V)'\theta|X = x] = E[\rho_n(V)'\theta \frac{dF_{(X,V)}}{dF_X \otimes F_V}(x, V)]$$

In the case of continuous X , by Assumption 5.3.iii, $\frac{dF_{(X,V)}}{dF_X \otimes F_V}(x, V) \in \Lambda_{s_5}^{\dim(X)}(c_5)$, so following steps analogous to those in Lemma 3.4 we get:

$$x \mapsto E[\rho_n(V)'\theta|X = x] \in \Lambda_{s_5}^{\dim(X)}(c_5\|Q_n^{1/2}\theta\|_2)$$

In the continuous- X case, using Assumption 5.1.ii, the above implies that uniformly over $\theta \in \mathbb{R}^{k(n)}$ and F_X -almost all x :

$$\frac{r_n(x)'\theta}{\|Q_n^{1/2}\theta\|_2} \lesssim \ell_{\chi,n}(s_5)$$

Which in turn implies:

$$\|r_n(x)Q_n^{-1/2}\|_2 \lesssim \ell_{\chi,n}(s_5)$$

In the case of X with finite discrete support Assumption 5.1.ii implies that for sufficiently large n , $\|r_n(x)Q_n^{-1/2}\|_2 = 0$. So let ζ_n equal $\ell_{\chi,n}(s_5)$ in the case of X continuous and zero otherwise (we take $a_n \lesssim 0$ to mean a_n equals zero for sufficiently large n). Thus we have:

$$\|r_n(x)Q_n^{-1/2}\|_2 \lesssim \zeta_n \quad (.37)$$

Recall that for any $\theta \in \mathbb{R}^{k(n)l(n)}$, $\phi_n(x, v)' \theta = \rho_n(v)' \iota(\theta) \chi_n(x)$, where ι is defined as in Lemma 3.3. Now decompose:

$$\alpha_n(x_1, x_2)' \theta = \chi_n(x_2)' \mu_n \iota(\theta) \chi(x_1) + r_n(x_2)' \iota(\theta) \chi(x_1)$$

Substituting the above and using the formulas for α_n and $\hat{\alpha}_n$ we get:

$$\begin{aligned} & (\alpha_n(x_1, x_2) - \hat{\alpha}_n(x_1, x_2))' \theta \\ &= r_n(x_2)' \iota(\theta) \chi_n(x_1) \\ & - \chi_n(x_2)' \hat{G}_{\lambda_{3,n}}^{-1} \frac{1}{n} \sum_{i=1}^n \varphi_{n,i} (\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x_1) \\ & - \chi_n(x_2)' \hat{G}_{\lambda_{3,n}}^{-1} \lambda_{3,n} \mu_n \iota(\theta) \chi_n(x_1) \end{aligned}$$

Where $r_{n,i} = r_n(X_i)$. By the triangle inequality, Cauchy-Schwartz, and the definition of the operator norm, the above gives:

$$\begin{aligned} & |(\alpha_n(x_1, x_2) - \hat{\alpha}_n(x_1, x_2))' \theta| \\ & \leq |r_n(x_2)' \iota(\theta) \chi_n(x_1)| \\ & + \left(\|G_n^{-1/2} \chi_n(x_2)\|_2 \|G_n^{1/2} \hat{G}_{\lambda_{3,n}}^{-1} G_n^{1/2}\|_{op} \right. \\ & \quad \left. \times \left\| \frac{1}{n} \sum_{i=1}^n G_n^{-1/2} \varphi_{n,i} (\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x_1) \right\|_2 \right) \\ & + |\chi_n(x_2)' \hat{G}_{\lambda_{3,n}}^{-1} \lambda_{3,n} \mu_n \iota(\theta) \chi_n(x_1)| \end{aligned} \quad (.38)$$

Lemma 3.3 implies $\|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2^2 \lesssim \xi_{\chi,n}^2 E[|\phi_n(X, V)' \theta|^2]$ uniformly. Combining with (.37) and applying Cauchy-Schwartz:

$$\begin{aligned} |r_n(x_2)' \iota(\theta) \chi_n(x_1)| & \leq \|r_n(x_2)Q_n^{-1/2}\|_2 \|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2 \\ & \lesssim E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{\chi,n} \zeta_n \end{aligned} \quad (.39)$$

Uniformly over $\theta \in \mathbb{R}^{k(n)l(n)}$ and F_X -almost all x . From (.39), for the first term we immediately get:

$$E[|r_n(x, Z)' \iota(\theta) \chi_n(x)|^2 | X = x]^{1/2} \leq E[|\phi_n(X, V)' \theta|^2]^{1/2} O(\xi_{\chi,n} \zeta_n)$$

In the discrete finite X case we had $\|r_n(x)Q_n^{-1/2}\|_2 = 0$ for large enough n and so the above term also equals zero for n sufficiently large. Next note that:

$$\begin{aligned}
& \left\| \frac{1}{n} \sum_{i=1}^n G_n^{-1/2} \varphi_{n,i} (\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x_1) \right\|_2 \\
& \leq \|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2 \left(\sum_{l=1}^{m(n)} \left\| \frac{1}{n} \sum_{i=1}^n [G_n^{-1/2} \varphi_{n,i}]_l (\epsilon_{n,i} + r_{n,i})' Q_n^{-1/2} \right\|_2^2 \right)^{1/2}
\end{aligned}$$

Where $[\varphi_{n,i}]_l$ is the l^{th} component of the vector $\varphi_{n,i}$. Note that:

$$\begin{aligned}
& E \sum_{l=1}^{m(n)} \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n [G_n^{-1/2} \varphi_{n,i}]_l (\epsilon_{n,i} + r_{n,i})' Q_n^{-1/2} \right\|_2^2 \\
& = E \left[\|G_n^{-1/2} \varphi_{n,i}\|_2^2 \|(\epsilon_{n,i} + r_{n,i}) Q_n^{-1/2}\|_2^2 \right] \\
& \leq 2E \left[\|G_n^{-1/2} \varphi_{n,i}\|_2^2 \|\epsilon_{n,i} Q_n^{-1/2}\|_2^2 \right] \\
& \quad + 2E \left[\|G_n^{-1/2} \varphi_{n,i}\|_2^2 \|Q_n^{-1/2} r_{n,i}\|_2^2 \right] \\
& \leq 2 \min \left\{ \xi_{\chi,n}^2 E \left[\|\epsilon_{n,i} Q_n^{-1/2}\|_2^2 \right], \xi_{\rho,n}^2 E \left[\|G_n^{-1/2} \varphi_{n,i}\|_2^2 \right] \right\} \\
& \quad + 2 \text{ess sup} \|Q_n^{-1/2} r_n(X)\|_2^2 E \left[\|G_n^{-1/2} \varphi_{n,i}\|_2^2 \right] \\
& \lesssim \min \{ \xi_{\chi,n}^2 k(n), \xi_{\rho,n}^2 l(n) \} + l(n) \zeta_n^2
\end{aligned}$$

Where the first equality follows because $E[\varphi_{n,i} (\epsilon_{n,i} + r_{n,i})'] = 0$ for all n and the data are iid. The first inequality follows by Young's inequality, the next by Holder inequality and then the rate by (.35), (.36), and (.37). By Markov's inequality, uniformly over $\theta \in \mathbb{R}^{k(n)l(n)}$ and F_X -almost all x :

$$\begin{aligned}
& \left\| \frac{1}{n} \sum_{i=1}^n G_n^{-1/2} \varphi_{n,i} (\epsilon_{n,i} + r_{n,i})' \iota(\theta) \chi_n(x_1) \right\|_2 \tag{.40} \\
& \lesssim_p \|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2 \min \{ \sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n} \} \\
& \quad + \|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2 \zeta_n \sqrt{l(n)/n}
\end{aligned}$$

Next let us consider the term $|\chi_n(x_2)' \hat{G}_{\lambda_{3,n}}^{-1} \lambda_{3,n} \mu_n \iota(\theta) \chi_n(x_1)|$. Note that:

$$\begin{aligned}
& |\chi_n(x_2)' \hat{G}_{\lambda_{3,n}}^{-1} \lambda_{3,n} \mu_n \iota(\theta) \chi_n(x_1)| \\
& \leq \lambda_{3,n} \xi_{\chi,n} \|G_n^{1/2} \hat{G}_{\lambda_{3,n}}^{-1} G_n^{1/2}\|_{op} \|G_n^{-1}\|_{op} \|G_n^{1/2} \mu_n \iota(\theta) \chi_n(x)\|_2
\end{aligned}$$

By Lemma 3.2 $\|G_n^{1/2} \hat{G}_{\lambda_{3,n}}^{-1} G_n^{1/2}\|_{op} \lesssim_p 1$. Furthermore:

$$\begin{aligned}
\|G_n^{1/2} \mu_n \iota(\theta) \chi_n(x)\|_2^2 &= E[|\chi_n(X)' \mu_n \iota(\theta) \chi_n(x)|^2] \\
&\leq E[E[\rho_n(V)|X]' \iota(\theta) \chi_n(x)]^2 \\
&\leq E[|\rho_n(V)' \iota(\theta) \chi_n(x)|^2] \\
&= \|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2^2
\end{aligned} \tag{.41}$$

Where the first inequality holds by the properties of least-squares projection, the second inequality by positivity of the variance. Thus we have uniformly over $\theta \in \mathbb{R}^{k(n)}$ and F_X -almost all x :

$$|\chi_n(x_2)' \hat{G}_{\lambda_{3,n}}^{-1} \lambda_{3,n} \mu_n \iota(\theta) \chi_n(x_1)| \lesssim_p \lambda_{3,n} \|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2 \|G_n^{-1}\|_{op} \xi_{\chi,n}$$

By Lemma 3.3, $\|Q_n^{1/2} \iota(\theta) \chi_n(x)\|_2 \lesssim E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{\chi,n}$, and finally note that $\|G_n^{-1/2} \chi_n(x_2)\|_2 \leq \xi_{\chi,n}$. Substituting these and also (.39), (.40), and (.41) into (.38) we get that uniformly over $\theta \in \mathbb{R}^{k(n)l(n)}$ and F_X -almost all x :

$$\begin{aligned}
&(\alpha_n(x_1, x_2) - \hat{\alpha}_n(x_1, x_2))' \theta \\
&\lesssim_p E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{\chi,n} (1 + \sqrt{\xi_{\chi,n}^2 l(n)/n}) \zeta_n \\
&\quad + E[|\phi_n(X, V)' \theta|^2]^{1/2} \xi_{\chi,n}^2 \min\{\sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n}\} \\
&\quad + \lambda_{3,n} \xi_{\chi,n} E[|\phi_n(X, V)' \theta|^2]^{1/2} \|G_n^{-1}\|_{op} \xi_{\chi,n}
\end{aligned}$$

□

Proposition 3.3. *Let Assumptions 5.1.ii, 5.2.i-ii, 5.3.iv, and 5.4.ii hold. Suppose $E[|\phi_n(X, V)' \theta_n|^2]$ is bounded above uniformly over n , and uniformly over $F_{(X,Z)}$ -almost all (x, z) :*

$$|g(x, z) - \pi_n(x, z)' \theta_n| \lesssim \kappa_n$$

Suppose in addition that $\lambda_{0,n}$ goes to zero sufficiently slowly, precisely:

$$\begin{aligned}
&\lambda_{0,n}^{1/2} \underline{\mu}_n^{-1/2} \\
&\lesssim_p (1 + \ell_{\psi,n}(s_3)) \sqrt{m(n)/n} + \ell_{\psi,n}(s_3) + \lambda_{1,n} \|\Omega_n^{-1}\|_{op} \\
&\quad + \xi_{\chi,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\
&\quad + \xi_{\chi,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \xi_{\chi,n} \ell_{\psi,n}(s_4) \\
&\quad + \kappa_n
\end{aligned}$$

Then uniformly over $\theta \in \mathbb{R}^{k(n)l(n)}$ and F_X -almost all x_1 and x_2 :

$$\begin{aligned} & (\alpha_n(x_1, x_2) - \hat{\alpha}_n(x_1, x_2))' \theta \\ & \lesssim_p \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 (1 + \sqrt{\xi_{\chi,n}^2 l(n)/n}) \zeta_n \\ & \quad + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 \min\{\sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n}\} \\ & \quad + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 \lambda_{3,n} \|\Omega_n^{-1}\|_{op} \end{aligned}$$

Where $\bar{\mu}_n = \sqrt{\mu_{\max}(Q_n)\mu_{\max}(G_n)}$, $\underline{\mu}_n = \sqrt{\mu_{\min}(Q_n)\mu_{\min}(G_n)}$, and $\zeta_n = 0$ if X has finite discrete support and otherwise $\zeta_n = \ell_{\chi,n}(s_5)$.

Proof. Under the conditions of this proposition, in Step 1 of Proposition 3.2 we show:

$$E_{(X,V)} [|\phi_n(X, V)'(\hat{\theta} - \theta_n)|^2]^{1/2} \lesssim_p \sqrt{\frac{\mu_{\max}(Q_n)\mu_{\max}(G_n)}{\mu_{\min}(Q_n)\mu_{\min}(G_n)}}$$

Combing with Lemma 3.6 immediately gives the result. \square

Proposition 3.4. *Let Assumptions 1, 2, 3, and 5 hold and Assumption 4 with $D(x_1)$ and $C(x_1, x_2)$ bounded uniformly over F_X -almost all x_1 and x_2 . Suppose that $\lambda_{0,n}$ goes to zero sufficiently slowly that:*

$$\begin{aligned} & \lambda_{0,n}^{1/2} \underline{\mu}_n^{-1/2} \\ & \lesssim_p (1 + \ell_{\psi,n}(s_3)) \sqrt{m(n)/n} + \ell_{\psi,n}(s_3) + \lambda_{1,n} \|\Omega_n^{-1}\|_{op} \\ & \quad + \xi_{\chi,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ & \quad + \xi_{\chi,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \xi_{\chi,n} \ell_{\psi,n}(s_4) \\ & \quad + \kappa_n \end{aligned}$$

where $\kappa_n = \ell_{\rho,n}(s_1)$ if X has finite discrete support, and otherwise $\kappa_n = \ell_{\rho,n}(s_1) + (\xi_{\rho,n} \ell_{\chi,n}(1))^{\tilde{s}}$, where $\tilde{s} = \frac{\min\{s_2, s_3, 1\}}{\min\{s_2, s_3, 1\} + 1}$. Let $\zeta_n = 0$ if X has finite discrete support and otherwise let $\zeta_n = \ell_{\chi,n}(s_5)$. Then uniformly over F_X -almost all x_1 and x_2 :

$$\begin{aligned}
& |\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| \\
& \lesssim_p \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 (1 + \sqrt{\xi_{\chi,n}^2 l(n)/n}) \zeta_n \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 \min\{\sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n}\} \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 \lambda_{3,n} \|G_n^{-1}\|_{op} \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} \xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} (\xi_{\Omega,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \ell_{\psi,n}(s_4)) \\
& + \xi_{\Omega,n} \lambda_{0,n}^{1/2} \underline{\mu}_n^{-1/2} \\
& + \kappa_n
\end{aligned}$$

Proof. By Proposition 3.1 there is a sequence $\{\theta_n\}_{n=1}^\infty$ with $E[|\phi_n(X, V)' \theta_n|^2]$ bounded above uniformly over n so that, uniformly over $F_{(X,Z)}$ -almost all (x, z) :

$$|g(x, z) - \pi_n(x, z)' \theta_n| \lesssim \kappa_n$$

where $\kappa_n = \ell_{\rho,n}(s_1)$ if X has finite discrete support, and otherwise $\kappa_n = \ell_{\rho,n}(s_1) + (\xi_{\rho,n} \ell_{\chi,n}(1))^{\bar{s}}$. Then by Propositions 3.2 and 3.3, if $\lambda_{0,n}$ goes to zero sufficiently slowly that:

$$\begin{aligned}
& \lambda_{0,n}^{1/2} \underline{\mu}_n^{-1/2} \\
& \lesssim_p (1 + \ell_{\psi,n}(s_3)) \sqrt{m(n)/n} + \ell_{\psi,n}(s_3) + \lambda_{1,n} \|\Omega_n^{-1}\|_{op} \\
& + \xi_{\chi,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\
& + \xi_{\chi,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \xi_{\chi,n} \ell_{\psi,n}(s_4) \\
& + \kappa_n
\end{aligned}$$

Then we have:

$$\begin{aligned}
& (\alpha_n(x_1, x_2) - \hat{\alpha}_n(x_1, x_2))' \theta \\
& \lesssim_p \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 (1 + \sqrt{\xi_{\chi,n}^2 l(n)/n}) \zeta_n \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 \min\{\sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n}\} \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 \lambda_{3,n} \|G_n^{-1}\|_{op}
\end{aligned}$$

And:

$$\begin{aligned}
& E_Z [|\pi_n(x_1, Z)'(\hat{\theta} - \theta_n)|^2 | X = x_1]^{1/2} \\
& \lesssim_p \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} \xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} (\xi_{\Omega,n} \lambda_{2,n} \|\Omega_n^{-1}\|_{op} + \ell_{\psi,n}(s_4)) \\
& + \xi_{\Omega,n} \lambda_{0,n}^{1/2} \underline{\mu}_n^{-1/2}
\end{aligned}$$

Substituting into Lemma 3.1 then gives the result. \square

Proof of Theorem 3.1. Recall the conclusion of Proposition 3.4. Suppose we set $\lambda_{0,n}$ so that the constraint on its rate in Theorem 3.1 binds and substitute this into the rate in Proposition 3.4. If we then set $\lambda_{1,n}$, $\lambda_{2,n}$, and $\lambda_{3,n}$ low enough that they disappear from the resulting expression we get (also removing redundant terms due to $\ell_{\psi,n}(s_3) < 1$ and $l(n) \gtrsim 1$ which implies $\xi_{\Omega,n}, \xi_{\chi,n} \gtrsim 1$):

$$\begin{aligned}
& |\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| \\
& \lesssim_p \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 \zeta_n \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n}^2 \min\{\sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n}\} \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} \xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\
& + \frac{\bar{\mu}_n}{\underline{\mu}_n} \xi_{\chi,n} \ell_{\psi,n}(s_4) + \xi_{\Omega,n} (\ell_{\psi,n}(s_3) + \kappa_n)
\end{aligned}$$

Where ζ_n is defined as in the statement of Proposition 3.3. Because s_1, s_2, \dots, s_5 are all equal to s and the bounds on the smallest and largest eigenvalues of Q_n and G_n implies $\frac{\bar{\mu}_n}{\underline{\mu}_n} \gtrsim 1$ we get:

$$\begin{aligned}
& |\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| \\
& \lesssim_p \xi_{\chi,n}^2 \zeta_n \\
& + \xi_{\chi,n}^2 \min\{\sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n}\} \\
& + \xi_{\chi,n} \xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\
& + \xi_{\chi,n} \ell_{\psi,n}(s) + \xi_{\Omega,n} (\ell_{\psi,n}(s) + \kappa_n)
\end{aligned}$$

If X is finite and discrete then it is easy to see that $l(n) = l(n) \gtrsim 1$, $\xi_{\Omega,n} \gtrsim 1$, $\xi_{\chi,n} \gtrsim 1$, $\xi_{\chi,n} \gtrsim 1$ and $\zeta_n = 0$ for n sufficiently large. Using these rates and

removing redundant terms we get that in the case of X finite and discrete:

$$\begin{aligned} & |\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| \\ & \lesssim_p \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ & \quad + \ell_{\psi,n}(s) + \ell_{\rho,n}(s) \end{aligned}$$

In the continuous case, using $\kappa_n = \ell_{\rho,n}(s) + (\xi_{\rho,n} \ell_{\chi,n}(1))^{\tilde{s}}$, and $\zeta_n = \ell_{\chi,n}(s)$ we get:

$$\begin{aligned} & |\bar{y}(x_1|x_2) - \hat{\alpha}_n(x_1, x_2)' \hat{\theta}| \\ & \lesssim_p \xi_{\chi,n}^2 \min\{\sqrt{\xi_{\chi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 l(n)/n}\} \\ & \quad + \xi_{\chi,n} \xi_{\Omega,n} \min\{\sqrt{\xi_{\psi,n}^2 k(n)/n}, \sqrt{\xi_{\rho,n}^2 m(n)/n}\} \\ & \quad + \xi_{\chi,n}^2 \ell_{\chi,n}(s) + \xi_{\Omega,n} \ell_{\psi,n}(s) + \xi_{\Omega,n} \ell_{\rho,n}(s) + \xi_{\Omega,n} (\xi_{\rho,n} \ell_{\chi,n}(1))^{\tilde{s}} \end{aligned}$$

Finally note that $s_2 = s_3 = s$ implies $\tilde{s} = \frac{\min\{s_2, s_3, 1\}}{\min\{s_2, s_3, 1\} + 1} = \min\{\frac{1}{2}, \frac{s}{s+1}\}$ \square