

# Simple Inference on Functionals of Set-Identified Parameters Defined by Linear Moments

(Preliminary and incomplete. All comments are welcome.)

JoonHwan Cho\*

Thomas M. Russell†

*University of Toronto*

*University of Toronto*

December 28, 2023

## Abstract

This paper considers uniformly valid (over a class of data generating processes) inference for linear functionals of partially identified parameters in cases where the identified set is defined by linear (in the parameter) moment inequalities. We propose a bootstrap procedure for constructing uniformly valid confidence sets for a linear functional of a partially identified parameter. The proposed method amounts to bootstrapping the value functions of a linear optimization problem, and subsumes subvector inference as a special case. In other words, this paper shows the conditions under which naively bootstrapping a linear program can be used to construct a confidence set with uniform correct coverage for a partially identified linear functional. Unlike other proposed subvector inference procedures, our procedure does not require the researcher to repeatedly invert a hypothesis test, and is extremely computationally efficient.

*Keywords:* Stochastic Programming, Subvector Inference, Partial Identification, Linear Programming

## 1 Introduction

This paper proposes a uniformly valid (over a large class of data generating processes) inference procedure for a linear functional  $\psi$  of a partially identified parameter vector  $\theta$  in models with linear (in  $\theta$ ) moment functions. In particular, the paper proposes a bootstrap procedure to approximate the distribution of the endpoints of the projected identified set, and discusses primitive conditions under which the procedure is uniformly valid.

---

A previous version of this paper was circulated under the title “Inference on Functionals of Set-Identified Parameters Defined by Convex Moments.” We are grateful to Victor Aguirregabiria, Bulat Gafarov, Christian Gourieroux, Ismael Mourifie, Jeffrey Negrea, Brennan Thompson, Stanislav Volgushev and Yuanyuan Wan for helpful comments and discussion. We are also grateful to participants at the 7th Annual Doctoral Workshop in Applied Econometrics at the University of Toronto, as well as participants at the 2019 North America Summer Meeting of the Econometric Society at the University of Washington. This research was supported by the Social Sciences and Humanities Research Council of Canada. All errors are our own.

\*JoonHwan Cho, Department of Economics, University of Toronto, 150 St. George Street, Toronto, Ontario, M5S3G7, Canada.

†Thomas M. Russell, Department of Economics, University of Toronto, 150 St. George Street, Toronto, Ontario, M5S3G7, Canada. Email: thomas.russell@mail.utoronto.ca.



The main idea is to use results from the Operations Research literature that allow the researcher to approximate the distribution of the value functions in linear programs with stochastic constraints using a functional delta method. The contribution of this paper is to use these results for stochastic programs as a proof device to show the uniform validity of a simple bootstrap procedure for constructing confidence sets for subvectors or functionals of the identified set in partially identified econometric models. Intuitively, bounding a linear functional over an identified set defined by linear moment functions amounts to solving two linear optimization problems: one maximization problem for the upper bound, and one minimization problem for the lower bound. Thus, the endpoints of the identified set for a functional of a partially-identified parameter can be viewed as *value functions* of two “stochastic programs.” An inference procedure is then constructed by decomposing perturbations in the optimal value function into perturbations arising from the objective function and perturbations arising from the constraint functions. By the envelope theorem, perturbations in the constraints are related to the value functions through the Lagrange multipliers. The total effect of perturbations in the constraints on the value function is then given by a weighted sum of the perturbations in all binding constraints, where the weights are determined by the Lagrange multipliers. Through this mechanism, we can relate the distribution of the binding moment functions to the distribution of the value function of a stochastic program.

To prove the validity of our procedure requires noticing that, under some conditions, the value functions in linear stochastic programs are *Hadamard directionally differentiable* with respect to perturbations in the underlying probability measure. However, this form of differentiability is not sufficient for *uniformly* valid confidence sets. This result relates to [Kasy \(2015\)](#), who emphasizes that failures of uniformity often result as failures of the uniform versions of the delta method. We demonstrate the conditions under which the value functions of a linear stochastic program satisfy the natural definition of *uniform* Hadamard directional differentiability with respect to perturbations in the underlying probability measure. The condition that emerges as being most important for our procedure is the existence and uniqueness of optimal solutions and Lagrange multipliers. For now, we impose high-level assumptions to ensure this holds, but plan to elaborate on more primitive conditions in future drafts.

Uniform Hadamard directional differentiability is sufficient for us to prove the validity of a simple uniformly valid bootstrap procedure to estimate the confidence set for a functional of interest. In the environment considered in this paper, bounds on the linear functional of interest can always be constructed by solving linear programs, and our bootstrap procedure amounts to repeatedly solving analogous “bootstrap linear program.” In this way, bootstrap distributions for the value functions of these linear programs are constructed, and then a confidence set for the partially identified functional of interest is constructed by selecting appropriate quantiles from these bootstrap distributions. In other words, this paper shows the conditions under which naively bootstrapping a linear program can be used to construct a confidence set with uniform correct coverage for a partially identified linear functional.

Interestingly, the use of Lagrange multipliers allows us to avoid rescaling the moment conditions by their



sample standard deviations in our procedure. This is in contrast to other comparable methods. Intuitively, any rescaling of the moment functions is countered by an equivalent (but opposite) rescaling of the Lagrange multipliers. This ensures our bootstrap procedure remains a sequence of linear—and thus easy to solve—optimization problems.

Subvector inference, or inference on functionals of the identified set, has recently been a topic of considerable interest in the partial identification literature. The earlier papers of [Andrews and Guggenberger \(2009\)](#) and [Andrews and Soares \(2010\)](#) propose to project confidence sets constructed for the entire parameter vector in order to obtain confidence sets for a particular subvector of interest. While these procedures are uniformly valid, they can be highly conservative when the dimension of the partially identified parameter vector is large (see the discussion in [Kaido et al. \(2017\)](#)). Both [Romano and Shaikh \(2008\)](#) and [Bugni et al. \(2017\)](#) consider inverting profiled test statistics in order to construct confidence sets for subvectors or functionals, where [Romano and Shaikh \(2008\)](#) construct critical values using subsampling and where [Bugni et al. \(2017\)](#) derive the asymptotic distribution for their profile test statistic for a large class of test functions. [Bugni et al. \(2017\)](#) show that their test dominates projection-based procedures in terms of asymptotic power, and they derive conditions under which it dominates the subsampling-based approach of [Romano and Shaikh \(2008\)](#). [Kaido et al. \(2017\)](#) provide a “calibrated projection” inference method for functionals of a partially identified parameter. Intuitively, this procedure suitably relaxes the model’s moment inequalities, and then solves two optimization problems subject to the relaxed constraints in order to obtain the endpoints of the confidence interval for the functional of interest. The relaxation of the constraints requires the correct calibration of a relaxation parameter in order to obtain uniformly correct coverage. [Kaido et al. \(2017\)](#) first linearize any nonlinear moment functions, and then propose an efficient algorithm to calibrate the relaxation parameter. This allows their procedure to be computationally attractive relative to other methods in nonlinear models. Similar to the method proposed here, the method of [Kaido et al. \(2017\)](#) does not invert a test statistic.

The overall approach to constructing confidence sets in this paper is most closely related to the approach in [Gafarov \(2018\)](#), who shows how to construct both pointwise and uniformly valid confidence sets for linear functionals of a partially identified parameter in an optimization framework. However, the optimization problems used to perform functional inference may have non-unique solutions, and it is well known from [Hirano and Porter \(2012\)](#) that it is impossible to obtain a locally unbiased estimator of the value function when the value function is nondifferentiable. To address these problems, [Gafarov \(2018\)](#) proposes including a regularization term in the objective function to ensure a unique optimal solution is selected. In contrast we impose a high level assumption on the class of DGPs and moment conditions that guarantees a unique solution, and avoids arbitrarily selecting a unique solution from a set of candidate solutions. Similar to [Gafarov \(2018\)](#), we also impose a linear independence constraint qualification to ensure uniqueness of the Lagrange multipliers. However, both our bootstrap procedure and our proof of uniform validity are very different. As mentioned above, we plan to provide more primitive conditions for uniqueness of solutions and Lagrange multipliers in future drafts, which should help to further differentiate our work from [Gafarov](#)



(2018).

The main proofs of this paper uses results from [Shapiro et al. \(2009\)](#) and [Fang and Santos \(2018\)](#). The main result used from [Shapiro et al. \(2009\)](#) is the proof of Hadamard directional differentiability of value functions for stochastic programs. However, we extend this result by showing the conditions under which the value functions for a stochastic program satisfy uniform Hadamard directional differentiability, which is sufficient to derive a uniform delta method result. Next, while we use some of the results in [Fang and Santos \(2018\)](#) related to bootstrap consistency and Hadamard directional differentiability, our main focus is on inference for the functional  $\psi$  of the true parameter  $\theta_0$  in partially identified models using stochastic programs, which is not considered in [Fang and Santos \(2018\)](#).

The remainder of the paper proceeds as follows. Section 2 provides a simplified overview of the main results, and examples to motivate the need for functional inference procedures in partially identified models. Section 3 develops the methodology in detail and contains the main results of this paper. Section 4 concludes. The proofs of all of the main results are provided in Appendix A.

Throughout the paper we use notation standard in empirical process theory; in particular, the expectation of a random element  $X_t$  with respect to a measure  $P$  is given by  $PX_t$ . If the random element  $X_t$  is a vector, then the expectation is interpreted element-wise. The random variables  $W_1, W_2, \dots, W_n$  are assumed to be coordinate projections from the product space  $(\mathcal{W}^n, \mathcal{A}^n, P^n)$ , where  $P^n = P \otimes P \otimes \dots \otimes P$ , and we will denote  $(\mathcal{W}^\infty, \mathcal{A}^\infty, P^\infty)$  as the infinite product space. The empirical measure is represented by  $\mathbb{P}_n$ , which is implicitly a function of the generating measure  $P_n$  at sample size  $n$ . We index estimated quantities by the empirical distribution; for example, rather than  $\hat{\theta}$ , we write  $\theta(\mathbb{P}_n)$ . This is done to emphasize the underlying measure relevant to the construction of the parameter, and becomes useful in both the discussion and the proofs of the main results. Finally, we use  $\|\cdot\|$  to denote the euclidean norm throughout. For the most part, we will avoid issues of measurability as much as possible, although all the proofs follow from the definition of weak convergence in the sense of [Hoffmann-Jørgensen \(1991\)](#), where the outer measure is denoted with a superscript  $*$  (c.f. [Van Der Vaart and Wellner \(1996\)](#) Chapters 1.1 and 1.2).

## 2 Overview of Results and Motivating Examples

### 2.1 Main Ideas

This subsection will discuss simplified versions of the main ideas in the paper before the technical details are introduced in the next section. Our main motivation is to construct uniformly valid confidence sets for the expectation of the random objective function  $\psi(W, \theta)$ , where  $W \in \mathcal{W}$  denotes the relevant finite-dimensional vector of random variables in the model, and where  $\theta$  is only partially identified, and constrained to lie in the identified set.

To this end, we suppose the identified set for  $\theta \in \Theta$  is defined by  $k$  moment (in)equalities where the moment function  $m_j(W, \theta) : \Theta \rightarrow \mathbb{R}$  is linear in  $\theta \in \Theta$  for  $j = 1, \dots, k$ . In this case, the identified set  $\Theta_I(P)$ —



indexed here by the true asymptotic distribution  $P$ —is compact and linear, and so the image of  $\Theta_I(P)$  under the expectation of any continuous functional  $P\psi(W, \theta) : \Theta \rightarrow \mathbb{R}$  will be an interval  $\Psi_I(P) = [\Psi_I^{\ell b}(P), \Psi_I^{ub}(P)]$ . In this framework, the endpoints of the interval  $\Psi_I(P)$  can be determined by solving two linear optimization problems:

- (i) minimize  $P\psi(W, \theta)$  over  $\theta \in \Theta_I(P)$  to determine  $\Psi_I^{\ell b}(P)$ ,
- (ii) maximize  $P\psi(W, \theta)$  over  $\theta \in \Theta_I(P)$  to determine  $\Psi_I^{ub}(P)$ .

Seen in this way,  $\Psi_I^{\ell b}(P)$  and  $\Psi_I^{ub}(P)$  are the *value functions* of two stochastic linear optimization problems. Now let  $\theta_0 \in \Theta$  denote the true value of the parameter, and consider the problem of constructing a confidence set  $C_n^\psi(1 - \alpha)$  that asymptotically covers  $P\psi(W, \theta_0)$  with probability at least  $1 - \alpha$  uniformly over  $(\theta, P) \in \{(\theta, P) : \theta \in \Theta_I(P), P \in \mathcal{P}\}$ , where  $\mathcal{P}$  is some large class of DGPs. In particular, we wish to construct a confidence set  $C_n^\psi(1 - \alpha)$  such that

$$\liminf_{n \rightarrow \infty} \inf_{\{(\psi, P) : \psi \in \Psi_I(P), P \in \mathcal{P}\}} P(\psi \in C_n^\psi(1 - \alpha)) \geq 1 - \alpha.$$

To construct such a set, we will approximate the distribution of the endpoints  $(\Psi_I^{\ell b}(P), \Psi_I^{ub}(P))$  of the identified set  $\Psi_I(P)$ . In particular, let  $\mathcal{F}$  denote the relevant class of functions (we define this class more precisely in Section 3). We show that under a constraint qualification condition, for any sequence  $\{P_n \in \mathcal{P}\}_{n=1}^\infty$  converging to a measure  $P \in \mathcal{P}$  in an appropriate sense (to be made precise), there exist continuous functionals  $(\Psi_I^{\ell b})'_P, (\Psi_I^{ub})'_P : \ell^\infty(\mathcal{F}) \rightarrow \mathbb{R}$  such that

$$\sqrt{n} (\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P_n)) \rightsquigarrow (\Psi_I^{\ell b})'_P(\mathbb{G}_P), \quad (2.1)$$

$$\sqrt{n} (\Psi_I^{ub}(\mathbb{P}_n) - \Psi_I^{ub}(P_n)) \rightsquigarrow (\Psi_I^{ub})'_P(\mathbb{G}_P), \quad (2.2)$$

where  $\mathbb{G}_P \in \ell^\infty(\mathcal{F})$  is the limit of the empirical process  $\mathbb{G}_{n, P_n} := \sqrt{n}(\mathbb{P}_n - P_n) \in \ell^\infty(\mathcal{F})$ , and  $(\Psi_I^{\ell b}(\mathbb{P}_n), \Psi_I^{ub}(\mathbb{P}_n))$  are suitable estimates of the value functions. Moreover, we show conditions under which:

$$\sqrt{n} (\Psi_I^{\ell b}(\mathbb{P}_n^b) - \Psi_I^{\ell b}(\mathbb{P}_n)) | \{W_i\}_{i=1}^n \rightsquigarrow (\Psi_I^{\ell b})'_P(\mathbb{G}_P), \quad (2.3)$$

$$\sqrt{n} (\Psi_I^{ub}(\mathbb{P}_n^b) - \Psi_I^{ub}(\mathbb{P}_n)) | \{W_i\}_{i=1}^n \rightsquigarrow (\Psi_I^{ub})'_P(\mathbb{G}_P), \quad (2.4)$$

uniformly over  $\mathcal{P}$ , where  $\mathbb{P}_n^b$  is the empirical bootstrap distribution. From here, our proposed confidence set takes the form:

$$C_n^\psi(1 - \alpha) := \left[ \Psi_I^{\ell b}(\mathbb{P}_n) - \frac{\hat{\Psi}_\alpha^{\ell b}}{\sqrt{n}}, \Psi_I^{ub}(\mathbb{P}_n) + \frac{\hat{\Psi}_\alpha^{ub}}{\sqrt{n}} \right],$$

where the quantiles  $\hat{\Psi}_\alpha^{\ell b}$  and  $\hat{\Psi}_\alpha^{ub}$  are selected from the bootstrap approximation to the distributions of  $(\Psi_I^{\ell b})'_P(\mathbb{G}_P)$  and  $(\Psi_I^{ub})'_P(\mathbb{G}_P)$  given by (2.3) and (2.4) in order to guarantee uniformly correct coverage.



After presenting some motivating examples, the next sections develop this methodology rigorously.

## 2.2 Examples

We now present some motivating examples that illustrate why inference procedures for functionals of partially identified parameters are needed.

**Example 1** (Missing Data). *Consider the simple missing data example as presented, for example, in [Chen et al. \(2016\)](#). In this example the researcher observes a sample  $\{Y_i D_i, D_i\}_{i=1}^n$ . For simplicity, suppose that  $Y_i, D_i \in \{0, 1\}$ . The parameter of interest is the unconditional average of the outcome variable:*

$$P\psi(W, \theta) = \psi(\theta) = \sum_y \sum_d \theta_{yd} \cdot y,$$

where  $\theta_{yd} := P(Y = y, D = d)$ . The constraints imposed by the observed distribution  $\mathbb{P}_n(YD, D)$  on the latent distribution  $\theta_{yd} = P(Y = y, D = d)$  are given by:

$$\mathbb{P}_n(YD = 0, D = 1) = \theta_{01}, \quad (2.5)$$

$$\mathbb{P}_n(YD = 1, D = 1) = \theta_{11}, \quad (2.6)$$

$$\mathbb{P}_n(YD = 0, D = 0) = \theta_{00} + \theta_{10}. \quad (2.7)$$

It is straightforward to see that point identification of  $\theta$  occurs only when  $\mathbb{P}_n(D = 0) = 0$ . The identified set for our function of interest,  $\Psi_I(\mathbb{P}_n) = [\Psi_I^{\ell b}(\mathbb{P}_n), \Psi_I^{ub}(\mathbb{P}_n)]$  can be obtained by solving the problems:

$$\Psi_I^{\ell b}(\mathbb{P}_n) = \min_{\theta \in \Theta_I(\mathbb{P}_n)} \psi(\theta), \quad \Psi_I^{ub}(\mathbb{P}_n) = \max_{\theta \in \Theta_I(\mathbb{P}_n)} \psi(\theta), \quad (2.8)$$

where  $\Theta_I(\mathbb{P}_n)$  is the set of  $\theta_{yd}$  satisfying the constraints (2.5)-(2.7). Note that the optimization problems in (2.8) are linear programs. This paper will attempt to exploit the structure of the optimization problems in (2.8) to propose an inference procedure that is easy to use for functionals of partially identified parameters. Here, note that  $\psi$  is a functional of the partially identified parameter  $\theta$ , where the identified set for  $\theta$  is given by  $\Theta_I(\mathbb{P}_n)$ .

**Example 2** (Linear Regression with Interval-Valued Dependent Variable). *Consider the example of linear regression with interval-valued dependent variable. We will follow closely the exposition in [Kaido et al. \(2017\)](#) Appendix C. In this example the model is given by  $Y = X^T \theta + \varepsilon$ , where  $X \in \mathbb{R}^d$  with  $R$  points of support. However, it is assumed that the dependent variable is interval-valued in the following way: although the value of  $Y$  is never observed, there exists two observable random variables  $Y^*$  and  $Y_*$  such that  $P(Y_* \leq Y \leq Y^*) = 1$ . The objective is then to construct bounds on the parameter  $\theta$  given that researcher observes a sample  $\{Y_i^*, Y_{*i}, X_i\}_{i=1}^n$ , and never directly observes the value of  $Y$ . Denoting the support points of  $X$  as  $\{x_1, \dots, x_r, \dots, x_R\}$ , as in [Kaido et al. \(2017\)](#) the identified set is given by:*

$$\Theta_I(P) := \{\theta : \mathbb{E}[Y_* | X = x_r] - x_r^T \theta \leq 0, \ x_r^T \theta - \mathbb{E}[Y^* | X = x_r] \leq 0, \ r = 1, \dots, R\}.$$



We now suppose that the researcher is interested in conducting inference only on the first component  $\theta_1$  of the parameter vector  $\theta$ . Then in our notation we can set  $\psi(W, \theta) = \psi(\theta) = \theta_1$ . Under some weak conditions we will have that the identified set for the functional  $\psi$  is an interval  $\Psi_I(\mathbb{P}_n) = [\Psi_I^{\ell b}(\mathbb{P}_n), \Psi_I^{ub}(\mathbb{P}_n)]$  with the endpoints determined by the program:

$$\Psi_I^{\ell b}(\mathbb{P}_n) = \min_{\theta \in \Theta_I(\mathbb{P}_n)} \psi(\theta), \quad \Psi_I^{ub}(\mathbb{P}_n) = \max_{\theta \in \Theta_I(\mathbb{P}_n)} \psi(\theta), \quad (2.9)$$

where  $\Theta_I(\mathbb{P}_n)$  is the estimate of the identified set obtained by replacing the moment conditions with their sample analogs. Note that since all moment conditions defining the identified set are linear in  $\theta$ , the optimization problems in (2.9) are linear programs. Again, this paper will propose an inference procedure for functionals of partially identified parameters that uses the special structure of the optimization problems in (2.9) that characterizes the functional bounding problem.

**Example 3** (Nonparametric State Dependence). Consider the model of nonparametric state dependence given in [Torgovitsky \(2016\)](#). In this model, the researcher observes a realization of a random sequence  $Y := (Y_0, \dots, Y_T)$  for each individual for  $T$  periods. As in [Torgovitsky \(2016\)](#), we consider for simplicity that each outcome  $Y_t$  is binary, so that  $Y \in \{0, 1\}^{T+1}$ . The sequence of observed outcomes  $Y$  are related to a sequence of unobserved potential outcomes  $U(0) := (U_1(0), \dots, U_T(0))$  and  $U(1) := (U_1(1), \dots, U_T(1))$  through the equation:

$$Y_t = Y_{t-1}U_t(0) + (1 - Y_{t-1})U_t(1).$$

The researcher may also have access to a sequence of covariates  $X := (X_0, \dots, X_T)$  for each individual. The object of interest for the researcher is assumed to be treatment effect parameters that depend on the unobserved potential outcomes  $(U_t(0), U_t(1))$  at time  $1 \leq t \leq T$ . Examples of such treatment effect parameters include the average treatment effect, given by  $ATE_t = P(U_t(0) = 0, U_t(1) = 1) - P(U_t(0) = 1, U_t(1) = 0)$ , or the voting criterion given by  $P(U_t(0) = 0, U_t(1) = 1)$  (or  $P(U_t(0) = 1, U_t(1) = 0)$ ).

To see how to bound these parameters, define the vector

$$\mathbf{u} := (u_0, u_1(0), \dots, u_T(0), u_1(1), \dots, u_T(1))',$$

where  $u_0$  is the initial (period 0) potential outcome. In addition, let  $U := (U_0, U(0), U(1))'$ , and let

$$\mathcal{U}^\dagger(\mathbf{y}) := \{\mathbf{u} : u_0 = y_0, y_t = y_{t-1}u_t(0) + (1 - y_{t-1})u_t(1), \forall t\},$$

which is the set of all vectors  $\mathbf{u}$  of potential outcomes that could rationalize an observed vector of outcomes  $\mathbf{y} = (y_0, \dots, y_T)'$ . Finally, let  $\mathbf{X} = (x_0, \dots, x_T)'$ . [Torgovitsky \(2016\)](#) shows that without any additional restrictions, the sharp set of constraints on the unobserved joint distribution  $\theta_{\mathbf{u}, \mathbf{x}} := P(U = \mathbf{u}, X = \mathbf{x})$  is



given by:

$$\mathbb{P}_n(Y = \mathbf{y}, X = \mathbf{x}) = \sum_{\mathbf{u} \in \mathcal{U}^\dagger(\mathbf{y})} \theta_{\mathbf{u}, \mathbf{x}}. \quad (2.10)$$

[Torgovitsky \(2016\)](#) shows how additional restrictions can also be imposed on the unobserved joint distribution  $\theta_{\mathbf{u}, \mathbf{x}}$ , such as monotone treatment response (MTR) constraints, stationarity (ST) constraints, monotone instrumental variable (MIV) constraints and monotone treatment selection (MTS) constraints. All of these constraints can be imposed on the optimization problem as moment-inequality constraints. Let  $\Theta_I(\mathbb{P}_n)$  denote the set of all joint distributions  $\theta$  satisfying the imposed constraints as well as the observational equivalence condition (2.10). Proposition 1 in [Torgovitsky \(2016\)](#) shows that if  $\psi : \Theta_I(\mathbb{P}_n) \rightarrow \mathbb{R}$  is a continuous treatment effect parameter, then the identified set for  $\psi$  can be estimated by  $\Psi_I(\mathbb{P}_n) = [\Psi_I^{\ell b}(\mathbb{P}_n), \Psi_I^{ub}(\mathbb{P}_n)]$ , and can be obtained by solving the problems:

$$\Psi_I^{\ell b}(\mathbb{P}_n) = \min_{\theta \in \Theta_I(\mathbb{P}_n)} \psi(\theta), \quad \Psi_I^{ub}(\mathbb{P}_n) = \max_{\theta \in \Theta_I(\mathbb{P}_n)} \psi(\theta), \quad (2.11)$$

Note that when  $T$  is large, there can be a large number of constraints defining the set  $\Theta_I(\mathbb{P}_n)$ , and the partially identified parameter  $\theta$  can be high-dimensional.

**Example 4** (Inference on Counterfactual Policies). In the setting of [Kasy \(2016\)](#), the researcher is interested in ranking counterfactual policies “A” and “B” which represent two competing proposals of assigning individuals to some treatment based on covariate values. It is assumed that the policy maker only has knowledge of the partially-identified parameters  $g_0(X) := \mathbb{E}[Y_0|X]$  and  $g_1(X) := \mathbb{E}[Y_1|X]$ , where  $Y_d$  is the partially-observed potential outcome for treatment state  $D = d$ .

We assume that the researcher’s object of interest is the linear functional  $\psi := \psi(f^A, f^B)$  where  $f^A$  is the distribution of the random variable  $Y^A$  representing the observed outcome under policy A, and  $f^B$  is the distribution of the random variable  $Y^B$  representing the observed outcome under policy B. Furthermore, let  $D^A$  be the random variable representing treatment assignment under policy A, and let  $D^B$  be the random variable representing treatment under assignment B, and assume that  $D^A, D^B \perp (Y_0, Y_1)|X$ . Some simple objective functions include  $\psi^A := \mathbb{E}[Y^A]$  (or  $\psi^B := \mathbb{E}[Y^B]$ ), which measures the average outcome under policy A, or  $\psi^{AB} := \mathbb{E}[Y^A - Y^B]$ , which measures the difference in average outcomes between policies A and B. Let  $\mathcal{G}_d$  denote the identified set for  $g_d(X)$ . Note that the objective function  $\psi^A$  can be decomposed as:

$$\begin{aligned} \psi^A &= \mathbb{E}[Y^A], \\ &= \mathbb{E}[\mathbb{E}[Y^A|D^A = 1, X] P(D^A = 1|X) + \mathbb{E}[Y^A|D^A = 0, X] (1 - P(D^A = 1|X))], \\ &= \mathbb{E}[\mathbb{E}[Y_0|X] + P(D^A = 1|X) (\mathbb{E}[Y_1|X] - \mathbb{E}[Y_0|X])], \\ &= \mathbb{E}[g_0(X) + h^A(X) (g_1(X) - g_0(X))]. \end{aligned}$$

where  $h^A(X) = P(D^A = 1|X)$ . Since  $g_0(\cdot)$  and  $g_1(\cdot)$  are only partially-identified,  $\psi^A$  will also be partially identified. Let  $\Psi_I^A(P) = [\Psi_{\ell b}^A(P), \Psi_{ub}^A(P)]$  denote the identified set for  $\psi^A = \mathbb{E}[Y^A]$ , where the endpoints of



$\Psi_I^A$  are determined by:

$$\Psi_{lb}^A(P) = \inf_{(g_0, g_1) \in \mathcal{G}_0 \times \mathcal{G}_1} \sum_{x \in \mathcal{X}} [g_0(x) + h^A(x) (g_1(x) - g_0(x))] P(X = x), \quad (2.12)$$

$$\Psi_{ub}^A(P) = \sup_{(g_0, g_1) \in \mathcal{G}_0 \times \mathcal{G}_1} \sum_{x \in \mathcal{X}} [g_0(x) + h^A(x) (g_1(x) - g_0(x))] P(X = x), \quad (2.13)$$

where  $P(X = x)$  is the probability  $X = x$  in the target population. Similarly, as in [Kasy \(2016\)](#), the objective function  $\psi^{AB}$  can be decomposed as:

$$\begin{aligned} \psi^{AB} &= \mathbb{E} [Y^A - Y^B], \\ &= \mathbb{E} [(h^A(X) - h^B(X)) (Y_1 - Y_0)], \\ &= \mathbb{E} [h^{AB}(X)g(X)]. \end{aligned}$$

where  $h^{AB}(X) = h^A(X) - h^B(X)$ ,  $h^A(X) = P(D^A = 1|X)$ ,  $h^B(X) = P(D^B = 1|X)$  and  $g(X) = g_1(X) - g_0(X)$ . Since  $g(X)$  is only partially identified, the objective function  $\psi^{AB}$  will also only be partially identified. Let  $\Psi_I^{AB}(P) = [\Psi_{lb}^{AB}(P), \Psi_{ub}^{AB}(P)]$  denote the identified set for  $\psi^{AB}$ , where the endpoints of  $\Psi_I^{AB}$  are given by:

$$\Psi_{lb}^{AB}(P) = \inf_{(g_0, g_1) \in \mathcal{G}_0 \times \mathcal{G}_1} \sum_{x \in \mathcal{X}} h^{AB}(x) (g_1(x) - g_0(x)) P(X = x), \quad (2.14)$$

$$\Psi_{ub}^{AB}(P) = \sup_{(g_0, g_1) \in \mathcal{G}_0 \times \mathcal{G}_1} \sum_{x \in \mathcal{X}} h^{AB}(x) (g_1(x) - g_0(x)) P(X = x), \quad (2.15)$$

where  $P(X = x)$  is the probability  $X = x$  in the target population. In this example, note that the partially identified parameter is  $\theta = (g_0, g_1)$  and the identified set is  $\Theta_I(P) = \mathcal{G}_0 \times \mathcal{G}_1$ .

**Remark 2.1.** In practice, the probabilities  $P(X = x)$  in the optimization problems (2.12) and (2.13), or (2.14) and (2.15), may need to be estimated, meaning that the objective functions in these optimization problems contain sampling uncertainty that must be accounted for when performing inference on either  $\Psi_I^A$  or  $\Psi_I^{AB}$  in addition to the sampling uncertainty inherent in the estimation of the sets  $\mathcal{G}_0$  and  $\mathcal{G}_1$ . Currently, we are unaware of any uniformly valid inference procedure in partially identified models that can handle these cases.

### 3 Methodology

In this section, we develop the ideas introduced in the previous section. We consider a setting when the identified set  $\Theta_I(P)$  is defined by moment equalities and inequalities that are satisfied at the true parameter  $\theta_0$ :

$$Pm_j(W, \theta_0) = 0, \quad \text{for } j = 1, \dots, r_1, \quad (3.1)$$

$$Pm_j(W, \theta_0) \leq 0, \quad \text{for } j = r_1 + 1, \dots, r_1 + r_2. \quad (3.2)$$



Note that we can always convert these moment equalities/inequalities defined above into  $k = 2r_1 + r_2$  equivalent moment inequalities given by:

$$Pm_j(W, \theta_0) \leq 0, \quad \text{for } j = 1, \dots, r_1, \quad (3.3)$$

$$-Pm_j(W, \theta_0) \leq 0, \quad \text{for } j = 1, \dots, r_1, \quad (3.4)$$

$$Pm_j(W, \theta_0) \leq 0, \quad \text{for } j = r_1 + 1, \dots, r_1 + r_2 \quad (3.5)$$

Thus, we will assume throughout most of the exposition that the model is defined only by  $k$  moment inequalities:

$$Pm_j(W, \theta_0) \leq 0, \quad \text{for } j = 1, \dots, k. \quad (3.6)$$

Only on rare occasions will it be necessary to know which of the moment inequalities correspond to moment equalities; in these cases we will simply refer back to the original formulation in (3.1) and (3.2).

We assume that the researcher is interested in bounding the expected value of a functional  $\psi : \mathcal{W} \times \Theta \rightarrow \mathbb{R}$ . Define the following class of functions:

$$\mathcal{F} := \left\{ (\psi(W, \theta), m_1(W, \theta), \dots, m_k(W, \theta))^T : \theta \in \Theta \right\}. \quad (3.7)$$

A typical element of  $\mathcal{F}$  will then be the vector-valued function:

$$f(W, \theta) = \left[ \psi(W, \theta), m_1(W, \theta), \dots, m_k(W, \theta) \right]^T.$$

Furthermore, we will equip this class of functions with a semimetric that depends on the probability measure  $P$ :

$$\rho_P(\theta, \theta') := \left\| \text{diag}(V_P(f(W, \theta) - f(W, \theta'))^{1/2}) \right\|, \quad (3.8)$$

for  $\theta, \theta' \in \Theta$ .<sup>1</sup> This semimetric was also considered in [Bugni et al. \(2015\)](#). Furthermore, define the class:

$$\mathcal{F}' := \{f - f' : f, f' \in \mathcal{F}\},$$

and let  $\mathcal{G} = \mathcal{F} \cup \mathcal{F}' \cup (\mathcal{F}')^2$ . Let  $\{P_n \in \mathcal{P}\}_{n \geq 1}$  be any sequence of data-generating measures. Throughout the text, we will interpret the statement  $P_n \rightarrow P$  as meaning  $\|P_n - P\|_{\mathcal{G}} \rightarrow 0$ , where  $\|\cdot\|_{\mathcal{G}}$  represents the sup-norm over the class of functions  $\mathcal{G}$ . This is the relevant notion of the convergence of drifting sequences of probability measures for the results in this paper; see [Sheehy and Wellner \(1992\)](#) and [Van Der Vaart and Wellner \(1996\)](#) Section 2.8.3. We will later impose assumptions that guarantee weak convergence over any such drifting sequence. Throughout the article we will switch freely between considering uniform convergence over  $\mathcal{P}$  and convergence over any convergent data-generating sequence  $\{P_n \in \mathcal{P}\}_{n=1}^\infty$  in the sense described above.

---

<sup>1</sup>Recall a semimetric satisfies (i)  $\rho(f, f) = 0$ , (ii)  $\rho(f, g) \leq \rho(f, h) + \rho(h, g)$  and (iii)  $\rho(x, y) = \rho(y, x)$ . However, unlike a metric, a semimetric can be equal to zero when evaluated at two distinct elements.



Let  $\mathcal{P}$  denote the collection of all probability measures on  $\mathcal{W}$ . We now impose the following assumptions:

**Assumption 3.1.** *The parameter space  $(\Theta, \mathcal{P})$  satisfies the following conditions:*

- (i)  $\Theta \subset \mathbb{R}^{d_\theta}$  is convex and compact.
- (ii)  $\mathcal{F}$  is a measurable class of functions.
- (iii) Each distribution  $P \in \mathcal{P} \subseteq \mathcal{P}$  satisfies:
  - (a)  $Pm_j(W, \theta_0) \leq 0$ , for  $j = 1, \dots, k$ .
  - (b) In a sample  $\{W_i\}_{i=1}^n$ ,  $W_i$  are independent and identically distributed according to  $P \in \mathcal{P}$ .
- (iv) There exists a bounded envelope function  $F$  for the class  $\mathcal{F}$  such that for some  $a > 0$ ,

$$\sup_{P \in \mathcal{P}} P||F(W)||^{2+a} < \infty.$$

**Remark 3.1.** In Assumption 3.1(ii), we call  $\mathcal{F}$  measurable if  $\mathcal{F}$  is  $P$ -measurable in the sense of [Van Der Vaart and Wellner \(1996\)](#) Definition 2.3.3 for all probability measures  $P \in \mathcal{P}$ .

Note that we can write the identified set  $\Theta_I(P)$  as:

$$\Theta_I(P) = \{\theta \in \Theta : Pm_j(W, \theta) \leq 0, \quad j = 1, \dots, k\}. \quad (3.9)$$

Now let  $\Theta_I(\mathbb{P}_n)$  denote the estimate of the identified set:

$$\Theta_I(\mathbb{P}_n) = \{\theta \in \Theta : \mathbb{P}_n m_j(W, \theta) \leq 0, \quad j = 1, \dots, k\}. \quad (3.10)$$

where  $\mathbb{P}_n$  denotes the empirical measure for the first  $n$  observations:

$$\mathbb{P}_n := \frac{1}{n} \sum_{i=1}^n \delta_{W_i}, \quad (3.11)$$

where  $\delta_{W_i}$  is the Dirac delta function. We restrict attention to a certain class of functionals of the identified set.

**Assumption 3.2.** (i) The functional of interest  $\psi(w, \theta) : \mathcal{W} \times \Theta \rightarrow \mathbb{R}$  is linear in  $\theta$ , and is continuous in  $w \in \mathcal{W}$ ; (ii) the functions  $m_j(w, \theta) : \mathcal{W} \times \Theta \rightarrow \mathbb{R}$  are linear in  $\theta$  and continuous in  $w \in \mathcal{W}$  for  $j = 1, \dots, k$ .

Denote the identified set for  $P\psi(W, \theta)$  to be  $\Psi_I(P)$ , and note that the identified set for  $P\psi(W, \theta)$  is the projection of  $\Theta_I(P)$  on the manifold generated by  $P\psi(W, \theta)$ . As such, under standard conditions (see Lemma A.4(ii)) the projection estimator  $\Psi_I(\mathbb{P}_n)$  will be a consistent estimator of  $\Psi_I(P)$ . Moreover, since  $P\psi(W, \theta)$  is continuous and  $\Theta_I(P)$  is convex and compact, the identified set  $\Psi_I(P)$  is an interval— $\Psi_I(P) = [\Psi_I^{\ell b}(P), \Psi_I^{ub}(P)]$ —with endpoints determined by:

$$\Psi_I^{\ell b}(P) := \inf_{\theta \in \Theta} P\psi(W, \theta) \quad \text{s.t.} \quad Pm_j(W, \theta) \leq 0, \quad j = 1, \dots, k, \quad (3.12)$$



$$\Psi_I^{ub}(P) := \sup_{\theta \in \Theta} P\psi(W, \theta) \quad \text{s.t.} \quad Pm_j(W, \theta) \leq 0, \quad j = 1, \dots, k. \quad (3.13)$$

However, since  $P$  is not known, the programs (3.12) and (3.13) will be approximated using the empirical distribution  $\mathbb{P}_n$  by replacing the population moment conditions and objective function with their sample counterparts:

$$\Psi_I^{\ell b}(\mathbb{P}_n) := \inf_{\theta \in \Theta} \mathbb{P}_n\psi(W, \theta) \quad \text{s.t.} \quad \mathbb{P}_nm_j(W, \theta) \leq 0, \quad j = 1, \dots, k, \quad (3.14)$$

$$\Psi_I^{ub}(\mathbb{P}_n) := \sup_{\theta \in \Theta} \mathbb{P}_n\psi(W, \theta) \quad \text{s.t.} \quad \mathbb{P}_nm_j(W, \theta) \leq 0, \quad j = 1, \dots, k. \quad (3.15)$$

After an estimate of the identified set is obtained, interest will lie in constructing uniformly valid confidence sets for the true parameter  $\psi_0 := P\psi(W, \theta_0)$ . To perform inference on the true parameter using the optimal values  $(\Psi_I^{\ell b}(P), \Psi_I^{ub}(P))$  in programs (3.12) and (3.13), we will approximate the distributions of  $\sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P))$  and  $\sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n) - \Psi_I^{ub}(P))$  by a simple bootstrap procedure, and will be particularly interested in proving the procedure is valid *uniformly over  $\mathcal{P}$* .

**Remark 3.2.** *As a technical note, the functions  $(\Psi_I^{\ell b}(\cdot), \Psi_I^{ub}(\cdot))$  will be seen as maps from  $\mathcal{P}_+$  to  $\mathbb{R}$ , where  $\mathcal{P}_+$  is defined as the collection of all measures  $\mathcal{P}$  as well as all finite empirical measures  $\mathbb{P}_n$  generated by a  $P \in \mathcal{P}$  (i.e.  $\mathcal{P}_+ = \text{span}(\mathcal{P}, \{\delta_w\}_{w \in \mathcal{W}})$ , where  $\{\delta_w\}_{w \in \mathcal{W}}$  is any finite collection of point masses). It will be useful to distinguish between the collections  $\mathcal{P}$  and  $\mathcal{P}_+$  throughout.*

### 3.1 Value Function Differentiability

We will begin by showing that the value functions satisfy an appropriate level of differentiability with respect to the underlying probability measure. Since the underlying probability measure is a possibly infinite-dimensional object, we must use a form of differentiability that is valid between metrizable topological spaces. In particular, it is well-known (e.g. Shapiro (1990), Shapiro (1991)) that under some conditions the functions  $(\Psi_I^{\ell b}(P), \Psi_I^{ub}(P))$  are *Hadamard directionally differentiable*. To introduce the differentiability concepts used in this paper in general form, let  $(\mathbb{D}, d_D)$  and  $(\mathbb{E}, d_E)$  be metric spaces.

**Definition 3.1** (Hadamard Directional Differentiability). *A map  $\phi : \mathbb{D}_\phi \subseteq \mathbb{D} \rightarrow \mathbb{E}$  is called Hadamard directionally differentiable at  $\zeta \in \mathbb{D}_\phi$  if there is a linear map  $\phi'_\zeta : \mathbb{D} \rightarrow \mathbb{E}$  such that*

$$\frac{\phi(\zeta + t_n h_n) - \phi(\zeta)}{t_n} \rightarrow \phi'_\zeta(h),$$

*for converging sequences  $\{t_n\} \subset \mathbb{R}_+$  with  $t_n \downarrow 0$  and  $h_n \rightarrow h$  such that  $\zeta + t_n h_n \in \mathbb{D}_\phi$  for every  $n$ . In addition, we say  $\phi$  is Hadamard directionally differentiable tangential to a set  $\mathbb{D}_0 \subseteq \mathbb{D}$  if we also require that the limit  $h \in \mathbb{D}_0$  in the above.*



While Hadamard directional differentiability can be used to justify an inference procedure in stochastic programs for a fixed data-generating measure  $P \in \mathcal{P}$  (c.f. [Shapiro \(1991\)](#)), it is not sufficient to construct an inference procedure for stochastic programs that is valid *uniformly* over  $\mathcal{P}$ . It is natural to wonder whether stochastic programs are uniformly Hadamard directionally differentiable, which is defined in the following:

**Definition 3.2** (Uniform Hadamard Directional Differentiability). *Let  $\phi : \mathbb{D}_\phi \subseteq \mathbb{D} \rightarrow \mathbb{E}$ ,  $\mathbb{D}_0 \subseteq \mathbb{D}$ , and  $\mathbb{D}_\zeta \subseteq \mathbb{D}_\phi$ . The map  $\phi : \mathbb{D}_\phi \subseteq \mathbb{D} \rightarrow \mathbb{E}$  is called uniformly Hadamard directionally differentiable in  $\zeta \in \mathbb{D}_\zeta$  if there is a continuous map  $\phi'_\zeta : \mathbb{D} \rightarrow \mathbb{E}$  such that*

$$\frac{\phi(\zeta_n + t_n h_n) - \phi(\zeta_n)}{t_n} \rightarrow \phi'_\zeta(h), \quad (3.16)$$

*for all converging sequences  $\zeta_n \rightarrow \zeta \in \mathbb{D}_\zeta$ ,  $\{t_n\} \subset \mathbb{R}_+$  with  $t_n \downarrow 0$ , and  $h_n \rightarrow h$  such that  $\zeta_n + t_n h_n \in \mathbb{D}_\phi$  for every  $n$ . In addition, we say  $\phi$  is uniformly Hadamard directionally differentiable tangential to a set  $\mathbb{D}_0 \subseteq \mathbb{D}$  if we also require that the limit  $h \in \mathbb{D}_0$  in the above.*

This definition is analogous to the extension of Hadamard differentiability to uniform Hadamard differentiability presented in [Belloni et al. \(2017\)](#), although our definition restricts  $t_n \rightarrow 0$  from above (providing a “direction”). It also allows the spaces involved to be metric spaces rather than normed linear spaces, and allows the derivative map  $\phi'_\zeta$  to be continuous rather than linear.

In addition, reflecting more the definition in [Van Der Vaart and Wellner \(1996\)](#) p. 379, we do not explicitly require that the derivative map  $(\zeta, h) \mapsto \phi'_\zeta(h)$  be continuous at every  $(\zeta, h)$ , as is done in the extension of Hadamard differentiability to uniform Hadamard differentiability in [Belloni et al. \(2017\)](#). However, similar to [Belloni et al. \(2017\)](#), we will use the flexibility provided by the above definition to allow  $\zeta_n$  to lie outside  $\mathbb{D}_\zeta$ .

As we will see, under some conditions the value functions of a stochastic program are differentiable in the sense of Definition 3.2. Our first main result requires non-emptiness of the identified set, the existence and uniqueness of Lagrange multipliers, and uniqueness of the optimal solutions in the programs (3.12) and (3.13). To guarantee these properties will require that a “uniform constraint qualification” holds for the linear programs. We impose such a constraint qualification in the next assumption.

Let  $\mathcal{A}(\theta, P)$  be an index set defined as:

$$\mathcal{A}(\theta, P) := \{j \in \{r_1 + 1, \dots, r_1 + r_2\} : Pm_j(W, \theta) = 0\} \quad (3.17)$$

i.e.  $\mathcal{A}(\theta, P)$  denotes the set indexing the binding moment inequalities at  $\theta$  for some probability measure  $P \in \mathcal{P}$ .

**Assumption 3.3.** *Let  $G(\theta, P)$  be the matrix formed by vertically stacking the row vectors  $\nabla_\theta \psi(W, \theta)$ ,  $\{\nabla_\theta Pm_j(W, \theta)\}_{j=1}^{r_1}$  and  $\{\nabla_\theta Pm_j(W, \theta)\}_{j \in \mathcal{A}(\theta, P)}$ . There exists  $\varepsilon > 0$  such that:*

- (i)  $\Theta_I(P) \neq \emptyset$  for all  $P \in \mathcal{P}^\varepsilon$ , where  $\mathcal{P}^\varepsilon := \{Q \in \mathcal{P}_+ : \|Q - P\|_{\mathcal{G}} \leq \varepsilon, P \in \mathcal{P}\}$ .



(ii) There exists a  $\kappa > 0$  such that:

$$\inf_{P \in \mathcal{P}^\varepsilon} \inf_{\theta \in \Theta_I(P)} \text{eig}(\mathbf{G}(\theta, P)\mathbf{G}(\theta, P)^T) \geq \kappa \quad (3.18)$$

where  $\text{eig}(A)$  denotes the minimum eigenvalue of  $A$ .

Assumption 3.3(i) implies that, for large enough  $n$ , the identified set is non-empty. Given non-emptiness of the identified set, Assumption 3.3(ii) implies a uniform version of the *linear independence constraint qualification* (LICQ), and is instrumental in ensuring the existence and uniqueness of optimal solutions and Lagrange multipliers in the programs (3.12) and (3.13). Constraint qualifications in various forms have appeared throughout the recent history of partial identification (e.g. Beresteanu and Molinari (2008), Pakes et al. (2011) Kaido and Santos (2014), Freyberger and Horowitz (2015), Kaido et al. (2017), Gafarov et al. (2018), and Gafarov (2018)). There are some cases where it may be easy to directly verify that the Assumption 3.3(ii) is satisfied, but in general Assumption 3.3 is a high-level condition.<sup>2</sup>

Assumption 3.3 is a regularity assumption that is important in the proof of uniform Hadamard directional differentiability. Specifically, it is necessary to restrict the optimal solutions and Lagrange multipliers in (3.12) and (3.13) to be unique for all  $P \in \mathcal{P}$ . To understand why, note that if the problems (3.12) and (3.13) admit multiple solutions there may be differences between the sets representing “the limiting optimal solutions” (over the sequence  $\{P_n \in \mathcal{P}\}_{n=1}^\infty$ ), and the sets representing “the optimal solutions at the limit” ( $P \in \mathcal{P}$ ). This is related to the Theorem of the Maximum, and the fact that the Theorem of the Maximum guarantees only that the solution correspondence is upper hemicontinuous, but not lower hemicontinuous (and thus, not continuous). In this case it is possible to show that the value functions  $\Psi_I^{\ell b}(\cdot)$  and  $\Psi_I^{ub}(\cdot)$  are Hadamard directionally differentiable, but not necessarily *uniformly* Hadamard directionally differentiable. The same intuition follows for the Lagrange multipliers. However, the only case when there are non-unique solutions in linear programs are when the gradient of the objective function is exactly zero, or is some multiple of the gradient of one of the moment conditions. Both these cases are ruled out by Assumption 3.3(ii). In addition, Wachsmuth (2013) shows that the LICQ—implied by Assumption 3.3—is the weakest constraint qualification under which the Lagrange multipliers are guaranteed to be unique. Since the LICQ is implied uniformly over  $\mathcal{P}^\varepsilon$  by Assumption 3.3(ii), existence and uniqueness of Lagrange multipliers also follows from Assumption 3.3(ii). Assumption 3.3(i) enables us to impose Assumption 3.3(ii) over  $\mathcal{P}^\varepsilon$ , but is also required to ensure that the uniform Hadamard directional derivative is well-defined, which would not be the case if the identified set was allowed to be empty for all  $n$ . Again, this is a high-level condition, and we hope to provide more primitive conditions in future drafts.

A few additional remarks about Assumption 3.3 are in order. First, Assumption 3.3 is one of the rare cases where it is useful to distinguish between moment equalities as in (3.1) and moment inequalities as in (3.2), since the assumption imposes different conditions on the two types of moments.

<sup>2</sup>For example, if the moment functions can be expressed as  $Pm_j(W, \theta) = P\tilde{m}_j(W) + a_j'\theta$ , where  $\tilde{m}_j$  is a function of the random variable  $W \in \mathcal{W}$ , and  $a_j \in \mathbb{R}^d$  is a vector, then it suffices to verify the Jacobian of the moment functions (w.r.t.  $\theta$ ) has full column rank.



Next note that it is possible to show that Assumption 3.3 implies that at every  $P \in \mathcal{P}$  there must be at least one interior point of the set defined by the moment inequalities at which all moment equalities are satisfied. The major restriction imposed by this implication is that the moment inequalities evaluated at the limiting  $P \in \mathcal{P}$  cannot point-identify the parameter of interest.<sup>3</sup> This condition is reminiscent of condition 4 in Theorem 2.1 in Shi and Shum (2015), and its discussion on page 499 of Shi and Shum (2015). Similar to their discussion, we note that in many cases this assumption will fail when two inequality constraints become equivalent, in which case the inequality constraints can be combined to form an equality constraint so that the assumption still holds.

Finally, note that this assumption is sufficient for our method to be uniformly valid, but is not necessary. However, the assumption is the most primitive assumption we are currently aware of, as it connects to the highly used constraint qualification assumptions in optimization literature while imposing minimal constraints on any sequence  $\{P_n \in \mathcal{P}_+\}_{n=1}^\infty$  required for uniformity.

The final assumption relates to the gradient of the objective function and the moments:

**Assumption 3.4.** *The gradients  $\{\nabla_\theta P\psi(W, \theta), \{\nabla_\theta Pm_j(W, \theta)\}_{j=1}^k\}$  are uniformly bounded over  $\mathcal{P}^\varepsilon$ .*

This assumption is required only to show that the Lagrange multipliers are uniformly bounded over  $\mathcal{P}^\varepsilon$ . Any other assumption that implies uniform boundedness of the Lagrange multipliers might then be safely substituted for Assumption 3.4.

Finally, as a piece of technical machinery, we define the tangent cone as:

$$\mathcal{T}_P(\mathcal{F}) = \{v \in UC_b(\mathcal{F}, \rho_P) : \forall t_n \downarrow 0, \forall \{P_n \in \mathcal{P}\}_{n=1}^\infty \rightarrow P \in \mathcal{P}, \exists \{Q_n \in \mathcal{P}_+\}_{n=1}^\infty \text{ s.t. } t_n^{-1}(Q_n - P_n) \rightarrow v\}, \quad (3.19)$$

where  $UC_b(\mathcal{F}, \rho_P) \subset \ell^\infty(\mathcal{F})$  denotes the space of bounded, and uniformly continuous functions with respect to the semimetric  $\rho_P$  defined in (3.8). While restricting the tangent cone to be a subset of  $UC_b(\mathcal{F}, \rho_P)$  might appear to be restrictive, under the Donsker-type assumptions to be introduced later almost all paths of the limit of the empirical process  $\sqrt{n}(\mathbb{P}_n - P_n)$  will be uniformly continuous; see Addendum 1.5.8 in Van Der Vaart and Wellner (1996). We now have the following result:

**Theorem 3.1.** *Suppose Assumptions 3.1 - 3.4 hold, and consider  $\Psi_I^{\ell b}, \Psi_I^{ub} : \mathcal{P}_+ \rightarrow \mathbb{R}$  defined by the programs (3.14) and (3.15). Then  $\Psi_I^{\ell b}(\cdot), \Psi_I^{ub}(\cdot)$  are uniformly Hadamard directionally differentiable tangential to  $\mathcal{T}_P(\mathcal{F})$ . In particular, for all converging sequences  $P_n \rightarrow P \in \mathcal{P}$ ,  $\{t_n\} \subset \mathbb{R}_+$  with  $t_n \downarrow 0$ , and  $h_n \rightarrow h \in \mathcal{T}_P(\mathcal{F})$  such that  $P_n + t_n h_n \in \mathcal{P}_+$  for every  $n$ , we have:*

$$(\Psi_I^{\ell b})'_P(h) := \lim_{n \rightarrow \infty} \frac{\Psi_I^{\ell b}(P_n + t_n h_n) - \Psi_I^{\ell b}(P_n)}{t_n} = h_1 \psi(W, \theta_{\ell b}^*(P)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P) h_{j+1} m_j(W, \theta_{\ell b}^*(P)), \quad (3.20)$$

---

<sup>3</sup>Note also that this condition rules out the case that the moment inequalities define an empty region. However, we do not consider this a ‘‘major restriction’’ of our method, since if the true identified set is empty then computing functionals over the identified set becomes a dubious exercise.



$$(\Psi_I^{ub})'_P(h) := \lim_{n \rightarrow \infty} \frac{\Psi_I^{ub}(P_n + t_n h_n) - \Psi_I^{ub}(P_n)}{t_n} = -h_1 \psi(W, \theta_{ub}^*(P)) + \sum_{j=1}^k \lambda_{ub,j}^*(P) h_{j+1} m_j(W, \theta_{ub}^*(P)), \quad (3.21)$$

where  $h_j f_j$  is the  $j^{\text{th}}$  component of  $h f$  for  $f \in \mathcal{F}$ ,  $\theta_{lb}^*(P)$  and  $\theta_{ub}^*(P)$  are the optimal solutions in the lower and upper bounding problems at  $P \in \mathcal{P}$ , and  $\{\lambda_{lb,j}^*(P)\}_{j=1}^k$  and  $\{\lambda_{ub,j}^*(P)\}_{j=1}^k$  are the Lagrange multipliers in the lower and upper bounding problems at  $P \in \mathcal{P}$ .

The uniform component of this theorem lies in the fact that it is valid over any generating sequence  $\{P_n \in \mathcal{P}\} \rightarrow P \in \mathcal{P}$ . This uniform version of differentiability turns out to be sufficient to apply the extended continuous mapping theorem (Theorem 1.11.1 in [Van Der Vaart and Wellner \(1996\)](#)) in order to relate this result to inference on the optimal value functions. This is exactly what is done in Lemma 3.1 in the next subsection.

### 3.2 From Differentiability to Weak Convergence

We now consider the asymptotic distribution of the properly rescaled and recentered value functions given in (3.14) and (3.15), which will make use of the uniform differentiability property given in Theorem 3.1. To cover the case of a drifting sequence of data-generating processes, which will be necessary to show uniformity, we impose additional assumptions.

**Assumption 3.5.** *The collections  $\mathcal{F}$  and  $\mathcal{P}$  satisfy the following:*

- (i) *The empirical process  $\mathbb{G}_{n,P} := \sqrt{n}(\mathbb{P}_n - P)$  is asymptotically equicontinuous uniformly over  $\mathcal{P}$ ; that is, for every  $\varepsilon > 0$ ,*

$$\lim_{\delta \downarrow 0} \limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} P_P^* \left( \sup_{\rho_P(\theta, \theta') < \delta} \|\mathbb{G}_{n,P} f(W, \theta) - \mathbb{G}_{n,P} f(W, \theta')\| > \varepsilon \right) = 0,$$

where  $\rho_P$  is as in (3.8).

- (ii) *The semimetric  $\rho_P$  satisfies:*

$$\lim_{\delta \downarrow 0} \sup_{\|(\theta_1, \theta'_1) - (\theta_2, \theta'_2)\| < \delta} \sup_{P \in \mathcal{P}} |\rho_P(\theta_1, \theta'_1) - \rho_P(\theta_2, \theta'_2)| = 0$$

- (iii) *Let  $\mathcal{A}(\theta, P) \subseteq \{r_1 + 1, \dots, r_1 + r_2\}$  denote the binding moment inequalities at  $(\theta, P)$ , let  $\mathcal{I}_{eq} = \{1, \dots, r_1\}$ , and let  $V_j(\theta) := \text{Var}(m_j(W, \theta))$ , for  $j = 1, \dots, r_1 + r_2$ . Then there exists a constant  $\underline{v} > 0$  such that for all  $P \in \mathcal{P}$ :*

$$\inf_{\theta \in \Theta_I(P)} \min_{j \in \mathcal{A}(\theta, P) \cup \mathcal{I}_{eq}} V_j(\theta) \geq \underline{v}$$

- (iv) *Let  $\mathcal{A}(\theta, P) \subseteq \{r_1 + 1, \dots, r_1 + r_2\}$  denote the binding moment inequalities at  $(\theta, P)$ , and let  $\underline{\sigma} > 0$  be a constant. One of the following two holds:*



(a) Let  $\mathbf{V}_P^m := \text{Var}_P \{ \{m_j(W, \theta)\}_{j=1}^{r_1}, \{m_j(W, \theta)\}_{j \in \mathcal{A}(\theta, P)} \}$ . The objective function  $\psi(w, \theta)$  is a trivial function of  $w$ , and we have:

$$\inf_{\theta \in \Theta_I(P)} \text{eig}(\mathbf{V}_P^m) \geq \underline{\sigma}$$

(b) Let  $\mathbf{V}_P := \text{Var}_P \{ \psi(W, \theta), \{m_j(W, \theta)\}_{j=1}^{r_1}, \{m_j(W, \theta)\}_{j \in \mathcal{A}(\theta, P)} \}$ . Then we have:

$$\inf_{\theta \in \Theta_I(P)} \text{eig}(\mathbf{V}_P) \geq \underline{\sigma}$$

(v) There exist positive constants  $C, \delta > 0$  such that  $\max_{j=1, \dots, k} |Pm_j(W, \theta)| \geq C \min(\delta, d_H(\theta, \Theta_I(P)))$  for every  $P \in \mathcal{P}$  and  $\theta \in \Theta$ , where  $d_H$  is the Hausdorff metric.

Assumption 3.5(i) is required to apply a uniform central limit theorem to the class of functions  $\mathcal{F}$ . Assumption 3.5(ii) is also required to prove the class of functions  $\mathcal{F}$  is uniformly Donsker over any sequence in  $\mathcal{P}$ . Assumption 3.5(iii) and 3.5(iv) are required to ensure a uniform multivariate central limit theorem holds for the moment functions. These assumptions are related to Assumption 4.3 in Kaido et al. (2017), and are also required for our bootstrap procedure to hold. The option (a) or (b) in Assumption 3.5(iv) splits the cases when the researcher's objective function depends on  $W$  (such as in Example 4) with the cases when the researcher's objective function does not depend on  $W$  (such as in subvector inference). Finally, Assumption 3.5(iv) is the partial identification condition given in Chernozhukov et al. (2007), equation (4.5), and is useful when establishing the Hausdorff consistency and the rate of convergence of the estimated identified set to the true identified set.

In the following lemma, for any sequence  $\{P_n \in \mathcal{P}\}_{n=1}^\infty$  converging to the Borel probability measure  $P \in \mathcal{P}$ , we let  $\mathbb{G}_{n, P_n} := \sqrt{n}(\mathbb{P}_n - P_n) \in \ell^\infty(\mathcal{F})$  denote the empirical process indexed by  $P_n$ . Adding Assumption 3.5, we have the following result:

**Lemma 3.1.** *Suppose Assumptions 3.1 - 3.5 hold. Then for any sequence  $\{P_n \in \mathcal{P}\}_{n=1}^\infty \rightarrow P \in \mathcal{P}$  we have  $\mathbb{G}_{n, P_n} \rightsquigarrow \mathbb{G}_P$  where  $\mathbb{G}_P$  is a tight Borel measurable element in  $\mathcal{T}_P(\mathcal{F})$ , and:*

$$\sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P_n)) \rightsquigarrow (\Psi_I^{\ell b})'_P(\mathbb{G}_P), \quad (3.22)$$

$$\sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n) - \Psi_I^{ub}(P_n)) \rightsquigarrow (\Psi_I^{ub})'_P(\mathbb{G}_P). \quad (3.23)$$

This result follows from the extended continuous mapping theorem (Theorem 1.11.1 in Van Der Vaart and Wellner (1996)) in combination with the result of Theorem 3.1. When combined with Theorem 3.1, Lemma 3.1 shows that the properly recentered and rescaled value functions converge in distribution to  $(\Psi_I^{\ell b})'_P(\mathbb{G}_P)$  and  $(\Psi_I^{ub})'_P(\mathbb{G}_P)$ , evaluated at the limiting empirical process  $\mathbb{G}_P$ , along any converging sequence  $\{P_n \in \mathcal{P}\}_{n=1}^\infty$  satisfying Assumptions 3.1-3.5. The next section shows that the objects on the right side of (3.22) and (3.23) can be approximated uniformly using a nonparametric bootstrap procedure.



### 3.3 The Bootstrap Version

This section proposes a bootstrap procedure that will allow us to consistently estimate the distributions of the value functions  $(\Psi_I^{\ell b}(P), \Psi_I^{ub}(P))$  uniformly over  $\mathcal{P}$ . In particular, we propose the following approximations:

$$\text{Lower Approximation: } \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n^b) - \Psi_I^{\ell b}(\mathbb{P}_n)), \quad (3.24)$$

$$\text{Upper Approximation: } \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n^b) - \Psi_I^{ub}(\mathbb{P}_n)). \quad (3.25)$$

We will use the distribution of (3.24) to approximate the distribution of  $(\Psi_I^{\ell b})'_P(\mathbb{G}_P)$ , and we will use the distribution of (3.25) to approximate the distribution of  $(\Psi_I^{ub})'_P(\mathbb{G}_P)$ .

**Remark 3.3.** *Note, unlike typical inference procedures, we do not standardize the moment conditions by their sample standard deviations. However, our procedure is still invariant to rescaling of the moment conditions by the fact that any rescaling will be reflected in the procedure as an equivalent (but opposite) rescaling of the Lagrange multipliers.*

We must be precise about the conditions under which the law of the approximations (3.24) and (3.25), conditional on the data  $\{W_i\}_{i=1}^n$ , can approximate the unconditional law of  $(\Psi_I^{\ell b})'_P(\mathbb{G}_P)$  and  $(\Psi_I^{ub})'_P(\mathbb{G}_P)$  uniformly over  $\mathcal{P}$ . Let  $\{\{W_i^b\}_{i=1}^n : b = 1, \dots, B\}$  denote the bootstrap samples. We maintain the following assumption:

**Assumption 3.6.** *The bootstrap samples  $\{W_i^b\}_{i=1}^n$  for  $b = 1, \dots, B$ , are drawn i.i.d. with replacement from the original sample  $\{W_i\}_{i=1}^n$ .*

The following lemma, which is necessary for our main result, shows that the proposed bootstrap procedure is uniformly valid:

**Lemma 3.2.** *Suppose that conditional on  $\{W_i\}_{i=1}^n$  we have that, uniformly over  $\mathcal{P}$ ,  $\mathbb{G}_n^b \rightsquigarrow \mathbb{G}_P$  where  $\mathbb{G}_P$  is a tight random element in  $\ell^\infty(\mathcal{F})$ . Then under Assumptions 3.1-3.6:*

$$\sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n^b) - \Psi_I^{\ell b}(\mathbb{P}_n)) | \{W_i\}_{i=1}^n \rightsquigarrow (\Psi_I^{\ell b})'_P(\mathbb{G}_P),$$

$$\sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n^b) - \Psi_I^{ub}(\mathbb{P}_n)) | \{W_i\}_{i=1}^n \rightsquigarrow (\Psi_I^{ub})'_P(\mathbb{G}_P).$$

A confidence set for the true parameter  $\psi_0$  can then be constructed using the quantiles of the bootstrapped distributions of (3.24) and (3.25). In particular, the confidence set  $C_n^\psi(1 - \alpha)$  with asymptotic coverage probability of  $1 - \alpha$  can be constructed as:

$$C_n^\psi(1 - \alpha) := \left[ \Psi_I^{\ell b}(\mathbb{P}_n) - \frac{\hat{\Psi}_\alpha^{\ell b}}{\sqrt{n}}, \Psi_I^{ub}(\mathbb{P}_n) + \frac{\hat{\Psi}_\alpha^{ub}}{\sqrt{n}} \right], \quad (3.26)$$



where the pair  $(\hat{\Psi}_\alpha^{\ell b}, \hat{\Psi}_\alpha^{ub})$  minimize the length of the confidence set  $C_n^\psi(1 - \alpha)$  subject to the constraints:

$$P_n^b \left( \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n^b) - \Psi_I^{\ell b}(\mathbb{P}_n)) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n^b) - \Psi_I^{ub}(\mathbb{P}_n)) + \sqrt{n}\Delta(\mathbb{P}_n) \right) \geq 1 - \alpha, \quad (3.27)$$

$$P_n^b \left( \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n^b) - \Psi_I^{\ell b}(\mathbb{P}_n)) - \sqrt{n}\Delta(\mathbb{P}_n) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n^b) - \Psi_I^{ub}(\mathbb{P}_n)) \right) \geq 1 - \alpha, \quad (3.28)$$

where  $P_n^b$  is the bootstrap distribution and  $\Delta$  is the length of the identified set. Note that under Assumption 3.3, we will rule out cases where length of the identified set can be drifting towards zero and thus we avoid issues of uniformity that occur in this scenario (see Stoye (2009)).

The following result verifies that under our assumptions, the confidence set given in (3.26) is uniformly asymptotically valid:

**Theorem 3.2.** *Under Assumptions 3.1 - 3.6,*

$$\liminf_{n \rightarrow \infty} \inf_{\{(\psi, P): \psi \in \Psi_I(P), P \in \mathcal{P}\}} P(\psi \in C_n^\psi(1 - \alpha)) \geq 1 - \alpha, \quad (3.29)$$

where  $C_n^\psi(1 - \alpha)$  is as in (3.26).

The confidence set  $C_n^\psi(1 - \alpha)$  is both conceptually simple and easy to implement. Indeed, computing the confidence set amounts to bootstrapping the value functions for the optimization problems that define the endpoints of the set  $\Psi_I(\cdot)$ . Calibrating the critical values  $\hat{\Psi}_\alpha^{\ell b}$  and  $\hat{\Psi}_\alpha^{ub}$  is then easily done once the bootstrap distribution has been recovered. In other words, Assumptions 3.1 - 3.6 are sufficient for a researcher to “naively” bootstrap the value functions of a linear program in order to construct a uniformly valid confidence set for a linear functional of interest.

## 4 Conclusion

This paper proposes a simple procedure for constructing confidence intervals for functionals of a partially identified parameter vector. The procedure approximates the distribution of the upper and lower bounds of the identified set for the functional of interest through a simple bootstrap procedure. In particular, we show that if the problem is sufficiently regular, a “naive” bootstrap procedure can be used, where the researcher (essentially) repeatedly solves a linear program, and computes confidence sets by taking appropriate quantiles of the bootstrap distribution of the value functions. Uniform validity of this naive procedure is proven by making connections to results in the Operations Research literature on stochastic programming, and in particular by appealing to the notion of uniform Hadamard directional differentiability. The most important condition for the validity of our procedure is found to be the existence and uniqueness of optimal solutions and Lagrange multipliers. In future drafts we plan to replace our high-level conditions with primitive conditions that are more easily verified by the applied researcher.



## Appendix A Proofs

Throughout this appendix we use the following notation: if  $X_n, X$  are maps in a metric space  $(\mathbb{D}, d)$  then:

- $X_n = o_{\mathcal{P}}(a_n)$  is used to denote uniform (over  $\mathcal{P}$ ) convergence in probability of the random element  $|X_n/a_n|$  to 0; i.e.  $\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} P_P^*(|X_n/a_n| > \varepsilon) = 0$  for every  $\varepsilon > 0$ ,
- $X_n = O_{\mathcal{P}}(a_n)$  is used to denote uniform (over  $\mathcal{P}$ ) stochastic boundedness of the random element  $|X_n/a_n|$ ; i.e. the fact that for any  $\varepsilon > 0$  there exists a finite  $M$  and an  $N$  such that  $\sup_{P \in \mathcal{P}} P_P^*(|X_n/a_n| > M) < \varepsilon$  for all  $n \geq N$ .

We will also rely on the following facts which are not proven here, but for which references are provided.

**Fact A.1.** *Suppose that  $\{P_n \in \mathcal{P}_+\}_{n=1}^\infty \rightarrow P \in \mathcal{P}$ . Under Assumption 3.1, 3.2 and 3.3 there exists an  $N$  such that for all  $n \geq N$  strong duality holds for  $P_n \in \mathcal{P}_+$ ; that is, if  $\mathcal{L}(\theta, \lambda)(P_n)$  is the Lagrangian at probability measure  $P$ , then*

$$\Psi_I^{\ell b}(P_n) = \inf_{\theta \in \Theta} \sup_{\lambda \geq 0} \mathcal{L}(\theta, \lambda)(P_n) = \sup_{\lambda \geq 0} \inf_{\theta \in \Theta} \mathcal{L}(\theta, \lambda)(P_n),$$

and

$$\Psi_I^{ub}(P_n) = \sup_{\theta \in \Theta} \inf_{\lambda \leq 0} \mathcal{L}(\theta, \lambda)(P_n) = \inf_{\lambda \leq 0} \sup_{\theta \in \Theta} \mathcal{L}(\theta, \lambda)(P_n).$$

This result is called *Lagrangian Duality in convex optimization*; see, for example, [Borwein and Lewis \(2010\)](#) Theorem 4.3.7. This follows since any sequence  $\{P_n \in \mathcal{P}_+\}_{n=1}^\infty \rightarrow P \in \mathcal{P}$  must eventually lie in  $\mathcal{P}^\varepsilon$ , so that Assumption 3.3 holds in the tails of any such sequence.

Before the next fact, some definitions:

**Definition A.1** (Upper Hemicontinuity). *For metric spaces  $\mathcal{X}$  and  $\mathcal{Y}$ , a correspondence  $G : \mathcal{X} \rightarrow \mathcal{Y}$  is said to be upper hemicontinuous at  $x \in \mathcal{X}$  if for every open subset  $S$  of  $\mathcal{Y}$  with  $G(x) \subseteq S$  there exists a  $\delta > 0$  such that  $G(B_\delta(x)) \subseteq S$ .*

**Definition A.2** (Compact-Valued). *For metric spaces  $\mathcal{X}$  and  $\mathcal{Y}$ , a correspondence  $G : \mathcal{X} \rightarrow \mathcal{Y}$  is said to be compact-valued if  $G(x)$  is a compact subset of  $\mathcal{Y}$  for each  $x \in \mathcal{X}$ .*

**Definition A.3** (Closed at  $x$ ). *For metric spaces  $\mathcal{X}$  and  $\mathcal{Y}$ , a correspondence  $G : \mathcal{X} \rightarrow \mathcal{Y}$  is said to be closed at  $x$  if for any sequence  $\{x_n\}$  and  $\{y_n\}$  with  $x_n \rightarrow x$  and  $y_n \rightarrow y$  we have that  $y \in G(x)$  whenever  $y_n \in G(x_n)$  for all  $n$ .*

**Fact A.2** (Proposition E.2 in [Ok \(2007\)](#)). *Let  $\mathcal{X}$  and  $\mathcal{Y}$  be two metric spaces and  $\Gamma : \mathcal{X} \rightarrow \mathcal{Y}$  a correspondence. If  $\Gamma$  is compact-valued and upper hemicontinuous at  $x \in \mathcal{X}$ , then for any sequence  $\{x_m\}_{m=1}^\infty \subset \mathcal{X}$  and  $\{y_m\}_{m=1}^\infty \subset \mathcal{Y}$  with  $x_m \rightarrow x$  and  $y_m \in \Gamma(x_m)$  for each  $m$ , there exists a subsequence  $\{y_{m_k}\}_{k=1}^\infty$  such that  $y_{m_k} \rightarrow y \in \Gamma(x)$ .*



Finally,  $\Theta_{\ell b}(P)$  and  $\Theta_{ub}(P)$  denote the set of optimal solutions to (3.12) and (3.13), and  $\Lambda_{\ell b}(P)$  and  $\Lambda_{ub}(P)$  denote the set of Lagrange multipliers for (3.12) and (3.13).

## A.1 Proof of Results in Main Text

**Remark A.1.** *The following proof follows similar steps to the proof of Theorem 7.24 in Shapiro et al. (2009), which shows Hadamard directional differentiability. However, the proof here establishes that this property holds “uniformly” over  $\mathcal{P}$  under the assumptions of Theorem 3.1. The proof of uniformity follows namely from (i) the assumption  $h_n \rightarrow h$  in the sup norm (and thus uniformly) where  $h$  is an operator that is uniformly continuous with respect to  $\theta$  (the latter is provided by Lemma A.5), (ii) boundedness of the Lagrangian (given by Lemma A.3) and (iii) continuity of the optimal solutions and Lagrange multipliers (given by Lemma A.2).*

*Proof of Theorem 3.1.* We can focus on the lower bound, since the upper bound can be treated analogously. Consider any converging sequences  $P_n \rightarrow P \in \mathcal{P}$ ,  $\{h_n\} \subset \ell^\infty(\mathcal{F})$  and  $\{t_n\} \subset \mathbb{R}_+$  with  $t_n \downarrow 0$  and  $h_n \rightarrow h \in \mathcal{T}_P(\mathcal{F})$  such that  $P_n + t_n h_n \in \mathcal{P}_+$  for all  $n \geq 1$ . Recall the Lagrangian at a probability measure  $P_n$  is given by:

$$\mathcal{L}(\theta, \lambda)(P_n) := P_n \psi(W, \theta) + \sum_{j=1}^k \lambda_j P_n m_j(W, \theta), \quad (\text{A.1})$$

where  $\lambda := (\lambda_1, \dots, \lambda_k)' \in \overline{\mathbb{R}}_+^k$  is a vector of Lagrange multipliers. Denote the “unperturbed” and “perturbed” programs respectively as:

$$\Psi_I^{\ell b}(P_n) := \inf_{\theta \in \Theta} \sup_{\lambda \in \overline{\mathbb{R}}_+^k} \mathcal{L}(\theta, \lambda)(P_n), \quad (\text{A.2})$$

$$\Psi_I^{\ell b}(P_n + t_n h_n) := \inf_{\theta \in \Theta} \sup_{\lambda \in \overline{\mathbb{R}}_+^k} \mathcal{L}(\theta, \lambda)(P_n + t_n h_n), \quad (\text{A.3})$$

where  $P_n + t_n h_n$  is interpreted elementwise. By Fact A.1 we have by Lemma A.1 that there exists an  $N$  such that for all  $n \geq N$ :

$$\Psi_I^{\ell b}(P_n) = \inf_{\theta \in \Theta} \sup_{\lambda \in \overline{\mathbb{R}}_+^k} \mathcal{L}(\theta, \lambda)(P_n) = \sup_{\lambda \in \overline{\mathbb{R}}_+^k} \inf_{\theta \in \Theta} \mathcal{L}(\theta, \lambda)(P_n), \quad (\text{A.4})$$

$$\Psi_I^{\ell b}(P_n + t_n h_n) = \inf_{\theta \in \Theta} \sup_{\lambda \in \overline{\mathbb{R}}_+^k} \mathcal{L}(\theta, \lambda)(P_n + t_n h_n) = \sup_{\lambda \in \overline{\mathbb{R}}_+^k} \inf_{\theta \in \Theta} \mathcal{L}(\theta, \lambda)(P_n + t_n h_n). \quad (\text{A.5})$$

Lemma A.1 implies there exists an optimal  $\theta_{\ell b}^*(P_n)$  for each  $n \geq N$ . Now consider the sequence  $\{\theta_{\ell b}^*(P_n)\}_{n=1}^\infty$



with  $\theta_{\ell b}^*(P_n)$  optimal for each  $n \geq N$ , and conclude that for all  $n \geq N$ :

$$\Psi_I^{\ell b}(P_n) = \sup_{\lambda \in \bar{\mathbb{R}}_+^k} \mathcal{L}(\theta_{\ell b}^*(P_n), \lambda)(P_n), \quad (\text{A.6})$$

$$\Psi_I^{\ell b}(P_n + t_n h_n) \leq \sup_{\lambda \in \bar{\mathbb{R}}_+^k} \mathcal{L}(\theta_{\ell b}^*(P_n), \lambda)(P_n + t_n h_n), \quad (\text{A.7})$$

where (A.6) follows from strong duality, and (A.7) follows from the fact that  $\theta_{\ell b}^*(P_n)$  is optimal for program (A.2) but not necessarily program (A.3).

By Lemma A.1, we have that there exists a optimal vector of Lagrange multipliers in (A.7) for  $n \geq N$ . Let  $\{\lambda_{\ell b}^*(P_n + t_n h_n)\}_{n=1}^\infty$  be a sequence with  $\lambda_{\ell b}^*(P_n + t_n h_n)$  optimal for each  $n \geq N$ . For any such sequence, note from (A.6) and (A.7) we have for all  $n \geq N$ :

$$\Psi_I^{\ell b}(P_n) \geq \mathcal{L}(\theta_{\ell b}^*(P_n), \lambda_{\ell b}^*(P_n + t_n h_n))(P_n), \quad (\text{A.8})$$

$$\Psi_I^{\ell b}(P_n + t_n h_n) \leq \mathcal{L}(\theta_{\ell b}^*(P_n), \lambda_{\ell b}^*(P_n + t_n h_n))(P_n + t_n h_n). \quad (\text{A.9})$$

Finally, also note that since  $h_n \rightarrow h \in \mathcal{T}_P(\mathcal{F})$  by assumption, we have that:

$$h_n = h + o(1). \quad (\text{A.10})$$

Thus, for all  $n \geq N$ :

$$\begin{aligned} & \Psi_I^{\ell b}(P_n + t_n h_n) - \Psi_I^{\ell b}(P_n), \\ & \leq \mathcal{L}(\theta_{\ell b}^*(P_n), \lambda_{\ell b}^*(P_n + t_n h_n))(P_n + t_n h_n) - \mathcal{L}(\theta_{\ell b}^*(P_n), \lambda_{\ell b}^*(P_n + t_n h_n))(P_n), \quad (\text{from (A.8) and (A.9)}) \\ & = t_n h_{n,1} \psi(W, \theta_{\ell b}^*(P_n)) + \sum_{j=1}^k \lambda_{\ell b,j}^*(P_n + t_n h_n) t_n h_{n,j+1} m_j(W, \theta_{\ell b}^*(P_n)), \quad (\text{by (A.1)}) \\ & = t_n \left( h_1 \psi(W, \theta_{\ell b}^*(P_n)) + \sum_{j=1}^k \lambda_{\ell b,j}^*(P_n + t_n h_n) h_{j+1} m_j(W, \theta_{\ell b}^*(P_n)) \right) + o(t_n), \quad (\text{by (A.10)}) \end{aligned}$$

where the final line follows from uniform boundedness of the Lagrangian from Lemma A.3(ii). Thus for any sequence  $\{\theta_{\ell b}^*(P_n)\}$ :

$$\limsup_{n \rightarrow \infty} \frac{\Psi_I^{\ell b}(P_n + t_n h_n) - \Psi_I^{\ell b}(P_n)}{t_n}$$



$$\begin{aligned}
&\leq \limsup_{n \rightarrow \infty} \left[ h_1 \psi(W, \theta_{\ell b}^*(P_n)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P_n + t_n h_n) h_{j+1} m_j(W, \theta_{\ell b}^*(P_n)) \right], \\
&= h_1 \psi(W, \theta_{\ell b}^*(P)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P) h_{j+1} m_j(W, \theta_{\ell b}^*(P)). \tag{A.11}
\end{aligned}$$

The last line follows by convergence of  $P_n \rightarrow P \in \mathcal{P}$  and  $t_n \downarrow 0$ , by uniform continuity of  $h$  with respect to  $\theta$  from Lemma A.5, and by convergence of the optimal solutions to a unique optimal solution (by Assumptions 3.2 and 3.3). This latter fact follows from continuity of the optimal solutions and optimal Lagrange multipliers, which follows from Lemma A.2.

For the reverse inequality, recall the “unperturbed” and “perturbed” problems given in (A.2) and (A.3) respectively. By Lemma A.1 the set of optimal solutions to program (A.3) is nonempty for all  $n \geq N$ . Thus, let  $\theta_{\ell b}^*(P_n + t_n h_n)$  be a sequence of optimal solutions to program (A.3). Furthermore, by Lemma A.1, the set of optimal Lagrange multipliers to program (A.2) is nonempty for all  $n \geq N$ . Now note for any  $\lambda_{\ell b}^*(P_n)$  we have:

$$\Psi_I^{\ell b}(P_n) \leq \mathcal{L}(\theta_{\ell b}^*(P_n + t_n h_n), \lambda_{\ell b}^*(P_n))(P_n), \tag{A.12}$$

$$\Psi_I^{\ell b}(P_n + t_n h_n) \geq \mathcal{L}(\theta_{\ell b}^*(P_n + t_n h_n), \lambda_{\ell b}^*(P_n))(P_n + t_n h_n). \tag{A.13}$$

It follows that for  $n \geq N$ :

$$\begin{aligned}
&\Psi_I^{\ell b}(P_n + t_n h_n) - \Psi_I^{\ell b}(P_n), \\
&\geq \mathcal{L}(\theta_{\ell b}^*(P_n + t_n h_n), \lambda_{\ell b}^*(P_n))(P_n + t_n h_n) - \mathcal{L}(\theta_{\ell b}^*(P_n + t_n h_n), \lambda_{\ell b}^*(P_n))(P_n), \quad (\text{by (A.12) and (A.13)}) \\
&= t_n h_{n,1} \psi(W, \theta_{\ell b}^*(P_n + t_n h_n)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P_n) t_n h_{n, j+1} m_j(W, \theta_{\ell b}^*(P_n + t_n h_n)), \quad (\text{by (A.1)}) \\
&= t_n \left( h_1 \psi(W, \theta_{\ell b}^*(P_n + t_n h_n)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P_n) h_{j+1} m_j(W, \theta_{\ell b}^*(P_n + t_n h_n)) \right) + o(t_n), \quad (\text{by (A.10)})
\end{aligned}$$

where the final line follows from uniform boundedness of the Lagrangian from Lemma A.3(ii). Thus,

$$\liminf_{n \rightarrow \infty} \frac{\Psi_I^{\ell b}(P_n + t_n h_n) - \Psi_I^{\ell b}(P_n)}{t_n}$$



$$\begin{aligned}
&\geq \liminf_{n \rightarrow \infty} \left[ h_1 \psi(W, \theta_{\ell b}^*(P_n + t_n h_n)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P_n) h_{j+1} m_j(W, \theta_{\ell b}^*(P_n + t_n h_n)) \right], \\
&= h_1 \psi(W, \theta_{\ell b}^*(P)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P) h_{j+1} m_j(W, \theta_{\ell b}^*(P)). \tag{A.14}
\end{aligned}$$

The last line follows by convergence of  $P_n \rightarrow P \in \mathcal{P}$  and  $t_n \downarrow 0$ , by uniform continuity of  $h$  with respect to  $\theta$  from Lemma A.5, and by convergence of the optimal solutions to a unique optimal solution (by Assumptions 3.2 and 3.3). This latter fact follows from continuity of the optimal solutions and optimal Lagrange multipliers, which follows from Lemma A.2.

Finally, combining inequalities we obtain:

$$\lim_{n \rightarrow \infty} \frac{\Psi_I^{\ell b}(P_n + t_n h_n) - \Psi_I^{\ell b}(P_n)}{t_n} = h_1 \psi(W, \theta_{\ell b}^*(P)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P) h_{j+1} m_j(W, \theta_{\ell b}^*(P)). \tag{A.15}$$

This completes the proof. ■

*Proof of Lemma 3.1.* Let  $\mathbb{G}_{n, P_n} = \sqrt{n}(\mathbb{P}_n - P_n)$ . By Lemma D.1(2) in Bugni et al. (2015) we have that, under Assumptions 3.1 and 3.5,  $\mathcal{F}$  is Donsker and pre-Gaussian, both uniformly over  $\mathcal{P}$ . By Theorem 2.8.7 in Van Der Vaart and Wellner (1996), we have that Assumption 3.1 and 3.5 imply that  $\mathbb{G}_{n, P_n} \rightsquigarrow \mathbb{G}_P$  in  $\ell^\infty(\mathcal{F})$ , which is a tight Gaussian process with sample paths that are almost all uniformly continuous. Let  $\tilde{\mathbb{G}}_P$  be a version of  $\mathbb{G}_P$  with all sample paths uniformly continuous. Let  $\mathbb{D} = \ell^\infty(\mathcal{F})$ ,  $\mathbb{D}_0 = \mathcal{T}_P(\mathcal{F})$ ,  $\mathbb{E} = \mathbb{R}$ , and define:

$$\mathbb{D}_n = \{h : P_n + n^{-1/2}h \in \mathcal{P}_+\},$$

Then  $\mathbb{D}_n \subset \mathbb{D}$  and  $\mathbb{D}_0 \subset \mathbb{D}$ . Now consider the maps  $g_n : \mathbb{D}_n \rightarrow \mathbb{E}$  and  $g : \mathbb{D}_0 \rightarrow \mathbb{E}$  defined as:

$$g_n(h_n) := \sqrt{n} \left\{ \Psi_I^{\ell b}(P_n + n^{-1/2}h_n) - \Psi_I^{\ell b}(P_n) \right\}, \quad h_n \in \mathbb{D}_n \tag{A.16}$$

$$g(h) := (\Psi_I^{\ell b})'_P(h), \quad h \in \mathbb{D} \tag{A.17}$$

By Theorem 3.1, if  $h_n \rightarrow h$  with  $h_n \in \mathbb{D}_n$  for every  $n$  and  $h \in \mathbb{D}_0$ , then  $g_n(h_n) \rightarrow g(h)$ , where  $g : \mathbb{D}_0 \rightarrow \mathbb{D}$ . Now note that  $\mathbb{G}_{n, P_n} \in \mathbb{D}_n$ . Using the fact that  $\tilde{\mathbb{G}}_P$  is a tight (and thus separable) Borel element with values in  $\mathbb{D}_0$ , combined with the extended continuous mapping theorem (Theorem 1.11.1 in Van Der Vaart and Wellner (1996)), we conclude that:

$$\sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P_n)) \rightsquigarrow (\Psi_I^{\ell b})'_P(\tilde{\mathbb{G}}_P),$$

as desired. An identical proof can be completed for the upper bound. Thus, this completes the proof. ■



*Proof of Lemma 3.2.* Let  $\tilde{\mathbb{G}}_P$  be a version of  $\mathbb{G}_P$  with all sample paths uniformly continuous. Let  $\mathbb{D} = \ell^\infty(\mathcal{F})$ ,  $\mathbb{D}_0 = \mathcal{T}_P(\mathcal{F})$ ,  $\mathbb{E} = \mathbb{R}$ , and define:

$$\mathbb{D}_n = \{h \in \mathbb{D} : \mathbb{P}_n + n^{-1/2}h \in \mathcal{P}_+\},$$

Then  $\mathbb{D}_n \subset \mathbb{D}$  and  $\mathbb{D}_0 \subset \mathbb{D}$ . Now consider the maps  $g_n : \mathbb{D}_n \rightarrow \mathbb{E}$  and  $g : \mathbb{D}_0 \rightarrow \mathbb{E}$  defined as:

$$g_n(h_n) := \sqrt{n} \left( \Psi_I^{\ell b} \left( \mathbb{P}_n + n^{-1/2}h_n \right) - \Psi_I^{\ell b}(\mathbb{P}_n) \right), \quad h_n \in \mathbb{D}_n \quad (\text{A.18})$$

$$g(h) := (\Psi_I^{\ell b})'_P(h), \quad h \in \mathbb{D} \quad (\text{A.19})$$

By Theorem 3.1, if  $h_n \rightarrow h$  with  $h_n \in \mathbb{D}_n$  for every  $n$  and  $h \in \mathbb{D}_0$ , then  $g_n(h_n) \rightarrow g(h)$ , where  $g : \mathbb{D}_0 \rightarrow \mathbb{D}$ . Now note that  $\mathbb{G}_n^b \in \mathbb{D}_n$ , and by assumption  $\mathbb{G}_n^b | \{W_i\}_{i=1}^n \rightsquigarrow \tilde{\mathbb{G}}_P$  uniformly over  $\mathcal{P}$ . Using the fact that  $\tilde{\mathbb{G}}_P$  is a tight (and thus separable) Borel element with values in  $\mathbb{D}_0$ , combined with the extended continuous mapping theorem (Theorem 1.11.1 in [Van Der Vaart and Wellner \(1996\)](#)), we conclude that:

$$\sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n^b) - \Psi_I^{\ell b}(\mathbb{P}_n)) | \{W_i\}_{i=1}^n \rightsquigarrow (\Psi_I^{\ell b})'_P(\tilde{\mathbb{G}}_P),$$

as desired. An identical proof can be completed for the upper bound. Thus, this completes the proof. ■

*Proof of Theorem 3.2.* By definition there exists a sequence  $(\psi_n, P_n) \in \{(\psi, P) : \psi \in \Psi_I(P), P \in \mathcal{P}\}$  satisfying:

$$\liminf_{n \rightarrow \infty} \inf_{\{(\psi, P) : \psi \in \Psi_I(P), P \in \mathcal{P}\}} P(\psi \in C_n^\psi(1 - \alpha)) = \liminf_{n \rightarrow \infty} P_n(\psi_n \in C_n^\psi(1 - \alpha)),$$

where  $\{\psi_n\}$  is a sequence with  $\psi_n \in [\Psi_I^{\ell b}(P_n), \Psi_I^{ub}(P_n)]$  for each  $n$ . For such a sequence, there exists a convergent subsequence indexed by  $n'$  such that:

$$\liminf_{n \rightarrow \infty} P_n(\psi_n \in C_n^\psi(1 - \alpha)) = \lim_{n' \rightarrow \infty} P_{n'}(\psi_{n'} \in C_{n'}^\psi(1 - \alpha)).$$

For the remainder of the proof we will argue along this subsequence, and abusing notation we will refer to this subsequence by  $n$  rather than  $n'$ . Since by construction we have  $\psi_n \in [\Psi_I^{\ell b}(P_n), \Psi_I^{ub}(P_n)]$  for each  $n$ , it suffices to establish that:

$$\lim_{n \rightarrow \infty} P_n(\Psi_I^{\ell b}(P_n) \in C_n^\psi(1 - \alpha)) \geq 1 - \alpha, \quad (\text{A.20})$$

$$\lim_{n \rightarrow \infty} P_n(\Psi_I^{ub}(P_n) \in C_n^\psi(1 - \alpha)) \geq 1 - \alpha. \quad (\text{A.21})$$

We can focus on (A.20) since (A.21) can be treated analogously. We have:

$$P_n(\Psi_I^{\ell b}(P_n) \in C_n^\psi(1 - \alpha))$$



$$\begin{aligned}
&= P_n \left( \Psi_I^{\ell b}(\mathbb{P}_n) - \hat{\Psi}_\alpha^{\ell b} / \sqrt{n} \leq \Psi_I^{\ell b}(P_n) \leq \Psi_I^{ub}(\mathbb{P}_n) + \hat{\Psi}_\alpha^{ub} / \sqrt{n} \right), \\
&= P_n \left( \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P_n)) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n) - \Psi_I^{\ell b}(P_n)) \right), \\
&= P_n \left( \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P_n)) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n) - \Psi_I^{ub}(P_n)) + \sqrt{n}\Delta(P_n) \right). \tag{A.22}
\end{aligned}$$

Decomposing this probability we have:

$$\begin{aligned}
&P_n \left( \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P_n)) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n) - \Psi_I^{ub}(P_n)) + \sqrt{n}\Delta(P_n) \right) \\
&= P_n^b \left( \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n^b) - \Psi_I^{\ell b}(\mathbb{P}_n)) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n^b) - \Psi_I^{ub}(\mathbb{P}_n)) + \sqrt{n}\Delta(\mathbb{P}_n) \right) \\
&+ \left[ P_n \left( \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P_n)) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n) - \Psi_I^{ub}(P_n)) + \sqrt{n}\Delta(P_n) \right) \right. \\
&\quad \left. - P_n \left( (\Psi_I^{\ell b})'_P(\mathbb{G}_P) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq (\Psi_I^{ub})'_P(\mathbb{G}_P) + \sqrt{n}\Delta(P_n) \right) \right] \tag{A.23}
\end{aligned}$$

$$\begin{aligned}
&+ \left[ P_n \left( (\Psi_I^{\ell b})'_P(\mathbb{G}_P) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq (\Psi_I^{ub})'_P(\mathbb{G}_P) + \sqrt{n}\Delta(P_n) \right) \right. \\
&\quad \left. - P_n \left( (\Psi_I^{\ell b})'_P(\mathbb{G}_P) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq (\Psi_I^{ub})'_P(\mathbb{G}_P) + \sqrt{n}\Delta(\mathbb{P}_n) \right) \right] \tag{A.24}
\end{aligned}$$

$$\begin{aligned}
&+ \left[ P_n \left( (\Psi_I^{\ell b})'_P(\mathbb{G}_P) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq (\Psi_I^{ub})'_P(\mathbb{G}_P) + \sqrt{n}\Delta(\mathbb{P}_n) \right) \right. \\
&\quad \left. - P_n^b \left( \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n^b) - \Psi_I^{\ell b}(\mathbb{P}_n)) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n^b) - \Psi_I^{ub}(\mathbb{P}_n)) + \sqrt{n}\Delta(\mathbb{P}_n) \right) \right]. \tag{A.25}
\end{aligned}$$

Note by construction we will have for all  $n$ :

$$P_n^b \left( \sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n^b) - \Psi_I^{\ell b}(\mathbb{P}_n)) \leq \hat{\Psi}_\alpha^{\ell b}, -\hat{\Psi}_\alpha^{ub} \leq \sqrt{n}(\Psi_I^{ub}(\mathbb{P}_n^b) - \Psi_I^{ub}(\mathbb{P}_n)) + \sqrt{n}\Delta(\mathbb{P}_n) \right) \geq 1 - \alpha,$$

so that it suffices to show that the terms (A.23), (A.24) and (A.25) converge to non-negative values. First consider (A.23). By Lemma 3.1 we have that:

$$\sqrt{n}(\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P_n)) \rightsquigarrow (\Psi_I^{\ell b})'_P(\mathbb{G}_P). \tag{A.26}$$

Assumptions 3.1 and 3.5 ensure the objective function (when it is a non-trivial function of  $w \in \mathcal{W}$ ) and moment functions are uniformly Donsker over  $\mathcal{P}$ . Thus, when combined with uniform boundedness of the Lagrange multipliers from Lemma A.3, this ensures continuity of the distribution of  $(\Psi_I^{\ell b})'_P(\mathbb{G}_P)$  at its  $\alpha$



quantile and  $(\Psi_I^{ub})'_P(\mathbb{G}_P)$  at its  $1 - \alpha$  quantile. Thus, convergence of (A.23) to zero follows from (A.26), Theorem 1.3.4(vi) in Van Der Vaart and Wellner (1996), and continuity of the distributions of  $(\Psi_I^{\ell b})'_P(\mathbb{G}_P)$  and  $(\Psi_I^{ub})'_P(\mathbb{G}_P)$ .

Next, note from Lemma A.4 that  $\Psi_I(\mathbb{P}_n)$  is Hausdorff consistent for  $\Psi_I(P)$  over  $\{P_n \in \mathcal{P}\}_{n=1}^\infty$ , which implies consistency of  $\Delta(\mathbb{P}_n)$  for  $\Delta(P)$ . Also note that Assumptions 3.2 and 3.3 imply that  $\Delta(P) > 0$  for all  $P \in \mathcal{P}$ , so that  $\sqrt{n}\Delta(P_n) \rightarrow \infty$ . However,  $\Delta(\mathbb{P}_n) = \Delta(P) + o_{P_n}(1)$  by Lemma A.4, so that (A.24) converges to zero, as desired.

Finally, (A.25) converges to zero w.p.a. 1, which follows from bootstrap consistency over the sequence  $\{P_n \in \mathcal{P}\}_{n=1}^\infty$  from Lemma 3.2, and again from continuity of the distributions of  $(\Psi_I^{\ell b})'_P(\mathbb{G}_P)$  and  $(\Psi_I^{ub})'_P(\mathbb{G}_P)$  described above. ■

## A.2 Proofs of Additional Results

**Lemma A.1.** *Under Assumptions 3.1 - 3.3,  $\Theta_I(P)$ ,  $\Theta_{\ell b}(P)$ ,  $\Theta_{ub}(P)$ ,  $\Lambda_{\ell b}(P)$  and  $\Lambda_{ub}(P)$  are nonempty for every  $P \in \mathcal{P}$ . Furthermore, if  $\{P_n\}_{n=1}^\infty$  is any sequence converging to  $P \in \mathcal{P}$ , then there exists an  $N$  such that  $\Theta_I(P_n)$ ,  $\Theta_{\ell b}(P_n)$ ,  $\Theta_{ub}(P_n)$ ,  $\Lambda_{\ell b}(P_n)$  and  $\Lambda_{ub}(P_n)$  are nonempty for all  $n \geq N$ .*

*Proof.* Nonemptiness of  $\Theta_I(P)$ ,  $\Theta_{\ell b}(P)$ ,  $\Theta_{ub}(P)$  follows from Assumption 3.1. Nonemptiness of  $\Lambda_{\ell b}(P)$  and  $\Lambda_{ub}(P)$  follows from 3.3(ii) and Wachsmuth (2013) Theorems 1 and 2.

The second claim can be established from 3.3(ii) and Wachsmuth (2013) Theorems 1 and 2 if we can show there exists an  $N$  such that  $\Theta_I(P_n)$  is nonempty for all  $n \geq N$ . This follows immediately from Assumption 3.3(i) and the definition of convergence of probability measures used in this paper. ■

**Lemma A.2.** *Under Assumptions 3.1, 3.2, and 3.3, we have:*

- (a)  $\theta_{\ell b}^*(P)$  and  $\theta_{ub}^*(P)$  are continuous at any  $P \in \mathcal{P}$ .
- (b)  $\lambda_{\ell b}^*(P)$  and  $\lambda_{ub}^*(P)$  are continuous at any  $P \in \mathcal{P}$ .
- (c)  $\Psi_I^{\ell b}(P)$  and  $\Psi_I^{ub}(P)$  are continuous at any  $P \in \mathcal{P}$ .

*Proof.* Let  $\|x - y\|_{\mathbb{R}_+^k} = \|\arctan(x) - \arctan(y)\|$ , where  $\|\cdot\|$  is the euclidean norm. Note that  $(\Theta, \|\cdot\|)$ ,  $(\mathbb{R}_+^k, \|\cdot\|_{\mathbb{R}_+^k})$  and  $(\mathcal{P}_+, \|\cdot\|_G)$  are all metric spaces. Focus first on the lower bound program in (3.12). Take any  $P \in \mathcal{P}$ . Define:

$$\Theta_I(P_n) := \{\theta \in \Theta : P_n m_j(W, \theta) = 0, j = 1, \dots, r_1, P_n m_j(W, \theta) \leq 0, j = r_1 + 1, \dots, r_1 + r_2\}.$$

By Lemma A.1, for any sequence  $P_n \rightarrow P \in \mathcal{P}$  (possibly with  $P_n \in \mathcal{P}_+$ ) we have that there exists an  $N$  such that  $\Theta_I(P_n)$  is nonempty for all  $n \geq N$ . By Assumption 3.1(i),  $\Theta_I(\cdot)$  is also a compact-valued correspondence



for all  $n \geq N$ . Recall the Lagrangian for problem (3.12):

$$\mathcal{L}(\theta, \lambda)(P) := P\psi(W, \theta) + \sum_{j=1}^k \lambda_j Pm_j(W, \theta).$$

By Assumption 3.2,  $\mathcal{L}(\theta, \lambda)(P)$  is continuous in  $(\theta, \lambda, P)$ . Define:

$$\Theta^*(\lambda, P) := \arg \min\{\mathcal{L}(\theta, \lambda, P) : \theta \in \Theta_I(P)\},$$

$$\mathcal{L}_\theta^*(\lambda, P) := \min\{\mathcal{L}(\theta, \lambda, P) : \theta \in \Theta_I(P)\}.$$

Note that  $\Theta^*(\lambda, P) \neq \emptyset$  and  $\mathcal{L}_\theta^*(\lambda, P) > -\infty$  by Lemma A.1. By the Theorem of the Maximum (Ok (2007), p. 306) we have that  $\Theta^*(\lambda, P)$  is compact-valued, upper-hemicontinuous, and closed, and the profiled-Lagrangian  $\mathcal{L}_\theta^*(\lambda, P)$  is continuous in  $(\lambda, P)$ . Now define:

$$\Lambda_\theta^*(P) := \arg \max\{\mathcal{L}_\theta^*(\lambda, P) : \lambda \in \overline{\mathbb{R}}_+^k\},$$

$$\mathcal{L}_{\theta, \lambda}^*(P) := \max\{\mathcal{L}_\theta^*(\lambda, P) : \lambda \in \overline{\mathbb{R}}_+^k\}.$$

Note that  $\Lambda_\theta^*(P) \neq \emptyset$  and  $\mathcal{L}_{\theta, \lambda}^*(P) < \infty$  by Lemma A.1. Applying the Theorem of the Maximum again, we have that  $\Lambda_\theta^*(P)$  is compact-valued, upper-hemicontinuous, and closed, and the profiled-Lagrangian  $\mathcal{L}_{\theta, \lambda}^*(P)$  is continuous in  $P$ . Similarly, define:

$$\Lambda^*(\theta, P) := \arg \max\{\mathcal{L}(\theta, \lambda, P) : \lambda \in \overline{\mathbb{R}}_+^k\},$$

$$\mathcal{L}_\lambda^*(\theta, P) := \max\{\mathcal{L}(\theta, \lambda, P) : \lambda \in \overline{\mathbb{R}}_+^k\},$$

$$\Theta_\lambda^*(P) := \arg \min\{\mathcal{L}_\lambda^*(\theta, P) : \theta \in \Theta_I(P)\},$$

$$\mathcal{L}_{\lambda, \theta}^*(P) := \min\{\mathcal{L}_\lambda^*(\theta, P) : \theta \in \Theta_I(P)\}.$$

I.e. reverse the order of profiling of the Lagrangian with respect to  $\lambda$  and  $\theta$ . Note this can be done by strong duality (Fact A.1) without affecting the optimal solution sets. Applying Lemma A.1 as above, we conclude that  $\Lambda^*(\theta, P) \neq \emptyset$ ,  $\mathcal{L}_\lambda^*(\theta, P) > -\infty$ ,  $\Theta_\lambda^*(P) \neq \emptyset$ , and  $\mathcal{L}_{\lambda, \theta}^*(P) < \infty$ . Applying the Theorem of the Maximum sequentially as above, we conclude that  $\Theta_\lambda^*(P)$  is compact-valued, upper-hemicontinuous, and closed, and the profiled-Lagrangian  $\mathcal{L}_{\lambda, \theta}^*(P)$  is continuous in  $P$ . Finally, by strong duality (Fact A.1) we conclude  $\Psi_I^{\ell b}(P) = \mathcal{L}_{\theta, \lambda}^*(P) = \mathcal{L}_{\lambda, \theta}^*(P)$ ,  $\Lambda_{\ell b}(P) = \Lambda_\theta^*(P)$ , and  $\Theta_{\ell b}(P) = \Theta_\lambda^*(P)$ . By Assumption 3.3, all of these sets are singletons. Repeating the exercise for the upper bound program, the proof is complete. ■



**Lemma A.3.** Under Assumptions 3.1, 3.2, 3.3 and 3.4,

(i) There exists constants  $L_{\ell b}, L_{ub} < \infty$  such that:

$$\sup_{P \in \mathcal{P}^\varepsilon} \|\lambda_{\ell b}^*(P)\| \leq L_{\ell b}, \quad (\text{A.27})$$

$$\sup_{P \in \mathcal{P}^\varepsilon} \|\lambda_{ub}^*(P)\| \leq L_{ub}. \quad (\text{A.28})$$

I.e. the Lagrange multipliers are uniformly bounded over  $\mathcal{P}$  in both the lower bound and upper bound programs.

(ii) There exist constants  $C_{\ell b}, C_{ub} < \infty$  such that:

$$\sup_{P \in \mathcal{P}^\varepsilon} \left| \psi(W, \theta_{\ell b}^*(P)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P) m_j(W, \theta_{\ell b}^*(P)) \right| \leq C_{\ell b}, \quad (\text{A.29})$$

$$\sup_{P \in \mathcal{P}^\varepsilon} \left| \psi(W, \theta_{ub}^*(P)) + \sum_{j=1}^k \lambda_{ub, j}^*(P) m_j(W, \theta_{ub}^*(P)) \right| \leq C_{ub}. \quad (\text{A.30})$$

I.e. the Lagrangian is uniformly bounded in both the lower bound and upper bound programs.

*Proof.* Part (i): We will focus on (A.27) since (A.28) follows analogously. By Assumption 3.2 and 3.3, we have the KKT conditions:

$$\nabla_\theta P \mathbf{m}(W, \theta_{\ell b}^*(P))^T \boldsymbol{\lambda}_{\ell b}^*(P) = -\nabla_\theta P \psi(W, \theta_{\ell b}^*(P))^T$$

where  $\nabla_\theta P \mathbf{m}(W, \theta_{\ell b}^*(P))$  is the  $(r_1 + r_2) \times d_\theta$  Jacobian matrix for the moment conditions. Let  $B$  denote the index set for the active constraints. Now let  $\boldsymbol{\lambda}_B(P)$  denote the subvector of  $\boldsymbol{\lambda}_{\ell b}^*(P)$  corresponding to the active constraints. Then clearly:

$$\nabla_\theta P \mathbf{m}(W, \theta_{\ell b}^*(P))^T \boldsymbol{\lambda}_{\ell b}^*(P) = G(\theta_{\ell b}^*(P), P)^T \boldsymbol{\lambda}_B(P) = -\nabla_\theta P \psi(W, \theta_{\ell b}^*(P))^T$$

Pre-multiplying by  $G(\theta_{\ell b}^*(P), P)$  and inverting we obtain:

$$\boldsymbol{\lambda}_B(P) = -[G(\theta_{\ell b}^*(P), P)G(\theta_{\ell b}^*(P), P)^T]^{-1} G(\theta_{\ell b}^*(P), P) \nabla_\theta P \psi(W, \theta_{\ell b}^*(P))^T$$

Denote:

$$A_1(P) := -[G(\theta_{\ell b}^*(P), P)G(\theta_{\ell b}^*(P), P)^T]^{-1} \quad (\text{A.31})$$

$$A_2(P) := G(\theta_{\ell b}^*(P), P) \nabla_\theta P \psi(W, \theta_{\ell b}^*(P))^T \quad (\text{A.32})$$



Now note:

$$\begin{aligned} \sup_{P \in \mathcal{P}^\varepsilon} \|A_1(P)\|_2 &\leq \frac{1}{\sqrt{\kappa}}, & (\text{by Assumption 3.3}), \\ \sup_{P \in \mathcal{P}^\varepsilon} \|A_2(P)\| &\leq \sqrt{\kappa} \cdot L_{\ell b}, & (\text{by Assumption 3.4}), \end{aligned}$$

where  $L_{\ell b} < \infty$  is some constant. Then:

$$\begin{aligned} \sup_{P \in \mathcal{P}^\varepsilon} \|\lambda_B(P)\| &= \sup_{P \in \mathcal{P}^\varepsilon} \|A_1(P)A_2(P)\| \\ &= \sup_{P \in \mathcal{P}^\varepsilon} \|A_1(P)A_2(P)\|_F \\ &\leq \sup_{P \in \mathcal{P}^\varepsilon} \|A_1(P)\|_2 \|A_2(P)\|_F \\ &\leq \left( \sup_{P \in \mathcal{P}^\varepsilon} \|A_1(P)\|_2 \right) \left( \sup_{P \in \mathcal{P}^\varepsilon} \|A_2(P)\|_F \right) \\ &\leq L_{\ell b} \end{aligned}$$

where  $\|\cdot\|_2$  denotes the 2-norm and  $\|\cdot\|_F$  denotes the Frobenius norm. This upper bound is also sufficient for the arctan norm, and completes the proof of the first part.

Part (ii): We will focus on (A.29) since (A.30) follows analogously. By Assumption 3.1(v) there exists a function  $F(w)$  such that  $\sup_{\theta \in \Theta} \|f(w, \theta)\| \leq \|F(w)\|$  for every  $w \in \mathcal{W}$ , and such that  $F(w)$  is uniformly bounded. Let  $C_F < \infty$  be a positive constant satisfying  $\|F(w)\| \leq C_F$  for all  $w \in \mathcal{W}$ . Then:

$$\begin{aligned} \sup_{P \in \mathcal{P}^\varepsilon} \left| \psi(W, \theta_{\ell b}^*(P)) + \sum_{j=1}^k \lambda_{\ell b, j}^*(P) m_j(W, \theta_{\ell b}^*(P)) \right| &\leq \sup_{P \in \mathcal{P}^\varepsilon} \|F(w)\| \cdot \|\lambda_{\ell b}^*(P)\| \\ &\leq C_F L_{\ell b}, \end{aligned}$$

where the first inequality follows from Cauchy-Schwarz, and the last inequality follows from part (i). Thus, taking  $C_{\ell b} = C_F L_{\ell b}$  the proof is complete.  $\blacksquare$

**Lemma A.4.** *Under Assumptions 3.1-3.5, we have that,*

$$(i) \quad d_H(\Theta_I(\mathbb{P}_n), \Theta_I(P)) = O_P(n^{-1/2}),$$

$$(ii) \quad d_H(\Psi_I(\mathbb{P}_n), \Psi_I(P)) = o_P(1),$$

$$(iii) \quad \text{For any } \varepsilon > 0,$$

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} P_P^* (\|\theta_{\ell b}^*(\mathbb{P}_n) - \theta_{\ell b}^*(P)\| > \varepsilon) = 0, \quad (\text{A.33})$$

and the analogous result for  $\theta_{ub}^*(\cdot)$ .

$$(iv) \quad \text{For any } \varepsilon > 0,$$

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} P_P^* (\|\lambda_{\ell b}^*(\mathbb{P}_n) - \lambda_{\ell b}^*(P)\|_{\mathbb{R}^k} > \varepsilon) = 0. \quad (\text{A.34})$$



and the analogous result for  $\lambda_{ub}^*(\cdot)$ .

*Proof of Lemma A.4.* Part (i): We follows closely the proof of Theorem 4.3(II) in [Kaido et al. \(2017\)](#). Define the set:

$$\Theta_I^\gamma(P) := \left\{ \theta \in \Theta : \max_{r_1+1 \leq j \leq r_1+r_2} Pm_j(W, \theta) \leq \gamma, \quad Pm_j(W, \theta) = 0, \quad j = 1, \dots, r_1 \right\},$$

for  $\gamma \in \mathbb{R}$ . First note that by Lemma D.1 in [Bugni et al. \(2015\)](#) Assumption 3.5 implies that  $\mathcal{F}$  is uniformly Donsker. In particular, we have that  $\|\mathbb{G}_{n,P}\|_{\mathcal{F}} = O_{\mathcal{P}}(1)$ . This implies:

$$\begin{aligned} \sup_{\theta \in \Theta_I^{-\varepsilon_n}(P)} \sqrt{n} \max_j |\mathbb{P}_n m_j(W, \theta)|_+ &\leq \sup_{\theta \in \Theta_I^{-\varepsilon_n}(P)} \sum_j \sqrt{n} |\mathbb{P}_n m_j(W, \theta)|_+, \\ &= \sup_{\theta \in \Theta_I^{-\varepsilon_n}(P)} \sum_j |\mathbb{G}_{n,P} m_j(W, \theta) + \sqrt{n} P m_j(W, \theta)|_+, \\ &\leq r_1 |O_{\mathcal{P}}(1)| + r_2 |O_{\mathcal{P}}(1) - \sqrt{n} \varepsilon_n|_+, \end{aligned}$$

from which we conclude that  $\Theta_I^{-\varepsilon_n}(P) \subseteq \Theta_I(\mathbb{P}_n)$  w.p.a. 1 for  $\varepsilon_n = O_{\mathcal{P}}(n^{-1/2})$ . Furthermore, by Assumption 3.5(iv) we can choose  $\delta(\varepsilon_n) > 0$  such that:

$$\begin{aligned} &\inf_{\theta \in \Theta \setminus \Theta_I^{\varepsilon_n}(P)} \sqrt{n} \max_j |\mathbb{P}_n m_j(W, \theta)|_+, \\ &= \inf_{\theta \in \Theta \setminus \Theta_I^{\varepsilon_n}(P)} \max_j |\mathbb{G}_{n,P} m_j(W, \theta) + \sqrt{n} P m_j(W, \theta)|_+, \\ &\geq \inf_{\theta \in \Theta \setminus \Theta_I^{\varepsilon_n}(P)} \frac{1}{J} \sum_j |\mathbb{G}_{n,P} m_j(W, \theta) + \sqrt{n} P m_j(W, \theta)|_+, \\ &\geq \inf_{\theta \in \Theta \setminus \Theta_I^{\varepsilon_n}(P)} \frac{1}{J} [(J-1) \cdot 0 + |O_{\mathcal{P}}(1) + \sqrt{n} C \min\{\delta(\varepsilon_n), d(\theta, \Theta_I(P))\}|_+], \\ &= \inf_{\theta \in \Theta \setminus \Theta_I^{\varepsilon_n}(P)} \frac{1}{J} |O_{\mathcal{P}}(1) + \sqrt{n} C \min\{\delta(\varepsilon_n), d(\theta, \Theta_I(P))\}|_+, \end{aligned}$$

from which we conclude that  $\Theta_I(\mathbb{P}_n) \cap (\Theta \setminus \Theta_I^{\varepsilon_n}(P)) = \emptyset$  w.p.a. 1 for  $\varepsilon_n = O_{\mathcal{P}}(n^{-1/2})$  (from the first line). Note that this concludes the proof of part (i).

Part (ii): It suffices to show consistency of the upper and lower bounds; i.e. that  $|\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P)| = o_{\mathcal{P}}(1)$  and that  $|\Psi_I^{ub}(\mathbb{P}_n) - \Psi_I^{ub}(P)| = o_{\mathcal{P}}(1)$ . We will focus on the lower bounds, since the upper bound proof is symmetric. First note that since  $\psi(W, \theta)$  is continuous with respect to  $\theta$  by Assumption 3.2, and that  $\Theta$  is compact by Assumption 3.1(i), we have that  $\psi(W, \theta)$  is uniformly continuous (w.r.t.  $\theta$ ) on  $\Theta$ . Thus, for every  $\varepsilon > 0$  there exists a  $\delta(\varepsilon) > 0$  such that  $|\mathbb{P}_n \psi(W, \theta) - \mathbb{P}_n \psi(W, \theta')| < \varepsilon$  whenever  $\|\theta - \theta'\| < \delta(\varepsilon)$ .

Now note that:

$$\begin{aligned} |\Psi_I^{\ell b}(\mathbb{P}_n) - \Psi_I^{\ell b}(P)| &= \left| \min_{\theta \in \Theta_I(\mathbb{P}_n)} \mathbb{P}_n \psi(W, \theta) - \min_{\theta \in \Theta_I(P)} P \psi(W, \theta) \right|, \\ &\leq \left| \min_{\theta \in \Theta_I(\mathbb{P}_n)} \mathbb{P}_n \psi(W, \theta) - \min_{\theta \in \Theta_I(P)} \mathbb{P}_n \psi(W, \theta) \right| + \left| \min_{\theta \in \Theta_I(P)} \mathbb{P}_n \psi(W, \theta) - \min_{\theta \in \Theta_I(P)} P \psi(W, \theta) \right|, \end{aligned}$$



$$\leq \sup_{\|\theta - \theta'\| \leq d_H(\Theta_I(\mathbb{P}_n), \Theta_I(P))} |\mathbb{P}_n \psi(W, \theta) - \mathbb{P}_n \psi(W, \theta')| + \sup_{\theta \in \Theta_I(P)} |\mathbb{P}_n \psi(W, \theta) - P \psi(W, \theta)|.$$

It suffices to show the two terms in the last line of the previous array converge to zero in probability uniformly. Note that by part (i) of this Lemma, we have  $d_H(\Theta_I(\mathbb{P}_n), \Theta_I(P)) = o_{\mathcal{P}}(1)$ . Thus by uniform continuity of  $\mathbb{P}_n \psi(W, \theta)$ :

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} P_P^* \left( \sup_{\|\theta - \theta'\| \leq d_H(\Theta_I(\mathbb{P}_n), \Theta_I(P))} |\mathbb{P}_n \psi(W, \theta) - \mathbb{P}_n \psi(W, \theta')| > \varepsilon \right) = 0.$$

Also, by the uniform Donsker property:

$$\sup_{\theta \in \Theta_I(P)} |\mathbb{P}_n \psi(W, \theta) - P \psi(W, \theta)| \leq \sup_{\theta \in \Theta} |\mathbb{P}_n \psi(W, \theta) - P \psi(W, \theta)| = o_{\mathcal{P}}(1).$$

This completes the proof.

Part (iii) + (iv): Using Lemma A.3, we can restrict  $\lambda$  to lie in the set  $\Lambda := \{\lambda : \|\lambda\| \leq \max\{L_{\ell b}, L_{ub}\}\}$ . Fix any  $\varepsilon, \eta > 0$ . By the uniform Donsker property we have:

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} \sup_{f \in \mathcal{F}} \|\mathbb{P}_n - P\|_{\mathcal{F}} = 0,$$

which implies the following inequalities hold w.p.a. 1:

$$\begin{aligned} \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(\mathbb{P}_n))(P) &< \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(\mathbb{P}_n))(\mathbb{P}_n) + \varepsilon/3, \\ \mathcal{L}(\theta_{\ell b}^*(P), \lambda_{\ell b}^*(\mathbb{P}_n))(\mathbb{P}_n) &< \mathcal{L}(\theta_{\ell b}^*(P), \lambda_{\ell b}^*(\mathbb{P}_n))(P) + \varepsilon/3, \\ \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(\mathbb{P}_n))(P) &> \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(\mathbb{P}_n))(\mathbb{P}_n) - \eta/3, \\ \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(P))(\mathbb{P}_n) &> \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(P))(P) - \eta/3. \end{aligned}$$

Furthermore, by optimality of  $\theta_{\ell b}^*(\mathbb{P}_n)$  and  $\lambda_{\ell b}^*(\mathbb{P}_n)$  we have:

$$\begin{aligned} \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(\mathbb{P}_n))(\mathbb{P}_n) &< \mathcal{L}(\theta_{\ell b}^*(P), \lambda_{\ell b}^*(\mathbb{P}_n))(\mathbb{P}_n) + \varepsilon/3, \\ \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(\mathbb{P}_n))(\mathbb{P}_n) &> \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(P))(\mathbb{P}_n) - \eta/3. \end{aligned}$$

Combining these inequalities we obtain w.p.a. 1:

$$\mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(\mathbb{P}_n))(P) < \mathcal{L}(\theta_{\ell b}^*(P), \lambda_{\ell b}^*(\mathbb{P}_n))(P) + \varepsilon \leq \mathcal{L}(\theta_{\ell b}^*(P), \lambda_{\ell b}^*(P))(P) + \varepsilon,$$

$$\mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(\mathbb{P}_n))(P) > \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(P))(P) - \eta \geq \mathcal{L}(\theta_{\ell b}^*(P), \lambda_{\ell b}^*(P))(P) - \eta.$$

Now let  $B_\theta$  and  $B_\lambda$  be any open balls around  $\theta_{\ell b}^*(P)$  and  $\lambda_{\ell b}^*(P)$ , respectively, and set:

$$\varepsilon = \inf_{\Theta \cap B_\theta^c} \mathcal{L}(\theta, \lambda_{\ell b}^*(P))(P) - \mathcal{L}(\theta_{\ell b}^*(P), \lambda_{\ell b}^*(P))(P),$$



$$\eta = \mathcal{L}(\theta_{\ell b}^*(P), \lambda_{\ell b}^*(P))(P) - \sup_{\Lambda \cap B_\lambda^c} \mathcal{L}(\theta_{\ell b}^*(P), \lambda)(P).$$

Note by Assumption 3.3, we have that the optimal solutions and Lagrange multipliers are unique, so that  $\varepsilon, \eta > 0$ . Combining with the results above we conclude that w.p.a. 1:

$$\sup_{\Lambda \cap B_\lambda^c} \mathcal{L}(\theta_{\ell b}^*(P), \lambda)(P) < \mathcal{L}(\theta_{\ell b}^*(\mathbb{P}_n), \lambda_{\ell b}^*(\mathbb{P}_n))(P) < \inf_{\Theta \cap B_\theta^c} \mathcal{L}(\theta, \lambda_{\ell b}^*(P))(P).$$

Furthermore at least one of the inequalities in the previous display is violated if either  $\theta_{\ell b}^*(\mathbb{P}_n) \notin B_\theta$  or  $\lambda_{\ell b}^*(\mathbb{P}_n) \notin B_\lambda$ , which concludes the proof.  $\blacksquare$

**Lemma A.5.** *Under Assumptions 3.1 and 3.5:*

- (i) *For every  $\varepsilon > 0$  there exists a  $\delta > 0$  such that  $\|\theta - \theta'\| < \delta$  implies  $\rho_P(\theta, \theta') < \varepsilon$  for all  $P \in \mathcal{P}$ .*
- (ii) *Any function  $h \in \ell^\infty(\mathcal{F})$  uniformly continuous in the sup-norm with respect to  $\rho_P$  is uniformly continuous in the sup-norm with respect to  $\|\cdot\|$ .*

*Proof.* Part (i): Recall that under Assumption 3.5 the semimetric  $\rho_P$  satisfies:

$$\lim_{\delta \downarrow 0} \sup_{\|(\theta_1, \theta'_1) - (\theta_2, \theta'_2)\| < \delta} \sup_{P \in \mathcal{P}} |\rho_P(\theta_1, \theta'_1) - \rho_P(\theta_2, \theta'_2)| = 0.$$

Now take  $(\theta_2, \theta'_2) = (\theta'_1, \theta_1)$  and obtain:

$$\lim_{\delta \downarrow 0} \sup_{\|\theta_1 - \theta'_1\| < \delta} \sup_{P \in \mathcal{P}} \rho_P(\theta_1, \theta'_1) = 0.$$

Thus, we conclude for any  $\varepsilon > 0$  there exists a  $\delta > 0$  such that:

$$\sup_{\|\theta_1 - \theta'_1\| < \delta} \sup_{P \in \mathcal{P}} \rho_P(\theta_1, \theta'_1) < \varepsilon.$$

In other words:

$$\{\theta, \theta' \in \Theta : \|\theta - \theta'\| < \delta\} \subseteq \{\theta, \theta' \in \Theta : \rho_P(\theta, \theta') < \varepsilon\}.$$

Part (ii): By uniform continuity of  $h$  we have for any  $\eta > 0$ , there exists a  $\varepsilon(\eta, P) > 0$  such that:

$$\sup_{\rho_P(\theta, \theta') < \varepsilon(\eta, P)} \|hf(w, \theta) - hf(w, \theta')\| < \eta.$$

However, for any such  $\varepsilon(\eta, P) > 0$ , by Part (i) there exists a  $\delta(\varepsilon(\eta, P)) > 0$  such that:

$$\{\theta, \theta' \in \Theta : \|\theta - \theta'\| < \delta\} \subseteq \{\theta, \theta' \in \Theta : \rho_P(\theta, \theta') < \varepsilon\}.$$

We conclude that for any  $\eta > 0$  there exists a  $\delta(\eta, P) > 0$  such that:

$$\sup_{\|\theta - \theta'\| < \delta(\eta, P)} \|hf(w, \theta) - hf(w, \theta')\| \leq \sup_{\rho_P(\theta, \theta') < \varepsilon(\eta, P)} \|hf(w, \theta) - hf(w, \theta')\| < \eta.$$



which completes the proof. ■



## References

- Andrews, D. W. and Guggenberger, P. (2009). Validity of subsampling and plug-in asymptotic inference for parameters defined by moment inequalities. *Econometric Theory*, 25(3):669–709.
- Andrews, D. W. and Soares, G. (2010). Inference for parameters defined by moment inequalities using generalized moment selection. *Econometrica*, 78(1):119–157.
- Belloni, A., Chernozhukov, V., Fernández-Val, I., and Hansen, C. (2017). Program evaluation and causal inference with high-dimensional data. *Econometrica*, 85(1):233–298.
- Beresteanu, A. and Molinari, F. (2008). Asymptotic properties for a class of partially identified models. *Econometrica*, 76(4):763–814.
- Borwein, J. and Lewis, A. S. (2010). *Convex analysis and nonlinear optimization: theory and examples*. Springer Science & Business Media.
- Bugni, F. A., Canay, I. A., and Shi, X. (2015). Specification tests for partially identified models defined by moment inequalities. *Journal of Econometrics*, 185(1):259–282.
- Bugni, F. A., Canay, I. A., and Shi, X. (2017). Inference for subvectors and other functions of partially identified parameters in moment inequality models. *Quantitative Economics*, 8(1):1–38.
- Chen, X., Christensen, T., O’Hara, K., and Tamer, E. (2016). Mcmc confidence sets for identified sets. *arXiv preprint arXiv:1605.00499*.
- Chernozhukov, V., Hong, H., and Tamer, E. (2007). Estimation and confidence regions for parameter sets in econometric models. *Econometrica*, 75(5):1243–1284.
- Fang, Z. and Santos, A. (2018). Inference on directionally differentiable functions. *The Review of Economic Studies*, 86(1):377–412.
- Freyberger, J. and Horowitz, J. L. (2015). Identification and shape restrictions in nonparametric instrumental variables estimation. *Journal of Econometrics*, 189(1):41–53.
- Gafarov, B. (2018). Inference on scalar parameters in setidentified affine models. *Working Paper*.
- Gafarov, B., Meier, M., and Olea, J. L. M. (2018). Delta-method inference for a class of set-identified svars. *Journal of Econometrics*, 203(2):316–327.
- Hirano, K. and Porter, J. R. (2012). Impossibility results for nondifferentiable functionals. *Econometrica*, 80(4):1769–1790.
- Hoffmann-Jørgensen, J. (1991). *Stochastic processes on Polish spaces*. Various publications series. Aarhus Universitet. Matematisk Institut.



- Kaido, H., Molinari, F., and Stoye, J. (2017). Confidence intervals for projections of partially identified parameters. *arXiv preprint arXiv:1601.00934*.
- Kaido, H. and Santos, A. (2014). Asymptotically efficient estimation of models defined by convex moment inequalities. *Econometrica*, 82(1):387–413.
- Kasy, M. (2015). Uniformity and the delta method. *arXiv preprint arXiv:1507.05731*.
- Kasy, M. (2016). Partial identification, distributional preferences, and the welfare ranking of policies. *Review of Economics and Statistics*, 98(1):111–131.
- Ok, E. A. (2007). *Real analysis with economic applications*, volume 10. Princeton University Press.
- Pakes, A., Porter, J., Ho, K., and Ishii, J. (2011). Moment inequalities and their applications, discussion paper, harvard.
- Romano, J. P. and Shaikh, A. M. (2008). Inference for identifiable parameters in partially identified econometric models. *Journal of Statistical Planning and Inference*, 138(9):2786–2807.
- Shapiro, A. (1990). On concepts of directional differentiability. *Journal of optimization theory and applications*, 66(3):477–487.
- Shapiro, A. (1991). Asymptotic analysis of stochastic programs. *Annals of Operations Research*, 30(1):169–186.
- Shapiro, A., Dentcheva, D., and Ruszczyński, A. (2009). *Lectures on stochastic programming: modeling and theory*. SIAM.
- Sheehy, A. and Wellner, J. A. (1992). Uniform donsker classes of functions. *The Annals of Probability*, pages 1983–2030.
- Shi, X. and Shum, M. (2015). Simple two-stage inference for a class of partially identified models. *Econometric Theory*, 31(3):493–520.
- Stoye, J. (2009). More on confidence intervals for partially identified parameters. *Econometrica*, 77(4):1299–1315.
- Torgovitsky (2016). Nonparametric inference on state dependence with applications to employment dynamics.
- Van Der Vaart, A. W. and Wellner, J. A. (1996). Weak convergence. In *Weak Convergence and Empirical Processes*, pages 16–28. Springer.
- Wachsmuth, G. (2013). On licq and the uniqueness of lagrange multipliers. *Operations Research Letters*, 41(1):78–80.