

CONTINUUM SCHROEDINGER OPERATORS FOR SHARPLY TERMINATED GRAPHENE-LIKE STRUCTURES

C.L. FEFFERMAN AND M. I. WEINSTEIN

ABSTRACT. We study the single electron model of a semi-infinite graphene sheet interfaced with the vacuum and terminated along a zigzag edge. The model is a Schroedinger operator acting on $L^2(\mathbb{R}^2)$: $H_{\text{edge}}^\lambda = -\Delta + \lambda^2 V_\sharp$, with a potential V_\sharp given by a sum of translates an atomic potential well, V_0 , of depth λ^2 , centered on a subset of the vertices of a discrete honeycomb structure with a zigzag edge. We give a complete analysis of the low-lying energy spectrum of H_{edge}^λ in the strong binding regime (λ large). In particular, we prove scaled resolvent convergence of H_{edge}^λ acting on $L^2(\mathbb{R}^2)$, to the (appropriately conjugated) resolvent of a limiting discrete tight-binding Hamiltonian acting in $l^2(\mathbb{N}_0; \mathbb{C}^2)$. We also prove the existence of *edge states*: solutions of the eigenvalue problem for H_{edge}^λ which are localized transverse to the edge and pseudo-periodic (propagating or plane-wave like) parallel to the edge. These edge states arise from a “flat-band” of eigenstates the tight-binding Hamiltonian.

1. INTRODUCTION

Tight binding models are discrete models which are central to the modeling of spatially periodic and more general crystalline structures in condensed matter physics. These models apply when the the quantum state of the system is well-approximated by superpositions of translates of highly-localized quantum states (orbitals) within deep atomic potential wells centered at lattice sites [3]. An important example is the tight-binding model of graphene, a planar honeycomb arrangement of carbon atoms with two atoms per unit cell. The two-band tight-binding model yields an explicit approximation for its lowest two dispersion surfaces, which touch conically at *Dirac points* over the vertices of the Brillouin zone [66]. Such Dirac points are central to the remarkable electronic properties of graphene [27, 50, 52, 69] and its artificial (electronic, photonic, acoustic, mechanical, . . .) analogues; see, for example, [8, 38, 44, 49, 57, 63] and the survey [54]. The existence of Dirac points for generic honeycomb Schroedinger operators was proved in [24, 25]; see also [6]. That the two-band tight-binding model gives an accurate approximation of the low-lying dispersion surfaces in the regime of strong binding was proved in [26]; see also Section 1.3. Other results on Dirac points for Schroedinger operators on \mathbb{R}^2 may be found in [1, 2, 14, 29, 40], coupled oscillator models [45] and on quantum graphs in [17, 39].

Edge states are modes which are propagating (plane-wave like) parallel to an interface and which are localized transverse to the interface. In condensed matter physics edge states describe the phenomenon of electrical conduction along an interface. Two types of interfaces of great physical interest are a sharp terminations of a bulk structure structure studied in this

Date: February 28, 2022.

Key words and phrases. Schrödinger equation, Dirac point, Floquet-Bloch spectrum, Topological insulator, protected edge state, Honeycomb lattice, strong binding regime, tight binding approximation.

article (see [15, 28, 47, 48]) and domain wall / line-defects within the bulk (see [8, 38, 51, 64] and studied, for example, in [18, 20, 22, 23, 43]). The role of edge or surface modes in the spectral theory of Schroedinger operators with potentials which model, for example, the interface between a general periodic medium and a vacuum is studied in *e.g.* [13, 37].

In this paper we study the low-lying energy spectrum (discrete and continuous spectrum) of a sharply terminated honeycomb structure, corresponding to a semi-infinite sheet of graphene joined to the vacuum along a sharp interface. We prove convergence of the operator resolvent to that of a discrete tight-binding model and construct the continuous spectrum of edge states.

Edge states in honeycomb structures such as graphene are of particular interest as foundational building blocks in the field topological insulators (TI). TI's are materials which are insulating in their bulk and conduction along boundaries, which is robust against large localized perturbations. When graphene is subjected to a magnetic field, its edge currents become unidirectional and acquire such robustness. This phenomenon has an explanation in terms of topological invariants associated with a bulk Floquet-Bloch vector bundle, which takes on non-trivial values when time-reversal symmetry is broken [31, 35, 36].

A key difference between the types of interfaces is that the sharply terminated structure has no spectral gap, resulting in certain edge orientations supporting edge states and others not. In contrast, the domain wall structures perturbations studied in [18, 20, 22, 23, 43] have edge states which localize along arbitrary rational edges. For a discussion of the roles played by edge orientation and the type of symmetry breaking in the existence and robustness of edge states for domain wall / line-defects, see [20].

Specifically, for the tight-binding model, edge states exist at sharp terminations along a zigzag edge for a subinterval of parallel quasi-momenta, $k_{\parallel} \in [0, 2\pi)$ associated with the direction of translation invariance parallel to the edge. They do not exist at the sharp termination along an armchair edge; see, for example, [15, 28, 47, 48] and Section 2. Such results may be interpreted as consequences of the non-vanishing of the Berry-Zak phase, $\mathcal{Z}(k_{\parallel})$, defined as the integral of the Berry connection over the one-dimensional Brillouin zone associated with the type of edge [15, 47].

1.1. Mathematical setup. In this paper we initiate a study of these phenomena in the context of the underlying continuum equations of quantum physics, in particular the single-electron model of bulk (infinite) graphene and its terminations. In particular, we study Schroedinger operators on \mathbb{R}^2 for a sharp termination of a honeycomb structure along a zigzag edge.

We denote the equilateral triangular lattice in \mathbb{R}^2 by .

$$(1.1) \quad \Lambda = \mathbb{Z}\mathbf{v}_1 \oplus \mathbb{Z}\mathbf{v}_2 ,$$

where \mathbf{v}_1 and \mathbf{v}_2 are given by

$$(1.2) \quad \mathbf{v}_1 = \begin{pmatrix} \frac{\sqrt{3}}{2} \\ \frac{1}{2} \end{pmatrix}, \quad \mathbf{v}_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix} .$$

The dual lattice, Λ^* , is given by

$$(1.3) \quad \Lambda^* = \mathbb{Z}\mathbf{r}_1 \oplus \mathbb{Z}\mathbf{r}_2 ,$$

where \mathfrak{K}_1 and \mathfrak{K}_2 are given by

$$(1.4) \quad \mathfrak{K}_1 = 2\pi \begin{pmatrix} \frac{2\sqrt{3}}{3} \\ 0 \end{pmatrix}, \quad \mathfrak{K}_2 = 2\pi \begin{pmatrix} -\frac{\sqrt{3}}{3} \\ 1 \end{pmatrix}.$$

Note that

$$(1.5) \quad \mathfrak{K}_i \cdot \mathbf{v}_m = 2\pi \delta_{im}.$$

To generate the honeycomb structure, we first fix base points in \mathbb{R}^2 :

$$(1.6) \quad \mathbf{v}_A = (0, 0), \quad \mathbf{v}_B = \left(1/2, 1/(2\sqrt{3})\right).$$

The honeycomb structure, \mathbb{H} , is the union of the two interpenetrating sublattices

$$(1.7) \quad \Lambda_A = \mathbf{v}_A + \Lambda, \quad \Lambda_B = \mathbf{v}_B + \Lambda :$$

$$(1.8) \quad \mathbb{H} = \Lambda_A \cup \Lambda_B .$$

Let $V_0(\mathbf{x})$ be an *atomic potential well* which may be considered, for the present discussion, to be radially symmetric, compactly supported with $\text{supp } V_0 \subset B_{r_0}(0)$, the open disc of radius r_0 about 0. We discuss more general and physically reasonable conditions on V_0 below in Section 3.

Our bulk Hamiltonian is the honeycomb Schroedinger operator:

$$(1.9) \quad H_{\text{bulk}}^\lambda = -\Delta + \lambda^2 V(\mathbf{x}) \text{ acting on } L^2(\mathbb{R}^2),$$

where $V(\mathbf{x})$ is a superposition identical atomic potential wells, centered at the vertices of \mathbb{H} :

$$(1.10) \quad V(\mathbf{x}) = \sum_{\mathbf{v} \in \mathbb{H}} V_0(\mathbf{x} - \mathbf{v}), \quad \mathbf{x} \in \mathbb{R}^2,$$

The potential $V(\mathbf{x})$ satisfies the conditions of a *honeycomb lattice potential* in the sense of Definition 2.1 of [25]. For all but a discrete subset of values of λ (including $\lambda = 0$), the operator H_{bulk}^λ has Dirac points at energy / quasi-momentum pairs, $(E_D^\lambda, \mathbf{K}_*)$, where \mathbf{K}_* , varies over the vertices of the Brillouin zone [24, 25]; see also [6]. Moreover, for λ large (strong binding), the low-lying Floquet-Bloch dispersion surfaces of H_{bulk}^λ , when rescaled, are uniformly approximated by the dispersion surfaces of the two-band tight-binding model [26].

Consider now a “half-plane” of vertices $\mathbb{H}_\sharp \subset \mathbb{H}$, whose extreme points trace out a zigzag pattern:

$$(1.11) \quad \mathbb{H}_\sharp \equiv \{\mathbf{v}_A + \mathbb{N}_0 \mathbf{v}_1 \oplus \mathbb{Z} \mathbf{v}_2\} \cup \{\mathbf{v}_B + \mathbb{N}_0 \mathbf{v}_1 \oplus \mathbb{Z} \mathbf{v}_2\}, \quad \mathbb{N}_0 = \{0, 1, 2, \dots\},$$

The set \mathbb{H}_\sharp is invariant with respect to translations by \mathbf{v}_2 and is the subset of sites in \mathbb{H} to the right of an infinite zigzag edge; see Figure 1. The set of zigzag edge (boundary) sites, also translation invariant by \mathbf{v}_2 , is given by: $\{\mathbf{v}_A + \mathbb{Z} \mathbf{v}_2\} \cup \{\mathbf{v}_B + \mathbb{Z} \mathbf{v}_2\}$.

We define the potential

$$(1.12) \quad V_\sharp(\mathbf{x}) = \sum_{\mathbf{v} \in \mathbb{H}_\sharp} V_0(\mathbf{x} - \mathbf{v}), \quad \mathbf{x} \in \mathbb{R}^2.$$

The operator

$$H_{\text{edge}}^\lambda = -\Delta + \lambda^2 V_\sharp(\mathbf{x})$$




FIGURE 1. (a) \mathbb{H} : Bulk honeycomb structure consists of all vertices (circles, light and dark). (b) \mathbb{H}_\sharp : Honeycomb structure terminated along a zigzag edge consists of vertices indicated by dark circles; see (1.11). (c) Ω_Σ : Indicated strip is a choice of fundamental cell for the cylinder $\Sigma = \mathbb{R}^2/\mathbb{Z}\mathbf{v}_2$. $\Omega_\Sigma = \Omega_{-1} \cup \Omega_0 \cup \Omega_1 \cup \dots \cup \Omega_n \cup \dots$. Sites: $\mathbf{v}_A^n, \mathbf{v}_B^n$ in finite parallelograms Ω_n , $n \geq 0$, are sites in \mathbb{H}_\sharp . Ω_{-1} denotes the infinite parallelogram containing no vertices of the terminated structure, \mathbb{H}_\sharp .

models a half-plane of graphene interfaced with the vacuum along a zigzag edge. Note the translation invariance: $V_\sharp(\mathbf{x} + \mathbf{v}_2) = V_\sharp(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^2$.

Let $(E_0^\lambda, p_0^\lambda(\mathbf{x}))$, with $p_0^\lambda > 0$ and L^2 -normalized, denote the ground state eigenpair of the *atomic Hamiltonian*

$$H_{\text{atom}}^\lambda = -\Delta + \lambda^2 V_0(\mathbf{x}).$$

Let ρ_λ denote the *hopping coefficient*, given by:

$$(1.13) \quad \rho_\lambda = \int_{|\mathbf{y}| < r_0} p_0^\lambda(\mathbf{y}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} - \mathbf{e}) d\mathbf{y},$$

where \mathbf{e} is any vector from one lattice site in \mathbb{H} to a nearest neighbor in \mathbb{H} , *e.g.* $\mathbf{v}_B - \mathbf{v}_A$. The potential $V_0(\mathbf{y})$ and ground state $p_0^\lambda(\mathbf{y})$ are localized around $\mathbf{y} = 0$, while $p_0^\lambda(\mathbf{y} - \mathbf{e})$, is localized at any nearest neighbor site $\mathbf{e} \in \mathbb{H}$. Recall that $\text{supp } V_0$ is contained in the set where $|\mathbf{x}| < r_0$. For λ large ρ_λ is exponentially small (see (3.3)) [26].

The key accomplishments of this paper are the following:

- (1) *Theorem 1.1 (Scaled resolvent convergence)*: We prove for $\lambda \geq \lambda_\star$ sufficiently large (the strong binding regime), that the re-centered and scaled resolvent $\left((H_{\text{edge}}^\lambda - E_0^\lambda)/\rho_\lambda - zI \right)^{-1}$ has a universal limit (in the uniform operator norm) described by a discrete (tight-binding) Hamiltonian, defined on a truncated honeycomb structure. The band structure of this limiting operator is displayed in Figure 2.

- (2) *Theorem 1.2 (Zigzag edge states)*: We construct a continuum of edge state modes. These are eigenstates of H_{edge}^λ , which are propagating (plane-wave like) parallel to and localized transverse to the zigzag edge. Upon appropriate λ -dependent rescaling, these edge-states are close to (and converge as λ tends to infinity to) the flat band of zero energy edge states of the tight-binding model; see Figure 2.
- (3) *Resolvent kernel bounds on arbitrary discrete sets*: The methods of this article go considerably beyond those of our previous article on the strong binding regime [26], which established convergence to the (universal) two-band tight binding spectrum for the bulk graphene-like structures. Since Theorems 1.1 and 1.2 involve convergence of operators and eigenstates on an infinite cylinder (Figure 1), we required pointwise decay properties of the resolvent kernel H_{edge}^λ for energies near E_0^λ . These bounds are stated in Theorem 10.1. In Proposition 10.15 we establish these kernel estimates for potentials which are a sum of atomic potentials centered on an arbitrary discrete set of lattice sites $\Gamma \subset \mathbb{R}^2$ (not necessarily translation invariant) whose minimal pairwise distance is Mr_0 , where r_0 is the radius of the support of V_0 and $M > 2$ is some positive constant. We then specialize to a translation invariant set to obtain Theorem 10.1. We believe the technique we have developed will be quite broadly applicable.

We next introduce the *edge state eigenvalue problem*. Associated with the translation invariance of $-\Delta + \lambda^2 V_\#(\mathbf{x})$ by \mathbf{v}_2 is a *parallel quasi-momentum*, denoted $k_\parallel \in [0, 2\pi)$. The condition that an edge state, Φ , is propagating parallel to the zigzag edge is:

$$(1.14) \quad \Phi(\mathbf{x} + \mathbf{v}_2) = e^{ik_\parallel} \Phi(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^2 .$$

We introduce the cylinder

$$(1.15) \quad \Sigma = \mathbb{R}^2 / \mathbb{Z}\mathbf{v}_2.$$

The space $L^2(\Sigma)$ consists of functions which are square integrable over a fundamental cell of Σ , *e.g.* the strip Ω_Σ shown in Figure 1, and which satisfy the periodic boundary condition with respect to \mathbf{v}_2 : $\phi(\mathbf{x} + \mathbf{v}_2) = \phi(\mathbf{x})$ for almost all $\mathbf{x} \in \Omega_\Sigma$ and all $\mathbf{v} \in \Lambda$.

We enforce the condition that (i) Φ is k_\parallel -pseudo-periodic parallel to the zigzag edge, (1.14), and (ii) decaying to zero transverse to the zigzag edge as \mathbf{x} tends to infinity by requiring

$$e^{-i\frac{k_\parallel}{2\pi}\mathfrak{K}_2 \cdot \mathbf{x}} \Phi(\mathbf{x}) \in L^2(\Sigma).$$

For such functions we write $\Phi \in L_{k_\parallel}^2(\Sigma)$ or just $\Phi \in L_{k_\parallel}^2$. We can now formulate the

k_\parallel -Zigzag Edge State Eigenvalue Problem for $H_{\text{edge}}^\lambda = -\Delta + V_\#(\mathbf{x})$:

$$(1.16) \quad H_{\text{edge}}^\lambda \Psi(\mathbf{x}) \equiv \left(-\Delta + \lambda^2 V_\#(\mathbf{x}) \right) \Psi(\mathbf{x}) = E \Psi(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^2, \quad \Psi \in L_{k_\parallel}^2(\Sigma).$$

Defining $\Psi(\mathbf{x}) = e^{i\frac{k_\parallel}{2\pi}\mathfrak{K}_2 \cdot \mathbf{x}} \psi(\mathbf{x})$, we may formulate (1.16) equivalently as:

$$(1.17) \quad H_{\text{edge}}^\lambda(k_\parallel)\psi \equiv \left(-\left(\nabla + i\frac{k_\parallel}{2\pi}\mathfrak{K}_2 \right)^2 + \lambda^2 V_\#(\mathbf{x}) \right) \psi(\mathbf{x}) = E \psi(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^2, \quad \psi \in L^2(\Sigma) .$$

We refer to non-trivial solutions of the eigenvalue problem (1.16) (equivalently (1.17)) as zigzag edge states.

Before stating our main results Theorems 1.1 and 1.2, we recall a key observation used in [26] to obtain the low-lying dispersion surfaces (energies near the atomic ground state energy, E_0^λ) of the bulk honeycomb Schroedinger operator, H_{bulk}^λ . That is, for λ large, the \mathbf{k} - pseudo-periodic Floquet-Bloch eigenmodes which are associated with the two lowest spectral bands of H_{bulk}^λ , acting in $L^2(\mathbb{R}^2)$, can be uniformly approximated by appropriate linear combinations of the two \mathbf{k} - pseudo-periodic functions: $P_{\mathbf{k},I}^\lambda(\mathbf{x})$, $I = A, B$. These functions are constructed as \mathbf{k} - pseudo-periodic weighted sums of translates, $p_0^\lambda(\mathbf{x} + \mathbf{v})$, of the atomic ground state, where \mathbf{v} varies over the sublattices: $\Lambda_I = \mathbf{v}_I + \Lambda$, $I = A, B$.

In the present work, to study the low-lying spectral bands associated with eigenvalue problem $H_{\text{edge}}^\lambda(k_\parallel)\psi_{k_\parallel}^\lambda = E^\lambda(k_\parallel)\psi_{k_\parallel}$, $\psi_{k_\parallel}^\lambda \in L^2(\Sigma)$ with $E^\lambda(k_\parallel)$ near E_0^λ and we find it very natural to approximate eigenstates by superpositions of the infinite family of functions

$$(1.18) \quad p_{I,k_\parallel}^\lambda[n](\mathbf{x}) \in L^2(\Sigma), \quad I = A, B, \quad n \geq 0,$$

which are constructed as k_\parallel - dependent and periodized (infinite) sums of translates of the ground state $p_0^\lambda(\mathbf{x})$ over the one-dimensional sublattices: $\mathbf{v}_I + n\mathbf{v}_1 + \mathbb{Z}\mathbf{v}_2$ of Λ_I , $I = A, B$ and $n \geq 0$; see (1.7). The states $p_{I,k_\parallel}^\lambda[n](\mathbf{x})$ are introduced in Definition 4.1 in Section 4. For λ sufficiently large, any $F \in L^2(\Sigma)$ has the expansion

$$(1.19) \quad F = \sum_{I=A,B} \sum_{n \geq 0} \alpha_n^I p_{I,k_\parallel}^\lambda[n](\mathbf{x}) + F_\perp,$$

where $\{\alpha_n^I\} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ and F_\perp is $L^2(\Sigma)$ - orthogonal to the span of the functions $p_{I,k_\parallel}^\lambda[n]$; see Proposition 4.4. The tight-binding (discrete) edge Hamiltonian, $H_\#^{\text{TB}}(k_\parallel)$ acting in $l^2(\mathbb{N}_0; \mathbb{C}^2)$, arises via translation and rescaling, of the operator whose matrix elements are $\left\langle p_{J,k_\parallel}^\lambda[m], H_{\text{edge}}^\lambda(k_\parallel) p_{I,k_\parallel}^\lambda[n] \right\rangle_{L_{k_\parallel}^2}$, for $J, I = A, B$ and $m, n \geq 0$. The tight-binding model is studied in Section 2 and its band spectrum is displayed in Figure 2.

1.2. Main results. The relation of $H_{\text{edge}}^\lambda(k_\parallel)$ to the tight-binding Hamiltonian $H_\#^{\text{TB}}(k_\parallel)$ is given by the following result on scaled resolvent convergence.

Theorem 1.1 (Scaled resolvent convergence). *As in Theorem 1.2, assume that E_0^λ , the ground state energy of the atomic Hamiltonian, $H_{\text{atom}}^\lambda = -\Delta + \lambda^2 V_0$, satisfies the conditions (GS) (3.4) and (EG) (3.6) on the ground state energy and energy-gap, respectively.*

Let \mathcal{C} denote a compact subset of $\mathbb{C} \setminus \sigma(H_\#^{\text{TB}}(k_\parallel))$, the resolvent set of $H_\#^{\text{TB}}(k_\parallel)$. There exist constants λ_\star , C_\star and c , which are independent of λ but which depend on \mathcal{C} and conditions (GS) and (EG), such that for all $\lambda > \lambda_\star$ the following holds:

$$(1.20) \quad \text{Let } J_{k_\parallel} : L^2(\Sigma) \mapsto l^2(\mathbb{N}_0; \mathbb{C}^2) \oplus \text{span}\{p_{I,k_\parallel}^\lambda[n]\}^\perp \text{ be defined, via (1.19), } F \mapsto (\{\alpha_n^I[F]\}, F_\perp)^\top. \text{ Then, uniformly in } k_\parallel \in [0, 2\pi], \text{ we have}$$

$$\left\| \left(\rho_\lambda^{-1} \left(H_{\text{edge}}^\lambda(k_\parallel) - E_0^\lambda \right) - z \text{Id} \right)^{-1} - J_{k_\parallel}^* \left(H_\#^{\text{TB}}(k_\parallel) - z \text{Id} \right)^{-1} J_{k_\parallel} \right\|_{L^2(\Sigma) \rightarrow L^2(\Sigma)} \leq C_\star e^{-c\lambda}.$$

In preparation for our theorem on edge states, we introduce the functions:

$$(1.21) \quad \zeta(k_{\parallel}) = 1 + e^{ik_{\parallel}}, \quad \delta_{\text{gap}}(k_{\parallel}) = \left| 1 - |\zeta(k_{\parallel})| \right| \geq 0, \quad \delta_{\text{max}}(k_{\parallel}) = 1 + |\zeta(k_{\parallel})|.$$

We note that for $k_{\parallel} \in [0, 2\pi]$ that $\delta_{\text{gap}}(k_{\parallel}) = 0$ if and only if $k_{\parallel} \in \{2\pi/3, 4\pi/3\}$.

Theorem 1.2 (Zigzag Edge States). *Assume that E_0^λ , the ground state energy of the atomic Hamiltonian, $H_{\text{atom}}^\lambda = -\Delta + \lambda^2 V_0$, satisfies the conditions (GS) (3.4) and (EG) (3.6) on the ground state energy and energy-gap, respectively. Let \mathcal{J} denote an arbitrary compact subinterval of quasi-momenta:*

$$(1.22) \quad \mathcal{J} \subset \subset (2\pi/3, 4\pi/3) \setminus \{\pi\}.$$

Thus, $\min_{k_{\parallel} \in \mathcal{J}} \delta_{\text{gap}}(k_{\parallel}) > 0$.

There exists $\lambda_* = \lambda_*(\mathcal{J}) > 0$ sufficiently large, such that for all $\lambda > \lambda_*$ the following holds:

- (1) There is a mapping $k_{\parallel} \in \mathcal{J} \mapsto (E^\lambda(k_{\parallel}), \psi_{k_{\parallel}}^\lambda)$, from parallel quasimomenta k_{\parallel} to simple eigenpairs of the family of the k_{\parallel} -edge state eigenvalue problem (1.16):

$$(1.23) \quad \begin{aligned} H_{\text{edge}}^\lambda(k_{\parallel}) \psi_{k_{\parallel}} &= E^\lambda(k_{\parallel}) \psi_{k_{\parallel}}^\lambda, \quad \psi_{k_{\parallel}} \in L^2(\Sigma) \\ E^\lambda(k_{\parallel}) &= E_0^\lambda + \rho_\lambda \Omega^\lambda(k_{\parallel}), \end{aligned}$$

where $|\Omega^\lambda(k_{\parallel})| \lesssim e^{-c\lambda}$ with $c > 0$ independent of λ . Correspondingly, the eigenvalue problem (1.16) is solved by the states $\Psi_{k_{\parallel}}^\lambda(\mathbf{x}) = e^{i\frac{k_{\parallel}}{2\pi} \mathbf{k}_2 \cdot \mathbf{x}} \psi_{k_{\parallel}}^\lambda(\mathbf{x})$.

- (2) The edge states $\psi_{k_{\parallel}}^\lambda \in L^2_{k_{\parallel}}(\Sigma)$ are approximated to within $\mathcal{O}(e^{-c\lambda})$ error in $L^2(\Sigma)$ as:

$$(1.24) \quad \psi_{k_{\parallel}}^\lambda(\mathbf{x}) = \sum_{n \geq 0} \alpha_A^n p_{A,k_{\parallel}}^\lambda[n](\mathbf{x}) + \sum_{n \geq 0} \alpha_B^n p_{B,k_{\parallel}}^\lambda[n](\mathbf{x}) + \mathcal{O}_{L^2(\Sigma)}(e^{-c\lambda}),$$

where $c > 0$ is independent of λ . Here, $\psi_{k_{\parallel}}^{\text{TB,bd}} \equiv \{(\alpha_A^n, \alpha_B^n)^\top\}_{n \geq 0} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$, $\|\psi_{k_{\parallel}}^{\text{TB,bd}}\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)} = 1$ is a zero energy normalized eigenstate of the limiting tight-binding edge Hamiltonian; $H_{\#}^{\text{TB}}(k_{\parallel}) \psi_{k_{\parallel}}^{\text{TB,bd}} = 0$. See Theorem 2.2 in Section 2.

Remark 1.3 (Symmetry of edge state curves). Let $k_{\parallel} \in [0, \pi]$. If $(E^\lambda(k_{\parallel}), \Psi_{k_{\parallel}}^\lambda(\mathbf{x}))$ is an eigenpair of the k_{\parallel} -edge state eigenvalue problem, then $(E^\lambda(k_{\parallel}), \overline{\Psi_{k_{\parallel}}^\lambda(\mathbf{x})})$ is an eigenpair of the $2\pi - k_{\parallel}$ edge state eigenvalue problem.

Remark 1.4 (Non-flatness of band). The large λ edge states of eigenfrequencies, $E^\lambda(k_{\parallel})$, in Theorem 1.2 arise from the *flat band* of edge states, $\Omega(k_{\parallel}) = 0$ for $2\pi/3 < k_{\parallel} < 4\pi/3$, of the tight-binding Hamiltonian, $H_{\#}^{\text{TB}}(k_{\parallel})$. Although $E^\lambda(k_{\parallel})$ has only exponentially small variation, we do not expect $E^\lambda(k_{\parallel})$ to be identically constant. Indeed, numerical simulations illustrate the weak variation in k_{\parallel} [65].

Remark 1.5 (Regularity). We do not address the question of smoothness of $k_{\parallel} \in \mathcal{J} \mapsto (E^\lambda(k_{\parallel}), \psi_{k_{\parallel}}^\lambda) \in \mathbb{R} \times L^2(\Sigma)$ in the present article. We believe however that the methods of [26] may be adapted to show that this mapping extends as an analytic mapping in a complex neighborhood of \mathcal{J} from which derivative bounds, e.g. on $E^\lambda(k_{\parallel})$ ($k_{\parallel} \in \mathcal{J}$) can be derived via Cauchy estimates.

Remark 1.6. In Theorem 2.2 we find: $\psi_{k_{\parallel}}^{\text{TB,bd}} = \sqrt{1 - |\zeta(k_{\parallel})|^2} \left([-\zeta(k_{\parallel})]^n, 0 \right)^{\top}$. Therefore, at leading order, $\Psi_{k_{\parallel}}^{\lambda}(\mathbf{x})$ is concentrated about the A - sublattice, Λ_A :

$$(1.25) \quad \Psi_{k_{\parallel}}^{\lambda}(\mathbf{x}) = \sqrt{1 - |\zeta(k_{\parallel})|^2} \sum_{n \geq 0} [-\zeta(k_{\parallel})]^n P_{A,k_{\parallel}}^{\lambda}[n](\mathbf{x}) + \mathcal{O}_{L_{k_{\parallel}}^2}(e^{-c\lambda}).$$

Remark 1.7. As noted in our discussion of the tight-binding model in Section 2 (Remark 2.3) the constraint of Theorem 1.2 on parallel quasimomenta: $k_{\parallel} \in (2\pi/3, 4\pi/3)$ ($|\zeta(k_{\parallel})| < 1$) corresponds to the non-vanishing of the *Zak phase*. This is discussed further in Remark 2.3.

Remark 1.8. In work in progress we show, for a sharp termination of the bulk honeycomb structure along an *armchair edge*, that there are no edge states in an energy range about E_0^{λ} . In this case, the relevant Zak phase vanishes for all $k_{\parallel} \in [0, 2\pi]$.

1.3. Relation to previous work. Tight-binding limits arising from general distributions of potential wells has been discussed in the book of [16] as well as [9, 53]. There is extensive related earlier work on the semiclassical limits and methods *e.g.* [10–12, 32, 33, 46, 60–62]. The above works are based on detailed semiclassical (WKB) approximations for potential wells which are assumed to have non-degenerate local minima. In contrast, in the present article our essential assumptions are only on the ground state energy (GS) and spectral gap (EG) of the atomic Hamiltonian, H_{atom}^{λ} for large λ . The relation of the continuum periodic Schroedinger operator with a magnetic field to tight-binding models, such as the Harper model, is studied for example in [34].

For convenience we have restricted attention here to C^{∞} potentials. However, we believe it will be easy to make small changes in our proofs, to apply our results to nonsmooth potentials of interest. In particular, the atomic potential may be taken to have a Coulomb singularity at the origin, or to have the form $V_0(x) = -1$ for \mathbf{x} in a ball B , $V_0(x) = 0$ otherwise. Here the radius of B is taken small enough to satisfy the hypothesis (PW_2) of Section 3.1 below. Examples of artificial graphene, in which experiments are performed, are periodic honeycomb arrays of identical microfeatures, say small discs, with one dielectric constant inside the discs and a second dielectric constant outside the discs. Hence, compactly supported atomic potentials are a natural model; see, for example, [8, 38, 44, 49, 54, 57, 63].

For smooth atomic potentials V_0 with nondegenerate minima, the general semiclassical works in [9, 16, 53] lead to an “interaction matrix”, which defines an operator. In the case of periodic potentials, this can be used to compute relevant dispersion surfaces modulo exponentially small errors. These works do not assert that Dirac points form; indeed, much of the work is in the setting of a square lattice, which does not give rise to Dirac points. However, we believe that these methods are powerful enough to deal with Dirac points of honeycomb lattice potentials, when they are combined with the consequences of special symmetry properties of the honeycomb. The essential requirement for the semiclassical analysis approach is that the atomic potential is smooth and has a nondegenerate minimum. Another aspect of the general work of this semiclassical work is that atomic potentials are not assumed to be of compact support and the interaction matrix (hopping coefficients) are obtained in terms of the Agmon metric. Finally, the consideration of edge states and the spectrum for honeycombs with line defects is not within the scope of [9, 16, 53].

Remark 1.9. A different class of line-defects of great interest in the study of topologically protected edge states is the class of *domain walls*. In our previous work, motivated by




FIGURE 2. Spectrum of tight-binding Hamiltonian $H_{\#}^{\text{TB}}(k_{\parallel})$, for $0 \leq k_{\parallel} \leq 2\pi$, described in Theorem 2.2. This spectrum contains a *flat band* of zero energy states; $H_{\#}^{\lambda}(k_{\parallel})$ has an isolated simple 0– energy eigenstate for $2\pi/3 \leq k_{\parallel} \leq 4\pi/3$. Shaded regions consist of continuous spectrum. For sufficiently large λ , the low-lying part of the spectrum of $-\Delta + \lambda^2 V_{\#} - E_0^{\lambda}$, after rescaling by ρ_{λ} , is approximated by spectrum of the 2-band model $H_{\#}^{\text{TB}}$; see Theorem 1.2.

[30, 56, 67], domain walls are realized by starting with two periodic structures at “ $+\infty$ ” and “ $-\infty$ ”, with a common spectral gap and phase-shifted from one another, and connecting them across a line-defect at which there is no phase-distortion. See the analytical work in 1D [19, 21, 24] and 2D [22, 23, 43] as well as theoretical and experimental work on photonic realizations [41, 42, 55].

Remark 1.10. Quantum graphs [7] are another class of discrete models in condensed matter, electromagnetic and other systems; see also, for example, [4, 5, 58]. An extensive discussion of edge states for nanotube structures in the setting of quantum graphs is given in [17, 39]. It would be of interest to investigate a relation between the edge modes of these models and continuum models.

1.4. Outline of the paper.

We present a brief outline. Section 2 discusses tight binding models; first, the tight binding model for bulk, and then the tight binding model for a honeycomb structure terminated along a zigzag edge.

Section 3 first introduces the atomic Hamiltonian $H_{\text{atom}}^\lambda = -\Delta + V_0$, where V_0 is a potential well whose support in a sufficiently small disc about the origin, and such that V_0 satisfies some basic general assumptions $(PW_1) - (PW_4)$. The bulk honeycomb structure is defined by $H_{\text{bulk}}^\lambda = -\Delta + \lambda^2 V$, where V is the periodic potential by summing V_0 over translates of the atomic potential, V_0 , over the honeycomb structure. Thus V consists of a potential well V_0 centered at each site of the honeycomb. Finally the edge Hamiltonian, $H_{\text{edge}} = -\Delta + \lambda^2 V_{\sharp}$, which acts on $L^2(\mathbb{R}^2)$, has potential V_{\sharp} which is identically equal to V on a half-space with a zigzag edge and zero on the other side of this zigzag edge. (We shall also work with the translated edge Hamiltonian $H_{\sharp}^\lambda = H_{\text{edge}} - E_0^\lambda$.) The edge state eigenvalue problem for parallel quasi-momentum k_{\parallel} is then stated on $L^2(\Sigma)$, where Σ is the infinite cylinder (1.15).

Section 4 introduces a natural basis for approximating the 2 lowest lying bands of H_{\sharp}^λ for λ sufficiently large. This basis consists of functions, $\{p_{I,k_{\parallel}}^\lambda[n](\mathbf{x}) : I = A, B, n \geq 0\}$ on Σ , which are pseudo-periodic (with respect to the direction parallel to the edge) infinite sums of atomic orbitals.

Section 5 establishes energy estimates on H_{\sharp}^λ which imply invertibility of H_{\sharp}^λ on $\mathcal{X}_{AB}(k_{\parallel})$, the orthogonal complement of the orbital subspace: $\text{span}\{p_{I,k_{\parallel}}^\lambda[n] : n \geq 0, I = A, B\}$. This implies that the resolvent of H_{\sharp}^λ is well-defined and bounded on $\mathcal{X}_{AB}(k_{\parallel})$.

Section 6 implements a Lyapunov-Schmidt / Schur complement reduction strategy: The spectral problem on $L^2(\Sigma) = \text{span}\{p_{I,k_{\parallel}}^\lambda[n] : n \geq 0, I = A, B\} \oplus \mathcal{X}_{AB}(k_{\parallel})$ is reduced, using the resolvent bounds on $\mathcal{X}_{AB}(k_{\parallel})$, to an equivalent problem on the space $\text{span}\{p_{I,k_{\parallel}}^\lambda[n] : n \geq 0, I = A, B\}$. This problem depends nonlinearly on the eigenvalue parameter $E = E_0^\lambda + \rho_\lambda \Omega$ and is of the form of an infinite algebraic system:

$$\sum_{I=A,B} \sum_{n \geq 0} \mathcal{M}_{JI}^{\lambda,k_{\parallel}}[m,n](\Omega, k_{\parallel}) \alpha_n^I = 0; \quad J = A, B, \quad m \geq 0$$

for (Ω, α) , where $\alpha = \{\alpha_n^I\}_{n \geq 0, I=A,B} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ are coordinates relative to the basis $\{p_{I,k_{\parallel}}^\lambda[n] : n \geq 0, I = A, B\}$.

Section 7 summarizes the required properties of $\mathcal{M}^\lambda(\Omega, k_{\parallel})$ acting in $l^2(\mathbb{N}_0; \mathbb{C}^2)$. We write $\mathcal{M}^{\lambda,k_{\parallel}}(\Omega, k_{\parallel}) = \mathcal{M}_{\text{lin}}^\lambda(\Omega, k_{\parallel}) - \mathcal{M}_{\text{nl}}^\lambda(\Omega, k_{\parallel})$, separating matrix elements contributions which are linear in $H_{\sharp}^\lambda(k_{\parallel})$ and those which are nonlinear in $H_{\sharp}^\lambda(k_{\parallel})$. We have $\mathcal{M}_{\text{lin}}^\lambda(\Omega, k_{\parallel}) = \rho_\lambda H_{\sharp}^{\text{TB}}(k_{\parallel}) + \mathcal{O}_{l^2 \rightarrow l^2}(\rho_\lambda e^{-c\lambda})$ (Proposition 7.1) and $\mathcal{M}_{\text{nl}}^\lambda(\Omega, k_{\parallel}) = \mathcal{O}_{l^2 \rightarrow l^2}(\rho_\lambda e^{-c\lambda})$ (Proposition 7.2). These propositions are proved in later sections.

Section 8 proves Theorem 1.2, the existence of edge states, bifurcating from the flat band of eigenstates of H_{\sharp}^{TB} , via our formulation of the eigenvalue in $l^2(\mathbb{N}_0; \mathbb{C}^2)$.

Section 9 proves Theorem 1.1, the convergence of a translation and scaling of the resolvent of $H_{\sharp}^{\lambda}(k_{\parallel})$ to that of $H_{\sharp}^{\text{TB}}(k_{\parallel})$.

Section 10 is the most technically involved and introduces techniques not present in our earlier work. Theorem 10.1 is a pointwise estimate on the resolvent kernel of $H_{\sharp}^{\lambda} - z = H_{\text{bulk}}^{\lambda} - (E_0^{\lambda} + z)$, z small, when restricted to the orthogonal complement of $\text{span}\{p_{I,k_{\parallel}}^{\lambda}[n] : n \geq 0, I = A, B\}$. These bounds are stated in Theorem 10.1. We first, in Proposition 10.15, establish these kernel estimates for potentials which are a sum of atomic potentials centered on an arbitrary discrete set of lattice sites $\Gamma \subset \mathbb{R}^2$ (not necessarily translation invariant), whose minimal pairwise distance is Mr_0 , where r_0 is the radius of the support of V_0 and $M > 2$ is some positive constant. We then specialize to a translation invariant set to obtain Theorem 10.1.

Section 11 expands the linear matrix elements, $\mathcal{M}_{\text{lin}}^{\lambda}(\Omega, k_{\parallel})$, in terms of $H_{\sharp}^{\text{TB}}(k_{\parallel})$ and estimates the corrections, proving Proposition 7.1.

Section 12 estimates the nonlinear matrix elements, $\mathcal{M}_{\text{nl}}^{\lambda}(\Omega, k_{\parallel})$, proving Proposition 7.2.

Finally, there are two appendices. Appendix A introduces a technical tool used to construct the resolvent of H_{\sharp}^{λ} on $\mathcal{X}_{AB}^{\lambda}(k_{\parallel})$. Appendix B contains general results on overlap integrals enabling expansion of $\mathcal{M}_{\text{lin}}^{\lambda}(\Omega, k_{\parallel})$, for λ large, estimate corrections.

1.5. Notation.

- (1) $\mathbb{N} = \{1, 2, 3, \dots\}$, $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$.
- (2) When we write the expression $g_{\varepsilon} = \mathcal{O}_X(\gamma_{\varepsilon})$ as $\varepsilon \rightarrow \varepsilon_0 \in \mathbb{R} \cup \{\infty\}$, we mean that there exists $C > 0$, independent of ε , such that $\|g_{\varepsilon}\|_X \leq C\gamma_{\varepsilon}$ as $\varepsilon \rightarrow \varepsilon_0$.
- (3) We shall be concerned with the asymptotic behavior of many expressions, $a(\lambda), b(\lambda), \dots$, in the regime where the parameter λ sufficiently large. The relation $a(\lambda) \lesssim b(\lambda)$ means that there is a constant C , which can be taken to be independent of λ , such that for all λ sufficiently large: $a(\lambda) \leq Cb(\lambda)$.
- (4) $\Lambda = \mathbb{Z}\mathbf{v}_1 \oplus \mathbb{Z}\mathbf{v}_2$, the equilateral triangular lattice, is generated by the basis vectors \mathbf{v}_1 and \mathbf{v}_2 , displayed in (1.2).
- (5) $\mathbf{m}\vec{\mathbf{v}} = m_1\mathbf{v}_1 + m_2\mathbf{v}_2$, where $\mathbf{m} = (m_1, m_2) \in \mathbb{Z}^2$.
- (6) $\Lambda^* = \mathbb{Z}\mathfrak{K}_1 \oplus \mathbb{Z}\mathfrak{K}_2$, the dual lattice, spanned by the dual basis vectors \mathfrak{K}_1 and \mathfrak{K}_2 , displayed in (1.4). Note that $\mathfrak{K}_{\ell} \cdot \mathbf{v}_{\ell'} = 2\pi\delta_{\ell\ell'}$.
- (7) We remark that alternative bases for Λ and Λ^* (used for example in [25, 26]) are:

$$\begin{aligned} \mathbf{v}_1 &= \mathbf{v}_1, & \mathbf{v}_2 &= \mathbf{v}_1 - \mathbf{v}_2 \\ \mathbf{k}_1 &= \mathfrak{K}_1 + \mathfrak{K}_2, & \mathbf{k}_2 &= -\mathfrak{K}_2. \end{aligned}$$

We have $\Lambda = \mathbb{Z}\mathbf{v}_1 \oplus \mathbb{Z}\mathbf{v}_2$, $\Lambda^* = \mathbb{Z}\mathbf{k}_1 \oplus \mathbb{Z}\mathbf{k}_2$ and $\mathbf{k}_{\ell} \cdot \mathbf{v}_{\ell'} = 2\pi\delta_{\ell\ell'}$.

- (8) \mathbb{H} , Honeycomb structure; see (1.8).
- (9) \mathbb{H}_{\sharp} , Zigzag-truncated honeycomb structure; see (1.11).
- (10) $\Sigma = \mathbb{R}^2/\mathbb{Z}\mathbf{v}_2$, the cylinder with Ω_{Σ} , a choice of fundamental cell for Σ ; see Figure 1.

(11) $L_{k_{\parallel}}^2 = L_{k_{\parallel}}^2(\Sigma)$, functions f such that $f(\mathbf{x} + \mathbf{v}_2) = e^{ik_{\parallel}} f(\mathbf{x})$ for almost all \mathbf{x} , and

$$\|f\|_{L_{k_{\parallel}}^2}^2 = \int_{\Omega_{\Sigma}} |f|^2 < \infty.$$

(12) $\mathcal{H}^{(\omega)} \equiv L^2(\mathbb{R}^2; e^{\gamma|\mathbf{x}-\omega|} d\mathbf{x})$, exponentially weighted L^2 space.

(13) $\mathcal{B}(X)$ denotes the space of bounded linear operators on X .

(14) $G_{\lambda}^{\text{free}}(\mathbf{x}, \mathbf{y})$ denotes the free Green's function defined in (10.3).

(15) $G_{\lambda}^{\text{atom}}(\mathbf{x}, \mathbf{y})$ denotes the atomic Green's function defined in (10.7).

(16) Hamiltonians:

$H_{\text{atom}}^{\lambda} = -\Delta + \lambda^2 V_0(\mathbf{x})$, the atomic Hamiltonian with ground state energy E_0^{λ}

$H_{\text{bulk}}^{\lambda} = -\Delta + \lambda^2 V(\mathbf{x})$ and $H_{\text{edge}}^{\lambda} = -\Delta + \lambda^2 V_{\sharp}(\mathbf{x})$, denote bulk and edge Hamiltonians

acting in $L^2(\mathbb{R}^2)$

$H_{\sharp}^{\lambda} = H_{\text{edge}}^{\lambda} - E_0^{\lambda}$, the centered edge Hamiltonian, acting in $L_{k_{\parallel}}^2$

$\tilde{H}_{\sharp}^{\lambda} = (\rho\lambda)^{-1} H_{\sharp}^{\lambda}$, the scaled and centered edge Hamiltonian acting in $L_{k_{\parallel}}^2$

$H_{\sharp}^{\text{TB}}(k_{\parallel})$, the tight-binding edge Hamiltonian, acting in $l^2(\mathbb{N}_0; \mathbb{C}^2)$; see Definition 2.6.

Acknowledgements: The authors wish to thank Gian Michele Graf and Alexis Drouot for very stimulating discussions. We would also like to thank Bernard Helffer for correspondence concerning previous general results on tight-binding limits. Part of this research was done while MIW was Bergman Visiting Professor at Stanford University. CLF and MIW wish to thank the Department of Mathematics at Stanford University for its hospitality. This research was supported in part by National Science Foundation grants DMS-1265524 (CLF) and DMS-1412560, DMS-1620418 and Simons Foundation Math + X Investigator Award #376319 (MIW).

2. TIGHT-BINDING

Consider a tiling of the entire plane, \mathbb{R}^2 , by parallelograms of the sort shown in Figure 1. Each parallelogram has exactly two points of \mathbb{H} . This is a particular *dimerization* of \mathbb{H} . We assign the label (n_1, n_2) to the parallelogram which contains $\mathbf{v}_A^{(n_1, n_2)} = \mathbf{v}_A + n_1 \mathbf{v}_1 + n_2 \mathbf{v}_2$ and $\mathbf{v}_B^{(n_1, n_2)} = \mathbf{v}_B + n_1 \mathbf{v}_1 + n_2 \mathbf{v}_2$. To the sites $\mathbf{v}_A^{(n_1, n_2)}$ and $\mathbf{v}_B^{(n_1, n_2)}$ we assign complex amplitudes ψ_{n_1, n_2}^A and ψ_{n_1, n_2}^B and form the tight binding wave function:

$$\psi_{n_1, n_2} = \begin{pmatrix} \psi_{n_1, n_2}^A \\ \psi_{n_1, n_2}^B \end{pmatrix}$$

2.1. $H_{\text{bulk}}^{\text{TB}}$, **the tight-binding bulk Hamiltonian.** The bulk tight binding Hamiltonian can be represented with respect to the above dimerization. Starting with any dimerization would give a unitarily equivalent operator on $l^2(\mathbb{Z}^2; \mathbb{C}^2)$. The nearest neighbor tight binding bulk Hamiltonian, relative to the dimerization of \mathbb{H} in Figure 1 is:

$$(2.1) \quad \left[H_{\text{bulk}}^{\text{TB}} \psi \right]_{n_1, n_2} = \begin{pmatrix} \left(H_{\text{bulk}}^{\text{TB}} \psi \right)_A^{n_1, n_2} \\ \left(H_{\text{bulk}}^{\text{TB}} \psi \right)_B^{n_1, n_2} \end{pmatrix} = \begin{pmatrix} \psi_{n_1-1, n_2}^B + \psi_{n_1, n_2-1}^B + \psi_{n_1, n_2}^B \\ \psi_{n_1+1, n_2}^A + \psi_{n_1, n_2+1}^A + \psi_{n_1, n_2}^A \end{pmatrix}$$

where $n_1, n_2 \in \mathbb{Z}$. The operator $H_{\text{bulk}}^{\text{TB}}$ is a bounded self-adjoint linear operator on $l^2(\mathbb{Z}^2; \mathbb{C}^2)$ and was introduced in [66]. The spectrum of $H_{\text{bulk}}^{\text{TB}}$ consists of two spectral bands which touch conically at Dirac points over the vertices of \mathcal{B} . The approximation and convergence as λ increases of the low-lying dispersion surfaces and the resolvent H_{bulk}^λ acting on $L^2(\mathbb{R}^2)$ to those of $H_{\text{bulk}}^{\text{TB}}$ acting on $l^2(\mathbb{Z}^2; \mathbb{C}^2)$ was studied in [26].

2.2. Tight-binding Hamiltonian for the zigzag edge.

Our goal in this section is to introduce a tight-binding edge Hamiltonian which will act on functions $\psi \in l^2((\mathbb{N}_0 \times \mathbb{Z}); \mathbb{C}^2)$ defined on the vertices of $\mathbb{H}_\#$. We shall do this by first expressing $H_{\text{bulk}}^{\text{TB}}$, as a direct integral over k_\parallel of fiber operators $H_{\text{bulk}}^{\text{TB}}(k_\parallel)$ acting on states which are “ k_\parallel -pseudo-periodic” with respect to one lattice direction and square-summable with respect to the other lattice direction. The edge Hamiltonian $H_\#^{\text{TB}}$ is then obtained from $H_{\text{bulk}}^{\text{TB}}(k_\parallel)$ by appropriate restriction to functions defined on $\mathbb{H}_\#$.

Since the truncated structure $\mathbb{H}_\#$ and its subset edge vertices are invariant with respect to translation by \mathbf{v}_2 , we introduce $k_\parallel \in S^1 = \mathbb{R}/2\pi\mathbb{Z}$, the parallel quasi-momentum associated with this translation invariance. For each $k_\parallel \in [0, 2\pi]$, we refer to a state as being k_\parallel -pseudo-periodic if:

$$(2.2) \quad \psi_{n_1, n_2+1} = e^{ik_\parallel} \psi_{n_1, n_2}, \quad n_1 \geq 0, \quad n_2 \in \mathbb{Z}.$$

Functions $\psi = \{\psi_{n_1, n_2}\} \in l^2(\mathbb{Z}; \mathbb{C}^2)$ may be expressed via the discrete Fourier transform as

$$(2.3) \quad \psi_{n_1, n_2} = (2\pi)^{-1} \int_0^{2\pi} e^{in_2 k_\parallel} \psi_{n_1}(k_\parallel) dk_\parallel,$$

as a superposition over states $\{e^{in_2 k_\parallel} \psi_{n_1}(k_\parallel)\}$ which are square-summable over \mathbb{Z} with respect to n_1 and which satisfy (2.2).

Therefore, the tight binding bulk Hamiltonian $H_{\text{bulk}}^{\text{TB}}$ may be reduced to the k_\parallel -dependent fiber (Bloch) Hamiltonians, $H_{\text{bulk}}^{\text{TB}}(k_\parallel) : l^2(\mathbb{Z}; \mathbb{C}^2) \rightarrow l^2(\mathbb{Z}; \mathbb{C}^2)$, defined by

$$(2.4) \quad \begin{aligned} \left[H_{\text{bulk}}^{\text{TB}}(k_\parallel) \psi \right]_{n_1} &\equiv \begin{pmatrix} \psi_{n_1-1}^B + (1 + e^{-ik_\parallel}) \psi_{n_1}^B \\ \psi_{n_1+1}^A + (1 + e^{+ik_\parallel}) \psi_{n_1}^A \end{pmatrix}, \\ &= \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \psi_{n_1-1}^A \\ \psi_{n_1-1}^B \end{pmatrix} + \begin{pmatrix} 0 & 1 + e^{-ik_\parallel} \\ 1 + e^{+ik_\parallel} & 0 \end{pmatrix} \begin{pmatrix} \psi_{n_1}^A \\ \psi_{n_1}^B \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \psi_{n_1+1}^A \\ \psi_{n_1+1}^B \end{pmatrix}. \end{aligned}$$

Finally, we define the tight-binding edge Hamiltonian, $H_\#^{\text{TB}}$. For $\psi = (\psi_0, \psi_1, \psi_2, \dots) \in l^2(\mathbb{N}_0; \mathbb{C}^2)$, introduce the extension operator:

$$\begin{aligned} \iota &: l^2(\mathbb{N}_0; \mathbb{C}^2) \rightarrow l^2(\mathbb{Z}; \mathbb{C}^2) \\ \iota \psi &= (\dots, 0, 0, 0, \psi_0, \psi_1, \psi_2, \dots) \in l^2(\mathbb{Z}; \mathbb{C}^2). \end{aligned}$$

The adjoint of ι is the restriction operator defined on $\phi = (\dots, \phi_{-2}, \phi_{-1}, \phi_0, \phi_1, \phi_2, \dots) \in l^2(\mathbb{Z}; \mathbb{C}^2)$ by:

$$\begin{aligned} \iota^* &: l^2(\mathbb{Z}; \mathbb{C}^2) \rightarrow l^2(\mathbb{N}_0; \mathbb{C}^2), \\ \iota^* \phi &= (\phi_0, \phi_1, \phi_2, \dots) \in l^2(\mathbb{N}_0; \mathbb{C}^2). \end{aligned}$$

Definition 2.1. The tight-binding edge fiber operators, $H_{\#}^{\text{TB}}(k_{\parallel})$, and edge Hamiltonian $H_{\#}^{\text{TB}}$ are given by

$$(2.5) \quad H_{\#}^{\text{TB}}(k_{\parallel}) = \iota^* H_{\text{bulk}}^{\text{TB}}(k_{\parallel}) \iota : l^2(\mathbb{N}_0; \mathbb{C}^2) \rightarrow l^2(\mathbb{N}_0; \mathbb{C}^2)$$

and

$$(2.6) \quad H_{\#}^{\text{TB}} = \int_{[0, 2\pi]}^{\oplus} H_{\#}^{\text{TB}}(k_{\parallel}) dk_{\parallel} : l^2(\mathbb{N}_0 \times \mathbb{Z}) \rightarrow l^2(\mathbb{N}_0 \times \mathbb{Z}) .$$

2.3. Spectrum of $H_{\#}^{\text{TB}}(k_{\parallel})$. Define, for $k_{\parallel} \in [0, 2\pi]$, the functions

$$(2.7) \quad \zeta(k_{\parallel}) \equiv 1 + e^{ik_{\parallel}} \quad ,$$

$$(2.8) \quad \delta_{\text{gap}}(k_{\parallel}) \equiv \min_{k_{\perp} \in [0, 2\pi]} \left| 1 + e^{ik_{\parallel}} + e^{ik_{\perp}} \right| = \left| 1 - |\zeta(k_{\parallel})| \right| ,$$

$$(2.9) \quad \delta_{\text{max}}(k_{\parallel}) \equiv 1 + |\zeta(k_{\parallel})| .$$

Note $\delta_{\text{gap}}(2\pi/3) = \delta_{\text{gap}}(4\pi/3) = 0$, $\delta_{\text{gap}}(k_{\parallel}) > 0$ otherwise in $[0, 2\pi]$, and that $|\zeta(k_{\parallel})| < 1$ for $k_{\parallel} \in (2\pi/3, 4\pi/3)$. We next prove that the spectrum of $H_{\#}^{\text{TB}}(k_{\parallel})$ is as displayed in Figure

2. Let us enumerate the coordinates of the vector in $l^2(\mathbb{N}_0; \mathbb{C}^2)$, $\psi = \left\{ \begin{pmatrix} \psi_n^A \\ \psi_n^B \end{pmatrix} \right\}_{n \geq 0}$, by $\psi = (\psi_0^A, \psi_0^B, \psi_1^A, \psi_1^B, \dots)^{\top}$. We denote the corresponding unit vectors by $\hat{\mathbf{e}}_1 = (1, 0, 0, \dots)$, $\hat{\mathbf{e}}_2 = (0, 1, 0, \dots)$, etc.

Theorem 2.2 ($\sigma(H_{\#}^{\text{TB}}(k_{\parallel}))$, the spectrum of $H_{\#}^{\text{TB}}(k_{\parallel})$ in $l^2(\mathbb{N}_0; \mathbb{C}^2)$).

For each $k_{\parallel} \in [0, 2\pi]$, $\sigma(H_{\#}^{\text{TB}}(k_{\parallel})) = \sigma_{\text{pt}}(\sigma(H_{\#}^{\text{TB}}(k_{\parallel}))) \cup \sigma_{\text{ess}}(\sigma(H_{\#}^{\text{TB}}(k_{\parallel})))$.

(1) Point spectrum of $H_{\#}^{\text{TB}}(k_{\parallel})$:

$$\sigma_{\text{pt}}(H_{\#}^{\text{TB}}(k_{\parallel})) = \begin{cases} \{0\} & \text{if } k_{\parallel} \in (2\pi/3, 4\pi/3) \\ \{-1, 0, 1\} & \text{if } k_{\parallel} = \pi \\ \emptyset & \text{if } k_{\parallel} \in [0, 2\pi] \setminus (2\pi/3, 4\pi/3) : \end{cases}$$

In particular,

$H_{\#}^{\text{TB}}$ has a zero energy “flat-band” of eigenstates over the range $2\pi/3 < k_{\parallel} < 4\pi/3$.

For $k_{\parallel} \in (2\pi/3, 4\pi/3) \setminus \{\pi\}$ the point spectrum, which consists eigenvalue $E = 0$ is simple. The corresponding normalized 0– energy eigenstate, $\psi^{\text{TB, bd}} = \{\psi_n^{\text{TB, bd}}\}_{n \geq 0}$, is given by

$$(2.10) \quad \psi_n^{\text{TB, bd}}(k_{\parallel}) = \sqrt{1 - |\zeta(k_{\parallel})|^2} \begin{pmatrix} (-\zeta(k_{\parallel}))^n \\ 0 \end{pmatrix}, \quad n \geq 0 .$$

For $k_{\parallel} = \pi$, $E = 0$ is a simple eigenvalue with corresponding normalized 0– energy eigenstate is given by:

$$(2.11) \quad \psi_0^{\text{TB, bd}}(\pi) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \psi_n^{\text{TB, bd}}(\pi) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad n \geq 1 .$$

The eigenvalues $E = +1$ and $E = -1$ have infinite multiplicity. The corresponding eigenspaces are:

$$\begin{aligned} \text{kernel}(H_{\#}^{\text{TB}}(\pi) - Id) &= \left\{ \frac{1}{\sqrt{2}} (\hat{\mathbf{e}}_{2j+1} + \hat{\mathbf{e}}_{2j+2}) : j = 0, 1, 2, \dots \right\}, \\ \text{kernel}(H_{\#}^{\text{TB}}(\pi) + Id) &= \left\{ \frac{1}{\sqrt{2}} (\hat{\mathbf{e}}_{2j+1} - \hat{\mathbf{e}}_{2j+2}) : j = 0, 1, 2, \dots \right\} \end{aligned}$$

(2) Essential spectrum of $H_{\#}^{\text{TB}}(k_{\parallel})$:

$$(2.12) \quad \sigma_{\text{ess}}(H_{\#}^{\text{TB}}(k_{\parallel})) = \begin{cases} \left\{ z \in \mathbb{R} : \delta_{\text{gap}}(k_{\parallel}) \leq |z| \leq \delta_{\text{max}}(k_{\parallel}) \right\}, & k_{\parallel} \in [0, 2\pi] \setminus \{\pi\} \\ \emptyset, & k_{\parallel} = \pi. \end{cases}$$

(3) Resolvent expansion:

(a) Let $k_{\parallel} \in (2\pi/3, 4\pi/3) \setminus \{\pi\}$. Then, for $z \in \mathbb{C} \setminus \sigma_{\text{ess}}(H_{\#}^{\text{TB}}(k_{\parallel}))$ and $z \neq 0$ we have

$$(2.13) \quad \left(H_{\#}^{\text{TB}}(k_{\parallel}) - zI \right)^{-1} f = \frac{1}{z} \langle \psi^{\text{TB,bd}}(k_{\parallel}), f \rangle_{l^2(\mathbb{N}_0; \mathbb{C}^2)} \psi^{\text{TB,bd}}(k_{\parallel}) + \mathcal{G}_{\text{reg}}(z; k_{\parallel}) f.$$

Here, $z \mapsto \mathcal{G}_{\text{reg}}(z; k_{\parallel})$ is an analytic mapping from $\mathbb{C} \setminus \sigma_{\text{ess}}(H_{\#}^{\text{TB}}(k_{\parallel}))$ to the space of bounded linear operators on $l^2(\mathbb{N}_0; \mathbb{C}^2)$. If (z, k_{\parallel}) varies over a compact set $\Upsilon \subset \mathbb{R} \times [0, 2\pi]$ for which $\text{distance}(z, \sigma_{\text{ess}}(H_{\#}^{\text{TB}}(k_{\parallel}))) \geq b > 0$, where b is a positive constant depending on Υ , then $\|\mathcal{G}_{\text{reg}}(z; k_{\parallel})\|_{\mathcal{B}(l^2(\mathbb{N}_0; \mathbb{C}^2))} < B(b) < \infty$.

(b) Let $k_{\parallel} = \pi$. Then, $\left(H_{\#}^{\text{TB}}(k_{\parallel}) - zI \right)^{-1} f$ has an expression analogous to (2.13) with poles at $z = 0$, $z = +1$ and $z = -1$.

(c) Let $k_{\parallel} \in [0, 2\pi] \setminus (2\pi/3, 4\pi/3)$. Then, for $z \in \mathbb{C} \setminus \sigma_{\text{ess}}(H_{\#}^{\text{TB}}(k_{\parallel}))$ we have

$$(2.14) \quad \left(H_{\#}^{\text{TB}}(k_{\parallel}) - zI \right)^{-1} f = \mathcal{G}_{\text{reg}}(z; k_{\parallel}) f,$$

where $z \mapsto \mathcal{G}_{\text{reg}}(z; k_{\parallel})$ is as in part (a).

(4) For $k_{\parallel} \in (2\pi/3, 4\pi/3)$, the equation $H_{\#}^{\text{TB}}(k_{\parallel})\psi = f$, where $f \in l^2(\mathbb{N}_0; \mathbb{C}^2)$, is solvable for $\psi \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ if and only if $\langle \psi^{\text{TB,bd}}(k_{\parallel}), f \rangle_{l^2(\mathbb{N}_0; \mathbb{C}^2)} = 0$.

Remark 2.3. We remark on the connection between the condition $k_{\parallel} \in (2\pi/3, 4\pi/3)$ (equivalently $|\zeta(k_{\parallel})| < 1$) and the non-vanishing of a winding number, known as the *Zak phase*. For fixed k_{\parallel} , consider the normalized bulk Floquet-Bloch modes of $H_{\text{bulk}}^{\text{TB}}(k_{\parallel})$; see (2.4). There are two families of eigenpairs: $(\mu^{\pm}(k_{\parallel}), U_{n_1}^{\pm}(k_{\perp}; k_{\parallel}))$, where

$$\begin{aligned} \mu^{\pm}(k_{\parallel}) &= \pm |\zeta(k_{\parallel}) + e^{ik_{\perp}}|, & (\zeta(k_{\parallel}) &= 1 + e^{ik_{\parallel}}), \\ U_{n_1}^{\pm}(k_{\perp}; k_{\parallel}) &= e^{ik_{\perp} n_1} \xi^{\pm}(k_{\perp}; k_{\parallel}), & \xi^{\pm}(k_{\perp}; k_{\parallel}) &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ \pm j(k_{\perp}) \end{pmatrix}, \\ j(e^{ik_{\perp}}) &= \frac{\zeta(k_{\parallel}) + e^{ik_{\perp}}}{|\zeta(k_{\parallel}) + e^{ik_{\perp}}|}, & j(z)\overline{j(z)} &= 1. \end{aligned}$$

For either family of modes (say +), we consider the *Berry connection* defined by $A(k_\perp; k_\parallel) \equiv \langle \xi(k_\perp; k_\parallel), \frac{1}{i} \partial_{k_\perp} \xi(k_\perp; k_\parallel) \rangle$ and the *Zak phase* by $\mathcal{Z}(k_\parallel) \equiv \int_0^{2\pi} A(k_\perp; k_\parallel) dk_\perp$. We have

$$\begin{aligned} \mathcal{Z}(k_\parallel) &= -i \int_0^{2\pi} \overline{j(e^{ik_\perp}; k_\parallel)} \frac{\partial}{\partial k_\perp} j(e^{ik_\perp}; k_\parallel) dk_\perp \\ &= -i \int_{|w|=1} \overline{j(w; k_\parallel)} \partial_z j(w; k_\parallel) dw \\ &= -i \int_{|w|=1} \frac{\partial_w j(w; k_\parallel)}{j(w; k_\parallel)} dw \\ &= 2\pi \times \text{Winding number of } w \in S^1 \mapsto j(w; k_\parallel) \in \mathbb{C} . \end{aligned}$$

If $|\zeta(k_\parallel)| < 1$, then $\mathcal{Z}(k_\parallel) = 2\pi$ and if $|\zeta(k_\parallel)| > 1$, then $\mathcal{Z}(k_\parallel) = 0$. This is an example of the *bulk-edge correspondence* (see, for example, [15, 28, 47]) and Theorem 1.2 establishes its validity in the strong-binding regime.

Proof of Theorem 2.2: Fix $k_\parallel \in [0, 2\pi)$ and set $\zeta = \zeta(k_\parallel) = 1 + e^{ik_\parallel}$. We study the operator $H_\#^{\text{TB}}(k_\parallel)$ in the Hilbert space $l^2(\mathbb{N}_0; \mathbb{C}^2)$. An energy z is in the point spectrum of $H_\#^{\text{TB}}(k_\parallel)$ if there exists $\psi \neq 0$, $\psi \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ such that $H_\#^{\text{TB}}(k_\parallel)\psi = z\psi$. Written out componentwise, the eigenvalue problem is:

$$(2.15) \quad \psi_{n-1}^B + \zeta^* \psi_n^B = z \psi_n^A, \quad n \geq 0,$$

$$(2.16) \quad \psi_{n+1}^A + \zeta \psi_n^A = z \psi_n^B, \quad n \geq 0,$$

and $\psi_n = \begin{pmatrix} \psi_n^A \\ \psi_n^B \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ for all $n \leq -1$.

We begin by showing that for $k_\parallel \in (2\pi/3, 4\pi/3)$, we have that $0 \in \sigma_{\text{pt}}(H_\#^{\text{TB}}(k_\parallel))$ and that for $k_\parallel \in [0, 2\pi] \setminus (2\pi/3, 4\pi/3)$, $z = 0$ is not in the point spectrum. Set $E = 0$ and observe that equations (2.15) and (2.16) become decoupled first order difference equations: $\psi_{n+1}^A = (-\zeta)\psi_n^A$, $n \geq 0$ and $\psi_{n-1}^B = (-\zeta^*)\psi_n^B$, $n \geq 0$.

The equation for ψ^A has the solution: $\psi_n^A = (-\zeta)^n \psi_0^A$, $n \geq 0$, where ψ_0^A can be set arbitrarily. If $k_\parallel \in (2\pi/3, 4\pi/3)$, then $|\zeta(k_\parallel)| < 1$ and hence $\psi_n^A \rightarrow 0$ exponentially as $n \rightarrow \infty$. Turning to ψ^B , let us first assume that $k_\parallel \neq \pi$ so that $\zeta(k_\parallel) \neq 0$. In this case, $\psi_n^B = (-\zeta^*)^{-1} \psi_{n-1}^B$, $n \geq 0$. Since $\psi_{-1}^B = 0$, we have $\psi_n^B = 0$ for all $n \geq 0$. If $k_\parallel = \pi$ then we have from (2.15) that $\psi_{n-1}^B = 0$ for all $n \geq 0$.

Now suppose $k_\parallel \in [0, 2\pi] \setminus (2\pi/3, 4\pi/3)$. Then, the above discussion also implies that if $\psi \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ solves the eigenvalue equation with $z = 0$, then $\psi \equiv 0$.

We conclude: $E = 0$ is a point eigenvalue of $H_\#^{\text{TB}}(k_\parallel)$ acting in $l^2(\mathbb{N}_0; \mathbb{C}^2)$ if and only if $k_\parallel \in (2\pi/3, 4\pi/3)$. For $k_\parallel \in (2\pi/3, 4\pi/3) \setminus \{\pi\}$, the $l^2(\mathbb{N}_0; \mathbb{C}^2)$ -normalized eigenstate is given by:

$$(2.17) \quad \psi_n^{\text{TB,bd}}(k_\parallel) = \sqrt{1 - |\zeta(k_\parallel)|^2} \begin{pmatrix} (-\zeta(k_\parallel))^n \\ 0 \end{pmatrix}, \quad n \geq 0$$

$$(2.18) \quad \zeta(k_\parallel) \equiv 1 + e^{ik_\parallel} .$$

For $k_{\parallel} = \pi$ ($\zeta(k_{\parallel}) = 0$), the eigenstate is given by the expression:

$$(2.19) \quad \psi_0^{\text{TB,bd}}(\pi) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \psi_n^{\text{TB,bd}}(\pi) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad n \geq 1,$$

and is supported strictly at the edge.

We now assume that z is complex and $z \neq 0$, and explore the invertibility of $H_{\sharp}^{\text{TB}}(k_{\parallel}) - zI$ on $l^2(\mathbb{N}_0; \mathbb{C}^2)$. Written out componentwise, the system $(H_{\sharp}^{\text{TB}}(k_{\parallel}) - zI)\psi = f$, where $f \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ is:

$$(2.20) \quad \psi_{n-1}^B + \zeta^* \psi_n^B = z\psi_n^A + f_n^A, \quad n \geq 0$$

$$(2.21) \quad \psi_{n+1}^A + \zeta \psi_n^A = z\psi_n^B + f_n^B, \quad n \geq 0,$$

$$(2.22) \quad \psi_n = \begin{pmatrix} \psi_n^A \\ \psi_n^B \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad f_n = \begin{pmatrix} f_n^A \\ f_n^B \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad \text{for all } n \leq -1$$

$$(2.23) \quad \text{and } |\psi_n| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

We focus on the case $k_{\parallel} \in [0, 2\pi] \setminus \{\pi\}$, so that $\zeta(k_{\parallel}) = 1 + e^{ik_{\parallel}} \neq 0$.

Remark 2.4. For $k_{\parallel} = \pi$, the system (2.22) is of the form $(H_{\sharp}^{\text{TB}}(\pi) - z)\psi = f$, where $\psi = (\psi_0^A, \psi_0^B, \psi_1^A, \psi_1^B, \dots)^{\top}$, $f = (f_0^A, f_0^B, f_1^A, f_1^B, \dots)^{\top}$ and $H_{\sharp}^{\text{TB}}(\pi)$ is a block-diagonal matrix consisting of a 1×1 block, 0 in the (1, 1) entry, followed by an infinite sequence of identical 2×2 blocks, each equal to $\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, filling out the diagonal. The statements in Theorem 2.2 on the spectrum of $H_{\sharp}^{\text{TB}}(\pi)$ and the mapping $z \mapsto (H_{\sharp}^{\text{TB}}(\pi) - z)^{-1}$ are easily verified.

For $k_{\parallel} \neq \pi$, we next rewrite (2.20)-(2.21) as a first order recursion. Consider (2.20) with n replaced by $n + 1$:

$$(2.24) \quad \psi_n^B + \zeta^* \psi_{n+1}^B = z\psi_{n+1}^A + f_{n+1}^A, \quad n \geq -1.$$

For $n = -1$, equation (2.24) implies the boundary condition at site $n = 0$:

$$(2.25) \quad \zeta^* \psi_0^B - z\psi_0^A = f_0^A.$$

For $n \geq 0$, we use $\zeta \neq 0$ and (2.21) in (2.24) and obtain:

$$(2.26) \quad \psi_{n+1}^B = \left(-\frac{\zeta}{\zeta^*} \right) z \psi_n^A + \frac{z^2 - 1}{\zeta^*} \psi_n^B + \frac{z}{\zeta^*} f_n^B + \frac{1}{\zeta^*} f_{n+1}^A, \quad n \geq 0$$

Summarizing, we have that the system: (2.20), (2.21) and (2.22) is equivalent to the first order system (2.21), (2.26) for $\psi_n = \begin{pmatrix} \psi_n^A \\ \psi_n^B \end{pmatrix}$, $n \geq 0$, with the boundary condition (2.25) at $n = 0$. We write this more compactly as:

$$(2.27) \quad \psi_{n+1} = M(z, \zeta) \psi_n + F_n(z, \zeta), \quad n \geq 0,$$

$$(2.28) \quad \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^{\top} \psi_0 \equiv \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^{\top} \begin{pmatrix} \psi_0^A \\ \psi_0^B \end{pmatrix} = f_0^A,$$

$$(2.29) \quad |\psi_m| \rightarrow 0, \quad m \rightarrow \infty.$$

where

$$(2.30) \quad M(z, \zeta) = \begin{pmatrix} -\zeta & z \\ -\frac{\zeta}{\zeta^*} z & \frac{z^2-1}{\zeta^*} \end{pmatrix},$$

$$(2.31) \quad F_n(z, \zeta; f) = \begin{pmatrix} f_n^B & \\ \frac{z}{\zeta^*} f_n^B & + \frac{1}{\zeta^*} f_{n+1}^A \end{pmatrix}, \quad n \geq 0.$$

We next solve (2.27)-(2.28) by diagonalizing the matrix $M(z, \zeta)$.

The eigenvalues λ of $M(z, \zeta)$ are solutions of the quadratic equation

$$(2.32) \quad \zeta^* \lambda^2 + (1 + |\zeta|^2 - z^2) \lambda + \zeta = 0,$$

whose solutions are:

$$(2.33) \quad \lambda_1(z, \zeta) = \frac{-(1 + |\zeta|^2 - z^2) + \sqrt{(1 + |\zeta|^2 - z^2)^2 - 4|\zeta|^2}}{2\zeta^*}$$

$$(2.34) \quad \lambda_2(z, \zeta) = \frac{-(1 + |\zeta|^2 - z^2) - \sqrt{(1 + |\zeta|^2 - z^2)^2 - 4|\zeta|^2}}{2\zeta^*}.$$

When convenient, we suppress the dependence of λ_1 and λ_2 on ζ and E and occasionally write λ_j or $\lambda_j(z)$. These expressions depend on k_{\parallel} through $\zeta(k_{\parallel}) = 1 + e^{ik_{\parallel}}$.

Note that $|\lambda_1 \lambda_2| = |\det M(z, \zeta)| = |\zeta/\zeta^*| = 1$ and hence $M(z, \zeta)$ may have at most one eigenvalue strictly inside the unit circle in \mathbb{C} .

Recall the definitions: $\delta_{\text{gap}}(k_{\parallel}) \equiv \left| 1 - |\zeta(k_{\parallel})| \right|$ and $\delta_{\text{max}}(k_{\parallel}) \equiv 1 + |\zeta(k_{\parallel})|$.

Remark 2.5. We shall see just below that for fixed $k_{\parallel} \neq 2\pi/3, \pi$ or $4\pi/3$: if (a) $|z| < \delta_{\text{gap}}(k_{\parallel})$ or (b) $|z| > \delta_{\text{max}}(k_{\parallel})$ then the discriminant in (2.33)-(2.34), $(1 + |\zeta(k_{\parallel})|^2 - z^2)^2 - 4|\zeta(k_{\parallel})|^2$, is strictly positive and uniformly bounded away from zero. Therefore, in each of these cases the expressions in (2.33)-(2.34) define single-valued functions $\lambda_1(z, \zeta)$ and $\lambda_2(z, \zeta)$. This property continues to hold for $k_{\parallel} \in \mathcal{J}_1 \subset \subset [0, 2\pi] \setminus \{2\pi/3, \pi, 4\pi/3\}$ and either (a') $|\Re z| < \delta_{\text{gap}}(k_{\parallel})$ and $|\Im z| < \eta(\mathcal{J}_1)$ or (b') $|\Re z| > \delta_{\text{max}}(k_{\parallel})$ and $|\Im z| < \eta(\mathcal{J}_1)$, for some $\eta(\mathcal{J}_1) > 0$ chosen sufficiently small. In the case where z is real and $\delta_{\text{gap}}(k_{\parallel}) \leq |z| \leq \delta_{\text{max}}(k_{\parallel})$ the discriminant is nonpositive and we do not distinguish between the roots of (2.32); they comprise a two element set on the unit circle in \mathbb{C} .

Lemma 2.6. *Assume $0 < |\zeta(k_{\parallel})| \neq 1$, i.e. $k_{\parallel} \neq 2\pi/3, \pi$ or $4\pi/3$. Then, the following hold:*

(1) *Let $z \in \mathbb{R}$ and assume that either*

$$(2.35) \quad |z| < \delta_{\text{gap}}(k_{\parallel}) \text{ or } |z| > \delta_{\text{max}}(k_{\parallel}).$$

Then, $M(z, \zeta(k_{\parallel}))$ has one eigenvalue inside the unit circle and one eigenvalue outside the unit circle.

(2) *Let $\lambda_1(z)$ and $\lambda_2(z)$ denote be the expressions for the eigenvalues of $M(z, \zeta(k_{\parallel}))$ displayed in (2.33)-(2.34).*

(i) *If $z \in \mathbb{R}$ and $|z| < \delta_{\text{gap}}(k_{\parallel})$, then $|\lambda_1(z; k_{\parallel})| < 1 < |\lambda_2(z; k_{\parallel})|$.*

(ii) *If $z \in \mathbb{R}$ and $|z| > \delta_{\text{max}}(k_{\parallel})$, then $|\lambda_2(z; k_{\parallel})| < 1 < |\lambda_1(z; k_{\parallel})|$.*

(iii) *If $z \in \mathbb{R}$ and $\delta_{\text{gap}}(k_{\parallel}) \leq |z| \leq \delta_{\text{max}}(k_{\parallel})$, then equation (2.32) has two roots, λ , satisfying $|\lambda| = 1$.*

(3) Let \mathcal{J}_1 denote a compact subset of $[0, 2\pi] \setminus \{2\pi/3, \pi, 4\pi/3\}$. There exists a constant $\eta > 0$, which depends on \mathcal{J}_1 , such that for all $k_{\parallel} \in \mathcal{J}_1$ the following hold:

(a) If z is in the complex open neighborhood

$$(2.36) \quad \mathcal{O}_0(k_{\parallel}) : \quad |\Re z| < \delta_{\text{gap}}(k_{\parallel}) \quad \text{and} \quad |\Im z| < \eta(\mathcal{J}_1),$$

then (2.35) holds. Moreover, $\lambda_1(z, \zeta)$ and $\lambda_2(z, \zeta)$ satisfy the strict inequalities of (2.i), and their magnitudes are uniformly bounded away from 1, provided z remains in a compact subset of $\mathcal{O}_0(k_{\parallel})$.

(b) If z is in the complex open neighborhood

$$(2.37) \quad \mathcal{O}_+(k_{\parallel}) : \quad |\Re z| > \delta_{\text{max}}(k_{\parallel}) \quad \text{and} \quad |\Im z| < \eta(\mathcal{J}_1),$$

then (2.35) holds and moreover $\lambda_1(z, \zeta)$ and $\lambda_2(z, \zeta)$ satisfy the inequalities of (2.ii) and their magnitudes are uniformly bounded away from 1, provided z remains in a compact subset of $\mathcal{O}_+(k_{\parallel})$.

Proof of Lemma 2.6: Part 3 of the Lemma follows from parts (1) and (2) and the expressions (2.33), (2.34) for $\lambda_1(z; k_{\parallel})$, and $\lambda_2(z; k_{\parallel})$. We now proceed with the proof of assertions (1) and (2), which assume $z \in \mathbb{R}$.

We consider the two cases delineated by the sign of the discriminant:

Case 1: $(1 + |\zeta|^2 - z^2)^2 - 4|\zeta|^2 > 0$ and **Case 2:** $(1 + |\zeta|^2 - z^2)^2 - 4|\zeta|^2 \leq 0$.

Case 1: In this case, $|1 + |\zeta|^2 - z^2| > 2|\zeta|$. There are two subcases:

(1a) $1 + |\zeta|^2 - z^2 > 2|\zeta|$ and (1b) $z^2 - 1 - |\zeta|^2 > 2|\zeta|$.

In subcase (1a), we have $z^2 < (1 - |\zeta|)^2$ and therefore $|z| < \delta_{\text{gap}}(k_{\parallel}) = |1 - |\zeta||$, where $\delta_{\text{gap}}(k_{\parallel}) > 0$ since $k_{\parallel} \neq 2\pi/3, 4\pi/3$. In this subcase we also have: $-(1 + |\zeta|^2 - z^2) < -2|\zeta| < 0$. Therefore,

$$\begin{aligned} 0 > (2\zeta^*)\lambda_1 &= -(1 + |\zeta|^2 - z^2) + \sqrt{(1 + |\zeta|^2 - z^2)^2 - 4|\zeta|^2} \\ &> -(1 + |\zeta|^2 - z^2) - \sqrt{(1 + |\zeta|^2 - z^2)^2 - 4|\zeta|^2} = (2\zeta^*)\lambda_2. \end{aligned}$$

Let $\lambda_1 = r_1/(2\zeta^*)$ and $\lambda_2 = r_2/(2\zeta^*)$. Therefore, $|r_1| = |(2\zeta^*)\lambda_1| < |(2\zeta^*)\lambda_2| = |r_2|$. Therefore, $|\lambda_1|/|\lambda_2| = |r_1|/|r_2| < 1$. Since $|\lambda_1| |\lambda_2| = 1$,

$$(2.38) \quad \text{in subcase (1a), we have } |z| < \delta_{\text{gap}}(k_{\parallel}), \text{ and } |\lambda_1(z)| < 1 < |\lambda_2(z)|.$$

In subcase (1b) we have $|z| > 1 + |\zeta(k_{\parallel})| = \delta_{\text{max}}(k_{\parallel})$. Hence, $1 + |\zeta|^2 - z^2 < 1 + |\zeta|^2 - (1 + |\zeta|)^2 = -2|\zeta| < 0$ since $k_{\parallel} \neq \pi$. Therefore,

$$(2.39) \quad \text{in subcase (1b), we have } |z| > \delta_{\text{max}}(k_{\parallel}) \text{ and } |\lambda_2(z)| < 1 < |\lambda_1(z)|.$$

Case 2: Here we have $\delta_{\text{gap}}(k_{\parallel}) \leq |z| \leq \delta_{\text{max}}(k_{\parallel})$. In this case, $\lambda_1 = (a + ib)/(2\zeta^*)$ and $\lambda_2 = (a - ib)/(2\zeta^*)$, where a and b are real. Therefore, $|\lambda_1|/|\lambda_2| = 1$ and hence $|\lambda_1| = |\lambda_2|$ implying that

$$(2.40) \quad \text{in case (2), we have } \delta_{\text{gap}}(k_{\parallel}) \leq |z| \leq \delta_{\text{max}}(k_{\parallel}) \text{ and } |\lambda_1(z)| = |\lambda_2(z)| = 1.$$

We note the assertions (2.38), (2.39) and (2.40), hold for any $k_{\parallel} \notin \{2\pi/3, \pi, 4\pi/3\}$.

The proof of Lemma 2.6 is now complete.

We continue now with the proof of Theorem 2.2. Assume that $k_{\parallel} \in [0, 2\pi] \setminus \{2\pi/3, \pi, 4\pi/3\}$, and hence $0 < |\zeta(k_{\parallel})| \neq 1$, so that Lemma 2.6 applies. Corresponding to the eigenvalues, $\lambda_1(z)$ and $\lambda_2(z)$ of $M(z, \zeta)$ we can take the corresponding eigenvectors to be of the form:

$$(2.41) \quad \xi_1(z) = \begin{pmatrix} z \\ \zeta + \lambda_1 \end{pmatrix}, \quad \xi_2(z) = \begin{pmatrix} z \\ \zeta + \lambda_2 \end{pmatrix}.$$

Due to the hypothesized constraints on k_{\parallel} , in particular that $k_{\parallel} \neq 2\pi/3, 4\pi/3$, we have $\zeta \neq 0$. For small z we find the following asymptotic expansions for $\lambda_j(z, \zeta)$, which are valid uniformly in k_{\parallel} varying over any prescribed compact subset, \mathcal{J}_1 , of $[0, 2\pi] \setminus \{2\pi/3, \pi, 4\pi/3\}$:

$$(2.42) \quad \begin{aligned} k_{\parallel} \in \mathcal{J}_1 \subset (2\pi/3, 4\pi/3) \setminus \{\pi\} & \quad (\text{hence, } 0 < |\zeta(k_{\parallel})| < 1) \\ \implies \begin{cases} \lambda_1 = \lambda_1(z, \zeta) = -\zeta + \mathcal{O}(|z|^2) \\ \lambda_2 = \lambda_2(z, \zeta) = -(\zeta^*)^{-1} + \mathcal{O}(|z|^2). \end{cases} \end{aligned}$$

and

$$(2.43) \quad \begin{aligned} k_{\parallel} \in \mathcal{J}_1 \subset [0, 2\pi] \setminus [2\pi/3, 4\pi/3] & \quad (\text{equivalently, } |\zeta(k_{\parallel})| > 1) \\ \implies \begin{cases} \lambda_1 = \lambda_1(z, \zeta) = -(\zeta^*)^{-1} + \mathcal{O}(|z|^2) \\ \lambda_2 = \lambda_2(z, \zeta) = -\zeta + \mathcal{O}(|z|^2). \end{cases} \end{aligned}$$

The resolvent $(H_{\sharp}^{\text{TB}}(k_{\parallel}) - z I)^{-1}$ **on** $l^2(\mathbb{N}_0; \mathbb{C}^2)$: Let us now restrict k_{\parallel} to vary over the set $(2\pi/3, 4\pi/3) \setminus \{\pi\}$, and assume $0 < |z| < \delta_{\text{gap}}(k_{\parallel})$; and construct the resolvent of $H_{\sharp}^{\text{TB}}(k_{\parallel})$ by solving (2.27), (2.28). The construction of the resolvent for $|z| > \delta_{\text{max}}(k_{\parallel})$ for all $k_{\parallel} \in [0, 2\pi]$ and all z such that $|z| < \delta_{\text{gap}}(k_{\parallel})$, where $k_{\parallel} \in [0, 2\pi] \setminus (2\pi/3, 4\pi/3)$ can be carried out similarly (see remarks below).

For $k_{\parallel} \in (2\pi/3, 4\pi/3) \setminus \{\pi\}$, the expansions (2.42) are valid and we have

$$\zeta + \lambda_1 = \mathcal{O}(|z|^2), \quad \zeta + \lambda_2 = \zeta - \frac{1}{\zeta^*} + \mathcal{O}(|z|^2),$$

and we have by (2.41) that the eigenvectors satisfy

$$(2.44) \quad \frac{1}{z}\xi_1(z) = \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \mathcal{O}_{\mathbb{C}^2}(|z|), \quad \xi_2(z) = \left(\zeta - \frac{1}{\zeta^*}\right) \begin{pmatrix} 0 \\ 1 \end{pmatrix} + \mathcal{O}_{\mathbb{C}^2}(|z|)$$

for all z small. Hence,

$$\left\{ \frac{1}{z}\xi_1(z), \xi_2(z) \right\} \text{ is a basis of } \mathbb{C}^2 \text{ for } 0 < |z| < \delta_{\text{gap}}(k_{\parallel}) \text{ and } k_{\parallel} \in (2\pi/3, 4\pi/3) \setminus \{\pi\}$$

which does not degenerate in the limit $z \rightarrow 0$. Indeed, by (2.41) for $z \neq 0$ this set is linearly independent if and only if $\lambda_1 \neq \lambda_2$. However, for $0 < |z| < \delta_{\text{gap}}(k_{\parallel})$ we have $|\lambda_1| < 1 < |\lambda_2|$.

To solve (2.27), (2.28) we next express $F_n = F_n(z, \zeta; f)$ in the non-degenerate basis (2.44). We shall, when convenient, suppress the dependence of F_n on ζ and f :

$$(2.45) \quad \begin{aligned} F_n(f; z, \zeta) &= \begin{pmatrix} f_n^B & \\ \frac{z}{\zeta^*} f_n^B & + \frac{1}{\zeta^*} f_{n+1}^A \end{pmatrix} \\ &= F_n^{(1)}(f; z, \zeta) \frac{1}{z} \xi_1(z) + F_n^{(2)}(f; E, \zeta) \xi_2(z). \end{aligned}$$

We also seek a solution as an expansion in the basis (2.44):

$$(2.46) \quad \psi_n = \psi_n^{(1)} \frac{1}{z} \xi_1(z) + \psi_n^{(2)} \xi_2(z),$$

where $\psi_n^{(1)} = \psi_n^{(1)}(z)$ and $\psi_n^{(2)} = \psi_n^{(2)}(z)$ are to be determined. Then, we obtain the two decoupled first order difference equations:

$$(2.47) \quad \psi_{n+1}^{(1)} = \lambda_1(z) \psi_n^{(1)} + F_n^{(1)}(z), \quad n \geq 0,$$

$$(2.48) \quad \psi_{n+1}^{(2)} = \lambda_2(z) \psi_n^{(2)} + F_n^{(2)}(z), \quad n \geq 0,$$

with boundary condition (2.28) to be expressed in terms of $\psi_0^{(j)}$, and $F_0^{(j)}$, $j = 1, 2$:

$$(2.49) \quad \frac{1}{z} \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^\top \xi_1(z) \psi_0^{(1)} + \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^\top \xi_2(z) \psi_0^{(2)} = f_0^A$$

We now proceed to solve the decoupled system (2.47)-(2.48) and then impose the boundary condition (2.49). Recall our assumption that $0 < |\zeta| < 1$, *i.e.* $k_{\parallel} \in (2\pi/3, 4\pi/3) \setminus \{\pi\}$ and therefore for z real and $|z| < \delta_{\text{gap}}(k_{\parallel})$, we have that $|\lambda_1(z)| < 1 < |\lambda_2(z)|$. In this case, the most general solution of (2.47), which decays as $n \rightarrow +\infty$ is:

$$(2.50) \quad \psi_n^{(1)}(z) = \sum_{j=0}^{n-1} (\lambda_1(z))^{n-1-j} F_j^{(1)}(z) + \mu (\lambda_1(z))^n.$$

where μ is an arbitrary constant to be determined and $F_j^{(1)}(f; z, \zeta)$, $F_j^{(2)}(f; z, \zeta)$ are defined by (2.45).

Furthermore, the most general solution of (2.48) which decays as $n \rightarrow +\infty$ is:

$$(2.51) \quad \psi_n^{(2)}(z) = - \sum_{j=n}^{\infty} (\lambda_2(z))^{n-j-1} F_j^{(2)}(z).$$

Finally, we now turn to the boundary condition (2.49). Using (2.50) and (2.51) for $n = 0$ in (2.49) we find:

$$(2.52) \quad \mu \frac{1}{z} \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^\top \xi_1(z) - \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^\top \xi_2(z) \sum_{j=0}^{\infty} (\lambda_2(z))^{-j-1} F_j^{(2)}(z, \zeta; f) = f_0^A.$$

By (2.32), the quadratic equation for the roots λ_j , we find:

$$(2.53) \quad \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^\top \xi_j(z) = \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^\top \begin{pmatrix} z \\ \zeta + \lambda_j(z) \end{pmatrix} = - \frac{\zeta + \lambda_j(z)}{\lambda_j(z)}, \quad j = 1, 2.$$

Claim: Assume $z \neq 0$ and $z \in \mathbb{R}$. If $\lambda(z)$ is any root of (2.32), then $\frac{\zeta + \lambda(z)}{\lambda(z)} \neq 0$.

It follows from this claim and (2.53) that the coefficient of μ in (2.52) is non-zero and hence if $z \neq 0$ we can solve (2.49) for $\mu = \mu(z, \zeta; f)$.

To prove the above Claim we first note that $\lambda \neq 0$. Indeed, if $\lambda = 0$ then (2.32) would then imply $\zeta = 1 + e^{ik_{\parallel}} = 0$; this contradicts our assumption that $k_{\parallel} \neq \pi$. Thus, $\lambda(z) \neq 0$. Furthermore, we claim that $\zeta + \lambda(z) \neq 0$. Again, using (2.32) we have that if $\zeta + \lambda = 0$ then $\zeta z^2 = 0$. This contradicts the assumptions that $z \neq 0$ and $\zeta \neq 0$.

It follows from this discussion that for $z \neq 0$ and $k_{\parallel} \neq \pi$:

$$(2.54) \quad \mu(f; z, \zeta) = -\frac{z \lambda_1(z)}{\zeta + \lambda_1(z)} \left[f_0^A - \frac{\zeta + \lambda_2(z)}{\lambda_2(z)} \sum_{j=0}^{\infty} (\lambda_2(z))^{-j-1} F_j^{(2)}(f; z, \zeta) \right]$$

Therefore if $0 < |z| < \delta_{\text{gap}}(k_{\parallel})$ and $k_{\parallel} \in (2\pi/3, 4\pi/3) \setminus \{\pi\}$, we can solve for $\mu = \mu(z, \zeta; f)$. We obtain for any $f \in l^2(\mathbb{N}_0; \mathbb{C}^2)$, the unique solution of (2.27), (2.28) and (2.29)

$\psi = \{\psi_n\}_{n \geq 0}$, with ψ_n tending to zero as $n \rightarrow \infty$, is given by

$$(2.55) \quad \begin{aligned} \psi_n = & \left[\sum_{j=0}^{n-1} (\lambda_1(z, \zeta))^{n-1-j} F_j^{(1)}(f; z, \zeta) + \mu(z, \zeta; f) (\lambda_1(z, \zeta))^n \right] \frac{1}{z} \xi_1(z, \zeta) \\ & - \left[\sum_{j=n}^{\infty} (\lambda_2(z, \zeta))^{n-j-1} F_j^{(2)}(f; z, \zeta) \right] \xi_2(z, \zeta), \quad n \geq 0, \end{aligned}$$

where $\mu = \mu(z, \zeta; f)$ is obtained from (2.52). By (2.45), we may express $F_j^{(1)}$ and $F_j^{(2)}$ as

$$(2.56) \quad F_j^{(1)} = \alpha_1(z, \zeta) f_j^B + \alpha_2(z, \zeta) f_{j+1}^A, \quad F_j^{(2)} = \beta_1(z, \zeta) f_j^B + \beta_2(z, \zeta) f_{j+1}^A,$$

where the coefficients are bounded and smooth over the ranges of z and k_{\parallel} under consideration.

Next, introduce the discrete vector-valued kernel, depending on parameters α and β :

$$(2.57) \quad \mathcal{K}(n, j; \alpha, \beta) = \begin{cases} \alpha \lambda_1(z, \zeta)^{n-1-j} \frac{1}{z} \xi_1(z, \zeta), & 0 \leq j \leq n-1 \\ -\beta \lambda_2(z, \zeta)^{n-1-j} \xi_2(z, \zeta), & n \leq j < \infty. \end{cases}$$

Then, we have

$$(2.58) \quad \begin{aligned} \psi_n = & \sum_{j=0}^{\infty} \mathcal{K}(n, j; \alpha_1, \beta_1) f_j^B + \sum_{j=0}^{\infty} \mathcal{K}(n, j; \alpha_2, \beta_2) f_{j+1}^A \\ & + \mu(f; z, \zeta) (\lambda_1(z, \zeta))^n \frac{1}{z} \xi_1(z, \zeta), \end{aligned}$$

where $\mu(f; z, \zeta)$ is given by the linear functional of f , displayed in (2.54).

Proposition 2.7. *Let \mathcal{J}_1 denote a compact subset of $(2\pi/3, 4\pi/3) \setminus \{\pi\}$ and let $\eta(\mathcal{J}_1) > 0$, denote the constant appearing in part (3) of Lemma 2.6.*

(1) *There is a constant, C , depending on \mathcal{J}_1 such that for all complex energies, $z \in \mathcal{O}_0(k_{\parallel}) \setminus \{0\}$ (see (2.36)), the resolvent operator:*

$$(2.59) \quad f \in l^2(\mathbb{N}_0; \mathbb{C}^2) \mapsto \psi = \{\psi_n\}_{n \geq 0} \equiv \left(H_{\sharp}^{\text{TB}}(k_{\parallel}) - z \right)^{-1} f,$$

given by the expression in (2.58), defines a bounded linear operator on $l^2(\mathbb{N}_0; \mathbb{C}^2)$ with

$$(2.60) \quad \left\| \left(H_{\#}^{\text{TB}}(k_{\parallel}) - z \right)^{-1} f \right\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)} \leq C \frac{1}{|z|} \|f\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)},$$

where the constant, C , is independent of depends on the compact set \mathcal{J}_1 .

(2) The mapping $z \mapsto \left(H_{\#}^{\text{TB}}(k_{\parallel}) - z \right)^{-1}$ is meromorphic for z varying in the open set $\mathcal{O}_0(k_{\parallel})$ into $\mathcal{B}(l^2(\mathbb{N}_0; \mathbb{C}^2))$, the space of bounded linear operators on $l^2(\mathbb{N}_0; \mathbb{C}^2)$, with only pole at $z = 0$. For $z \in \mathcal{O}_0(k_{\parallel}) \setminus \{0\}$ we have

$$(2.61) \quad \left(H_{\#}^{\text{TB}}(k_{\parallel}) - zI \right)^{-1} f = \frac{1}{z} \langle \psi^{\text{TB, bd}}(k_{\parallel}), f \rangle_{l^2(\mathbb{N}_0; \mathbb{C}^2)} \psi^{\text{TB, bd}}(k_{\parallel}) + \mathcal{G}_{\text{reg}}(z; k_{\parallel}) f,$$

where $z \mapsto \mathcal{G}_{\text{reg}}(z; k_{\parallel})$ is an analytic map from $\mathcal{O}_0(k_{\parallel})$ to $\mathcal{B}(l^2(\mathbb{N}_0; \mathbb{C}^2))$.

(3) $H_{\#}^{\text{TB}}(k_{\parallel})\psi = f \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ has a solution in the space $l^2(\mathbb{N}_0; \mathbb{C}^2)$ if and only if $\langle \psi^{\text{TB, bd}}(k_{\parallel}), f \rangle_{l^2(\mathbb{N}_0; \mathbb{C}^2)} = 0$.

Proof of Proposition 2.7: We fix $\mathcal{J}_1 \subset (2\pi/3, 4\pi/3) \setminus \{\pi\}$ and take $E \in \mathcal{O}_0(k_{\parallel}) \setminus \{0\}$. To bound the resolvent we estimate the expression in $\{\psi_n\}_{n \geq 0}$ displayed in (2.58) in $l^2(\mathbb{N}_0; \mathbb{C}^2)$. We begin with an estimate of the latter term in (2.58): $\mu(f; z, \zeta) (\lambda_1(z, \zeta))^n \frac{1}{z} \xi_1(z, \zeta)$. From the expression for μ in (2.54) and the definition of $F_j^{(2)}$ in (2.45) (recall $F_j^{(1)}$ and $F_j^{(2)}$ are coordinates of $F_j \in \mathbb{C}^2$, also given in (2.45)) with respect to the basis $\{\frac{1}{z}\xi_1(z), \xi_2(z)\}$, we have that $|\mu(f; z, \zeta)| \lesssim |f_0^A| + \sum_{j=0}^{\infty} |\lambda_2|^{-j-1} (|f_j^B| + |f_{j+1}^A|) \leq C_1(z, \zeta) \|f\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)}$, where $C_1(z, \zeta)$ is a finite constant which depends on z and ζ in the ranges specified above. The constant $C_1(z, \zeta)$ is bounded for z bounded away from $z = 0$ and $k_{\parallel} \in \mathcal{J}_1$. As we shall see below, for $k_{\parallel} \in \mathcal{J}_1$, there is pole of order one as $E \rightarrow 0$.

Therefore, applying Young's inequality to the first two terms in (2.58) we obtain:

$$\|\psi\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)} \leq \left(C(\mathcal{K}, z, \zeta) + C_1(z, \zeta) \right) \|f\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)},$$

where

$$(2.62) \quad C(\mathcal{K}, z, \zeta) = \max_{r=1,2} \left(\sup_{n \geq 0} \sum_{j=0}^{\infty} |\mathcal{K}(n, j, \alpha_r, \beta_r)| + \sup_{j \geq 0} \sum_{n=0}^{\infty} |\mathcal{K}(n, j, \alpha_r, \beta_r)| \right),$$

and we recall from (2.56) that α_r and β_r are smooth and bounded functions of z and ζ . Estimating the first sum in (2.62), we have for $r = 1, 2$:

$$(2.63) \quad \begin{aligned} \sum_{j=0}^{\infty} |\mathcal{K}(n, j, \alpha_r, \beta_r)| &\lesssim |\alpha_r(z, \zeta)| \sum_{j=0}^{n-1} |\lambda_1(z, \zeta)|^{n-1-j} + |\beta_r(z, \zeta)| \sum_{j=n}^{\infty} |\lambda_2(z, \zeta)|^{n-1-j} \\ &\lesssim |\alpha_r(z, \zeta)| (1 - |\lambda_1(z, \zeta)|)^{-1} + |\beta_r(z, \zeta)| (|\lambda_2(z, \zeta)| - 1)^{-1} \end{aligned}$$

The bound (2.63) holds, for $r = 1, 2$ and any fixed $z \in \mathcal{O}_0(k_{\parallel}) \setminus \{0\}$, uniform in $k_{\parallel} \in \mathcal{J}_1$. The second sum in (2.62) is bounded similarly. Therefore, we have for all $k_{\parallel} \in \mathcal{J}_1$ and any $z \in \mathcal{O}_0(k_{\parallel})$, the resolvent operator: $f \mapsto \left(H_{\#}^{\text{TB}}(k_{\parallel}) - z \right)^{-1} f$ (see (2.59)) is a bounded linear operator on $l^2(\mathbb{N}_0; \mathbb{C}^2)$. The next step in the proof of Proposition 2.7 requires us to consider the resolvent for small complex z in $\mathcal{O}_0(k_{\parallel}) \setminus \{0\}$.

2.4. The resolvent $(H_{\#}^{\text{TB}}(k_{\parallel}) - z I)^{-1}$ for z near zero energy. Since there is a simple zero energy eigenstate for each $k_{\parallel} \in (2\pi/3, 4\pi/3)$, we expect a simple pole of the resolvent at $z = 0$. We now make this explicit by expanding the resolvent in a neighborhood of $z = 0$ for $k_{\parallel} \in (2\pi/3, 4\pi/3)$. In order to work with the above detailed calculations, we restrict our discussion to the case where $k_{\parallel} \neq \pi$ ($\zeta \neq 0$). Consider first the relation (2.52), which determined the free parameter $\mu = \mu(f; z, \zeta)$. We shall simplify (2.52) using the following expansions which hold for $|z|$ small:

(2.64)

$$\begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^{\top} \frac{1}{z} \xi_1(z) = \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^{\top} \frac{1}{z} \begin{pmatrix} z \\ \zeta + \lambda_1(z) \end{pmatrix} = -\frac{1}{z} \frac{\zeta + \lambda_1(z)}{\lambda_1(z)} = \frac{z}{|\zeta|^2 - 1} + \mathcal{O}(|z|^3)$$

(2.65)

$$\begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^{\top} \xi_2(z) = \begin{pmatrix} -z \\ \zeta^* \end{pmatrix}^{\top} \begin{pmatrix} z \\ \zeta + \lambda_2(z) \end{pmatrix} = -\frac{\zeta + \lambda_2(z)}{\lambda_2(z)} = |\zeta|^2 - 1 + \mathcal{O}(|z|^2)$$

We also have from (2.45) that

$$\begin{aligned} F_n(f; z, \zeta) &= \begin{pmatrix} f_n^B \\ \frac{z}{\zeta^*} f_n^B + \frac{1}{\zeta^*} f_{n+1}^A \end{pmatrix} \\ &= f_n^B \frac{1}{z} \xi_1(z) + f_{n+1}^A \frac{1}{\zeta^*} \cdot \left(\zeta - \frac{1}{\zeta^*} \right)^{-1} \xi_2(z) + \mathcal{O}(|z| [|f_n| + |f_{n+1}|]). \end{aligned}$$

Therefore, for $|z|$ small

$$\begin{aligned} F_n^{(1)}(f; z, \zeta) &= f_n^B + \mathcal{O}(|z| [|f_n| + |f_{n+1}|]), \\ F_n^{(2)}(f; z, \zeta) &= \frac{1}{|\zeta|^2 - 1} f_{n+1}^A + \mathcal{O}(|z| [|f_n| + |f_{n+1}|]). \end{aligned} \quad (2.66)$$

Substitution of the expansions (2.64), (2.65) and (2.66) into (2.52), we obtain:

$$\begin{aligned} \frac{z}{|\zeta|^2 - 1} \mu - (|\zeta|^2 - 1) \sum_{j=0}^{\infty} \left(-\frac{1}{\zeta^*} \right)^{-(j+1)} \frac{1}{|\zeta|^2 - 1} f_{j+1}^A \\ + \mathcal{O}(|z| \|f\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)}) + \mathcal{O}(|z| |\mu|) = f_0^A. \end{aligned} \quad (2.67)$$

Hence,

$$\begin{aligned} \frac{z}{|\zeta|^2 - 1} \mu &= f_0^A + \sum_{j=0}^{\infty} \left(-\frac{1}{\zeta^*} \right)^{-(j+1)} f_{j+1}^A + \mathcal{O}(|z| \|f\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)}) \\ &= \sum_{j=0}^{\infty} (-\zeta^*)^j f_j^A + \mathcal{O}(|z| \|f\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)}) . \end{aligned} \quad (2.68)$$

Recall that we have assumed $k_{\parallel} \in \mathcal{J}_1 \subset (2\pi/3, 4\pi/3) \setminus \{\pi\}$ (thus $|\zeta(k_{\parallel})|^2 - 1 \neq 0$) and $z \in \mathcal{O}_0(k_{\parallel}) \setminus \{0\}$. Solving (2.68) for $\mu(f; z, \zeta)$ and using the expression for $\{\psi_j^{\text{TB}, \text{bd}}(k_{\parallel})\}_{j \geq 0}$,

the zero energy eigenstate of H_{\sharp}^{TB} in (2.17), we obtain:

$$\begin{aligned}
\mu(z, \zeta; f) &= \frac{1}{z} \sqrt{1 - |\zeta|^2} \sum_{j=0}^{\infty} \overline{\psi_j^{\text{TB, bd}}(k_{\parallel})} f_j^A + \mathcal{O}(\|f\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)}) \\
(2.69) \quad &= \frac{1}{z} \sqrt{1 - |\zeta|^2} \langle \psi^{\text{TB, bd}}(k_{\parallel}), f \rangle_{l^2(\mathbb{N}_0; \mathbb{C}^2)} + \mathcal{O}(\|f\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)}),
\end{aligned}$$

The error bound in (2.69) is uniform in $k_{\parallel} \in \mathcal{J}_1 \setminus \{\pi\}$ and bounds an expression which is analytic in $z \in \mathcal{O}_0(k_{\parallel}) \setminus \{0\}$. From the previous discussion we conclude the following. Fix any $k_{\parallel} \in \mathcal{J}_1 \subset \subset (2\pi/3, 4\pi/3) \setminus \{\pi\}$. Let $\mathcal{O}_0(k_{\parallel})$ denote the open neighborhood in \mathbb{C} defined in (2.36). Then, for all z in $\mathcal{O}_0(k_{\parallel})$, the mapping

$$z \in \mathcal{O}_0(k_{\parallel}) \mapsto \left(H_{\sharp}^{\text{TB}}(k_{\parallel}) - zI \right)^{-1} \text{ is meromorphic with values in } l^2(\mathbb{N}_0; \mathbb{C}^2)$$

with only one pole, located at $z = 0$. Moreover, for $z \in \mathcal{O}_0(k_{\parallel}) \setminus \{0\}$ we have

$$(2.70) \quad \left(H_{\sharp}^{\text{TB}}(k_{\parallel}) - zI \right)^{-1} f = \frac{1}{z} \langle \psi^{\text{TB, bd}}(k_{\parallel}), f \rangle_{l^2(\mathbb{N}_0; \mathbb{C}^2)} \psi^{\text{TB, bd}}(k_{\parallel}) + \mathcal{G}_{\text{reg}}(z; k_{\parallel})f,$$

where $z \mapsto \mathcal{G}_{\text{reg}}(z; k_{\parallel})$ is an analytic map from $\mathcal{O}_0(k_{\parallel})$ to $\mathcal{B}(l^2(\mathbb{N}_0; \mathbb{C}^2))$. Thus we have proved part (3a) of Theorem 2.2, except for the case $k_{\parallel} = \pi$. We leave this as an exercise for the reader.

Note that for all $k_{\parallel} \in (2\pi/3, 4\pi/3)$, we have that

$$(2.71) \quad H_{\sharp}^{\text{TB}}(k_{\parallel})\psi = f \in l^2(\mathbb{N}_0; \mathbb{C}^2) \text{ is solvable in } l^2(\mathbb{N}_0; \mathbb{C}^2) \iff \langle \psi^{\text{TB, bd}}(k_{\parallel}), f \rangle_{l^2(\mathbb{N}_0; \mathbb{C}^2)} = 0.$$

Thus we have proved all assertions of Theorem 2.2 for $k_{\parallel} \in \mathcal{J}_1$ (\mathcal{J}_1 arbitrary compact subset of $(2\pi/3, 4\pi/3)$), and all E in the open complex neighborhood $\mathcal{O}_0(k_{\parallel})$, defined in (2.36).

It remains to address the cases:

- (A) $k_{\parallel} \in [0, 2\pi] \setminus (2\pi/3, 4\pi/3)$ and $z \in \mathcal{O}_0(k_{\parallel})$, defined in (2.36) and
- (B) $k_{\parallel} \in [0, 2\pi]$ and $z \in \mathcal{O}_+(k_{\parallel})$, defined in (2.37).

In case (A), Lemma 2.6 tells us that $|\lambda_1(z)| < 1 < |\lambda_2(z)|$. Hence, the construction of the resolvent is as above, and gives the map $f \mapsto \psi$ defined by (2.55). However now, since $z = 0$ is not an eigenvalue, $\mu = \mu(f; z, \zeta)$ does not have a pole, as was the case in for for $k_{\parallel} \in (2\pi/3, 4\pi/3)$; see (2.54).

In case (B), Lemma 2.6 tells us that $|\lambda_2(z)| < 1 < |\lambda_1(z)|$. The construction of the resolvent is analogous with the roles of the eigenpairs: (λ_1, ξ_1) and (λ_2, ξ_2) interchanged. Since in $\mathcal{O}_+(k_{\parallel})$ $|z| > |\Re z| > \delta_{\max}(k_{\parallel}) \geq 1$ and the only possible eigenvalue is at $z = 0$, the analogue of the $\mu(f; z, \zeta)$ - term in (2.55) does not have a pole in this case as well. Therefore, in both cases (A) and (B) the mapping $z \mapsto \left(H_{\sharp}^{\text{TB}} - zI \right)^{-1}$ is analytic with values in $\mathcal{B}(l^2(\mathbb{N}_0; \mathbb{C}^2))$.

Finally, using part (2) of Lemma 2.6, one can check that $H_{\sharp}^{\text{TB}}(k_{\parallel}) - zI$ is not invertible for $\delta_{\text{gap}}(k_{\parallel}) \leq |z| \leq \delta_{\max}(k_{\parallel})$ since the eigenvalues of $M(z, \zeta)$ satisfy: $|\lambda_1(z, \zeta)| = |\lambda_2(z, \zeta)| = 1$. Such energies z comprise the essential spectrum of $H_{\sharp}^{\text{TB}}(k_{\parallel})$, $\sigma_{\text{ess}}\left(H_{\sharp}^{\text{TB}}(k_{\parallel})\right)$. The details are left to the reader. This completes the proof of Theorem 2.2.

3. SETUP FOR THE CONTINUUM PROBLEM; ZIGZAG EDGE HAMILTONIAN AND THE ZIGZAG EDGE-STATE EIGENVALUE PROBLEM

In this section we begin our detailed formulation and discussion of the continuum edge state eigenvalue problem. For this we must first discuss the atomic, bulk and edge Hamiltonians: H_{atom}^λ , H_{bulk}^λ and H_{\sharp}^λ .

3.1. The atomic Hamiltonian and its ground state. We work with the class of “atomic potential wells” introduced in [26]. Fix a smooth potential $V_0(\mathbf{x})$ on \mathbb{R}^2 with the following properties.

$$(PW_1) \quad -1 \leq V_0(\mathbf{x}) \leq 0, \quad \mathbf{x} \in \mathbb{R}^2.$$

$$(PW_2) \quad \text{supp } V_0 \subset \{\mathbf{x} \in \mathbb{R}^2 : |\mathbf{x}| < r_0\}, \text{ where } r_0 < r_{\text{cr}}. \text{ Here, } r_{\text{cr}} \text{ is a universal constant defined in [26] satisfying } 0.33|\mathbf{e}| \leq r_{\text{cr}} < 0.5|\mathbf{e}|, \text{ and } |\mathbf{e}| = |\mathbf{v}_B - \mathbf{v}_A| = 1/\sqrt{3} \text{ is the distance between one vertex in } \mathbb{H} \text{ and any nearest neighbor.}$$

$$(PW_3) \quad V_0(\mathbf{x}) \text{ is invariant under a } 2\pi/3 \text{ (120}^\circ\text{) rotation about the origin, } \mathbf{x} = 0.$$

$$(PW_4) \quad V_0(\mathbf{x}) \text{ is inversion-symmetric with respect to the origin; } V_0(-\mathbf{x}) = V_0(\mathbf{x}).$$

Consider the “atomic” Hamiltonian: $H_{\text{atom}}^\lambda = -\Delta + \lambda^2 V_0(\mathbf{x})$ acting in $L^2(\mathbb{R}^2)$. Let $p_0^\lambda(\mathbf{x})$, E_0^λ , respectively, be the ground state eigenfunction and its strictly negative ground state eigenvalue:

$$(3.1) \quad (-\Delta + \lambda^2 V_0(\mathbf{x}) - E_0^\lambda) p_0^\lambda(\mathbf{x}) = 0, \quad p_0^\lambda \in L^2(\mathbb{R}^2), \quad E_0^\lambda < 0.$$

This eigenpair is simple and, by the symmetries of $V_0(\mathbf{x})$, the ground state $p_0^\lambda(\mathbf{x})$ is invariant under a $\pi/3$ (60°) rotation about the origin. We choose $p_0^\lambda(\mathbf{x})$ so that $p_0^\lambda(\mathbf{x}) > 0$ for all $\mathbf{x} \in \mathbb{R}^2$ and $\int_{\mathbb{R}^2} |p_0^\lambda(\mathbf{x})|^2 d\mathbf{x} = 1$. Since $V_0 \in L^\infty(\mathbb{R}^2)$ and $-\Delta p_0^\lambda = (E_0^\lambda - \lambda^2 V_0) p_0^\lambda$, it follows that $p_0^\lambda \in H^2(\mathbb{R}^2)$.

Recall the hopping coefficient ρ_λ given by:

$$(3.2) \quad \rho_\lambda = \int_{|\mathbf{y}| < r_0} p_0^\lambda(\mathbf{y}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} - \mathbf{e}) d\mathbf{y}.$$

By Proposition 4.1 of [26] we have, under hypotheses $(PW_1), \dots, (PW_4)$ on $V_0(\mathbf{x})$ the upper and lower bounds for large λ :

$$(3.3) \quad e^{-c_- \lambda} \lesssim \rho_\lambda \lesssim e^{-c_+ \lambda}$$

for some constants: $0 < c_+ < c_-$ which depend on V_0 but not on λ .

Remark 3.1. The edge states we construct will have energies $E^\lambda = E_0^\lambda + \Omega^\lambda$, with $\rho_\lambda^{-1} |\Omega^\lambda| \ll 1$. In preparation for our later discussion, it is useful at this stage to introduce a positive constant, \hat{c} , such that $\hat{c} > c_-$ (see (3.3)) and to observe that

$$|\Omega^\lambda| < e^{-\hat{c}\lambda} \implies \rho_\lambda^{-1} |\Omega^\lambda| < e^{-(\hat{c}-c_-)\lambda} \downarrow 0 \text{ as } \lambda \uparrow \infty.$$

In addition to hypotheses $(PW_1), \dots, (PW_4)$ on $V_0(\mathbf{x})$, we assume the following two spectral properties of $H_{\text{atom}}^\lambda = -\Delta + \lambda^2 V_0$ acting on $L^2(\mathbb{R}^2)$:

(GS) Ground state energy upper bound: For λ large, E_0^λ , the ground state energy of $-\Delta + \lambda^2 V_0(\mathbf{x})$, satisfies the upper bound

$$(3.4) \quad E_0^\lambda \leq -c_{\text{gs}} \lambda^2.$$

Here, c_{gs} is a strictly positive constant depending on V_0 . A simple consequence of the variational characterization of E_0^λ is the lower bound $E_0^\lambda \geq -\|V_0\|_{L^\infty} \lambda^2 = -\lambda^2$. However, the upper bound (3.4) requires further restrictions on V_0 . Using the condition (GS), we can show that p_0^λ , satisfies the following pointwise bound:

$$(3.5) \quad |p_0^\lambda(\mathbf{x})| \leq C_1 \left(\lambda \mathbf{1}_{|\mathbf{x}| < r_0 + \delta_0} + e^{-c_1 \lambda |\mathbf{x}|} \right)$$

where $\text{supp}(V_0) \subset B(0, r_0)$, $\delta_0 > 0$ is arbitrary, and C_1 and c_1 are constants that depend on V_0 , r_0 and δ_0 ; see Corollary 15.5 of [26].

(EG) Energy gap property: For $\lambda > 0$ sufficiently large, there exists $c_{\text{gap}} > 0$, independent of λ , such that if $\psi \in H^2(\mathbb{R}^2)$ and $\langle p_0^\lambda, \psi \rangle_{L^2(\mathbb{R}^2)} = 0$, then

$$(3.6) \quad \langle (-\Delta + \lambda^2 V_0 - E_0^\lambda) \psi, \psi \rangle_{L^2(\mathbb{R}^2)} \geq c_{\text{gap}} \|\psi\|_{L^2(\mathbb{R}^2)}^2.$$

In Section 4.1 of [26] we discuss examples of potentials for which $-\Delta + \lambda^2 V_0$ satisfies (GS) and (EG).

3.2. Review of terminology and formulation. We conclude this section with a review of some terminology and the formulation of the edge state eigenvalue problem.

(1) *Continuum bulk Hamiltonian, H_{bulk}^λ :*

$$(3.7) \quad H_{\text{bulk}}^\lambda \equiv -\Delta + \lambda^2 V(\mathbf{x}) \quad \text{acting on } L^2(\mathbb{R}^2).$$

Here, $V(\mathbf{x})$, the *bulk periodic potential*, is defined to be the sum of all translates of atomic wells, $V_0(\mathbf{x} - \mathbf{v})$, where \mathbf{v} ranges over \mathbb{H} : $V(\mathbf{x}) = \sum_{\mathbf{v} \in \mathbb{H}} V_0(\mathbf{x} - \mathbf{v})$; see (1.10).

The potential $V(\mathbf{x})$ is a honeycomb lattice potential in the sense of Definition 2.1 of [25]; V is real-valued, and with respect to an origin placed at the center of a regular hexagon of the tiling of \mathbb{R}_x^2 : V is inversion symmetric and rotationally invariant by $2\pi/3$.

(2) *Continuum zigzag edge Hamiltonian, H_{edge}^λ :* The potential for a honeycomb structure interfaced with the vacuum along a sharp interface with direction $\mathbf{v}_2 \in \Lambda$ (parallel to the zigzag edge) is obtained by summing translates of V_0 over the truncated structure, \mathbb{H}_\sharp , defined in (1.11):

$$(3.8) \quad V_\sharp(\mathbf{x}) = \sum_{\mathbf{v} \in \mathbb{H}_\sharp} V_0(\mathbf{x} - \mathbf{v}).$$

The Hamiltonian for the truncated structure is given by

$$(3.9) \quad H_{\text{edge}}^\lambda \equiv -\Delta + \lambda^2 V_\sharp(\mathbf{x}), \quad \text{acting on } L^2(\mathbb{R}^2),$$

and its centering at the ground state energy, E_0^λ , of H_{atom}^λ is denoted:

$$(3.10) \quad H_\sharp^\lambda \equiv -\Delta + \lambda^2 V_\sharp(\mathbf{x}) - E_0^\lambda \quad \text{acting on } L^2(\mathbb{R}^2).$$

Since H_{edge}^λ and H_\sharp^λ are invariant under the translation invariance: $\mathbf{x} \mapsto \mathbf{x} + \mathbf{v}_2$, these operators act in $L_{k_\parallel}^2(\Sigma)$, $\Sigma = \mathbb{R}^2 / \mathbb{Z}\mathbf{v}_2$.

(3) The k_\parallel -dependent Edge Hamiltonian, $H_\sharp^\lambda(k_\parallel)$, acting in $L^2(\Sigma)$ is given by:

$$(3.11) \quad H_\sharp^\lambda(k_\parallel) \equiv - \left(\nabla + i \frac{k_\parallel}{2\pi} \mathfrak{R}_2 \right)^2 + \lambda^2 V_\sharp(\mathbf{x}) - E_0^\lambda.$$

Finally we recall that the *Zigzag Edge state Eigenvalue Problem* is given by (1.16), or equivalently, (1.17). With $E = E_0^\lambda + \Omega$, we have:

$$(3.12) \quad (H_{\sharp}^\lambda(k_{\parallel}) - \Omega) \psi = 0, \quad \psi \in L_{k_{\parallel}}^2.$$

4. A NATURAL SUBSPACE OF $L_{k_{\parallel}}^2(\Sigma)$

Define, for all $n \geq 0$ ¹

$$(4.1) \quad \mathbf{v}_A^n \equiv \mathbf{v}_A + n\mathbf{v}_1, \quad \mathbf{v}_B^n \equiv \mathbf{v}_B + n\mathbf{v}_1,$$

where $\mathbf{v}_A^0 = \mathbf{v}_A$ and $\mathbf{v}_B^0 = \mathbf{v}_B$. The cylinder $\Sigma = \mathbb{R}^2 / \mathbb{Z}\mathbf{v}_2$ has fundamental domain $\Omega_\Sigma \subset \mathbb{R}^2$, which may be expressed as the union of parallelograms:

$$(4.2) \quad \Omega_\Sigma = \cup_{n \geq 0} \Omega_n \cup \Omega_{-1} \quad \text{as in Figure 1.}$$

Each parallelogram Ω_n with $n \geq 0$ contains two atomic sites: \mathbf{v}_A^n and \mathbf{v}_B^n . The infinite parallelogram, Ω_{-1} , contains no atomic sites. A fundamental cell of the cylinder Σ , Ω_Σ , and its decomposition into parallelograms Ω_n , for $n \geq -1$ is depicted in Figure 1. The zigzag sharp truncation of \mathbb{H} may be expressed as a union over ‘‘vertical translates’’ (translates with respect to \mathbf{v}_2) of sites within Ω_Σ :

$$\mathbb{H}_{\sharp} = \cup_{n_2 \in \mathbb{Z}} \cup_{n_1 \geq 0} \left\{ \mathbf{v}_A^{n_1} + n_2 \mathbf{v}_2, \mathbf{v}_B^{n_1} + n_2 \mathbf{v}_2 \right\}.$$

We next introduce approximate k_{\parallel} -pseudo-periodic solutions of $H_{\sharp}^\lambda \Psi = 0$ via k_{\parallel} -pseudo-periodization of the atomic ground state, p_0^λ :

Definition 4.1. Fix $k_{\parallel} \in [0, 2\pi]$ and $I = A, B$. For each $n \in \mathbb{N}_0 \equiv \{0, 1, 2, \dots\}$, define

$$(4.3) \quad \begin{aligned} p_{k_{\parallel}, I}^\lambda[n](\mathbf{x}) &\equiv \sum_{m_2 \in \mathbb{Z}} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2) e^{-i \frac{k_{\parallel}}{2\pi} \mathbf{R}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2)} \\ &= e^{-i \frac{k_{\parallel}}{2\pi} \mathbf{R}_2 \cdot (\mathbf{x} - \mathbf{v}_I)} \sum_{m_2 \in \mathbb{Z}} e^{i k_{\parallel} m_2} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2) \end{aligned}$$

and

$$(4.4) \quad P_{k_{\parallel}, I}^\lambda[n](\mathbf{x}) \equiv e^{i \frac{k_{\parallel}}{2\pi} \mathbf{R}_2 \cdot (\mathbf{x} - \mathbf{v}_I)} p_{k_{\parallel}, I}^\lambda[n](\mathbf{x}) = \sum_{m_2 \in \mathbb{Z}} e^{i k_{\parallel} m_2} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2).$$

The function $\mathbf{x} \mapsto p_{k_{\parallel}, I}^\lambda[n](\mathbf{x})$ is defined on the cylinder Σ , *i.e.* $p_{k_{\parallel}, I}^\lambda[n](\mathbf{x} + \mathbf{v}_2) = p_{k_{\parallel}, I}^\lambda[n](\mathbf{x})$. To see this, replace \mathbf{x} by $\mathbf{x} + \mathbf{v}_2$ and redefine the summation index. Furthermore, we note that: $P_{k_{\parallel}, I}^\lambda[n](\mathbf{x} + \mathbf{v}_2) = e^{i k_{\parallel}} P_{k_{\parallel}, I}^\lambda[n](\mathbf{x})$.

The functions: $p_{k_{\parallel}, I}^\lambda[n]$, $I = A, B$, $n \geq 0$, form a nearly orthonormal set in $L^2(\Sigma)$ for large λ . In particular, we have:

Proposition 4.2. Fix $k_{\parallel} \in [0, 2\pi]$ and $\lambda > 0$.

(1) For all $n \in \mathbb{N}_0$, we have $p_{k_{\parallel}, I}^\lambda[n] \in L^2(\Sigma)$ and $P_{k_{\parallel}, I}^\lambda[n] \in L_{k_{\parallel}}^2$.

Furthermore, there exist constants $\lambda_*, c > 0$ such that for all $\lambda \geq \lambda_*$:

¹The labeling convention of A - points and B - sublattice points used in the present article differs from that used in [26]. This has no effect on the results in this article or in [26].

(2) For $n \in \mathbb{N}_0$, $I = A, B$

$$(4.5) \quad \left| \left\langle p_{k_{\parallel}, I}^{\lambda}[n], p_{k_{\parallel}, J}^{\lambda}[n] \right\rangle_{L^2(\Sigma)} - \delta_{IJ} \right| \lesssim e^{-c\lambda},$$

where δ_{IJ} denotes the Kronecker delta symbol.

(3) For $I = A, B$, $m, n \in \mathbb{N}_0$ with $m \neq n$ and all $\lambda > 0$ sufficiently large:

$$(4.6) \quad \left| \left\langle p_{k_{\parallel}, I}^{\lambda}[m], p_{k_{\parallel}, J}^{\lambda}[n] \right\rangle_{L^2(\Sigma)} \right| \lesssim e^{-c\lambda|m-n|}.$$

Assertions (4.5) and (4.6) hold as well with $p_{k_{\parallel}, I}^{\lambda}[m]$ replaced by $P_{k_{\parallel}, I}^{\lambda}[m]$, defined in (4.4), and with $L^2(\Sigma)$ replaced by $L_{k_{\parallel}}^2(\mathbb{R}^2)$. Here, λ_{\star} depends only on V .

This proposition follows from the normalization and decay properties of the atomic ground state, p_0^{λ} ; the details are omitted.

We conclude this section by showing that the functions $p_{k_{\parallel}, I}^{\lambda}[n]$, $I = A, B$, $n \geq 0$, are nearly annihilated by $H_{\sharp}^{\lambda}(k_{\parallel})$.

Proposition 4.3. *There exist positive constants λ_{\star} (large) and $c > 0$, such that for all $\lambda > \lambda_{\star}$ and all $I = A, B$ and $n \geq 0$:*

$$(4.7) \quad \left| H_{\sharp}^{\lambda}(k_{\parallel}) p_{k_{\parallel}, I}^{\lambda}[n](\mathbf{x}) \right| \lesssim e^{-c|\mathbf{x} - n\mathbf{v}_1|} e^{-c\lambda}, \quad \mathbf{x} \in \Omega_{\Sigma}$$

$$(4.8) \quad \left\| H_{\sharp}^{\lambda}(k_{\parallel}) p_{k_{\parallel}, I}^{\lambda}[n] \right\|_{L^2(\Sigma)} \lesssim e^{-c\lambda}.$$

Proof of Proposition 4.3: We first note that (4.8) follows from (4.7) by integrating the square of bound (4.7) over a fundamental domain (strip), Ω_{Σ} . Thus we focus on the pointwise bound (4.7). The identity $\nabla_{\mathbf{x}} = e^{i\frac{k_{\parallel}}{2\pi}\mathfrak{R}_2 \cdot \mathbf{x}} (\nabla + i\frac{k_{\parallel}}{2\pi}\mathfrak{R}_2) e^{-i\frac{k_{\parallel}}{2\pi}\mathfrak{R}_2 \cdot \mathbf{x}}$ and (3.1) imply that for arbitrary $\hat{\mathbf{v}} \in \mathbb{R}^2$:

$$(4.9) \quad \left(- \left(\nabla + i\frac{k_{\parallel}}{2\pi}\mathfrak{R}_2 \right)^2 + \lambda^2 V_0(\mathbf{x} - \hat{\mathbf{v}}) - E_0^{\lambda} \right) e^{-i\frac{k_{\parallel}}{2\pi}\mathfrak{R}_2 \cdot (\mathbf{x} - \hat{\mathbf{v}})} p_0^{\lambda}(\mathbf{x} - \hat{\mathbf{v}}) = 0;$$

we shall apply (4.9) for $\hat{\mathbf{v}} \in \mathbb{H}_{\sharp}$.

As a first step toward obtaining the bound (4.7) for $H_{\sharp}^{\lambda}(k_{\parallel}) p_{k_{\parallel}, I}^{\lambda}[n](\mathbf{x})$, we observe that

$$\text{for } \mathbf{x} \in \Omega_{\Sigma}, \quad V_{\sharp}(\mathbf{x}) = \sum_{J=A,B} \sum_{n_1 \geq 0} V_0(\mathbf{x} - \mathbf{v}_J - n_1 \mathbf{v}_1).$$

Therefore, for $\mathbf{x} \in \Omega_\Sigma$ we have

$$\begin{aligned}
H_\#^\lambda(k_\parallel) p_{k_\parallel, I}^\lambda[n](\mathbf{x}) &= \sum_{m_2 \in \mathbb{Z}} H_\#^\lambda(k_\parallel) e^{-i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2)} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2) \\
&= H_\#^\lambda(k_\parallel) e^{-i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n)} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n) \\
&\quad + \sum_{m_2 \in \mathbb{Z} \setminus \{0\}} \left(- \left(\nabla + i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \right)^2 - E_0^\lambda \right) e^{-i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2)} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2) \\
&\quad + \sum_{m_2 \in \mathbb{Z} \setminus \{0\}} \lambda^2 V_\#(\mathbf{x}) e^{-i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2)} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2) .
\end{aligned}$$

In the second equality just above we have split off the $m_2 = 0$ and $m_2 \neq 0$ contributions. The first term of the $m_2 \neq 0$ contribution vanishes identically for $\mathbf{x} \in \Omega_\Sigma$. Indeed, equation (4.9) for p_0^λ implies that this term is a sum of terms, each containing a factor $\lambda^2 V_0(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2)$ for some $m_2 \in \mathbb{Z} \setminus \{0\}$. Each of these terms vanishes since the constraint: $m_2 \neq 0$ implies they are all supported outside of Ω_Σ . Therefore,

$$\begin{aligned}
(4.10) \quad H_\#^\lambda(k_\parallel) p_{k_\parallel, I}^\lambda[n](\mathbf{x}) &= H_\#^\lambda(k_\parallel) e^{-i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n)} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n) \\
&\quad + \sum_{m_2 \in \mathbb{Z} \setminus \{0\}} \lambda^2 V_\#(\mathbf{x}) e^{-i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2)} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2) .
\end{aligned}$$

We may now use (4.9) with $\hat{\mathbf{v}} = \mathbf{v}_I^n = \mathbf{v}_I + n \mathbf{v}_1$ to simplify the first term on the right hand side of the previous equation. For all $\mathbf{x} \in \Omega_\Sigma$ with $n \geq 0$ and $I, J \in \{A, B\}$ with $I \neq J$, we obtain:

$$\begin{aligned}
H_\#^\lambda(k_\parallel) p_{k_\parallel, I}^\lambda[n](\mathbf{x}) &= \left(\lambda^2 \sum_{\substack{n_1 \geq 0 \\ n_1 \neq n}} V_0(\mathbf{x} - \mathbf{v}_I^{n_1}) \right) e^{-i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n)} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n) \\
&\quad + \left(\lambda^2 \sum_{n_1 \geq 0} V_0(\mathbf{x} - \mathbf{v}_J^{n_1}) \right) e^{-i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n)} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n) \\
&\quad + \sum_{m_2 \in \mathbb{Z} \setminus \{0\}} \lambda^2 V_\#(\mathbf{x}) e^{-i \frac{k_\parallel}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2)} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2) .
\end{aligned}$$

Thus,

$$\begin{aligned}
&\left| H_\#^\lambda(k_\parallel) p_{k_\parallel, I}^\lambda[n](\mathbf{x}) \right| \\
&\leq \left(\lambda^2 \sum_{\substack{n_1 \geq 0 \\ n_1 \neq n}} |V_0(\mathbf{x} - \mathbf{v}_I^{n_1})| \right) p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n) + \left(\lambda^2 \sum_{n_1 \geq 0} |V_0(\mathbf{x} - \mathbf{v}_J^{n_1})| \right) p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n) \\
&\quad + \sum_{m_2 \in \mathbb{Z} \setminus \{0\}} \lambda^2 |V_\#(\mathbf{x})| p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2) \\
(4.11) \quad &\equiv T_1(\mathbf{x}; n) + T_2(\mathbf{x}; n) + T_3(\mathbf{x}; n) .
\end{aligned}$$

To bound the first term of (4.11), we note that for $n_1 \neq n$

$$\begin{aligned} |V_0(\mathbf{x} - \mathbf{v}_I^{n_1})| p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n) &\leq \|V_0\|_\infty \mathbf{1}_{|\mathbf{x} - \mathbf{v}_I^{n_1}| < r_0} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n) \\ &\lesssim \mathbf{1}_{|\mathbf{x} - \mathbf{v}_I^{n_1}| < r_0} e^{-c\lambda|\mathbf{x} - \mathbf{v}_I^n|} \\ &\lesssim \mathbf{1}_{|\mathbf{x} - \mathbf{v}_I^{n_1}| < r_0} e^{-\frac{c}{2}\lambda|\mathbf{x} - \mathbf{v}_I^n|} e^{-\tilde{c}\lambda|n_1 - n|}. \end{aligned}$$

Summing over $n_1 \geq 0$ with $n_1 \neq n$ we obtain $T_1(\mathbf{x}; n) \lesssim e^{-c'\lambda} e^{-c''\lambda|\mathbf{x} - \mathbf{v}_I^n|}$. Very similarly we obtain: $T_2(\mathbf{x}; n) \lesssim e^{-c'\lambda} e^{-c''\lambda|\mathbf{x} - \mathbf{v}_I^n|}$. We finally consider $T_3(\mathbf{x}; n)$. For $\mathbf{x} \in \Omega_\Sigma$,

$$T_3(\mathbf{x}; n) \lesssim \lambda^2 \|V_0\|_\infty \sum_{m_2 \in \mathbb{Z} \setminus \{0\}} e^{-c\lambda|\mathbf{x} - \mathbf{v}_I^n|} e^{-c\lambda|m_2|} \lesssim e^{-c'\lambda} e^{-c''\lambda|\mathbf{x} - \mathbf{v}_I^n|}.$$

This completes the proof of Proposition 4.3.

4.1. The subspace $\mathcal{X}_{AB}^\lambda(k_\parallel)$. We introduce the closed subspace of $L^2(\Sigma)$:

$$(4.12) \quad \mathcal{X}_{AB}^\lambda(k_\parallel) = \text{the orthogonal complement in } L^2(\Sigma) \text{ of } \text{span} \left\{ p_{k_\parallel, I}^{\lambda, I}[n] : I = A, B; n \geq 0 \right\}.$$

We shall sometimes suppress the dependence on λ and write $\mathcal{X}_{AB}(k_\parallel)$. The space $L^2(\Sigma)$ may be decomposed as the orthogonal sum of subspaces:

$$(4.13) \quad L^2(\Sigma) = \text{span} \left\{ p_{k_\parallel, I}^\lambda[n] : I = A, B; n \geq 0 \right\} \oplus \mathcal{X}_{AB}(k_\parallel).$$

We also introduce the orthogonal projection onto $\mathcal{X}_{AB}(k_\parallel)$:

$$(4.14) \quad \Pi_{AB} = \Pi_{AB}(k_\parallel) : L^2(\Sigma) \rightarrow \mathcal{X}_{AB}(k_\parallel).$$

Since the set $\left\{ p_{k_\parallel, I}^\lambda[n] : I = A, B; n \geq 0 \right\}$ is only nearly-orthonormal for λ large (Proposition 4.2), we make use of the following:

Proposition 4.4. *There exists $\lambda_\star > 0$ such that for all $\lambda > \lambda_\star$ the following holds. Fix $k_\parallel \in [0, 2\pi]$.*

(1) *Then, for $F \in L^2(\Sigma)$ we have that*

$$F \equiv 0 \iff \Pi_{AB}(k_\parallel)F = 0 \text{ and } \left\langle p_{k_\parallel, I}^\lambda[n], F \right\rangle_{L^2(\Sigma)} = 0, \quad n \geq 0, I = A, B.$$

(2) *Any $\psi \in L^2(\Sigma)$ may be expressed in the form:*

$$(4.15) \quad \psi = \sum_{J=A,B} \sum_{n \geq 0} \alpha_n^J p_{k_\parallel, J}^\lambda[n] + \tilde{\psi},$$

where $\alpha = \{(\alpha_n^A, \alpha_n^B)^\top\}_{n \geq 0} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ and $\Pi_{AB}(k_\parallel)\tilde{\psi} = \tilde{\psi} \in \mathcal{X}_{AB}^\lambda(k_\parallel)$.

The proof is similar to that of Lemma 8.2 on page 31 of [26] and is omitted.

5. ENERGY ESTIMATES AND THE RESOLVENT

The following proposition concerns the invertibility of $\Pi_{AB}(k_{\parallel}) (H_{\sharp}^{\lambda}(k_{\parallel}) - \Omega) \Pi_{AB}(k_{\parallel})$ on $\mathcal{X}_{AB}(k_{\parallel})$ for λ sufficiently large. This will facilitate reduction of the edge state eigenvalue problem, (1.16) or (1.17), to a problem on the linear space $\text{span}\{p_{k_{\parallel},J}^{\lambda}[n] : I = A, B, n \geq 0\}$; see (4.12). The proof uses arguments analogous to those in [26]. The necessary modifications in the strategy are discussed at the end of this section.

Proposition 5.1. *There exist constants $\lambda_{\star} > 0$ (sufficiently large) and $c' > 0$ (sufficiently small), such that for all $\lambda > \lambda_{\star}$, $k_{\parallel} \in [0, 2\pi]$ and $|\Omega| \leq c'$ the following hold:*

(1) *For all $\varphi \in \mathcal{X}_{AB}(k_{\parallel})$, the equation*

$$(5.1) \quad \Pi_{AB}(k_{\parallel}) (H_{\sharp}^{\lambda}(k_{\parallel}) - \Omega) \psi = \varphi,$$

has a unique solution

$$\psi \equiv \mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel})\varphi \in \mathcal{X}_{AB} \cap H^2(\Sigma).$$

Thus, $\mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel})$ is the inverse of $\Pi_{AB}(k_{\parallel}) (H_{\sharp}^{\lambda}(k_{\parallel}) - \Omega) \Pi_{AB}(k_{\parallel})$ or equivalently $\Pi_{AB}(k_{\parallel}) (H_{\sharp}^{\lambda}(k_{\parallel}) - \Omega)$ acting on \mathcal{X}_{AB} .

(2) *The mapping $\varphi \mapsto \mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel})\varphi$ is a bounded linear operator :*

$$(5.2) \quad \mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel}) : \mathcal{X}_{AB}(k_{\parallel}) \rightarrow H^2(\Sigma) \cap \mathcal{X}_{AB}(k_{\parallel}).$$

(3) *We have the following operator norm bounds on $\mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel})$:*

$$(5.3) \quad \left\| \mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel}) \right\|_{\mathcal{X}_{AB} \rightarrow \mathcal{X}_{AB}} \lesssim 1$$

$$(5.4) \quad \lambda^{-1} \left\| \nabla_{\mathbf{x}} \mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel}) \right\|_{\mathcal{X}_{AB} \rightarrow \mathcal{X}_{AB}} \lesssim 1$$

$$(5.5) \quad \left\| \mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel}) \right\|_{\mathcal{X}_{AB} \rightarrow H^2(\Sigma) \cap \mathcal{X}_{AB}} \leq C(\lambda, k_{\parallel}).$$

(4) *Furthermore, this mapping depends analytically on $\Omega \in \mathbb{C}$ for $|\Omega| < c'$, and for all such Ω :*

$$(5.6) \quad \left\| \partial_{\Omega} \mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel}) \right\|_{\mathcal{X}_{AB} \rightarrow \mathcal{X}_{AB}} \lesssim 1.$$

(5) *For real $\Omega \in (-c', c')$, $\mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel})$ is self-adjoint on the Hilbert space \mathcal{X}_{AB} , endowed with the $L^2(\Sigma)$ inner product.*

A key step to proving Proposition 5.1 is the following energy estimate on the space $\mathcal{X}_{AB}(k_{\parallel})$:

Proposition 5.2 (Energy Estimate). *Fix $k_{\parallel} \in [0, 2\pi]$. There exists $\lambda_{\star} > 0$, independent of k_{\parallel} , and a constant $C_{\star} > 0$ such that the following holds for all $\lambda \geq \lambda_{\star}$. Let $\psi \in \mathcal{X}_{AB}(k_{\parallel}) \cap H^2(\Sigma)$. That is,*

$$(5.7) \quad \left\langle p_{k_{\parallel},J}^{\lambda}[n], \psi \right\rangle_{L^2(\Sigma)} = 0, \quad n \geq 0, \quad J = A, B.$$

Then,

$$(5.8) \quad \left\| H_{\sharp}^{\lambda}(k_{\parallel})\psi \right\|_{L^2(\Sigma)}^2 \geq c_{\star} \left(\|\psi\|_{L^2(\Sigma)}^2 + \lambda^{-2} \|\nabla\psi\|_{L^2(\Sigma)}^2 \right).$$

The constant c_* can be taken independent of k_{\parallel} but it does depend on properties of the atomic potential, V_0 , in particular on the constants c_{gs} and c_{gap} ; see (3.4) and (3.6).

The proof of Proposition 5.1 follows the general structure of the proof of the energy estimates in [26]. We now discuss the modifications in these arguments, which are required to prove Propositions 5.2 and 5.1. We follow the discussion of Section 9 of [26] with $\Sigma = \mathbb{R}^2/\mathbb{Z}\mathbf{v}_2$ playing the role of \mathbb{R}^2/Λ , and with the approximate eigenfunctions $p_{k_{\parallel},I}^{\lambda}[n] \in L^2(\Sigma)$ playing the role of $p_{k,I}^{\lambda} \in L^2(\mathbb{R}^2/\Lambda)$ in [26].

For $n \geq 0$, let \mathbf{x}_I^n , $I = A, B$ denote the two atomic sites in Ω_n , where $n \geq 0$. Recall Ω_{Σ} is the union, for $n \geq -1$, over all Ω_n ; see Figure 1. In place of the partitions of unity (9.11) in [26] on \mathbb{R}^2/Λ , we introduce here analogous partitions on Σ :

$$1 = \Theta_0^2 + \sum_{\substack{n \geq 0 \\ I=A,B}} \Theta_{n,I}^2, \quad 1 = \tilde{\Theta}_0^2 + \sum_{\substack{n \geq 0 \\ I=A,B}} \tilde{\Theta}_{n,I}^2$$

where $\Theta_{n,I}$ and $\tilde{\Theta}_{n,I}$ are supported near \mathbf{x}_I^n . All the arguments in Sections 9.1 through 9.4 of [26] go through in the above setting, with minimal changes. This gives Proposition 5.2.

We seek to show that the inverse of $\Pi_{AB}(k_{\parallel}) (H_{\#}^{\lambda}(k_{\parallel}) - \Omega) \Pi_{AB}(k_{\parallel})$, is a bounded linear operator on $\mathcal{X}_{AB}^{\lambda}(k_{\parallel})$, satisfying the bound (5.3) and (5.4) and furthermore that $\mathcal{K}_{\#}^{\lambda}(\Omega, k_{\parallel})$ maps $\mathcal{X}_{AB}^{\lambda}(k_{\parallel})$ to $H^2(\Sigma) \cap \mathcal{X}_{AB}^{\lambda}(k_{\parallel})$ and satisfies the operator bound (5.5).

To adapt Section 9.5 of [26] to our setting requires an additional argument which we now supply. Suppose we have $\Pi_{AB}(k_{\parallel}) [H_{\#}^{\lambda}(k_{\parallel}) - \Omega I] \psi = f$, where $\psi \in L^2(\Sigma) \cap \mathcal{X}_{AB}^{\lambda}(k_{\parallel})$ and $f \in L^2(\Sigma)$. Then, for some $\{\alpha_{I,n}\}$, ($I = A, B$ $n \geq 0$), in $l^2(\mathbb{N}_0; \mathbb{C}^2)$:

$$(5.9) \quad [H_{\#}^{\lambda}(k_{\parallel}) - \Omega I] \psi = f + \sum_{\substack{I=A,B \\ n \geq 0}} \alpha_{I,n} p_{k_{\parallel},I}^{\lambda}[n],$$

where the right hand sum is convergent in $L^2(\Sigma)$ and the left hand side is interpreted as a distribution on Σ . Taking the inner product in $L^2(\Sigma)$ of (9.1) with $p_{k_{\parallel},J}^{\lambda}[m]$, we find that

$$(5.10) \quad \sum_{\substack{I=A,B \\ n \geq 0}} \alpha_{I,n} \left\langle p_{k_{\parallel},J}^{\lambda}[m], p_{k_{\parallel},I}^{\lambda}[n] \right\rangle = \xi_{k_{\parallel},J}^{\lambda}[m], \quad \text{where}$$

$$\xi_{k_{\parallel},J}^{\lambda}[m] \equiv \left\langle H_{\#}^{\lambda}(k_{\parallel}) p_{k_{\parallel},J}^{\lambda}[m], \psi \right\rangle - \left\langle p_{k_{\parallel},J}^{\lambda}[m], f \right\rangle.$$

We have

$$(5.11) \quad \left| \xi_{k_{\parallel},J}^{\lambda}[m] \right|^2 \lesssim \left| \left\langle H_{\#}^{\lambda}(k_{\parallel}) p_{k_{\parallel},J}^{\lambda}[m], \psi \right\rangle \right|^2 + \left| \left\langle p_{k_{\parallel},J}^{\lambda}[m], f \right\rangle \right|^2$$

and summing over $J = A, B$ and $m \geq 0$ yields

$$(5.12) \quad \sum_{\substack{J=A,B \\ m \geq 0}} \left| \xi_{k_{\parallel},J}^{\lambda}[m] \right|^2 \lesssim \sum_{\substack{J=A,B \\ m \geq 0}} \left| \left\langle H_{\#}^{\lambda}(k_{\parallel}) p_{k_{\parallel},J}^{\lambda}[m], \psi \right\rangle \right|^2 + \sum_{\substack{J=A,B \\ m \geq 0}} \left| \left\langle p_{k_{\parallel},J}^{\lambda}[m], f \right\rangle \right|^2.$$

In order to bound the second term on the right in (5.12), note that the near-orthonormality of the set $\{p_{k_{\parallel},J}^{\lambda}[m] : J = A, B, m \geq 0\}$ for λ large (Proposition 4.2) implies the Bessel-type

inequality:

$$\sum_{\substack{J=A,B \\ m \geq 0}} \left| \left\langle p_{k_{\parallel},J}^{\lambda}[m], f \right\rangle \right|^2 \lesssim \|f\|_{L^2(\Sigma)}^2.$$

Consider next the first term on the right in (5.12). Thanks to the pointwise bound on $H_{\sharp}^{\lambda}(k_{\parallel})p_{k_{\parallel},J}^{\lambda}[m](\mathbf{x})$ from Proposition 4.3, a Young-type inequality yields:

$$\sum_{\substack{J=A,B \\ m \geq 0}} \left| \left\langle H_{\sharp}^{\lambda}(k_{\parallel})p_{k_{\parallel},J}^{\lambda}[m], \psi \right\rangle \right|^2 \lesssim e^{-c\lambda} \|\psi\|_{L^2(\Sigma)}^2.$$

Again, by Proposition 4.2, we have

$$(5.13) \quad \sum_{\substack{J=A,B \\ m \geq 0}} |\alpha_m^J|^2 \lesssim \sum_{\substack{J=A,B \\ m \geq 0}} |\xi_{k_{\parallel},J}^{\lambda}[m]|^2 \lesssim e^{-c\lambda} \|\psi\|_{L^2(\Sigma)}^2 + C \|f\|_{L^2(\Sigma)}^2.$$

And finally one more application of Proposition 4.2 gives

$$(5.14) \quad \left\| \sum_{\substack{I=A,B \\ n \geq 0}} \alpha_{k_{\parallel},I}^{\lambda}[n] p_{k_{\parallel},I}^{\lambda}[n] \right\|_{L^2(\Sigma)} \lesssim e^{-c\lambda} \|\psi\|_{L^2(\Sigma)} + C \|f\|_{L^2(\Sigma)}.$$

The estimates (5.13) and (5.14) allow us to argue as in Section 9.5 of [26], using our energy estimates, that the operator $\mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel})$, the inverse of $\Pi_{AB}(k_{\parallel}) (H_{\sharp}^{\lambda}(k_{\parallel}) - \Omega) \Pi_{AB}(k_{\parallel})$, is a bounded linear operator on $\mathcal{X}_{AB}^{\lambda}(k_{\parallel})$, satisfying the bounds (5.3) and (5.4).

To complete the proof of Proposition 5.1 must show that $\mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel})$ maps $\mathcal{X}_{AB}^{\lambda}(k_{\parallel})$ to $H^2(\mathbb{R}^2) \cap \mathcal{X}_{AB}^{\lambda}(k_{\parallel})$. To bound $\|\Delta\psi\|_{L^2(\Sigma)}$, we use (9.1) to obtain an expression for $\Delta\psi$ in terms of ψ and $\nabla\psi$. Then, the energy estimate for $\|\psi\|_{L^2(\Sigma)}$ and $\|\nabla\psi\|_{L^2(\Sigma)}$, and the bound (5.14) imply that for λ sufficiently large, the $L^2(\Sigma)$ norm of each term in the expression $\Delta\psi$ can be bounded by $C(\lambda) \times \|f\|_{L^2(\Sigma)}$, where $C(\lambda)$ denotes a λ -dependent constant.

6. LYAPUNOV-SCHMIDT REDUCTION; FORMULATION AS A PROBLEM IN $\mathcal{X}_{AB}^{\lambda}(k_{\parallel})$

The resolvent bounds of Proposition 5.1 ensure that on the subspace $\mathcal{X}_{AB}(k_{\parallel})$, the operator $H_{\sharp}^{\lambda}(k_{\parallel}) - \Omega$ is invertible in a neighborhood of $\Omega = 0$, *i.e.* the spectrum of $\Pi_{AB}(k_{\parallel})H_{\sharp}^{\lambda}(k_{\parallel})\Pi_{AB}(k_{\parallel})$ is bounded away from zero, uniformly in $\lambda \gg 1$. In this section, we make use of this spectral separation to obtain a reduction of the $L_{k_{\parallel}}^2$ eigenvalue problem to a problem on the subspace of $L^2(\Sigma)$ given by: $\text{span}\{p_{k_{\parallel}}^{\lambda,I}[n] : I = A, B; n \geq 0\}$.

Consider the eigenvalue problem:

$$(6.1) \quad \left(- \left(\nabla + i \frac{k_{\parallel}}{2\pi} \mathfrak{R}_2 \right)^2 + \lambda^2 V_{\sharp}(\mathbf{x}) \right) \psi = E\psi, \quad \psi \in H^2(\Sigma).$$

Let

$$(6.2) \quad E = E_0^{\lambda} + \Omega$$

Recall the centered edge-Hamiltonian:

$$(6.3) \quad H_{\#}^{\lambda}(k_{\parallel}) = - \left(\nabla + i \frac{k_{\parallel}}{2\pi} \mathfrak{K}_2 \right)^2 + \lambda^2 V_{\#}(\mathbf{x}) - E_0^{\lambda} ;$$

see also (3.10). Then, the eigenvalue problem may be rewritten as:

$$(6.4) \quad (H_{\#}^{\lambda}(k_{\parallel}) - \Omega) \psi = 0, \quad \psi \in H^2(\Sigma) .$$

By Proposition 4.4 any $\psi \in H^2(\Sigma)$ may be written in the form:

$$(6.5) \quad \psi = \sum_{I=A,B} \sum_{n \geq 0} \alpha_n^I p_{k_{\parallel},I}^{\lambda}[n] + \tilde{\psi},$$

where $\alpha = \{(\alpha_n^A, \alpha_n^B)^{\top}\}_{n \geq 0} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ and $\Pi_{AB}(k_{\parallel})\tilde{\psi} = \tilde{\psi}$. We adopt the convention

$$\alpha_n^I = 0, \quad n \leq -1, \quad I = A, B.$$

Substitution of (6.5) into (6.4) yields:

$$(6.6) \quad \sum_{I=A,B} \sum_{n \geq 0} \alpha_n^I (H_{\#}^{\lambda}(k_{\parallel}) - \Omega) p_{k_{\parallel},I}^{\lambda}[n] + (H_{\#}^{\lambda}(k_{\parallel}) - \Omega) \tilde{\psi} = 0 .$$

By part (1) of Proposition 4.4, the eigenvalue problem (6.4) is seen to be equivalent to the system obtained by: (i) applying the orthogonal projection $\Pi_{AB}(k_{\parallel})$ to (6.6):

$$(6.7) \quad \Pi_{AB}(k_{\parallel}) (H_{\#}^{\lambda}(k_{\parallel}) - \Omega) \tilde{\psi} + \sum_{I=A,B} \sum_{n \geq 0} \alpha_n^I \Pi_{AB}(k_{\parallel}) (H_{\#}^{\lambda}(k_{\parallel}) - \Omega) p_{k_{\parallel},I}^{\lambda}[n] = 0$$

and (ii) taking the inner product of (6.6) with the states: $p_{k_{\parallel},J}^{\lambda}[m]$; $m \geq 0$, $J = A, B$:

$$(6.8) \quad \left\langle p_{k_{\parallel},J}^{\lambda}[m], \sum_{I=A,B} \sum_{n \geq 0} \alpha_n^I (H_{\#}^{\lambda}(k_{\parallel}) - \Omega) p_{k_{\parallel},I}^{\lambda}[n] \right\rangle + \left\langle (H_{\#}^{\lambda}(k_{\parallel}) - \bar{\Omega}) p_{k_{\parallel},I}^{\lambda}[m], \tilde{\psi} \right\rangle = 0$$

where $m = 0, 1, 2, \dots$

Using Proposition 5.1 we solve (6.7) for $\tilde{\psi}$ as a function of $\alpha = (\alpha^A, \alpha^B)^{\top} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$:

$$(6.9) \quad \tilde{\psi} = - \sum_{I=A,B} \sum_{n \geq 0} \alpha_n^I \mathcal{K}_{\#}^{\lambda}(\Omega, k_{\parallel}) \Pi_{AB}(k_{\parallel}) H_{\#}^{\lambda}(k_{\parallel}) p_{k_{\parallel},I}^{\lambda}[n] .$$

Here we have used that $\Pi_{AB}(k_{\parallel}) p_{k_{\parallel},I}^{\lambda}[n] = 0$. Substitution of (6.9) into (6.8) yields

$$(6.10) \quad \sum_{I=A,B} \sum_{n \geq 0} \mathcal{M}_{JI}^{\lambda, k_{\parallel}}[m, n](\Omega, k_{\parallel}) \alpha_n^I = 0; \quad J = A, B, \quad m \geq 0 ,$$

where

$$\begin{aligned}
& \mathcal{M}_{JI}^\lambda[m, n](\Omega, k_\parallel) \\
& \equiv \left\langle p_{k_\parallel, J}^\lambda[m], (H_\#^\lambda(k_\parallel) - \Omega) p_{k_\parallel, I}^\lambda[n] \right\rangle_{L^2(\Sigma)} \\
& \quad - \left\langle H_\#^\lambda(k_\parallel) p_{k_\parallel, J}^\lambda[m], \Pi_{AB}(k_\parallel) \mathcal{K}_\#^\lambda(\Omega, k_\parallel) \Pi_{AB}(k_\parallel) H_\#^\lambda(k_\parallel) p_{k_\parallel, I}^\lambda[n] \right\rangle_{L^2(\Sigma)}.
\end{aligned} \tag{6.11}$$

Remark 6.1. For fixed $J = A$ or B and fixed $m \geq 0$, the equation (6.10) expresses the interaction of all atomic A - and B - sites within the cylinder, Σ , with the atomic site J in cell m . In particular, the $\mathcal{M}_{JA}[m, n]$ are interaction coefficients between site J in Ω_m and all sites \mathbf{v}_A^n , $n \geq 0$, and $\mathcal{M}_{JB}[m, n]$ are interaction coefficients between site J in cell Ω_m and all sites \mathbf{v}_B^n , $n \geq 0$.

Due to their dependence on the Hamiltonian, $H_\#^\lambda$, we refer to the first term on the right in (6.11) as the linear matrix elements, $\mathcal{M}^{\lambda, \text{lin}}[m, n](\Omega, k_\parallel)$ and second term on the right in (6.11) as the non-linear matrix elements, $\mathcal{M}^{\lambda, \text{nl}}[m, n](\Omega, k_\parallel)$. Thus,

$$\mathcal{M}^\lambda[m, n](\Omega, k_\parallel) \equiv \mathcal{M}^{\lambda, \text{lin}}[m, n](\Omega, k_\parallel) - \mathcal{M}^{\lambda, \text{nl}}[m, n](\Omega, k_\parallel). \tag{6.12}$$

In the subsequent sections we compute highly accurate approximations to the linear (Section 7) and non-linear (Section 12) matrix elements. This will enable us to recast and solve (6.10) as a perturbation of a tight-binding model for λ sufficiently large (Section 8).

7. MATRIX ELEMENTS $\mathcal{M}_{JI}^{\lambda, \text{lin}}[m, n](\Omega, k_\parallel)$ AND $\mathcal{M}_{JI}^{\lambda, \text{nl}}[m, n](\Omega, k_\parallel)$

In this section we provide expansions of the matrix entries of $\mathcal{M}_{JI}^{\lambda, \text{lin}}[m, n](\Omega, k_\parallel)$. Recall that

$$P_{k_\parallel, I}^\lambda[n](\mathbf{x}) \equiv e^{i \frac{k_\parallel}{2\pi} \mathfrak{R}_2 \cdot (\mathbf{x} - \mathbf{v}_I)} p_{k_\parallel, I}^\lambda[n](\mathbf{x}) = \sum_{m_2 \in \mathbb{Z}} e^{ik_\parallel m_2} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^n - m_2 \mathbf{v}_2); \tag{7.1}$$

(see also (4.4)) and that $H_\#^\lambda = -\Delta + \lambda^2 V_\#(\mathbf{x}) - E_0^\lambda$.

In preparation for our expansions, introduce the *nearest-neighbor hopping coefficient*:

$$\rho_\lambda = \int_{B_{r_0}(0)} p_0^\lambda(\mathbf{y}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} + \mathbf{e}) d\mathbf{y} = \int_{\mathbb{R}^2} p_0^\lambda(\mathbf{y}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} + \mathbf{e}) d\mathbf{y}, \tag{7.2}$$

where $\mathbf{e} = \mathbf{v}_B - \mathbf{v}_A$. The latter equality holds since V_0 has compact support in $B_{r_0}(0)$. We further recall the bounds (3.3) :

$$e^{-c_- \lambda} \lesssim \rho_\lambda \lesssim e^{-c_+ \lambda} \tag{7.3}$$

for some constants $c_-, c_+ > 0$ and all $\lambda > 0$ sufficiently large; this was proved in [26].

The main results of this section (Propositions 7.1 and 7.2) are the following two propositions which (i) isolate the dominant (nearest neighbor) behavior of the linear matrix elements and provide estimates on the corrections, and (ii) estimate the nonlinear matrix elements.

Proposition 7.1 (Expansion of linear matrix elements).

For all $\lambda > \lambda_*$ (sufficiently large), and all $k_\parallel \in [0, 2\pi]$, we have:

(1) For $m \geq 0$,

(7.4)

$$\left\langle P_{k_{\parallel},B}^{\lambda}[m], H_{\sharp}^{\lambda} P_{k_{\parallel},A}^{\lambda}[m] \right\rangle_{L^2(\Sigma)} = \left\langle p_{k_{\parallel},B}^{\lambda}[m], H_{\sharp}^{\lambda}(k_{\parallel}) p_{k_{\parallel},A}^{\lambda}[m] \right\rangle_{L^2(\Sigma)} = -\rho_{\lambda} (1 + e^{ik_{\parallel}}) + \mathcal{O}(e^{-c\lambda} \rho_{\lambda}),$$

(7.5)

$$\left\langle P_{k_{\parallel},A}^{\lambda}[m], H_{\sharp}^{\lambda} P_{k_{\parallel},B}^{\lambda}[m] \right\rangle_{L^2(\Sigma)} = \overline{\left\langle P_{k_{\parallel},B}^{\lambda}[m], H_{\sharp}^{\lambda} P_{k_{\parallel},A}^{\lambda}[m] \right\rangle_{L^2(\Sigma)}} = -\rho_{\lambda} (1 + e^{-ik_{\parallel}}) + \mathcal{O}(e^{-c\lambda} \rho_{\lambda}).$$

(2) For $m \geq 0$,

$$(7.6) \quad \left\langle P_{k_{\parallel},B}^{\lambda}[m], H_{\sharp}^{\lambda} P_{k_{\parallel},A}^{\lambda}[m+1] \right\rangle_{L^2(\Sigma)} = -\rho_{\lambda} + \mathcal{O}(e^{-c\lambda} \rho_{\lambda}),$$

and for $m \geq 1$

$$(7.7) \quad \left\langle P_{k_{\parallel},A}^{\lambda}[m], H_{\sharp}^{\lambda} P_{k_{\parallel},B}^{\lambda}[m-1] \right\rangle_{L^2(\Sigma)} = -\rho_{\lambda} + \mathcal{O}(e^{-c\lambda} \rho_{\lambda}).$$

(3)

$$(7.8) \quad \left\langle P_{k_{\parallel},B}^{\lambda}[m], H_{\sharp}^{\lambda} P_{k_{\parallel},A}^{\lambda}[n] \right\rangle_{L^2(\Sigma)} = \mathcal{O}(e^{-c\lambda|m-n|} \rho_{\lambda}), \quad m, n \geq 0, \quad n \neq m, m+1,$$

$$(7.9) \quad \left\langle P_{k_{\parallel},A}^{\lambda}[m], H_{\sharp}^{\lambda} P_{k_{\parallel},B}^{\lambda}[n] \right\rangle_{L^2(\Sigma)} = \mathcal{O}(e^{-c\lambda|m-n|} \rho_{\lambda}), \quad m, n \geq 0, \quad n \neq m, m-1.$$

(4) For $m, n \geq 0$ and $I = A$ or B

$$(7.10) \quad \left\langle P_{k_{\parallel},I}^{\lambda}[m], H_{\sharp}^{\lambda} P_{k_{\parallel},I}^{\lambda}[n] \right\rangle_{L^2(\Sigma)} = \mathcal{O}(e^{-c\lambda} e^{-c\lambda|m-n|} \rho_{\lambda}).$$

The implied constants in the $\mathcal{O}(\cdot)$ estimates and the constants λ_{\star} and c are independent of k_{\parallel} .

We note, by part (4) of Proposition 5.1, that the function

$$\Omega \mapsto \left\langle H_{\sharp}^{\lambda}(k_{\parallel}) p_{k_{\parallel},J}^{\lambda}[n], \Pi_{AB}^{\lambda}(k_{\parallel}) \mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel}) \Pi_{AB}^{\lambda}(k_{\parallel}) H_{\sharp}^{\lambda}(k_{\parallel}) p_{k_{\parallel},I}^{\lambda}[m] \right\rangle_{L^2(\Sigma)}$$

is analytic for $|\Omega| < c'$.

Proposition 7.2 (Estimation of nonlinear matrix element contributions). *There exists $\lambda > \lambda_{\star}$ (sufficiently large), such that for all $k_{\parallel} \in [0, 2\pi]$ and $|\Omega| \leq e^{-c'\lambda}$ (c' , a sufficiently small constant determined by V_0) the following holds for $j = 0, 1$:*

$$(7.11) \quad \left| \left\langle H_{\sharp}^{\lambda}(k_{\parallel}) p_{k_{\parallel},J}^{\lambda}[n], \Pi_{AB}^{\lambda}(k_{\parallel}) \partial_{\Omega}^j \mathcal{K}_{\sharp}^{\lambda}(\Omega, k_{\parallel}) \Pi_{AB}^{\lambda}(k_{\parallel}) H_{\sharp}^{\lambda}(k_{\parallel}) p_{k_{\parallel},I}^{\lambda}[m] \right\rangle_{L^2(\Sigma)} \right| \lesssim \rho_{\lambda} e^{-c\lambda} e^{-c|n-m|}.$$

The implied constants in the $\mathcal{O}(\cdot)$ estimates and the constants λ_{\star} and c are independent of k_{\parallel} .

Proposition 7.1 is proved in Section 11 and Proposition 7.2 in Section 12. The proof of Proposition 7.2 requires detailed information on the resolvent, which we need to control in weighted spaces. We obtain this control by constructing the resolvent kernel and obtaining pointwise bounds for it. The construction is carried out in Section 10.

8. EXISTENCE OF ZIGZAG EDGE STATES IN THE STRONG BINDING REGIME

In this section we apply Propositions 7.1 and 7.2 to rewrite the edge state eigenvalue problem as a perturbation of the eigenvalue problem for the tight-binding limiting operator studied in Section 2. We then use this reformulation to construct zigzag edge states for arbitrary $\lambda > \lambda_*$, where λ_* is fixed and sufficiently large.

Recall our reduction, for $k_{\parallel} \in \mathcal{J} \subset \subset (2\pi/3, 4\pi/3)$, of the edge state eigenvalue problem for $H_{\sharp}^{\lambda}(k_{\parallel})$ to the discrete eigenvalue problem for $\{(\alpha_m^A, \alpha_m^B)\}_{m \geq 0}$ in $l^2(\mathbb{N}_0; \mathbb{C}^2)$:

$$(8.1) \quad \sum_{I=A,B} \sum_{n \geq 0} \mathcal{M}_{JI}^{\lambda}[m, n](\Omega, k_{\parallel}) \alpha_n^I = 0; \quad J = A, B, \quad m \geq 0,$$

Let's cast (8.1) in a form in which the tight-binding operator $H_{\sharp}^{\text{TB}}(k_{\parallel})$ is made explicit. First, (8.1) is equivalent to the following system for $m \geq 0$:

$$(8.2) \quad \begin{aligned} \sum_{n \geq 0} \mathcal{M}_{AA}^{\lambda}[m, n](\Omega, k_{\parallel}) \alpha_n^A + \sum_{n \geq 0} \mathcal{M}_{AB}^{\lambda}[m, n](\Omega, k_{\parallel}) \alpha_n^B &= 0, \\ \sum_{n \geq 0} \mathcal{M}_{BA}^{\lambda}[m, n](\Omega, k_{\parallel}) \alpha_n^A + \sum_{n \geq 0} \mathcal{M}_{BB}^{\lambda}[m, n](\Omega, k_{\parallel}) \alpha_n^B &= 0. \end{aligned}$$

To isolate the dominant terms (see Propositions 7.1 and 7.2), we rearrange the expressions and obtain for $m \geq 0$:

$$(8.3) \quad \begin{aligned} &\mathcal{M}_{AB}^{\lambda}[m, m-1](\Omega, k_{\parallel}) \alpha_{m-1}^B + \mathcal{M}_{AB}^{\lambda}[m, m](\Omega, k_{\parallel}) \alpha_m^B + \mathcal{M}_{AA}^{\lambda}[m, m](\Omega, k_{\parallel}) \alpha_m^A \\ &= - \sum_{\substack{n \geq 0 \\ n \neq m, m-1}} \mathcal{M}_{AB}^{\lambda}[m, n](\Omega, k_{\parallel}) \alpha_n^B - \sum_{\substack{n \geq 0 \\ n \neq m}} \mathcal{M}_{AA}^{\lambda}[m, n](\Omega, k_{\parallel}) \alpha_n^A \\ &\mathcal{M}_{BA}^{\lambda}[m, m](\Omega, k_{\parallel}) \alpha_m^A + \mathcal{M}_{BA}^{\lambda}[m, m+1](\Omega, k_{\parallel}) \alpha_{m+1}^A + \mathcal{M}_{BB}^{\lambda}[m, m](\Omega, k_{\parallel}) \alpha_m^B \\ &= - \sum_{\substack{n \geq 0 \\ n \neq m, m+1}} \mathcal{M}_{BA}^{\lambda}[m, n](\Omega, k_{\parallel}) \alpha_n^A - \sum_{\substack{n \geq 0 \\ n \neq m}} \mathcal{M}_{BB}^{\lambda}[m, n](\Omega, k_{\parallel}) \alpha_n^B. \end{aligned}$$

Here, $\mathcal{M}_{JI}^{\lambda}[m, n]$ is given by (6.11), where we take $\mathcal{M}_{BA}^{\lambda}[m, m-1] = 0$ for $m = 0$. The system (8.3) is equivalent to (8.1).

Our next step will be to express the matrix elements on the left hand side of (8.3), using Proposition 4.2, Proposition 7.1 and Proposition 7.2. Since the leading order expressions are proportional to ρ_{λ} , it is natural to introduce the rescaled energy:

$$(8.4) \quad \Omega \equiv \rho_{\lambda} \tilde{\Omega}.$$

Recall our general upper and lower bounds on ρ_{λ} : $e^{-c-\lambda} \lesssim \rho_{\lambda} \lesssim e^{-c+\lambda}$ (see (7.3) or (3.3)) and let $\hat{c} > c_- > 0$ denote the positive constant introduced in Remark 3.1. We now constrain Ω to satisfy $|\Omega| < e^{-\hat{c}\lambda}$. Then, $|\tilde{\Omega}| = |\rho_{\lambda}^{-1}\Omega| \leq e^{-(\hat{c}-c_-)\lambda} < e^{-c''\lambda}$, where c'' is a small positive constant, for any finite λ sufficiently large.

Using Proposition 4.2, Proposition 7.1 and Proposition 7.2 in (8.3) we obtain after dividing by $-\rho_\lambda$:

$$\begin{aligned}
& (-1 + \mathcal{O}(e^{-c\lambda})) \alpha_{m-1}^B + (-1 + e^{-ik_\parallel}) + \mathcal{O}(e^{-c\lambda}) \alpha_m^B + (-1 + \mathcal{O}(e^{-c\lambda})) \tilde{\Omega} \alpha_m^A \\
&= \sum_{\substack{n \geq 0 \\ n \neq m, m-1}} \mathcal{O}(e^{-c\lambda} e^{-c|m-n|}) \alpha_n^B + \sum_{n \geq 0} \mathcal{O}(e^{-c\lambda} e^{-c|m-n|}) \alpha_n^A, \\
(8.5)
\end{aligned}$$

where $\alpha_{m-1}^B = 0$ for $m = 0$, and

$$\begin{aligned}
& (-1 + e^{ik_\parallel}) + \mathcal{O}(e^{-c\lambda}) \alpha_m^A + (-1 + \mathcal{O}(e^{-c\lambda})) \alpha_{m+1}^A + (-1 + \mathcal{O}(e^{-c\lambda})) \tilde{\Omega} \alpha_m^B \\
&= \sum_{\substack{n \geq 0 \\ n \neq m, m+1}} \mathcal{O}(e^{-c\lambda} e^{-c|m-n|}) \alpha_n^A + \sum_{n \geq 0} \mathcal{O}(e^{-c\lambda} e^{-c|m-n|}) \alpha_n^B, \\
(8.6)
\end{aligned}$$

where $|\tilde{\Omega}| < c''$.

Remark 8.1. By Proposition 5.1 (part 4) and Proposition 7.2, the expressions in (8.5)-(8.6) of the form $\mathcal{O}(g(\lambda))$ are analytic functions of $\tilde{\Omega}$ for $\tilde{\Omega}$ varying in the open subset of \mathbb{C} : $|\Omega| < e^{-\hat{c}\lambda}$. Moreover, these expressions are all uniformly bounded by $g(\lambda)$ for all $\tilde{\Omega}$ such that $|\tilde{\Omega}| < c''$, a small positive constant.

We obtain, for $m \geq 0$ and $|\tilde{\Omega}| < c''$:

$$\begin{aligned}
& -\alpha_{m-1}^B - (1 + e^{-ik_\parallel}) \alpha_m^B - \tilde{\Omega} \alpha_m^A \\
(8.7) \quad &= \sum_{n \geq 0} \mathcal{O}(e^{-c\lambda} e^{-c|m-n|}) \alpha_n^B + \sum_{n \geq 0} \mathcal{O}(e^{-c\lambda} e^{-c|m-n|}) \alpha_n^A, \\
& \text{where } \alpha_{m-1}^B = 0 \text{ for } m = 0, \text{ and}
\end{aligned}$$

$$\begin{aligned}
& -(1 + e^{ik_\parallel}) \alpha_m^A - \alpha_{m+1}^A - \tilde{\Omega} \alpha_m^B \\
(8.8) \quad &= \sum_{n \geq 0} \mathcal{O}(e^{-c\lambda} e^{-c|m-n|}) \alpha_n^B + \sum_{n \geq 0} \mathcal{O}(e^{-c\lambda} e^{-c|m-n|}) \alpha_n^A.
\end{aligned}$$

Again we remark, as in Remark 8.1, that in (8.7)-(8.8) expressions of the form $\mathcal{O}(g(\lambda))$ are analytic in $\tilde{\Omega}$ and uniformly bounded by $g(\lambda)$ for $|\tilde{\Omega}| < c''$.

The system (8.7)-(8.8) is of the form:

$$(8.9) \quad \left[\left(H_\#^{\text{TB}}(k_\parallel) - \tilde{\Omega} \right) \begin{pmatrix} \alpha^A \\ \alpha^B \end{pmatrix} \right]_m = \left[\mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \begin{pmatrix} \alpha^A \\ \alpha^B \end{pmatrix} \right]_m, \quad \text{for } m \geq 0,$$

where $H_{\#}^{\text{TB}}(k_{\parallel})$ is the tight binding Hamiltonian for a zigzag termination of \mathbb{H} , studied in Section 2; see, in particular, (2.4), (2.5) ² Note that our definition of $H_{\#}^{\text{TB}}(k_{\parallel})$ implies that the scaled spectral parameter, $\tilde{\Omega}$, appears with a plus rather than a minus sign in (8.9).

Furthermore, using that the mapping $\{\gamma_m\}_{m \geq 0} \mapsto \left\{ \sum_{n \geq 0} e^{-c|m-n|} \gamma_n \right\}_{m \geq 0}$ is bounded on $l^2(\mathbb{N}_0)$, we have that the mapping $\tilde{\Omega} \mapsto \mathcal{P}(\lambda; \rho_{\lambda} \tilde{\Omega})$ is an analytic mapping for $|\tilde{\Omega}| < c''$ with values in the space of bounded linear operators on $l^2(\mathbb{N}_0; \mathbb{C}^2)$. We also have, for all $|\tilde{\Omega}| \leq c'$, ($c' < c''$):

$$(8.10) \quad \|\mathcal{P}(\lambda; \rho_{\lambda} \tilde{\Omega})\|_{l^2 \rightarrow l^2} \lesssim e^{-c\lambda},$$

where the implied constant is independent of $\tilde{\Omega}$, but depends on c' . Recall that k_{\parallel} varies in a compact subinterval of $(2\pi/3, 4\pi/3)$, where $\delta_{\text{gap}}(k_{\parallel}) = \left|1 - |\zeta(k_{\parallel})|\right| = \left|1 - |1 + e^{ik_{\parallel}}|\right| > 0$. We will further restrict $\tilde{\Omega}$ to satisfy $|\tilde{\Omega}| < c' < \delta_{\text{gap}}(k_{\parallel})$.

Our goal is to construct, for all λ sufficiently large, a solution of (8.9):

$$(8.11) \quad \begin{aligned} \lambda &\mapsto \vec{\alpha}(\lambda) = (\alpha^A(\lambda), \alpha^B(\lambda)) \in l^2(\mathbb{N}_0; \mathbb{C}^2) \\ \lambda &\mapsto \tilde{\Omega}(\lambda), \quad \text{such that } |\tilde{\Omega}(\lambda)| \lesssim e^{-c\lambda} \leq c'. \end{aligned}$$

Given the mappings (8.11), equations (6.5), (6.9) and the relation $E = E_0^{\lambda} + \rho_{\lambda} \tilde{\Omega}$ define a solution to the $L_{k_{\parallel}}^2(\Sigma)$ edge state eigenvalue problem, $\Psi_{k_{\parallel}}^{\lambda}(\mathbf{x}) = e^{i\frac{k_{\parallel}}{2\pi} \mathfrak{R}^2 \cdot \mathbf{x}} \psi_{k_{\parallel}}^{\lambda}(\mathbf{x})$, where

$$(8.12) \quad \begin{aligned} \psi_{k_{\parallel}}^{\lambda}(\mathbf{x}) &= \sum_{I=A,B} \sum_{n \geq 0} \alpha_n^I(\lambda) p_{k_{\parallel}, I}^{\lambda}[n](\mathbf{x}) + \tilde{\psi}[\vec{\alpha}(\lambda)](\mathbf{x}), \\ E^{\lambda}(k_{\parallel}) &= E_0^{\lambda} + \rho_{\lambda} \tilde{\Omega}(\lambda; k_{\parallel}), \end{aligned}$$

and the map $\vec{\alpha} \mapsto \tilde{\psi}[\vec{\alpha}](\mathbf{x})$ is given in (6.9). We shall succeed in this construction for $k_{\parallel} \in \mathcal{J} \subset \subset (2\pi/3, 4\pi/3)$ and $\lambda > \lambda_{\star}(\mathcal{J})$ sufficiently large.

The first step in this construction is to note that as λ tends to infinity the system (8.7)-(8.8) formally reduces to the edge state eigenvalue problem for the tight-binding Hamiltonian, $H_{\#}^{\text{TB}}$ (see (2.1), (2.4)) given by:

$$(8.13) \quad \begin{aligned} \alpha_{m-1}^B + (1 + e^{-ik_{\parallel}}) \alpha_m^B - \tilde{\Omega} \alpha_m^A &= 0, \quad m \geq 0 \\ (1 + e^{ik_{\parallel}}) \alpha_m^A + \alpha_{m+1}^A - \tilde{\Omega} \alpha_m^B &= 0, \quad m \geq 0, \quad \text{with } \alpha_{-1}^B = 0. \end{aligned}$$

By Theorem 2.2, if $k_{\parallel} \in (2\pi/3, 4\pi/3)$ the system (8.13) has an isolated and simple eigenvalue at $\tilde{\Omega}^{\text{TB}} = 0$ with corresponding vector $\vec{\alpha}^{\text{TB}} = \{\alpha_m^{\text{TB}}\}_{m \geq 0} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ given by:

$$(8.14) \quad \alpha_m^{\text{TB}} = \begin{pmatrix} \alpha_{\text{TB}, A} \\ \alpha_{\text{TB}, B} \end{pmatrix}_m = \gamma_{\star} \begin{pmatrix} (-1)^m (1 + e^{ik_{\parallel}})^m \\ 0 \end{pmatrix}, \quad \text{for } m \geq 0,$$

where we take $\gamma_{\star} = \sqrt{1 - |\zeta(k_{\parallel})|^2} \neq 0$ so that $\vec{\alpha}^{\text{TB}}$ has $l^2(\mathbb{N}_0; \mathbb{C}^2)$ - norm equal to one.

² Actually, the operator which emerges in (8.7)-(8.8) is $-H_{\#}^{\text{TB}}(k_{\parallel})$, minus one times the operator studied in Section 2. However, since $\sigma_2 H_{\#}^{\text{TB}}(k_{\parallel}) \sigma_2 = -H_{\#}^{\text{TB}}(k_{\parallel})$, the spectrum of $H_{\#}^{\text{TB}}(k_{\parallel})$ is symmetric about zero energy and $-H_{\#}^{\text{TB}}(k_{\parallel}) - z\text{Id}$ has the same invertibility properties of $H_{\#}^{\text{TB}}(k_{\parallel}) - z\text{Id}$. Hence, in this and the following section we take $H_{\#}^{\text{TB}}(k_{\parallel})$ to denote the negative of the operator studied in Section 2.

To prove that (8.9) has a solution in $l^2(\mathbb{N}_0, \mathbb{C}^2)$ which for λ large is approximately equal to $\vec{\alpha}^{\text{TB}}$, we seek a solution of (8.9) of the form:

$$(8.15) \quad \begin{aligned} \vec{\alpha}(\lambda) &= \vec{\alpha}^{\text{TB}} + \vec{\beta}(\lambda) = \begin{pmatrix} \alpha^{\text{TB,A}} \\ \alpha^{\text{TB,B}} \end{pmatrix} + \begin{pmatrix} \beta^A(\lambda) \\ \beta^B(\lambda) \end{pmatrix}, \\ \tilde{\Omega} &= \tilde{\Omega}(\lambda), \quad \text{where we take } \langle \vec{\alpha}^{\text{TB}}, \vec{\beta} \rangle_{l^2(\mathbb{N}_0; \mathbb{C}^2)} = 0. \end{aligned}$$

Introduce the orthogonal projection $\Pi_0^{\text{TB}} : l^2(\mathbb{N}_0; \mathbb{C}^2) \rightarrow \left(\text{span}\{\vec{\alpha}^{\text{TB}}\} \right)^\perp$. Substituting (8.15) into (8.9) and projecting onto $\text{span}\{\vec{\alpha}^{\text{TB}}\}$ and its orthogonal complement, we obtain the equivalent system for $\vec{\beta}$ and $\tilde{\Omega}$:

$$(8.16) \quad \left(H_\#^{\text{TB}}(k_\parallel) - \tilde{\Omega} \right) \vec{\beta} = \Pi_0^{\text{TB}} \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \vec{\alpha}^{\text{TB}} + \Pi_0^{\text{TB}} \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \vec{\beta},$$

$$(8.17) \quad \tilde{\Omega} + \left\langle \vec{\alpha}^{\text{TB}}, \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \vec{\alpha}^{\text{TB}} \right\rangle + \left\langle \vec{\alpha}^{\text{TB}}, \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \vec{\beta} \right\rangle = 0.$$

Let $\mathcal{R}^{\text{TB}}(\tilde{\Omega}; k_\parallel)$ denote the inverse of $\Pi_0^{\text{TB}} \left(H_\#^{\text{TB}}(k_\parallel) - \tilde{\Omega} \right) \Pi_0^{\text{TB}}$, which for $|\tilde{\Omega}| < c'$ is well-defined as a bounded operator on the $l^2(\mathbb{N}_0; \mathbb{C}^2)$ -orthogonal complement of $\text{span}\{\vec{\alpha}^{\text{TB}}(k_\parallel)\}$. Moreover, $\|\mathcal{R}^{\text{TB}}(\tilde{\Omega}; k_\parallel)\| \lesssim 1$ for $|\tilde{\Omega}| < c' < \delta(k_\parallel)$, by Theorem 2.2. For λ sufficiently large we may solve (8.16) for $\vec{\beta}[\tilde{\Omega}; \lambda] \in \text{Range } \Pi_0^{\text{TB}}$ and obtain:

$$(8.18) \quad \begin{aligned} \vec{\beta}[\tilde{\Omega}; \lambda] &= \left[I - \mathcal{R}^{\text{TB}}(\tilde{\Omega}; k_\parallel) \Pi_0^{\text{TB}} \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \right]^{-1} \Pi_0^{\text{TB}} \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \vec{\alpha}^{\text{TB}} \\ &\equiv \mathcal{A}(\tilde{\Omega}; \lambda) \Pi_0^{\text{TB}} \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \vec{\alpha}^{\text{TB}}. \end{aligned}$$

This follows by the bound $\|\mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega})\|_{l^2 \rightarrow l^2} \lesssim e^{-c\lambda}$; see (8.10). Therefore, the construction of $\vec{\beta}(\lambda)$, $\tilde{\Omega}(\lambda)$ (see (8.11)) boils down to solving the following scalar nonlinear equation for $\tilde{\Omega}$ as a function of λ :

$$(8.19) \quad \tilde{\Omega} + \left\langle \vec{\alpha}^{\text{TB}}, \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \vec{\alpha}^{\text{TB}} \right\rangle + \left\langle \vec{\alpha}^{\text{TB}}, \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \mathcal{A}(\tilde{\Omega}; \lambda) \Pi_0^{\text{TB}} \mathcal{P}(\lambda; \rho_\lambda \tilde{\Omega}) \vec{\alpha}^{\text{TB}} \right\rangle = 0.$$

Using analyticity in $\tilde{\Omega}$ and previous bounds, we may write (8.19) as

$$(8.20) \quad \tilde{\Omega} + \left\langle \vec{\alpha}^{\text{TB}}, \mathcal{P}(\lambda; 0) \vec{\alpha}^{\text{TB}} \right\rangle + \mathcal{G}(\tilde{\Omega}; \lambda) = 0.$$

Here, $\mathcal{G}(\tilde{\Omega}; \lambda)$ is analytic with $|\partial_{\tilde{\Omega}}^j \mathcal{G}(\tilde{\Omega}; \lambda)| \lesssim e^{-c\lambda}$ ($j = 1, 2$) for all $\tilde{\Omega}$ in the complex neighborhood of zero, $|\tilde{\Omega}| < c'$. Since $\left| \left\langle \vec{\alpha}^{\text{TB}}, \mathcal{P}(\lambda; 0) \vec{\alpha}^{\text{TB}} \right\rangle \right| \leq e^{-c\lambda}$, for λ sufficiently large, equation (8.20) may be solved for $\tilde{\Omega}(\lambda)$ by using a contraction mapping argument on the disc: $|\tilde{\Omega}| \leq 2Ce^{-c\lambda}$. Therefore, modulo Propositions 7.1 and 7.2 which are proved in Sections 10, 11 and 12, we have proved our main result, Theorem 1.2.

9. RESOLVENT CONVERGENCE; PROOF OF THEOREM 1.1

We study the scaled resolvent:

$$\left(\rho_\lambda^{-1} H_\#^\lambda - z\text{Id} \right)^{-1} = \left(\rho_\lambda^{-1} (-\Delta + V_\# - E_0^\lambda) - z\text{Id} \right)^{-1}$$

as an operator on $L^2(\mathbb{R}^2)$. We consider the scaled non-homogeneous equation

$$(9.1) \quad (\rho_\lambda^{-1} H_\#^\lambda(k_\parallel) - z \text{Id}) \psi = \varphi, \quad \varphi \in L^2(\Sigma).$$

or equivalently

$$(9.2) \quad (H_\#^\lambda(k_\parallel) - \rho_\lambda z \text{Id}) \psi = \rho_\lambda \varphi, \quad \varphi \in L^2(\Sigma).$$

We express φ as:

$$(9.3) \quad \varphi = \sum_{J=A,B} \sum_{n \geq 0} \beta_n^J p_{k_\parallel, J}^\lambda[n] + \tilde{\varphi}, \quad \Pi_{AB}(k_\parallel) \tilde{\varphi} = \tilde{\varphi}$$

and seek a solution of (9.1) in the form

$$(9.4) \quad \psi = \sum_{I=A,B} \sum_{n \geq 0} \alpha_n^I p_{k_\parallel, I}^\lambda[n] + \tilde{\psi}, \quad \Pi_{AB}(k_\parallel) \tilde{\psi} = \tilde{\psi}$$

where $\alpha = \{(\alpha_n^A, \alpha_n^B)^\top\}_{n \geq 0} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ and $\tilde{\psi} = \Pi_{AB}(k_\parallel) \tilde{\psi} \in \mathcal{X}_{AB}^\lambda(k_\parallel)$.

Substitution of (9.3) and (9.4) into (9.2) and projecting the resulting equation with $\Pi_{AB}(k_\parallel)$ and $I - \Pi_{AB}(k_\parallel)$ (whose range is $\text{span}\{p_{k_\parallel, I}^\lambda[n] : I = A, B, n \geq 0\}$), yields the coupled system for $\alpha = \{\alpha_n^I : n \geq 0, I = A, B\} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ and $\tilde{\psi} \in \mathcal{X}_{AB}^\lambda(k_\parallel)$:

$$(9.5) \quad \Pi_{AB}(k_\parallel) (H_\#^\lambda(k_\parallel) - \rho_\lambda z \text{Id}) \tilde{\psi} = - \sum_{I,n} \alpha_n^I \Pi_{AB}(k_\parallel) H_\#^\lambda(k_\parallel) p_{k_\parallel, I}^\lambda[n] + \rho_\lambda \tilde{\varphi}$$

$$(9.6) \quad \sum_{I,n} \left\langle p_{k_\parallel, J}^\lambda[m], (H_\#^\lambda(k_\parallel) - \rho_\lambda z \text{Id}) p_{k_\parallel, I}^\lambda[n] \right\rangle \alpha_n^I + \left\langle H_\#^\lambda(k_\parallel) p_{k_\parallel, J}^\lambda[m], \tilde{\psi} \right\rangle \\ = \rho_\lambda \sum_{I,n} \left\langle p_{k_\parallel, J}^\lambda[m], p_{k_\parallel, I}^\lambda[n] \right\rangle \beta_n^I, \text{ for } J = A, B \text{ and } m \geq 0,$$

where the sums $\sum_{I,n}$ are over $I = A, B$ and $n \geq 0$.

We next use Proposition 5.1 to solve (9.5) for $\tilde{\psi} \in L^2(\Sigma)$ and obtain:

$$(9.7) \quad \tilde{\psi} = - \sum_{I,n} \alpha_n^I \mathcal{K}_\#^\lambda(\rho_\lambda z, k_\parallel) H_\#^\lambda(k_\parallel) p_{k_\parallel, I}^\lambda[n] + \rho_\lambda \mathcal{K}_\#^\lambda(\rho_\lambda z, k_\parallel) \tilde{\varphi}.$$

Substitution of the expression in (9.7) for $\tilde{\psi}$ into the left hand side of (9.6) yields the closed non-homogeneous system for $\alpha \in l^2(\mathbb{N}_0; \mathbb{C}^2)$:

$$(9.8) \quad \sum_{I,n} \mathcal{M}_{JI}^\lambda[m, n] \alpha_n^I = \rho_\lambda \left[\sum_{I,n} \left\langle p_{k_\parallel, J}^\lambda[m], p_{k_\parallel, I}^\lambda[n] \right\rangle \beta_n^I - \left\langle H_\#^\lambda(k_\parallel) p_{k_\parallel, J}^\lambda[m], \mathcal{K}_\#^\lambda(\rho_\lambda z, k_\parallel) \tilde{\varphi} \right\rangle \right],$$

for each $J = A, B$ and $m \geq 0$. The matrix elements $\mathcal{M}_{JI}^\lambda[m, n]$ are displayed in (6.11). As in our study of the edge state eigenvalue problem (Section 8) we expand the $\mathcal{M}_{JI}^\lambda[m, n]$ using

Proposition 7.1 and obtain the following system, which is equivalent to (9.8)³ :

$$(9.9) \quad \begin{aligned} & \left[\left(H_{\#}^{\text{TB}}(k_{\parallel}) - z \text{Id} - \mathcal{P}(\lambda; \rho_{\lambda} z) \right) \begin{pmatrix} \alpha^A \\ \alpha^B \end{pmatrix} \right]_m \\ &= - \begin{pmatrix} \sum_{I,n} \langle p_{k_{\parallel},A}^{\lambda}[m], p_{k_{\parallel},I}^{\lambda}[n] \beta_n^I \rangle \\ \sum_{I,n} \langle p_{k_{\parallel},B}^{\lambda}[m], p_{k_{\parallel},I}^{\lambda}[n] \beta_n^I \rangle \end{pmatrix} + \begin{pmatrix} \langle H_{\#}^{\lambda}(k_{\parallel}) p_{k_{\parallel},A}^{\lambda}[m], \mathcal{K}_{\#}^{\lambda}(\rho_{\lambda} z, k_{\parallel}) \tilde{\varphi} \rangle \\ \langle H_{\#}^{\lambda}(k_{\parallel}) p_{k_{\parallel},B}^{\lambda}[m], \mathcal{K}_{\#}^{\lambda}(\rho_{\lambda} z, k_{\parallel}) \tilde{\varphi} \rangle \end{pmatrix}, \quad m \geq 0. \end{aligned}$$

Recalling the bound $\|\mathcal{P}(\lambda; \rho_{\lambda} \tilde{\Omega})\|_{l^2 \rightarrow l^2} \lesssim e^{-c\lambda}$ (see (8.10)), Proposition 4.2 and Proposition 4.3 we solve for α^{λ} and find

$$(9.10) \quad \begin{aligned} \alpha^{\lambda} &= \left(H_{\#}^{\text{TB}}(k_{\parallel}) - z \text{Id} \right)^{-1} \beta + \alpha_1^{\lambda}, \quad \text{where} \\ \|\alpha_1^{\lambda}\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)} &\lesssim e^{-c\lambda} \left(\|\beta\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)} + \|\Pi_{AB}(k_{\parallel})\varphi\|_{L^2(\Sigma)} \right) \end{aligned}$$

We therefore have that $\psi = \left(\rho_{\lambda}^{-1} H_{\#}^{\lambda}(k_{\parallel}) - z \text{Id} \right)^{-1} \varphi \in L^2(\Sigma)$ is given by:

$$(9.11) \quad \begin{aligned} \left(\rho_{\lambda}^{-1} H_{\#}^{\lambda}(k_{\parallel}) - z \text{Id} \right)^{-1} \varphi &= \sum_{I,n} \left[\left(H_{\#}^{\text{TB}}(k_{\parallel}) - z \text{Id} \right)^{-1} \beta + \alpha_1^{\lambda} \right] p_{k_{\parallel},I}^{\lambda}[n] \\ &+ \mathcal{O}_{L^2(\Sigma)} \left(e^{-c\lambda} \|\beta\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)} + e^{-c\lambda} \|\Pi_{AB}(k_{\parallel})\varphi\|_{L^2(\Sigma)} \right) \end{aligned}$$

Introduce $H_{\#,k_{\parallel}}^{\lambda}$, the restriction of $H_{\#}^{\lambda}$, to the space $H_{k_{\parallel}}^2$. Since $H_{\#}^{\lambda}$ commutes with $\mathbf{x} \mapsto \mathbf{x} + \mathbf{v}_2$ it follows that $H_{\#,k_{\parallel}}^{\lambda}$ maps the space $H_{k_{\parallel}}^2$ into $L_{k_{\parallel}}^2$. Let $\mathcal{P}_{AB,k_{\parallel}}$ denote the projection of $L_{k_{\parallel}}^2$ onto the orthogonal complement of the subspace of $L_{k_{\parallel}}^2$ spanned by the states: $P_{k_{\parallel},I}^{\lambda}[n] = e^{i\frac{k_{\parallel}}{2\pi}(\mathbf{x} - \mathbf{v}_I)} p_{k_{\parallel},I}^{\lambda}[n] \in L_{k_{\parallel}}^2$, where $I = A, B$ and $n \geq 0$; see (4.4). Therefore, for any $F \in L_{k_{\parallel}}^2$:

$$(9.12) \quad \begin{aligned} \left(\rho_{\lambda}^{-1} H_{\#,k_{\parallel}}^{\lambda} - z \text{Id} \right)^{-1} F &= \sum_{I,n} \left[\left(H_{\#}^{\text{TB}}(k_{\parallel}) - z \text{Id} \right)^{-1} \beta + \alpha_1^{\lambda} \right] P_{k_{\parallel},I}^{\lambda}[n] \\ &+ \mathcal{O}_{L^2(\Sigma)} \left(e^{-c\lambda} \|\beta\|_{l^2(\mathbb{N}_0; \mathbb{C}^2)} + e^{-c\lambda} \|\mathcal{P}_{AB,k_{\parallel}} F\|_{L_{k_{\parallel}}^2} \right). \end{aligned}$$

Any $F \in L_{k_{\parallel}}^2$ has the representation $F = \sum_{I,n} \alpha_n^I[F] P_{k_{\parallel},I}^{\lambda}[n] + F_{\perp}$, where $\{\alpha_n^I[F]\}_{I,n} \in l^2(\mathbb{N}_0; \mathbb{C}^2)$ and $F_{\perp} \in \text{Range}(\mathcal{P}_{AB,k_{\parallel}})$. Define the map $J_{k_{\parallel}} : L_{k_{\parallel}}^2 \rightarrow l^2(\mathbb{N}_0; \mathbb{C}^2) \oplus \text{Range}(\mathcal{P}_{AB,k_{\parallel}})$ by:

$$(9.13) \quad J_{k_{\parallel}} : F \mapsto \begin{pmatrix} \{\alpha_n^I[F]\} \\ F_{\perp} \end{pmatrix} = \begin{pmatrix} \left\{ \langle P_{k_{\parallel},I}^{\lambda}[n], F \rangle + \mathcal{O}(e^{-c\lambda} \|F\|_{L_{k_{\parallel}}^2}) \right\} \\ F_{\perp} \end{pmatrix}$$

³As in Section 8 (see the footnote after (8.9)), based on the observation $\sigma_2 H_{\#}^{\text{TB}}(k_{\parallel}) \sigma_2 = -H_{\#}^{\text{TB}}(k_{\parallel})$ we let $H_{\#}^{\text{TB}}(k_{\parallel})$ denote the negative of the operator studied in Section 2.

We therefore have from (9.12) that

$$\left(\rho_\lambda^{-1} H_{\sharp, k_\parallel}^\lambda - z \text{Id} \right)^{-1} - J_{k_\parallel}^* \begin{pmatrix} \left(H_{\sharp}^{\text{TB}}(k_\parallel) - z \text{Id} \right)^{-1} & 0 \\ 0 & 0 \end{pmatrix} J_{k_\parallel} = \mathcal{O}_{L_{k_\parallel}^2 \rightarrow L_{k_\parallel}^2}(e^{-c\lambda}).$$

This completes the proof of Theorem 1.1.

10. THE RESOLVENT KERNEL AND WEIGHTED RESOLVENT BOUNDS

It remains for us to prove Propositions 7.1 and 7.2 on the expansion and estimation of matrix elements. The proof of Proposition 7.1 concerning the linear matrix elements uses the energy estimates on the resolvent obtained in Section 5.

To prove Proposition 7.2 we require exponentially weighted estimates, which we obtain by constructing the resolvent kernel and obtaining pointwise bounds on it. We carry this out in the present section. In Section 11 we then give the proof of Proposition 7.1 and in Section 12 we prove Proposition 7.2.

In Section 5 we obtained energy estimates for $\mathcal{K}_\sharp^\lambda(\Omega, k_\parallel)$, the inverse of

$$\Pi_{AB}(k_\parallel) \left(H_\sharp^\lambda(k_\parallel) - \Omega \right) \Pi_{AB}(k_\parallel) = \Pi_{AB} \left[- \left(\nabla_{\mathbf{x}} + i \frac{k_\parallel}{2\pi} \mathfrak{R}_2 \right)^2 + \lambda^2 V_\sharp(\mathbf{x}) - E_0^\lambda - \Omega \right] \Pi_{AB},$$

defined as a bounded operator from $\mathcal{X}_{AB}(k_\parallel)$ to $\mathcal{X}_{AB}(k_\parallel) \cap H^2(\Sigma)$; see Proposition 5.1, which holds for all $|\Omega| < c'$, where c' is a sufficiently small positive constant. We may extend $\mathcal{K}_\sharp^\lambda(\Omega, k_\parallel)$ to an operator acting on all of $L^2(\Sigma)$, not just $\mathcal{X}_{AB}(k_\parallel)$, by composing it with $\Pi_{AB}(k_\parallel)$, *i.e.* we require $\mathcal{K}_\sharp^\lambda(\Omega, k_\parallel)\psi = 0$ if $\Pi_{AB}(k_\parallel)\psi = 0$.

In this section we shall prove, under the more stringent restriction on Ω : $|\Omega| \leq e^{-c\lambda}$ for some $c > 0$ and $\lambda \gg 1$, that this operator derives from a kernel $\mathcal{K}_\sharp^\lambda(\mathbf{x}, \mathbf{y}; \Omega, k_\parallel)$. Specifically, we have

Theorem 10.1. *There exist constants $\lambda_\star, c > 0$ such that for $\lambda \geq \lambda_\star$, $|\Omega| \leq e^{-c\lambda}$ and for each $k_\parallel \in [0, 2\pi]$ the following holds for the operator $\mathcal{K}_\sharp^\lambda(\Omega, k_\parallel)$, which is bounded on $L^2(\Sigma)$:*

(1) $\mathcal{K}_\sharp^\lambda(\Omega, k_\parallel)$ is arises from an integral kernel $\mathcal{K}_\sharp^\lambda(\mathbf{x}, \mathbf{y}; \Omega, k_\parallel)$:

$$(10.1) \quad \mathcal{K}_\sharp^\lambda(\Omega, k_\parallel)[f](\mathbf{x}) = \Pi_{AB}^\lambda \mathcal{K}_\sharp^\lambda(\Omega, k_\parallel) \Pi_{AB}^\lambda[f](\mathbf{x}) = \int_{\Omega_\Sigma} \mathcal{K}_\sharp^\lambda(\mathbf{x}, \mathbf{y}; \Omega, k_\parallel) f(\mathbf{y}) d\mathbf{y} .$$

(2) *The integral kernel $\mathcal{K}_\sharp^\lambda(\mathbf{x}, \mathbf{y}; \Omega, k_\parallel)$ satisfies the following bound: there exist positive constants R, C_1, C_2 , independent of k_\parallel and Ω , such that for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$:*

$$(10.2) \quad \left| \mathcal{K}_\sharp^\lambda(\mathbf{x}, \mathbf{y}; \Omega, k_\parallel) \right| \leq C_1 \left[\lambda^4 + \left| \log |\mathbf{x} - \mathbf{y}| \right| \right] \mathbf{1}_{|\mathbf{x} - \mathbf{y}| \leq R} + C_2 e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|} .$$

Theorem 10.1 is at the heart of the proof of Proposition 7.2, which provides bounds on the nonlinear matrix elements of $\mathcal{M}^\lambda(\Omega, k_\parallel)$. The remainder of this section is devoted to the proof of Theorem 10.1. The construction and estimation $\mathcal{K}_\sharp^\lambda$ is based on a strategy, in which we piece together localized atomic Green's functions with appropriate corrections.

10.1. The free Green's function and bounds on the atomic ground state. Denote by $G_\lambda^{\text{free}}(\mathbf{x})$ the fundamental solution of $-\Delta - E_0^\lambda$:

$$(10.3) \quad (-\Delta_{\mathbf{x}} - E_0^\lambda) G_\lambda^{\text{free}}(\mathbf{x}) = \delta(\mathbf{x}),$$

where $\delta(\mathbf{x})$ is the Dirac delta function. Here, E_0^λ denotes the ground state of $H_{\text{atom}}^\lambda = -\Delta + \lambda^2 V_0$; see hypothesis (GS), (3.4). Note that $G_\lambda^{\text{free}}(\mathbf{x}) = G^{\text{free}}\left(\sqrt{|E_0^\lambda|} \mathbf{x}\right)$, where $G^{\text{free}}(\mathbf{x})$ satisfies $(-\Delta_{\mathbf{x}} + 1) G^{\text{free}}(\mathbf{x}) = \delta(\mathbf{x})$, $\mathbf{x} \in \mathbb{R}^2$. $G^{\text{free}}(\mathbf{x}) = K_0(|\mathbf{x}|)$ is the modified Bessel function of order zero, which decays to zero exponentially as $|\mathbf{x}| \rightarrow \infty$ [68]. The following lemma summarizes important standard properties of $G_\lambda^{\text{free}}(\mathbf{x})$; see [26, 59]

Lemma 10.2. *For $\mathbf{x} \in \mathbb{R}^2$,*

- (1) $G^{\text{free}}(\mathbf{x}) = G^{\text{free}}(|\mathbf{x}|)$ is positive and strictly decreasing for $|\mathbf{x}| \geq 0$.
- (2) There exist entire functions f and g and constants C_1, c_2 , such that

$$(10.4) \quad G^{\text{free}}(\mathbf{x}) = f(|\mathbf{x}|) \log |\mathbf{x}| + g(|\mathbf{x}|),$$

where $f(0) = -1/2\pi$ and $|\partial_s^j f(s)|, |\partial_s^j g(s)| \leq C_1 e^{-c_2 s}$, for $j = 0, 1$ and all $s \in [0, \infty)$.

- (3) $G^{\text{free}}(\mathbf{x}) \lesssim |\mathbf{x}|^{-\frac{1}{2}} e^{-|\mathbf{x}|}$ for $|\mathbf{x}|$ large.

The bounds on $f(s)$ and $g(s)$ are proved, for the case $j = 0$, in [59]. This proof can be extended to a derivation of the bounds for $j = 1$. Alternatively, these bounds may be deduced directly from the integral representation for $G^{\text{free}}(\mathbf{x})$ used in the proof of Lemma 15.3 of [26].

We shall apply the following consequence of Lemma 10.2 and (3.4):

There exist $c, c' > 0$, and for each $R > 0$, additional constants $C_R, C'_R > 0$, such that

$$(10.5) \quad 0 < G_\lambda^{\text{free}}(\mathbf{x}) = G^{\text{free}}\left(\sqrt{|E_0^\lambda|} \mathbf{x}\right) \leq C_R e^{-c\lambda|\mathbf{x}|} \left(\left| \log(\lambda|\mathbf{x}|) \right| \mathbf{1}_{\{\lambda|\mathbf{x}| \leq R\}} + 1 \right), \quad \mathbf{x} \in \mathbb{R}^2.$$

$$(10.6) \quad |\nabla_{\mathbf{x}} G_\lambda^{\text{free}}(\mathbf{x})| \leq C'_R e^{-c'\lambda|\mathbf{x}|} \left(\frac{1}{\lambda|\mathbf{x}|} \mathbf{1}_{\{\lambda|\mathbf{x}| \leq R\}} + 1 \right)$$

10.2. The atomic Green's function.

In this section we establish bounds (integral and then pointwise) on the Green's function associated with $H_{\text{atom}}^\lambda - E_0^\lambda = -\Delta + \lambda^2 V_0(\mathbf{x}) - E_0^\lambda$. Since H_{atom}^λ has a one dimensional kernel spanned by $p_0^\lambda(\mathbf{x})$, and a spectral gap (see (3.6)), the operator $H_{\text{atom}}^\lambda - E_0^\lambda$ is invertible on the orthogonal complement of $\text{span}\{p_0^\lambda\}$.

We denote by $G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})$ the associated Green's kernel, which solves

$$(10.7) \quad (-\Delta_{\mathbf{x}} + \lambda^2 V_0(\mathbf{x}) - E_0^\lambda) G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) = \delta(\mathbf{x} - \mathbf{y}) - p_0^\lambda(\mathbf{x}) p_0^\lambda(\mathbf{y})$$

and which satisfies

$$(10.8) \quad \int_{\mathbb{R}^2} G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) p_0^\lambda(\mathbf{y}) d\mathbf{y} = 0, \quad \text{for all } \mathbf{x} \in \mathbb{R}^2,$$

$$(10.9) \quad G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) = G_\lambda^{\text{atom}}(\mathbf{y}, \mathbf{x}) \quad \text{for } \mathbf{x}, \mathbf{y} \in \mathbb{R}^2 \text{ with } \mathbf{x} \neq \mathbf{y}.$$

For fixed \mathbf{x} , the function $\mathbf{y} \mapsto G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})$ belongs to $L^2(\mathbb{R}_y^2)$, and we have for any $f \in L^2(\mathbb{R}^2)$ that the function

$$(10.10) \quad u(\mathbf{x}) = \int_{\mathbb{R}^2} G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y}$$

solves

$$(10.11) \quad (-\Delta + \lambda^2 V_0(\mathbf{x}) - E_0^\lambda) u(\mathbf{x}) = f(\mathbf{x}) - \langle p_0^\lambda, f \rangle_{L^2(\mathbb{R})} p_0^\lambda(\mathbf{x}),$$

$$(10.12) \quad \langle p_0^\lambda, u \rangle_{L^2(\mathbb{R}^2)} = 0.$$

10.2.1. L^2 bounds on $\mathbf{x} \mapsto G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})$ and $\mathbf{y} \mapsto G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})$.

By the spectral gap hypothesis on H_{atom}^λ , (3.6), we have that u satisfies the bound:

$$(10.13) \quad \|u\|_{L^2(\mathbb{R}^2)} \leq C \|f\|_{L^2(\mathbb{R}^2)}.$$

We may next obtain pointwise bounds on $u(\mathbf{x})$ in terms of $\|f\|_{L^2(\mathbb{R}^2)}$. In particular, we claim that

$$(10.14) \quad |u(\mathbf{x})| \leq C \lambda^2 \|f\|_{L^2(\mathbb{R}^2)}.$$

We prove this as follows:

$$\begin{aligned} |u(\mathbf{x})| &\leq C \left(\|\Delta u\|_{L^2(B_1(\mathbf{x}))} + \|u\|_{L^2(B_1(\mathbf{x}))} \right) \\ &\leq C \left(\left\| (E_0^\lambda - \lambda^2 V_0)u + f - \langle p_0^\lambda, f \rangle p_0^\lambda \right\|_{L^2(B_1(\mathbf{x}))} + \|u\|_{L^2(B_1(\mathbf{x}))} \right) \\ &\leq C \lambda^2 \|f\|_{L^2(\mathbb{R}^2)} \end{aligned}$$

which implies the bound (10.14).

Therefore, by (10.10), for all $f \in L^2(\mathbb{R}^2)$:

$$(10.15) \quad \left| \int_{\mathbb{R}^2} G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y} \right| \leq C \lambda^2 \|f\|_{L^2(\mathbb{R}^2)}.$$

Consequently,

$$(10.16) \quad \left(\int_{\mathbb{R}^2} |G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})|^2 d\mathbf{y} \right)^{\frac{1}{2}} \leq C \lambda^2, \quad \mathbf{x} \in \mathbb{R}^2$$

and by symmetry of G_λ^{atom}

$$(10.17) \quad \left(\int_{\mathbb{R}^2} |G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})|^2 d\mathbf{x} \right)^{\frac{1}{2}} \leq C \lambda^2, \quad \mathbf{y} \in \mathbb{R}^2.$$

We now use these L^2 bounds on $G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})$ to obtain pointwise bounds.

10.2.2. *Pointwise bounds on $G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})$.* Recall that $\text{supp} V_0 \subset B_{r_0}(0)$.

Theorem 10.3 (Pointwise bounds on $G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})$). *(1) For all $R > 0$, there exist $\lambda_0 = \lambda_0(R)$ and positive constants c, C_R and D_R such that for all $\lambda > \lambda_0$:*

$$(10.18) \quad \left| G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) - c_0 \log |\mathbf{x} - \mathbf{y}| \right| \leq C_R \lambda^4 \quad \text{for } |\mathbf{x} - \mathbf{y}| \leq R,$$

where $c_0 = -(2\pi)^{-1}$.

(2) There exist $R > 10r_0$ and positive constants λ' , C and c , which depend on R but not on λ , such that for all $\lambda > \lambda'(R)$:

$$(10.19) \quad |G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})| \leq C e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{y}|}, \quad |\mathbf{x} - \mathbf{y}| \geq R.$$

(3) Choose r_j , $j = 1, 2, 3$, such that $r_0 < r_1 < r_2 < r_3 < \frac{1}{10}R$. Assume $\mathbf{y} \in B_{r_1}(0)$ and $\mathbf{x} \notin B_{r_3}(0)$. Then,

$$(10.20) \quad \left| G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) \right| \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{y}|},$$

where the implied constants depend on r_0, r_1, r_2 and r_3 .

Proof of bound (10.18):

Fix $\mathbf{y} \in \mathbb{R}^2$. By (10.7) we have

$$(10.21) \quad \begin{aligned} -\Delta_{\mathbf{x}} G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) &= \delta(\mathbf{x} - \mathbf{y}) - p_0^\lambda(\mathbf{x}) p_0^\lambda(\mathbf{y}) + (E_0^\lambda - \lambda^2 V_0(\mathbf{x})) G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) \\ &= -\Delta_{\mathbf{x}} c_0 \log |\mathbf{x} - \mathbf{y}| - p_0^\lambda(\mathbf{x}) p_0^\lambda(\mathbf{y}) + (E_0^\lambda - \lambda^2 V_0(\mathbf{x})) G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}). \end{aligned}$$

Hence,

$$(10.22) \quad -\Delta_{\mathbf{x}} \left[G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) - c_0 \log |\mathbf{x} - \mathbf{y}| \right] = -p_0^\lambda(\mathbf{x}) p_0^\lambda(\mathbf{y}) + (E_0^\lambda - \lambda^2 V_0(\mathbf{x})) G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}).$$

Therefore, using that $|f(\mathbf{x})| \lesssim \|\Delta f(\mathbf{z})\|_{L^2(B_1(\mathbf{x}); d\mathbf{z})} + \|f(\mathbf{z})\|_{L^2(B_1(\mathbf{x}); d\mathbf{z})}$ we have for arbitrary fixed $\mathbf{y} \in \mathbb{R}^2$ and all $\mathbf{x} \in \mathbb{R}^2$ satisfying $|\mathbf{x} - \mathbf{y}| \leq R$:

$$(10.23) \quad \begin{aligned} &\left| G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) - c_0 \log |\mathbf{x} - \mathbf{y}| \right| \\ &\leq \left\| -p_0^\lambda(\mathbf{z}) p_0^\lambda(\mathbf{y}) + (E_0^\lambda - \lambda^2 V_0(\mathbf{z})) G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y}) \right\|_{L^2(B_1(\mathbf{x}); d\mathbf{z})} \\ &\quad + \left\| G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y}) - c_0 \log |\mathbf{z} - \mathbf{y}| \right\|_{L^2(B_1(\mathbf{x}); d\mathbf{z})}. \end{aligned}$$

To continue this bound, we use that

$$(10.24) \quad \begin{aligned} |p_0^\lambda(\mathbf{y})| &\lesssim \lambda \text{ (see (3.5))}, \quad \|p_0^\lambda\|_{L^2} = 1, \quad |E_0^\lambda - \lambda^2 V_0(\mathbf{z})| \lesssim \lambda^2, \\ \|G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y})\|_{L^2(B_1(\mathbf{x}); d\mathbf{z})} &\lesssim \lambda^2 \text{ and } \|\log |\mathbf{z} - \mathbf{y}|\|_{L^2(B_1(\mathbf{x}); d\mathbf{z})} \leq C'_R. \end{aligned}$$

The bounds (10.24) follow since $|E_0^\lambda| \lesssim \lambda^2$ (since $\|V_0\|_\infty < \infty$) and by (3.5) and (10.17). We obtain for any $R > 0$ that there exists $C_R < \infty$ such that

$$(10.25) \quad \left| G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) - c_0 \log |\mathbf{x} - \mathbf{y}| \right| \leq C_R \lambda^4, \quad \text{for all } |\mathbf{x} - \mathbf{y}| \leq R, \text{ with } \mathbf{x} \neq \mathbf{y}.$$

Proof of bound (10.19): Recall that the support of V_0 is contained in $B_{r_0}(0)$. Assume $|\mathbf{x} - \mathbf{y}| > R$, and choose constants:

$$(10.26) \quad r_0 < r_1 < r_2 < r_3 < \frac{1}{10}R.$$

Thus, we require $R > 10r_0$. Without any loss of generality, we assume $|\mathbf{y}| \leq |\mathbf{x}|$. Therefore, $R < |\mathbf{x} - \mathbf{y}| \leq |\mathbf{x}| + |\mathbf{y}| \leq 2|\mathbf{x}|$ and therefore

$$(10.27) \quad |\mathbf{x}| \geq \frac{1}{2}|\mathbf{x} - \mathbf{y}| > \frac{1}{2}R > r_3.$$

Let $\Theta_{\text{out}} = \Theta_{\text{out}}(\mathbf{x})$ denote a smooth function of $r = |\mathbf{x}|$, defined for all $\mathbf{x} \in \mathbb{R}^2$, such that $0 \leq \Theta_{\text{out}}(\mathbf{x}) \leq 1$ and

$$(10.28) \quad \Theta_{\text{out}}(\mathbf{x}) \equiv \begin{cases} 1, & |\mathbf{x}| \geq r_2 \\ 0, & |\mathbf{x}| \leq r_1 \end{cases}$$

We note that $\Theta_{\text{out}} \cdot V_0 \equiv 0$.

Using the defining equation for G_λ^{atom} , (10.7), we obtain:

$$(10.29) \quad \begin{aligned} & (-\Delta_{\mathbf{z}} - E_0^\lambda) [\Theta_{\text{out}}(\mathbf{z}) G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y})] \\ &= \Theta_{\text{out}}(\mathbf{z}) \{ -p_0^\lambda(\mathbf{z}) p_0^\lambda(\mathbf{y}) \} + \Theta_{\text{out}}(\mathbf{z}) \cdot \delta(\mathbf{z} - \mathbf{y}) \\ &+ 2\nabla_{\mathbf{z}} \Theta_{\text{out}}(\mathbf{z}) \cdot \nabla_{\mathbf{z}} G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y}) + (\Delta_{\mathbf{z}} \Theta_{\text{out}}(\mathbf{z})) G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y}) . \end{aligned}$$

We next use the Green's function G_λ^{free} (see (10.3)) to represent $\Theta_{\text{out}}(\mathbf{x}) G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y})$. Multiplication of (10.29) by $G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z})$ and integration with respect to \mathbf{z} yields

$$\begin{aligned} \Theta_{\text{out}}(\mathbf{x}) G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) &= \int_{\mathbb{R}^2} G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z}) (-\Delta_{\mathbf{z}} - E_0^\lambda) [\Theta_{\text{out}}(\mathbf{z}) G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y})] d\mathbf{z} \\ &= \Theta_{\text{out}}(\mathbf{y}) G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) - \int_{\mathbb{R}^2} G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z}) \Theta_{\text{out}}(\mathbf{z}) p_0^\lambda(\mathbf{z}) d\mathbf{z} p_0^\lambda(\mathbf{y}) \\ &+ 2 \int_{\mathbb{R}^2} G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z}) \nabla_{\mathbf{z}} \Theta_{\text{out}}(\mathbf{z}) \cdot \nabla_{\mathbf{z}} G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y}) d\mathbf{z} \\ &+ \int_{\mathbb{R}^2} G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z}) (\Delta_{\mathbf{z}} \Theta_{\text{out}}(\mathbf{z})) G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y}) d\mathbf{z} , \end{aligned}$$

which, since $\Theta_{\text{out}}(\mathbf{x}) = 1$ for $|\mathbf{x}| > r_2$, we write as

$$(10.30) \quad G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) = \Theta_{\text{out}}(\mathbf{y}) G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) + \text{Term}_1(\mathbf{x}, \mathbf{y}) + \text{Term}_2(\mathbf{x}, \mathbf{y}) + \text{Term}_3(\mathbf{x}, \mathbf{y}) .$$

Since $|\mathbf{x} - \mathbf{y}| > R$, by (10.5) we have $|\Theta_{\text{out}}(\mathbf{y}) G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y})| \lesssim e^{-c\lambda|\mathbf{x}-\mathbf{y}|}$. We next estimate the latter three terms in (10.30) individually.

Bound on $\text{Term}_1(\mathbf{x}, \mathbf{y})$ of (10.30): Consider the integral

$$(10.31) \quad \text{Term}_1(\mathbf{x}, \mathbf{y}) \equiv - \int_{\mathbb{R}^2} G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z}) \Theta_{\text{out}}(\mathbf{z}) p_0^\lambda(\mathbf{z}) d\mathbf{z} p_0^\lambda(\mathbf{y}) .$$

Due to the factor of $\Theta_{\text{out}}(\mathbf{z})$ in the integrand of (10.31), only \mathbf{z} such that $|\mathbf{z}| \geq r_1$ are relevant. On this set we have $p_0^\lambda(\mathbf{z}) \lesssim e^{-c_1\lambda} e^{-c\lambda|\mathbf{z}|}$ by (3.5), for some constants $c_1, c > 0$. Furthermore, by (10.5), there exists $c' > 0$ such that $G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z}) \lesssim e^{-c'\lambda|\mathbf{x}-\mathbf{z}|} \left(\left| \log \lambda |\mathbf{x} - \mathbf{z}| \right| \mathbf{1}_{\{|\mathbf{x}-\mathbf{z}| \leq 1\}} + 1 \right)$.

Therefore, for some constant \tilde{c} (smaller than the minimum of c_1, c, c') we have

$$\begin{aligned}
\left| \text{Term}_1(\mathbf{x}, \mathbf{y}) \right| &\lesssim e^{-\tilde{c}\lambda} \int_{|\mathbf{z}| \geq r_1} e^{-\tilde{c}\lambda|\mathbf{x}-\mathbf{z}|} \left(\left| \log \lambda |\mathbf{x}-\mathbf{z}| \right| \mathbf{1}_{\{|\mathbf{x}-\mathbf{z}| \leq 1\}} + 1 \right) e^{-\tilde{c}\lambda|\mathbf{z}|} d\mathbf{z} p_0^\lambda(\mathbf{y}) \\
&= e^{-\tilde{c}\lambda} \int_{|\mathbf{z}| \geq r_1} e^{-\frac{\tilde{c}}{2}\lambda(|\mathbf{x}-\mathbf{z}|+|\mathbf{z}|)} e^{-\frac{\tilde{c}}{2}\lambda(|\mathbf{x}-\mathbf{z}|+|\mathbf{z}|)} \left(\left| \log \lambda |\mathbf{x}-\mathbf{z}| \right| \mathbf{1}_{\{|\mathbf{x}-\mathbf{z}| \leq 1\}} + 1 \right) d\mathbf{z} p_0^\lambda(\mathbf{y}) \\
&\leq e^{-\tilde{c}\lambda} e^{-\frac{\tilde{c}}{2}\lambda|\mathbf{x}|} \int_{|\mathbf{z}| \geq r_1} e^{-\frac{\tilde{c}}{2}\lambda(|\mathbf{x}-\mathbf{z}|+|\mathbf{z}|)} \left(\left| \log \lambda |\mathbf{x}-\mathbf{z}| \right| \mathbf{1}_{\{|\mathbf{x}-\mathbf{z}| \leq 1\}} + 1 \right) d\mathbf{z} p_0^\lambda(\mathbf{y}) \\
&\lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}|} p_0^\lambda(\mathbf{y}) .
\end{aligned}$$

For $|\mathbf{y}| < r_0 + \delta_0$, with small $\delta_0 > 0$, we have $p_0^\lambda(\mathbf{y}) \lesssim \lambda$. For such \mathbf{y} , $|\mathbf{x}| = |\mathbf{x} - \mathbf{y} + \mathbf{y}| \geq |\mathbf{x} - \mathbf{y}| - r_0 - \delta_0 \geq \frac{1}{2}|\mathbf{x} - \mathbf{y}| + \frac{R}{2} - r_0 - \delta_0 \geq \frac{1}{2}|\mathbf{x} - \mathbf{y}|$. Therefore, for $|\mathbf{x} - \mathbf{y}| > R$ and $|\mathbf{y}| < r_0 + \delta_0$ we have $\left| \text{Term}_1(\mathbf{x}, \mathbf{y}) \right| \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}|} p_0^\lambda(\mathbf{y}) \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}|} \lambda \lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{x}-\mathbf{y}|}$.

Therefore, for $|\mathbf{y}| \geq r_0 + \delta_0$ and $|\mathbf{x} - \mathbf{y}| > R$, we have $\left| \text{Term}_1(\mathbf{x}, \mathbf{y}) \right| \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}|} p_0^\lambda(\mathbf{y}) \lesssim e^{-c\lambda} e^{-c\lambda(|\mathbf{x}|+|\mathbf{y}|)} \lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{x}-\mathbf{y}|}$.

Bound on $\text{Term}_2(\mathbf{x}, \mathbf{y})$ of (10.30): We first note that $\nabla_{\mathbf{z}}\Theta_{\text{out}}(\mathbf{z}) = 0$ for $|\mathbf{z}| > r_2$. Since $|\mathbf{x}| > \frac{1}{2}R > r_2$, the integrand of $\text{Term}_2(\mathbf{x}, \mathbf{y})$ is supported away from $\mathbf{z} = \mathbf{x}$. Integration by parts yields

$$(10.32) \quad \text{Term}_2(\mathbf{x}, \mathbf{y}) = -2 \int_{\mathbb{R}^2} \nabla_{\mathbf{z}} \cdot \left[G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z}) \nabla_{\mathbf{z}}\Theta_{\text{out}}(\mathbf{z}) \right] G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y}) d\mathbf{z} .$$

We note this integration by parts can be justified even though there is a weak singularity of the integrand at $\mathbf{z} = \mathbf{y}$, and we remark on this at the conclusion of the proof. Bounding $\text{Term}_2(\mathbf{x}, \mathbf{y})$ using the Cauchy-Schwarz inequality we obtain:

$$\left| \text{Term}_2(\mathbf{x}, \mathbf{y}) \right| \leq 2 \left(\int_{\mathbb{R}^2} \left| \nabla_{\mathbf{z}} \cdot \left[G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z}) \nabla_{\mathbf{z}}\Theta_{\text{out}}(\mathbf{z}) \right] \right|^2 d\mathbf{z} \right)^{\frac{1}{2}} \cdot \left(\int_{\mathbb{R}^2} \left| G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y}) \right|^2 d\mathbf{z} \right)^{\frac{1}{2}} .$$

The second factor is bounded by a constant times λ^2 thanks to the L^2 bound on G_λ^{atom} given in (10.17). To bound the first factor note, due to the properties of $\Theta_{\text{out}}(\mathbf{z})$, that the support of the integrand is contained in: $r_1 \leq |\mathbf{z}| \leq r_2$ and $|\mathbf{x}| \geq r_3$. Therefore, $|\mathbf{x} - \mathbf{z}| \geq ||\mathbf{x}| - |\mathbf{z}|| \geq r_3 - r_2 > 0$. Therefore, by (10.5) and (10.6), for all $|\mathbf{x}| \geq r_3$:

$$\left| \nabla_{\mathbf{z}} \cdot \left[G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{z}) \nabla_{\mathbf{z}}\Theta_{\text{out}}(\mathbf{z}) \right] \right| \lesssim e^{-c\lambda|\mathbf{x}-\mathbf{z}|} \mathbf{1}_{\{r_1 \leq |\mathbf{z}| \leq r_2\}} \lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{x}|} .$$

It follows from (10.27) that

$$\begin{aligned}
\left| \text{Term}_2(\mathbf{x}, \mathbf{y}) \right| &\lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{x}|} \left(\int_{|\mathbf{z}| \leq r_2} \left| G_\lambda^{\text{atom}}(\mathbf{z}, \mathbf{y}) \right|^2 d\mathbf{z} \right)^{\frac{1}{2}} \\
(10.33) \quad &\lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{x}|} \lambda^2 \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{y}|} .
\end{aligned}$$

The bound on $\text{Term}_3(\mathbf{x}, \mathbf{y})$ is obtained in a manner similar to the bound on $\text{Term}_2(\mathbf{x}, \mathbf{y})$, but there is no need to integrate by parts.

We conclude the proof of (10.19) by remarking on the technical point raised above concerning the integration by parts leading to (10.32). Recall that

$$(-\Delta_{\mathbf{z}} + \lambda^2 V_0(\mathbf{z})) G_{\lambda}^{\text{atom}}(\mathbf{z}, \mathbf{y}) = \delta(\mathbf{z} - \mathbf{y}) + E_0^{\lambda} G_{\lambda}^{\text{atom}}(\mathbf{z}, \mathbf{y}) - p_0^{\lambda}(\mathbf{z}) p_0^{\lambda}(\mathbf{y}).$$

For fixed \mathbf{y} , $\mathbf{z} \mapsto p_0^{\lambda}(\mathbf{z}) p_0^{\lambda}(\mathbf{y})$ is C^{∞} by elliptic regularity because $(-\Delta_{\mathbf{z}} + \lambda^2 V_0(\mathbf{z}) - E_0^{\lambda}) p_0^{\lambda}(\mathbf{z}) = 0$ and $V_0 \in C^{\infty}$. Furthermore,

$$(-\Delta_{\mathbf{z}} + \lambda^2 V_0(\mathbf{z}) - E_0^{\lambda}) [G_{\lambda}^{\text{atom}}(\mathbf{z}, \mathbf{y}) - G_{\lambda}^{\text{free}}(\mathbf{z} - \mathbf{y})] = -\lambda^2 V_0(\mathbf{z}) G_{\lambda}^{\text{free}}(\mathbf{z}, \mathbf{y}) - p_0^{\lambda}(\mathbf{z}) p_0^{\lambda}(\mathbf{y}).$$

Since $V_0 \in C^{\infty}$, $\mathbf{z} \mapsto G_{\lambda}^{\text{free}}(\mathbf{z}, \mathbf{y}) \in H^{1-\varepsilon}(\mathbb{R}^2)$ ($\varepsilon > 0$ arbitrary), we have by elliptic regularity that $G_{\lambda}^{\text{atom}}(\mathbf{z}, \mathbf{y}) - G_{\lambda}^{\text{free}}(\mathbf{z} - \mathbf{y}) \in H_{\text{loc}}^{3-\varepsilon}(\mathbb{R}^2)$. Furthermore by (10.4), for fixed \mathbf{y}

$$G_{\lambda}^{\text{atom}}(\mathbf{z}, \mathbf{y}) = c_0 \log |\mathbf{z} - \mathbf{y}| + j(\mathbf{z}, \mathbf{y}) \quad \text{for } \mathbf{z} \text{ near } \mathbf{y},$$

where $\mathbf{z} \mapsto j(\mathbf{z}, \mathbf{y}) \in H_{\text{loc}}^{2-\varepsilon}(\mathbb{R}^2)$. This makes it easy to justify the integration by parts. For example, replace $G_{\lambda}^{\text{atom}}(\mathbf{z}, \mathbf{y})$ by $\frac{1}{2} c_0 \log [|\mathbf{z} - \mathbf{y}|^2 + \tau^2] + j(\mathbf{z}, \mathbf{y})$, integrate by parts and pass to the limit $\tau \rightarrow 0^+$. This concludes the proof of (10.19). Since the proof of the bound (10.20) follows from a very similar argument, we omit it. This completes the proof of Theorem 10.3.

10.3. Kernels. Our goal will be to construct the Green's kernel for a Hamiltonian $H_{\Gamma}^{\lambda} = -\Delta + V_{\Gamma}^{\lambda}(\mathbf{x}) - E_0^{\lambda}$, with potential V_{Γ}^{λ} defined via superposition involving translates of the atomic potential, V_0 , centered at the sites of a discrete set Γ . The construction of this Green's function, $G_{\lambda}^{\Gamma}(\mathbf{x}, \mathbf{y})$ makes use of some technical tools developed in this section.

We work with integral operators of the form

$$(10.34) \quad f \mapsto A_{\lambda}[f](\mathbf{x}) \equiv \int_{\mathbb{R}^2} A_{\lambda}(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y}.$$

We shall use the notation $A_{\lambda}f$ and $A_{\lambda}[f]$ to denote such operators and occasionally omit the λ dependence.

Definition 10.4 (Main Kernel). The function $A_{\lambda}(\mathbf{x}, \mathbf{y}) : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is called a main kernel if there exist positive constants R, c, C_1, C_2 and λ_0 such that for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$ with $\mathbf{x} \neq \mathbf{y}$ we have

$$(10.35) \quad |A_{\lambda}(\mathbf{x}, \mathbf{y})| \leq C_1 \left[\lambda^4 + \left| \log |\mathbf{x} - \mathbf{y}| \right| \right] \mathbf{1}_{|\mathbf{x} - \mathbf{y}| \leq R} + C_2 e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|}$$

for all $\lambda \geq \lambda_0$.

By Theorem 10.3, the atomic Green's function $G_{\lambda}^{\text{atom}}(\mathbf{x}, \mathbf{y})$ is a main kernel.

Definition 10.5 (Error Kernel). The function $\mathcal{E}_{\lambda}(\mathbf{x}, \mathbf{y}) : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is called a *error kernel* if there exist positive constants c, C and λ_0 such that for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$

$$(10.36) \quad |\mathcal{E}_{\lambda}(\mathbf{x}, \mathbf{y})| \leq C e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|}$$

for all $\lambda \geq \lambda_0$.

If A and B are operators with kernels given by $A(\mathbf{x}, \mathbf{y})$ and $B(\mathbf{x}, \mathbf{y})$, respectively, then AB is defined to be the operator with kernel $(AB)(\mathbf{x}, \mathbf{y})$ given by

$$(10.37) \quad (AB)(\mathbf{x}, \mathbf{y}) \equiv \int_{\mathbb{R}^2} A(\mathbf{x}, \mathbf{z}) B(\mathbf{z}, \mathbf{y}) d\mathbf{z}$$

Remark 10.6. If $\mathcal{E}(\mathbf{x}, \mathbf{y})$ is an error kernel, then $\lambda^p \mathcal{E}(\mathbf{x}, \mathbf{y})$ is an error kernel for any $p \geq 0$. To see this, replace the constant c in (10.36) by a slightly smaller positive constant, c' .

Lemma 10.7.

Let K_λ arise from a main kernel and \mathcal{E}_λ arise from an error kernel.

(1) Then,

$$(10.38) \quad \tilde{\mathcal{E}}_\lambda = I - (I - \mathcal{E}_\lambda)^{-1} = \sum_{l \geq 1} \mathcal{E}_\lambda^l$$

arises from an error kernel.

(2) The operators $\mathcal{E}_\lambda K_\lambda$ and $K_\lambda \mathcal{E}_\lambda$ arise from error kernels.

(3) The operator $e^{-c\lambda} K_\lambda^2$, where $c > 0$, arises from an error kernel.

The proof of Lemma 10.7 is presented in Appendix A

10.4. Green's kernel for a set of atoms centered on points of a discrete set, Γ . Let Γ denote a discrete subset of \mathbb{R}^2 , which we refer to as a set of *nuclei*. The set Γ may be finite or infinite. We assume that

$$(10.39) \quad \inf\{|\mathbf{v} - \mathbf{w}| : \mathbf{v}, \mathbf{w} \in \Gamma, \mathbf{v} \neq \mathbf{w}\} \geq r_{\min} > 2r_0.$$

At sites $\omega \in \Gamma$ we center identical *atoms* described by the atomic potential V_0 :

$$(10.40) \quad V_\Gamma^\lambda(\mathbf{x}) = \sum_{\omega \in \Gamma} \lambda^2 V_\omega(\mathbf{x}), \quad \text{where } V_\omega(\mathbf{x}) \equiv V_0(\mathbf{x} - \omega).$$

Example 10.8. Some choices of Γ which are of interest to us are:

(1) $\Gamma = \mathbb{H} = \Lambda_A \cup \Lambda_B$, the bulk honeycomb structure.

(2) $\Gamma = \Lambda_I$, $I = A, B$, the A - and B - sublattices.

(3) $\Gamma = \mathbb{H}_\# = \{\mathbf{v}_I + n_1 \mathbf{v}_1 + n_2 \mathbf{v}_2 : n_1 \geq 0, n_2 \in \mathbb{Z}\}$, the set of lattice points in a zigzag-terminated honeycomb structure.

Our goal will be to construct the Green's kernel $G_\Gamma^\lambda(\mathbf{x}, \mathbf{y})$ associated with the operator

$$(10.41) \quad H_\Gamma^\lambda = -\Delta + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda,$$

where E_0^λ is the ground state energy of $H_{\text{atom}}^\lambda = -\Delta + \lambda^2 V_0$; see (3.4).

Recall G_λ^{atom} which satisfies

$$\begin{aligned} (-\Delta + \lambda^2 V_0(\mathbf{x}) - E_0^\lambda) G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) &= \delta(\mathbf{x} - \mathbf{y}) - p_0^\lambda(\mathbf{x}) p_0^\lambda(\mathbf{y}), \\ \int_{\mathbb{R}^2} G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) p_0^\lambda(\mathbf{x}) d\mathbf{x} &= 0, \\ G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) &= G_\lambda^{\text{atom}}(\mathbf{y}, \mathbf{x}). \end{aligned}$$

Recalling $r_j, j = 1, 2, 3$ specified in (10.26), we further introduce r_4 such that

$$(10.42) \quad 0 < r_0 < r_1 < r_2 < r_3 < r_4 < \frac{1}{2} r_{\min}, \quad (r_{\min} > 2r_0),$$

where r_{\min} is a lower bound for the minimum distance between points in Γ ; see (10.39).

Introduce the smooth cutoff function $\Theta_0(\mathbf{x})$ satisfying:

$0 \leq \Theta_0 \leq 1$ on \mathbb{R}^2 , $\Theta_0(\mathbf{x}) = 1$ for $\mathbf{x} \in B_{r_3}(0)$, and $\Theta_0(\mathbf{x}) = 0$ for $\mathbf{x} \notin B_{r_4}(0)$.

For $\omega \in \Gamma$, define $\Theta_\omega(\mathbf{x}) = \Theta_0(\mathbf{x} - \omega)$. Finally, let

$$(10.43) \quad \Theta_{\text{free}}(\mathbf{x}) \equiv 1 - \sum_{\omega \in \Gamma} \Theta_\omega(\mathbf{x}).$$

Then, $0 \leq \Theta_{\text{free}} \leq 1$ on \mathbb{R}^2 ; Θ_{free} is smooth and supported away from Γ . In particular for all $\omega \in \Gamma$, $\Theta_{\text{free}} = 0$ in $B_{r_3}(\omega)$.

We write $p_\omega^\lambda(\mathbf{x}) \equiv p_0^\lambda(\mathbf{x} - \omega)$, where $p_0^\lambda(\mathbf{x})$ is the ground state of $H_{\text{atom}}^\lambda = -\Delta + \lambda^2 V_0(\mathbf{x})$. Thus, $p_\omega^\lambda(\mathbf{x})$ is the ground state of $-\Delta + \lambda^2 V_\omega(\mathbf{x})$. We also express the translated atomic Green's kernel as

$$(10.44) \quad G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) = G_\lambda^{\text{atom}}(\mathbf{x} - \omega, \mathbf{y} - \omega).$$

For any $f \in L^2(\mathbb{R}^2)$ we may write:

$$(10.45) \quad f(\mathbf{x}) = \sum_{\omega \in \Gamma} (\Theta_\omega f)(\mathbf{x}) + (\Theta_{\text{free}} f)(\mathbf{x}),$$

and for each $\omega \in \Gamma$, we have by (10.7)

$$(10.46) \quad \begin{aligned} \Theta_\omega(\mathbf{x})f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + \lambda^2 V_\omega(\mathbf{x}) - E_0^\lambda \right) \int_{\mathbb{R}^2} G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) (\Theta_\omega(\mathbf{y})f(\mathbf{y})) d\mathbf{y} \\ &+ \langle p_\omega^\lambda, \Theta_\omega f \rangle_{L^2(\mathbb{R}^2)} p_\omega^\lambda(\mathbf{x}), \end{aligned}$$

and by (10.8)

$$(10.47) \quad \int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{x}) \left[\int_{\mathbb{R}^2} G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) (\Theta_\omega(\mathbf{y})f(\mathbf{y})) d\mathbf{y} \right] d\mathbf{x} = 0.$$

Next we express V_Γ^λ as:

$$V_\Gamma^\lambda(\mathbf{x}) = \lambda^2 V_\omega(\mathbf{x}) + \sum_{\omega' \in \Gamma \setminus \{\omega\}} \lambda^2 V_{\omega'}(\mathbf{x}),$$

and therefore by (10.46)

$$(10.48) \quad \begin{aligned} \Theta_\omega(\mathbf{x})f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda \right) \int_{\mathbb{R}^2} G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) \Theta_\omega(\mathbf{y}) \cdot f(\mathbf{y}) d\mathbf{y} \\ &- \sum_{\omega' \in \Gamma \setminus \{\omega\}} \lambda^2 V_{\omega'}(\mathbf{x}) \int_{\mathbb{R}^2} G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) \Theta_\omega(\mathbf{y}) \cdot f(\mathbf{y}) d\mathbf{y} + \langle p_\omega^\lambda, \Theta_\omega f \rangle_{L^2(\mathbb{R}^2)} p_\omega^\lambda(\mathbf{x}). \end{aligned}$$

Similarly,

$$(10.49) \quad \begin{aligned} \Theta_{\text{free}}(\mathbf{x})f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} - E_0^\lambda \right) \int_{\mathbb{R}^2} G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) (\Theta_{\text{free}}(\mathbf{y})f(\mathbf{y})) d\mathbf{y} \\ &= \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda \right) \int_{\mathbb{R}^2} G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) \Theta_{\text{free}}(\mathbf{y}) \cdot f(\mathbf{y}) d\mathbf{y} \\ &- V_\Gamma^\lambda(\mathbf{x}) \int_{\mathbb{R}^2} G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) \Theta_{\text{free}}(\mathbf{y}) \cdot f(\mathbf{y}) d\mathbf{y}. \end{aligned}$$

We note that $V_\Gamma^\lambda(\mathbf{x}) \equiv 0$ on the support of Θ_{free} .

Now summing (10.48) over $\omega \in \Gamma$ and adding the result to (10.49), we have by (10.45) the following:

$$\begin{aligned}
(10.50) \quad f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + V_{\Gamma}^{\lambda}(\mathbf{x}) - E_0^{\lambda} \right) \cdot \\
&\quad \int_{\mathbb{R}^2} \left[\sum_{\omega \in \Gamma} G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) \Theta_{\omega}(\mathbf{y}) + G_{\lambda}^{\text{free}}(\mathbf{x} - \mathbf{y}) \Theta_{\text{free}}(\mathbf{y}) \right] \cdot f(\mathbf{y}) \, d\mathbf{y} \\
&\quad - \int_{\mathbb{R}^2} \left[\sum_{\substack{\omega, \omega' \in \Gamma \\ \omega \neq \omega'}} \lambda^2 V_{\omega'}(\mathbf{x}) G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) \Theta_{\omega}(\mathbf{y}) + V_{\Gamma}^{\lambda}(\mathbf{x}) G_{\lambda}^{\text{free}}(\mathbf{x} - \mathbf{y}) \Theta_{\text{free}}(\mathbf{y}) \right] \cdot f(\mathbf{y}) \, d\mathbf{y} \\
&\quad + \sum_{\omega \in \Gamma} \langle \Theta_{\omega} p_{\omega}^{\lambda}, f \rangle_{L^2(\mathbb{R}^2)} p_{\omega}^{\lambda}(\mathbf{x}) .
\end{aligned}$$

Introduce the kernels K_0^{λ} and \mathcal{E}_0^{λ} :

$$(10.51) \quad K_0^{\lambda}(\mathbf{x}, \mathbf{y}) \equiv \sum_{\omega \in \Gamma} G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) \Theta_{\omega}(\mathbf{y}) + G_{\lambda}^{\text{free}}(\mathbf{x} - \mathbf{y}) \Theta_{\text{free}}(\mathbf{y})$$

$$(10.52) \quad \mathcal{E}_0^{\lambda}(\mathbf{x}, \mathbf{y}) \equiv \sum_{\substack{\omega, \omega' \in \Gamma \\ \omega \neq \omega'}} \lambda^2 V_{\omega'}(\mathbf{x}) G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) \Theta_{\omega}(\mathbf{y}) + V_{\Gamma}^{\lambda}(\mathbf{x}) G_{\lambda}^{\text{free}}(\mathbf{x} - \mathbf{y}) \Theta_{\text{free}}(\mathbf{y}) .$$

Equation (10.50) is equivalent to

$$\begin{aligned}
(10.53) \quad f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + V_{\Gamma}^{\lambda}(\mathbf{x}) - E_0^{\lambda} \right) \int_{\mathbb{R}^2} K_0^{\lambda}(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) \, d\mathbf{y} \\
&\quad + \sum_{\omega \in \Gamma} \langle \Theta_{\omega} p_{\omega}^{\lambda}, f \rangle_{L^2(\mathbb{R}^2)} p_{\omega}^{\lambda}(\mathbf{x}) - \int_{\mathbb{R}^2} \mathcal{E}_0^{\lambda}(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) \, d\mathbf{y} .
\end{aligned}$$

and in any even more compact form :

$$\begin{aligned}
(10.54) \quad f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + V_{\Gamma}^{\lambda}(\mathbf{x}) - E_0^{\lambda} \right) K_0^{\lambda}[f](\mathbf{x}) - \mathcal{E}_0^{\lambda}[f](\mathbf{x}) \\
&\quad + \sum_{\omega \in \Gamma} \langle \Theta_{\omega} p_{\omega}^{\lambda}, f \rangle_{L^2(\mathbb{R}^2)} p_{\omega}^{\lambda}(\mathbf{x}) .
\end{aligned}$$

Proposition 10.9. $K_0^{\lambda}(\mathbf{x}, \mathbf{y})$ is a main kernel in the sense of Definition 10.4 and $\mathcal{E}_0^{\lambda}(\mathbf{x}, \mathbf{y})$ is an error kernel in the sense of (10.5).

Proof of Proposition 10.9: We first prove that $K_0^{\lambda}(\mathbf{x}, \mathbf{y})$, displayed in (10.51), is a main kernel. Note that for each $\mathbf{y} \in \mathbb{R}^2$ there is at most one $\omega = \omega_{\mathbf{y}} \in \Gamma$ with $\mathbf{y} \in \text{supp } \Theta_{\omega} \subset \{\mathbf{y} : |\mathbf{y} - \omega| \leq r_4\}$. Therefore, for the first term in (10.51) we have by Theorem 10.3 the bound

$$\begin{aligned}
\left| \sum_{\omega \in \Gamma} G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) \Theta_{\omega}(\mathbf{y}) \right| &\leq \left| G_{\lambda, \omega_{\mathbf{y}}}^{\text{atom}}(\mathbf{x}, \mathbf{y}) \right| \\
&\lesssim C \left[\lambda^4 + |\log |\mathbf{x} - \mathbf{y}| | \right] \mathbf{1}_{\{|\mathbf{x} - \mathbf{y}| \leq R\}} + e^{-c\lambda} e^{-c\lambda |\mathbf{x} - \mathbf{y}|} .
\end{aligned}$$

Furthermore by (10.5), the second term in (10.51) satisfies the bound

$$\begin{aligned} \left| G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) \Theta_{\text{free}}(\mathbf{y}) \right| &\leq \left| G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) \right| \\ &\lesssim C \left[\lambda^4 + |\log |\mathbf{x} - \mathbf{y}|| \right] \mathbf{1}_{\{|\mathbf{x} - \mathbf{y}| \leq R\}} + e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|} \end{aligned}$$

Adding the two previous bounds we conclude that $K_0^\lambda(\mathbf{x}, \mathbf{y})$ is a main kernel.

We now prove that $\mathcal{E}_0^\lambda(\mathbf{x}, \mathbf{y})$ given by (10.52) is an error kernel. Consider the sum in (10.52). This sum is non-zero at $(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^2 \times \mathbb{R}^2$, if there are distinct points $\omega'_x, \omega_y \in \Gamma$ with $\mathbf{x} \in \text{supp } V_{\omega'_x}$ and $\mathbf{y} \in \text{supp } \Theta_{\omega_y}$. The choice of points $\omega'_x, \omega_y \in \Gamma$ is unique. We have $\mathbf{y} \in B_{r_4}(\omega_y)$ and $\mathbf{x} \notin B_{r_4 + \delta_1}(\omega_y)$, where $\delta_1 > 0$. Therefore, part (3) of Theorem 10.3 implies

$$\begin{aligned} \left| \sum_{\substack{\omega, \omega' \in \Gamma \\ \omega \neq \omega'}} \lambda^2 V_{\omega'}(\mathbf{x}) G_{\lambda, \omega}^{\text{atom}}(\mathbf{x}, \mathbf{y}) \Theta_\omega(\mathbf{y}) \right| &\leq \lambda^2 |V_{\omega'_x}(\mathbf{x})| |G_{\lambda, \omega_y}^{\text{atom}}(\mathbf{x}, \mathbf{y})| \Theta_{\omega_y}(\mathbf{y}) \\ &\leq \lambda^2 e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|} \lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{x} - \mathbf{y}|}. \end{aligned}$$

For the second term in (10.52), if $\mathbf{x} \in \text{supp } V_\Gamma$ and $\mathbf{y} \in \text{supp } \Theta_{\text{free}}$, then $|\mathbf{x} - \mathbf{y}| \geq r_3 - r_0 > 0$. Therefore, $G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) \lesssim e^{-c\lambda|\mathbf{x} - \mathbf{y}|} \lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{x} - \mathbf{y}|}$. It follows that for some $\omega = \omega_x \in \Gamma$:

$$\left| V_\Gamma^\lambda(\mathbf{x}) G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) \Theta_{\text{free}}(\mathbf{y}) \right| \lesssim \lambda^2 \left| V_{\omega_x}(\mathbf{x}) G_\lambda^{\text{free}}(\mathbf{x} - \mathbf{y}) \right| \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|}.$$

The latter two bounds imply that $\mathcal{E}_0^\lambda(\mathbf{x}, \mathbf{y})$, defined in (10.52), is an error kernel. The proof of Proposition 10.9 is now complete.

Remark 10.10. At this stage we wish to remark that if Γ is translation invariant by some vector, then K_0^λ and \mathcal{E}_0^λ inherit this invariance. In particular, for $\Gamma = \mathbb{H}_\#$, the zigzag truncation of the honeycomb \mathbb{H} , we have $K_0^\lambda(\mathbf{x} + \mathbf{v}_2, \mathbf{y} + \mathbf{v}_2) = K_0^\lambda(\mathbf{x}, \mathbf{y})$ and $\mathcal{E}_0^\lambda(\mathbf{x} + \mathbf{v}_2, \mathbf{y} + \mathbf{v}_2) = \mathcal{E}_0^\lambda(\mathbf{x}, \mathbf{y})$.

Introduce the orthogonal subspaces \mathcal{X}_Γ :

$$(10.55) \quad \mathcal{X}_\Gamma \equiv \text{span} \left\{ p_\omega^\lambda : \omega \in \Gamma \right\}^\perp = \left\{ f \in L^2(\mathbb{R}^2) : \langle p_\omega^\lambda, f \rangle_{L^2(\mathbb{R}^2)} = 0, \omega \in \Gamma \right\},$$

and the orthogonal projections:

$$(10.56) \quad \Pi_\Gamma^\lambda : L^2(\mathbb{R}^2) \rightarrow \mathcal{X}_\Gamma, \quad \tilde{\Pi}_\Gamma^\lambda = I - \Pi_\Gamma^\lambda : L^2(\mathbb{R}^2) \rightarrow \text{span} \left\{ p_\omega^\lambda : \omega \in \Gamma \right\}.$$

We seek the integral kernel for the inverse of the operator $\Pi_\Gamma^\lambda (H_\Gamma^\lambda - E_0^\lambda - \Omega) \Pi_\Gamma^\lambda$ on \mathcal{X}_Γ .

The operator $f \mapsto K_0^\lambda f$ (see (10.51), (10.53)) defines an approximate inverse of $H_\Gamma^\lambda - E_0^\lambda - \Omega$ on the range of Π_Γ^λ but we do not have that $\Pi_\Gamma^\lambda K_0^\lambda[f] = K_0^\lambda[f]$. Our next step is to correct K_0^λ in order to achieve the desired projection.

Recall that the set $\{p_\omega^\lambda : \omega \in \Gamma\}$ is not orthonormal, but only nearly so; see Proposition 4.2. The following lemma gives a representation for $\tilde{\Pi}_\Gamma^\lambda$, defined in (10.56).

Lemma 10.11. $\tilde{\Pi}_\Gamma^\lambda = I - \Pi_\Gamma^\lambda$, the orthogonal projection of $L^2(\mathbb{R}^2)$ onto $\text{span}\{p_\omega^\lambda : \omega \in \Gamma\}$, is given by

$$(10.57) \quad \tilde{\Pi}_\Gamma^\lambda[g](\mathbf{x}) = \sum_{\omega, \hat{\omega} \in \Gamma} M^{\omega, \hat{\omega}} \langle p_{\hat{\omega}}^\lambda, g \rangle p_\omega^\lambda(\mathbf{x}),$$

where $M^{\omega, \hat{\omega}}$ satisfies the estimate

$$(10.58) \quad | M^{\omega, \hat{\omega}} - \delta_{\omega, \hat{\omega}} | \lesssim e^{-c\lambda} e^{-c\lambda |\hat{\omega} - \omega|} .$$

Proof of Lemma 10.11: If we define $\tilde{\Pi}_\Gamma^\lambda[g]$ by (10.57), then for all $g \in L^2(\mathbb{R}^2)$

$$(10.59) \quad \begin{aligned} \langle p_{\omega'}^\lambda, g \rangle &= \langle p_{\omega'}^\lambda, \tilde{\Pi}_\Gamma^\lambda[g] \rangle = \sum_{\omega, \hat{\omega} \in \Gamma} M^{\omega, \hat{\omega}} \langle p_{\hat{\omega}}^\lambda, g \rangle \langle p_{\omega'}^\lambda, p_\omega^\lambda \rangle \\ &= \sum_{\omega, \hat{\omega} \in \Gamma} \left(\sum_{\omega \in \Gamma} \langle p_{\omega'}^\lambda, p_\omega^\lambda \rangle M^{\omega, \hat{\omega}} \right) \langle p_{\hat{\omega}}^\lambda, g \rangle \end{aligned}$$

Therefore, $\tilde{\Pi}_\Gamma^\lambda$ is as required provided:

$$\sum_{\omega \in \Gamma} \langle p_{\omega'}^\lambda, p_\omega^\lambda \rangle M^{\omega, \hat{\omega}} = \delta_{\omega' \hat{\omega}} .$$

We claim that if $\omega', \omega \in \Gamma$ are distinct, then

$$(10.60) \quad | \langle p_{\omega'}^\lambda, p_\omega^\lambda \rangle | \lesssim e^{-c'\lambda|\omega - \omega'|} e^{-c'\lambda} .$$

Indeed, if $\omega \neq \omega'$

$$\begin{aligned} | \langle p_{\omega'}^\lambda, p_\omega^\lambda \rangle | &\leq \int_{B_{r_4}(\omega)} p_{\omega'}^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{x}) d\mathbf{x} + \int_{B_{r_4}(\omega')} p_{\omega'}^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{x}) d\mathbf{x} \\ &\quad + \int_{\mathbb{R}^2 \setminus B_{r_4}(\omega) \cup B_{r_4}(\omega')} p_{\omega'}^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{x}) d\mathbf{x} \\ &\leq \int_{B_{r_4}(\omega)} [e^{-c\lambda|\mathbf{x} - \omega'|}] \cdot [\lambda^2] d\mathbf{x} + \int_{B_{r_4}(\omega')} [\lambda^2] \cdot [e^{-c\lambda|\mathbf{x} - \omega|}] d\mathbf{x} \\ &\quad + \int_{\mathbb{R}^2 \setminus B_{r_4}(\omega) \cup B_{r_4}(\omega')} e^{-c\lambda|\mathbf{x} - \omega|} \cdot e^{-c\lambda|\mathbf{x} - \omega'|} d\mathbf{x} \lesssim e^{-c'\lambda|\omega - \omega'|} e^{-c'\lambda} . \end{aligned}$$

Since also $p_\omega^\lambda(\mathbf{x}) = p_0^\lambda(\mathbf{x} - \omega)$ is normalized in $L^2(\mathbb{R}^2)$, we have

$$(10.61) \quad | \langle p_{\omega'}^\lambda, p_\omega^\lambda \rangle - \delta_{\omega, \omega'} | \lesssim e^{-c\lambda} e^{-c\lambda|\omega - \omega'|} .$$

Let $P = (\langle p_{\omega'}^\lambda, p_\omega^\lambda \rangle)_{\omega, \omega' \in \Gamma}$ and for any $\nu \in \mathbb{R}^2$, $|\nu| = 1$, let $D = (e^{\bar{c}\lambda\nu \cdot \omega} \delta_{\omega, \omega'})_{\omega, \omega' \in \Gamma}$, with \bar{c} smaller than the constant c appearing in (10.61). Then, $D P D^{-1} = (e^{\bar{c}\lambda\nu \cdot (\omega - \omega')} \langle p_{\omega'}^\lambda, p_\omega^\lambda \rangle)_{\omega, \omega' \in \Gamma} = (\tilde{p}_{\omega, \omega'})$ with

$$| \tilde{p}_{\omega, \omega'} - \delta_{\omega, \omega'} | \lesssim e^{-c'\lambda|\omega - \omega'|} e^{-c'\lambda} .$$

by (10.61). Hence, $D P^{-1} D^{-1} = (D P D^{-1})^{-1}$ has an (ω, ω') -entry that differs from $\delta_{\omega, \omega'}$ by at most $e^{-\bar{c}\lambda}$. That is, $| [e^{\bar{c}\lambda\nu \cdot (\omega - \omega')} M^{\omega, \omega'}] - \delta_{\omega, \omega'} | \lesssim e^{-c\lambda}$ and hence

$$| e^{\bar{c}\lambda\nu \cdot (\omega - \omega')} [M^{\omega, \omega'} - \delta_{\omega, \omega'}] | \lesssim e^{-c\lambda}$$

for all $\omega, \omega' \in \Gamma$ and all unit vectors $\nu \in \mathbb{R}^2$. Optimizing over ν gives

$$| M^{\omega, \omega'} - \delta_{\omega, \omega'} | \lesssim e^{-c\lambda} e^{-\bar{c}\lambda|\omega - \omega'|} .$$

This completes the proof of Lemma 10.11.

By (10.54), after subtracting and adding $\tilde{\Pi}_\Gamma^\lambda K_0^\lambda$, we have

$$\begin{aligned}
(10.62) \quad f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda \right) \left[K_0^\lambda[f](\mathbf{x}) - \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right) [f] \right] \\
&\quad + \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda \right) \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right) [f] \\
&\quad - \mathcal{E}_0^\lambda[f](\mathbf{x}) + \sum_{\omega \in \Gamma} \langle \Theta_\omega p_\omega^\lambda, f \rangle_{L^2(\mathbb{R}^2)} p_\omega^\lambda(\mathbf{x}) .
\end{aligned}$$

Here, we have arranged for the expression within the square brackets in (10.62):

$$(10.63) \quad K_1^\lambda[f] \equiv K_0^\lambda[f] - \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right) [f],$$

to be orthogonal to the translated atomic ground states p_ω^λ , for all $\omega \in \Gamma$. Our next task is to show that the remaining terms in (10.62) comprise an error kernel.

Proposition 10.12. *The operators $\tilde{\Pi}_\Gamma^\lambda K_0^\lambda$ and $\left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda \right) \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)$ derive from error kernels in the sense of Definition 10.5.*

Proof of Proposition 10.12: By (10.57)

$$\begin{aligned}
(10.64) \quad \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right) [f](\mathbf{x}) &= \sum_{\omega, \hat{\omega} \in \Gamma} M^{\omega, \hat{\omega}} \langle p_\omega^\lambda, K_0^\lambda[f] \rangle p_{\hat{\omega}}^\lambda(\mathbf{x}) \\
&= \sum_{\omega, \hat{\omega} \in \Gamma} M^{\omega, \hat{\omega}} \int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{y}) \int_{\mathbb{R}^2} K_0^\lambda(\mathbf{y}, \mathbf{z}) f(\mathbf{z}) d\mathbf{z} d\mathbf{y} p_{\hat{\omega}}^\lambda(\mathbf{x}) \\
&= \sum_{\omega, \hat{\omega} \in \Gamma} M^{\omega, \hat{\omega}} \int_{\mathbb{R}^2} \left[\int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{y}) K_0^\lambda(\mathbf{y}, \mathbf{z}) d\mathbf{y} p_{\hat{\omega}}^\lambda(\mathbf{x}) \right] f(\mathbf{z}) d\mathbf{z} \\
&= \int_{\mathbb{R}^2} \left[\sum_{\omega, \hat{\omega} \in \Gamma} M^{\omega, \hat{\omega}} \int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{y}) K_0^\lambda(\mathbf{y}, \mathbf{z}) d\mathbf{y} p_{\hat{\omega}}^\lambda(\mathbf{x}) \right] f(\mathbf{z}) d\mathbf{z}
\end{aligned}$$

Thus,

$$\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right) (\mathbf{x}, \mathbf{z}) = \int_{\mathbb{R}^2} \left[\sum_{\omega, \hat{\omega} \in \Gamma} M^{\omega, \hat{\omega}} p_{\hat{\omega}}^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{y}) \right] K_0^\lambda(\mathbf{y}, \mathbf{z}) d\mathbf{y} ,$$

where K_0^λ is given by (10.51):

$$(10.65) \quad K_0^\lambda(\mathbf{y}, \mathbf{z}) \equiv \sum_{\omega' \in \Gamma} G_{\lambda, \omega'}^{\text{atom}}(\mathbf{y}, \mathbf{z}) \Theta_{\omega'}(\mathbf{z}) + G_\lambda^{\text{free}}(\mathbf{y} - \mathbf{z}) \Theta_{\text{free}}(\mathbf{z}) .$$

Now decompose $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right) (\mathbf{x}, \mathbf{z})$ has follows:

$$\begin{aligned}
\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right) (\mathbf{x}, \mathbf{z}) &= \int_{\mathbb{R}^2} \left[\sum_{\substack{\omega, \hat{\omega} \in \Gamma \\ \omega \neq \hat{\omega}}} M^{\omega, \hat{\omega}} p_{\hat{\omega}}^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{y}) \right] K_0^\lambda(\mathbf{y}, \mathbf{z}) d\mathbf{y} \\
&\quad + \int_{\mathbb{R}^2} \left[\sum_{\omega \in \Gamma} M^{\omega, \omega} p_\omega^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{y}) \right] K_0^\lambda(\mathbf{y}, \mathbf{z}) d\mathbf{y} \\
(10.66) \quad &\equiv \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_1 (\mathbf{x}, \mathbf{z}) + \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_2 (\mathbf{x}, \mathbf{z}) .
\end{aligned}$$

We prove that each term in (10.66) is an error kernel, *i.e.* $\left| \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_j (\mathbf{x}, \mathbf{z}) \right| \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{z}|}$ for $j = 1, 2$. For $\omega \neq \hat{\omega}$ we have by (10.58) that

$$|M^{\omega, \hat{\omega}}| \lesssim e^{-c'\lambda|\omega-\hat{\omega}|} e^{-c'\lambda}.$$

We may therefore write:

$$(10.67) \quad |M^{\omega, \hat{\omega}} p_{\hat{\omega}}^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{y})| \leq e^{-c'\lambda|\omega-\hat{\omega}|} e^{-\tilde{c}\lambda} p_{\hat{\omega}}^\lambda(\mathbf{x}) \cdot e^{-\tilde{c}\lambda} p_\omega^\lambda(\mathbf{y}).$$

Next, using (3.5) we bound $e^{-\tilde{c}\lambda} p_{\hat{\omega}}^\lambda(\mathbf{x})$ and $e^{-\tilde{c}\lambda} p_\omega^\lambda(\mathbf{y})$ as follows:

$$\begin{aligned}
e^{-\tilde{c}\lambda} p_{\hat{\omega}}^\lambda(\mathbf{x}) &\lesssim \left(e^{-c'\lambda} \mathbf{1}_{\{|\mathbf{x}-\hat{\omega}| \leq r_1\}} + e^{-c'\lambda} e^{-c\lambda|\mathbf{x}-\hat{\omega}|} \right) \\
(10.68) \quad &\lesssim \left(e^{-\frac{c'}{2}\lambda} e^{-\frac{c'}{2r_1}\lambda|\mathbf{x}-\hat{\omega}|} \mathbf{1}_{\{|\mathbf{x}-\hat{\omega}| \leq r_1\}} + e^{-c'\lambda} e^{-c\lambda|\mathbf{x}-\hat{\omega}|} \right)
\end{aligned}$$

Therefore, $e^{-\tilde{c}\lambda} p_{\hat{\omega}}^\lambda(\mathbf{x}) \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\hat{\omega}|}$ and similarly $e^{-\tilde{c}\lambda} p_\omega^\lambda(\mathbf{y}) \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{y}-\omega|}$. Substituting these bounds into (10.67), we obtain for some $c > 0$

$$\begin{aligned}
|M^{\omega, \hat{\omega}} p_{\hat{\omega}}^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{y})| &\lesssim e^{-c\lambda} e^{-c\lambda|\omega-\hat{\omega}|} e^{-c\lambda|\mathbf{x}-\hat{\omega}|} e^{-c\lambda|\mathbf{y}-\omega|} \\
&\lesssim e^{-c\lambda} e^{-\frac{c}{2}\lambda|\mathbf{x}-\mathbf{y}|} \times e^{-\frac{c}{2}\lambda|\omega-\hat{\omega}|} e^{-\frac{c}{2}\lambda|\mathbf{x}-\hat{\omega}|} e^{-\frac{c}{2}\lambda|\mathbf{y}-\omega|},
\end{aligned}$$

since $|\mathbf{x}-\mathbf{y}| \leq |\mathbf{x}-\hat{\omega}| + |\omega-\hat{\omega}| + |\mathbf{y}-\omega|$. Therefore, for some c' which is independent of λ :

$$\sum_{\substack{\omega, \hat{\omega} \in \Gamma \\ \omega \neq \hat{\omega}}} |M^{\omega, \hat{\omega}} p_{\hat{\omega}}^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{y})| \lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{x}-\mathbf{y}|}$$

and therefore $\sum_{\substack{\omega, \hat{\omega} \in \Gamma \\ \omega \neq \hat{\omega}}} M^{\omega, \hat{\omega}} p_{\hat{\omega}}^\lambda(\mathbf{x}) p_\omega^\lambda(\mathbf{y})$ is therefore an error kernel. And since K_0^λ is a main kernel we have, by the expression for $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_1 (\mathbf{x}, \mathbf{z})$ in (10.66), and by part 2 of Lemma 10.7, that $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_1 (\mathbf{x}, \mathbf{z})$ is an error kernel.

We next prove that $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda\right)_2(\mathbf{x}, \mathbf{z})$, defined in (10.66) is an error kernel. Using (10.65) we have

$$\begin{aligned}
\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda\right)_2(\mathbf{x}, \mathbf{z}) &\equiv \sum_{\omega \in \Gamma} M^{\omega, \omega} p_\omega^\lambda(\mathbf{x}) \int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{y}) K_0^\lambda(\mathbf{y}, \mathbf{z}) d\mathbf{y} \\
&= \sum_{\omega \in \Gamma} M^{\omega, \omega} p_\omega^\lambda(\mathbf{x}) \int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{y}) \left[\sum_{\omega' \in \Gamma \setminus \{\omega\}} G_{\lambda, \omega'}^{\text{atom}}(\mathbf{y}, \mathbf{z}) \Theta_{\omega'}(\mathbf{z}) \right] d\mathbf{y} \\
&\quad + \sum_{\omega \in \Gamma} M^{\omega, \omega} p_\omega^\lambda(\mathbf{x}) \int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{y}) G_\lambda^{\text{free}}(\mathbf{y} - \mathbf{z}) \Theta_{\text{free}}(\mathbf{z}) d\mathbf{y} \\
(10.69) \quad &\equiv \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda\right)_{2a}(\mathbf{x}, \mathbf{z}) + \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda\right)_{2b}(\mathbf{x}, \mathbf{z}).
\end{aligned}$$

Note the absence of the $\omega' = \omega$ term in the inner sum just above since the atomic Green's function, $G_{\lambda, \omega'}^{\text{atom}}$, projects onto the orthogonal complement of the function p_ω^λ .

We prove that the kernels $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda\right)_{2a}(\mathbf{x}, \mathbf{z})$ and $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda\right)_{2b}(\mathbf{x}, \mathbf{z})$, defined in (10.69) are both bounded in absolute value by $e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{z}|}$. We first recall the following relations and definitions:

$$\begin{aligned}
G_{\lambda, \omega}^{\text{atom}}(\mathbf{y}, \mathbf{z}) &= G_\lambda^{\text{atom}}(\mathbf{x} - \omega, \mathbf{y} - \omega), \\
(H_{\text{atom}}^\lambda - E_0^\lambda) G_\lambda^{\text{atom}}(\mathbf{x}, \mathbf{y}) &= \delta(\mathbf{x} - \mathbf{y}) - p_0^\lambda(\mathbf{x}) p_0^\lambda(\mathbf{y}) \\
\Theta_0(\mathbf{x}) &\equiv \begin{cases} 1, & |\mathbf{x}| \leq r_3 \\ 0, & |\mathbf{x}| \geq r_4 \end{cases}, \quad \text{and} \\
\Theta_\omega(\mathbf{x}) &= \Theta(\mathbf{x} - \omega), \quad \text{for } \omega \in \Gamma, \text{ and } \Theta_{\text{free}}(\mathbf{x}) = 1 - \sum_{\omega \in \Gamma} \Theta_\omega(\mathbf{x}).
\end{aligned}$$

Estimation of $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda\right)_{2a}(\mathbf{x}, \mathbf{z})$; see (10.69): Suppose first that $|\mathbf{z} - \omega'| \geq r_4$, for all $\omega' \in \Gamma \setminus \{\omega\}$. Then, \mathbf{z} is outside the support of $\Theta_{\omega'}(\mathbf{z})$ for all $\omega' \in \Gamma \setminus \{\omega\}$. and we have: $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda\right)_{2a}(\mathbf{x}, \mathbf{z}) \equiv 0$.

Suppose now that \mathbf{z} is such that $|\mathbf{z} - \omega'| \leq r_4$ for some $\omega' = \omega'_z \in \Gamma \setminus \{\omega\}$. Therefore, the bracketed expression in the definition of $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda\right)_{2a}$ (see (10.69)) is given by: $[\dots](\mathbf{y}, \mathbf{z}) = G_{\lambda, \omega'_z}^{\text{atom}}(\mathbf{y}, \mathbf{z}) \Theta_{\omega'_z}(\mathbf{z})$. Therefore, for $|\mathbf{z} - \omega'_z| \leq r_4$, we have

$$\begin{aligned}
\int p_\omega^\lambda(\mathbf{y}) [\dots](\mathbf{y}, \mathbf{z}) d\mathbf{y} &= \int p_\omega^\lambda(\mathbf{y}) G_{\lambda, \omega'_z}^{\text{atom}}(\mathbf{y}, \mathbf{z}) \Theta_{\omega'_z}(\mathbf{z}) d\mathbf{y} \\
&\leq \int_{|\mathbf{y}-\omega| \leq r_1} p_\omega^\lambda(\mathbf{y}) G_{\lambda, \omega'_z}^{\text{atom}}(\mathbf{y}, \mathbf{z}) d\mathbf{y} \\
(10.70) \quad &\quad + \int_{|\mathbf{y}-\omega| \geq r_1} p_\omega^\lambda(\mathbf{y}) G_{\lambda, \omega'_z}^{\text{atom}}(\mathbf{y}, \mathbf{z}) d\mathbf{y}.
\end{aligned}$$

We bound the latter two integrals individually by using the pointwise bounds on $p_\omega^\lambda(\mathbf{y}) = p_0^\lambda(\mathbf{y} - \omega)$ given in (3.5) and the pointwise bounds on $G_{\lambda, \omega'_z}^{\text{atom}}(\mathbf{y}, \mathbf{z}) = G_\lambda^{\text{atom}}(\mathbf{y} - \omega'_z, \mathbf{z} - \omega'_z)$ of Theorem 10.3.

With $|\mathbf{z} - \omega'_z| \leq r_4$, we first consider the integral over the set $|\mathbf{y} - \omega| \leq r_1$. For such \mathbf{y} , we have by (3.5): $|p_\omega^\lambda(\mathbf{y})| \lesssim \lambda^2$. Furthermore, note that $|\mathbf{y} - \omega'_z| \geq |\omega - \omega'_z| - |\mathbf{y} - \omega| \geq r_{\min} - r_1 > r_4$; see (10.42). Because $|\mathbf{y} - \omega'_z| > r_{\min} - r_1$, while $|\mathbf{z} - \omega'_z| < r_4$, it follows from (10.20) (part 3 of Theorem 10.3) that $|G_{\lambda, \omega'_z}^{\text{atom}}(\mathbf{y}, \mathbf{z})| \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{y}-\mathbf{z}|}$. The first integral in (10.70) therefore satisfies

$$\int_{|\mathbf{y}-\omega| \leq r_1} p_\omega^\lambda(\mathbf{y}) G_{\lambda, \omega'_z}^{\text{atom}}(\mathbf{y}, \mathbf{z}) d\mathbf{y} \lesssim \lambda^2 \int_{|\mathbf{y}-\omega| \leq r_1} e^{-c\lambda} e^{-c\lambda|\mathbf{y}-\mathbf{z}|} d\mathbf{y} \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{z}-\omega|}.$$

Next, with $|\mathbf{z} - \omega'_z| \leq r_4$, we consider the integral over the set $|\mathbf{y} - \omega| \geq r_1$. On this set, we have $|p_\omega^\lambda(\mathbf{y})| \lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{y}-\omega|}$ and, by the bounds of Theorem 10.3:

$$\begin{aligned} & \int_{|\mathbf{y}-\omega| \geq r_1} p_\omega^\lambda(\mathbf{y}) G_{\lambda, \omega'_z}^{\text{atom}}(\mathbf{y}, \mathbf{z}) d\mathbf{y} \\ & \lesssim \int_{|\mathbf{y}-\omega| \geq r_1} e^{-c'\lambda} e^{-c'\lambda|\mathbf{y}-\omega|} [(c_0 |\log|\mathbf{z} - \mathbf{y}|| + \lambda^4) \mathbf{1}_{|\mathbf{y}-\mathbf{z}| \leq R} + e^{-c\lambda} e^{-c\lambda|\mathbf{z}-\mathbf{y}|}] d\mathbf{y} \\ & \lesssim e^{-\tilde{c}\lambda} e^{-\tilde{c}\lambda|\mathbf{z}-\omega|}. \end{aligned}$$

Therefore, the integral expression in the definition of $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_{2a}(\mathbf{x}, \mathbf{z})$ satisfies the bound:

$$\int p_\omega^\lambda(\mathbf{y}) [\dots](\mathbf{y}, \mathbf{z}) d\mathbf{y} = \int p_\omega^\lambda(\mathbf{y}) G_{\lambda, \omega'_z}^{\text{atom}}(\mathbf{y}, \mathbf{z}) \Theta_{\omega'_z}(\mathbf{z}) d\mathbf{y} \lesssim e^{-\tilde{c}\lambda} e^{-\tilde{c}\lambda|\mathbf{z}-\omega|} = e^{-\tilde{c}\lambda} e^{-\frac{1}{2}\tilde{c}\lambda|\mathbf{z}-\omega|} e^{-\frac{1}{2}\tilde{c}\lambda|\mathbf{z}-\omega|}.$$

We next multiply this estimate by $p_\omega^\lambda(\mathbf{x})$ and once again use the pointwise bound (3.5):

$$\begin{aligned} p_\omega^\lambda(\mathbf{x}) \int p_\omega^\lambda(\mathbf{y}) [\dots](\mathbf{y}, \mathbf{z}) d\mathbf{y} & \lesssim (\lambda^2 \mathbf{1}_{|\mathbf{x}-\omega| \leq R} + e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\omega|}) e^{-\tilde{c}\lambda} e^{-\frac{1}{2}\tilde{c}\lambda|\mathbf{z}-\omega|} e^{-\frac{1}{2}\tilde{c}\lambda|\mathbf{z}-\omega|} \\ & \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{z}|} e^{-\frac{1}{2}\tilde{c}\lambda|\mathbf{z}-\omega|}. \end{aligned}$$

Finally, we multiply the previous bound by $M^{\omega, \omega} = 1 + \mathcal{O}(e^{-c\lambda})$ (see (10.58)) and sum over all $\omega \in \Gamma$ to obtain:

$$\begin{aligned} \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_{2a}(\mathbf{x}, \mathbf{z}) & = \sum_{\omega \in \Gamma} p_\omega^\lambda(\mathbf{x}) \int p_\omega^\lambda(\mathbf{y}) [\dots](\mathbf{y}, \mathbf{z}) d\mathbf{y} \\ & \lesssim (1 + \mathcal{O}(e^{-c\lambda})) e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{z}|} \sum_{\omega \in \Gamma} e^{-\frac{1}{2}\tilde{c}\lambda|\mathbf{z}-\omega|} \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{z}|}. \end{aligned}$$

Therefore, the contribution to $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_2(\mathbf{x}, \mathbf{z})$ from $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_{2a}(\mathbf{x}, \mathbf{z})$ is an error kernel.

Estimation of $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_{2b}(\mathbf{x}, \mathbf{z})$; see (10.69): From the expression (10.69) we need only consider $\mathbf{z} \in \text{supp}(\Theta_{\text{free}})$, that is \mathbf{z} bounded away from the all sites $\omega \in \Gamma$; in particular,

$|\mathbf{z} - \omega| \geq r_3$ for all $\omega \in \Gamma$. By (3.5) and (10.5):

$$\begin{aligned} & p_\omega^\lambda(\mathbf{y}) G_\lambda^{\text{free}}(\mathbf{y} - \mathbf{z}) \Theta_{\text{free}}(\mathbf{z}) \\ & \lesssim \left(\lambda^2 \mathbf{1}_{|\mathbf{y} - \omega| \leq r_1} + e^{-c\lambda} e^{-c\lambda|\mathbf{y} - \omega|} \right) \cdot e^{-c\lambda|\mathbf{y} - \mathbf{z}|} \cdot \left(1 + \left| \log \lambda |\mathbf{y} - \mathbf{z}| \right| \right) \Theta_{\text{free}}(\mathbf{z}) \\ & \lesssim \left(e^{-c'\lambda|\mathbf{y} - \mathbf{z}|} \mathbf{1}_{|\mathbf{y} - \omega| \leq r_1} + e^{-c\lambda|\mathbf{y} - \mathbf{z}|} e^{-c\lambda|\mathbf{y} - \omega|} \mathbf{1}_{|\mathbf{y} - \omega| \geq r_1} \right) \cdot \left(1 + \left| \log \lambda |\mathbf{y} - \mathbf{z}| \right| \right) \Theta_{\text{free}}(\mathbf{z}) . \end{aligned}$$

Integrating over \mathbb{R}^2 with respect to \mathbf{y} , we find that

$$\int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{y}) G_\lambda^{\text{free}}(\mathbf{y} - \mathbf{z}) \Theta_{\text{free}}(\mathbf{z}) d\mathbf{y} \lesssim e^{-c\lambda|\mathbf{z} - \omega|} \Theta_{\text{free}}(\mathbf{z}) .$$

Now multiply this bound by $M^{\omega, \omega} p_\omega^\lambda(\mathbf{x})$ and apply the pointwise bound for $p_\omega^\lambda(\mathbf{x})$, implied by (3.5), and the expansion $M^{\omega, \omega} = 1 + \mathcal{O}(e^{-c\lambda})$ of (10.58), to obtain

$$\begin{aligned} & M^{\omega, \omega} p_\omega^\lambda(\mathbf{x}) \int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{y}) G_\lambda^{\text{free}}(\mathbf{y} - \mathbf{z}) \Theta_{\text{free}}(\mathbf{z}) d\mathbf{y} \\ & \lesssim \left(\lambda^2 \mathbf{1}_{|\mathbf{x} - \omega| \leq r_1} e^{-\frac{1}{4}c\lambda|\mathbf{z} - \omega|} e^{-\frac{1}{4}c\lambda|\mathbf{z} - \omega|} + e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \omega|} e^{-\frac{1}{2}c\lambda|\mathbf{z} - \omega|} \right) \Theta_{\text{free}}(\mathbf{z}) e^{-\frac{1}{2}c\lambda|\mathbf{z} - \omega|} \\ & \lesssim \left(\mathbf{1}_{|\mathbf{x} - \omega| \leq r_1} e^{-\tilde{c}\lambda|\mathbf{x} - \omega|} e^{-\tilde{c}\lambda|\mathbf{z} - \omega|} + e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \omega|} e^{-\frac{1}{2}c\lambda|\mathbf{z} - \omega|} \right) \Theta_{\text{free}}(\mathbf{z}) e^{-\frac{1}{2}c\lambda|\mathbf{z} - \omega|} \\ & \lesssim e^{-c'\lambda|\mathbf{x} - \mathbf{z}|} \Theta_{\text{free}}(\mathbf{z}) e^{-\frac{1}{2}c\lambda|\mathbf{z} - \omega|} . \end{aligned}$$

Summing over $\omega \in \Gamma$ and using that on the support of $\Theta_{\text{free}}(\mathbf{z})$, \mathbf{z} is uniformly bounded away from Γ , we have that

$$\sum_{\omega \in \Gamma} M^{\omega, \omega} p_\omega^\lambda(\mathbf{x}) \int_{\mathbb{R}^2} p_\omega^\lambda(\mathbf{y}) G_\lambda^{\text{free}}(\mathbf{y} - \mathbf{z}) \Theta_{\text{free}}(\mathbf{z}) d\mathbf{y} \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{z}|} .$$

Hence, the contribution to $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_2(\mathbf{x}, \mathbf{z})$ of $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_{2b}(\mathbf{x}, \mathbf{z})$ is also an error kernel. Therefore, $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_2(\mathbf{x}, \mathbf{z})$ is an error kernel, and since we have already verified that $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)_1(\mathbf{x}, \mathbf{z})$ is an error kernel, we conclude that $\left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)(\mathbf{x}, \mathbf{z})$ is an error kernel. Furthermore, it is straightforward to show by arguments similar to those above that $H_\Gamma^\lambda \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right)(\mathbf{x}, \mathbf{z})$ is an error kernel, where H_Γ^λ is defined in (10.41). Indeed, we just replace $p_\omega^\lambda(\mathbf{x})$ by $H_\Gamma^\lambda p_\omega^\lambda(\mathbf{x})$ in the previous discussion. Note that $H_\Gamma^\lambda p_\omega^\lambda(\mathbf{x}) = \lambda^2 \sum_{\omega' \in \Gamma \setminus \{\omega\}} V_0(\mathbf{x} - \omega') p_0^\lambda(\mathbf{x} - \omega)$ and therefore $|H_\Gamma^\lambda p_\omega^\lambda(\mathbf{x})| \lesssim \lambda^2 \|V_0\|_{L^\infty} p_\omega^\lambda(\mathbf{x})$. Hence, the estimates lose at worst one power of λ^2 , which can be absorbed by our exponentials $e^{-c\lambda}$. This completes the proof of Proposition 10.12.

From (10.62), Proposition 10.9 and Proposition 10.12 we have

$$\begin{aligned} (10.71) \quad f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda \right) K_1^\lambda[f](\mathbf{x}) \\ &+ \sum_{\omega \in \Gamma} \langle \Theta_\omega p_\omega^\lambda, f \rangle_{L^2(\mathbb{R}^2)} p_\omega^\lambda(\mathbf{x}) + \mathcal{E}_1^\lambda[f](\mathbf{x}) , \end{aligned}$$

where

$$(10.72) \quad K_1^\lambda \equiv K_0^\lambda - \tilde{\Pi}_\Gamma^\lambda K_0^\lambda = \Pi_\Gamma^\lambda K_0^\lambda \quad \text{is a main kernel ,}$$

$$(10.73) \quad \langle p_\omega^\lambda, K_1^\lambda[f] \rangle = 0, \quad \text{for all } \omega \in \Gamma,$$

and

$$(10.74) \quad \mathcal{E}_1^\lambda = -\mathcal{E}_0^\lambda + \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda \right) \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right) = -\mathcal{E}_0^\lambda + H_\Gamma^\lambda \left(\tilde{\Pi}_\Gamma^\lambda K_0^\lambda \right).$$

is derived from an error kernel.

Now let $|\Omega| < e^{-\hat{c}\lambda}$, where \hat{c} is a constant that was introduced in Remark 3.1, and thus $(\rho_\lambda)^{-1}|\Omega| \leq e^{-(\hat{c}-c)\lambda} \rightarrow 0$, as $\lambda \rightarrow \infty$. Then, from (10.71) we have

$$(10.75) \quad \begin{aligned} f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda - \Omega \right) K_1^\lambda[f](\mathbf{x}) \\ &+ \sum_{\omega \in \Gamma} \langle \Theta_\omega p_\omega^\lambda, f \rangle_{L^2(\mathbb{R}^2)} p_\omega^\lambda(\mathbf{x}) + \left(\mathcal{E}_1^\lambda + \Omega K_1^\lambda \right) [f](\mathbf{x}) \end{aligned}$$

and hence

$$(10.76) \quad \begin{aligned} \left(I - \left(\mathcal{E}_1^\lambda + \Omega K_1^\lambda \right) \right) f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda - \Omega \right) K_1^\lambda[f](\mathbf{x}) \\ &+ \sum_{\omega \in \Gamma} \langle \Theta_\omega p_\omega^\lambda, f \rangle_{L^2(\mathbb{R}^2)} p_\omega^\lambda(\mathbf{x}). \end{aligned}$$

For λ large, the operator $\mathcal{E}_1^\lambda + \Omega K_1^\lambda$ has small norm as a bounded operator on $L^2(\mathbb{R}^2)$. Hence, $I - (\mathcal{E}_1^\lambda + \Omega K_1^\lambda)$ is invertible. Applying (10.75) to $\tilde{f} = \left(I - (\mathcal{E}_1^\lambda + \Omega K_1^\lambda) \right)^{-1} f$ yields

$$(10.77) \quad \begin{aligned} f(\mathbf{x}) &= \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda - \Omega \right) \left(K_1^\lambda \left(I - \left(\mathcal{E}_1^\lambda + \Omega K_1^\lambda \right) \right)^{-1} \right) [f](\mathbf{x}) \\ &+ \sum_{\omega \in \Gamma} \langle \Theta_\omega p_\omega^\lambda, \tilde{f} \rangle_{L^2(\mathbb{R}^2)} p_\omega^\lambda(\mathbf{x}). \end{aligned}$$

From (10.77) we see that for all $f \in L^2(\mathbb{R}^2)$ and $|\Omega| \lesssim e^{-\hat{c}\lambda}$

$$(10.78) \quad \begin{aligned} \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda - \Omega \right) \left(K_1^\lambda \left(I - \left(\mathcal{E}_1^\lambda + \Omega K_1^\lambda \right) \right)^{-1} \right) f &= f \\ \text{modulo the span of } \{p_\omega^\lambda : \omega \in \Gamma\}. \end{aligned}$$

Here, K_1^λ , defined in (10.63), is derived from a main kernel, \mathcal{E}_1^λ is derived from an error kernel.

Proposition 10.13. *For λ sufficiently large and Ω such that $|\Omega| \lesssim e^{-\hat{c}\lambda}$,*

$$(10.79) \quad K_2^\lambda \equiv K_1^\lambda \left(I - \left(\mathcal{E}_1^\lambda + \Omega K_1^\lambda \right) \right)^{-1} \equiv K_1^\lambda + \mathcal{E}_2^\lambda.$$

Here, K_1^λ is derived from a main kernel, \mathcal{E}_2^λ from an error kernel and therefore K_2^λ is derived from a main kernel. Moreover, for all $f \in L^2(\mathbb{R}^2)$:

$$(10.80) \quad \begin{aligned} \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda - \Omega \right) K_2^\lambda f &= f, \\ \text{modulo the span of } \{p_\omega^\lambda : \omega \in \Gamma\}, \end{aligned}$$

$$(10.81) \quad K_2^\lambda[f] \perp \text{span}\{p_\omega^\lambda : \omega \in \Gamma\}.$$

Proof. Proof of Proposition 10.13: Set $A = \Omega K_1^\lambda + \mathcal{E}_1^\lambda$, where λ is taken sufficiently large. First note that by Lemma 10.7 that the operator A^2 is derived from an error kernel. As an operator on $L^2(\mathbb{R}^2)$ we have $(I - A)^{-1} = (I + A)(I - A^2)^{-1} = (I + A)(I + A_1)$, where A_1 is an error kernel, again by Lemma 10.7. Therefore, $(I - A)^{-1} = I + A + A_2 = I + \Omega K_1^\lambda + A_3$, where A_j ($j = 2, 3$) arise from error kernels. Another application of Lemma 10.7 completes the proof that \mathcal{E}_2^λ is derived from an error kernel. That (10.80)-(10.81) hold follows from (10.78) and (10.73). \square

Recall the subspace \mathcal{X}_Γ , the orthogonal complement of $\text{span}\{p_\omega^\lambda : \omega \in \Gamma\}$:

$$(10.82) \quad \mathcal{X}_\Gamma \equiv \text{span}\{p_\omega^\lambda : \omega \in \Gamma\}^\perp = \left\{ f \in L^2(\mathbb{R}^2) : \langle p_\omega^\lambda, f \rangle_{L^2(\mathbb{R}^2)} = 0, \omega \in \Gamma \right\},$$

and the orthogonal projections: $\Pi_\Gamma^\lambda : L^2(\mathbb{R}^2) \rightarrow \mathcal{X}_\Gamma$ and $\tilde{\Pi}_\Gamma^\lambda : L^2(\mathbb{R}^2) \rightarrow \text{span}\{p_\omega^\lambda : \omega \in \Gamma\}$; see (10.55). We now write

$$K_2^\lambda = K_3^\lambda + \mathcal{E}_3^\lambda$$

where

$$(10.83) \quad K_3^\lambda \equiv K_2^\lambda \Pi_\Gamma^\lambda, \quad \text{and} \quad \mathcal{E}_3^\lambda \equiv K_2^\lambda \tilde{\Pi}_\Gamma^\lambda$$

Note that

$$K_3^\lambda[f] = 0 \text{ in } L^2(\mathbb{R}^2) \text{ if } f \in \text{span}\{p_\omega^\lambda : \omega \in \Gamma\},$$

and by Proposition 10.13:

$$(10.84) \quad \left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda - \Omega \right) \mathcal{E}_3^\lambda \in \text{span}\{p_\omega^\lambda : \omega \in \Gamma\}.$$

Hence, for all $f \in L^2(\mathbb{R}^2)$:

$$\left(-\Delta_{\mathbf{x}} + V_\Gamma^\lambda(\mathbf{x}) - E_0^\lambda - \Omega \right) K_3^\lambda f = f \quad \text{modulo the span of } \{p_\omega^\lambda : \omega \in \Gamma\}.$$

We therefore have

Proposition 10.14. *Let $|\Omega| \leq e^{-c\lambda}$ with λ chosen sufficiently large. Then, the operator $\Pi_\Gamma^\lambda(H_\Gamma^\lambda - E_0^\lambda - \Omega) = \Pi_\Gamma^\lambda(-\Delta + V_\Gamma^\lambda - E_0^\lambda - \Omega)$ is invertible on \mathcal{X}_Γ , the orthogonal complement of $\text{span}\{p_\omega^\lambda : \omega \in \Gamma\}$. Its inverse is given by $K_3^\lambda|_{\mathcal{X}_\Gamma}$ and we write*

$$\mathcal{K}_\Gamma^\lambda(\Omega)|_{\mathcal{X}_\Gamma} \equiv K_3^\lambda|_{\mathcal{X}_\Gamma} : \mathcal{X}_\Gamma \rightarrow \mathcal{X}_\Gamma.$$

The following proposition characterizes the operator kernel we seek:

Proposition 10.15. *Let $|\Omega| \lesssim e^{-\varepsilon\lambda}$ with λ chosen sufficiently large. Then, $\mathcal{K}_\Gamma^\lambda(\Omega)$ defined in Proposition 10.14 satisfies the following properties:*

(1)

$$(10.85) \quad \mathcal{K}_\Gamma^\lambda(\Omega)[f] = 0 \text{ in } L^2(\mathbb{R}^2) \text{ if } f \in \text{span}\{p_\omega^\lambda : \omega \in \Gamma\}.$$

(2)

$$(10.86) \quad \mathcal{K}_\Gamma^\lambda(\Omega)[f] \perp \text{span}\{p_\omega^\lambda : \omega \in \Gamma\} \text{ in } L^2(\mathbb{R}^2).$$

(3)

$$(10.87) \quad \Pi_{\Gamma}^{\lambda} (-\Delta + V_{\Gamma}^{\lambda} - E_0^{\lambda} - \Omega) \mathcal{K}_{\Gamma}^{\lambda}(\Omega)[f] = f .$$

(4) *The operator $\mathcal{K}_{\Gamma}^{\lambda}(\Omega)$ is derived from a kernel:*

$$(10.88) \quad \mathcal{K}_{\Gamma}^{\lambda}(\Omega)[f](\mathbf{x}) = \int_{\mathbb{R}^2} \mathcal{K}_{\Gamma}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega) f(\mathbf{y}) d\mathbf{y} \quad \text{for all } f \in L^2(\mathbb{R}^2), \quad \text{where}$$

$$(10.89) \quad \left| \mathcal{K}_{\Gamma}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega) \right| \leq C \left[\left| \log |\mathbf{x} - \mathbf{y}| \right| + \lambda^6 \right] \mathbf{1}_{|\mathbf{x} - \mathbf{y}| \leq C} + e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|}$$

for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$.

The only assertion in Proposition 10.15 that requires proof is part (4). Recall that $\mathcal{K}_{\Gamma}^{\lambda}(\Omega) = K_3^{\lambda} = K_2^{\lambda} \Pi_{\Gamma}^{\lambda} = K_2^{\lambda} - K_2^{\lambda} \tilde{\Pi}_{\Gamma}^{\lambda}$. Since K_2^{λ} is derived from a main kernel, it suffices to study the kernel of $K_2^{\lambda} \tilde{\Pi}_{\Gamma}^{\lambda}$. We begin with a bound on the kernel of $\tilde{\Pi}_{\Gamma}^{\lambda}$, which we derive using Lemma 10.11. The kernel of $\tilde{\Pi}_{\Gamma}^{\lambda}$, $K_{\tilde{\Pi}}^{\lambda}(\mathbf{x}, \mathbf{y})$, is given by (see (10.57)):

$$(10.90) \quad K_{\tilde{\Pi}}^{\lambda}(\mathbf{x}, \mathbf{y}) = \sum_{\omega, \omega'} M^{\omega, \omega'} p_{\omega}^{\lambda}(\mathbf{x}) p_{\omega'}^{\lambda}(\mathbf{y}),$$

and we have

$$(10.91) \quad \tilde{\Pi}_{\Gamma}^{\lambda}[g](\mathbf{x}) = \int_{\mathbb{R}^2} K_{\tilde{\Pi}}^{\lambda}(\mathbf{x}, \mathbf{y}) g(\mathbf{y}) d\mathbf{y}.$$

Our goal is to bound

$$(10.92) \quad \mathcal{K}_{\Gamma}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega) = K_2^{\lambda}(\mathbf{x}, \mathbf{y}) - \left(K_2^{\lambda} \circ K_{\tilde{\Pi}}^{\lambda} \right)(\mathbf{x}, \mathbf{y}) = K_2^{\lambda}(\mathbf{x}, \mathbf{y}) - \int_{\mathbb{R}^2} K_2^{\lambda}(\mathbf{x}, \mathbf{z}) K_{\tilde{\Pi}}^{\lambda}(\mathbf{z}, \mathbf{y}) d\mathbf{y}$$

Note that

$$(10.93) \quad K_{\tilde{\Pi}}^{\lambda}(\mathbf{x}, \mathbf{y}) = \sum_{\omega} p_{\omega}^{\lambda}(\mathbf{x}) p_{\omega}^{\lambda}(\mathbf{y}) + \sum_{\omega, \omega'} \left[M^{\omega, \omega'} - \delta_{\omega, \omega'} \right] p_{\omega}^{\lambda}(\mathbf{x}) p_{\omega'}^{\lambda}(\mathbf{y}).$$

Recall from (10.58) that $\left| M^{\omega, \omega'} - \delta_{\omega, \omega'} \right| \lesssim e^{-c\lambda} e^{-c\lambda|\omega - \omega'|}$. Also, from the pointwise bounds, (3.5), on p_0^{λ} we have:

$$\left| p_{\omega}^{\lambda}(\mathbf{x}) \right| \lesssim \lambda \mathbf{1}_{|\mathbf{x} - \omega| \leq R} + e^{-c\lambda|\mathbf{x} - \omega|}, \quad \left| p_{\omega'}^{\lambda}(\mathbf{y}) \right| \lesssim \lambda \mathbf{1}_{|\mathbf{y} - \omega| \leq R} + e^{-c\lambda|\mathbf{y} - \omega'|},$$

which it follows that

$$\left| \sum_{\omega} p_{\omega}^{\lambda}(\mathbf{x}) p_{\omega}^{\lambda}(\mathbf{y}) \right| \lesssim \lambda^2 \mathbf{1}_{|\mathbf{y} - \omega| \leq 2R} + e^{-c'|\mathbf{x} - \mathbf{y}|},$$

$$\left| \sum_{\omega, \omega'} \left[M^{\omega, \omega'} - \delta_{\omega, \omega'} \right] p_{\omega}^{\lambda}(\mathbf{x}) p_{\omega'}^{\lambda}(\mathbf{y}) \right| \lesssim e^{-c'\lambda} \left[\mathbf{1}_{|\mathbf{x} - \mathbf{y}| \leq 2R} + e^{-c'\lambda|\mathbf{x} - \mathbf{y}|} \right]$$

Substitution into (10.93), we obtain

$$(10.94) \quad \left| K_{\tilde{\Pi}}^{\lambda}(\mathbf{x}, \mathbf{y}) \right| \lesssim \mathbf{1}_{|\mathbf{x} - \mathbf{y}| \leq 2R} \lambda^2 + e^{-c'\lambda|\mathbf{x} - \mathbf{y}|} .$$

Now since $K_2(\mathbf{x}, \mathbf{y}; \Omega)$ is a main kernel we have

$$(10.95) \quad \left| K_2(\mathbf{x}, \mathbf{y}; \Omega) \right| \lesssim \left[\lambda^4 + \left| \log |\mathbf{x} - \mathbf{y}| \right| \right] \mathbf{1}_{|\mathbf{x} - \mathbf{y}| \leq R} + e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|}$$

Inserting the bounds (10.94) and (10.95) into (10.92) we find that $\mathcal{K}_\Gamma^\lambda(\mathbf{x}, \mathbf{y}; \Omega)$ satisfies the bound:

$$(10.96) \quad |\mathcal{K}_\Gamma(\mathbf{x}, \mathbf{y}; \Omega)| \lesssim \left[\lambda^6 + \left| \log |\mathbf{x} - \mathbf{y}| \right| \right] \mathbf{1}_{|\mathbf{x} - \mathbf{y}| \leq 3R} + e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|}$$

The proof is complete of Proposition 10.15 is complete.

10.5. $\mathcal{K}_\Gamma^\lambda(\Omega)$ for the case where Γ , the set of nuclei, is translation invariant. We now suppose that our discrete set of nuclei, Γ , is translation invariant by a vector $\mathbf{v}_2 \in \mathbb{R}^2$. Of course, we have in mind, $\Gamma = \mathbb{H}_\sharp$, the zigzag truncation of \mathbb{H} ; see (1.11). But our arguments would apply to other *rational truncations* of \mathbb{H} , for example along an armchair edge. For the particular choice $\Gamma = \mathbb{H}_\sharp$, we have $V_\Gamma(\mathbf{x}) = V_\sharp(\mathbf{x})$ and

$$H_\Gamma^\lambda = H_\sharp^\lambda \equiv -\Delta + \lambda^2 V_\sharp(\mathbf{x}) - E_0^\lambda.$$

As commented upon in Remark 10.10, all our constructions of integral operators and kernels respect that translation invariance. Thus, at each stage our integral kernels $A(\mathbf{x}, \mathbf{y})$ satisfy: $A(\mathbf{x} + \mathbf{v}_2, \mathbf{y} + \mathbf{v}_2) = A(\mathbf{x}, \mathbf{y})$. It follows that

$$(10.97) \quad \mathcal{K}_\Gamma^\lambda(\mathbf{x} + \mathbf{v}_2, \mathbf{y} + \mathbf{v}_2) = \mathcal{K}_\Gamma^\lambda(\mathbf{x}, \mathbf{y}) \quad \text{for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^2.$$

10.5.1. $\mathcal{K}_\Gamma^\lambda$ as a bounded operator acting on $L_{k_\parallel}^2(\Sigma)$.

Let Γ be invariant under translation by \mathbf{v}_2 . We recall the setting discussed earlier. Associated with this translation invariance is a parallel quasi-momentum, $k_\parallel \in [0, 2\pi)$. We define the cylinder $\Sigma = \mathbb{R}^2 / \mathbb{R}\mathbf{v}_2$ and let Ω_Σ denote a fundamental domain for Σ . The space $L^2(\Sigma)$ consists of functions f such that $f(\mathbf{x} + \mathbf{v}_2) = f(\mathbf{x})$ for almost all $\mathbf{x} \in \mathbb{R}^2$ and such that $\|f\|_{L^2(\Sigma)} \equiv \left(\int_{\Omega_\Sigma} |f(\mathbf{x})|^2 d\mathbf{x} \right)^{\frac{1}{2}} < \infty$. The space $L_{k_\parallel}^2(\Sigma)$ consists of functions f such that $g(\mathbf{x}) \equiv f(\mathbf{x}) e^{-i\frac{k_\parallel}{2\pi} \mathbf{v}_2 \cdot \mathbf{x}}$ satisfies $g(\mathbf{x} + \mathbf{v}_2) = g(\mathbf{x})$ almost everywhere in \mathbf{x} and $g \in L^2(\Sigma)$.

We now show that $\mathcal{K}_\Gamma^\lambda$ also gives rise to a bounded operator $L_{k_\parallel}^2(\Sigma)$. For any $f \in L_{k_\parallel}^2(\Sigma)$, we define

$$(10.98) \quad \mathcal{K}_\Gamma^\lambda[f](\mathbf{x}) = \int_{\mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y}.$$

Similarly, Π_Γ^λ may be defined on $L_{k_\parallel}^2(\Sigma)$ using Lemma 10.11.

By our bounds on $\mathcal{K}_\Gamma^\lambda(\mathbf{x}, \mathbf{y})$, $\mathcal{K}_\Gamma^\lambda[f]$ is well-defined for all $f \in L_{k_\parallel}^2(\Sigma)$. Using (10.97) and our assumption that $f(\mathbf{x} + \mathbf{v}_2) = e^{ik_\parallel} f(\mathbf{x})$ almost everywhere, we obtain by change of variables:

$$(10.99) \quad \begin{aligned} \mathcal{K}_\Gamma^\lambda[f](\mathbf{x} + \mathbf{v}_2) &= \int_{\mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\mathbf{x} + \mathbf{v}_2, \mathbf{y}) f(\mathbf{y}) d\mathbf{y} = \int_{\mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\mathbf{x} + \mathbf{v}_2, \mathbf{y} + \mathbf{v}_2) f(\mathbf{y} + \mathbf{v}_2) d\mathbf{y} \\ &= \int_{\mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\mathbf{x}, \mathbf{y}) f(\mathbf{y} + \mathbf{v}_2) d\mathbf{y} = e^{ik_\parallel} \int_{\mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y} = e^{ik_\parallel} \mathcal{K}_\Gamma^\lambda[f](\mathbf{x}). \end{aligned}$$

Hence, $e^{-i\frac{k_\parallel}{2\pi} \mathbf{v}_2 \cdot \mathbf{x}} \mathcal{K}_\Gamma^\lambda[f](\mathbf{x})$ is a function defined on the cylinder Σ . Similarly, one shows easily that Π_Γ^λ maps $L^2(\Sigma)$ into itself. Furthermore, we have

$$(10.100) \quad \left(\Pi_\Gamma^\lambda (H_\Gamma^\lambda - E_D^0 - \Omega) \Pi_\Gamma^\lambda \right) \circ \mathcal{K}_\Gamma^\lambda f = \Pi_\Gamma^\lambda f, \quad \mathcal{K}_\Gamma^\lambda f \in L_{k_\parallel}^2(\Sigma)$$

thanks to Proposition 10.15. That $e^{-i\frac{\mathfrak{K}_2 \cdot \mathbf{x}}{2\pi} k_{\parallel}} \mathcal{K}_{\Gamma}^{\lambda} f \in L^2(\Sigma)$ is a consequence of the kernel bounds on $\mathcal{K}_{\Gamma}^{\lambda}(\mathbf{x}, \mathbf{y})$ and Young's inequality. Therefore, we have

Proposition 10.16. *Let $|\Omega| \leq e^{-\hat{c}\lambda}$ with λ chosen sufficiently large. Let the discrete set Γ be invariant under translation by the vector \mathbf{v}_2 . Then, the kernel $\mathcal{K}_{\Gamma}^{\lambda}(\Omega)(\mathbf{x}, \mathbf{y})$, defined in Proposition 10.15 and (10.98), gives rise to a bounded operator on $L^2_{k_{\parallel}}(\Sigma)$. Furthermore, the operator*

$$(10.101) \quad \mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel}) \equiv e^{-i\frac{\mathfrak{K}_2 \cdot \mathbf{x}}{2\pi} k_{\parallel}} \mathcal{K}_{\Gamma}^{\lambda}(\Omega) e^{i\frac{\mathfrak{K}_2 \cdot \mathbf{x}}{2\pi} k_{\parallel}}$$

is a bounded operator on $L^2(\Sigma)$.

10.5.2. *The operator $\mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel})$ acting on periodized sums.*

Let Γ be invariant under translates by integer multiples of \mathbf{v}_2 . We are interested in $\mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel}) : L^2(\Sigma) \rightarrow L^2(\Sigma)$ (see (10.101)) applied to a sum over all \mathbf{v}_2 -integer-translates of

$$(10.102) \quad p_{k_{\parallel}, \omega}^{\lambda}(\mathbf{x}) = e^{i\frac{k_{\parallel}}{2\pi} \mathfrak{K}_2 \cdot (\mathbf{x} - \omega)} p_0(\mathbf{x} - \omega).$$

For $\omega \in \Gamma$, let $[\omega]$ denote the equivalence class of all translates of ω by integer multiples of \mathbf{v}_2 . The set of such equivalence classes is

$$(10.103) \quad \Lambda_{\Sigma} \equiv \{[\omega] : \omega \in \Gamma\}$$

For any $[\omega] \in \Lambda_{\Sigma}$ we set

$$(10.104) \quad p_{k_{\parallel}, [\omega]}^{\lambda}(\mathbf{x}) \equiv \sum_{m \in \mathbb{Z}} p_{k_{\parallel}, \omega}^{\lambda}(\mathbf{x} + m\mathbf{v}_2).$$

Our estimates on $p_{\omega}^{\lambda} \in L^2(\mathbb{R}^2)$ imply that $p_{k_{\parallel}, [\omega]}^{\lambda} \in L^2(\Sigma)$, and by our discussion of the previous subsection $\mathcal{K}_{\Gamma}^{\lambda}[p_{k_{\parallel}, [\omega]}^{\lambda}] \in L^2(\Sigma)$. Furthermore, we have

Proposition 10.17. *Let $|\Omega| \leq e^{-\hat{c}\lambda}$ with λ chosen sufficiently large.*

- (1) $\mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel})[f] = 0$ in $L^2(\Sigma)$ for all $f \in \text{span}\{p_{k_{\parallel}, [\omega]}^{\lambda} : \omega \in \Gamma\}$.
- (2) For all $\omega \in \Gamma$ and $f \in L^2(\Sigma)$, we have $\left\langle \mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel})[f], p_{k_{\parallel}, [\omega]}^{\lambda} \right\rangle_{L^2(\Sigma)} = 0$.

Proof of claim (1) of Proposition 10.17: We claim in fact for any $\omega \in \Gamma$, and for any $\mathbf{x} \in \mathbb{R}^2$, we have $\mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel})[p_{k_{\parallel}, [\omega]}^{\lambda}](\mathbf{x}) = 0$. Indeed,

$$\begin{aligned} \mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel})[p_{k_{\parallel}, [\omega]}^{\lambda}](\mathbf{x}) &= \int_{\mathbb{R}^2} \mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel})(\mathbf{x}, \mathbf{y}) \sum_{m \in \mathbb{Z}} p_{k_{\parallel}, \omega - m\mathbf{v}_2}^{\lambda}(\mathbf{y}) \, d\mathbf{y} \\ &= \lim_{N \rightarrow \infty} \int_{\mathbb{R}^2} \mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel})(\mathbf{x}, \mathbf{y}) \sum_{|m| \leq N} p_{k_{\parallel}, \omega - m\mathbf{v}_2}^{\lambda}(\mathbf{y}) \, d\mathbf{y} \\ &= \lim_{N \rightarrow \infty} \sum_{|m| \leq N} \int_{\mathbb{R}^2} \mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel})(\mathbf{x}, \mathbf{y}) p_{k_{\parallel}, \omega - m\mathbf{v}_2}^{\lambda}(\mathbf{y}) \, d\mathbf{y} \\ &= \lim_{N \rightarrow \infty} \sum_{|m| \leq N} \mathcal{K}_{\Gamma}^{\lambda}(\Omega, k_{\parallel})[p_{k_{\parallel}, \omega - m\mathbf{v}_2}^{\lambda}](\mathbf{x}) = 0, \end{aligned}$$

by property (10.85) of Proposition 10.15. These formal manipulations are easily justified thanks to our estimates on $\mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)(\mathbf{x}, \mathbf{y})$ and $p_\omega^\lambda(\mathbf{x})$. This completes the proof of the first claim of Proposition 10.17.

Proof of claim (2) of Proposition 10.17: Let $\omega \in \Gamma$ and $f \in L^2(\Sigma)$. Then,

$$\begin{aligned} \left\langle \mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)[f], p_{k_\parallel, [\omega]}^\lambda \right\rangle_{L^2(\Sigma)} &= \int_{\mathbf{x} \in \Omega_\Sigma} \sum_{m \in \mathbb{Z}} p_{k_\parallel, \omega}^\lambda(\mathbf{x} + m\mathbf{v}_2) \cdot \int_{\mathbf{y} \in \mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) \, d\mathbf{y} \, d\mathbf{x} \\ &= \sum_{m \in \mathbb{Z}} \int_{\mathbf{x} \in \Omega_\Sigma} p_{k_\parallel, \omega}^\lambda(\mathbf{x} + m\mathbf{v}_2) \cdot \int_{\mathbf{y} \in \mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) \, d\mathbf{y} \, d\mathbf{x} \\ &= \sum_{m \in \mathbb{Z}} \int_{\mathbf{x} \in \Omega_\Sigma} p_{k_\parallel, \omega}^\lambda(\mathbf{x} + m\mathbf{v}_2) \cdot \int_{\mathbf{y} \in \mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)(\mathbf{x} + m\mathbf{v}_2, \mathbf{y} + m\mathbf{v}_2) f(\mathbf{y} + m\mathbf{v}_2) \, d\mathbf{y} \, d\mathbf{x} \end{aligned}$$

The latter equality holds by properties of $\mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)$ and f under translation by \mathbf{v}_2 . Continuing, we have

$$\begin{aligned} \left\langle \mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)[f], p_{k_\parallel, [\omega]}^\lambda \right\rangle_{L^2(\Sigma)} &= \sum_{m \in \mathbb{Z}} \int_{\Omega_\Sigma + m\mathbf{v}_2} p_{k_\parallel, \omega}^\lambda(\mathbf{x}') \int_{\mathbf{y}' \in \mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)(\mathbf{x}', \mathbf{y}') f(\mathbf{y}') \, d\mathbf{y}' \, d\mathbf{x}' \\ &= \int_{\mathbf{x} \in \mathbb{R}^2} p_{k_\parallel, \omega}^\lambda(\mathbf{x}) \int_{\mathbf{y} \in \mathbb{R}^2} \mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) \, d\mathbf{y} \, d\mathbf{x} \\ &= \lim_{N \rightarrow \infty} \int_{\mathbf{x} \in \mathbb{R}^2} p_{k_\parallel, \omega}^\lambda(\mathbf{x}) \int_{|\mathbf{y}| \leq N} \mathcal{K}_\Gamma^\lambda(\Omega, k_\parallel)(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) \, d\mathbf{y} \, d\mathbf{x} = 0 \end{aligned}$$

by property (10.86) of Proposition 10.15. Again, the formal manipulations are easily justified. This completes the proof of Proposition 10.17.

10.6. Green's kernel.

We recall the cylinder $\Sigma = \mathbb{R}^2 / \mathbb{R}\mathbf{v}_2$ and the choice of fundamental domain $\Omega_\Sigma \subset \mathbb{R}^2$, given as the union of finite parallelograms, Ω_n , $n \geq 0$ together with one unbounded parallelogram, Ω_{-1} , $\Omega_\Sigma = \cup_{n \geq 0} \Omega_n \cup \Omega_{-1}$; see (4.2). In each finite parallelogram, Ω_n , $n \geq 0$, are two lattice points of $\mathbb{H}_\sharp^{(n)}$: $\mathbf{v}_A^{(n)}$ and $\mathbf{v}_B^{(n)}$. As our discrete set we take $\Gamma = \mathbb{H}_\sharp$, our potential $V_\sharp(\mathbf{x})$ and our Hamiltonian H_\sharp^λ acting on $L_{k_\parallel}^2(\Sigma)$.

Next recall the subspace of $L^2(\Sigma)$ (see (4.12)):

$$\mathcal{X}_{AB}^\lambda(k_\parallel) = \text{orthogonal complement in } L^2(\Sigma) \text{ of span} \left\{ p_{k_\parallel, I}^\lambda[n] : n \geq 0, I = A, B \right\}$$

with orthogonal projection:

$$\Pi_{AB}^\lambda(k_\parallel) : L^2(\Sigma) \rightarrow \mathcal{X}_{AB}^\lambda(k_\parallel) .$$

By definition

$$p_{k_\parallel, [\mathbf{v}_I^{(n)}]}^\lambda(\mathbf{x}) = p_{k_\parallel, I}^\lambda[n](\mathbf{x}), \quad I = A, B,$$

where $p_{k_\parallel, I}^\lambda[n]$ is defined in (4.3).

Recall that $\mathcal{K}_\#^\lambda(\Omega, k_\parallel)$, the inverse of $\Pi_{AB}(k_\parallel) (H_\#^\lambda(k_\parallel) - \Omega) \Pi_{AB}(k_\parallel)$ (equivalently $\Pi_{AB}(k_\parallel) \circ (H_\#^\lambda(k_\parallel) - \Omega)$) acting on $\mathcal{X}_{AB}^\lambda(k_\parallel)$; see Proposition 5.1. By Propositions 10.15 and 10.16 this inverse is given by an integral operator

$$(10.105) \quad f \mapsto \mathcal{K}_\#^\lambda(\Omega, k_\parallel)[f] \equiv \int_{\mathbb{R}^2} \mathcal{K}_\#^\lambda(\mathbf{x}, \mathbf{y}; \Omega, k_\parallel) f(\mathbf{y}) d\mathbf{y},$$

with kernel

$$(10.106) \quad \mathcal{K}_\#^\lambda(\mathbf{x}, \mathbf{y}; \Omega, k_\parallel) = e^{-i\frac{\mathbf{R}_2 \cdot \mathbf{x}}{2\pi} k_\parallel} \mathcal{K}_\#^\lambda(\mathbf{x}, \mathbf{y}, \Omega) e^{i\frac{\mathbf{R}_2 \cdot \mathbf{y}}{2\pi} k_\parallel}$$

which satisfies the pointwise bounds:

$$(10.107) \quad |\mathcal{K}_\#^\lambda(\mathbf{x}, \mathbf{y}; \Omega, k_\parallel)| \leq C \left[|\log |\mathbf{x} - \mathbf{y}| | + \lambda^6 \right] \mathbf{1}_{|\mathbf{x} - \mathbf{y}| \leq C} + e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|}$$

for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$.

Now applying Proposition 10.17 we obtain:

Proposition 10.18. *Let $|\Omega| \leq e^{-\hat{c}\lambda}$ with λ chosen sufficiently large.*

(1) $\mathcal{K}_\#^\lambda(\Omega, k_\parallel)[f] = 0$ in $L^2(\Sigma)$ for all $f \in \text{span}\{p_{k_\parallel, I}^\lambda[n] : I = A, B, n \geq 0\}$.

(2) Assume $f \in L^2(\Sigma)$. Then, for all $n \geq 0$ and $I = A, B$, we have

$$\left\langle \mathcal{K}_\#^\lambda(\Omega, k_\parallel)[f], p_{k_\parallel, I}^\lambda[n] \right\rangle_{L^2(\Sigma)} = 0.$$

(3) $[H^\lambda(k_\parallel) - \Omega]\mathcal{K}_\#^\lambda(\Omega, k_\parallel)[f] = f$ modulo $\text{span}\{p_{k_\parallel, I}^\lambda[n] : I = A, B, n \geq 0\}$.

A consequence of the forgoing discussion is:

Corollary 10.19. *Let $|\Omega| \leq e^{-\hat{c}\lambda}$ with λ chosen sufficiently large. The operator $\mathcal{K}_\#^\lambda(\Omega, k_\parallel)$, the inverse of $\Pi_{AB}^\lambda(k_\parallel) (H_\#^\lambda(k_\parallel) - \Omega) \Pi_{AB}^\lambda(k_\parallel)$, arises from a kernel satisfying (10.105)-(10.107). $\mathcal{K}_\#^\lambda(\Omega, k_\parallel)$ is a bounded linear operator on $L^2(\Sigma)$.*

11. EXPANSION AND ESTIMATION OF LINEAR MATRIX ELEMENTS: PROOF OF PROPOSITION 7.1

Our first step in the proof of Proposition 7.1 is to expand the inner products:

$$\left\langle P_{k_\parallel, I}^\lambda[m], H_\#^\lambda P_{k_\parallel, J}^\lambda[n] \right\rangle_{L^2(\Sigma)} = \left\langle p_{k_\parallel, I}^\lambda[m], H_\#^\lambda(k_\parallel) p_{k_\parallel, J}^\lambda[n] \right\rangle_{L^2(\Sigma)},$$

where $m, n \in \mathbb{N}_0$, in terms of overlap integrals of translates of the atomic potential, V_0 , and the atomic ground state, p_0^λ . We have, by the definition of the $L^2(\Sigma)$ inner product:

$$\left\langle P_{k_\parallel, I}^\lambda[m], H_\#^\lambda P_{k_\parallel, J}^\lambda[n] \right\rangle_{L^2(\Sigma)} = \int_{\Omega_\Sigma} \overline{P_{k_\parallel, I}^\lambda[m](\mathbf{x})} H_\#^\lambda P_{k_\parallel, J}^\lambda[n](\mathbf{x}) d\mathbf{x}.$$

We first simplify the integrand: $\overline{P_{k_\parallel, I}^\lambda[m]} H_\#^\lambda P_{k_\parallel, J}^\lambda[n]$. We recall the definition of $H_\#^\lambda$ (see (1.16)) and introduce the notation:

$$(11.1) \quad J' = A \text{ if } J = B \text{ and } J' = B \text{ if } J = A.$$

For $\mathbf{x} \in \Omega_\Sigma$, the fundamental domain (see Figure 1), we have for $J = A, B$:

$$\begin{aligned}
H_\#^\lambda P_{k_\parallel, J}^\lambda[n](\mathbf{x}) &= \sum_{\tilde{m}_2 \in \mathbb{Z}} e^{i\tilde{m}_2 k_\parallel} \\
&\quad \cdot \left[-\Delta + \lambda^2 \sum_{n_1 \geq 0} V_0(\mathbf{x} - \mathbf{v}_J^{n_1}) + \lambda^2 \sum_{n_1 \geq 0} V_0(\mathbf{x} - \mathbf{v}_{J'}^{n_1}) - E_0^\lambda \right] p_0^\lambda(\mathbf{x} - \mathbf{v}_J^n - \tilde{m}_2 \mathbf{v}_2) \\
&= \lambda^2 \left[\sum_{\substack{n_1 \geq 0 \\ n_1 \neq n}} V_0(\mathbf{x} - \mathbf{v}_J^{n_1}) + \sum_{n_1 \geq 0} V_0(\mathbf{x} - \mathbf{v}_{J'}^{n_1}) \right] \cdot \left[\sum_{\tilde{m}_2 \in \mathbb{Z}} e^{i\tilde{m}_2 k_\parallel} p_0^\lambda(\mathbf{x} - \mathbf{v}_J^n - \tilde{m}_2 \mathbf{v}_2) \right] \\
(11.2) \quad &+ \lambda^2 V_0(\mathbf{x} - \mathbf{v}_J^n) \sum_{\tilde{m}_2 \in \mathbb{Z} \setminus \{0\}} e^{i\tilde{m}_2 k_\parallel} p_0^\lambda(\mathbf{x} - \mathbf{v}_J^n - \tilde{m}_2 \mathbf{v}_2).
\end{aligned}$$

To obtain (11.2) we use that $(-\Delta_{\mathbf{x}} + \lambda^2 V_0(\mathbf{x}) - E_0^\lambda) p_0^\lambda(\mathbf{x}) = 0$ and therefore $(-\Delta_{\mathbf{x}} + \lambda^2 V_0(\mathbf{x} - \mathbf{v}) - E_0^\lambda) p_0^\lambda(\mathbf{x} - \mathbf{v}) = 0$ for all $\mathbf{v} \in \mathbb{H}$. From (11.2) we obtain:

$$\begin{aligned}
&\overline{P_{k_\parallel, I}^\lambda[m](\mathbf{x})} H_\#^\lambda P_{k_\parallel, J}^\lambda[n](\mathbf{x}) \\
&= \sum_{m_2 \in \mathbb{Z}} \sum_{\tilde{m}_2 \in \mathbb{Z}} e^{i(\tilde{m}_2 - m_2) k_\parallel} \\
&\quad \cdot p_0^\lambda(\mathbf{x} - \mathbf{v}_I^m - m_2 \mathbf{v}_2) \left[\sum_{\substack{n_1 \geq 0 \\ n_1 \neq n}} \lambda^2 V_0(\mathbf{x} - \mathbf{v}_J^{n_1}) + \sum_{n_1 \geq 0} \lambda^2 V_0(\mathbf{x} - \mathbf{v}_{J'}^{n_1}) \right] p_0^\lambda(\mathbf{x} - \mathbf{v}_J^n - \tilde{m}_2 \mathbf{v}_2) \\
&\quad + \sum_{m_2 \in \mathbb{Z}} \sum_{\tilde{m}_2 \in \mathbb{Z} \setminus \{0\}} e^{i(\tilde{m}_2 - m_2) k_\parallel} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_J^n) p_0^\lambda(\mathbf{x} - \mathbf{v}_J^n - \tilde{m}_2 \mathbf{v}_2),
\end{aligned}$$

for all $\mathbf{x} \in \Omega_\Sigma$. Integrating the previous identity over Ω_Σ , we obtain:

$$\begin{aligned}
&\left\langle P_{k_\parallel, I}^\lambda[m](\mathbf{x}), H_\#^\lambda P_{k_\parallel, J}^\lambda[n](\mathbf{x}) \right\rangle_{L^2(\Sigma)} \\
&= \sum_{m_2, \tilde{m}_2 \in \mathbb{Z}} \sum_{\substack{n_1 \geq 0 \\ n_1 \neq n}} e^{i(\tilde{m}_2 - m_2) k_\parallel} \int_{\Omega_\Sigma} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_J^{n_1}) p_0^\lambda(\mathbf{x} - \mathbf{v}_J^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x} \\
&\quad + \sum_{m_2, \tilde{m}_2 \in \mathbb{Z}} \sum_{n_1 \geq 0} e^{i(\tilde{m}_2 - m_2) k_\parallel} \int_{\Omega_\Sigma} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_{J'}^{n_1}) p_0^\lambda(\mathbf{x} - \mathbf{v}_J^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x} \\
&\quad + \sum_{m_2 \in \mathbb{Z}} \sum_{\tilde{m}_2 \in \mathbb{Z} \setminus \{0\}} e^{i(\tilde{m}_2 - m_2) k_\parallel} \int_{\Omega_\Sigma} p_0^\lambda(\mathbf{x} - \mathbf{v}_I^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_J^n) p_0^\lambda(\mathbf{x} - \mathbf{v}_J^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x} \\
(11.3) \quad &\equiv S_1^{IJ}(m, n) + S_2^{IJ}(m, n) + S_3^{IJ}(m, n),
\end{aligned}$$

where the three expressions $S_1^{IJ}(m, n)$, $S_2^{IJ}(m, n)$, and $S_3^{IJ}(m, n)$ denote the three sums in (11.3). The dependence on λ and k_\parallel has been suppressed. We recall that in the expression for $S_2^{IJ}(m, n)$, the index J' is defined in (11.1).

We now provide a general lemma, which will facilitate our determination of the leading terms and estimation of the error terms in the above sums. In preparation for the statement of this lemma we introduce some terminology.

Definition 11.1. (1) For $I_1, J_1 \in \{A, B\}$, we write $\mathbf{v}_{I_1} - \mathbf{v}_{J_1} = \sigma(\mathbf{v}_B - \mathbf{v}_A) = \sigma\mathbf{e}$, where $\sigma = 1$ if $I_1 = B$ and $J_1 = A$, and $\sigma = -1$ if $I_1 = A$ and $J_1 = B$. We therefore write:

$$(11.4) \quad \sigma(B, A) = +1, \quad \sigma(A, B) = -1, \quad \text{and we define } \sigma(I_1, I_1) = 0.$$

(2) For $\sigma = +1, -1, 0$ we define $N_b(\sigma) = \{\mathbf{r} = (r_1, r_2) \in \mathbb{Z}^2 : |\sigma\mathbf{e} + \mathbf{r}\vec{\mathbf{v}}| = |\mathbf{e}|\}$. Therefore $N_b(+1) \equiv \{(0, 0), (-1, 0), (0, -1)\}$, $N_b(-1) \equiv \{(0, 0), (1, 0), (0, 1)\}$, and $N_b(0) \equiv \emptyset$.

Note that if $\mathbf{m} = (m_1, m_2) \in N_b(\sigma)$ with $\sigma = \pm 1$, then there exists $l \in \{0, 1, 2\}$ such that

$$(11.5) \quad \sigma\mathbf{e} + m_1\mathbf{v}_1 + m_2\mathbf{v}_2 = R^l\mathbf{e}$$

where R denotes the 2×2 rotation in \mathbb{R}^2 by $2\pi/3$.

Lemma 11.2. For $I_1, J_1, \tilde{I}_1 \in \{A, B\}$, $m, n, n_1 \geq 0$ and $m_2, \tilde{m}_2 \in \mathbb{Z}$, consider the overlap integral

$$(11.6) \quad \mathcal{J}_\# \equiv \int p_0^\lambda(\mathbf{x} - \mathbf{v}_{I_1}^m - m_2\mathbf{v}_2) \lambda^2 |V_0(\mathbf{x} - \mathbf{v}_{J_1}^{n_1})| p_0^\lambda(\mathbf{x} - \mathbf{v}_{\tilde{I}_1}^n - \tilde{m}_2\mathbf{v}_2) d\mathbf{x}.$$

Recall the hopping coefficient defined by: $\rho_\lambda = \int p_0^\lambda(\mathbf{y}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} - \mathbf{e}) d\mathbf{y}$. Then we have the bound

$$(11.7) \quad \mathcal{J}_\# \lesssim e^{-c\lambda(|m-n_1| + |m_2| + |n-n_1| + |\tilde{m}_2|)} \rho_\lambda,$$

except in the following cases of exceptional indices $(m, n, n_1, m_2, \tilde{m}_2)$:

(a) $I_1 = \tilde{I}_1 = J_1$, $m = n = n_1$ and $m_2 = \tilde{m}_2 = 0$. This case does not arise in the proof of Proposition 7.1 so we say nothing further about it.

(b) $\tilde{I}_1 = J_1$, $I_1 \neq J_1$, $(m - n_1, m_2) \in N_b(\sigma(I_1, J_1))$, $n = n_1$ and $\tilde{m}_2 = 0$, in which case $\mathcal{J}_\# = \rho_\lambda$.

(c) $I_1 = J_1$, $\tilde{I}_1 \neq J_1$, $(n - n_1, \tilde{m}_2) \in N_b(\sigma(\tilde{I}_1, J_1))$, $m = n_1$ and $m_2 = 0$, in which case $\mathcal{J}_\# = \rho_\lambda$.

Furthermore, if $I_1 \neq J_1$, $\tilde{I}_1 \neq J_1$, then for all $m, n, n_1, m_2, \tilde{m}_2$:

$$(11.8) \quad \mathcal{J}_\# \lesssim e^{-c\lambda} e^{-c\lambda(|m-n_1| + |m_2| + |n-n_1| + |\tilde{m}_2|)} \rho_\lambda.$$

Lemma 11.2 is proved in Appendix B. It makes repeated use of the following pointwise decay estimates for the atomic ground state, p_0^λ :

Lemma 11.3 (See Lemma 15.6 of [26]). *There exists a constant c such that for $\mathbf{y} \in \text{supp}(V_0) \subset B_{r_0}(\mathbf{0})$, i.e. $|\mathbf{y}| \leq r_0$, we have:*

$$(11.9) \quad p_0^\lambda(\mathbf{y} - \mathbf{n}\vec{\mathbf{v}}) \lesssim e^{-c|\mathbf{n}|\lambda} p_0^\lambda(\mathbf{y}), \quad \mathbf{n} \in \mathbb{Z}^2,$$

$$(11.10) \quad p_0^\lambda(\mathbf{y} - (\sigma\mathbf{e} + \mathbf{n}\vec{\mathbf{v}})) \lesssim e^{-c|\mathbf{n}|\lambda} p_0^\lambda(\mathbf{y} - \sigma\mathbf{e}), \quad \mathbf{n} \notin N_b(\sigma), \quad \sigma = \pm 1,$$

$$(11.11) \quad p_0^\lambda(\mathbf{y} - \sigma\mathbf{e}) \lesssim e^{-c\lambda} p_0^\lambda(\mathbf{y}), \quad \sigma = \pm 1, \quad \text{and}$$

$$(11.12) \quad p_0^\lambda(\mathbf{y} - \mathbf{n}\vec{\mathbf{v}}) \lesssim e^{-c\lambda|\mathbf{n}|} p_0^\lambda(\mathbf{y} - \sigma\mathbf{e}), \quad \mathbf{n} \in \mathbb{Z}^2 \setminus \{(0, 0)\}.$$

Remark 11.4. In [26], Lemma 11.3 was proved for all r_0 satisfying $0 < r_0 < r_{\text{critical}}$, where $0.33|\mathbf{e}| \leq r_{\text{critical}} < 0.5|\mathbf{e}|$, and $|\mathbf{e}| = |\mathbf{v}_B - \mathbf{v}_A| = 1/\sqrt{3}$.

To prove Proposition 7.1, we now apply Lemma 11.2 to the expansion of the matrix elements: $\left\langle P_{k_{\parallel},I}^{\lambda}[m](\mathbf{x}), H_{\sharp}^{\lambda} P_{k_{\parallel},J}^{\lambda}[n] \right\rangle_{L^2(\Sigma)}$, where $I, J = A, B$ and $m, n \in \mathbb{N}_0$, for large λ .

11.1. Expansion of the inner product $\left\langle P_{k_{\parallel},B}^{\lambda}[m](\mathbf{x}), H_{\sharp}^{\lambda} P_{k_{\parallel},A}^{\lambda}[n] \right\rangle_{L^2(\Sigma)}$.

We consider the summations $S_j^{IJ}(m, n), j = 1, 2, 3$ in order (see (11.3)) with $I = B$ and $J = A$.

Estimation of $S_1^{BA}(m, n)$: The expression to be summed over $m_2, \tilde{m}_2 \in \mathbb{Z}$ and $n_1 \geq 0, n_1 \neq n$ is:

$$(11.13) \quad e^{i(\tilde{m}_2 - m_2)k_{\parallel}} \int_{\mathbb{R}^2} p_0^{\lambda}(\mathbf{x} - \mathbf{v}_B^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_A^{n_1}) p_0^{\lambda}(\mathbf{x} - \mathbf{v}_A^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x}.$$

We apply Lemma 11.2 with $I_1 = B, J_1 = A$ and $\tilde{I}_1 = A$. All summands (11.13) of $S_1^{BA}(m, n)$, except for exceptional indices in case (b), defined by $\tilde{I}_1 = J_1, I_1 \neq J_1$, are bounded by $e^{-c\lambda(|m-n_1|+|m_2|+|n-n_1|+|\tilde{m}_2|)} \rho_{\lambda}$. The exceptional indices are characterized by the relations: $(m - n_1, m_2) \in N_b(\sigma(B, A)) = N_b(+1), n = n_1$ and $\tilde{m}_2 = 0$. Since the sum in the definition of $S_1^{BA}(m, n)$ is over $n_1 \geq 0$ with $n_1 \neq n$, there are no relevant exceptional indices and we conclude for all $m, n \geq 0$:

$$(11.14) \quad |S_1^{BA}(m, n)| \lesssim \rho_{\lambda} \sum_{m_2, \tilde{m}_2 \in \mathbb{Z}} \sum_{\substack{n_1 \geq 0 \\ n_1 \neq n}} e^{-c\lambda(|m-n_1|+|m_2|+|n-n_1|+|\tilde{m}_2|)} \lesssim e^{-c'\lambda} e^{-c'\lambda|m-n|} \rho_{\lambda},$$

for some strictly positive constant c' .

Expansion of $S_2^{BA}(m, n)$: Since $I = B, J = A$ and $J' = B$, the expression to be summed over $m_2, \tilde{m}_2 \in \mathbb{Z}$ and $n_1 \geq 0$ is:

$$(11.15) \quad e^{i(\tilde{m}_2 - m_2)k_{\parallel}} \int_{\mathbb{R}^2} p_0^{\lambda}(\mathbf{x} - \mathbf{v}_B^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_B^{n_1}) p_0^{\lambda}(\mathbf{x} - \mathbf{v}_A^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x}.$$

We apply Lemma 11.2 with $I_1 = B, J_1 = B$ and $\tilde{I}_1 = A$. All summands (11.15) of $S_2^{BA}(m, n)$, except for exceptional indices in case (c), defined by $I_1 = J_1$ and $\tilde{I}_1 \neq J_1$, are bounded by $e^{-c\lambda(|m-n_1|+|m_2|+|n-n_1|+|\tilde{m}_2|)} \rho_{\lambda}$. The exceptional indices are characterized by the relations: $(n - n_1, \tilde{m}_2) \in N_b(\sigma(\tilde{I}_1, J_1)) = N_b(\sigma(A, B)) = N_b(-1) = \{(0, 0), (1, 0), (0, 1)\}, m = n_1$ and $m_2 = 0$. We next simplify the expression (11.15) in each of these three exceptional cases.

$(n - n_1, \tilde{m}_2) = (0, 0), m = n_1, m_2 = 0$: We have $n_1 = m = n$ and $m_2 = \tilde{m}_2 = 0$. For this case, the expression in (11.15) is equal to $-\rho_{\lambda}$ and contributes to $S_2^{BA}(m, m)$.

$(n - n_1, \tilde{m}_2) = (0, 1), m = n_1, m_2 = 0$: We have $n_1 = n = m, m_2 = 0$ and $\tilde{m}_2 = 1$. For this case, the expression (11.15) is equal to $-e^{ik_{\parallel}} \rho_{\lambda}$ and contributes to $S_2^{BA}(m, m)$.

$(n - n_1, \tilde{m}_2) = (1, 0), m = n_1, m_2 = 0$: We have $n_1 = m, n = m + 1, m_2 = \tilde{m}_2 = 0$. For this case, the expression in (11.15) is equal to $-\rho_{\lambda}$ and contributes to $S_2^{BA}(m, m + 1)$.

We conclude from the above discussion of $S_2^{BA}(m, n)$ that:

$$(11.16) \quad S_2^{BA}(m, m) = -(1 + e^{ik_{\parallel}}) \rho_{\lambda} + \mathcal{O}(e^{-c\lambda} \rho_{\lambda}), \quad (n = m)$$

$$(11.17) \quad S_2^{BA}(m, m+1) = -\rho_{\lambda} + \mathcal{O}(e^{-c\lambda} \rho_{\lambda}), \quad (n = m+1)$$

$$(11.18) \quad S_2^{BA}(m, n) = \mathcal{O}(e^{-c\lambda} e^{-c\lambda|m-n|} \rho_{\lambda}), \quad \text{if } n \neq m, m+1.$$

The $\mathcal{O}(\cdot)$ error terms are bounds on contributions to $S_2^{BA}(m, n)$ arising from the summation over $m_2, \tilde{m}_2 \in \mathbb{Z}$ and $n_1 \geq 0$ of the bound $e^{-c\lambda(|m-n_1|+|m_2|+|n-n_1|+|\tilde{m}_2|)} \rho_{\lambda}$ for non-exceptional indices (as in (11.14)).

Expansion of $S_3^{BA}(m, n)$: Since $I = B$ and $J = A$, the expression to be summed over $m_2 \in \mathbb{Z}$ and $\tilde{m}_2 \in \mathbb{Z} \setminus \{0\}$ is:

$$(11.19) \quad e^{i(\tilde{m}_2 - m_2)k_{\parallel}} \int_{\mathbb{R}^2} p_0^{\lambda}(\mathbf{x} - \mathbf{v}_B^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_A^n) p_0^{\lambda}(\mathbf{x} - \mathbf{v}_A^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x}.$$

We apply Lemma 11.2 with $I_1 = B$, $J_1 = A$, $\tilde{I}_1 = A$ and $n_1 = n$. All summands (11.19) of $S_3^{BA}(m, n)$, except for exceptional indices in case (b), defined by $I_1 \neq J_1$ and $\tilde{I}_1 = J_1$, are bounded by $e^{-c\lambda(|m-n|+|m_2|+|\tilde{m}_2|)} \rho_{\lambda}$ ($n_1 = n$). Now exceptional indices in case (b) of Lemma 11.2 are such that $\tilde{m}_2 = 0$. However, in $S_3^{BA}(m, n)$ we sum over $\tilde{m}_2 \neq 0$. Hence, there are no relevant exceptional indices and therefore all expressions (11.19) are bounded by $e^{-c\lambda(|m-n|+|m_2|+|\tilde{m}_2|)} \rho_{\lambda}$. Summing over $m_2 \in \mathbb{Z}$ and $\tilde{m}_2 \in \mathbb{Z} \setminus \{0\}$ we obtain:

$$(11.20) \quad |S_3^{BA}(m, n)| \lesssim e^{-c\lambda} e^{-c\lambda|m-n|}, \quad m, n \geq 0.$$

Putting together the expression (11.3) for the inner product $\left\langle P_{k_{\parallel}, B}^{\lambda}[m](\mathbf{x}), H_{\sharp}^{\lambda} P_{k_{\parallel}, A}^{\lambda}[n] \right\rangle_{L^2(\Sigma)}$ with the expansions and bounds in (11.14), (11.16), (11.17), (11.18) and (11.20) we obtain:

$$(11.21) \quad \left\langle P_{k_{\parallel}, B}^{\lambda}[m](\mathbf{x}), H_{\sharp}^{\lambda} P_{k_{\parallel}, A}^{\lambda}[n] \right\rangle_{L^2(\Sigma)} = \begin{cases} -(1 + e^{ik_{\parallel}}) \rho_{\lambda} + \mathcal{O}(e^{-c\lambda} e^{-c\lambda|m-n|} \rho_{\lambda}), & n = m \\ -\rho_{\lambda} + \mathcal{O}(e^{-c\lambda} e^{-c\lambda|m-n|} \rho_{\lambda}), & n = m+1 \\ \mathcal{O}(e^{-c\lambda} e^{-c\lambda|m-n|} \rho_{\lambda}), & n \neq m, m+1. \end{cases}$$

By self-adjointness,

$$(11.22) \quad \left\langle P_{k_{\parallel}, A}^{\lambda}[m](\mathbf{x}), H_{\sharp}^{\lambda} P_{k_{\parallel}, B}^{\lambda}[n] \right\rangle_{L^2(\Sigma)} = \begin{cases} -(1 + e^{-ik_{\parallel}}) \rho_{\lambda} + \mathcal{O}(e^{-c\lambda} e^{-c\lambda|m-n|} \rho_{\lambda}), & n = m \\ -\rho_{\lambda} + \mathcal{O}(e^{-c\lambda} e^{-c\lambda|m-n|} \rho_{\lambda}), & n = m-1 \\ \mathcal{O}(e^{-c\lambda} e^{-c\lambda|m-n|} \rho_{\lambda}), & n \neq m, m-1. \end{cases}$$

Equations (11.22) and (11.21) imply assertions (1), (2) and (3) of Proposition 7.1.

Finally, we turn to the proof of part (4) of Proposition 7.1. By (11.3), we have for $I = A, B$:

$$\left\langle P_{k_{\parallel}, I}^{\lambda}[m](\mathbf{x}), H_{\sharp}^{\lambda} P_{k_{\parallel}, I}^{\lambda}[n] \right\rangle_{L^2(\Sigma)} = S_1^{II}(m, n) + S_2^{II}(m, n) + S_3^{II}(m, n).$$

We claim that $|S_j^{II}(m, n)| \lesssim e^{-c\lambda} e^{-c\lambda|m-n|}$ for $j = 1, 2, 3$ and $I = A, B$. We consider the case $I = A$. The case $I = B$ is essentially the same.

Estimation of $S_1^{AA}(m, n)$: The expression to be summed over $m_2, \tilde{m}_2 \in \mathbb{Z}$ for $n_1 \geq 0, n_1 \neq n$ is:

$$(11.23) \quad e^{i(\tilde{m}_2 - m_2)k_{\parallel}} \int_{\mathbb{R}^2} p_0^\lambda(\mathbf{x} - \mathbf{v}_A^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_A^{n_1}) p_0^\lambda(\mathbf{x} - \mathbf{v}_A^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x} .$$

We apply Lemma 11.2 with $I_1 = A, J_1 = A$ and $\tilde{I}_1 = A$. All summands in the expression for $S_1^{AA}(m, n)$, except for exceptional indices are bounded by $e^{-c\lambda(|m-n_1|+|m_2|+|n-n_1|+|\tilde{m}_2|)} \rho_\lambda$. The only possible exceptional indices are of case (a) in Lemma 11.2. This case requires $n_1 = n$ and since the summation in $S_1^{AA}(m, n)$ is over $n_1 \geq 0$ with $n_1 \neq n$, there are no relevant exceptional indices. We conclude for all $m, n \geq 0$:

$$(11.24) \quad |S_1^{AA}(m, n)| \lesssim \rho_\lambda \sum_{m_2, \tilde{m}_2 \in \mathbb{Z}} \sum_{\substack{n_1 \geq 0 \\ n_1 \neq n}} e^{-c\lambda(|m-n_1| + |m_2| + |n-n_1| + |\tilde{m}_2|)} \lesssim e^{-c'\lambda} e^{-c'\lambda|m-n|} \rho_\lambda ,$$

for some strictly positive constant c' .

Estimation of $S_2^{AA}(m, n)$: The expression to be summed over $m_2, \tilde{m}_2 \in \mathbb{Z}$ for $n_1 \geq 0$ is

$$e^{i(\tilde{m}_2 - m_2)k_{\parallel}} \int p_0^\lambda(\mathbf{x} - \mathbf{v}_A^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_B^{n_1}) p_0^\lambda(\mathbf{x} - \mathbf{v}_A^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x} .$$

Since $I_1 = A, J_1 = B$ and $\tilde{I}_1 = A$, we have that $I_1 \neq J_1$ and $\tilde{I}_1 \neq J_1$. Hence, the bound (11.8) applies. Thus, all summands in the expression for $S_2^{AA}(m, n)$ are bounded by $e^{-c\lambda} e^{-c\lambda(|m-n_1|+|m_2|+|n-n_1|+|\tilde{m}_2|)} \rho_\lambda$. Summing over all relevant indices we have:

$$(11.25) \quad |S_2^{AA}(m, n)| \lesssim e^{-c\lambda} \sum_{m_2, \tilde{m}_2 \in \mathbb{Z}} \sum_{n_1 \geq 0} e^{-c\lambda(|m-n_1|+|m_2|+|n-n_1|+|\tilde{m}_2|)} \rho_\lambda \lesssim e^{-c'\lambda} e^{-c'\lambda|m-n|} \rho_\lambda ,$$

for some strictly positive constant c' .

Estimation of $S_3^{AA}(m, n)$: The expression to be summed over $m_2 \in \mathbb{Z}$ and $\tilde{m}_2 \in \mathbb{Z} \setminus \{0\}$ for $n \geq 0$ is

$$e^{i(\tilde{m}_2 - m_2)k_{\parallel}} \int p_0^\lambda(\mathbf{x} - \mathbf{v}_A^m - m_2 \mathbf{v}_2) \lambda^2 V_0(\mathbf{x} - \mathbf{v}_A^n) p_0^\lambda(\mathbf{x} - \mathbf{v}_A^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x} .$$

Since $I_1 = J_1 = \tilde{I}_1 = A$, the only possible exceptional case is case (a). However, note that $\tilde{m}_2 = 0$ is omitted in the summation and hence there are no relevant exceptional cases. Thus, summands in the expression for $S_3^{AA}(m, n)$ are bounded by $e^{-c\lambda(|m-n|+|m_2|+|\tilde{m}_2|)} \rho_\lambda$, and we have:

$$(11.26) \quad |S_3^{AA}(m, n)| \lesssim \sum_{m_2 \in \mathbb{Z}} \sum_{\tilde{m}_2 \in \mathbb{Z} \setminus \{0\}} e^{-c\lambda(|m-n|+|m_2|+|\tilde{m}_2|)} \rho_\lambda \lesssim e^{-c'\lambda} e^{-c'\lambda|m-n|} \rho_\lambda ,$$

for some strictly positive constant c' .

Finally, summing the bounds (11.24), (11.25) and (11.26) implies the bound (7.10). This completes the proof of Proposition 7.1.

12. ESTIMATION OF THE NONLINEAR MATRIX ELEMENTS; PROOF OF PROPOSITION 7.2

Recall our decomposition of $\mathcal{M}^\lambda[m, n](\Omega, k_\parallel)$ into its *linear* and *nonlinear* contributions:

$$(12.1) \quad \mathcal{M}^\lambda[m, n](\Omega, k_\parallel) = \mathcal{M}^{\lambda,1}[m, n](\Omega; k_\parallel) - \mathcal{M}^{\lambda, \text{nl}}[m, n](\Omega; k_\parallel),$$

where the latter nonlinear matrix elements are given by (see (6.11)):

$$(12.2) \quad \begin{aligned} & \mathcal{M}_{JI}^{\lambda, \text{nl}}[m, n](\Omega; k_\parallel) \\ & \equiv \left\langle H_\sharp^\lambda(k_\parallel) p_{k_\parallel, J}^\lambda[m], \Pi_{AB}(k_\parallel) \mathcal{K}_\sharp^\lambda(\Omega, k_\parallel) \Pi_{AB}(k_\parallel) H_\sharp^\lambda(k_\parallel) p_{k_\parallel, I}^\lambda[n] \right\rangle_{L^2(\Sigma)}. \end{aligned}$$

Here, we recall (from Section 4.1) $\Pi_{AB}(k_\parallel)$ denotes the projection onto

$$\mathcal{X}_{AB}(k_\parallel) = \text{the orthogonal complement in } L^2(\Sigma) \text{ of } \text{span} \left\{ p_{k_\parallel, I}^\lambda[n] : I = A, B, n \geq 0 \right\},$$

and $\Pi_{AB}(k_\parallel) \mathcal{K}_\sharp^\lambda(\Omega, k_\parallel) \Pi_{AB}(k_\parallel) : \mathcal{X}_{AB}(k_\parallel) \rightarrow \mathcal{X}_{AB}(k_\parallel)$ is the inverse of

$$\Pi_{AB}(k_\parallel) \left(- \left(\nabla_{\mathbf{x}} + i \frac{k_\parallel}{2\pi} \mathfrak{R}_2 \right)^2 + V_\sharp - E_0^\lambda - \Omega \right) \Pi_{AB}(k_\parallel).$$

Furthermore, the operator $\Pi_{AB}(k_\parallel) \mathcal{K}_\sharp^\lambda(\Omega, k_\parallel) \Pi_{AB}(k_\parallel)$ arises from a kernel $\mathcal{K}_\sharp^\lambda(\mathbf{x}, \mathbf{y}, \Omega, k_\parallel)$; see Corollary 10.19. And finally we recall the projection operators Π_Γ^λ (see (10.82)) which projects onto the orthogonal complement of the set of atomic ground states, centered at nuclei of the discrete set Γ ,

$$\mathcal{X}_\Gamma \equiv \text{span} \left\{ p_\omega^\lambda : \omega \in \Gamma \right\}^\perp$$

and $\tilde{\Pi}_\Gamma^\lambda = I - \Pi_\Gamma^\lambda$; see (10.82) and Proposition 10.15. In the following discussion we shall be interested in the choice $\Gamma = \mathbb{H}_\sharp$, the zigzag truncation of \mathbb{H} . Finally, we recall the notation: $F_\omega(\mathbf{x}) = F(\mathbf{x} - \omega)$.

Given $F(\mathbf{x})$, a rapidly decaying function on \mathbb{R}^2 , define

$$(12.3) \quad F_{[\omega]}(\mathbf{x}) \equiv \sum_{n \in \mathbb{Z}} F(\mathbf{x} - \omega + n\mathbf{v}_2) = \sum_{n \in \mathbb{Z}} F_\omega(\mathbf{x} + n\mathbf{v}_2).$$

The functions $p_{k_\parallel, J}^\lambda[m]$ in (12.2) are of this type and we now seek to bound inner products in $L^2(\Sigma)$ of the form (12.2).

For a constant $\gamma > 0$ to be fixed, we introduce the weighted $L^2(\mathbb{R}^2)$ - spaces:

$$(12.4) \quad \mathcal{H}^{(\omega)} \equiv L^2(\mathbb{R}^2; e^{\gamma|\mathbf{x}-\omega|} d\mathbf{x}).$$

Proposition 12.1. *Fix $\Gamma = \mathbb{H}_\sharp$, which is translation-invariant by the vector $\mathbf{v}_2 \in \mathbb{H}$. Let $[\omega], [\omega']$ denote equivalence classes (see (10.103) with $\Gamma = \mathbb{H}_\sharp$), and $\omega_0 \in [\omega] \cap \Omega_\Sigma$ and $\omega'_0 \in [\omega'] \cap \Omega_\Sigma$.*

(1) *For any rapidly decaying functions F and G on \mathbb{R}^2 we have*

$$(12.5) \quad \begin{aligned} & \left\langle F_{[\omega]}, \Pi_{AB}(k_\parallel) \mathcal{K}_\sharp^\lambda(\Omega, k_\parallel) \Pi_{AB}(k_\parallel) G_{[\omega']} \right\rangle_{L^2(\Sigma)} \\ & = \sum_{l \in \mathbb{Z}} \int_{\mathbf{x} \in \mathbb{R}^2} F_{\omega_0}(\mathbf{x}) \int_{\mathbf{y} \in \mathbb{R}^2} \mathcal{K}_\sharp^\lambda(\mathbf{x}, \mathbf{y} + l\mathbf{v}_2; \Omega, k_\parallel) G_{\omega'_0}(\mathbf{y}) d\mathbf{y} d\mathbf{x}. \end{aligned}$$

(2) The expression in (12.5) may be bounded in exponentially weighted norms as follows:

$$(12.6) \quad \left| \left\langle F_{[\omega]}, \Pi_{AB}(k_{\parallel}) \mathcal{K}_{\#}^{\lambda}(\Omega, k_{\parallel}) \Pi_{AB}(k_{\parallel}) G_{[\omega']} \right\rangle_{L^2(\Sigma)} \right| \\ \leq \left[\sum_{l \in \mathbb{Z}} \|\mathcal{K}_{\#}^{\lambda, \omega_0, \omega'_0, l}(\Omega, k_{\parallel})\|_{L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2)} \right] \|F_{\omega_0}\|_{\mathcal{H}(\omega_0)} \|G_{\omega'_0}\|_{\mathcal{H}(\omega'_0)}.$$

where

$$(12.7) \quad (\mathcal{K}_{\#}^{\lambda, \omega_0, \omega'_0, l} f)(\mathbf{x}) = \int_{\mathbb{R}^2} e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} \mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y} + l\mathbf{v}_2) e^{-\frac{\gamma}{2}|\mathbf{y}-\omega'_0|} f(\mathbf{y}) d\mathbf{y}.$$

Note: The above may be formulated for an arbitrary discrete set Γ satisfying $\inf\{|\omega - \omega'| : \omega, \omega' \in \Gamma \text{ distinct}\} > r_4$, which is translation invariant by the vector \mathbf{v}_2 .

Proof of Proposition 12.1: By Corollary 10.19 we have that the operator $\Pi_{AB}(k_{\parallel}) \mathcal{K}_{\#}^{\lambda}(\Omega, k_{\parallel}) \Pi_{AB}$ arises from a kernel $\mathcal{K}_{\Gamma}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega, k_{\parallel})$. We have

$$\begin{aligned} & \left\langle F_{[\omega]}, \Pi_{AB}(k_{\parallel}) \mathcal{K}_{\#}^{\lambda}(\Omega, k_{\parallel}) \Pi_{AB}(k_{\parallel}) G_{[\omega']} \right\rangle_{L^2(\Sigma)} \\ &= \int_{\Omega_{\Sigma}} F_{[\omega]}(\mathbf{x}) \int_{\mathbf{y} \in \mathbb{R}^2} \mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega, k_{\parallel}) G_{[\omega']}(\mathbf{y}) d\mathbf{y} d\mathbf{x} \\ &= \int_{\Omega_{\Sigma}} \sum_{n \in \mathbb{Z}} F(\mathbf{x} - \omega_0 + n\mathbf{v}_2) \int_{\mathbf{y} \in \mathbb{R}^2} \mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega) \sum_{n' \in \mathbb{Z}} G(\mathbf{y} - \omega'_0 + n'\mathbf{v}_2) d\mathbf{y} d\mathbf{x} \\ &= \sum_{n, n' \in \mathbb{Z}} \int_{\Omega_{\Sigma}} F_{\omega_0}(\mathbf{x} + n\mathbf{v}_2) \int_{\mathbf{y} \in \mathbb{R}^2} \mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega, k_{\parallel}) G_{\omega'_0}(\mathbf{y} + n'\mathbf{v}_2) d\mathbf{y} d\mathbf{x} \\ &= \left[\begin{array}{l} \tilde{\mathbf{x}} = \mathbf{x} + n\mathbf{v}_2 \\ \tilde{\mathbf{y}} = \mathbf{y} + n'\mathbf{v}_2 \end{array} \right] \sum_{n, n' \in \mathbb{Z}} \int_{\tilde{\mathbf{x}} \in \Omega_{\Sigma} + n\mathbf{v}_2} F_{\omega_0}(\tilde{\mathbf{x}}) \int_{\tilde{\mathbf{y}} \in \mathbb{R}^2} \mathcal{K}_{\#}^{\lambda}(\tilde{\mathbf{x}} - n\mathbf{v}_2, \tilde{\mathbf{y}} - n'\mathbf{v}_2; \Omega, k_{\parallel}) G_{\omega'_0}(\tilde{\mathbf{y}}) d\tilde{\mathbf{y}} d\tilde{\mathbf{x}} \\ &= \text{by equation (10.97)} \\ & \quad \sum_{n, n' \in \mathbb{Z}} \int_{\tilde{\mathbf{x}} \in \Omega_{\Sigma} + n\mathbf{v}_2} F_{\omega_0}(\tilde{\mathbf{x}}) \int_{\tilde{\mathbf{y}} \in \mathbb{R}^2} \mathcal{K}_{\#}^{\lambda}(\tilde{\mathbf{x}}, \tilde{\mathbf{y}} + (n - n')\mathbf{v}_2; \Omega, k_{\parallel}) G_{\omega'_0}(\tilde{\mathbf{y}}) d\tilde{\mathbf{y}} d\tilde{\mathbf{x}} \\ &= \sum_{n \in \mathbb{Z}} \int_{\tilde{\mathbf{x}} \in \Omega_{\Sigma} + n\mathbf{v}_2} F_{\omega_0}(\tilde{\mathbf{x}}) \sum_{n' \in \mathbb{Z}} \int_{\tilde{\mathbf{y}} \in \mathbb{R}^2} \mathcal{K}_{\#}^{\lambda}(\tilde{\mathbf{x}}, \tilde{\mathbf{y}} + (n - n')\mathbf{v}_2; \Omega, k_{\parallel}) G_{\omega'_0}(\tilde{\mathbf{y}}) d\tilde{\mathbf{y}} d\tilde{\mathbf{x}} \\ &= \int_{\tilde{\mathbf{x}} \in \mathbb{R}^2} F_{\omega_0}(\tilde{\mathbf{x}}) \int_{\tilde{\mathbf{y}} \in \mathbb{R}^2} \sum_{l \in \mathbb{Z}} \mathcal{K}_{\#}^{\lambda}(\tilde{\mathbf{x}}, \tilde{\mathbf{y}} + l\mathbf{v}_2; \Omega, k_{\parallel}) G_{\omega'_0}(\tilde{\mathbf{y}}) d\tilde{\mathbf{y}} d\tilde{\mathbf{x}} \\ &= \sum_{l \in \mathbb{Z}} \int_{\tilde{\mathbf{x}} \in \mathbb{R}^2} F_{\omega_0}(\tilde{\mathbf{x}}) \int_{\tilde{\mathbf{y}} \in \mathbb{R}^2} \mathcal{K}_{\#}^{\lambda}(\tilde{\mathbf{x}}, \tilde{\mathbf{y}} + l\mathbf{v}_2; \Omega, k_{\parallel}) G_{\omega'_0}(\tilde{\mathbf{y}}) d\tilde{\mathbf{y}} d\tilde{\mathbf{x}} \end{aligned}$$

This completes the proof of part (1) of Proposition 12.1. To prove part (2) of Proposition 12.1, we bound the expression in (12.5). Write $\mathcal{K}_{\#}^{\lambda, \omega_0, \omega'_0, l}$ for the operator:

$$(12.8) \quad (\mathcal{K}_{\#}^{\lambda, \omega_0, \omega'_0, l} f)(\mathbf{x}) = \int_{\mathbb{R}^2} e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} \mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y} + l\mathbf{v}_2) e^{-\frac{\gamma}{2}|\mathbf{y}-\omega'_0|} f(\mathbf{y}) d\mathbf{y}.$$

Then, by part (1) of Proposition 12.1, we have

$$\begin{aligned}
& \left| \left\langle F_{[\omega]}, \Pi_{AB}(k_{\parallel}) \mathcal{K}_{\#}^{\lambda}(\Omega, k_{\parallel}) \Pi_{AB}(k_{\parallel}) G_{[\omega']} \right\rangle_{L^2(\Sigma)} \right| \\
& \leq \sum_{l \in \mathbb{Z}} \left| \int_{\mathbf{x} \in \mathbb{R}^2} \left[e^{\frac{\gamma}{2}|\mathbf{x}-\omega_0|} F(\mathbf{x} - \omega_0) \right] \right. \\
& \quad \left. \int_{\mathbf{y} \in \mathbb{R}^2} \left[e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} \mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y} + l\mathbf{v}_2; \Omega, k_{\parallel}) e^{-\frac{\gamma}{2}|\mathbf{y}-\omega_0|} \right] \left[e^{-\frac{\gamma}{2}|\mathbf{y}-\omega'_0|} G_{\omega'_0}(\mathbf{y}) \right] dy dx \right| \\
& \leq \sum_{l \in \mathbb{Z}} \|F_{\omega_0}\|_{\mathcal{H}(\omega_0)} \|\mathcal{K}_{\#}^{\lambda, \omega_0, \omega'_0, l}\|_{L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2)} \|G_{\omega'_0}\|_{\mathcal{H}(\omega'_0)} \\
& = \left[\sum_{l \in \mathbb{Z}} \|\mathcal{K}_{\#}^{\lambda, \omega_0, \omega'_0, l}\|_{L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2)} \right] \|F_{\omega_0}\|_{\mathcal{H}(\omega_0)} \|G_{\omega'_0}\|_{\mathcal{H}(\omega'_0)}.
\end{aligned}$$

This completes the proof of part (2) of Proposition 12.1.

We shall apply conclusion (2) of Proposition 12.1 with $F_{[\omega]} = H_{\#}^{\lambda} p_{k_{\parallel}, J}^{\lambda}[n]$ and $G_{[\omega']} = \Pi_{AB}^{\lambda}(k_{\parallel}) H_{\#}^{\lambda} p_{k_{\parallel}, I}^{\lambda}[m]$, $J, I \in \{A, B\}$. Two more tasks remain in this section:

- (1) Bound the sum of norms on the right hand side of (12.6) using our pointwise kernel bounds, (10.107), on $\mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega, k_{\parallel})$, and
- (2) Bound $\|F_{\omega_0}\|_{\mathfrak{H}(\omega_0)}$ and $\|G_{\omega'_0}\|_{\mathfrak{H}(\omega'_0)}$, where $F_{\omega_0} = H_{\#}^{\lambda} p_{\omega_0}^{\lambda}$ and $G_{\omega'_0} = H_{\#}^{\lambda} p_{\omega'_0}^{\lambda}$.

This will enable us to bound the nonlinear contributions to matrix $\mathcal{M}[m, n](\Omega, k_{\parallel})$, displayed in (12.2), thereby proving Proposition 7.2.

The following two propositions will do the trick:

Proposition 12.2. *Let ω_0 and ω'_0 be as in the statement of Proposition 12.1. There exist constants $\lambda_1 > 0$ and $c > 0$ such that for all $\lambda \geq \lambda_1$ and $|\Omega| \leq e^{-c\lambda}$:*

$$(12.9) \quad \sum_{l \in \mathbb{Z}} \|\mathcal{K}_{\#}^{\lambda; \omega_0, \omega'_0, l}(\Omega, k_{\parallel})\|_{L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2)} \lesssim \lambda^{10} e^{-c|\omega_0 - \omega'_0|}.$$

Proposition 12.3. *Then,*

$$\|H_{\#}^{\lambda} p_{\omega_0}^{\lambda}\|_{\mathfrak{H}(\omega_0)} \leq e^{-c\lambda} \sqrt{\rho\lambda} \quad \text{and} \quad \|H_{\#}^{\lambda} p_{\omega'_0}^{\lambda}\|_{\mathfrak{H}(\omega'_0)} \lesssim e^{-c\lambda} \sqrt{\rho\lambda}.$$

The proofs of Propositions 12.2 and 12.3 are presented in the following two subsections. We first apply them to conclude the proof of Proposition 7.2, which gives our bound on nonlinear matrix elements. Estimate (12.6) with $F_{\omega_0} = H_{\#}^{\lambda} p_{\omega_0}^{\lambda}$ and $G_{\omega'_0} = H_{\#}^{\lambda} p_{\omega'_0}^{\lambda}$ implies

$$\begin{aligned}
(12.10) \quad & \left| \left\langle H_{\#}^{\lambda} p_{k_{\parallel}, J}^{\lambda}[n], \Pi_{AB}^{\lambda}(k_{\parallel}) \mathcal{K}_{\#}^{\lambda}(\Omega, k_{\parallel}) \Pi_{AB}^{\lambda}(k_{\parallel}) H_{\#}^{\lambda} p_{k_{\parallel}, I}^{\lambda}[m] \right\rangle_{L^2(\Sigma)} \right| \\
& \leq \left[\sum_{l \in \mathbb{Z}} \|\mathcal{K}_{\#}^{\lambda; \omega_0, \omega'_0, l}\|_{L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2)} \right] \cdot \|H_{\#}^{\lambda} p_{\omega_0}^{\lambda}\|_{\mathfrak{H}(\omega_0)} \cdot \|H_{\#}^{\lambda} p_{\omega'_0}^{\lambda}\|_{\mathfrak{H}(\omega'_0)}.
\end{aligned}$$

Now apply Propositions 12.2 and 12.3 to obtain

$$\begin{aligned}
(12.11) \quad & \left| \left\langle H_{\#}^{\lambda} p_{k_{\parallel}, J}^{\lambda}[n], \Pi_{AB}^{\lambda}(k_{\parallel}) \mathcal{K}_{\#}(\Omega, k_{\parallel}) \Pi_{AB}^{\lambda}(k_{\parallel}) H_{\#}^{\lambda} p_{k_{\parallel}, I}^{\lambda}[m] \right\rangle_{L^2(\Sigma)} \right| \\
& \lesssim \lambda^{10} e^{-c|\omega_0 - \omega'_0|} \cdot e^{-c\lambda} \sqrt{\rho_{\lambda}} \cdot e^{-c\lambda} \sqrt{\rho_{\lambda}} \\
& \lesssim \rho_{\lambda} e^{-c\lambda} e^{-c|\omega_0 - \omega'_0|}.
\end{aligned}$$

We have proved Proposition 7.2 for the case $j = 0$. From this, the case $j = 1$ follows by analytic dependence of the inner product on Ω ; see the remark just prior to the statement of Proposition 7.2. This completes the proof of Proposition 7.2.

12.0.1. *Proof of Proposition 12.2:* From the expression for the integral kernel, displayed in (12.7), we have

$$\begin{aligned}
(12.12) \quad & \|\mathcal{K}_{\#}^{\lambda, \omega_0, \omega'_0, l}\|_{L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2)} \leq \sup_{\mathbf{x} \in \mathbb{R}^2} \int_{\tilde{\mathbf{y}} \in \mathbb{R}^2} e^{-\frac{\gamma}{2}|\mathbf{x} - \omega_0|} |\mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \tilde{\mathbf{y}} + l\mathbf{v}_2)| e^{-\frac{\gamma}{2}|\tilde{\mathbf{y}} - \omega'_0|} d\tilde{\mathbf{y}} \\
& \quad + \sup_{\tilde{\mathbf{y}} \in \mathbb{R}^2} \int_{\mathbf{x} \in \mathbb{R}^2} e^{-\frac{\gamma}{2}|\mathbf{x} - \omega_0|} |\mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \tilde{\mathbf{y}} + l\mathbf{v}_2)| e^{-\frac{\gamma}{2}|\tilde{\mathbf{y}} - \omega'_0|} d\mathbf{x} \\
& = \sup_{\mathbf{x} \in \mathbb{R}^2} \int_{\tilde{\mathbf{y}} \in \mathbb{R}^2} e^{-\frac{\gamma}{2}|\mathbf{x} - \omega_0|} |\mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y})| e^{-\frac{\gamma}{2}|\mathbf{y} - l\mathbf{v}_2 - \omega'_0|} d\mathbf{y} \\
& \quad + \sup_{\tilde{\mathbf{y}} \in \mathbb{R}^2} \int_{\mathbf{x} \in \mathbb{R}^2} e^{-\frac{\gamma}{2}|\mathbf{x} - \omega_0|} |\mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y})| e^{-\frac{\gamma}{2}|\mathbf{y} - l\mathbf{v}_2 - \omega'_0|} d\mathbf{x} \\
& = \sup_{\mathbf{x} \in \mathbb{R}^2} \mathcal{J}^{\lambda}(\mathbf{x}; l) + \sup_{\mathbf{y} \in \mathbb{R}^2} \mathcal{J}^{\lambda}(\mathbf{y}; l)
\end{aligned}$$

Recall that the kernel $\mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega, k_{\parallel})$ satisfies the pointwise bound (10.107):

$$(12.13) \quad \left| \mathcal{K}_{\#}^{\lambda}(\mathbf{x}, \mathbf{y}; \Omega, k_{\parallel}) \right| \leq C \left[|\log|\mathbf{x} - \mathbf{y}|| + \lambda^{10} \right] \mathbf{1}_{|\mathbf{x} - \mathbf{y}| \leq R} + e^{-c\lambda} e^{-c\lambda|\mathbf{x} - \mathbf{y}|}$$

for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$.

The bounds on $\sup_{\mathbf{x} \in \mathbb{R}^2} \mathcal{J}^{\lambda}(\mathbf{x}; l)$ and $\sup_{\mathbf{y} \in \mathbb{R}^2} \mathcal{J}^{\lambda}(\mathbf{y}; l)$ are obtained very similarly. We present the argument for $\sup_{\mathbf{x} \in \mathbb{R}^2} \mathcal{J}^{\lambda}(\mathbf{x}; l)$. To bound $\mathcal{J}^{\lambda}(\mathbf{x}; l)$, we bound the $d\mathbf{y}$ integral over \mathbb{R}^2 separately over the sets $|\mathbf{x} - \mathbf{y}| \leq R$ and $|\mathbf{x} - \mathbf{y}| \geq R$. Call these parts: $\mathcal{J}_{\leq R}^{\lambda}(\mathbf{x}; l)$ and $\mathcal{J}_{\geq R}^{\lambda}(\mathbf{x}; l)$.

First assume $|\mathbf{x} - \mathbf{y}| \leq R$. By (12.13)

$$\begin{aligned}
\mathcal{J}_{\leq R}^\lambda(\mathbf{x}; l) &\leq e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} \int_{|\mathbf{x}-\mathbf{y}| \leq R} |\mathcal{K}_\#^\lambda(\mathbf{x}, \mathbf{y}; \Omega, k_\parallel)| e^{-\frac{\gamma}{2}|\mathbf{y}-l\mathbf{v}_2-\omega'_0|} d\mathbf{y} \\
&\lesssim e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} \int_{|\mathbf{x}-\mathbf{y}| \leq R} \left[|\log |\mathbf{x} - \mathbf{y}| | + \lambda^{10} \right] e^{-\frac{\gamma}{2}|\mathbf{y}-l\mathbf{v}_2-\omega'_0|} d\mathbf{y} \\
&\lesssim e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} \int_{|\mathbf{z}| \leq R} \left[|\log |\mathbf{z}| | + \lambda^{10} \right] e^{-\frac{\gamma}{2}|\mathbf{x}-\mathbf{z}-l\mathbf{v}_2-\omega'_0|} d\mathbf{z} \\
&\lesssim e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} \int_{0 \leq |\mathbf{z}| < \rho} \left[|\log |\mathbf{z}| | + \lambda^{10} \right] e^{-\frac{\gamma}{2}|\mathbf{x}-\mathbf{z}-l\mathbf{v}_2-\omega'_0|} d\mathbf{z} \\
&\quad + e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} \int_{\rho \leq |\mathbf{z}| \leq R} \left[|\log |\mathbf{z}| | + \lambda^{10} \right] e^{-\frac{\gamma}{2}|\mathbf{x}-\mathbf{z}-l\mathbf{v}_2-\omega'_0|} d\mathbf{z} \\
&\lesssim e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} e^{-c_1|\mathbf{x}-l\mathbf{v}_2-\omega'_0|} \int_{0 \leq |\mathbf{z}| \leq \rho} \left[|\log |\mathbf{z}| | + \lambda^{10} \right] d\mathbf{z} \\
&\quad + e^{-\frac{\gamma}{2}|\mathbf{x}-\omega_0|} \left[C_{\rho, R} + \lambda^{10} \right] \int_{\rho \leq |\mathbf{z}| \leq R} e^{-\frac{\gamma}{2}|\mathbf{x}-\mathbf{z}-l\mathbf{v}_2-\omega'_0|} d\mathbf{z} .
\end{aligned}$$

The latter two terms are each $\lesssim \lambda^{10} e^{-c_2|\omega_0-\omega'_0|} e^{-c_3|l|}$. Therefore,

$$(12.14) \quad \sup_{\mathbf{x} \in \mathbb{R}^2} \mathcal{J}_{\leq R}^\lambda(\mathbf{x}; l) \lesssim \lambda^{10} e^{-c_2|\omega_0-\omega'_0|} e^{-c_3|l|} .$$

A similar argument yields a bound of this type for $\sup_{\mathbf{x} \in \mathbb{R}^2} \mathcal{J}_{\leq R}^\lambda(\mathbf{x}; l)$.

Next assume $|\mathbf{x} - \mathbf{y}| \geq R$. By (12.13),

$$\mathcal{J}_{\geq R}^\lambda(\mathbf{x}; l) \lesssim e^{-\frac{\gamma}{2}\lambda} e^{-c|\mathbf{x}-\omega_0|} \int_{|\mathbf{x}-\mathbf{y}| \geq R} e^{-c\lambda|\mathbf{x}-\mathbf{y}|} e^{-\frac{\gamma}{2}|\mathbf{y}-l\mathbf{v}_2-\omega'_0|} d\mathbf{y} .$$

Note that $|\mathbf{x} - \omega_0| + |\mathbf{y} - l\mathbf{v}_2 - \omega'_0| \geq |(\mathbf{x} - \omega_0) - (\mathbf{y} - l\mathbf{v}_2 - \omega'_0)| = |\mathbf{x} - \mathbf{y} - (\omega_0 - \omega'_0) + l\mathbf{v}_2| \geq c_3(|\omega_0 - \omega'_0| + |l|) - |\mathbf{x} - \mathbf{y}|$. Thus,

$$(12.15) \quad \mathcal{J}_{\geq R}^\lambda(\mathbf{x}; l) \lesssim e^{-c\lambda} \int_{|\mathbf{x}-\mathbf{y}| \geq R} e^{-c_4\lambda|\mathbf{x}-\mathbf{y}|} d\mathbf{y} e^{-c_3|\omega_0-\omega'_0|} e^{-c_3|l|} \lesssim e^{-c\lambda} e^{-c_3|\omega_0-\omega'_0|} e^{-c_3|l|} .$$

The bounds (12.14) and (12.15) imply that

$$\sup_{\mathbf{x} \in \mathbb{R}^2} \mathcal{J}^\lambda(\mathbf{x}; l) \leq e^{-c_3|\omega_0-\omega'_0|} e^{-c_3|l|} \lambda^{10} .$$

and similarly

$$\sup_{\mathbf{y} \in \mathbb{R}^2} \mathcal{J}^\lambda(\mathbf{y}; l) \leq e^{-c_3|\omega_0-\omega'_0|} e^{-c_3|l|} \lambda^{10} .$$

Therefore, by (12.12) it follows that $\|\mathcal{K}_\#^{\lambda; \omega_0, \omega'_0, l}\|_{L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2)} \lesssim e^{-c_3|\omega_0-\omega'_0|} e^{-c_3|l|} \lambda^{10}$. Finally, summing over $l \in \mathbb{Z}$ we deduce (12.9). The proof of Proposition 12.2 is now complete.

12.0.2. *Proof of Proposition 12.3.* We need to verify that there are constants $\gamma, \lambda_1 > 0$, such that for all $\omega \in \mathbb{H}_\#$ and all $\lambda \geq \lambda_1$:

$$(12.16) \quad \|\mathcal{H}_\#^\lambda p_\omega^\lambda\|_{\mathcal{H}(\omega)} = \|e^{\frac{\gamma}{2}|\mathbf{x}-\omega|} (-\Delta + V_\#^\lambda(\mathbf{x}) - E_0^\lambda) p_\omega^\lambda(\mathbf{x})\|_{L^2(\mathbb{R}_\#^2)} \lesssim e^{-c\lambda} \sqrt{\rho_\lambda}.$$

Since $(-\Delta + \lambda^2 V_\omega(\mathbf{x})) p_\omega^\lambda(\mathbf{x}) = E_0^\lambda p_\omega^\lambda(\mathbf{x})$, it follows that

$$\mathcal{H}_\#^\lambda p_\omega^\lambda(\mathbf{x}) \equiv (-\Delta + V_\#^\lambda(\mathbf{x}) - E_0^\lambda) p_\omega^\lambda(\mathbf{x}) = \sum_{\omega' \in \mathbb{H}_\# \setminus \{\omega\}} \lambda^2 V_0(\mathbf{x} - \omega') p_\omega^\lambda(\mathbf{x}).$$

By invariance of $H_\#^\lambda$ under translation by \mathbf{v}_2 , we may assume $\omega \in \Omega_\Sigma$. Thus, $\omega = \mathbf{v}_I + n\mathbf{v}_1$ for $I = A$ or B and $n \geq 0$. Fix $I = A$; the argument for $I = B$ is similar. Then, $p_\omega^\lambda(\mathbf{x}) = p_0^\lambda(\mathbf{x} - \mathbf{v}_A - n\mathbf{v}_1)$. Recall, for $I = A, B$ and $n_1, n_2 \in \mathbb{Z}$: $\mathbf{v}_I^{n_1, n_2} = \mathbf{v}_I + n_1\mathbf{v}_1 + n_2\mathbf{v}_2$. Thus,

$$\begin{aligned} H_\#^\lambda p_\omega^\lambda(\mathbf{x}) &= \sum_{n_1 \geq 0, n_2 \in \mathbb{Z}} \lambda^2 V_0(\mathbf{x} - \mathbf{v}_B^{n_1, n_2}) p_0^\lambda(\mathbf{x} - \mathbf{v}_A - n\mathbf{v}_1) \\ &\quad + \sum_{\substack{n_1 \geq 0, n_2 \in \mathbb{Z} \\ (n_1, n_2) \neq (n, 0)}} \lambda^2 V_0(\mathbf{x} - \mathbf{v}_A^{n_1, n_2}) p_0^\lambda(\mathbf{x} - \mathbf{v}_A - n\mathbf{v}_1). \end{aligned}$$

For the $\mathcal{H}(\omega)$ norm ($\omega = \mathbf{v}_I + n\mathbf{v}_1$) we have

$$\begin{aligned} (12.17) \quad \|\mathcal{H}_\#^\lambda p_\omega^\lambda\|_{\mathcal{H}(\omega)} &= \left\| e^{\frac{\gamma}{2}|\mathbf{x}-\omega|} H_\#^\lambda p_\omega^\lambda(\mathbf{x}) \right\|_{L^2(\mathbb{R}_\#^2)} \\ &\leq \lambda^2 \sum_{n_1 \geq 0, n_2 \in \mathbb{Z}} \left(\int e^{\gamma|\mathbf{x}-(\mathbf{v}_A+n\mathbf{v}_1)|} |V_0(\mathbf{x} - \mathbf{v}_B^{n_1, n_2})|^2 |p_0^\lambda(\mathbf{x} - \mathbf{v}_A - n\mathbf{v}_1)|^2 d\mathbf{x} \right)^{\frac{1}{2}} \\ &\quad + \lambda^2 \sum_{\substack{n_1 \geq 0, n_2 \in \mathbb{Z} \\ (n_1, n_2) \neq (n, 0)}} \left(\int e^{\gamma|\mathbf{x}-(\mathbf{v}_A+n\mathbf{v}_1)|} |V_0(\mathbf{x} - \mathbf{v}_A^{n_1, n_2})|^2 |p_0^\lambda(\mathbf{x} - \mathbf{v}_A - n\mathbf{v}_1)|^2 d\mathbf{x} \right)^{\frac{1}{2}} \\ &\equiv \sum_{n_1 \geq 0, n_2 \in \mathbb{Z}} A_{n_1, n_2}^\lambda + \sum_{\substack{n_1 \geq 0, n_2 \in \mathbb{Z} \\ (n_1, n_2) \neq (n, 0)}} B_{n_1, n_2}^\lambda. \end{aligned}$$

Consider A_{n_1, n_2}^λ , for any fixed $n_1 \geq 0$ and $n_2 \in \mathbb{Z}$.

$$\begin{aligned} |A_{n_1, n_2}^\lambda|^2 &= \lambda^4 \int_{|\mathbf{x}-\mathbf{v}_B^{n_1, n_2}| \leq r_0} e^{\gamma|\mathbf{x}-(\mathbf{v}_A+n\mathbf{v}_1)|} |V_0(\mathbf{x} - \mathbf{v}_B^{n_1, n_2})|^2 |p_0^\lambda(\mathbf{x} - \mathbf{v}_A - n\mathbf{v}_1)|^2 d\mathbf{x} \\ &= \lambda^4 \int_{|\mathbf{y}| \leq r_0} e^{\gamma|\mathbf{y}+\mathbf{v}_B^{n_1, n_2}-\mathbf{v}_A^0|} |V_0(\mathbf{y})|^2 |p_0^\lambda(\mathbf{y} + \mathbf{v}_B^{n_1, n_2} - \mathbf{v}_A^0)|^2 d\mathbf{y} \\ &= \lambda^4 \int_{|\mathbf{y}| \leq r_0} e^{\gamma|\mathbf{y}+\mathbf{v}_B-\mathbf{v}_A+(n_1-n)\mathbf{v}_1+n_2\mathbf{v}_2|} |V_0(\mathbf{y})|^2 |p_0^\lambda(\mathbf{y} + \mathbf{v}_B - \mathbf{v}_A + (n_1 - n)\mathbf{v}_1 + n_2\mathbf{v}_2)|^2 d\mathbf{y} \\ &= \lambda^4 \int_{|\mathbf{y}| \leq r_0} e^{\gamma|\mathbf{y}+\mathbf{e}+(n_1-n)\mathbf{v}_1+n_2\mathbf{v}_2|} |V_0(\mathbf{y})|^2 |p_0^\lambda(\mathbf{y} - [-\mathbf{e} + (n - n_1)\mathbf{v}_1 - n_2\mathbf{v}_2])|^2 d\mathbf{y}. \end{aligned}$$

As in Section 11 we divide index pairs $(n - n_1, -n_2)$ into those in the set $N_b(-1) = \{(0, 0), (1, 0), (0, 1)\}$ and those not in $N_b(-1)$. Those in $N_b(-1)$, “bad index pairs”, correspond to the cases: (i) $(n_1, n_2) = (n - 1, 0)$ with $n \geq 1$, (ii) $(n_1, n_2) = (n, 0)$ with $n \geq 0$ or (iii) $(n_1, n_2) = (n, -1)$ with $n \geq 0$. By the remark immediately following Definition 11.1, we then have for some $l = 0, 1$ or 2

$$p_0^\lambda(\mathbf{y} - [-\mathbf{e} + (n - n_1)\mathbf{v}_1 - n_2\mathbf{v}_2]) = p_0^\lambda(\mathbf{y} - [-R^l\mathbf{e}]),$$

where R is a $2\pi/3$ rotation matrix. Therefore, by orthogonality of the matrix R and symmetry assumptions on V_0 , we have:

$$\begin{aligned} |A_{n_1, n_2}^\lambda|^2 &= \lambda^4 \int_{|\mathbf{y}| \leq r_0} e^{2c|\mathbf{y} - [-R^l\mathbf{e}]|} |V_0(\mathbf{y})|^2 |p_0^\lambda(\mathbf{y} - [-R^l\mathbf{e}])|^2 d\mathbf{y} \\ &= \lambda^4 \int_{|\mathbf{y}| \leq r_0} e^{2c|R^{-l}\mathbf{y} + \mathbf{e}|} |V_0(R^{-l}\mathbf{y})|^2 |p_0^\lambda(R^{-l}\mathbf{y} + \mathbf{e})|^2 d\mathbf{y} \\ &= \lambda^4 \int_{|\mathbf{z}| \leq r_0} e^{2c|\mathbf{z} + \mathbf{e}|} |V_0(\mathbf{z})|^2 |p_0^\lambda(\mathbf{z} + \mathbf{e})|^2 d\mathbf{z}. \end{aligned}$$

Next, applying the bound (11.11) to one factor of $p_0^\lambda(\mathbf{z} + \mathbf{e})$ yields

$$\begin{aligned} |A_{n_1, n_2}^\lambda|^2 &\lesssim \lambda^4 \|V_0\|_\infty \int_{|\mathbf{z}| \leq r_0} e^{2c|\mathbf{z} + \mathbf{e}| - c\lambda} |V_0(\mathbf{z})| p_0^\lambda(\mathbf{z}) p_0^\lambda(\mathbf{z} + \mathbf{e}) d\mathbf{z} \\ &\lesssim e^{-c'\lambda} \rho_\lambda. \end{aligned}$$

Next consider $n_1 \geq 0$ and $n_2 \in \mathbb{Z}$, for which $(n - n_1, -n_2) \notin N_{\text{bad}}(-1)$. By Proposition 11.3, in particular (11.10), we have

$$(12.18) \quad p_0^\lambda(\mathbf{y} - [-\mathbf{e} + (n - n_1)\mathbf{v}_1 - n_2\mathbf{v}_2]) \lesssim e^{-c\lambda(|n - n_1| + |n_2|)} p_0^\lambda(\mathbf{y} + \mathbf{e}).$$

Therefore, for $|\mathbf{y}| \leq r_0$ and λ sufficiently large:

$$\begin{aligned} (12.19) \quad &e^{\gamma|\mathbf{y} - [-\mathbf{e} + (n - n_1)\mathbf{v}_1 - n_2\mathbf{v}_2]|} p_0^\lambda(\mathbf{y} - [-\mathbf{e} + (n - n_1)\mathbf{v}_1 - n_2\mathbf{v}_2]) \\ &\lesssim e^{\gamma|\mathbf{y} - [-\mathbf{e} + (n - n_1)\mathbf{v}_1 - n_2\mathbf{v}_2]|} e^{-c\lambda(|n_1 - n| + |n_2|)} p_0^\lambda(\mathbf{y} + \mathbf{e}) \\ &\lesssim e^{-c'\lambda(|n_1 - n| + |n_2|)} p_0^\lambda(\mathbf{y} + \mathbf{e}) \lesssim e^{-c'\lambda(|n_1 - n| + |n_2|)} p_0^\lambda(\mathbf{y}), \end{aligned}$$

where the last inequality uses (11.11). Therefore, for good index pairs $(n - n_1, -n_2)$ we have

$$\begin{aligned} |A_{n_1, n_2}^\lambda|^2 &\lesssim \lambda^4 \|V_0\|_\infty e^{-c\lambda(|n - n_1| + |n_2|)} \int |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y}) p_0^\lambda(\mathbf{y} + \mathbf{e}) d\mathbf{y} \\ &\lesssim e^{-c\lambda(|n - n_1| + |n_2|)} \rho_\lambda. \end{aligned}$$

Taking the square root and summing over good index pairs (n_1, n_2) we have:

$$(12.20) \quad \sum_{\substack{n_1, n_2 \\ (n - n_1, -n_2) \text{ good}}} A_{n_1, n_2}^\lambda \lesssim e^{-c\lambda} \sqrt{\rho_\lambda}.$$

Taken together with our bound on $|A_{n_1, n_2}^\lambda|$ for the three cases of bad indices, this tells us that

$$(12.21) \quad \sum_{n_1 \geq 0, n_2 \in \mathbb{Z}} A_{n_1, n_2}^\lambda \lesssim e^{-c\lambda} \sqrt{\rho_\lambda}.$$

The proof that

$$(12.22) \quad \sum_{n_1 \geq 0, n_2 \in \mathbb{Z}} B_{n_1, n_2}^\lambda \lesssim e^{-c\lambda} \sqrt{\rho\lambda}.$$

is similar, so this completes the proof of (12.3).

APPENDIX A. ERROR AND MAIN KERNELS; PROOF OF LEMMA 10.7

We prove that if \mathcal{E} is an operator derived from an error kernel $\mathcal{E}(\mathbf{x}, \mathbf{y})$ in the sense of Definition 10.5, then $\tilde{\mathcal{E}} = I - (I - \mathcal{E})^{-1}$ is an operator derived from an error kernel $\tilde{\mathcal{E}}(\mathbf{x}, \mathbf{y})$.

A.1. Elementary integrals in 1d. Let $f \in L^1(\mathbb{R})$. We define $f^{*0} = \delta$, the Dirac delta function and $f^{*1} = f$. Let f^{*n} denote the n -fold convolution of f with itself:

For f and g in $L^1(\mathbb{R})$,

$$(A.1) \quad (f + g)^{*n} = \sum_{k=0}^n \binom{n}{k} f^{*k} g^{*(n-k)}.$$

Let $f(t) = ae^{-\gamma|t|}$, where a and γ are positive constants with $\gamma > a$. We may write

$$f(t) = f_+(t) + f_-(t), \quad f_+(t) = ae^{-\gamma t} \mathbf{1}_{\{t>0\}}, \quad f_-(t) = ae^{-\gamma|t|} \mathbf{1}_{\{t<0\}}.$$

Induction on k gives:

$$f_+^{*k}(t) = a^k e^{-\gamma t} \frac{t^{k-1}}{(k-1)!} \mathbf{1}_{\{t>0\}} \leq ae^{-\gamma t} \sum_{l=0}^{\infty} \frac{(at)^l}{l!} \mathbf{1}_{\{t>0\}} = ae^{-(\gamma-a)t} \mathbf{1}_{\{t>0\}}, \quad k \geq 1.$$

A similar bound holds for f_- . Therefore, for all $0 < a < \gamma$:

$$f_+^{*k}(t) \leq a e^{-(\gamma-a)t} \mathbf{1}_{\{t>0\}} \quad \text{and} \quad f_-^{*k}(t) \leq a e^{-(\gamma-a)|t|} \mathbf{1}_{\{t<0\}}, \quad k \geq 1.$$

Therefore, for $m \geq 1$, we have from (A.1) that

$$(A.2) \quad \begin{aligned} f^{*m}(t) &= (f_+ + f_-)^{*m}(t) \leq a^2 \sum_{k=1}^{m-1} \binom{m}{k} [e^{-(\gamma-a)t} \mathbf{1}_{\{t>0\}} \star e^{-(\gamma-a)|t|} \mathbf{1}_{\{t<0\}}] (t) \\ &\quad + ae^{-(\gamma-a)t} \mathbf{1}_{\{t>0\}} + ae^{-(\gamma-a)|t|} \mathbf{1}_{\{t<0\}}. \end{aligned}$$

The last two terms, which sum to $ae^{-(\gamma-a)|t|}$, correspond to $k=0$ and $k=m$ in the binomial formula. We calculate the convolution in (A.2). For $t > 0$,

$$\begin{aligned} &[e^{-(\gamma-a)|t|} \mathbf{1}_{\{t>0\}} \star e^{-(\gamma-a)|t|} \mathbf{1}_{\{t<0\}}] (t) \\ &= \int_0^\infty e^{-(\gamma-a)s} e^{-(\gamma-a)|t-s|} \mathbf{1}_{\{t-s<0\}} ds = \int_t^\infty e^{-2(\gamma-a)s} e^{(\gamma-a)t} ds = \frac{e^{-(\gamma-a)t}}{2(\gamma-a)}. \end{aligned}$$

Similarly, if $t < 0$ then this convolution is $\frac{e^{-(\gamma-a)|t|}}{2(\gamma-a)}$. Therefore,

$$[e^{-(\gamma-a)|t|} \mathbf{1}_{\{t>0\}} \star e^{-(\gamma-a)|t|} \mathbf{1}_{\{t<0\}}] (t) = \frac{e^{-(\gamma-a)|t|}}{2(\gamma-a)}, \quad \text{for all } t \in \mathbb{R}.$$

Substituting into (A.2) we have

$$f^{*m}(t) = (ae^{-\gamma|t|})^{*m}(t) \leq a^2 \frac{e^{-(\gamma-a)|t|}}{2(\gamma-a)} \sum_{k=1}^{m-1} \binom{m}{k} + ae^{-(\gamma-a)|t|} \leq \left[a + \frac{2^m a^2}{2(\gamma-a)} \right] e^{-(\gamma-a)|t|}.$$

Therefore,

$$(A.3) \quad \left(\frac{a}{4} e^{-\gamma|t|} \right)^{*m} (t) \leq \left[4^{-m} a + \frac{2^{-m} a^2}{2(\gamma - a)} \right] e^{-(\gamma-a)|t|} \quad \text{for } m \geq 1.$$

A.2. Elementary integrals in n dimensions. For $(x_1, \dots, x_n) \in \mathbb{R}^n$, let

$$K(x_1, \dots, x_n) = \frac{a^n}{4^n} e^{-\gamma(|x_1| + \dots + |x_n|)} \quad \text{with } 0 < a < \gamma.$$

We now apply (A.3) to the l -fold convolution of $K(x_1, \dots, x_n)$:

$$(A.4) \quad \begin{aligned} K^{*l}(x_1, \dots, x_n) &\equiv \underbrace{K \star K \star \dots \star K}_{l \text{ times}}(x_1, \dots, x_n) \leq \prod_{j=1}^n \left\{ \left[4^{-l} a + \frac{2^{-l} a^2}{2(\gamma - a)} \right] e^{-(\gamma-a)|x_j|} \right\} \\ &= \left[4^{-l} a + \frac{2^{-l} a^2}{2(\gamma - a)} \right]^n e^{-(\gamma-a)(|x_1| + \dots + |x_n|)}. \end{aligned}$$

A.3. Proof of part (1) of Lemma 10.7. For $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$ we write $|\mathbf{x}|_{l^1}$ to denote $|x_1| + \dots + |x_n|$. Suppose that $E(\mathbf{x}, \mathbf{y})$ satisfies the bound:

$$(A.5) \quad |E(\mathbf{x}, \mathbf{y})| \leq (a/4)^n e^{-\gamma|\mathbf{x}-\mathbf{y}|_{l^1}}, \quad \text{for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^n$$

and gives rise to the integral operator:

$$(A.6) \quad (Ef)(\mathbf{x}) = \int_{\mathbb{R}^n} E(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y},$$

then for all $l \geq 1$ the l^{th} power of the operator $E: f \mapsto E^l[f]$, is given by

$$E^l[f](\mathbf{x}) = \int_{\mathbb{R}^n} E_l(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y},$$

where by (A.4), E_l satisfies the bound

$$|E_l(\mathbf{x}, \mathbf{y})| \leq \left[4^{-l} a + \frac{2^{-l} a^2}{2(\gamma - a)} \right]^n e^{-(\gamma-a)|\mathbf{x}-\mathbf{y}|_{l^1}}.$$

If $\gamma > 2a$, then $\frac{a^2}{2(\gamma-a)} \leq \frac{a}{2}$. Therefore, for $l \geq 1$:

$$\left[4^{-l} a + \frac{2^{-l} a^2}{2(\gamma - a)} \right] \leq \left[4^{-l} a + 2^{-l} \frac{a}{2} \right] = 2^{-l} a [2^{-l} + 2^{-1}] \leq 2^{-l} a.$$

Hence,

$$|E_l(\mathbf{x}, \mathbf{y})| \leq 2^{-ln} a^n e^{-(\gamma-a)|\mathbf{x}-\mathbf{y}|_{l^1}}, \quad l \geq 1.$$

Let's now apply these observations to $E(\mathbf{x}, \mathbf{y}) = \mathcal{E}(\mathbf{x}, \mathbf{y})$, where $\mathcal{E}(\mathbf{x}, \mathbf{y})$ is an error kernel which by Definition 10.5 satisfies $|\mathcal{E}(\mathbf{x}, \mathbf{y})| \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{y}|}$ for $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$; here $n = 2$. Note that $e^{-c'\lambda|\mathbf{x}-\mathbf{y}|_{l^1}} \leq e^{-c\lambda|\mathbf{x}-\mathbf{y}|} \leq e^{-c'\lambda|\mathbf{x}-\mathbf{y}|_{l^1}}$. Therefore, $|\mathcal{E}(\mathbf{x}, \mathbf{y})| \lesssim e^{-c\lambda} e^{-c'\lambda|\mathbf{x}-\mathbf{y}|_{l^1}}$. It follows that $\mathcal{E}(\mathbf{x}, \mathbf{y})$ satisfies the bound (A.5) with $n = 2$, $(a/4)^2 = e^{-c\lambda}$ and $\gamma = c'\lambda$. Therefore, the operator \mathcal{E}^l is given by a kernel $\mathcal{E}_l(\mathbf{x}, \mathbf{y})$:

$$\mathcal{E}^l[f](\mathbf{x}) = \int_{\mathbb{R}^n} \mathcal{E}_l(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y},$$

where \mathcal{E}_l satisfies the bound

$$(A.7) \quad |\mathcal{E}_l(\mathbf{x}, \mathbf{y})| \leq 2^{-2l} e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{y}|}, \quad l \geq 1$$

for some $c > 0$, which is independent of l . Consequently, $f \mapsto \tilde{\mathcal{E}}f = (I - (I - \mathcal{E})^{-1})f = \sum_{l \geq 1} \mathcal{E}^l f$ is given by the kernel $\tilde{\mathcal{E}}(\mathbf{x}, \mathbf{y}) = \sum_{l \geq 1} \mathcal{E}_l(\mathbf{x}, \mathbf{y})$, which by (A.7) satisfies the bound $|\tilde{\mathcal{E}}(\mathbf{x}, \mathbf{y})| \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{y}|}$. Thus, $\tilde{\mathcal{E}}$ is an error kernel and

$$(A.8) \quad \tilde{\mathcal{E}}f(\mathbf{x}) = \int_{\mathbb{R}^2} \tilde{\mathcal{E}}(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y} .$$

The proof of part (1) of Lemma 10.7 is now complete.

A.3.1. *Proof of part (2) of Lemma 10.7.* We need to prove that if \mathcal{E}^λ derives from an error kernel and K^λ from a main kernel, then $K^\lambda \mathcal{E}^\lambda$ and $\mathcal{E}^\lambda K^\lambda$ derive from error kernels $(K^\lambda \mathcal{E}^\lambda)(\mathbf{x}, \mathbf{y})$ and $(\mathcal{E}^\lambda K^\lambda)(\mathbf{x}, \mathbf{y})$. We begin with the following bounds on $\mathcal{E}^\lambda(\mathbf{x}, \mathbf{z})$ and $K^\lambda(\mathbf{z}, \mathbf{y})$:

$$\begin{aligned} |\mathcal{E}^\lambda(\mathbf{x}, \mathbf{z})| &\lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{z}|} \\ |K^\lambda(\mathbf{z}, \mathbf{y})| &\lesssim [\lambda^4 + |\log|\mathbf{z}-\mathbf{y}||] \mathbf{1}_{\{|\mathbf{z}-\mathbf{y}| \leq R\}} + e^{-c\lambda} e^{-c\lambda|\mathbf{z}-\mathbf{y}|} . \end{aligned}$$

Thus,

$$\begin{aligned} |(\mathcal{E}^\lambda K^\lambda)(\mathbf{x}, \mathbf{y})| &\lesssim \int_{|\mathbf{z}-\mathbf{y}| \leq R} (\lambda^4 + |\log|\mathbf{z}-\mathbf{y}||) e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{z}|} d\mathbf{z} + \int_{|\mathbf{z}-\mathbf{y}| \geq R} e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{z}|} e^{-c\lambda|\mathbf{z}-\mathbf{y}|} d\mathbf{z} \\ &\lesssim e^{-c'\lambda} e^{-c'\lambda|\mathbf{x}-\mathbf{y}|} . \end{aligned}$$

Thus, $(\mathcal{E}^\lambda K^\lambda)(\mathbf{x}, \mathbf{y})$ is an error kernel. A similar bound shows that $(K^\lambda \mathcal{E}^\lambda)(\mathbf{x}, \mathbf{y})$ is an error kernel.

A.3.2. *Proof of part (3) of Lemma 10.7.* We show that if K_λ arises from a main kernel, then $e^{-c\lambda} K_\lambda^2$ arises from an error kernel. Since $K_\lambda(\mathbf{x}, \mathbf{y})$ is bounded by the sum of a first term: $\sim (\lambda^4 + |\log|\mathbf{x}-\mathbf{y}||) \mathbf{1}_{|\mathbf{x}-\mathbf{y}| < R}$ and a second term $\lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{y}|}$ (an error kernel), by part (2) we need only consider the contribution to $e^{-c\lambda} (K_\lambda^2)(\mathbf{x}, \mathbf{z}) = e^{-c\lambda} \int K_\lambda(\mathbf{x}, \mathbf{y}) K_\lambda(\mathbf{y}, \mathbf{z}) d\mathbf{y}$ arising from the first term. The size of this contribution is $\lesssim \lambda^8 e^{-c\lambda} \mathbf{1}_{|\mathbf{x}-\mathbf{z}| < 2R} \lesssim e^{-c'\lambda} \mathbf{1}_{|\mathbf{x}-\mathbf{z}| < 2R}$. Hence,

$$e^{-c\lambda} (K_\lambda^2)(\mathbf{x}, \mathbf{z}) \lesssim e^{-c'\lambda} \mathbf{1}_{|\mathbf{x}-\mathbf{z}| < 2R} + e^{-c\lambda} e^{-c\lambda|\mathbf{x}-\mathbf{z}|} \lesssim e^{-c''\lambda} e^{-c''\lambda|\mathbf{x}-\mathbf{z}|} .$$

Hence, $e^{-c\lambda} K_\lambda^2$ derives from an error kernel.

APPENDIX B. OVERLAP INTEGRALS; PROOF OF LEMMA 11.2

In this section we prove Lemma 11.2, which we restate here for convenience: For $I_1, J_1, \tilde{I}_1 \in \{A, B\}$, $m, n, n_1 \geq 0$ and $\tilde{m}_2 \in \mathbb{Z}$, consider the overlap integral

$$(B.1) \quad \mathcal{J}_\# \equiv \int p_0^\lambda(\mathbf{x} - \mathbf{v}_{I_1}^m - m_2 \mathbf{v}_2) \lambda^2 |V_0(\mathbf{x} - \mathbf{v}_{J_1}^{n_1})| p_0^\lambda(\mathbf{x} - \mathbf{v}_{\tilde{I}_1}^n - \tilde{m}_2 \mathbf{v}_2) d\mathbf{x} .$$

Note that the overlap integral in (B.1), although taken over \mathbb{R}^2 , has an integrand supported on the disc $B_{r_0}(\mathbf{v}_{J_1}^{n_1})$. Recall the hopping coefficient defined by:

$$\rho_\lambda = \int p_0^\lambda(\mathbf{y}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} - \mathbf{e}) d\mathbf{y} .$$

We also recall from Lemma 11.1 that for $I, J \in \{A, B\}$, we define $\sigma(I, J)$ so that: $\mathbf{v}_I - \mathbf{v}_J = \sigma(\mathbf{v}_B - \mathbf{v}_A) \equiv \sigma \mathbf{e}$. Thus, $\sigma(A, B) = -1$, $\sigma(B, A) = 1$, and $\sigma(A, A) = \sigma(B, B) = 0$.

Further, for $\sigma = +1, -1, 0$ we define $N_b(\sigma) = \{(r_1, r_2) : |\sigma \mathbf{e} + r_1 \mathbf{v}_1 + r_2 \mathbf{v}_2| = |\mathbf{e}|\}$. Hence, $N_b(+1) \equiv \{(0, 0), (-1, 0), (0, -1)\}$, $N_b(-1) \equiv \{(0, 0), (1, 0), (0, 1)\}$, and $N_b(0) \equiv \emptyset$.

Lemma 11.2 asserts the bound:

$$(B.2) \quad \mathcal{J}_\sharp \lesssim e^{-c\lambda(|m-n_1| + |m_2| + |n-n_1| + |\tilde{m}_2|)} \rho_\lambda,$$

except in the following cases of exceptional indices $(m, n, n_1, m_2, \tilde{m}_2)$:

- (a) $I_1 = \tilde{I}_1 = J_1$, $m = n = n_1$ and $m_2 = \tilde{m}_2 = 0$. This case does not arise in the proof of Proposition 7.1, so we say nothing further about it.
- (b) $\tilde{I}_1 = J_1$, $I_1 \neq J_1$, $(m - n_1, m_2) \in N_b(\sigma(I_1, J_1))$, $n = n_1$ and $\tilde{m}_2 = 0$, in which case $\mathcal{J}_\sharp = \rho_\lambda$.
- (c) $I_1 = J_1$, $\tilde{I}_1 \neq J_1$, $(n - n_1, \tilde{m}_2) \in N_b(\sigma(\tilde{I}_1, J_1))$, $m = n_1$ and $m_2 = 0$, in which case $\mathcal{J}_\sharp = \rho_\lambda$.

Lemma 11.2 further asserts that if $I_1 \neq J_1$, $\tilde{I}_1 \neq J_1$, then

$$(B.3) \quad \mathcal{J}_\sharp \lesssim e^{-c\lambda} e^{-c\lambda(|m-n_1| + |m_2| + |n-n_1| + |\tilde{m}_2|)} \rho_\lambda,$$

We shall occasionally use the notation: $\mathbf{m}\vec{\mathbf{v}} = m_1 \mathbf{v}_1 + m_2 \mathbf{v}_2$, where $\mathbf{m} = (m_1, m_2) \in \mathbb{Z}^2$.

To prove Lemma 11.2 we begin with a change of variables: $\mathbf{y} = \mathbf{x} - \mathbf{v}_{J_1}^{n_1}$. Therefore,

$$(B.4) \quad \mathcal{J}_\sharp \equiv \int p_0^\lambda(\mathbf{y} - [\sigma(I_1, J_1)\mathbf{e} + (m - n_1)\mathbf{v}_1 + m_2\mathbf{v}_2]) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} - [\sigma(\tilde{I}_1, J_1)\mathbf{e} + (n - n_1)\mathbf{v}_1 + \tilde{m}_2\mathbf{v}_2]) d\mathbf{y}.$$

Thus, our task is to consider integrals of the form

$$(B.5) \quad \mathcal{J} = \int p_0^\lambda(\mathbf{y} - [\sigma \mathbf{e} + r_1 \mathbf{v}_1 + r_2 \mathbf{v}_2]) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} - [\tilde{\sigma} \mathbf{e} + \tilde{r}_1 \mathbf{v}_1 + \tilde{r}_2 \mathbf{v}_2]) d\mathbf{y}.$$

Lemma B.1. *Consider the overlap integral (B.5), which depends on $\sigma, \tilde{\sigma} \in \{0, +1, -1\}$ and $\mathbf{r} = (r_1, r_2)$, $\tilde{\mathbf{r}} = (\tilde{r}_1, \tilde{r}_2) \in \mathbb{Z}^2$. The expression \mathcal{J} satisfies the bound:*

$$(B.6) \quad \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda(|r_1|+|r_2|+|\tilde{r}_1|+|\tilde{r}_2|)} \rho_\lambda$$

except in the following cases:

$$(\alpha) \quad \sigma = \tilde{\sigma} = 0, \mathbf{r} = \mathbf{0}, \tilde{\mathbf{r}} = \mathbf{0}.$$

This case does not arise in the proof of Proposition 7.1 so we say nothing further about it.

$$(\beta) \quad \tilde{\sigma} = 0, \sigma \neq 0, \mathbf{r} \in N_b(\sigma), \tilde{\mathbf{r}} = \mathbf{0}, \text{ in which case } \mathcal{J} = \rho_\lambda.$$

$$(\gamma) \quad \tilde{\sigma} \neq 0, \sigma = 0, \tilde{\mathbf{r}} \in N_b(\tilde{\sigma}), \mathbf{r} = \mathbf{0}, \text{ in which case } \mathcal{J} = \rho_\lambda.$$

We shall also make use of

Lemma B.2. *Suppose $\tilde{\sigma} \neq 0$ and $\sigma \neq 0$. Then,*

(1) *If $\mathbf{r} \in N_b(\sigma)$ and $\tilde{\mathbf{r}} \in N_b(\tilde{\sigma})$, then*

$$(B.7) \quad \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda} \rho_\lambda .$$

(2) *If $\mathbf{r} \in N_b(\sigma)$ and $\tilde{\mathbf{r}} \notin N_b(\tilde{\sigma})$, then*

$$(B.8) \quad \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda} e^{-c\lambda(|\tilde{r}_1|+|\tilde{r}_2|)} \rho_\lambda .$$

The analogous bound holds with \mathbf{r} and $\tilde{\mathbf{r}}$ interchanged.

(3) *If $\mathbf{r} \notin N_b(\sigma)$ and $\tilde{\mathbf{r}} \notin N_b(\tilde{\sigma})$ (and therefore $\mathbf{r}, \tilde{\mathbf{r}} \neq (0, 0)$), then*

$$(B.9) \quad \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c'\lambda} e^{-c'\lambda(|r_1|+|r_2|+|\tilde{r}_1|+|\tilde{r}_2|)} \rho_\lambda .$$

Note that Lemma 11.2 is an immediate consequence of Lemma B.1 and Lemma B.2 since $\mathcal{J}_\# = \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}})$ (see (B.5)), for the choices: $\sigma = \sigma(I_1, J_1)$, $\tilde{\sigma} = \sigma(\tilde{I}_1, J_1)$, $(r_1, r_2) = (m - n_1, m_2)$ and $(\tilde{r}_1, \tilde{r}_2) = (n - n_1, \tilde{m}_2)$. Hence it suffices to prove Lemma B.1 and Lemma B.2.

B.1. Proof of Lemma B.1 and Lemma B.2: We estimate the overlap integral (B.5) by considering the two cases: **Case 1:** $\tilde{\sigma} = 0$ and **Case 2:** $\tilde{\sigma} \neq 0$, and a number of subcases within each.

Case 1: $\tilde{\sigma} = 0$. In this case, for all $\mathbf{y} \in B_{r_0}(0)$, we have by (11.9):

$$(B.10) \quad p_0^\lambda(\mathbf{y} - \tilde{\sigma}\mathbf{e} - \tilde{r}_1\mathbf{v}_1 - \tilde{r}_2\mathbf{v}_2) = p_0^\lambda(\mathbf{y} - \tilde{r}_1\mathbf{v}_1 - \tilde{r}_2\mathbf{v}_2) \lesssim e^{-c\lambda(|\tilde{r}_1|+|\tilde{r}_2|)} p_0^\lambda(\mathbf{y}) .$$

Thus,

$$(B.11) \quad \begin{aligned} \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) &= \int p_0^\lambda(\mathbf{y} - [\sigma\mathbf{e} + r_1\mathbf{v}_1 + r_2\mathbf{v}_2]) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} - [\tilde{r}_1\mathbf{v}_1 + \tilde{r}_2\mathbf{v}_2]) d\mathbf{y} \\ &\lesssim e^{-c\lambda(|\tilde{r}_1|+|\tilde{r}_2|)} \int p_0^\lambda(\mathbf{y} - [\sigma\mathbf{e} + r_1\mathbf{v}_1 + r_2\mathbf{v}_2]) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y}) d\mathbf{y} . \end{aligned}$$

We next consider two subcases:

Subcase 1A: $\tilde{\sigma} = 0$ and $\sigma = 0$ and Subcase 1B: $\tilde{\sigma} = 0$ and $\sigma \neq 0$

Subcase 1A: $\tilde{\sigma} = 0$ and $\sigma = 0$ For any $(r_1, r_2) \neq (0, 0)$, we have by (11.12)

$$(B.12) \quad p_0^\lambda(\mathbf{y} - \sigma\mathbf{e} - r_1\mathbf{v}_1 - r_2\mathbf{v}_2) = p_0^\lambda(\mathbf{y} - [r_1\mathbf{v}_1 + r_2\mathbf{v}_2]) \lesssim e^{-c\lambda(|r_1|+|r_2|)} p_0^\lambda(\mathbf{y} - \mathbf{e}) .$$

Therefore, in subcase 1A we have after substitution of (B.12) into (B.11), that

$$(B.13) \quad \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda(|r_1|+|r_2|+|\tilde{r}_1|+|\tilde{r}_2|)} \rho_\lambda .$$

Interchanging the roles of \mathbf{r} and $\tilde{\mathbf{r}}$ in the case where $\tilde{\sigma} = \sigma = 0$, we also have that (B.13) holds unless $\tilde{\mathbf{r}} = 0$. Hence when $\sigma = \tilde{\sigma} = 0$, we have (B.13) unless $r_1 = r_2 = \tilde{r}_1 = \tilde{r}_2 = 0$.

Subcase 1B, $\tilde{\sigma} = 0$ and $\sigma \neq 0$: Then, by (11.10) we have

$$(B.14) \quad p_0^\lambda(\mathbf{y} - \sigma\mathbf{e} - r_1\mathbf{v}_1 - r_2\mathbf{v}_2) \lesssim e^{-c\lambda(|r_1|+|r_2|)} p_0^\lambda(\mathbf{y} - \sigma\mathbf{e})$$

unless $(r_1, r_2) \in N_b(\sigma)$. Substituting (B.14) into (B.11), we obtain the bound (B.13) unless $(r_1, r_2) \in N_b(\sigma)$.

Now consider the case where $(r_1, r_2) \in N_b(\sigma)$. Then, for some $l \in \{0, 1, 2\}$ which depends on σ, r_1 and r_2 we have: $p_0^\lambda(\mathbf{y} - (\sigma\mathbf{e} + r_1\mathbf{v}_1 + r_2\mathbf{v}_2)) = p_0^\lambda(\mathbf{y} - \sigma R^{-l}\mathbf{e})$, where $l = 0, 1$ or 2

and R is a $2\pi/3$ rotation matrix. Substituting into (B.11), we conclude that $\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda(|\tilde{r}_1|+|\tilde{r}_2|)}\rho_\lambda$. Indeed, using symmetry we obtain for $(r_1, r_2) \in N_b(\sigma)$:

$$\begin{aligned}
\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) &\lesssim e^{-c\lambda(|\tilde{r}_1|+|\tilde{r}_2|)} \int p_0^\lambda(\mathbf{y})\lambda^2|V_0(\mathbf{y})|p_0^\lambda(R^l\mathbf{y} - \sigma\mathbf{e}) d\mathbf{y} \\
&= e^{-c\lambda(|\tilde{r}_1|+|\tilde{r}_2|)} \int p_0^\lambda(R^l\mathbf{y})\lambda^2|V_0(R^l\mathbf{y})|p_0^\lambda(R^l\mathbf{y} - \sigma\mathbf{e}) d\mathbf{y} \\
\text{(B.15)} \quad &= e^{-c\lambda(|\tilde{r}_1|+|\tilde{r}_2|)} \int p_0^\lambda(\mathbf{z})\lambda^2|V_0(\mathbf{z})|p_0^\lambda(\mathbf{z} - \sigma\mathbf{e}) d\mathbf{z} = e^{-c\lambda(|\tilde{r}_1|+|\tilde{r}_2|)} \rho_\lambda.
\end{aligned}$$

Since $|r_1| + |r_2| = 0$ or 1 for $(r_1, r_2) \in N_b(\sigma)$, it follows that (B.13) holds (with a smaller constant, also denoted c , than appearing on the right hand side of (B.15)), unless $\tilde{r}_1 = \tilde{r}_2 = 0$. Therefore, if $\tilde{\sigma} = 0$ and $\sigma \neq 0$, the bound (B.13) holds provided $(\tilde{r}_1, \tilde{r}_2) \neq (0, 0)$.

Now consider the case where $\tilde{\sigma} = 0, \sigma \neq 0, (r_1, r_2) \in N_b(\sigma)$ and $(\tilde{r}_1, \tilde{r}_2) = (0, 0)$. Then,

$$\begin{aligned}
\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) &= \int p_0^\lambda(\mathbf{y} - [\sigma\mathbf{e} + r_1\mathbf{v}_1 + r_2\mathbf{v}_2]) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y}) d\mathbf{y} \\
&= \int p_0^\lambda(\mathbf{y} - [\sigma R^{-l}\mathbf{e}]) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y}) d\mathbf{y} = \rho_\lambda,
\end{aligned}$$

where R is a $2\pi/3$ rotation matrix and we have used the symmetry assumptions on V_0 .

Summarizing, for Case 1 we have proved:

Claim 1: $\tilde{\sigma} = 0$, then (B.13) holds unless

(1) $\sigma = 0$ and $r_1 = r_2 = \tilde{r}_1 = \tilde{r}_2 = 0$, a case we address no further since it does not arise in the proof of Proposition 7.1

or

(2) $\sigma \neq 0$ and $\tilde{r}_1 = \tilde{r}_2 = 0, (r_1, r_2) \in N_b(\sigma)$, in which case $\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) = \rho_\lambda$.

Furthermore, because $\tilde{\sigma}$ and σ play symmetric roles as do \mathbf{r} and $\tilde{\mathbf{r}}$, we have

Claim 2: if $\sigma = 0$, then the bound (B.13) on $\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}})$ holds unless

(1) $\tilde{\sigma} = 0$ and $r_1 = r_2 = \tilde{r}_1 = \tilde{r}_2 = 0$, a case we address no further since it does not arise in the proof of Proposition 7.1

or

(2) $\tilde{\sigma} \neq 0$ and $r_1 = r_2 = 0, (\tilde{r}_1, \tilde{r}_2) \in N_b(\tilde{\sigma})$, in which case $\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) = \rho_\lambda$.

We now turn to bound on $\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}})$ in

Case 2: $\sigma \neq 0$ and $\tilde{\sigma} \neq 0$

Case 2a: $\mathbf{r} \in N_b(\sigma)$ and $\tilde{\mathbf{r}} \in N_b(\tilde{\sigma})$: We claim that

$$\text{(B.16)} \quad \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda} \rho_\lambda \quad \text{for } \mathbf{r} \in N_b(\sigma), \tilde{\mathbf{r}} \in N_b(\tilde{\sigma}).$$

By (11.5), there exist $l, \tilde{l} \in \{0, 1, 2\}$ such that $p_0^\lambda(\mathbf{y} - [\sigma\mathbf{e} + \mathbf{r}\tilde{\mathbf{v}}]) = p_0^\lambda(\mathbf{y} - \sigma R^l \mathbf{e})$ and $p_0^\lambda(\mathbf{y} - [\tilde{\sigma}\mathbf{e} + \tilde{\mathbf{r}}\tilde{\mathbf{v}}]) = p_0^\lambda(\mathbf{y} - \tilde{\sigma} R^{\tilde{l}} \mathbf{e})$. Therefore,

$$\begin{aligned} \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) &= \int p_0^\lambda(\mathbf{y} - \sigma R^l \mathbf{e}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y} - \tilde{\sigma} R^{\tilde{l}} \mathbf{e}) d\mathbf{y} \\ &\lesssim e^{-c\lambda} \int p_0^\lambda(\mathbf{y} - \sigma R^l \mathbf{e}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y}) d\mathbf{y} \quad (\text{by (11.11)}) \\ &\lesssim e^{-c\lambda} \int p_0^\lambda(R^{-l}\mathbf{y} - \sigma\mathbf{e}) \lambda^2 |V_0(R^{-l}\mathbf{y})| p_0^\lambda(R^{-l}\mathbf{y}) d\mathbf{y} = e^{-c\lambda} \rho_\lambda. \end{aligned}$$

Case 2b: $\mathbf{r} \in N_b(\sigma)$ and $\tilde{\mathbf{r}} \notin N_b(\tilde{\sigma})$: We claim that

$$(B.17) \quad \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda} e^{-c\lambda|\tilde{\mathbf{r}}|} \rho_\lambda \quad \text{for } \mathbf{r} \in N_b(\sigma), \tilde{\mathbf{r}} \notin N_b(\tilde{\sigma}).$$

By (11.5) $p_0^\lambda(\mathbf{y} - [\sigma\mathbf{e} + \mathbf{r}\tilde{\mathbf{v}}]) = p_0^\lambda(\mathbf{y} - \sigma R^l \mathbf{e})$, and by (11.10) and (11.11) $p_0^\lambda(\mathbf{y} - [\tilde{\sigma}\mathbf{e} + \tilde{\mathbf{r}}\tilde{\mathbf{v}}]) \lesssim e^{-c|\tilde{\mathbf{r}}|\lambda} p_0^\lambda(\mathbf{y} - \tilde{\sigma}\mathbf{e}) \lesssim e^{-c\lambda} e^{-c|\tilde{\mathbf{r}}|\lambda} p_0^\lambda(\mathbf{y})$. These observations together with symmetry imply:

$$\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda} e^{-c\lambda|\tilde{\mathbf{r}}|} \int p_0^\lambda(\mathbf{y} - \sigma R^l \mathbf{e}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y}) d\mathbf{y} = e^{-c\lambda} e^{-c\lambda|\tilde{\mathbf{r}}|} \rho_\lambda.$$

This proves (B.17). Similarly, if $\mathbf{r} \notin N_b(\sigma)$ and $\tilde{\mathbf{r}} \in N_b(\tilde{\sigma})$ we have $\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{r}|} \rho_\lambda$.

Case 2c: $\mathbf{r} \notin N_b(\sigma)$ and $\tilde{\mathbf{r}} \notin N_b(\tilde{\sigma})$: We claim that

$$(B.18) \quad \mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c\lambda} e^{-c\lambda(|\mathbf{r}|+|\tilde{\mathbf{r}}|)} \rho_\lambda \quad \text{for } \mathbf{r} \notin N_b(\sigma), \tilde{\mathbf{r}} \notin N_b(\tilde{\sigma}).$$

By (11.10) and (11.11), $p_0^\lambda(\mathbf{y} - [\tilde{\sigma}\mathbf{e} + \tilde{\mathbf{r}}\tilde{\mathbf{v}}]) \lesssim e^{-c|\tilde{\mathbf{r}}|\lambda} p_0^\lambda(\mathbf{y} - \tilde{\sigma}\mathbf{e})$ and $p_0^\lambda(\mathbf{y} - [\sigma\mathbf{e} + \mathbf{r}\tilde{\mathbf{v}}]) \lesssim e^{-c\lambda} e^{-c\lambda|\mathbf{r}|} p_0^\lambda(\mathbf{y})$. Therefore,

$$\mathcal{J}(\sigma, \mathbf{r}, \tilde{\sigma}, \tilde{\mathbf{r}}) \lesssim e^{-c|\tilde{\mathbf{r}}|\lambda} e^{-c\lambda} e^{-c\lambda|\mathbf{r}|} \int p_0^\lambda(\mathbf{y} - \tilde{\sigma}\mathbf{e}) \lambda^2 |V_0(\mathbf{y})| p_0^\lambda(\mathbf{y}) d\mathbf{y} = e^{-c\lambda} e^{-c\lambda(|\mathbf{r}|+|\tilde{\mathbf{r}}|)} \rho_\lambda.$$

The bounds (B.16), (B.17) and (B.18) imply Lemma B.2, and together with Claim 1 and Claim 2 above Lemma B.1 follows. This also completes the proof of Lemma 11.2.

REFERENCES

- [1] M. J. Ablowitz, C. W. Curtis, and Y. Zhu, *On tight-binding approximations in optical lattices*, Studies in Applied Mathematics **129** (2012), no. 4, 362–388.
- [2] M J Ablowitz, S D Nixon, and Y Zhu, *Conical diffraction in honeycomb lattices*, Physical Review A **79** (2009), no. 5, 053830.
- [3] N.W. Ashcroft and N.D. Mermin, *Solid state physics*, Saunders College Publishing, 1976.
- [4] S. Becker, R. Han, and S. Jitomirskaya, *Cantor spectrum of graphene in magnetic fields* (<https://arxiv.org/abs/1803.00988>).
- [5] S. Becker and M. Zworski, *Magnetic oscillations in a model of graphene* (<https://arxiv.org/abs/1801.01931>).
- [6] G. Berkolaiko and A. Comech, *Symmetry and Dirac points in graphene spectrum*, Journal of Spectral Theory **8** (2018), no. 3, 1099–1147.
- [7] G. Berkolaiko and P. Kuchment, *Introduction to quantum graphs*, Mathematical Surveys and Monographs, vol. 186, American Mathematical Society, 2013.
- [8] O. Painter C. Brendel V. Peano and F. Marquardt, *Snowflake topological insulator for sound waves*, <https://arxiv.org/abs/1708.08888> (2017).
- [9] U. Carlsson, *An infinite number of wells in the semi-classical limit*, Asympt. Anal. **3** (1990), 189–214.

- [10] K. Chantelau, *Coulombic potentials in the semi-classical limit*, Lett. Math. Phys. **19** (1990), no. 4, 285–298.
- [11] F. Daumer, *équations de hartree-fock dans l’approximation du tight-binding*, Helv. Phys. Acta **67** (1994), no. 3, 237–256.
- [12] ———, *équations de schrödinger avec potentiels singuliers et à longue portée dans l’approximation de liaison forte*, Ann. Inst. H. Poincaré Phys. Théor. **64** (1996), no. 1, 1–31.
- [13] E. B. Davies and B. Simon, *Scattering theory for systems with different spatial asymptotics on the left and right*, Commun. Math. Phys. **63** (1978), 277–301.
- [14] Y. Colin de Verdière, *Sur les singularités de van hove génériques*, Mémoires de la S. M. F. série 2 **46** (1991), 99–109.
- [15] P. Delplace, D. Ullmo, and G. Montambaux, *Zak phase and the existence of edge states in graphene*, Physical Review B **84** (2011), no. 19, 195452.
- [16] M. Dimassi and J. Sjostrand, *Spectral asymptotics in the semi-classical limit*, London Mathematical Society Lecture Note Series, vol. 268, Cambridge University Press, 1999.
- [17] N. T. Do and P. Kuchment, *Quantum graph spectra of a graphyne structure*, Nanoscale Systems: Mathematical Modeling, Theory and Applications **2** (2013), 107–123.
- [18] A. Drouot, *Characterization of edge states in perturbed honeycomb structures*, Pure and Applied Analysis **1** (2019), no. 385–445.
- [19] A. Drouot, C. L. Fefferman, and M. I. Weinstein, *Edge states for one-dimensional asymptotically periodic systems* (preprint).
- [20] A. Drouot and M.I. Weinstein, *Edge states and the valley hall effect* (<https://arxiv.org/abs/1910.03509>).
- [21] C. L. Fefferman, J. P. Lee-Thorp, and M. I. Weinstein, *Topologically protected states in one-dimensional continuous systems and Dirac points*, Proceedings of the National Academy of Sciences (2014), 07391.
- [22] ———, *Bifurcations of edge states – topologically protected and non-protected – in continuous 2d honeycomb structures*, 2D Mater. **3** (2016), 014008.
- [23] ———, *Edge states in honeycomb structures*, Annals of PDE **2** (2016), no. 12.
- [24] ———, *Topologically protected states in one-dimensional systems*, Memoirs of the American Mathematical Society **247** (2017), no. 1173.
- [25] C. L. Fefferman and M. I. Weinstein, *Honeycomb lattice potentials and Dirac points*, J. Amer. Math. Soc. **25** (2012), no. 4, 1169–1220.
- [26] C.L. Fefferman, J.P. Lee-Thorp, and M.I. Weinstein, *Honeycomb Schroedinger operators in the strong-binding regime*, Comm. Pure Appl. Math. **71** (2018), no. 6.
- [27] A. K. Geim and K. S. Novoselov, *The rise of graphene*, Nature materials **6** (2007), no. 3, 183–191.
- [28] G. M. Graf and M. Porta, *Bulk-edge correspondence for two-dimensional topological insulators*, Comm. Math. Phys. **324** (2012), no. 3, 851–895.
- [29] V. V. Grushin, *Multiparameter perturbation theory of Fredholm operators applied to Bloch functions*, Mathematical Notes **86** (2009), no. 6, 767–774.
- [30] F. D. M. Haldane and S. Raghu, *Possible realization of directional optical waveguides in photonic crystals with broken time-reversal symmetry*, Phys. Rev. Lett. **100** (2008), no. 1, 013904.
- [31] M. Z. Hasan and C. L. Kane, *Colloquium: Topological Insulators*, Reviews of Modern Physics **82** (2010), 3045.
- [32] B. Helffer, *Semi-classical analysis for the schrödinger operator and applications*, Springer Lecture Notes in Mathematics, Springer-Verlag, 1988.
- [33] B. Helffer and J. Sjostrand, *Multiple wells in the semi-classical limit I.*, Comm. in P.D.E. **9** (1984), no. 4, 337–408.
- [34] ———, *Analyse semi-classique pour l’équation de Harper (avec application à l’équation de schrödinger avec champ magnétique)*, Mémoire de la Société Mathématique de France **34** (1988).
- [35] C. L. Kane and E. J. Mele, *\mathbb{Z}_2 topological order and the quantum spin Hall effect*, Phys. Rev. Lett. **95** (2005), 146802.
- [36] ———, *Quantum spin hall effect in graphene*, Physical Review Letters **95** (2005), no. 22, 226801.
- [37] Y. E. Karpeshina, *Perturbation Theory for the Schroedinger Operator with a Periodic Potential*, Springer Lecture Notes in Mathematics, vol. 1663, Springer, 1997.

- [38] A. B. Khanikaev and G. Shvets, *Two-dimensional topological photonics*, Nature Photonics **11** (2017), 763–773.
- [39] P. Kuchment and O. Post, *On the spectra of carbon nano-structures*, Comm. Math. Phys. **275** (2007), 805–826.
- [40] M. Lee, *Dirac cones for point scatterers on a honeycomb lattice*, SIAM J. Math. Anal. **48** (2016), no. 2, 1459–1488.
- [41] J. P. Lee-Thorp, I. Vukićević, X. Xu, J. Yang, C. L. Fefferman, C. W. Wong, and M. I. Weinstein, *Photonic realization of topologically protected bound states in waveguide arrays*, submitted.
- [42] J. P. Lee-Thorp, I. Vukicevic, X. Xu, J. Yang, C. L. Fefferman, C. W. Wong, and M. I. Weinstein, *Photonic realization of topologically protected bound states in domain-wall waveguide arrays*, Cleo, 2016.
- [43] J. P. Lee-Thorp, M. I. Weinstein, and Y. Zhu, *Elliptic operators with honeycomb symmetry; Dirac points, edge states and applications to photonic graphene*, Arch. Rational Mech. Anal. (2018).
- [44] L. Lu, J. D. Joannopoulos, and M. Soljačić, *Topological photonics*, Nature Photonics **8** (2014), 821–829.
- [45] M. Makwana and R. V. Craster, *Homogenization for hexagonal lattices and honeycomb structures*, Q. Jl. Mech. Appl. Math. (2014).
- [46] A. Mohamed, *Estimations semi-classiques pour l'opérateur de schrödinger á potentiel de type coulombien et avec champ magnétique*, Asympt. Anal. **4** (1991), no. 3, 235–255.
- [47] R.S.K. Mong and V. Shivamoggi, *Edge states and the bulk-boundary correspondence in dirac hamiltonians*, Physical Review B **83** (2011), no. 12, 125109.
- [48] K. Nakada, M. Fujita, G. Dresselhaus, and M.S. Dresselhaus, *Edge state in graphene ribbons: Nanometer size effect and edge shape dependence*, Phys. Rev. B **54** (1996), no. 24.
- [49] L. M. Nash, D. Kleckner, A. Read, V. Vitelli, A. M. Turner, and W. T. M. Irvine, *Topological mechanics of gyroscopic materials*, Proceedings of the National Academy of Sciences **112** (2015), no. 47, 14495–14500.
- [50] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, *The electronic properties of graphene*, Reviews of Modern Physics **81** (2009), 109–162.
- [51] J. Noh, S. Huang, K. P. Chen, and M. C. Rechtsman, *Observation of photonic topological valley Hall edge states*, Phys. Rev. Lett. **120** (2018), 063902.
- [52] K. S. Novoselov, *Nobel lecture: Graphene: Materials in the flatland*, Reviews of Modern Physics **83** (2011), no. 3, 837.
- [53] A. Outassourt, *Comportement semi-classique pour l'opérateur de Schrödinger a potentiel périodique*, J. Funct. Anal. **72** (1987), 65–93.
- [54] T. Ozawa, H. M. Price, A. Amo, N. Goldman, M. Hafezi, L. Lu, M. Rechtsman, D. Schuster, J. Simon, O. Zilberberg, and I. Carusotto, *Topological photonics* (preprint).
- [55] Y. Poo, J. P. Lee-Thorp, Y. Tan, R. Wu, M. I. Weinstein, and Z. Yu, *Observation of highly robust phase-defect induced photonic states*, Cleo, 2016.
- [56] S. Raghu and F. D. M. Haldane, *Analogs of quantum-Hall-effect edge states in photonic crystals*, Phys. Rev. A **78** (2008), no. 3, 033834.
- [57] M. C. Rechtsman, J. M. Zeuner, Y. Plotnik, Y. Lumer, D. Podolsky, F. Dreisow, S. Nolte, M. Segev, and A. Szameit, *Photonic Floquet topological insulators*, Nature **496** (2013), 196.
- [58] S. Shipman, *Reducible Fermi surfaces for non-symmetric bilayer quantum-graph operators* (DOI: 10.4171/JST/285).
- [59] B. Simon, *The bound state of weakly coupled Schrödinger operators in one and two dimensions*, Annals of Physics **97** (1976), no. 2, 279–288.
- [60] ———, *Semiclassical analysis of low lying eigenvalues, I. Non-degenerate minima: asymptotic expansions*, Ann. I.H.P. **A38** (1983), no. 3, 295–308.
- [61] ———, *Semiclassical analysis of low lying eigenvalues, II. tunneling*, Ann. Math. **120** (1984), 89–118.
- [62] ———, *Semiclassical analysis of low lying eigenvalues, III. width of the ground state band in strongly coupled solids*, Ann. Phys. **158** (1984), 415–420.
- [63] A. Singha, M. Gibertini, B. Karmakar, S. Yuan, M. Polini, G. Vignale, M. I. Kastnelson, A. Pinczuk, L. N. Pfeiffer, K. W. West, and V. Pellegrini, *Two-dimensional Mott-Hubbard electrons in an artificial honeycomb lattice*, Science **332** (2011), 1176.

- [64] S.S. Sunku, G.X. Ni, B.Y. Jiang, H. Yoo, A. Sternbach, A. S. McLeod, T. Stauber, L. Xiong, T. Taniguchi, K. Wantanabe, P. Kim, M. Fogler, and D.N. Basov, *Photonic crystals for nano-light in moiré graphene superlattices*, *Science* **362** (2018), 1153–1156.
- [65] K. Thicke, A. Watson, and J. Lu, *Computing edge states without hard truncation* (<https://arxiv.org/pdf/1810.07082v2.pdf>).
- [66] P.R. Wallace, *The band theory of graphite*, *Phys. Rev.* **71** (1947), 622.
- [67] Z. Wang, Y. D. Chong, J. D. Joannopoulos, and M. Soljacic, *Reflection-free one-way edge modes in a gyromagnetic photonic crystal*, *Phys. Rev. Lett.* **100** (2008), 013905.
- [68] E. T. Whittaker and G. Watson, *A course of modern analysis*, Cambridge University Press, 1902.
- [69] Y. Zhang, Y-W Tan, H. L. Stormer, and P. Kim, *Experimental observation of the quantum Hall effect and Berry's phase in graphene*, *Nature* **438** (2005), 201–204.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ, USA
E-mail address: cf@math.princeton.edu

DEPARTMENT OF APPLIED PHYSICS AND APPLIED MATHEMATICS AND DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY, USA
E-mail address: miw2103@columbia.edu