

The Incidental Parameters Problem in Testing for Remaining Cross-section Correlation *

Artūras Juodis ^{†1} and Simon Reese ^{‡2}

¹University of Groningen

²Lund University

October 29, 2019

Abstract

In this paper we consider the properties of the Pesaran (2004, 2015) CD test for cross-section correlation when applied to residuals obtained from panel data models with many estimated parameters. We show that the presence of period-specific parameters leads the CD test statistic to diverge as length of the time dimension of the sample grows. This result holds even if cross-section dependence is correctly accounted for and hence constitutes an example of the Incidental Parameters Problem. The relevance of this problem is investigated both for the classical Time Fixed Effects estimator as well as the Common Correlated Effects estimator of Pesaran (2006). We suggest a weighted CD test statistic which re-establishes standard normal inference under the null hypothesis. Given the widespread use of the CD test statistic to test for remaining cross-section correlation, our results have far reaching implications for empirical researchers.

Keywords. Incidental Parameters Problem, Panel Data, Cross-section Dependence, Factor Model, U-statistic.

JEL Classifications. C12, C23, C33.

*We would like to thank participants at AMES (Seoul, 2018 and Xiamen, 2019), IPDC (Seoul, 2018 and Vilnius, 2019), IAAE (Montreal, 2018), Nordic Econometric Meeting (Stockholm, 2019), and seminar participants at Copenhagen Business School, Copenhagen University, Erasmus University Rotterdam, KU Leuven, Monash University, New Economic School, Queen Mary University London, University of Amsterdam, University of Bern, University of Birmingham, University of Groningen, University of Southern California, and Uppsala University. We express special thanks to Geert Dhaene, Jiti Gao, Cheng Hsiao, Frank Kleibergen, Blaise Melly, Roger Moon, Andreas Pick, Geert Ridder, Vasilis Sarafidis, Peter Schmidt, Yongcheol Shin, Vanessa Smith, Bas Werker, and Joakim Westerlund for helpful comments.

[†]Corresponding author. a.juodis@rug.nl. Faculty of Economics and Business, University of Groningen, The Netherlands. Financial support from the Netherlands Organization for Scientific Research (NWO) under research grant number 451-17-002 is gratefully acknowledged.

[‡]simon.reese@nek.lu.se. Department of Economics, Lund University, Sweden. Simon Reese was a Postdoctoral Scholar Research Associate at USC Dornsife INET while working on this project.

1 Introduction

Given that economic agents rarely act entirely independently of each other, modeling cross-section dependence plays a prominent role in panel data econometrics. Time fixed effects are probably the simplest way of addressing this issue and allow controlling for a common trend whose effect is homogenous across cross-section units. During the last decade, interactive fixed effects models have become a popular, more general alternative. They assume the presence of a factor error structure, i.e. a small number of unobserved common trends interacted with entity-specific slope coefficients. Using either of the two modeling possibilities begs the question whether cross-section dependence is adequately accounted for.

In this paper, we show that the application of tests for cross-section dependence to regression residuals obtained from two-way fixed effects models or interactive effects models is problematic. We use the popular CD test statistic of Pesaran (2004, 2015) as an example and show that the inclusion of period-specific parameters introduces a bias term of order \sqrt{T} into the CD test statistic. In order to avoid erroneous rejection of the null hypothesis of no unaccounted cross-section dependence, we suggest an modified test statistic that re-weights cross-section covariances with Rademacher distributed weights. This weighted CD test statistic is asymptotically standard normal and has very good size under appropriate regularity conditions on the chosen weights. Additionally, we develop a refinement based on the power enhancement method of Fan et al. (2015) which considerably improves rejection rates under the alternative hypothesis.

The issue of cross-section dependence in panel data has stirred attention to the degree that now most advanced modern textbooks on panel data analysis dedicate at least one chapter to the consequences of ignoring cross-section correlation as well as the econometric techniques that have been developed to account for it, see e.g. Hsiao (2014), and Pesaran (2015b). Especially in panel data studies using macro-economic data, considerable effort has been exerted in estimating and testing appropriate model specifications capable of controlling for comovements across panel units. In this regard, the methods of Pesaran (2006) and Bai (2009) are most well-known.

Recently, applied econometricians have begun to use tests for cross-section dependence as an ex-post diagnostic tool, applied to residuals from a regression model that explicitly allows for cross-section dependence.¹ For example, Holly et al. (2010), Everaert and Pozzi (2014), Bailey et al. (2016), Eberhardt and Presbitero (2015), Mastromarco et al. (2016), and Chudik et al. (2017) (among others) apply the CD test of Pesaran (2004, 2015) to residuals obtained from estimation of their model with Pesaran's (2006) Common Correlated Effects (CCE) estimator. In some applications, the CD test statistic is explicitly used as a model-selection tool, interpreting a reduction in the absolute value of the CD test statistic as an indication of a *better* model. In other cases, only specifications not rejected by the CD test statistic (given some significance

¹For example, as implemented by the `xtcd/xtcdf` commands in Stata.

level) were considered. For example, Eberhardt et al. (2013, p. 444) partially motivate the choice of the empirical specification by noting that “[...] *curiously the residual CD tests for cross-section independence seem to reject in case of CCEP estimators. [...] The CCEP models fail to address the concerns for which they were developed, namely to account for all cross-section dependencies.*”

In microeconomic applications, the dominant modeling strategy has long been to consider a two-way fixed effects model. An example of the current practice is given by the earning dynamics literature which predominantly uses data from the Panel Study of Income Dynamics (PSID) (see e.g. Bonhomme and Robin 2010, Arellano et al. 2017, among others). Rather than analyzing the raw earnings data, empirical studies focus on the properties of residuals from a linear regression of log wages on a rich set of regressors and year specific time dummies. The post-estimation analysis is based on the implicit assumption that the model specification is sufficiently general to account for all sorts of cross-section dependence. While this assumption is (so far) rarely tested, the validity of two-way fixed effects models in applied microeconometrics and the policy evaluation literature has recently been called into question. For example, Hsiao et al. (2012b) and Gobillon and Magnac (2016) have recently emphasized the relevance of considering interactive fixed effects models rather than more restrictive time fixed effects or two-way fixed effects models when using difference-in-differences for treatment effect analysis. This sets the stage for an application of tests for cross-section dependence to residuals from two-way fixed effects models.

To the best of our knowledge, there are currently no theoretical results in the literature that could justify the use of tests for cross-section dependence as a misspecification test for regression models that account for cross-section dependence. Furthermore, this issue is usually completely ignored in any of the large scale Monte Carlo studies integral to theoretical and empirical papers in the field. Only very recently, Mao (2018) reported the size of three tests for cross-section dependence in a subset of his simulation experiments. However, despite clear evidence of excessive over-rejections, these results are neither discussed nor investigated theoretically. Therefore, this study is the first one to investigate the properties of the cross-section dependence tests applied to residuals, i.e. as a post-estimation diagnostic tool. In particular, given its popularity in the applied panel data literature, we restrict our attention to the CD test statistic of Pesaran (2004, 2015). Our interest lies in the class of residuals obtained from the models that characterize cross-section dependence driven by a small number of unobservable factors, entering the model either as time fixed effects in an additive error component model or in terms of a multifactor error structure, interacted with unit-specific slope coefficients. The results that we obtain are summarized as follows:

1. The application of the CD test to residuals obtained from a model where common factors enter either as time fixed effects or through a multifactor error structure renders the test statistic biased for any fixed T , and divergent as $T \rightarrow \infty$.
2. In addition to the mean of the CD test statistic, even its variance may be affected. This

can result in an asymptotically degenerate distribution of the test statistic.

3. Power may be reduced in small samples since divergent components resulting from the estimation of time fixed effects and the presence of unaccounted cross-section correlation can nearly cancel out. However, from a large sample perspective, the rate at which the CD test statistic diverges is generally unaffected.
4. A simple way of eliminating bias is to construct the CD test statistic from specifically weighted cross-section covariances rather than correlations. This leads to a valid test statistic for remaining cross-section correlation with good small sample properties in simulations.

The degeneracy of the CD statistics can be seen as a manifestation of the Incidental Parameters Problem (IPP) of Neyman and Scott (1948). In this respect, this paper contributes to this branch of the literature. So far, major focus in the panel data literature has been related to the IPP stemming from estimated individual specific effects. Our paper is the first one documenting non-trivial impact of estimated time specific common parameters on asymptotic properties. Furthermore, since the CD statistic can be seen as a time-series average of second degree (*degenerate*) U-statistics, our results shed some light on the potential impact of the IPP beyond simple cross-sectional averages. Lastly, while this article only considers linear models, the average correlation approach to testing for cross-section dependence was extended to nonlinear and nonparametric panel data models by Hsiao et al. (2012a) and Chen et al. (2012), respectively. Hence, problems documented in this paper carry over to post-estimation properties of non-linear models discussed in e.g. Chen et al. (2014), Fernández-Val and Weidner (2016), or Boneva and Linton (2017).

The remainder of this paper is structured as follows: Section 2 introduces the testing problem. In Sections 3 we present asymptotic results for stylized models with additive and multifactor error structure. In Section 4 we discuss standard approaches for bias correction and propose a weighted CD test statistic that achieves this goal. Sections 5 and 6 illustrate the problem documented in this paper by means of simulated and real data. Section 7 concludes.

Notation: I_m denotes an $m \times m$ identity matrix and the subscript is sometimes disregarded from for the sake of simplicity. $\mathbf{0}$ denotes a vector of zeros while \mathbf{O} stands for a matrix of zeros. s_m denotes a selection vector all of whose elements are zero except for element m which is one. $\mathbf{1}$ is a vector entirely consisting of ones. The dimension of these latter vectors and matrices is generally suppressed for the sake of simplicity and needs to be inferred from context. For a generic $m \times n$ matrix \mathbf{A} , $\mathbf{P}_A = \mathbf{A}(\mathbf{A}'\mathbf{A})^{-1}\mathbf{A}'$ projects onto the space spanned by the columns of \mathbf{A} and $\mathbf{M}_A = \mathbf{I}_m - \mathbf{P}_A$. Furthermore, $\text{rk}(\mathbf{A})$ denotes the rank of \mathbf{A} , $\text{tr}(\mathbf{A})$ its trace and $\|\mathbf{A}\| = (\text{tr}(\mathbf{A}'\mathbf{A}))^{1/2}$ the Frobenius norm of \mathbf{A} . For a set of $m \times n$ matrices $\{\mathbf{A}_1, \dots, \mathbf{A}_N\}$, $\bar{\mathbf{A}} = N^{-1} \sum_{i=1}^N \mathbf{A}_i$. δ and M stand for a small and large positive real number, respectively. For two real numbers a and b , $a \vee b = \max\{a, b\}$. Lastly, $\mathcal{O}_P(\cdot)$ and $o_P(\cdot)$ express stochastic order relations.

2 The testing problem

Let z_i be a T -dimensional data vector observed over N cross-sectional units indexed by i . Combining all z_i we obtain a two-dimensional data array of panel data (or longitudinal data). In empirical research it is common to investigate whether z_i can be regarded as independent over i in order to select a model that can properly characterize the statistical properties of the data. In particular, researchers might be interested in the statistical hypothesis

$$H_0 : z_i \perp z_j, \quad \text{for all } i, j = 1, \dots, N, \quad (1)$$

where we use the \perp notation to denote independence. Most often z_1, \dots, z_N contain residuals obtained from a regression model that does not allow for cross-section dependence, e.g. an entity fixed effects model or plain OLS.

By far the most widely used test for cross-section dependence is the CD test of Pesaran (2004, 2015), which is based on a simple rescaled sum of all pairwise cross-section correlation coefficients,² formally denoted

$$CD = \sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \hat{\rho}_{ij} = \sqrt{\frac{TN(N-1)}{2}} \bar{\hat{\rho}}. \quad (2)$$

Here,

$$\hat{\rho}_{ij} = \frac{T^{-1} \sum_{t=1}^T (z_{i,t} - \bar{z}_i)(z_{j,t} - \bar{z}_j)}{\sqrt{T^{-1} \sum_{t=1}^T (z_{i,t} - \bar{z}_i)^2} \sqrt{T^{-1} \sum_{t=1}^T (z_{j,t} - \bar{z}_j)^2}} \quad (3)$$

is the pairwise sample correlation coefficient between units i and j . Obviously, computing the CD test statistic involves obtaining $N(N-1)/2$ parameter estimates, each of which converges to the true parameter value at rate \sqrt{T} only. These circumstances are reminiscent of the panel data setup considered in e.g. Phillips and Moon (1999), Hahn and Kuersteiner (2002), where estimation of many incidental parameters in linear regression models turns out to have distributional effects on the asymptotic properties of common parameters. That is, they cause the incidental parameter problem. By contrast, the asymptotic distribution of the CD test statistic is unaffected by the estimation of all $N(N-1)/2$ cross-section correlation coefficients involved in its construction. In fact, applied to the residuals of a linear regression model with strictly exogenous regressors, and individual specific means, the CD test statistic is asymptotically normal as long as $N, T \rightarrow \infty$ (see Pesaran, 2015, Theorem 2), under H_0 of independence (or even limited local dependence). Hence, the IPP in the original setup disappears very rapidly as both dimensions increase. In short, for the setup of Pesaran (2004, 2015) the CD test statistic does not suffer from the (first order) IPP. However, as shown below, this result does not hold when the model specification includes period-specific parameters.

²Note that the CD test is designed for testing correlation, and not dependence in general. Only under assumption of joint normality these two concepts coincide.

2.1 Heuristic discussion of the incidental parameter problem

Consider a linear model where cross-section dependence is due to time fixed effects, so that unobserved heterogeneity over cross-sections and time enters the model additively. That is the relation between the $T \times 1$ vector \mathbf{y}_i and the $T \times m$ matrix \mathbf{X}_i is formally denoted

$$\mathbf{y}_i = \mathbf{X}_i\boldsymbol{\beta} + \boldsymbol{\tau} + \nu\mu_i + \boldsymbol{\varepsilon}_i, \quad i = 1, \dots, N. \quad (4)$$

Here, μ_i and $\boldsymbol{\tau}$ denote an entity-specific intercept and a $T \times 1$ vector of time-specific common parameters $\boldsymbol{\tau}$, respectively. $\boldsymbol{\varepsilon}_i$ is a vector of idiosyncratic error terms, independent across cross-sectional units. In the example of a Difference-in-Differences framework, $\boldsymbol{\tau}$ is the common trend affecting both treated and untreated individuals while the treatment indicator as well as other covariates are contained in \mathbf{X}_i . For the sake of simplicity, we assume $\boldsymbol{\beta}$ and μ_i to be known, so that $\boldsymbol{\beta} = \mathbf{0}$ and $\mu_i = 0 \forall i$ can be imposed without loss of generality. This highly restrictive assumption leaves the leading terms in the analysis below unaffected and is hence innocuous for the expository purpose of this section. Model (4) trivially reduces to

$$y_{i,t} = \tau_t + \varepsilon_{i,t}, \quad i = 1, \dots, N, \quad t = 1, \dots, T. \quad (5)$$

and we additionally assume that the variance of $\varepsilon_{i,t}$ is known and fixed to $\sigma_i^2 = 1$. In this setup, $y_{i,t}$ are clearly cross-sectionally dependent because of τ_t ; however we are interested in testing whether $\varepsilon_{i,t}$ are cross-sectionally uncorrelated. Given that $\varepsilon_{i,t}$ are unobserved, the common effects τ_t need to be estimated. The most natural approach is to estimate τ_t by OLS so that $\hat{\tau}_t = \bar{y}_t = \frac{1}{N} \sum_{i=1}^N y_{i,t}$. Using the residuals

$$\hat{\varepsilon}_{i,t} = y_{i,t} - \hat{\tau}_t, \quad (6)$$

the CD test statistic is given by

$$\begin{aligned} CD &= \sqrt{\frac{2}{TN(N-1)}} \sum_{t=1}^T \sum_{i=2}^N \sum_{j=1}^{i-1} \hat{\varepsilon}_{i,t} \hat{\varepsilon}_{j,t} \\ &= \sqrt{\frac{1}{2TN(N-1)}} \sum_{t=1}^T \left(\sum_{i=1}^N \hat{\varepsilon}_{i,t} \right)^2 - \sqrt{\frac{1}{2TN(N-1)}} \sum_{t=1}^T \sum_{i=1}^N \hat{\varepsilon}_{i,t}^2. \end{aligned} \quad (7)$$

Given this definition,

$$\sum_{i=1}^N \hat{\varepsilon}_{i,t} = \sum_{i=1}^N y_{i,t} - \sum_{i=1}^N y_{i,t} = 0,$$

implying that the first term in (7) cancels out. The expression $\sum_{t=1}^T \sum_{i=1}^N \hat{\varepsilon}_{i,t}^2$ in the second term is nothing more than the Sum of Squared Residuals (SSR) of the estimated model, a statistic that is of order $\mathcal{O}_P(NT)$. Consequently,

$$CD = \mathcal{O}_P(\sqrt{T})$$

even though $\varepsilon_{i,t}$ are cross-sectionally independent. Hence, the procedure commonly used by practitioners is prone to finding spurious cross-sectional dependence in the data.

The results shown for a model with additive unobserved heterogeneity are not coincidental. In fact, they carry over to a model where cross-section dependence is generated by a multifactor error structure. Following Pesaran (2006), this model is described by the $T \times 1$ vector \mathbf{y}_i and the $T \times m$ matrix \mathbf{X}_i defined as

$$\begin{aligned}\mathbf{y}_i &= \mathbf{X}_i\boldsymbol{\beta} + \mathbf{F}\boldsymbol{\lambda}_i + \boldsymbol{\varepsilon}_i \\ \mathbf{X}_i &= \mathbf{F}\boldsymbol{\Lambda}_i + \mathbf{E}_i.\end{aligned}\tag{8}$$

The most popular estimator designed to estimate the parameter vector $\boldsymbol{\beta}$ in this specific model is the CCE estimator of Pesaran (2006) which amounts to augmenting a linear regression model with cross-section averages of potentially all variables available to the researcher in order to account for the effect of unobservable common factors. In a model with homogenous slope coefficients, we have

$$\hat{\boldsymbol{\beta}}^{CCE} = \left(\sum_{i=1}^N \mathbf{X}_i' \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{X}_i \right)^{-1} \left(\sum_{i=1}^N \mathbf{X}_i' \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{y}_i \right)$$

where

$$\hat{\mathbf{F}} = \begin{bmatrix} \bar{\mathbf{y}} & \bar{\mathbf{X}} \end{bmatrix} = \left(\mathbf{F} \begin{bmatrix} \bar{\boldsymbol{\lambda}} & \bar{\boldsymbol{\Lambda}} \end{bmatrix} + \begin{bmatrix} \bar{\boldsymbol{\varepsilon}} & \bar{\mathbf{E}} \end{bmatrix} \right) \begin{bmatrix} 1 & \mathbf{0} \\ \boldsymbol{\beta} & \mathbf{I} \end{bmatrix} = \mathbf{F}\bar{\mathbf{C}} + \bar{\mathbf{U}}.\tag{9}$$

While the CCE estimator is agnostic about the true number of factors that affect the data, it has been shown that consistent estimation of the parameters of interest requires that the number of cross-section averages is at least as large as the true number of factors that drive the data (Westerlund and Urbain 2013).³ For this reason, an empirical practice of reporting the CD test statistic applied to CCE residuals has recently emerged, aimed at providing evidence on the amount of remaining cross-section dependence that is not accounted for by including cross-section averages into the model specification. A heuristic discussion of the exact mechanisms that lead to the failure of this procedure is provided in Appendix A.1. In the following section, we shall proceed to formally describing the consequences of the incidental parameters problem both for a model with multifactor error structure and the previously discussed model with additive error components.

3 Asymptotic Results

In this section we provide formal asymptotic results for CD statistic based on the residuals obtained from either the two-way FE model or a model with multifactor error structure. For

³An exception is the special case where the slope coefficients on the unobserved, common factors are random and uncorrelated.

the sake of simplicity, we continue to assume that β, μ_i and σ_i are known. However, we do not impose homogeneity of the individual-specific error variances. The heuristic discussion in Section 2 suggests that these assumptions, while being highly restrictive, do not affect the theoretical results presented in this section. To appreciate this claim, note the following two points: First, if individual fixed effects were unknown, they would be estimated via time averages of each cross-section. In contrast to cross-section averages, which eliminated the leading term of the CD test statistic (7), time averages do not interfere with any of its components. Second, if the vector of slope coefficients β were unknown, its estimator would converge to the true parameter value at the conventional rate \sqrt{NT} . By contrast, the documented effects on the first two moments of the CD test statistic resulted from the fact that estimates of common, period-specific parameters converge at a rate *slower* than \sqrt{NT} . Hence, effects similar to those arising from the estimation of time effects or latent common factors are not to be expected.

Additionally, the case of known slope coefficients nests the case where $\beta = \mathbf{0}$ is known, turning the models (6) and (8) into pure (restricted) factor models. This case is considered by Bailey et al. (2016, p.255) who suggest applying the CD test statistic on residuals of pure factor models as a general diagnostic technique. The issue is also of relevance in the context of PANIC panel unit root tests (Bai and Ng 2004)⁴ which investigate the common and the idiosyncratic components of the data separately. Given the problems that are associated with the use of information criteria for the number of common factors, an applied researcher might be tempted to use tests for cross-section dependence on the idiosyncratic errors of an estimated factor model in first differences and to increase the number of factors as long as remaining cross-section dependence is detected in the model residuals. The results reported below suggest that this method will not provide reliable information as to whether the factor model is correctly specified or not.

3.1 Additive error components

As in Section 2.1 we assume that the true model is given by equation (4). Additionally, we make the following assumptions on the model errors.

Assumption 1 (Errors).

1. Let $\varepsilon_{i,t} = \sigma_i \eta_{i,t}$ where $\eta_{i,t}$ is independently and identically distributed across both i and t with $E[\eta_{i,t}] = 0$, $E[\eta_{i,t}^2] = 1$ and $E[\eta_{i,t}^8] < M$ for some $M < \infty$.
2. σ_i is defined over an interval $[\delta; M]$ with $0 < \delta < M < \infty$. It is independently and identically distributed across i , and σ_i is independent of $\eta_{i,t}$ for all i and t .

For technical reasons, we assume that all stochastic variables in this paper have finite eighth moments. This is a sufficient condition, which facilitates proving joint convergence of the test

⁴Or alternatively the related PANICCA framework of Reese and Westerlund (2016).

statistics considered in this paper, see also Demetrescu and Hogg (2016). Assumption (1) is general enough to cover several models of conditional heteroscedasticity in $\varepsilon_{i,t}$.⁵ For example, natural examples for σ_i are either a standard exponential skedastic function

$$\sigma_i = \exp(\alpha + \gamma\mu_i), \quad (10)$$

or a location-scale model with

$$\sigma_i = \alpha + \gamma\mu_i. \quad (11)$$

Both satisfy the required restrictions, e.g. if μ_i has a bounded support. Furthermore, we denote different cross-sectional averages of σ_i by $\bar{\sigma}^k = N^{-1} \sum_{i=1}^N \sigma_i^k$ for $k = \{-2, -1, 1, 2\}$, and the corresponding population quantities by $E[\sigma_i^k]$. Assumption 1 guarantees that these quantities are well defined for all finite k .

Using these definitions the CD test statistic obtained from a model with unknown period-specific effects can be characterized as follows:

Theorem 1. *Suppose that the model parameters β , μ_i and σ_i are known. Under Assumption 1,*

$$\begin{aligned} CD &= \sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T \varepsilon_{i,t} \varepsilon_{j,t} \left(\sigma_i^{-1} - \bar{\sigma}^{-1} \right) \left(\sigma_j^{-1} - \bar{\sigma}^{-1} \right) + \sqrt{T} \Xi \\ &+ \mathcal{O}_P(\sqrt{TN}^{-1}), \end{aligned} \quad (12)$$

where

$$\Xi = \sqrt{\frac{N}{2(N-1)}} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 \left(\left(\bar{\sigma}^{-1} \right)^2 - 2\bar{\sigma}^{-1} \sigma_i^{-1} \right).$$

Furthermore, let $\Omega = \left(1 - 2E[\sigma_i]E[\sigma_i^{-1}] + E[\sigma_i^2](E[\sigma_i^{-1}])^2 \right)^2$. Then,

$$CD - \sqrt{T}\Xi \xrightarrow{d} N(0, \Omega) \quad (13)$$

as $N, T \rightarrow \infty$ jointly subject to the restriction $\sqrt{TN}^{-1} \rightarrow 0$.

Theorem 1 shows that the inclusion of time fixed effects into the model specification has an asymptotically non-negligible effect on the first two moments of the CD test statistic. Via expansion with appropriate functions of $\sigma_1^2, \dots, \sigma_N^2$, it can be shown that Ξ indicates the presence of a deterministic bias of order \sqrt{T} . Abstracting from this bias, (13) is dominated by an expression that reflects the CD test statistic obtained from the true model errors but imposing an incorrect normalization.

⁵Alternatively this assumption can be formulated in terms of unconditional variances, where σ_i are fixed numbers. However, in addition to having severe conceptual shortcomings (as discussed in (Gagliardini et al. 2016)) such an approach would lead to incorrect conclusions concerning the power of a CD test statistic for remaining cross-section correlation. See Section 3.3 for a detailed discussion.

The behavior of the CD test statistic in a model with time fixed effects can be seen as an example of the incidental parameters problem (IPP) of Neyman and Scott (1948),⁶ since the bias of the CD test statistic is due to the estimation of T period-specific intercepts τ_t . Interestingly, in the context of estimating linear dynamic panel data models, estimation of the time effects τ_t does not introduce any asymptotic bias into the FE estimator with strictly or weakly exogenous regressors (see e.g. Hahn and Moon 2006). In non-linear models, estimation of the time effects τ_t affects the asymptotic mean of the estimator for slope parameters associated with explanatory variables (see e.g. Fernández-Val and Weidner 2016) with the corresponding bias being proportional to $\sqrt{TN^{-1}}$. In this sense, our result adds new insights into the literature in that it highlights a scenario where the inclusion of time fixed effects into the model has non-standard implications for the asymptotic distribution of the statistic of interest.

The results of Theorem 1 suggest that asymptotically standard normal inference can be recovered by bias-correcting and rescaling (12). Before considering this remedy in Section 4, it is important to emphasise the following special case:

Corollary 1. *Under Assumption 1 and given $P(\sigma_i = \sigma) = 1$,*

$$CD = -\sqrt{\left(T - \frac{T}{N}\right) / 2} - \frac{1}{\sqrt{2N}} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T \left(\frac{\varepsilon_{i,t}^2}{\sigma^2} - 1 \right) \right) + o_P(N^{-1/2})$$

Corollary 1 provides more intuition about the approximate value of the bias term $\sqrt{T}(\Xi_1 - 2\Xi_2)$, suggesting that it should be reasonably close to $-\sqrt{T}/2$. More importantly, the result reveals that the leading stochastic component in the CD test statistic, the first term on the right-hand side of equation (12), cancels out when error variances are homogeneous across i . Instead, random variation around the bias term $-\sqrt{\frac{T}{2}}$ is of order $\mathcal{O}_P(N^{-1/2})$, rendering the distribution of a modified version of the test statistic asymptotically degenerate.

The special case of homogeneous error variances hence entails consequences for the CD test statistic that are qualitatively different from those in the more general case where σ_i may differ across i . Again, it would be possible to allow for asymptotically normal inference by adequately rescaling the modified test statistic. However, the resulting statistic would be of little practical use since the main source of variation is not related to error covariances across cross-sections, but is simply driven by variance of the idiosyncratic components.

3.2 Multifactor error structure

In analogy with the previous sections, we disregard from estimation error around the true slope coefficients of model (8). As shown by Pesaran (2006) and Juodis et al. (2017), the corresponding term $\hat{\beta}^{CCE} - \beta_0$ is of order $\mathcal{O}_P((NT)^{-1/2}) + \mathcal{O}_P(N^{-1})$. Hence, convergence to the true parameter values is fast enough to ensure that any terms present in CD involving

⁶See also Lancaster (2000).

$\hat{\beta}^{\text{CCE}} - \beta_0$ are asymptotically negligible. We shall not attempt to prove this claim formally but rely on Monte Carlo results of Section 5 to support this conjecture.

Given that the factor estimator \hat{F} , defined in equation (9), is constructed based on observed data, restrictions on the DGP of both $y_{i,t}$ and $x_{i,t}$ need to be imposed.

Assumption 2.

1. Let $\varepsilon_{i,t} = \sigma_i \eta_{i,t}$ where $\eta_{i,t}$ is independently and identically distributed across both i and t with $E[\eta_{i,t}] = 0$, $E[\eta_{i,t}^2] = 1$ and $E[\eta_{i,t}^8] < M$ for some $M < \infty$.
2. σ_i is defined over an interval $[\delta; M]$. It is independently and identically distributed across i .
3. The $m \times 1$ random vector $e_{i,t}$ is independently distributed across both i and t with $E[e_{i,t}] = \mathbf{0}$, $E[e_{i,t}e'_{i,t}] = \Sigma$ with the latter being a positive definite matrix and $E[\|e_{i,t}\|^8] < M$.

Assumption 3. f_t is a covariance stationary $r \times 1$ random vector with positive definite covariance matrix Σ_F , absolutely summable autocovariances and $E[\|f_t\|^4] < M$.

Assumption 4. λ_i is iid over i with $E[\lambda_i] = \mu_\lambda$ and $E[\|\lambda_i\|^4] < \infty$. Furthermore, Λ_i is iid over i with $E[\Lambda_i] = \mu_\Lambda$ and $E[\|\Lambda_i\|^4] < \infty$.

Assumption 5. $f_t, \{\lambda_i, \Lambda_i\}, \eta_{i',t'}, e_{i'',t''}$ and $\sigma_{i''}$ are mutually independent for all i, i', i'', i''', t, t' and t''

Assumption 6. $\text{rk}([\mu_\lambda, \mu_\Lambda]) = r = m + 1$.

This set of assumptions above are a slightly more restrictive version of the framework considered in Pesaran (2006). For example, the assumption of common Σ can be straightforwardly relaxed. However, unlike σ_i , Σ plays no major role for asymptotic results of this paper. Most importantly, we rule out the presence of serial correlation as this is in line with the assumptions made for the CD test to work. Furthermore, we restrict ourselves to a classical panel data regression model instead of considering heterogeneous slope coefficients. Moreover, any higher-order dependence between $\varepsilon_{i,t}$ and $e_{i,t}$ is assumed away in order to allow for a tractable proof of the main theoretical result in this paper. Lastly, the fact that we assume $\text{rk}([\mu_\lambda, \mu_\Lambda]) = r = m + 1$ to hold, suggest that our we consider an ideal setup where none of the rank condition related problems documented in Karabiyik et al. (2017), or Juodis et al. (2017) apply.

We begin our asymptotic analysis, by noting that in the model with assumed (known) homogeneous σ the result follows directly as in the model with time effects only. In particular, while it is not generally emphasized in the CCE literature, the residuals from CCE estimation which are formally given by

$$\hat{\varepsilon}_i = \varepsilon_i - P_{\hat{F}}\varepsilon_i - (P_{\hat{F}} - P_F)F\lambda_i \tag{14}$$

satisfy

$$\sum_{i=1}^N \hat{\varepsilon}_{i,t} = 0.$$

In that respect, the standard two-way error component FE estimator is similar to CCE estimator. More formally we formulate the following result

Proposition 1. *Under Assumptions 2–6 and $P(\sigma_i = \sigma) = 1$:*

$$CD = -\sqrt{\left(T - \frac{T}{N}\right)}/2 + \mathcal{O}_P(R_{N,T}), \quad R_{N,T} = (N^{-1/2} \vee T^{-1/2} \vee N^{-1}\sqrt{T}).$$

Proposition 1 shows that the result we derived previously for additive model in Corollary 1 continues to hold for models with a factor error structure, as long as $N \rightarrow \infty$. Note that remainder terms in Proposition 1 are $\mathcal{O}_P(1)$ only if both $N, T \rightarrow \infty$. This approximation error is a direct by-product of the CCE approach, as the idiosyncratic component $e_{i,t}$ enters the equation of interest indirectly via the factor proxies (see also Theorem 2).

It is worth mentioning that the above order effect is only valid if \hat{F} contains cross-sectional averages of the regressand as well as all regressors. If either of those variables is omitted (without affecting the rank condition in Assumption 6), the zero mean residual condition in (14) is violated, and consequently the CD test will have a $\mathcal{O}_P(1)$ term. However, this result is of limited empirical importance as in most cases researchers include all available cross-sectional averages. While this practice ensures that the estimator is invariant to β_0 , inclusion of too many cross-sectional averages can potentially have detrimental effects on the asymptotic properties of the estimator, see the corresponding discussion in Juodis et al. (2017).

In the homoscedastic case additive and multiplicative error component models have similar asymptotic effects on CD test statistic. The conclusions of the next theorem, which is the main result of this paper, indicate that this equivalence does not hold in the heteroscedastic case. In order to proceed, we introduce some useful notation. For $t = 1, \dots, T$, let

$$\left(\bar{\mathcal{C}}'\right)^{-1} \hat{f}_t - f_t = \frac{1}{N} \sum_{i=1}^N \psi_{i,t}, \tag{15}$$

where $\psi_{i,t} = \left(\bar{\mathcal{C}}'\right)^{-1} \mathbf{u}_{i,t}$ is the influence function of the corresponding factor estimator, in this case cross-section averages of $y_{i,t}$ and $\mathbf{x}_{i,t}$. Generally, the influence function depends on the joint process $[y_{i,t}, \mathbf{x}_{i,t}]'$, as long as all observed variables are used to form cross-sectional averages. Equipped with this notation we formulate the main result of this paper.

Theorem 2. *Under Assumptions 2–6,*

$$CD = \sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T \xi_{iN,t} \xi_{jN,t} + \sqrt{T} \Phi_1 - 2\sqrt{T} \Phi_2 + \mathcal{O}_P(R_{N,T}),$$

where

$$\zeta_{iN,t} = \sigma_i^{-1} \varepsilon_{i,t} - \left(\frac{1}{N} \sum_{i=1}^N \sigma_i^{-1} \lambda_i \right)' \psi_{i,t},$$

and

$$\begin{aligned} \Phi_1 &= \sqrt{\frac{N}{2(N-1)}} \left(\frac{1}{N} \sum_{i=1}^N \sigma_i^{-1} \lambda_i \right)' \left(\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \psi_{i,t} \psi_{i,t}' \right) \left(\frac{1}{N} \sum_{i=1}^N \sigma_i^{-1} \lambda_i \right), \\ \Phi_2 &= \sqrt{\frac{N}{2(N-1)}} \left(\frac{1}{N} \sum_{i=1}^N \sigma_i^{-1} \lambda_i \right)' \left(\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \psi_{i,t} \sigma_i^{-1} \varepsilon_{i,t} \right). \end{aligned}$$

Here, as previously, $R_{N,T} = (N^{-1/2} \vee T^{-1/2} \vee N^{-1} \sqrt{T})$. In line with all previous results, the CD test statistic in this setup has two diverging components. However, unlike all previous results, in particular Theorem 1, these bias terms are not solely non-linear functions of σ_i . Instead, they are also influenced by the first (rescaled) moments of factor loadings in $y_{i,t}$ and $x_{i,t}$, as well as corresponding variances of the idiosyncratic components in $x_{i,t}$. Thus the influence function $\psi_{i,t}$ directly alters distributional properties of the CD statistic. The result is qualitatively similar to any parametric two-step estimation procedure with a plug-in first-step estimator. Thus by including cross-sectional averages as factor proxies, one is implicitly testing that both $\varepsilon_{i,t}$ and $\psi_{i,t}$ are jointly cross-sectionally uncorrelated.⁷

Remark 1. One can easily see that expressions for $\zeta_{iN,t}$, Φ_1 , Φ_2 are the same if one assumes that λ_i is known. Thus it is only the influence function of common factors, and not those of factor loadings, that has an impact on the asymptotic properties of the test statistic. This conclusion is the same as in the additive model.

As alluded in Proposition 1, the leading $\mathcal{O}_p(1)$ is degenerate if $\sigma_i = \sigma$, as in this case:

$$\sum_{i=1}^N \zeta_{iN,t} = 0, \quad \forall t = 1, \dots, T. \quad (16)$$

One can easily see that the expressions for Φ_1 and Φ_2 derived for CCE coincide with the corresponding terms of Theorem 1 (up to a negligible remainder term), upon setting $\lambda_i = 1$ (thus $\bar{C}^{-1} = 1$), and $\psi_{i,t} = \varepsilon_{i,t}$. This situation is equivalent to using $\hat{f}_t = \bar{y}_t - \bar{x}_t' \beta_0$ as factor proxies.⁸

Recall that result in Theorem 2 considers only the CCE setup where the number of observable factor proxies equals the number of the true factors. If Assumption 6 is relaxed and there are more observables than factors, then following Karabiyik et al. (2017) and Juodis et al. (2017), one can show that the expressions for Φ_1 and Φ_2 will contain additional terms related to

⁷While $\varepsilon_{i,t}$ is usually assumed to be uncorrelated over t , the same is not true for $u_{i,t}$, e.g. if $x_{i,t} = y_{i,t-1}$. Thus if Assumption 2 is appropriately relaxed, then $\zeta_{iN,t}$ will be serially correlated.

⁸E.g. as suggested by Westerlund et al. (2017) in the context of predictability testing with cross-sectional dependence.

this discrepancy. In particular, Φ_1 , Φ_2 , and $\zeta_{iN,t}$ are functions of an unknown rotation matrix, which cannot be consistently estimated from the data.⁹ We will come back to this issue in Section 4.1. Finally, our results remain completely silent on the behavior of the test statistic where $m + 1 \rightarrow \infty$, a scenario for which the CCE estimator for panel data models with heterogeneous coefficients has been studied by Chudik and Pesaran (2015) and Chudik et al. (2017).

Remark 2. Given that the asymptotic bias of the CD test statistic is driven by estimated time specific variables τ_t (or f_t), and not by individual specific variables, it is natural to expect that the same type of results also applies to group-specific estimates of τ_t estimated at rate \sqrt{N} , e.g. as in Bonhomme and Manresa (2015), Su et al. (2016), or Bonhomme et al. (2017). Analogous results can also be derived for the CD test statistic applied to residuals obtained using the interactive fixed effects estimator of Bai (2009) Similarly, analogous results can be derived for test statistic based on Principal Components based residuals based on PC as in Bai (2009) or Westerlund and Urbain (2015). The expressions in Theorem 2 can be modified accordingly based on corresponding influence functions $\psi_{i,t}$ in each setup. Section S.1.2 in the online supplement accompanying this article provides an informal discussion of these extensions.

Remark 3. As it was discussed previously, our results are informative about general properties of the second degree U-statistics. In particular, given that the LM test statistic as in Baltagi et al. (2012), is also a U-statistic of the same degree, similar divergence patterns can be expected. Similarly, more powerful versions of *CD*, as recently proposed by Demetrescu and Homm (2016) and Mao (2016), share similar asymptotic properties.

3.3 Power analysis

The previous two subsections showed that sample estimates of population correlation coefficients in model errors play only a secondary role in a CD test statistic that is applied to two-way fixed effects residuals or CCE residuals. This raises the question as to whether any component in *CD* reliably indicates the presence of unaccounted cross-section correlation if neither two-way fixed effects or CCE estimation eliminate all sources of co-movements across cross-sections from the data.

To simplify the discussion, we disregard from the effect of covariates on the variable of interest $y_{i,t}$ and assume that the true model is given by a pure static factor model, amounting to model (8) with $\beta = \mathbf{0}$. Extending our results to the case of general β is possible, although formally cumbersome.¹⁰ In a first instance, suppose that a researcher erroneously assumes

⁹For the above reason Juodis et al. (2017) suggest the use of non-parametric bias correction procedure for CCE estimator.

¹⁰To appreciate this point note that, even if the unobserved heterogeneity driving the data is misspecified, least squares estimates of β are consistent for the pseudo-true parameter vector $\delta = \text{plim}_{N,T \rightarrow \infty} \left[\left(\sum_{i=1}^N \check{X}_i' \check{X}_i \right)^{-1} \left(\sum_{i=1}^N \check{X}_i' \check{y}_i \right) \right]$ with \check{X}_i and \check{y}_i being orthogonal to estimates of the assumed sources of unobserved heterogeneity.

cross-section dependence to stem from time fixed effects. The corresponding mis-specified model is formally given by

$$y_{i,t} = \mu_{y,t} + v_{i,t}, \quad (17)$$

where $\mu_{y,t} = \mathbf{f}'_t \mathbb{E}[\boldsymbol{\lambda}_i]$ and $v_{i,t} = \mathbf{f}'_t (\boldsymbol{\lambda}_i - \mathbb{E}[\boldsymbol{\lambda}_i]) + \tilde{\varepsilon}_{i,t}$. Deviations of the data from their cross-section averages can accordingly be written as

$$\begin{aligned} \hat{v}_{i,t} &= y_{i,t} - N^{-1} \sum_{i=1}^N y_{i,t} \\ &= \mathbf{f}'_t \tilde{\boldsymbol{\lambda}}_i + \tilde{\varepsilon}_{i,t} \end{aligned} \quad (18)$$

where $\tilde{\boldsymbol{\lambda}}_i = \boldsymbol{\lambda}_i - N^{-1} \sum_{i=1}^N \boldsymbol{\lambda}_i$ and $\tilde{\varepsilon}_{i,t} = \varepsilon_{i,t} - N^{-1} \sum_{i=1}^N \varepsilon_{i,t}$. These deviations are sample equivalents of the composite error term $v_{i,t}$ in the mis-specified time fixed effects model (17). Let the variance of this composite error be defined as $\varsigma_{v,i}^2 = \mathbb{E} \left[v_{i,t}^2 \mid \sigma_i, \boldsymbol{\lambda}_i \right] = (\boldsymbol{\lambda}_i - \mathbb{E}[\boldsymbol{\lambda}_i])' \boldsymbol{\Sigma}_F (\boldsymbol{\lambda}_i - \mathbb{E}[\boldsymbol{\lambda}_i]) + \sigma_i^2$ and let $\overline{\varsigma}_v^k = N^{-1} \sum_{i=1}^N \varsigma_{v,i}^k$, $k \in \{-2, -1, 1, 2\}$ be shorthand notation for cross-section averages of $\varsigma_{v,i}^k$. For the sake of simplicity, we continue to assume that $\overline{\varsigma}_v^2$ is known and used to standardize regression residuals when constructing the CD test statistic. This greatly simplifies the proofs of the following propositions while leaving the main results of this section qualitatively unaffected. We confirm this conjecture with corresponding simulations in Section 5, allowing for both unknown error variances and unknown, general slope coefficients $\boldsymbol{\beta}$. Given our current setup, let the CD test statistic constructed from $\hat{v}_{i,t}$ be expressed by

$$CD_{\mathbb{H}_1} = \sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^N \sum_{t=1}^T \varsigma_{v,i}^{-1} \hat{v}_{i,t} \hat{v}_{j,t} \varsigma_{v,j}^{-1}. \quad (19)$$

The properties of this test statistic are characterized as follows:

Proposition 2. *Suppose that the true model is given by (6) with $\boldsymbol{\beta} = \mathbf{0}$ and that its components satisfy Assumptions 2–6. Let the CD test statistic $CD_{\mathbb{H}_1}$ be constructed from the cross-sectionally demeaned data (18). Then,*

$$\begin{aligned} CD_{\mathbb{H}_1} &= \sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \left(\varsigma_{v,i}^{-1} - \mathbb{E} \left[\varsigma_{v,i}^{-1} \right] \right) \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \left(\varsigma_{v,j}^{-1} - \mathbb{E} \left[\varsigma_{v,j}^{-1} \right] \right) + \sqrt{T} \Xi_{\mathbb{H}_1} \\ &\quad + O_p \left(N^{-1} \sqrt{T} \right), \end{aligned} \quad (20)$$

where

$$\Xi_{\mathbb{H}_1} = \sqrt{\frac{N}{2(N-1)}} (NT)^{-1} \sum_{i=1}^N \boldsymbol{\nu}'_i \boldsymbol{\nu}_i \left[\left(\overline{\varsigma}_v^{-1} \right)^2 - 2 \overline{\varsigma}_v^{-1} \overline{\varsigma}_{v,i}^{-1} \right].$$

Furthermore, the leading stochastic term $\sqrt{\frac{1}{2TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \left(\varsigma_{v,i}^{-1} - \mathbb{E} \left[\varsigma_{v,i}^{-1} \right] \right) \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \left(\varsigma_{v,j}^{-1} - \mathbb{E} \left[\varsigma_{v,j}^{-1} \right] \right)$ has expected value $N\sqrt{T} \text{cov} \left[\varsigma_{v,1}^{-1}, \boldsymbol{\lambda}'_1 \right] \boldsymbol{\Sigma}_F \text{cov} \left[\boldsymbol{\lambda}_1, \varsigma_{v,1}^{-1} \right]$. In the special case $\text{cov} \left[\boldsymbol{\lambda}_1, \varsigma_{v,1}^{-1} \right] = \mathbf{0}$ the order of this leading term is determined by its variance which diverges at rate T .

Proposition 2 establishes a decomposition of the CD test statistic that is almost identical to that obtained under the null hypothesis: Analogous to Theorem 1 CD_{H_1} consists of a leading stochastic component which reflects a CD test statistic with incorrect normalization as well as an equivalent to the bias term in equation (12) which we denote by $\sqrt{T}\Xi_{H_1}$. The key difference is that both components are functions of the composite model error $v_{i,t} = \mathbf{f}'_t(\boldsymbol{\lambda}_i - \mathbb{E}[\boldsymbol{\lambda}_i]) + \varepsilon_{i,t}$ rather than the true errors $\varepsilon_{i,t}$.

Noticeable differences with regards to Theorem (12) are given as far as the leading stochastic component of CD_{H_1} , the first term in equation (20), is concerned. Its mean is a positive-definite quadratic form and will therefore generally diverge to $+\infty$ at rate $N\sqrt{T}$. This result is interesting in the context of Sarafidis et al. (2009) who conjecture “[...] that the CD test will have poor power properties when it is applied to a regression with time dummies or on cross-sectionally demeaned data.” First, the cited statement does not acknowledge the bias term Ξ_{H_1} which generally leads CD_{H_1} to diverge. However, it is reasonable not to consider this term as a source of power towards the alternative hypothesis since it does not involve sample estimates of cross-section covariances (see Remark 5 below). Second, Proposition 2 indicates that the CD test indeed has power since remaining sources of co-movements across cross-sections generally affect the mean of its leading stochastic component. The failure to appreciate this property is rooted in a convention to treat error variances as fixed parameters, thereby ruling out any correlation between them and factor loadings. Extending this understanding of error variances to the variance of the composite error term $v_{i,t}$ (i.e. $\zeta_{v,i}^2$) leads to the special case $\text{cov}[\boldsymbol{\lambda}_1, \zeta_{v,1}^{-1}] = \mathbf{0}$ in which the power of CD is indeed considerably reduced.

Still, while the rate at which CD_{H_1} diverges is in general unaffected by the inclusion of time fixed effects, power in small samples may be compromised because of the presence of two diverging components with opposite signs in expression (20). While the mean of the leading stochastic component dominates in large samples, it may be canceled out by Ξ_{H_1} if the number of observations available to the researcher is rather small.

Remark 4. It is straightforward to extend the results of Proposition 2 to a model specification without time fixed effects and to address a common concern about the power of CD that has repeatedly been made in the literature. As argued in Pesaran et al. (2008) and Sarafidis et al. (2009), the power of this test is reduced substantially if unaccounted sources of cross-section correlation average out, as for example in a factor model with zero mean factor loadings. This property is given by construction for two-way fixed effects or time fixed effects residuals by the mere fact that the within transformation involves cross-sectionally demeaning the data. While the bias term in Proposition 2 and the scaling effect on the leading stochastic component in CD_{H_1} are exclusively a consequence of accounting for time fixed effects, the defining role of $\text{cov}[\boldsymbol{\lambda}_1, \zeta_{v,1}^{-1}]$ for the rate at which CD_{H_1} diverges is a general property in models with zero mean loadings. It follows that existing claims about the power losses of CD in the presence of factor with zero mean loadings merely address the special case with $\text{cov}[\boldsymbol{\lambda}_1, \zeta_{v,1}^{-1}] = \mathbf{0}$. In fact,

Monte Carlo results that have been reported to support these claims, as e.g. in Pesaran (2004, 2015), Pesaran et al. (2008), Sarafidis et al. (2009) and Baltagi et al. (2012), all use specifications which entail a zero covariance between factor loadings and the inverse error variances.¹¹ The CD test statistic is very likely to have much better power properties in simulation experiments that enforce a non-zero correlation between factor loadings and error variances. A simple way of achieving this would be to impose a constant ratio between the magnitude of common variation and that of idiosyncratic variation which holds for all cross-sections. See (Parker and Sul 2016, equation (27)) or Section 5 in this study for a formal definition.

Remark 5. An additional similarity of Proposition 2 with Theorem 1 is that the term Ξ_{H_1} does not involve sample estimates of cross-section covariances, implying that this term is not indicative of the degree of cross-section co-movement in the data. Moreover, it can be shown that $\Xi_{H_1} \xrightarrow{p} \left(E \left[\zeta_{v,i}^{-1} \right] \right)^2 E \left[\zeta_{v,i}^2 \right] - 2 E \left[\zeta_{v,i}^{-1} \right] E \left[\zeta_{v,i} \right]$, so that the leading term in $\sqrt{T}\Xi_{H_1}$ would approach $-\sqrt{T/2}$ as $\zeta_{v,i} \rightarrow \zeta_v$, for some constant $\zeta_v > 0$.

The observations made for models that assume an additive error component structure apply almost identically for mis-specified common latent factor models estimated by the CCE procedure, see Appendix A.2.

4 Re-establishing standard normal inference

The diverging bias in the CD test statistic applied to residual from models with common, period-specific parameters is fundamentally problematic for its use in the context of testing for remaining cross-section correlation. Still, the literature review in Section 1 suggests that the underlying question of correct model specification is of high relevance for empirical researchers. For this reason it is relevant not to discard completely the CD test statistic but instead to discuss possible modifications aimed at ensuring an asymptotically standard normal inference under null hypothesis. Thus, methods aimed at addressing the IPP detailed in Sections 2 and 3 need to be considered.

4.1 Analytic bias correction

Given that parametric expressions for the bias of *CD* have been derived in Theorems 1 and 2, analytic bias correction is feasible. As detailed in Section S.1.1 in the online supplement accompanying this article, estimates of the unknown model components that constitute either Ξ or Φ_1 and Φ_2 can be obtained in order to eliminate the diverging component in *CD*. In addition, Propositions 2 and 3 suggest that these estimated bias terms would eliminate equivalent terms

¹¹This follows because all cited articles draw λ_i and σ_i independently of each other and assume a symmetric distribution for λ_i . Using the integer representation of the expected value it is possible to show that $\text{cov} \left[\lambda_1, \zeta_{v,1}^{-1} \right] = \mathbf{0}$ in this case.

Ξ_{H_1} or $\Phi_{H_1,1}$ and $\Phi_{H_1,2}$ under the alternative hypothesis without necessarily reducing the rate at which CD diverges.

Still, problems with its implementation in practice lead us to discard analytic bias correction and to consider it merely as a benchmark approach when evaluating our weighted CD test statistic, as introduced below, in Monte Carlo experiments. In particular, the construction of plug-in estimates of Φ_1 and Φ_2 proves to be tedious since both terms are constructed from estimates of $\sigma_1^2, \dots, \sigma_N, \lambda_1, \dots, \lambda_N$ and β . Additionally, estimation of the cross-section sum $N^{-1} \sum_{i=1}^N \sigma_i^{-1} \lambda_i$ poses a trade-off between generality and accuracy of the plug-in estimate. To be specific, independence between factor loadings λ_i and error variances σ_i^2 needs to be imposed to ensure that estimation error in $\hat{\lambda}_i$ and $\hat{\sigma}_i^2$ does not dominate the small-sample distribution of the bias-corrected version of CD . The need to impose further assumptions in order to improve the accuracy of the bias estimates is effectively a consequence of the fact that the unknown bias itself diverges at rate \sqrt{T} . This means that any error in the estimation of the two aforementioned components is scaled by a factor that increases as $N, T \rightarrow \infty$. Still, despite improving the approximation of plug-in estimates for Φ_1 and Φ_2 the assumption of independence between factor loadings and error variances can lead to size distortions in DGPs where it is violated.¹²

4.2 Bias correction via weighted cross-section covariances

The method for bias correction favored in this article is to construct a CD test statistic from estimated cross-section covariances that are weighted with individual-specific, random draws from a Rademacher distribution.¹³ This approach is based on noticing that the cross-section correlation estimator $\hat{\rho}_{ij}$, as defined in (3) with $z_{i,t} = \hat{\varepsilon}_{i,t}$, is merely a unit-specific weighting of $\widehat{\text{cov}} \left[\hat{\varepsilon}'_{i,t}, \hat{\varepsilon}_{j,t} \right]$ with advantageous properties: Under the assumptions made in Pesaran (2004), it ensures that the CD test statistic has unit variance under the null hypothesis of no cross-section correlation, allowing for standard normal inference without the need to obtain a variance estimate. However, the case is different in models with unknown, period-specific parameters. As shown in Theorems 1 and 2, studentization of the model residuals fails to ensure a test statistic with a variance of one, even when the true error variances are known. This justifies the use of an alternative weighting scheme which we construct as a weighted average of individual specific covariances,

$$\sqrt{\frac{2}{TN(N-1)}} \sum_{t=1}^T \sum_{i=2}^N \sum_{j=1}^{i-1} (w_i \hat{\varepsilon}_{i,t}) (w_j \hat{\varepsilon}_{j,t}). \quad (21)$$

¹²See the corresponding simulation results in Section 5 for an illustration of the problem.

¹³A Rademacher distributed random variable assumes the values 1 and -1 with probability 0.5 each.

for some set of weights w_1, \dots, w_N . Expression (21) would coincide with the CD test statistic of Pesaran (2004) for $w_i = \hat{\sigma}_i^{-1} \forall i$.¹⁴ This is not the case we consider here. Instead we make the following additional assumption:

Assumption 7 (Weights). w_1, \dots, w_N are identically and independently Rademacher distributed. Furthermore, w_i is independent of $\lambda_i, \sigma_i, \eta_{i,t}$ and $e_{i,t}$ for all i and t .

Rademacher distributed weights amount to random sample splitting, a method for breaking dependence that is not new to econometrics.¹⁵ Its effects are most obvious in a simple modification of Theorem 2 where we replace σ_i^{-1} with w_i . Under Assumption 7, the expected value of $N^{-1} \sum_{\ell=1}^N \lambda_i w_i$ can be conveniently split into the expectations of its two constituents. Given the zero expected value of w_i , an appropriate LLN applies and leads the cross-section average of $\lambda_i w_i$ to converge to zero. In an even simpler form the same reasoning applies to the cross-section averages involving w_i in an adapted version of Theorem 1. In both cases, this reduces the order of the leading bias components Ξ, Φ_1 and Φ_2 by a factor of N and entails asymptotic unbiasedness of the weighted CD test statistic, subject to the restriction $\sqrt{T}/N \rightarrow 0$. Additional rescaling with the asymptotic variance of expression (21) results in a weighted CD test statistic which, as formally stated in Theorem 3 below, converges to a standard normal distribution.

Theorem 3. Consider the weighted CD test statistic

$$CD_W = \left(\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \hat{\varepsilon}_{i,t}^2 w_i^2 \right)^{-1} \left(\sqrt{\frac{2}{TN(N-1)}} \sum_{t=1}^T \sum_{i=2}^N \sum_{j=1}^{i-1} w_i \hat{\varepsilon}_{i,t} w_j \hat{\varepsilon}_{j,t} \right)$$

and assume that either of the following two points hold.

1. The data are generated by the time fixed effects model (4) such that Assumption 1 holds. $\hat{\varepsilon}_{i,t}$ is given by (6).
2. The data are generated by the latent common factor model (8) such that Assumptions 2-6 hold. $\hat{\varepsilon}_{i,t}$ is defined by (28).

Under either of the two sets of assumptions above as well as Assumption 7, it holds that

$$CD_W \xrightarrow{d} N(0, 1),$$

as $N, T \rightarrow \infty$ jointly subject to the restriction $\sqrt{T}N^{-1} \rightarrow 0$.

¹⁴Note that the weighted CD statistic should not be confused with the ‘‘local CD’’ statistic discussed in Pesaran (2015b). In the latter case, the weights are pair specific, i.e. w_{ij} , and are usually motivated using the spatial (distance) structure of the data, see e.g. Robinson (2008). In our case the spatial structure is not needed, thus the weighted test is global and not local.

¹⁵Sample splitting has already been considered an *old* concept in articles as early as (Altonji and Segal 1996, see p. 358)

As stated by Theorem 3, the use of independent Rademacher distributed weights, analogous to weights drawn from many other distributions with zero mean, re-establishes asymptotic standard normal inference under the null hypothesis of the CD test statistic. However, as is often the case for bias-corrected statistics, asymptotic unbiasedness of CD_W comes at the cost of power. More specifically, our approach to bias correction centers the leading components of CD_W around zero, irrespective of whether cross-section correlation in the data is completely controlled for or not. As a consequence, only increases in the variance of CD_W under its alternative hypothesis lead to power against the null hypothesis of this test. This point can be formally illustrated by a minor modification of Proposition 2 which consists of replacing $\zeta_i^{-1} - E[\zeta_i^{-1}]$ with w_i . Given random weights, the leading stochastic component in expression (20),

$$\sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} w_i \boldsymbol{\nu}'_i \boldsymbol{\nu}_j w_j, \quad (22)$$

has an asymptotic mean of zero if $N^{-1}\sqrt{T} \rightarrow 0$ since independent Rademacher distributed weights ensure that $E[w_i \boldsymbol{\nu}_{i,t} \boldsymbol{\nu}'_{j,t} w_j] = 0$ holds for all $i \neq j$. Thus, remaining sources of cross-section co-movement in the composite error term $\boldsymbol{\nu}_{i,t}$ will not shift the location of expression (22). By contrast, it can be shown that the variance of this term continues to diverge at rate T . This entails that CD_W diverges at rate \sqrt{T} under its alternative hypothesis, a property that holds under the conditions of Proposition 2 as well.

To improve the power properties of CD_W we suggest a refinement of this test which follows the power enhancement approach of Fan et al. (2015). The authors suggest improving the power of high-dimensional cross-sectional tests by adding to the test statistic of interest a screening statistic. This screening statistic is equal to zero with probability approaching one under the null hypothesis of the test, but diverges at a fast rate under the alternative. In our case we choose the absolute sum of thresholded cross-section correlation coefficients. This results in a power-enhanced weighted CD test statistic which is defined as

$$CD_{W+} = CD_W + \sum_{i=2}^N \sum_{j=1}^{i-1} |\hat{\rho}_{ij}| \mathbf{1} \left(|\hat{\rho}_{ij}| > 2\sqrt{\ln(N)/T} \right), \quad (23)$$

where $\hat{\rho}_{ij}$ is as in (3) with $z_{i,t} = \hat{\boldsymbol{\epsilon}}_{i,t}$ and where $\mathbf{1}(A)$ is the indicator function for event A . The screening statistic on the right-hand side of (23) has an asymptotically negligible effect on the size of CD_{W+} because individual cross-section correlation coefficients are still consistent for their true value, despite the problems that arise when their values are averaged across all possible combinations of two different cross-sections. Formally, it can be shown that under null hypothesis

$$\hat{\rho}_{ij} = 0 + \mathcal{O}_P(T^{-1/2}) + \mathcal{O}_P((NT)^{-1/2}) + \mathcal{O}_P(N^{-1}), \quad (24)$$

for all pairs i, j . The threshold $2\sqrt{\ln(N)/T}$ is justified by the behavior of the maximum out of $N(N-1)/2$ estimated correlation coefficients in a model with $\rho_{ij} = 0$ for all i, j . Under the

assumption that a CLT holds for $\hat{\rho}_{ij}$ and that $N^{-1}\sqrt{T} \rightarrow 0$ as $N, T, \rightarrow \infty$, this maximum should diverge at rate $2\sqrt{\ln(N)/T}$. Multiplication of each indicator function in the screening statistic $\sum_{i=2}^N \sum_{j=1}^{i-1} |\hat{\rho}_{ij}| \mathbf{1}\left(|\hat{\rho}_{ij}| > 2\sqrt{\ln(N)/T}\right)$ with the absolute value of its corresponding correlation coefficient then ensures that the power enhancement component in (23) converges to 0 if $\rho_{ij} = 0$ for all i, j , subject to additional tail regularity conditions.

Remark 6. An approach to reducing the dependence of CD_W on a specific set of random weights would consist of averaging several weighted CD test statistics. Denote by $CD_W^{(g)}$ the CD test statistic obtained for a given draw of N Rademacher distributed weights, the latter being indexed by g . For a total of G different draws, an averaged weighted CD test can be constructed as

$$\overline{CD}_W = \frac{1}{\sqrt{G}} \sum_{g=1}^G CD_W^{(g)},$$

where the total number of draws G should be chosen sufficiently small (e.g. $G = 30$) to avoid size distortions that may arise from scaling up lower-order terms in $CD_W^{(g)}$.

Remark 7. An initially appealing alternative to external random numbers as weights for a bias-corrected CD test statistic would be a set of N statistics derived from the dataset available to the researcher. When considering such *internal weights*, it is desirable to opt for functions of the data that are optimal in the sense that they maximize the rate at which a weighted CD test statistic diverges for a wide number of alternatives. In A.3 we argue that such improved weights suggest the use of a different test statistic for each alternative. Hence, as far as a bias-corrected version of CD is preferred to the use of a different test, obtaining the former by weighting cross-section covariances with external random weights is the best possible alternative.

5 Monte carlo study

We investigate the properties of the CD test as well as its weighted alternatives in a small set of simulation experiments. Following the exposition of Section 2.1, we simulate the common latent factor model

$$y_{i,t} = \beta x_{i,t} + \boldsymbol{\lambda}'_i \mathbf{f}_t + \sqrt{\sigma_{i,y}^2} \varepsilon_{i,t} \quad (25)$$

$$x_{i,t} = \boldsymbol{\Lambda}'_i \mathbf{f}_t + e_{i,t}. \quad (26)$$

We consider two alternative specifications for the factor loadings on the regressand $y_{i,t}$. We define $\boldsymbol{\lambda}_i = \boldsymbol{\nu}_r + \tilde{\boldsymbol{\lambda}}_i$ where the r elements of $\tilde{\boldsymbol{\lambda}}_i$ are drawn (I) from $U(-0.75, 0.75)$ or (II) from a standardized $\chi^2(2)$ distribution that has zero mean and a variance of $1/6$. The latter case is designed to match the first two moments of $\tilde{\boldsymbol{\lambda}}_i$ in case (I). Concerning the factor loadings on the regressors $x_{i,t}$ we draw element $\Lambda^{(r')}$ of the $r \times 1$ vector $\boldsymbol{\Lambda}_i$ from $U(-0.5, 0.5)$ if $r' = 1$ and from

$U(0.5, 1, 5)$ otherwise. Concerning the latent factors, we let $\mathbf{f}_t \sim N(\mathbf{0}, \mathbf{I}_r)$. When considering the two-way fixed effects model instead of the common latent factor model, we set $r = 2$ and define $\mathbf{f}_t = [f_t^{(1)}, 1]'$ as well as $\boldsymbol{\lambda}_i = [1, \lambda_i^{(2)}]'$ and $\boldsymbol{\Lambda}_i = [1, \Lambda_i^{(2)}]'$ with the construction of $f_t^{(1)}, \lambda_i^{(2)}$ and $\Lambda_i^{(2)}$ being unchanged.

Analogous to our treatment of factor loadings we consider two cases for $\varepsilon_{i,t}$, namely that they are drawn independently over i and t from (i) a standard normal distribution or (ii) a standardized $\chi^2(2)$ distribution that has zero mean and unit variance. The scalar random variable $e_{i,t}$ is generated in the same way as $\varepsilon_{i,t}$. Lastly, error variances are obtained in two different fashions: We set (a) $\sigma_i^2 = c_\sigma (\zeta_i^2 - 2) / \sqrt{24} + 1$ where $\zeta_i^2 \sim \chi^2(2)$ or (b) $\sigma_i^2 = 0.5 + d_\sigma T^{-1} \sum_{t=1}^T (\boldsymbol{\lambda}_i \mathbf{f}_t)^2$. The scaling c_σ , which is only included in setup (a), is chosen manually as discussed below. This construction entails that $E[\sigma_i^2] = 1$ and $\text{var}[\sigma_i^2] = c_\sigma^2/6$. The normalizing constant d_σ , which appears in case (b) only, is $d_\sigma = \sqrt{2}$ in model error case (i) and $d_\sigma = (1/3)^{-1/2}$ in case (b). This scaling ensures that the variance of σ_i^2 across cross-sections is approximately of the same magnitude as in case (a) with $c_\sigma = 1$.

In all following simulations we extend our theoretical framework and assume that β is unknown and needs to be estimated. If the fixed effects estimator is estimated, individual fixed effects are assumed to be unknown as well so that a within transformation for the two-way fixed effects model is applied. Lastly, whenever model residuals are studentized, this is done using estimated error variances instead of their true values.

5.1 Standard CD statistic

In a first instance, we consider the properties of the original CD test statistic when applied to either 2WFE or CCE residuals in a scenario where all sources of cross-section correlation are adequately accounted for. We do this for the setup (i)-(a)-(I) as described above, meaning that errors are normally distributed, error variances independent of factor loadings (or individual fixed effects) and that the latter are symmetrically distributed around their mean. Results for a model with errors from a standardized χ^2 distribution are almost identical and reported in Section S.4 of the online supplement. Furthermore, we consider different degrees of heterogeneity among the individual-specific error variances by reporting results for the values $c_\sigma \in \{0.1, 0.5, 1, 1.5\}$.

Table 1 reports the first two moments of CD when applied either to 2WFE residuals when the true model is one with two-way fixed effects or to CCE residuals in a model with multi-factor error structure and two factors. The results for both cases are identical and show that CD has a bias that diverges towards $-\infty$ as $T \rightarrow \infty$ and a variance that is considerably below 1. The bias term is reasonably close to the benchmark value of $-\sqrt{T/2}$ indicated by Proposition 1 but tends towards zero as heterogeneity among the individual-specific error variances is amplified.

Given that the theoretical results of Sections 3.1 and 3.2 are confirmed by Table 1, we turn

Table 1: Sample moments of original CD test statistic for remaining CSD under H_0

| Part A: Application of CD to 2WFE residuals | | | | | | | | | |
|--|-----------------|-------------|--------|-------|-------|---|------|------|-------|
| N | c_σ T | 0.1 | 0.5 | 1 | 1.5 | 0.1 | 0.5 | 1 | 1.5 |
| | | Mean | | | | Variance $\times 100$ | | | |
| 25 | 25 | -3.53 | -3.50 | -3.41 | -3.24 | 0.09 | 0.20 | 0.82 | 2.96 |
| 25 | 50 | -5.05 | -5.01 | -4.88 | -4.63 | 0.03 | 0.14 | 0.94 | 4.07 |
| 25 | 100 | -7.18 | -7.12 | -6.94 | -6.57 | 0.02 | 0.15 | 1.33 | 6.10 |
| 25 | 200 | -10.18 | -10.09 | -9.83 | -9.30 | 0.01 | 0.21 | 2.12 | 10.48 |
| 200 | 25 | -3.47 | -3.44 | -3.34 | -3.14 | 0.05 | 0.11 | 0.43 | 1.67 |
| 200 | 50 | -4.96 | -4.92 | -4.78 | -4.49 | 0.01 | 0.05 | 0.32 | 1.48 |
| 200 | 100 | -7.05 | -6.99 | -6.79 | -6.39 | 0.00 | 0.04 | 0.37 | 1.85 |
| 200 | 200 | -10.00 | -9.91 | -9.63 | -9.06 | 0.00 | 0.04 | 0.42 | 2.25 |

| Part B: Application of CD to CCE residuals | | | | | | | | | |
|---|-----------------|-------------|--------|-------|-------|---|------|------|------|
| N | c_σ T | 0.1 | 0.5 | 1 | 1.5 | 0.1 | 0.5 | 1 | 1.5 |
| | | Mean | | | | Variance $\times 100$ | | | |
| 25 | 25 | -3.53 | -3.50 | -3.41 | -3.25 | 0.11 | 0.23 | 0.87 | 3.00 |
| 25 | 50 | -5.05 | -5.01 | -4.88 | -4.64 | 0.04 | 0.15 | 0.89 | 3.71 |
| 25 | 100 | -7.18 | -7.12 | -6.95 | -6.61 | 0.02 | 0.15 | 1.25 | 5.58 |
| 25 | 200 | -10.18 | -10.10 | -9.85 | -9.37 | 0.01 | 0.21 | 2.10 | 9.77 |
| 200 | 25 | -3.46 | -3.43 | -3.33 | -3.14 | 0.06 | 0.13 | 0.48 | 1.77 |
| 200 | 50 | -4.96 | -4.91 | -4.77 | -4.50 | 0.02 | 0.06 | 0.35 | 1.63 |
| 200 | 100 | -7.05 | -6.99 | -6.79 | -6.40 | 0.00 | 0.04 | 0.34 | 1.79 |
| 200 | 200 | -10.00 | -9.91 | -9.63 | -9.08 | 0.00 | 0.04 | 0.42 | 2.27 |

Notes. $\varepsilon_{i,t}$ and $e_{i,t}$ are $N(0,1)$. $\sigma_{i,y}^2 = c_\sigma (\zeta_{i,y}^2 - 2) / 4 + 1$; $\zeta_{i,y}^2$ is $\chi^2(2)$. In part B, $r = 2$ and loadings λ_i are drawn from $U(0.5, 1.5)$. Λ_i has first element from $U(0.5, 1.5)$ and second from $U(-0.5, 0.5)$. Factors f_t drawn from $N(0, 1)$. In part A, we restrict $f_t = [f_t^{(1)}, 1]'$, $\lambda_i = [1, \lambda_i^{(2)}]'$ and $\Lambda_i = [1, \Lambda_i^{(2)}]'$.

to the properties of CD under its alternative hypothesis. For this purpose, we simulate a model with multifactor error structure and three factors, implying that neither 2WFE nor CCE estimation completely accounts for all sources of cross-section correlation in the simulated data. Heterogeneity among unit-specific error variances is kept constant by only considering $c_\sigma = 1$. Instead we consider cases (I) and (II) for factor loadings as well as cases (a) and (b) for error variances. Again, results are presented only for normally distributed model errors. The moments of CD are very similar when model errors are drawn from a standardized $\chi^2(2)$ distribution and corresponding tables can be found in Section S.4 in the online supplement accompanying this article.

Tables 2 and 3 report the mean and variance of CD in the presence of remaining cross-section correlation. Both are very similar to the numbers reported in Table 1 as long as factor loadings are symmetrically distributed, an observation which is in line with Propositions 2 and 3. Since symmetric loadings center the leading stochastic component of CD around zero, the latter is dominated by a *bias equivalent* that diverges towards $-\infty$ as $T \rightarrow \infty$. Furthermore, as suggested in Remark 5, the mean of CD is relatively close to the benchmark value $-\sqrt{T/2}$. Qualitatively very different results can be observed when factor loadings are drawn from a skewed distribution, particularly when applying the CD test statistic to 2WFE residuals. As one can observe in columns two and four of Table 2, the strong negative divergence of CD as T increases is mitigated and it is reasonable to assume that it will be turned into positive divergence for a given $N > 200$. This pattern, which is amplified if error variances are a function of factor loadings, reflects the presence of a nonzero mean in the leading stochastic component of CD that diverges to $+\infty$ at rate $N\sqrt{T}$. In the case of 2WFE residuals, its magnitude is sufficiently large to counteract the negative bias equivalent already for $N = 200$. By contrast, columns two and four in Table 3 suggest that this term is considerably smaller when the CD test is applied to CCE residuals since the mean of CD continues to diverge towards $-\infty$ for $N = 200$. We conjecture that the same sign reversal will eventually be obtained even if this case, but that the required cross-section dimension for this to happen is higher.

5.2 Weighted CD statistic

Having documented the first two moments of the CD test statistic, we proceed with investigating the properties of our weighted CD test statistic. As previously, we keep heterogeneity across unit-specific error variances fixed at $c_\sigma = 1$ and consider two different specifications each for factor loadings and error variances. We report size and power for our weighted test statistic CD_W as well as the power-enhanced refinement CD_{W+} . As a benchmark test, we include a CD test statistic with analytic bias correction which corrects CD with sample equivalents of the asymptotic bias terms in Theorems 1 and 2. Details on its implementation can be found in Section S.1.1 in the online supplement. The original CD test is left out for the sake of saving space and in particular since rejection rates are 100% for most cases.

Table 2: Sample moments of original CD test statistic applied to 2WFE residuals under \mathbb{H}_1

| N | λ_i : σ_i^2 : T | Mean | | | | Variance | | | |
|-----|--------------------------------------|-------------------|----------------|-------------------|----------------|-------------------|----------------|-------------------|----------------|
| | | symmetric | | skewed | | symmetric | | skewed | |
| | | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ |
| 25 | 25 | -3.45 | -3.23 | -3.41 | -2.83 | 0.01 | 0.07 | 0.01 | 0.35 |
| 25 | 50 | -4.93 | -4.62 | -4.87 | -4.03 | 0.01 | 0.11 | 0.01 | 0.59 |
| 25 | 100 | -7.00 | -6.56 | -6.92 | -5.72 | 0.01 | 0.21 | 0.01 | 1.12 |
| 25 | 200 | -9.92 | -9.30 | -9.80 | -8.12 | 0.01 | 0.36 | 0.02 | 2.14 |
| 50 | 25 | -3.41 | -3.05 | -3.37 | -2.41 | 0.00 | 0.09 | 0.01 | 0.46 |
| 50 | 50 | -4.87 | -4.36 | -4.81 | -3.44 | 0.00 | 0.15 | 0.01 | 0.78 |
| 50 | 100 | -6.93 | -6.19 | -6.83 | -4.89 | 0.01 | 0.27 | 0.01 | 1.45 |
| 50 | 200 | -9.82 | -8.80 | -9.69 | -6.95 | 0.01 | 0.48 | 0.01 | 2.61 |
| 100 | 25 | -3.40 | -2.75 | -3.35 | -1.68 | 0.00 | 0.17 | 0.01 | 0.73 |
| 100 | 50 | -4.85 | -3.95 | -4.78 | -2.39 | 0.00 | 0.26 | 0.01 | 1.19 |
| 100 | 100 | -6.89 | -5.61 | -6.80 | -3.39 | 0.01 | 0.42 | 0.01 | 2.17 |
| 100 | 200 | -9.76 | -7.95 | -9.63 | -4.78 | 0.01 | 0.76 | 0.01 | 4.01 |
| 200 | 25 | -3.39 | -2.20 | -3.34 | -0.23 | 0.00 | 0.36 | 0.01 | 1.52 |
| 200 | 50 | -4.83 | -3.16 | -4.77 | -0.34 | 0.00 | 0.49 | 0.01 | 2.23 |
| 200 | 100 | -6.87 | -4.48 | -6.78 | -0.45 | 0.00 | 0.82 | 0.01 | 3.78 |
| 200 | 200 | -9.74 | -6.38 | -9.60 | -0.66 | 0.01 | 1.35 | 0.01 | 6.92 |

Notes. The model has a factor error structure with 3 factors. $\varepsilon_{i,t}$ and $e_{i,t}$ are $N(0,1)$. " σ_i^2 : $\perp \lambda_i$ " means that $\sigma_{i,y}^2 = (\zeta_{i,y}^2 - 2) / 4 + 1$ where $\zeta_{i,y}^2$ is $\chi^2(2)$. For " σ_i^2 : $f(\lambda_i)$ ", we let $\sigma_{i,y}^2 = d_\sigma T^{-1} \sum_{t=1}^T (\lambda_i' \mathbf{f}_t)^2$. The case " λ_i : symmetric" corresponds to drawing λ_i from $U(0.5, 1.5)$. " λ_i : skewed" loadings are from a standardized $\chi^2(2)$ distribution with mean and variance equal to 1. Λ_i has first element from $U(0.5, 1.5)$ and all others from $U(-0.5, 0.5)$. Factors \mathbf{f}_t are drawn from $N(0, 1)$.

Table 3: Sample moments of original CD test statistic applied to CCE residuals under \mathbb{H}_1

| | | Mean | | | | Variance | | | |
|----------------|-----|-------------------|----------------|-------------------|----------------|-------------------|----------------|-------------------|----------------|
| λ_i : | | symmetric | | skewed | | symmetric | | skewed | |
| σ_i^2 : | | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ |
| N | T | | | | | | | | |
| 25 | 25 | -3.42 | -3.38 | -3.42 | -3.21 | 0.01 | 0.02 | 0.01 | 0.11 |
| 25 | 50 | -4.89 | -4.85 | -4.89 | -4.58 | 0.01 | 0.02 | 0.01 | 0.18 |
| 25 | 100 | -6.95 | -6.89 | -6.95 | -6.52 | 0.01 | 0.03 | 0.01 | 0.35 |
| 25 | 200 | -9.86 | -9.77 | -9.85 | -9.23 | 0.02 | 0.05 | 0.01 | 0.68 |
| 50 | 25 | -3.38 | -3.34 | -3.38 | -3.17 | 0.01 | 0.01 | 0.01 | 0.07 |
| 50 | 50 | -4.83 | -4.78 | -4.83 | -4.52 | 0.01 | 0.01 | 0.01 | 0.12 |
| 50 | 100 | -6.87 | -6.80 | -6.86 | -6.43 | 0.01 | 0.02 | 0.01 | 0.23 |
| 50 | 200 | -9.74 | -9.65 | -9.74 | -9.12 | 0.01 | 0.03 | 0.01 | 0.39 |
| 100 | 25 | -3.36 | -3.32 | -3.35 | -3.15 | 0.01 | 0.01 | 0.01 | 0.05 |
| 100 | 50 | -4.80 | -4.76 | -4.80 | -4.50 | 0.00 | 0.01 | 0.00 | 0.08 |
| 100 | 100 | -6.83 | -6.76 | -6.83 | -6.40 | 0.01 | 0.01 | 0.00 | 0.14 |
| 100 | 200 | -9.68 | -9.59 | -9.68 | -9.07 | 0.01 | 0.02 | 0.01 | 0.25 |
| 200 | 25 | -3.35 | -3.31 | -3.35 | -3.14 | 0.00 | 0.01 | 0.00 | 0.05 |
| 200 | 50 | -4.79 | -4.74 | -4.79 | -4.48 | 0.00 | 0.01 | 0.00 | 0.06 |
| 200 | 100 | -6.81 | -6.74 | -6.80 | -6.38 | 0.00 | 0.01 | 0.00 | 0.10 |
| 200 | 200 | -9.65 | -9.56 | -9.65 | -9.05 | 0.01 | 0.02 | 0.01 | 0.19 |

Notes. See Table 2.

Table 4 reports the size and power of all three bias corrected test statistic when applied to 2WFE residuals. The size of CD_W and CD_{W+} is very close to the nominal level of 5% as long as $N \geq T$. Size distortions are given in cases where T is considerably large than N and the effect of power enhancement on size is generally negligible. The analytically bias-corrected CD test statistic CD_{BC} exhibits hardly any tendency to over-reject but is in general rather conservative, in particular when T is small.

Panel B in Table 4 report results on power. Here, we see that the power of CD_W is in general low and increases only in T . This is improved upon considerably by power enhancement, leading the refined test statistic CD_{W+} to reliably reject when it should as long as the number of time periods is large enough. The performance of CD_{W+} is considerably above that of the benchmark statistic CD_{BC} when factor loadings are drawn from a symmetric distribution and is on par in most other cases. An exception is the case of skewed loadings with small T and large N where CD_{BC} has markedly higher rejection rates. It can also be noted that the performance of CD_{BC} largely depends on whether factor loadings are drawn from a symmetric distribution or not. If this is the case, analytic bias correction leads to rejection rates under the alternative hypothesis that are worst among all three tests considered.

When testing for remaining cross-section correlation in CCE residuals we observe similar results as in Table 4. Interestingly, it can be observed that the benchmark statistic CD_{BC} exhibits size distortions in samples with large T when loadings are drawn from a skewed distribution

and if factor loadings and error variances are dependent. This results from the nature of our bias correction which assumes independence between σ_i^2 and λ_i to considerably improve the accuracy of the bias estimate.

The power properties of CD_W and CD_{W+} when either of these tests is applied to CCE residuals mirror those seen in the 2WFE case, even though rejection rates are generally somewhat lower. However, the power-enhanced statistic CD_{W+} now performs best in all cases without ever being inferior to CD_{BC} .

6 Empirical illustration: R& D investments

In this section we illustrate the applicability of the standard and the weighted CD statistics using the R&D investments data of Eberhardt et al. (2013). In this application serial correlation is important from an economic point of view. Hence, Section S.1.3 in the online supplement outlines how the CD statistic can be modified to account for serial correlation, given a set of known weights w_i .

Eberhardt et al. (2013) question whether R&D can be estimated in a standard Griliches-type “knowledge production function” framework ignoring the potential presence of knowledge spillovers between cross-sectional units as well as other cross-section dependencies. Among other things they document a “[...] strong evidence for cross-sectional dependence and the presence of a common factor structure in the data, which we interpret as indicative for the presence of knowledge spillovers and additional unobserved cross-sectional dependencies” (Eberhardt et al. 2013, p.437). Cross-sectional dependence was measured by means of a CD test.

The original dataset used information on up to twelve manufacturing industries in ten countries¹⁶ over a time period of up to 26 years from 1980 to 2005. All of the results presented assume the country-industry pair as the unit of analysis (panel group member i), of which we have $N = 119$. This panel is unbalanced where for Germany, Portugal, and Sweden the length of the available time-series is substantially shorter than for other countries.

Some of our results can be extended to cover unbalanced setups. However in order to simplify our analysis we disregard from using the observations for Germany, Portugal, Sweden in the construction of the test statistics. Additionally, we remove two sectors from the Great Britain where observations for $t = 2004, 2005$ are missing. This way we are left with $N = 82$ and $T = 25$.

In this section we will primarily revisit some of the results in Table 5 of Eberhardt et al. (2013) for pooled (static) production function estimates. Our goal is to investigate how the divergent properties of the CD test statistic might have influenced the choice between the First Difference (FD) estimator with yearly dummies and Pooled CCE (CCEP) estimators, as presented in columns 3 and 4, respectively, of the table mentioned above. Which of the two

¹⁶Denmark, Finland, Germany, Italy, Japan, Netherlands, Portugal, Sweden, United Kingdom, and the US.

Table 4: Rejection rates for weighted CD test statistic when applied to 2WFE residuals.

Part A: Size

| N | λ_i : σ_i^2 : T | symmetric | | | | | | skewed | | | | | |
|-----|--------------------------------------|-------------------|-----------|-----------|----------------|-----------|-----------|-------------------|-----------|-----------|----------------|-----------|-----------|
| | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | |
| | | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} |
| 25 | 25 | 5.3 | 5.5 | 2.7 | 6.2 | 6.4 | 2.8 | 5.3 | 5.5 | 3.1 | 5.2 | 5.3 | 2.5 |
| 25 | 50 | 6.6 | 6.9 | 4.3 | 5.8 | 6.1 | 4.1 | 6.2 | 6.8 | 4.0 | 6.3 | 6.4 | 2.7 |
| 25 | 100 | 7.8 | 8.5 | 3.7 | 7.0 | 7.1 | 4.6 | 6.5 | 6.9 | 4.5 | 6.4 | 6.6 | 2.2 |
| 25 | 200 | 9.8 | 10.3 | 5.6 | 9.0 | 10.0 | 6.1 | 9.3 | 9.6 | 6.1 | 8.9 | 9.4 | 3.3 |
| 50 | 25 | 4.6 | 4.7 | 2.8 | 4.7 | 4.7 | 3.7 | 4.6 | 4.6 | 2.9 | 5.9 | 5.9 | 2.9 |
| 50 | 50 | 6.1 | 6.2 | 3.4 | 5.2 | 5.4 | 5.0 | 5.4 | 5.3 | 4.4 | 4.5 | 4.6 | 2.7 |
| 50 | 100 | 5.5 | 5.9 | 4.7 | 5.4 | 5.8 | 4.8 | 5.8 | 6.0 | 4.2 | 5.1 | 5.6 | 3.1 |
| 50 | 200 | 6.0 | 6.5 | 4.6 | 6.1 | 6.3 | 4.9 | 6.4 | 6.2 | 4.7 | 5.7 | 5.7 | 3.4 |
| 100 | 25 | 5.0 | 5.0 | 2.8 | 5.2 | 5.2 | 3.4 | 5.9 | 5.9 | 3.8 | 4.7 | 4.7 | 2.9 |
| 100 | 50 | 5.0 | 5.1 | 3.5 | 4.6 | 4.8 | 3.9 | 4.4 | 4.5 | 4.0 | 5.0 | 5.1 | 2.9 |
| 100 | 100 | 5.3 | 5.3 | 5.1 | 5.1 | 5.1 | 3.9 | 4.8 | 5.0 | 3.7 | 5.5 | 5.5 | 3.5 |
| 100 | 200 | 4.5 | 4.5 | 4.7 | 5.4 | 5.3 | 4.5 | 5.8 | 5.6 | 4.0 | 6.2 | 6.3 | 3.1 |
| 200 | 25 | 5.5 | 5.5 | 3.3 | 5.5 | 5.5 | 3.6 | 4.7 | 4.7 | 3.4 | 4.7 | 4.7 | 3.2 |
| 200 | 50 | 5.4 | 5.5 | 4.4 | 5.5 | 5.6 | 3.2 | 5.5 | 5.6 | 4.2 | 4.7 | 4.8 | 3.6 |
| 200 | 100 | 4.3 | 4.3 | 4.6 | 6.1 | 6.2 | 4.4 | 4.5 | 4.5 | 4.6 | 4.9 | 4.9 | 3.2 |
| 200 | 200 | 4.5 | 4.8 | 4.3 | 5.1 | 5.3 | 4.6 | 5.1 | 5.1 | 4.7 | 6.2 | 6.3 | 3.4 |

Part B: Power

| N | λ_i : σ_i^2 : T | symmetric | | | | | | skewed | | | | | |
|-----|--------------------------------------|-------------------|-----------|-----------|----------------|-----------|-----------|-------------------|-----------|-----------|----------------|-----------|-----------|
| | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | |
| | | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} |
| 25 | 25 | 12.4 | 18.5 | 8.6 | 10.3 | 11.4 | 7.7 | 11.5 | 16.3 | 17.8 | 12.5 | 14.9 | 27.4 |
| 25 | 50 | 21.9 | 81.7 | 16.3 | 17.0 | 38.9 | 11.1 | 18.9 | 52.9 | 35.1 | 24.2 | 56.9 | 50.7 |
| 25 | 100 | 36.2 | 100 | 25.9 | 27.2 | 92.5 | 22.1 | 31.0 | 90.6 | 50.7 | 38.5 | 95.8 | 71.1 |
| 25 | 200 | 51.4 | 100 | 41.7 | 43.9 | 100 | 33.5 | 44.3 | 99.4 | 67.0 | 52.8 | 99.9 | 83.1 |
| 50 | 25 | 12.3 | 15.5 | 8.5 | 10.8 | 11.1 | 6.7 | 12.3 | 17.7 | 42.3 | 14.2 | 14.5 | 65.1 |
| 50 | 50 | 20.3 | 96.3 | 15.9 | 16.7 | 39.9 | 12.5 | 19.8 | 79.1 | 65.6 | 22.8 | 66.6 | 89.1 |
| 50 | 100 | 34.7 | 100 | 26.7 | 27.9 | 99.2 | 24.2 | 32.2 | 99.8 | 85.0 | 37.1 | 99.9 | 97.2 |
| 50 | 200 | 52.9 | 100 | 41.4 | 43.1 | 100 | 36.9 | 47.0 | 100 | 92.0 | 54.7 | 100 | 99.2 |
| 100 | 25 | 10.7 | 11.0 | 8.4 | 9.3 | 9.4 | 6.7 | 11.3 | 14.9 | 81.1 | 13.3 | 13.5 | 96.8 |
| 100 | 50 | 20.3 | 98.9 | 14.9 | 17.2 | 40.5 | 12.8 | 18.1 | 97.1 | 96.6 | 20.8 | 81.8 | 99.9 |
| 100 | 100 | 33.4 | 100 | 26.4 | 27.9 | 100 | 22.8 | 29.1 | 100 | 99.6 | 34.2 | 100 | 100 |
| 100 | 200 | 50.2 | 100 | 41.5 | 41.9 | 100 | 39.2 | 47.3 | 100 | 100 | 52.4 | 100 | 100 |
| 200 | 25 | 11.6 | 11.6 | 8.8 | 11.3 | 11.3 | 6.9 | 11.5 | 12.2 | 98.6 | 12.4 | 12.4 | 100 |
| 200 | 50 | 19.2 | 99.6 | 16.2 | 15.2 | 34.2 | 13.4 | 18.5 | 99.9 | 100 | 23.8 | 87.0 | 100 |
| 200 | 100 | 34.9 | 100 | 26.4 | 25.5 | 100 | 22.8 | 30.3 | 100 | 100 | 37.8 | 100 | 100 |
| 200 | 200 | 49.8 | 100 | 44.3 | 40.7 | 100 | 38.5 | 45.9 | 100 | 100 | 52.3 | 100 | 100 |

Notes. In Part A the model has two factors which are restricted as noted in Table 1. Part B corresponds to a model with factor error structure and 3 factors. For details on all other model parameters, see Table 2.

CD_W is the weighted CD test statistic introduced in Theorem 3. CD_{W+} is its power-enhanced refinement. CD_{BC} is a CD test statistic with analytic bias correction.

Table 5: Rejection rates for weighted CD test statistic when applied to CCE residuals.

| Part A: Size | | | | | | | | | | | | | |
|--------------|--------------------------------------|-------------------|-----------|-----------|----------------|-----------|-----------|-------------------|-----------|-----------|----------------|-----------|-----------|
| N | λ_i : σ_i^2 : T | symmetric | | | | | | skewed | | | | | |
| | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | |
| | | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} |
| 25 | 25 | 5.9 | 5.9 | 4.2 | 5.3 | 5.6 | 4.6 | 5.3 | 5.4 | 4.3 | 5.9 | 6.3 | 5.8 |
| 25 | 50 | 6.4 | 6.9 | 4.5 | 5.7 | 6.2 | 5.3 | 6.2 | 6.8 | 4.9 | 7.6 | 8.1 | 7.3 |
| 25 | 100 | 8.3 | 9.0 | 6.1 | 7.9 | 8.4 | 6.6 | 7.9 | 9.0 | 7.0 | 9.2 | 9.9 | 12.9 |
| 25 | 200 | 11.2 | 12.2 | 6.8 | 10.3 | 11.3 | 8.2 | 11.4 | 11.7 | 9.5 | 12.4 | 14.0 | 22.0 |
| 50 | 25 | 5.9 | 6.1 | 4.1 | 6.2 | 6.4 | 5.0 | 6.3 | 6.4 | 4.4 | 6.6 | 6.6 | 4.3 |
| 50 | 50 | 6.4 | 6.7 | 4.4 | 5.8 | 5.9 | 4.0 | 5.5 | 5.7 | 3.9 | 6.2 | 6.4 | 5.3 |
| 50 | 100 | 6.1 | 6.3 | 5.4 | 5.9 | 6.0 | 5.9 | 6.1 | 6.1 | 4.7 | 5.7 | 5.9 | 10.4 |
| 50 | 200 | 7.3 | 7.2 | 5.0 | 6.4 | 6.3 | 6.5 | 7.2 | 7.6 | 6.3 | 7.6 | 7.9 | 19.5 |
| 100 | 25 | 5.2 | 5.2 | 4.5 | 5.5 | 5.5 | 4.2 | 5.4 | 5.4 | 4.2 | 5.3 | 5.3 | 4.2 |
| 100 | 50 | 5.0 | 5.1 | 3.8 | 4.8 | 4.9 | 4.9 | 5.3 | 5.5 | 5.9 | 5.2 | 5.2 | 5.6 |
| 100 | 100 | 5.6 | 5.8 | 4.9 | 6.0 | 5.9 | 5.7 | 4.6 | 4.8 | 4.0 | 7.0 | 7.0 | 8.6 |
| 100 | 200 | 5.4 | 5.5 | 4.9 | 5.9 | 5.9 | 6.2 | 5.7 | 6.0 | 5.1 | 5.8 | 5.8 | 18.6 |
| 200 | 25 | 5.3 | 5.3 | 3.8 | 5.1 | 5.1 | 3.9 | 5.3 | 5.3 | 4.5 | 5.4 | 5.4 | 5.1 |
| 200 | 50 | 4.3 | 4.4 | 4.4 | 4.9 | 4.9 | 4.9 | 5.3 | 5.3 | 4.2 | 5.0 | 5.0 | 5.3 |
| 200 | 100 | 4.5 | 4.6 | 4.5 | 5.6 | 5.6 | 4.4 | 5.2 | 5.4 | 5.5 | 6.3 | 6.4 | 9.7 |
| 200 | 200 | 4.4 | 4.7 | 5.1 | 5.8 | 5.8 | 5.1 | 5.2 | 5.3 | 5.2 | 5.0 | 5.1 | 16.5 |

| Part B: Power | | | | | | | | | | | | | |
|---------------|--------------------------------------|-------------------|-----------|-----------|----------------|-----------|-----------|-------------------|-----------|-----------|----------------|-----------|-----------|
| N | λ_i : σ_i^2 : T | symmetric | | | | | | skewed | | | | | |
| | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | |
| | | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} |
| 25 | 25 | 10.0 | 11.8 | 5.4 | 8.9 | 9.5 | 6.2 | 9.0 | 12.2 | 7.4 | 10.5 | 11.4 | 7.4 |
| 25 | 50 | 13.8 | 39.6 | 10.4 | 12.5 | 18.9 | 9.4 | 13.4 | 31.4 | 13.6 | 16.5 | 29.6 | 16.1 |
| 25 | 100 | 21.3 | 81.9 | 15.4 | 17.1 | 44.4 | 14.6 | 19.0 | 62.2 | 20.2 | 24.5 | 63.1 | 28.1 |
| 25 | 200 | 31.8 | 97.5 | 22.9 | 26.1 | 85.1 | 20.8 | 30.3 | 84.6 | 28.8 | 36.5 | 90.2 | 43.2 |
| 50 | 25 | 9.6 | 11.2 | 5.9 | 8.0 | 8.3 | 6.9 | 9.0 | 11.2 | 7.8 | 10.6 | 10.8 | 9.3 |
| 50 | 50 | 14.3 | 54.2 | 8.6 | 10.3 | 16.0 | 8.4 | 11.7 | 52.8 | 12.7 | 15.2 | 34.1 | 20.0 |
| 50 | 100 | 19.6 | 98.5 | 13.7 | 16.5 | 59.5 | 15.3 | 17.8 | 90.5 | 23.1 | 22.2 | 86.4 | 29.7 |
| 50 | 200 | 29.4 | 100 | 23.0 | 20.8 | 98.3 | 24.5 | 27.8 | 99.2 | 34.5 | 32.7 | 100 | 45.2 |
| 100 | 25 | 8.2 | 8.1 | 5.1 | 8.4 | 8.4 | 5.3 | 8.1 | 9.7 | 7.0 | 9.7 | 9.7 | 10.9 |
| 100 | 50 | 12.3 | 64.8 | 9.4 | 10.9 | 15.5 | 10.0 | 11.2 | 74.7 | 15.3 | 13.4 | 38.6 | 20.5 |
| 100 | 100 | 20.2 | 99.9 | 15.7 | 15.5 | 76.4 | 15.3 | 16.6 | 99.4 | 24.9 | 20.2 | 97.3 | 35.6 |
| 100 | 200 | 29.3 | 100 | 23.8 | 22.8 | 100 | 24.8 | 26.0 | 100 | 40.3 | 32.0 | 100 | 50.4 |
| 200 | 25 | 8.9 | 8.9 | 7.0 | 8.2 | 8.2 | 5.9 | 8.8 | 9.0 | 8.6 | 9.9 | 10.0 | 10.8 |
| 200 | 50 | 13.7 | 72.9 | 8.3 | 10.8 | 15.0 | 10.5 | 11.7 | 90.4 | 15.8 | 14.1 | 39.6 | 18.5 |
| 200 | 100 | 17.5 | 100 | 14.6 | 13.4 | 86.7 | 14.5 | 17.0 | 100 | 27.5 | 20.4 | 99.7 | 32.7 |
| 200 | 200 | 29.7 | 100 | 22.8 | 22.5 | 100 | 25.3 | 25.7 | 100 | 45.4 | 32.1 | 100 | 52.6 |

Notes. The model has as factor error structure with two (Part A) or three (Part B) factors. For details on all other model parameters, see Table 2. For an explanation of the tests, see Table 4.

models is considered to be correctly specified has important consequences for the conclusions that can be drawn from the entire table. With regard to the coefficient of private R&D investments, Eberhardt et al. (2013) report that a significance test for the corresponding slope coefficient cannot reject the null hypothesis in the FD model while it can in the CCE model.

Table 6: Cross-sectional dependence testing with R& D data

| Estimator | Serial Correlation | CD | $q_{0.1}(CD_W)$ | $q_{0.5}(CD_W)$ | $q_{0.9}(CD_W)$ | \overline{CD}_W | \overline{CD}_{W+} |
|-----------|--------------------|-------|-----------------|-----------------|-----------------|-------------------|----------------------|
| FD | No | -1.90 | -1.41 | 0.04 | 0.78 | -0.23 | 1.49 |
| | Yes | NA | -1.24 | 0.04 | 0.69 | -0.20 | 1.52 |
| CCEP | No | -2.95 | -1.72 | -0.54 | 1.72 | -1.30 | 7.38 |
| | Yes | NA | -0.91 | -0.29 | 0.91 | -0.69 | 7.98 |

Notes. All results are obtained for Weighted Covariance Bias-corrected CD statistic based on $G = 30$ Rademacher draws of w_i . CD_{BC} denotes the analytical bias-corrected statistic. Serial Correlation with "Yes" option stands for the adjusted test statistic robust to the serial correlation, as described in Section S.1.3 in the online supplement. $q_\tau(CD_W)$ denotes the τ 'th sample quantile of CD_W from $G = 30$ draws.

As the results for CD_W statistic allowing for serial-correlation correction, are similar to those without serial correlation, we only focus on the latter option. First of all, as we can see from Table 6 the conclusions that we can draw from the original CD test are almost identical to those Eberhardt et al. (2013), despite adjustments made in terms of the sample size. In particular, while the value of CD statistic based on CCE residuals imply rejection of the null hypothesis, no such conclusion is implied by FD (at least at a 5% significance level). However, as given that the original test statistic suffers from the IPP, we further investigate if this conclusion also holds after proper bias-corrections. Motivated by finite sample evidence in Section 5, we consider only the CD_W statistic.¹⁷

Taking our attention to the proposed CD_W statistic based on random Rademacher weights, we can see that the corresponding values of \overline{CD}_W indicate that both the FD and CCEP models generate residuals that are cross-sectionally uncorrelated. As we discussed in Section 5, this conclusion might be partially attributed to the fact that for small values of T , the proposed statistic might lack power. For this reason, we also make use of the power-enhanced version \overline{CD}_{W+} of our test statistic. Notice that for this dataset $2\sqrt{\ln(N)/T} \approx 0.84$, thus power-enhancement will pick up only very large values of the correlation coefficients that otherwise might be averaged away by the original CD statistic. As we can see from the corresponding column in Table 6, power enhancement does not alter the conclusion we draw for the FD residuals, as there are only 2 coefficients $\hat{\rho}_{ij}$ above the threshold. However, for CCEP residuals the effect of power enhancement is non-negligible as there are 10 values of $\hat{\rho}_{ij}$ above the threshold.

These results indicate that after appropriate adjustments, the empirical evidence presented in Eberhardt et al. (2013) that favour simple First-difference estimator, as opposed to the CCEP

¹⁷The CD_{BC} option can be easily obtained for the FD estimator without serial-correlation adjustment, with the corresponding value of 1.15.

estimator, remain in place. As we can see from Table 6, this conclusion is unaltered after appropriate serial-correlation adjustments, thus serial-correlation cannot be the main factor affecting this ranking (unlike the discussion in Eberhardt et al. (2013)). Instead, our results might indicate that the underlying CCE restrictions (rank condition and regressors with finite factor structure) might be violated, as these restrictions are irrelevant for the FD estimator with time-effects only.

7 Conclusion

This article documents how the estimation of common time-specific parameters using panel data causes the CD test of Pesaran (2004, 2015) to break down. Using commonly used additive and multiplicative specifications for individual- and time-specific components in the model errors, we show that the CD test statistic applied to residuals of correctly specified regression models is divergent under null hypothesis of cross-sectional independence. We can find an equivalent term under the alternative hypothesis which may balance out the leading diverging component of CD and can lead to low power in small samples. The results documented in this article are interpreted as a manifestation of the incidental parameter problem (IPP) since they ultimately follow from the estimation of T period-specific parameters. Our main theorems illustrate the pervasive nature of the IPP in this setup, given that the consequences of estimating time specific parameters do not disappear as the sample size increases.

Our analyzed weighted CD test statistic achieves our primary goal of re-establishing asymptotic standard normal inference and hence constitutes an alternative to popular bias correction methods which circumvents the problems these approaches have in the present context. Our results have far reaching implications for empirical panel data analysis, where CD test has been widely used as a model selection/diagnostic tool. An illustration of how our theoretical results translate into applications is given via simulations and real datasets.

Finally, in this paper we assumed that the parameters in the linear model are estimated using a least squares objective function. If one deviates from this setup, and instead uses an (over-identified) GMM criterion function to estimate parameters, the usual GMM J-statistic is readily available for the purpose of testing residual cross-sectional correlation. Examples in a fixed- T framework are given by Sarafidis et al. (2009), Ahn et al. (2013), and Juodis (2018) among others. In the large N, T setup average J-statistic as a model specification tool was explicitly used e.g. by Everaert and Pozzi (2014) for the CCE-GMM estimator. Hence given the scale of the problems with CD statistic documented in this paper, these alternative procedures (if applicable) are more appropriate.

Appendix

Notation

Extending the paragraph on notation in introduction in the main text, we will use the following notation in this appendix.

- I_m denotes an $m \times m$ identity matrix and the subscript is sometimes disregarded from for the sake of simplicity. $\mathbf{0}$ denotes a column vector of zeros while \mathbf{O} stands for a matrix of zeros. s_m denotes a selection vector all of whose elements are zero except for element m which is one. $\mathbf{1}$ is a vector entirely consisting of ones. The dimension of these latter vectors and matrices is generally suppressed for the sake of simplicity and needs to be inferred from context.
- For a generic $m \times n$ matrix A , $P_A = A(A'A)^{-1}A'$ projects onto the space spanned by the columns of A and $M_A = I_m - P_A$. $\dim(A)$, $\text{col}(A)$ and $\text{ker}(A)$ refer to the dimension, column space and kernel of A . Moreover, $\text{rk}(A)$ denotes the rank of A , $\text{tr}(A)$ its trace and $\|A\| = (\text{tr}(A'A))^{1/2}$ the Frobenius norm of A .
- For a set of $m \times n$ matrices $\{A_1, \dots, A_N\}$, $\bar{A} = N^{-1} \sum_{i=1}^N A_i$. Multiple sums are generally abbreviated, so that $\sum_{i,j}^N$ is shorthand notation for $\sum_{i=1}^N \sum_{j=1}^N$.
- δ and M stand for a small and large positive real number, respectively. For two real numbers a and b , $a \vee b = \max\{a, b\}$.
- For some random variable $\varepsilon_{i,t}$, $\kappa_4[\varepsilon_{i,t}]$ denotes its fourth-order cumulant. Moreover, $\mathcal{O}(\cdot)$ and $\text{o}(\cdot)$ express order of magnitude relations whereas $\mathcal{O}_p(\cdot)$ and $\text{o}_p(\cdot)$ denote stochastic order relations (see e.g. White 2001, Definitions 2.5 and 2.33).

A Additional results for a model with multifactor error structure

A.1 Heuristics

As in the last paragraph of Section 2.1 consider the $T \times 1$ vector \mathbf{y}_i and the $T \times m$ matrix \mathbf{X}_i defined as

$$\begin{aligned} \mathbf{y}_i &= \mathbf{X}_i \boldsymbol{\beta} + \mathbf{F} \boldsymbol{\lambda}_i + \boldsymbol{\varepsilon}_i \\ \mathbf{X}_i &= \mathbf{F} \boldsymbol{\Lambda}_i + \mathbf{E}_i. \end{aligned} \tag{27}$$

We assume that the unknown vector of slope coefficients β is estimated using the Pooled Common Correlated Effects (CCE) estimator, given by

$$\hat{\beta}^{CCE} = \left(\sum_{i=1}^N \mathbf{X}'_i \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{X}_i \right)^{-1} \left(\sum_{i=1}^N \mathbf{X}'_i \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{y}_i \right)$$

with

$$\hat{\mathbf{F}} = \begin{bmatrix} \bar{\mathbf{y}} & \bar{\mathbf{X}} \end{bmatrix} = \left(\mathbf{F} \begin{bmatrix} \bar{\boldsymbol{\lambda}} & \bar{\mathbf{A}} \end{bmatrix} + \begin{bmatrix} \bar{\boldsymbol{\varepsilon}} & \bar{\mathbf{E}} \end{bmatrix} \right) \begin{bmatrix} \mathbf{1} & \mathbf{0} \\ \boldsymbol{\beta} & \mathbf{I} \end{bmatrix} = \mathbf{F}\bar{\mathbf{C}} + \bar{\mathbf{U}}.$$

The CCE estimator is consistent and asymptotically normal under appropriate assumptions, see e.g. Pesaran (2006). Moreover, even though the factor estimates are effectively *generated regressors*, asymptotic inference about the true value of β is not affected since the effect of having to estimate the unobserved true factors is negligible as the sample size increases. More specifically,

$$\hat{\mathbf{f}}_t - \bar{\mathbf{C}}' \mathbf{f}_t = \frac{1}{N} \sum_{i=1}^N \mathbf{u}_{i,t} = \mathcal{O}_P(N^{-1/2})$$

holds for each row of $\hat{\mathbf{F}}$ by application of an appropriate Central Limit Theorem (CLT). However, despite not being a problem for testing hypotheses about the values of β in model (27), this result leads the CD test statistic, constructed from CCE residuals, to diverge at rate \sqrt{T} . To appreciate this result, note that the residuals obtained from CCE estimation of model (27) are given by

$$\begin{aligned} \hat{\boldsymbol{\varepsilon}}_i &= \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{y}_i - \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{X}_i \hat{\boldsymbol{\beta}}^{CCE} \\ &= \boldsymbol{\varepsilon}_i - \mathbf{P}_{\hat{\mathbf{F}}} \boldsymbol{\varepsilon}_i - (\mathbf{P}_{\hat{\mathbf{F}}} - \mathbf{P}_{\mathbf{F}}) \mathbf{F} \boldsymbol{\lambda}_i - \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{X}_i (\hat{\boldsymbol{\beta}}^{CCE} - \boldsymbol{\beta}_0). \end{aligned} \quad (28)$$

This representation involves two terms that are exclusively due to the fact that the model of interest is augmented with estimates of the unobserved common factors. The first of these two terms is a projection of the true errors onto the space spanned by the estimated factors. Given that the estimated factors are a consistent estimator of a rotation of the true common factors, and knowing that \mathbf{F} and $\boldsymbol{\varepsilon}_i$ are independent by standard assumptions in the literature, $\mathbf{P}_{\hat{\mathbf{F}}} \boldsymbol{\varepsilon}_i$ is asymptotically negligible.

The same reasoning can be applied to $(\mathbf{P}_{\hat{\mathbf{F}}} - \mathbf{P}_{\mathbf{F}}) \mathbf{F} \boldsymbol{\lambda}_i$, the direct impact of the true factors on \mathbf{y}_i that remains because only an imperfect estimate of the factors can be used to account for their effect. Again, pointwise consistency of $\hat{\mathbf{F}}$ suggests that this term is asymptotically negligible. However, the rate of convergence of $\hat{\mathbf{F}}$ to $\mathbf{F}\bar{\mathbf{C}}$ is not fast enough to ensure asymptotic negligibility of this term when constructing a test statistic for cross-section dependence. In particular, the CD test statistic (2) contains a term of the form

$$\sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} T^{-1} \left(\frac{\boldsymbol{\lambda}_i}{\sigma_i} \right)' \mathbf{F}' (\mathbf{P}_{\hat{\mathbf{F}}} - \mathbf{P}_{\mathbf{F}}) \mathbf{F} \left(\frac{\boldsymbol{\lambda}_j}{\sigma_j} \right).$$

This term can be equivalently written

$$\sqrt{T} \operatorname{tr} \left[\left(T^{-1} \mathbf{F}' (\mathbf{P}_{\hat{\mathbf{F}}} - \mathbf{P}_{\mathbf{F}}) \mathbf{F} \right) \left(\sqrt{\frac{2}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \left(\frac{\lambda_j}{\sigma_j} \right) \left(\frac{\lambda_i}{\sigma_i} \right)' \right) \right].$$

Since the second component in parentheses is general of order $\mathcal{O}_p(N)$,¹⁸ the whole expression converges to zero only if the middle term

$$T^{-1} \mathbf{F}' (\mathbf{P}_{\hat{\mathbf{F}}} - \mathbf{P}_{\mathbf{F}}) \mathbf{F} = T^{-1} \mathbf{F}' \mathbf{P}_{\hat{\mathbf{F}}} \mathbf{F} - T^{-1} \mathbf{F}' \mathbf{F} \quad (29)$$

is of order $o_p(N^{-1}T^{-1/2})$. However, since $\hat{\mathbf{F}}$ converges to $\mathbf{F}\bar{\mathbf{C}}$ pointwise at rate \sqrt{N} , it is reasonable to conjecture that the order of the difference (29) is not smaller than $\mathcal{O}_p(N^{-1})$. Hence, the CD test statistic diverges at rate $\mathcal{O}_p(\sqrt{T})$. As in the model with additive time effects, this is a materialization of the IPP due to the presence of T unknown parameters that converge at the relatively slow rate of \sqrt{N} each.

A.2 Power of CD statistic with CCE residuals

A mis-specified latent common factor model can be characterized by equation (8) such that Assumptions 2–5 as well as the following Assumption 8 hold.

Assumption 8. $\operatorname{rk}([\boldsymbol{\mu}_{\lambda}, \boldsymbol{\mu}_{\Lambda}]) = m + 1 < r$.

Assumption 8 enforces a failure of the rank condition set up by Pesaran (2006) to ensure that the space spanned by factor estimates can be consistent for the space spanned by the true factors. Under failure of the rank condition, a fraction of the common variation affecting the dependent variable $y_{i,t}$ remains asymptotically unaccounted for. As proved in Lemma 5, we can assume without loss of generality that accounted and unaccounted sources of cross-section dependence are due to two uncorrelated sets of unobserved factors. Formally, we decompose

$$\mathbf{F} \boldsymbol{\lambda}_i = \mathbf{F}^{(1)} \boldsymbol{\lambda}_i^{(1)} + \mathbf{F}^{(2)} \boldsymbol{\lambda}_i^{(2)}, \quad (30)$$

where a rotation of the $m + 1$ factors $\mathbf{F}^{(1)}$ is consistently estimated by cross-section averages $\hat{\mathbf{F}} = [\bar{\mathbf{y}}, \bar{\mathbf{X}}]$ whereas the remaining $r - m - 1$ factors $\mathbf{F}^{(2)}$ are asymptotically orthogonal to $\hat{\mathbf{F}}$. An analogous decomposition can be applied to the matrix product $\mathbf{F} \boldsymbol{\Lambda}_i$, allowing us to split the loadings matrix into two blocks $\boldsymbol{\Lambda}_i^{(1)}$ and $\boldsymbol{\Lambda}_i^{(2)}$. Most importantly, this decomposition implies that $\mathbb{E}[\boldsymbol{\lambda}_i^{(2)}] = \mathbf{0}$ and $\mathbb{E}[\boldsymbol{\Lambda}_i^{(2)}] = \mathbf{O}$.

We can accordingly express the defactored data as

$$\begin{aligned} \hat{\boldsymbol{v}}_i &= \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{y}_i \\ &= \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{F}^{(1)} \boldsymbol{\lambda}_i^{(1)} + \mathbf{M}_{\hat{\mathbf{F}}} \mathbf{F}^{(2)} \boldsymbol{\lambda}_i^{(2)} + \mathbf{M}_{\hat{\mathbf{F}}} \boldsymbol{\varepsilon}_i. \end{aligned} \quad (31)$$

¹⁸Here we implicitly use the standard assumption of independently and identically distributed loadings with non-zero mean.

where Assumption 8 and Lemma 5 ensure that $M_{\hat{F}}\mathbf{F}^{(2)}\boldsymbol{\lambda}_i^{(2)}$ does not vanish as the sample size increases. Again, we assume a pure static factor model for $y_{i,t}$ by imposing the restriction $\boldsymbol{\beta} = \mathbf{0}$ and denote the variance of the composite error term $v_{i,t} = \mathbf{f}_t^{(2)'}\boldsymbol{\lambda}_i^{(2)} + \varepsilon_{i,t}$ by $\zeta_{i,v}^2 = \mathbb{E}\left[v_{i,t}^2 \mid \boldsymbol{\lambda}_i^{(2)}, \sigma_i^2\right] = \boldsymbol{\lambda}_i^{(2)'}\boldsymbol{\Sigma}_{F,22}\boldsymbol{\lambda}_i^{(2)} + \sigma_i^2$, where the $r - m - 1$ -dimensional square matrix $\boldsymbol{\Sigma}_{F,22}$ is the lower-right block of $\boldsymbol{\Sigma}_F$. The order in probability of the CD test statistic can then be characterized as follows.

Proposition 3. *Suppose that the true model is given by (6) with $\boldsymbol{\beta} = \mathbf{0}$ and that its components satisfy Assumptions 2–5 and 8. Let the CD test statistic $CD_{\mathbb{H}_1}$, defined in (19), be constructed from defactored data (31). Then,*

$$CD_{\mathbb{H}_1} = \sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \boldsymbol{\omega}'_i \boldsymbol{\omega}_j + \sqrt{T} (\Phi_{1,\mathbb{H}_1} - 2\Phi_{2,\mathbb{H}_1}) + \mathcal{O}_P(N^{-1/2}\sqrt{T}) + \mathcal{O}_P(T^{-1/2}) \quad (32)$$

where

$$\begin{aligned} \boldsymbol{\omega}_i &= \left[\zeta_{i,v}^{-1} \boldsymbol{\nu}_i - \mathbf{D}_i (\overline{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,v}^{-1} \right) \right], \\ \Phi_{1,\mathbb{H}_1} &= \left(N^{-1} \sum_{\ell=1}^N \zeta_{\ell,v}^{-1} \boldsymbol{\lambda}_\ell^{(1)'} \right) (\overline{\mathbf{C}}^{(1)'})^{-1} \left(N^{-1} \sum_{i=1}^N \mathbf{D}'_i \mathbf{D}_i \right) (\overline{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,v}^{-1} \right), \\ \Phi_{2,\mathbb{H}_1} &= N^{-1} \sum_{i=1}^N \zeta_{i,v}^{-1} \boldsymbol{\nu}'_i \mathbf{D}_i (\overline{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,v}^{-1} \right), \end{aligned}$$

where $\overline{\mathbf{C}}^{(1)} = [\overline{\boldsymbol{\lambda}}^{(1)}, \overline{\boldsymbol{\Lambda}}^{(1)}]$ and $\mathbf{D}_i = \mathbf{F} [\boldsymbol{\lambda}_i^{(2)}, \boldsymbol{\Lambda}_i^{(2)}] + [\boldsymbol{\varepsilon}_i, \mathbf{e}_i]$. Furthermore, it holds that

$$\text{plim}_{N,T \rightarrow \infty} N^{-1} T^{-1/2} \sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \boldsymbol{\omega}'_i \boldsymbol{\omega}_j = \text{cov} \left[\zeta_{1,v}^{-1}, \boldsymbol{\lambda}_1^{(2)'} \right] \boldsymbol{\Sigma}_{F,22} \text{cov} \left[\boldsymbol{\lambda}_1^{(2)}, \zeta_{1,v}^{-1} \right].$$

In the special case $\text{cov} \left[\zeta_{v,1}^{-1}, \boldsymbol{\lambda}_1^{(2)'} \right] = \mathbf{0}$, the term $\sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \boldsymbol{\omega}'_i \boldsymbol{\omega}_j$ is asymptotically centered around zero and has a variance that diverges at rate T .

Proof. In the Supplementary Appendix. □

The remarks made concerning Proposition 2 apply in an identical fashion to Proposition 3. That is, the CD test statistic allows for the same type of decomposition under the null hypothesis as it does under the alternative, differing only in that the expressions in Proposition 3 contain the composite error term $v_{i,t}$ rather than the true model errors $\varepsilon_{i,t}$.

A.3 Optimal weights

Propositions 2 and 3 stated that the original CD test statistic diverges at rate $N\sqrt{T}$ if certain conditions on the dependence between error variances and loadings associated with unaccounted common factors hold. A simple generalization of Proposition 3 to the case of general weights w_1, \dots, w_N suggest that the same rate of divergence can be achieved under the condition $\text{cov} \left[w_i, \lambda_i^{(2)} \right] \neq \mathbf{0}$.¹⁹ Consequently, a set of weights that generally leads to high power would be given by functions of the data that are estimates of $\lambda_i^{(2)}$ under \mathbb{H}_1 , such as $w_i = \sum_{t=1}^T \hat{v}_{i,t}$ or $w_{ij} = \sum_{t=1}^T \hat{v}_{i,t} \hat{v}_{j,t}$ for a more general statistic with index-pair specific weights. However, the latter weight suggestion directly results in a different test statistic, namely one based on *squared* cross-section covariances. This new test statistic, as well as one based on the first suggestion $w_i = \sum_{t=1}^T \hat{v}_{i,t}$, is closer to an existing, separate literature on LM tests for cross-section dependence than it is to the CD test and its extant modifications. The fact that a test statistic based on summing squared cross-section covariances is not centered around zero under its null hypothesis, and that it hence needs additional recentering, further emphasizes its relation to LM tests.

As an alternative to estimates of $\lambda_i^{(2)}$, estimates of other model components may be used to set up a weighted CD test statistic. In this regard, the most sensible choice is an estimate of loadings associated with unaccounted factors in the covariates of our regression model of interest, i.e. the parameter matrix $\Lambda_i^{(2)}$ in

$$\mathbf{X}_i = \mathbf{F}^{(1)} \Lambda_i^{(1)} + \mathbf{F}^{(2)} \Lambda_i^{(2)} + \mathbf{E}_i.$$

This choice directs the power of CD_W towards alternatives under which loadings associated with unaccounted factors are correlated in the sense of Westerlund and Urbain (2013) and Kapetanios et al. (2019). Following the theoretical results in these two studies, a CD test statistic that weights cross-section covariances with estimates of $\Lambda_i^{(2)}$ is effectively a test for inconsistency of either the two-way fixed effects or CCE estimator of β . It is generally possible to direct the CD test statistic towards these more specific hypothesis. However, if one is to test whether an estimate of the slope coefficients β on \mathbf{X}_i are inconsistent, it is simpler and more sensible to do this using a test statistic that addresses the moment condition $\text{cov} \left[\lambda_i^{(2)}, \Lambda_i^{(2)} \right] = \mathbf{0}$ directly.

¹⁹Analogous results hold for a generalization of Proposition 2

B Proofs: Additive model

Proof of Theorem 1.

At first we show how the test statistic in this case can be expressed in terms of four U statistics, and two additional terms that contribute to the bias. First, let

$$\begin{aligned} CD_\varepsilon &= \sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \frac{1}{T} \sum_{t=1}^T \varepsilon_{i,t} \varepsilon_{j,t}, \\ CD_{\varepsilon/\sigma} &= \sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \frac{1}{T} \sum_{t=1}^T \frac{\varepsilon_{i,t} \varepsilon_{j,t}}{\sigma_i \sigma_j}, \\ CD_{\varepsilon+} &= \sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \frac{1}{T} \sum_{t=1}^T \varepsilon_{i,t} \varepsilon_{j,t} (\sigma_i^{-1} + \sigma_j^{-1}), \\ CD_{\varepsilon++} &= \sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \frac{1}{T} \sum_{t=1}^T \varepsilon_{i,t} \varepsilon_{j,t} (\sigma_i^{-2} + \sigma_j^{-2}). \end{aligned}$$

Let us further denote by $k_{N,T} = \sqrt{\frac{1}{2TN(N-1)}}$. Observe that the original test statistic can be expressed as follows:

$$\begin{aligned} CD &= \sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T \frac{(\varepsilon_{i,t} - \bar{\varepsilon}_t)(\varepsilon_{j,t} - \bar{\varepsilon}_t)}{\sigma_i \sigma_j} \\ &= k_{N,T} \sum_{t=1}^T \left(\sum_{i=1}^N \frac{(\varepsilon_{i,t} - \bar{\varepsilon}_t)}{\sigma_i} \right)^2 - k_{N,T} \sum_{i=1}^N \sum_{t=1}^T \left(\frac{\varepsilon_{i,t} - \bar{\varepsilon}_t}{\sigma_i} \right)^2 \\ &= I - II. \end{aligned}$$

Let us now consider each term separately. First,

$$\begin{aligned} I &= k_{N,T} \sum_{t=1}^T \left(\sum_{i=1}^N \frac{\varepsilon_{i,t}}{\sigma_i} \right)^2 - 2k_{N,T} \overline{\sigma^{-1}} \sum_{t=1}^T \left[\left(\sum_{i=1}^N \frac{\varepsilon_{i,t}}{\sigma_i} \right) \left(\sum_{i=1}^N \varepsilon_{i,t} \right) \right] + k_{N,T} (\overline{\sigma^{-1}})^2 \sum_{t=1}^T \left(\sum_{i=1}^N \varepsilon_{i,t} \right)^2 \\ &= CD_{\varepsilon/\sigma} - \overline{\sigma^{-1}} CD_{\varepsilon+} + (\overline{\sigma^{-1}})^2 CD_\varepsilon \\ &\quad + k_{N,T} \sum_{i=1}^N \sum_{t=1}^T \left(\frac{\varepsilon_{i,t}}{\sigma_i} \right)^2 - 2k_{N,T} \overline{\sigma^{-1}} \sum_{i=1}^N \sum_{t=1}^T \sigma_i \left(\frac{\varepsilon_{i,t}}{\sigma_i} \right)^2 + k_{N,T} (\overline{\sigma^{-1}})^2 \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2. \end{aligned}$$

Similarly, for the second term we have

$$\begin{aligned} II &= k_{N,T} \sum_{t=1}^T \sum_{i=1}^N \left(\frac{\varepsilon_{i,t}}{\sigma_i} \right)^2 - 2k_{N,T} N^{-1} \sum_{t=1}^T \left[\left(\sum_{i=1}^N \frac{\varepsilon_{i,t}^2}{\sigma_i^2} \right) \left(\sum_{i=1}^N \varepsilon_{i,t} \right) \right] + k_{N,T} N^{-1} \overline{\sigma^{-2}} \sum_{t=1}^T \left(\sum_{i=1}^N \varepsilon_{i,t} \right)^2 \\ &= -N^{-1} CD_{\varepsilon++} + N^{-1} \overline{\sigma^{-2}} CD_\varepsilon \\ &\quad + k_{N,T} \sum_{t=1}^T \sum_{i=1}^N \left(\frac{\varepsilon_{i,t}}{\sigma_i} \right)^2 - 2k_{N,T} N^{-1} \sum_{i=1}^N \sum_{t=1}^T \left(\frac{\varepsilon_{i,t}}{\sigma_i} \right)^2 + k_{N,T} N^{-1} \overline{\sigma^{-2}} \sum_{i=1}^N \sum_{t=1}^T (\varepsilon_{i,t})^2. \end{aligned}$$

Combining these two expressions yields the expression in terms of four U-statistics and two bias terms referred to above:

$$\begin{aligned}
CD &= I - II \\
&= CD_{\varepsilon/\sigma} - 2\overline{\sigma^{-1}}CD_{\varepsilon+} - 2N^{-1}CD_{\varepsilon++} + \left((\overline{\sigma^{-1}})^2 - N^{-1}\overline{\sigma^{-2}} \right) CD_{\varepsilon} \\
&\quad + k_{N,T} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 \left((\overline{\sigma^{-1}})^2 - 2\overline{\sigma^{-1}}\sigma_i^{-1} \right) - N^{-1}k_{N,T} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 \left((\overline{\sigma^{-1}})^2 - 2\sigma_i^{-2} \right). \quad (33)
\end{aligned}$$

Concerning the second bias term in the last line above, we note that $(NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 = \mathcal{O}_P(1)$ and $(NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 \sigma_i^{-2} = \mathcal{O}_P(1)$ by application of Markov's inequality. It follows that

$$N^{-1}k_{N,T} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 \left((\overline{\sigma^{-1}})^2 - 2\sigma_i^{-2} \right) = \mathcal{O}_P(N^{-1}\sqrt{T}).$$

Concerning the four U-statistics, note that under Assumption 1 the error variances σ_i^2 are bounded, implying that all averages involving σ_i are in general of order $\mathcal{O}_P(1)$ by Markov's inequality. Lemma 1 can hence be used to show that all four U-statistics CD_{ε} , $CD_{\varepsilon/\sigma}$, $CD_{\varepsilon+}$ and $CD_{\varepsilon++}$ are of order $\mathcal{O}_P(1)$.²⁰ As a consequence, all terms in the second line of (33) involving N^{-1} are $\mathcal{O}_P(N^{-1})$. Additionally, a standard Lindeberg-Levi CLT implies that

$$\overline{\sigma^{-1}} = \mathbb{E}[\sigma_i^{-1}] + \mathcal{O}_P(N^{-1/2}).$$

This allows us to write

$$\begin{aligned}
&CD_{\varepsilon/\sigma} - 2\overline{\sigma^{-1}}CD_{\varepsilon+} + (\overline{\sigma^{-1}})^2 CD_{\varepsilon} \\
&= k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T \varepsilon_{i,t} \varepsilon_{j,t} \left(\sigma_i^{-1} - \mathbb{E}[\sigma_i^{-1}] \right) \left(\sigma_j^{-1} - \mathbb{E}[\sigma_j^{-1}] \right) + \mathcal{O}_P(N^{-1/2}).
\end{aligned}$$

Combining our results on the components of equation (33), we conclude that

$$CD = \sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T \varepsilon_{i,t} \varepsilon_{j,t} \left(\sigma_i^{-1} - \mathbb{E}[\sigma_i^{-1}] \right) \left(\sigma_j^{-1} - \mathbb{E}[\sigma_j^{-1}] \right) + \sqrt{T}\Xi + \mathcal{O}_P(N^{-1/2}),$$

where

$$\Xi = k_{N,T} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 \left((\overline{\sigma^{-1}})^2 - 2\overline{\sigma^{-1}}\sigma_i^{-1} \right).$$

It remains to prove weak convergence of the leading stochastic component in CD . For this purpose, set $a_{i,t} = \varepsilon_{i,t}(\sigma_i^{-1} - \mathbb{E}[\sigma_i^{-1}])$ and $q_i = \sigma_i(\sigma_i^{-1} - \mathbb{E}[\sigma_i^{-1}])$. We can then apply Lemma 1 and conclude that

$$CD - \sqrt{T}\Xi \xrightarrow{d} N(0, \Omega), \quad (34)$$

²⁰Strictly speaking, Lemma 1 does not apply to $CD_{\varepsilon+}$ and $CD_{\varepsilon++}$. However, it can be straightforwardly extended to accommodate the specific structure of the U-statistics in both expressions.

as $N, T \rightarrow \infty$ and $\sqrt{TN}^{-1} \rightarrow 0$, and where $\Omega = (\mathbb{E}[q_i^2])^2 = (\mathbb{E}[(1 - \sigma_i \mathbb{E}[\sigma_i^{-1}])^2])^2$. \square

Proof of Corollary 1.

Using the decomposition in equation (7) for a general value of σ^2 , we have

$$CD = \frac{1}{\sigma^2} \sqrt{\frac{1}{2TN(N-1)}} \sum_{t=1}^T \left(\sum_{i=1}^N \hat{\varepsilon}_{i,t} \right)^2 - \frac{1}{\sigma^2} \sqrt{\frac{1}{2TN(N-1)}} \sum_{t=1}^T \sum_{i=1}^N \hat{\varepsilon}_{i,t}^2. \quad (35)$$

The first component on the right-hand side above is 0 by construction. The second can be rewritten as

$$\begin{aligned} CD &= \frac{1}{N} CD_{\varepsilon/\sigma} - \frac{1}{\sigma^2} \sqrt{\frac{2}{TN(N-1)}} \frac{N-1}{2N} \sum_{t=1}^T \left(\sum_{i=1}^N \varepsilon_{i,t}^2 \right) \\ &= \frac{1}{N} CD_{\varepsilon/\sigma} - \sqrt{\frac{N-1}{2N^2}} \frac{1}{\sqrt{NT}} \sum_{t=1}^T \sum_{i=1}^N \left(\left(\frac{\varepsilon_{i,t}}{\sigma} \right)^2 - 1 \right) - \sqrt{\left(T - \frac{T}{N} \right) / 2}, \\ &= -\sqrt{\left(T - \frac{T}{N} \right) / 2} - \sqrt{\frac{1}{2N}} \frac{1}{\sqrt{NT}} \sum_{t=1}^T \sum_{i=1}^N \left(\left(\frac{\varepsilon_{i,t}}{\sigma} \right)^2 - 1 \right) + o_P(N^{-1/2}). \end{aligned}$$

Concerning the first expression in the last line, we can use the restriction $\sqrt{TN} \rightarrow 0$

Here, $CD_{\varepsilon/\sigma}$ is the usual CD test statistic based on raw error terms $\varepsilon_{i,t}/\sigma$. Lemma 1 allows us to conclude that $CD_{\varepsilon/\sigma} = \mathcal{O}_P(1)$. The final result follows after observing that an appropriate double index CLT for iid data (e.g. Theorem 3 in Phillips and Moon (1999)) implies

$$\frac{1}{\sqrt{NT}} \sum_{t=1}^T \sum_{i=1}^N \left(\left(\frac{\varepsilon_{i,t}}{\sigma} \right)^2 - 1 \right) = \frac{1}{\sqrt{NT}} \sum_{t=1}^T \sum_{i=1}^N (\eta_{i,t}^2 - 1) = \mathcal{O}_P(1), \quad (36)$$

given $\mathbb{E}[\eta_{i,t}^8] < \infty$. \square

Proof of Proposition 2.

The same steps as in the proof of Theorem 1 allow us to arrive at the decomposition

$$\begin{aligned} CD_{H_1} &= \sqrt{\frac{1}{2TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \left(\zeta_{v,i}^{-1} - \mathbb{E} \left[\zeta_{v,i}^{-1} \right] \right) \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \left(\zeta_{v,j}^{-1} - \mathbb{E} \left[\zeta_{v,j}^{-1} \right] \right) + \sqrt{T} \mathbb{E}_{H_1} \\ &\quad + \frac{\sqrt{T}}{N} \sqrt{\frac{N}{2(N-1)}} \left((NT)^{-1} \sum_{i,j}^N \frac{\boldsymbol{\nu}'_i \boldsymbol{\nu}_j}{\zeta_{v,i}^2} - \overline{\zeta_v^{-2}} (NT)^{-1} \sum_{i,j}^N \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \right), \end{aligned}$$

where

$$\mathbb{E}_{H_1} = \sqrt{\frac{N}{2(N-1)}} (NT)^{-1} \sum_{i=1}^N \boldsymbol{\nu}'_i \boldsymbol{\nu}_i \left[\left(\overline{\zeta_v^{-1}} \right)^2 - 2 \overline{\zeta_v^{-1}} \overline{\zeta_{v,i}} \right].$$

The terms in parentheses in the second line are stochastically bounded, entailing that the entire second line is $\mathcal{O}_P\left(N^{-1}\sqrt{T}\right)$. Next, consider the leading stochastic component of CD_{H_1} , namely $\sqrt{\frac{1}{2TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right) \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \left(\varsigma_{v,j}^{-1} - \mathbb{E}\left[\varsigma_{v,j}^{-1}\right]\right)$. Here, we have

$$\begin{aligned} & \mathbb{E}\left[\left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right) \nu_{i,t} \nu_{j,t} \left(\varsigma_{v,j}^{-1} - \mathbb{E}\left[\varsigma_{v,j}^{-1}\right]\right)\right] \\ &= \mathbb{E}\left[\left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right) \left(\boldsymbol{\lambda}_i - \mathbb{E}\left[\boldsymbol{\lambda}_i\right]\right)' \boldsymbol{\Sigma}_F \mathbb{E}\left[\left(\boldsymbol{\lambda}_j - \mathbb{E}\left[\boldsymbol{\lambda}_j\right]\right) \left(\varsigma_{v,j}^{-1} - \mathbb{E}\left[\varsigma_{v,j}^{-1}\right]\right)\right]\right] + 0 \\ &= \text{cov}\left[\varsigma_{v,i}^{-1}, \boldsymbol{\lambda}'_i\right] \boldsymbol{\Sigma}_F \text{cov}\left[\boldsymbol{\lambda}_j, \varsigma_{v,j}^{-1}\right] \\ &= \text{cov}\left[\varsigma_{v,1}^{-1}, \boldsymbol{\lambda}'_1\right] \boldsymbol{\Sigma}_F \text{cov}\left[\boldsymbol{\lambda}_1, \varsigma_{v,1}^{-1}\right], \end{aligned}$$

where we use *i.i.d.*-ness of $\boldsymbol{\lambda}_i$ and σ_i^2 to conclude that the covariance $\text{cov}\left[\varsigma_{v,i}^{-1}, \boldsymbol{\lambda}'_i\right]$ is the same across cross-sections.

Lastly, consider the variance of $\sqrt{\frac{1}{2TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right) \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \left(\varsigma_{v,j}^{-1} - \mathbb{E}\left[\varsigma_{v,j}^{-1}\right]\right)$ in the case where $\text{cov}\left[\varsigma_{v,i}^{-1}, \boldsymbol{\lambda}'_i\right] = 0$. We have

$$\begin{aligned} & \text{var}\left[\sum_{i=2}^N \sum_{j=1}^{i-1} \left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right) \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \left(\varsigma_{v,j}^{-1} - \mathbb{E}\left[\varsigma_{v,j}^{-1}\right]\right)\right] \\ &= \mathbb{E}\left\{\left[\sum_{i=2}^N \sum_{j=1}^{i-1} \left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right) \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \left(\varsigma_{v,j}^{-1} - \mathbb{E}\left[\varsigma_{v,j}^{-1}\right]\right)\right]^2\right\} - 0 \\ &= \sum_{i=2}^N \sum_{j=1}^{i-1} \mathbb{E}\left\{\left[\left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right) \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \left(\varsigma_{v,j}^{-1} - \mathbb{E}\left[\varsigma_{v,j}^{-1}\right]\right)\right]^2\right\}. \end{aligned}$$

Here, $(\boldsymbol{\nu}'_i \boldsymbol{\nu}_j)^2 = [(\boldsymbol{\lambda}'_i \mathbf{F}' + \varepsilon'_i) (\mathbf{F} \boldsymbol{\lambda}_j + \varepsilon_j)]^2$, where we only consider the leading term $\text{tr}\left(\boldsymbol{\lambda}_i \boldsymbol{\lambda}'_i \mathbf{F}' \mathbf{F} \boldsymbol{\lambda}_j \boldsymbol{\lambda}'_j \mathbf{F}' \mathbf{F}\right)$ further. Here, it is possible to show that

$$\mathbb{E}\left[\text{tr}\left(\boldsymbol{\lambda}_i \boldsymbol{\lambda}'_i \mathbf{F}' \mathbf{F} \boldsymbol{\lambda}_j \boldsymbol{\lambda}'_j \mathbf{F}' \mathbf{F}\right) \mid \boldsymbol{\lambda}_i, \boldsymbol{\lambda}_j, \sigma_i^2, \sigma_j^2\right] \leq T^2 M \text{tr}\left(\boldsymbol{\lambda}_i \boldsymbol{\lambda}'_i \boldsymbol{\lambda}_j \boldsymbol{\lambda}'_j\right),$$

since the fourth-order moments of \mathbf{f}_t are bounded. By application of the Law of Iterated expectations (LIE), it then follows that

$$\begin{aligned} & \sum_{i=2}^N \sum_{j=1}^{i-1} \mathbb{E}\left\{\left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right)^2 \text{tr}\left(\boldsymbol{\lambda}_i \boldsymbol{\lambda}'_i \mathbf{F}' \mathbf{F} \boldsymbol{\lambda}_j \boldsymbol{\lambda}'_j \mathbf{F}' \mathbf{F}\right) \left(\varsigma_{v,j}^{-1} - \mathbb{E}\left[\varsigma_{v,j}^{-1}\right]\right)^2\right\} \\ &= M T^2 \sum_{i=2}^N \sum_{j=1}^{i-1} \mathbb{E}\left\{\left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right)^2 \text{tr}\left(\boldsymbol{\lambda}_i \boldsymbol{\lambda}'_i \boldsymbol{\lambda}_j \boldsymbol{\lambda}'_j\right) \left(\varsigma_{v,j}^{-1} - \mathbb{E}\left[\varsigma_{v,j}^{-1}\right]\right)^2\right\} \\ &= \mathcal{O}\left(N^2 T^2\right), \end{aligned}$$

assuming that the higher-order moment $\mathbb{E}\left\{\left(\varsigma_{v,i}^{-1} - \mathbb{E}\left[\varsigma_{v,i}^{-1}\right]\right)^2 \text{tr}\left(\boldsymbol{\lambda}_i \boldsymbol{\lambda}'_i\right)\right\}$ exists. Involving the

square of the scaling $\sqrt{\frac{1}{2TN(N-1)}}$, we can conclude that the variance of

$$\sqrt{\frac{1}{2TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \left(\varsigma_{v,i}^{-1} - \mathbb{E} \left[\varsigma_v^{-1} \right] \right) \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \left(\varsigma_{v,j}^{-1} - \mathbb{E} \left[\varsigma_v^{-1} \right] \right) \quad (37)$$

diverges at rate T . □

Proof of Theorem 3 (Time fixed effects part).

The decomposition of CD provided in Theorem 1 is purely algebraic and holds even with general weights w_i . Hence we can write

$$\begin{aligned} & \sqrt{\frac{T}{2N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} T^{-1} \hat{\boldsymbol{\varepsilon}}'_i \hat{\boldsymbol{\varepsilon}}_j w_i w_j \\ &= k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \boldsymbol{\varepsilon}'_i \boldsymbol{\varepsilon}_j (w_i - \bar{w}) (w_j - \bar{w}) + \sqrt{T} (\Xi_{W,1} + \Xi_{W,2}), \end{aligned}$$

where

$$\begin{aligned} \Xi_{W,1} &= \sqrt{\frac{N}{2(N-1)}} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 \left((\bar{w})^2 - 2\bar{w}w_i \right) \\ \Xi_{W,2} &= \sqrt{\frac{1}{2N(N-1)}} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 \left((\bar{w})^2 - 2w_i^2 \right). \end{aligned}$$

Here, we consider the leading bias term $\Xi_{1,W}$ where $(NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 = \mathcal{O}_p(1)$, $\bar{w} = \mathcal{O}_p(N^{-1/2})$ and $N^{-1/2} T^{-1} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 w_i^2 = \mathcal{O}_p(1)$ straightforwardly hold by Markov's and Chebyshev's inequalities and uniform boundedness of the error variances. It follows that $\Xi_{1,W} = \mathcal{O}_p(N^{-1})$. For the second remainder term $\Xi_{W,2}$, we note that $\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 w_i^2 = \mathcal{O}_p(1)$ which together with the previous intermediary results implies $\Xi_{W,2} = \mathcal{O}_p(N^{-2})$.

We continue with the leading stochastic term in CD_W , given by

$$\begin{aligned} & \sqrt{\frac{T}{2N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \varepsilon_{i,t} \varepsilon_{j,t} (w_i - \bar{w}) (w_j - \bar{w}) \\ &= \sqrt{\frac{T}{2N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \varepsilon_{i,t} \varepsilon_{j,t} w_i w_j \\ & - (\bar{w} - N^{-1}) \left(\sqrt{\frac{T}{2N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \varepsilon_{i,t} \varepsilon_{j,t} (w_i + w_j) \right) \\ & + (\bar{w}^2 - N^{-1} \bar{w}^2) \left(\sqrt{\frac{T}{2N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \varepsilon_{i,t} \varepsilon_{j,t} \right). \end{aligned}$$

Given that $E[w_i] = 0$, the second and third components are of orders $\mathcal{O}_P(N^{-1/2})$ and $\mathcal{O}_P(N^{-1})$, respectively. This follows from the application of Lemma 1 on corresponding U-statistics. The remaining non-negligible term is of the form:

$$\sqrt{\frac{T}{2N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \varepsilon_{i,t} \varepsilon_{j,t} w_i w_j.$$

From Lemma 1 it follows that:

$$\sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \varepsilon_{i,t} \varepsilon_{j,t} w_i w_j \xrightarrow{d} N(0, \Omega) \quad (38)$$

as $N, T \rightarrow \infty$ jointly, where $\Omega = E[\sigma_i^2 w_i^2]$. The last step in this proof is to show consistency of the standard deviation estimator $\hat{\Omega} = (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \hat{\varepsilon}_{i,t}^2 w_i^2$ so that convergence in law of CD_W to a standard normal distribution follows by Slutsky's Theorem. Note here that an equivalent expression is given by term *II* in the proof of Theorem 1. Drawing from corresponding intermediary results in this theorem, adapted for general weights w_i , we conclude that

$$(NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \hat{\varepsilon}_{i,t}^2 w_i^2 = (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 w_i^2 + \mathcal{O}_P(N^{-1} \sqrt{T}).$$

While the second term on the right-hand side above is asymptotically negligible under the established restrictions on the relative expansion rate of N and T , we need to show convergence of the first term to Ω . This is done by using Corollary 1 in Phillips and Moon (1999), letting $Q_{i,T} = T^{-1} \sum_{t=1}^T \varepsilon_{i,t}^2 w_i^2$, $Q_i = \sigma_i^2 w_i^2$ as well as $C_i = 1$. Sufficient conditions of that lemma are satisfied given the existence of the eight moments of $\eta_{i,t}, \sigma_i$ and w_i . It follows that $CD_W \xrightarrow{d} N(0, 1)$ \square

C Proofs: CCE

The proofs in this section frequently employ the rotation and rescaling matrix

$$\mathbf{B} = \begin{bmatrix} 1, & \mathbf{0}' \\ \boldsymbol{\beta}, & \mathbf{I}_m \end{bmatrix}$$

which relates the $T \times (m+1)$ matrix $[\mathbf{y}_i, \mathbf{X}_i]$ (or alternatively the average over all i) to common and idiosyncratic variation affecting each of the observed variables directly. This relation is given by

$$\begin{aligned} [\mathbf{y}_i, \mathbf{X}_i] &= \left(\mathbf{F} [\boldsymbol{\lambda}_i, \mathbf{A}_i] + [\boldsymbol{\varepsilon}_i, \mathbf{E}_i] \right) \mathbf{B} \\ &= \mathbf{F} \mathbf{C}_i + \mathbf{U}_i \end{aligned}$$

In the following, it is assumed that $\boldsymbol{\beta}$ is bounded so that $\|\mathbf{B}\| < \infty$. Furthermore, \mathbf{B} is non-singular by construction, implying that \mathbf{B}^{-1} exists. Given these properties, it trivially follows from the properties of $\boldsymbol{\lambda}_i$ and \mathbf{A}_i that $\|\overline{\mathbf{C}}\| = \mathcal{O}_P(1)$ and $\|\overline{\mathbf{C}}^{-1}\| = \mathcal{O}_P(1)$. The same holds for the matrix $\overline{\mathbf{C}}^{(1)}$ which is considered in the proof of Proposition 3.

Proof of Proposition 1.

As previously, denote by $k_{N,T} = \sqrt{\frac{1}{2TN(N-1)}}$ the scaling that enters the CD statistic. Under the assumptions of this proposition, the CD test statistic is given by

$$\begin{aligned} CD &= \frac{1}{\sigma^2} 2k_{N,T} \sum_{i=1}^{N-1} \sum_{j=i+1}^N \hat{\boldsymbol{\varepsilon}}_i' \hat{\boldsymbol{\varepsilon}}_j = \frac{1}{\sigma^2} k_{N,T} \sum_{i,j} \hat{\boldsymbol{\varepsilon}}_i' \hat{\boldsymbol{\varepsilon}}_j - \frac{1}{\sigma^2} k_{N,T} \sum_{i=1}^N \hat{\boldsymbol{\varepsilon}}_i' \hat{\boldsymbol{\varepsilon}}_i \\ &= I - II. \end{aligned}$$

We begin by proving that $I = 0$. Note that

$$\sum_{i,j} \hat{\boldsymbol{\varepsilon}}_i' \hat{\boldsymbol{\varepsilon}}_j = \sum_{t=1}^T \left(\sum_{i=1}^N \hat{\boldsymbol{\varepsilon}}_{i,t} \right)^2.$$

Furthermore, for each time period t the sum of residuals can be expressed as

$$\sum_{i=1}^N \hat{\boldsymbol{\varepsilon}}_{i,t} = N \left(\bar{\mathbf{y}}_t - \bar{\boldsymbol{\lambda}}' \hat{\mathbf{f}}_t \right), \quad (39)$$

where the average estimated loading $\bar{\boldsymbol{\lambda}}$ is implicitly defined by projection off the space spanned by $\hat{\mathbf{f}}_t$. That is, it can be written

$$\bar{\boldsymbol{\lambda}} = \left(\sum_{t=1}^T \hat{\mathbf{f}}_t \hat{\mathbf{f}}_t' \right)^{-1} \sum_{t=1}^T \hat{\mathbf{f}}_t \bar{\mathbf{y}}_t = \left(\sum_{t=1}^T \hat{\mathbf{f}}_t \hat{\mathbf{f}}_t' \right)^{-1} \sum_{t=1}^T \hat{\mathbf{f}}_t \hat{\mathbf{f}}_t' [1, \mathbf{0}'_m]' = [1, \mathbf{0}'_m]', \quad (40)$$

such that $\bar{\lambda}' \hat{f}_t = \bar{y}_t$. Combining these results we conclude:

$$\sum_{i=1}^N \hat{\varepsilon}_{i,t} = N \left(\bar{y}_t - \bar{\lambda}' \hat{f}_t \right) = 0, \quad (41)$$

therefore also $I = 0$. Next, we show that $II = \mathcal{O}_P(\sqrt{T})$. Using corresponding results from the proof of Theorem 2 with $w_i = \sigma^{-1} \forall i$, we can write

$$\begin{aligned} II &= k_{N,T} \frac{1}{\sigma^2} \sum_{i=1}^N \sum_{t=1}^T \varepsilon_{i,t}^2 + \mathcal{O}_P(N^{-1/2} \vee T^{-1/2}) + \mathcal{O}_P(N^{-1} \sqrt{T}) \\ &= NT \sqrt{\frac{1}{2TN(N-1)}} + \frac{1}{\sqrt{2(N-1)}} \frac{1}{\sqrt{NT}} \left(\sum_{i=1}^N \sum_{t=1}^T \eta_{i,t}^2 - 1 \right) + \mathcal{O}_P(N^{-1/2} \vee T^{-1/2}) + \mathcal{O}_P(N^{-1} \sqrt{T}), \\ &= \sqrt{\frac{TN}{2(N-1)}} + \mathcal{O}_P(N^{-1/2} \vee T^{-1/2}) + \mathcal{O}_P(N^{-1} \sqrt{T}), \\ &= \sqrt{\frac{T - T/N}{2}} + \mathcal{O}_P(N^{-1/2} \vee T^{-1/2}) + \mathcal{O}_P(N^{-1} \sqrt{T}). \end{aligned}$$

Here the second line follows from Theorem 3 Phillips and Moon (1999) applied to the sequence $\eta_{i,t}^2 - 1$. Combining our results on expressions I and II we conclude

$$CD = -\sqrt{\left(T - \frac{T}{N}\right)} / 2 + \mathcal{O}_P(N^{-1/2}) + \mathcal{O}_P(T^{-1/2}) + \mathcal{O}_P(N^{-1} \sqrt{T}). \quad (42)$$

□

Proof of Theorem 2.

To simplify notation and to emphasize the generality of our result, we prove this theorem for a generic set of random weights $\{w_1, w_2, \dots, w_N\}$ which has the same properties as spelled out in Assumption 2.2. Taking some conflicting notation with respect to Theorem 3 into account, let the CD test statistic based on these generic weights be denoted

$$\begin{aligned} CD_W &= 2k_{N,T} \sum_{i=1}^{N-1} \sum_{j=i+1}^N \hat{\varepsilon}'_i \hat{\varepsilon}_j w_i w_j = k_{N,T} \sum_{i,j} \hat{\varepsilon}'_i \hat{\varepsilon}_j w_i w_j - k_{N,T} \sum_{i=1}^N \hat{\varepsilon}'_i \hat{\varepsilon}_i w_i^2 \\ &= I - II, \end{aligned}$$

where

$$\begin{aligned} \hat{\varepsilon}_i &= M_{\hat{F}} (F \lambda_i + \varepsilon_i) \\ &= M_{\hat{F}} \varepsilon_i - M_{\hat{F}} (\hat{F} - F \bar{C}) (\bar{C}^{-1}) \lambda_i \\ &= M_{\hat{F}} \varepsilon_i - M_{\hat{F}} \bar{U} (\bar{C}^{-1}) \lambda_i. \end{aligned}$$

Now let

$$\begin{aligned}
I &= k_{N,T} \sum_{i,j}^N \left(\varepsilon_i - \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_i \right)' \left(\varepsilon_j - \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_j \right) w_i w_j \\
&\quad - k_{N,T} \sum_{i,j}^N \left(\varepsilon_i - \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_i \right)' P_{\hat{\mathbf{F}}} \left(\varepsilon_j - \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_j \right) w_i w_j \\
&= I_1 - I_2.
\end{aligned}$$

Here, analogous to how we proceeded in the time fixed effects model, we write

$$\sum_{i=1}^N \left(\varepsilon_i - \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_i \right) w_i = \sum_{i=1}^N \varepsilon_i w_i - \sum_{i=1}^N U_i \left(\bar{\mathcal{C}}^{-1} \right) \left(N^{-1} \sum_{\ell=1}^N \lambda_\ell w_\ell \right).$$

Consequently, letting $\bar{\lambda}_w = N^{-1} \sum_{\ell=1}^N \lambda_\ell w_\ell$ and $\zeta_i = \varepsilon_i w_i - U_i \left(\bar{\mathcal{C}}^{-1} \right) \bar{\lambda}_w$

$$\begin{aligned}
I_1 &= k_{N,T} \sum_{i,j}^N \zeta_i' \zeta_j \\
&= 2k_{N,T} \sum_{i=1}^{N-1} \sum_{j=i+1}^N \zeta_i' \zeta_j + k_{N,T} \sum_{i=1}^N \zeta_i' \zeta_i,
\end{aligned} \tag{43}$$

where

$$\begin{aligned}
k_{N,T} \sum_{i=1}^N \zeta_i' \zeta_i &= k_{N,T} \sum_{i=1}^N \varepsilon_i' \varepsilon_i w_i^2 + k_{N,T} \bar{\lambda}_w' \left(\bar{\mathcal{C}}^{-1} \right)' \left(\sum_{i=1}^N U_i' U_i \right) \left(\bar{\mathcal{C}}^{-1} \right) \bar{\lambda}_w \\
&\quad - 2k_{N,T} \bar{\lambda}_w' \left(\bar{\mathcal{C}}^{-1} \right)' \left(\sum_{i=1}^N U_i' \varepsilon_i w_i \right)
\end{aligned}$$

is a linear function of asymptotically non-negligible terms. Lemma 3 implies that $I_2 = o_p(1)$ if both $\sqrt{T}/N \rightarrow 0$ and $T \rightarrow \infty$. Hence, proceed to

$$\begin{aligned}
II &= k_{N,T} \sum_{i=1}^N \left(\varepsilon_i - \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_i \right)' \left(\varepsilon_i - \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_i \right) w_i^2 \\
&\quad - k_{N,T} \sum_{i=1}^N \left(\varepsilon_i - \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_i \right)' P_{\hat{\mathbf{F}}} \left(\varepsilon_i - \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_i \right) w_i^2 \\
&= II_1 - II_2.
\end{aligned}$$

As shown in Lemma 4, II_2 is $o_p(1)$, if $N^{-1} \sqrt{T} \rightarrow 0$ and $T \rightarrow \infty$. For this reason, consider the first term above instead. We can write

$$II_1 = k_{N,T} \sum_{i=1}^N \varepsilon_i' \varepsilon_i w_i^2 + k_{N,T} \sum_{i=1}^N \left\| \bar{U} \left(\bar{\mathcal{C}}^{-1} \right) \lambda_i \right\|^2 w_i^2 - 2k_{N,T} \sum_{i=1}^N \lambda_i' \left(\bar{\mathcal{C}}^{-1} \right)' \bar{U}' \varepsilon_i w_i^2,$$

where

$$\begin{aligned}
& k_{N,T} \sum_{i=1}^N \left\| \bar{\mathbf{U}} (\bar{\mathbf{C}}^{-1}) \boldsymbol{\lambda}_i \right\|^2 w_i^2 \\
& \leq k_{N,T} \frac{T}{N} \sum_{i=1}^N w_i^2 \boldsymbol{\lambda}_i' \boldsymbol{\lambda}_i \operatorname{tr} \left((N/T) \bar{\mathbf{U}}' \bar{\mathbf{U}} \right) \|\bar{\mathbf{C}}\|^2 M \\
& = \mathcal{O}_P(N^{-1} \sqrt{T})
\end{aligned}$$

by using result (S.22) and Markov's inequality on $\frac{1}{N} w_i^2 \sum_{i=1}^N \boldsymbol{\lambda}_i' \boldsymbol{\lambda}_i$. Next,

$$2k_{N,T} \left| \sum_{i=1}^N \boldsymbol{\lambda}_i' (\bar{\mathbf{C}}^{-1})' \bar{\mathbf{U}}' \boldsymbol{\varepsilon}_i w_i^2 \right| \leq 2k_{N,T} \sqrt{N} \left(N^{-1} \sum_{i=1}^N \left\| \boldsymbol{\lambda}_i' (\bar{\mathbf{C}}^{-1})' \right\|^2 \right)^{1/2} \left(\sum_{i=1}^N \left\| \bar{\mathbf{U}}' \boldsymbol{\varepsilon}_i w_i^2 \right\|^2 \right)^{1/2}.$$

As noted above, $N^{-1} \sum_{i=1}^N \left\| \boldsymbol{\lambda}_i' (\bar{\mathbf{C}}^{-1})' \right\|^2 = \mathcal{O}_P(1)$. In order to proceed in with the second term above, we use the identity $\bar{\mathbf{U}} = [\bar{\boldsymbol{\varepsilon}}; \bar{\mathbf{E}}] \mathbf{B}$ and write

$$\begin{aligned}
\mathbb{E} \left[\sum_{i=1}^N \left\| \bar{\mathbf{U}}' \boldsymbol{\varepsilon}_i w_i^2 \right\|^2 \right] & \leq \|\mathbf{B}\|^2 N^{-2} \sum_{i,i',i''}^N \sum_{t,t'}^T \mathbb{E} [\boldsymbol{\varepsilon}_{i,t} \boldsymbol{\varepsilon}_{i',t} \boldsymbol{\varepsilon}_{i'',t'} \boldsymbol{\varepsilon}_{i,t'}] \mathbb{E} [w_i^4] \\
& \quad + \|\mathbf{B}\|^2 N^{-2} \sum_{i,i',i''}^N \sum_{t,t'}^T \mathbb{E} [\boldsymbol{\varepsilon}_{i,t} \boldsymbol{\varepsilon}_{i,t'}] \mathbb{E} [\boldsymbol{e}'_{i',t} \boldsymbol{e}_{i'',t'}] \mathbb{E} [w_i^4] \\
& = \mathcal{O}(T) + \mathcal{O}(N^{-1} T^2),
\end{aligned}$$

which follows from combining indexes such that only nonzero expectations remain. Together with uniform boundedness of w_i^4 and boundedness of $\|\mathbf{B}\|^2$, we have

$$\begin{aligned}
2k_{N,T} \left| \sum_{i=1}^N \boldsymbol{\lambda}_i' (\bar{\mathbf{C}}^{-1})' \bar{\mathbf{U}}' \boldsymbol{\varepsilon}_i w_i^2 \right| & = \sqrt{\frac{2}{TN(N-1)}} \sqrt{N} \mathcal{O}_P(1) \left(\mathcal{O}_P(\sqrt{T}) + \mathcal{O}_P(N^{-1/2} T) \right) \\
& = \mathcal{O}_P(N^{-1/2}) + \mathcal{O}_P(N^{-1} \sqrt{T}).
\end{aligned} \tag{44}$$

Hence, we can conclude that

$$II_1 = k_{N,T} \sum_{i=1}^N \boldsymbol{\varepsilon}_i' \boldsymbol{\varepsilon}_i w_i^2 + \mathcal{O}_P(N^{-1/2}) + \mathcal{O}_P(T^{-1/2}) + \mathcal{O}_P(N^{-1} \sqrt{T}) \tag{45}$$

Combining the results on I_1 , I_2 , II_1 and II_2 , we have

$$\begin{aligned}
CD_W & = I_1 - I_2 - II_1 + II_2 \\
& = 2k_{N,T} \sum_{i=1}^{N-1} \sum_{j=i+1}^N \zeta_i' \zeta_j + k_{N,T} \bar{\boldsymbol{\lambda}}_w' (\bar{\mathbf{C}}^{-1})' \left(\sum_{i=1}^N \mathbf{U}_i' \mathbf{U}_i \right) (\bar{\mathbf{C}}^{-1}) \bar{\boldsymbol{\lambda}}_w \\
& \quad - 2k_{N,T} \bar{\boldsymbol{\lambda}}_w' (\bar{\mathbf{C}}^{-1})' \left(\sum_{i=1}^N \mathbf{U}_i' \boldsymbol{\varepsilon}_i w_i \right) + \mathcal{O}_P(T^{-1/2}) + \mathcal{O}_P(N^{-1/2}) + \mathcal{O}_P(N^{-1} \sqrt{T}),
\end{aligned}$$

from which the main result to be proven here follows by setting $w_i = \sigma_i^{-1}$, by referring to the resulting CD test statistic as CD and by defining the influence function as $\psi_{i,t} = (\bar{\mathbf{C}}')^{-1} \mathbf{u}_{i,t}$. \square

Proof of Theorem 3 (Multifactor error part).

Proceeding from the decomposition given in Theorem 2, we consider the two bias terms $\Phi_{1,W}$ and $\Phi_{2,W}$. For both terms, it is instructive to note that

$$\mathbb{E} \left[\left\| N^{-1} \sum_{i=1}^N w_i \boldsymbol{\lambda}_i \right\|^2 \right] = N^{-2} \sum_{i,j} \mathbb{E} [w_i w_j] \mathbb{E} [\boldsymbol{\lambda}_i' \boldsymbol{\lambda}_j] = N^{-2} \sum_{i=1}^N \text{var} [w_i] \mathbb{E} [\boldsymbol{\lambda}_i' \boldsymbol{\lambda}_i] = \mathcal{O} (N^{-1}),$$

so that

$$\left\| N^{-1} \sum_{i=1}^N w_i \boldsymbol{\lambda}_i \right\| = \mathcal{O}_P (N^{-1/2}). \quad (46)$$

Concerning $\Phi_{1,W}$, we also need to take into account

$$\begin{aligned} \left\| (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \boldsymbol{\psi}_{i,t} \boldsymbol{\psi}_{i,t}' \right\| &\leq (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \mathbf{u}_{i,t}' \mathbf{u}_{i,t} \left\| \bar{\mathbf{C}}^{-1} \right\|^2 \\ &= \mathcal{O}_P (1), \end{aligned}$$

which follows from Markov's inequality. Using this result, we have

$$\begin{aligned} \Phi_{1,W} &\leq \sqrt{\frac{N}{2(N-1)}} \left\| (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \boldsymbol{\psi}_{i,t} \boldsymbol{\psi}_{i,t}' \right\| \left\| N^{-1} \sum_{i=1}^N w_i \boldsymbol{\lambda}_i \right\|^2 \\ &= \mathcal{O}_P (N^{-1}). \end{aligned}$$

Next, consider $\tilde{\Phi}_{2,W}$ where

$$\left\| (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \boldsymbol{\psi}_{i,t} w_i \boldsymbol{\varepsilon}_{i,t} \right\| = \mathcal{O}_P (N^{-1/2})$$

by isolating $\bar{\mathbf{C}}^{-1}$ from the definition of $\boldsymbol{\psi}_{i,t}$ and by using result (S.24) on the remaining part. Consequently,

$$\begin{aligned} |\Phi_{2,W}| &\leq \sqrt{\frac{N}{2(N-1)}} \left\| N^{-1} \sum_{i=1}^N w_i \boldsymbol{\lambda}_i \right\| \left\| (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \boldsymbol{\psi}_{i,t} w_i \boldsymbol{\varepsilon}_{i,t} \right\| \\ &= \mathcal{O}_P (N^{-1}). \end{aligned}$$

Thus, the two bias terms affecting CD_W in Theorem 2 are negligible for weights satisfying Assumption 7 as long as $N^{-1}\sqrt{T} \rightarrow 0$. We continue with the leading stochastic term of the weighted CD test statistic, given by

$$\sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \zeta_{iN,t} \zeta_{jN,t}$$

with $\zeta_{iN,t} = \left(w_i \varepsilon_{i,t} - \left(N^{-1} \sum_{\ell=1}^N w_\ell \boldsymbol{\lambda}'_\ell \right) \boldsymbol{\psi}_{i,t} \right)$ and $\boldsymbol{\psi}_{i,t} = \overline{\mathbf{C}}^{-1} \mathbf{u}_{i,t}$. Analogous to the time fixed effects part of this proof, we can apply Lemma 1 with $a_{i,t} = w_i \varepsilon_{i,t}$ and $q_i = w_i \sigma_i$ to obtain

$$\sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \varepsilon_{i,t} \varepsilon_{j,t} w_i w_j \xrightarrow{d} N(0, (\sigma_w^2 \mathbb{E}[\sigma_i^2])^2)$$

as $N, T \rightarrow \infty$ subject to $N^{-1}\sqrt{T} \rightarrow 0$. Now note that by equation (45) in the proof of Theorem 2,

$$(NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \hat{\varepsilon}_{i,t}^2 w_i^2 = (NT)^{-1} \sum_{i=1}^N \varepsilon'_i \varepsilon_i w_i^2 + \mathcal{O}_P((NT)^{-1/2}) + \mathcal{O}_P(N^{-1}) + \mathcal{O}_P(T^{-1})$$

where $(NT)^{-1} \sum_{i=1}^N \varepsilon'_i \varepsilon_i w_i^2 \xrightarrow{p} \sigma_w^2 \mathbb{E}[\sigma_i^2]$ as argued in the time fixed effects part of the proof of this theorem. By application of Slutsky's Theorem, it then follows that

$$\left((NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \hat{\varepsilon}_{i,t}^2 w_i^2 \right)^{-1} \sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \varepsilon_{i,t} \varepsilon_{j,t} \xrightarrow{d} N(0, 1).$$

Next, consider the sum

$$\sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T \mathbf{u}_{i,t} \mathbf{u}'_{j,t} = \mathbf{B}' \left(\sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T (\varepsilon_{i,t}, \mathbf{e}'_{j,t})' (\varepsilon_{i,t}, \mathbf{e}'_{j,t}) \right) \mathbf{B}. \quad (47)$$

Given that $\mathbf{e}_{i,t}$ has properties similar to $\varepsilon_{i,t}$, it can be shown analogously the reasoning leading to (38) that a CLT holds for every element of the $(m+1) \times (m+1)$ matrix in parentheses in (47) by Lemma 1. This allows us to conclude that the elements of (47) are stochastically bounded. Thus, recalling result (46),

$$\begin{aligned} \left(N^{-1} \sum_{\ell=1}^N w_\ell \boldsymbol{\lambda}'_\ell \right) \left(\sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T \boldsymbol{\psi}_{i,t} \boldsymbol{\psi}'_{j,t} \right) \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell w_\ell \right) &= \mathcal{O}_P(N^{-1}), \\ \left(\sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T \varepsilon_{i,t} \boldsymbol{\psi}'_{j,t} \right) \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell w_\ell \right) &= \mathcal{O}_P(N^{-1/2}), \\ \left(N^{-1} \sum_{\ell=1}^N w_\ell \boldsymbol{\lambda}'_\ell \right) \left(\sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T \boldsymbol{\psi}_{i,t} \varepsilon_{i,t} \right) &= \mathcal{O}_P(N^{-1/2}), \end{aligned}$$

so that

$$\left((NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \hat{\varepsilon}_{i,t}^2 w_i^2 \right)^{-1} \sqrt{\frac{2T}{N(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T T^{-1} \zeta_{iN,t} \zeta_{jN,t} \xrightarrow{d} N(0, 1),$$

which leads to the central result of this theorem. \square

References

- AHN, S. C., Y. H. LEE, AND P. SCHMIDT (2013): "Panel Data Models with Multiple Time-varying Individual Effects," *Journal of Econometrics*, 174, 1–14.
- ALTONJI, J. G., AND L. M. SEGAL (1996): "Small-sample bias in GMM estimation of covariance structures," *Journal of Business & Economic Statistics*, 14, 353–366.
- ARELLANO, M., R. BLUNDELL, AND S. BONHOMME (2017): "Earnings and Consumption Dynamics: A Nonlinear Panel Data Framework," *Econometrica*, 85, 693–734.
- BAI, J. (2003): "Inferential Theory for Factor Models of Large Dimensions," *Econometrica*, 71, 135–171.
- (2009): "Panel Data Models With Interactive Fixed Effects," *Econometrica*, 77, 1229–1279.
- BAI, J. AND S. NG (2004): "A PANIC Attack on Unit Roots and Cointegration," *Econometrica*, 72, 1127–1177.
- BAILEY, N., S. HOLLY, AND M. H. PESARAN (2016): "A Two-Stage Approach to Spatio-Temporal Analysis with Strong and Weak Cross-Sectional Dependence," *Journal of Applied Econometrics*, 31, 249–280.
- BALTAGI, B. H., Q. FENG, AND C. KAO (2012): "A Lagrange Multiplier Test for Cross-sectional Dependence in a Fixed Effects Panel Data Model," *Journal of Econometrics*, 170, 164 – 177.
- BONEVA, L. AND O. LINTON (2017): "A Discrete Choice Model for Large Heterogeneous Panels with Interactive Fixed Effects with an Application to the Determinants of Corporate Bond Issuance," *Journal of Applied Econometrics*, 32, 1226–1243.
- BONHOMME, S., T. LAMADON, AND E. MANRESA (2017): "Discretizing Unobserved Heterogeneity," Mimeo.
- BONHOMME, S. AND E. MANRESA (2015): "Grouped Patterns of Heterogeneity in Panel Data," *Econometrica*, 83, 1147–1184.
- BONHOMME, S. AND J.-M. ROBIN (2010): "Generalized Non-Parametric Deconvolution with an Application to Earnings Dynamics," *The Review of Economic Studies*, 77, 491–533.
- CHEN, J., J. GAO, AND D. LI (2012): "A New Diagnostic Test for Cross-section Uncorrelatedness in Nonparametric Panel Data Models," *Econometric Theory*, 28, 1144–1163.
- CHEN, M., I. FERNÁNDEZ-VAL, AND M. WEIDNER (2014): "Nonlinear Panel Models with Interactive Effects," Mimeo.

- CHUDIK, A., K. MOHADDES, M. H. PESARAN, AND M. RAISSI (2017): "Is There a Debt-Threshold Effect on Output Growth?" *The Review of Economics and Statistics*, 99, 135–150.
- CHUDIK, A. AND M. H. PESARAN (2015): "Common Correlated Effects Estimation of Heterogeneous Dynamic Panel Data Models with Weakly Exogenous Regressors," *Journal of Econometrics*, 188, 393–420.
- DEMETRESCU, M. AND U. HOMM (2016): "Directed Tests of No Cross-Sectional Correlation in Large-N Panel Data Models," *Journal of Applied Econometrics*, 31, 4–31.
- EBERHARDT, M., C. HELMERS, AND H. STRAUSS (2013): "Do Spillovers Matter When Estimating Private Returns to R&D?" *The Review of Economics and Statistics*, 95, 436–448.
- EBERHARDT, M. AND A. F. PRESBITERO (2015): "Public Debt and Growth: Heterogeneity and Non-linearity," *Journal of International Economics*, 97, 45 – 58.
- EVERAERT, G. AND L. POZZI (2014): "The Predictability of Aggregate Consumption Growth in OECD Countries: A Panel Data Analysis," *Journal of Applied Econometrics*, 29, 431–453.
- FAN, J., Y. LIAO AND J. YAO, (2015): "Power Enhancement in High-Dimensional Cross-Sectional Tests," *Econometrica*, 83, 1497–1541.
- FERNÁNDEZ-VAL, I. AND M. WEIDNER (2016): "Individual and Time Effects in Nonlinear Panel Models with Large N, T," *Journal of Econometrics*, 192, 291 – 312.
- FOERSTER, A. T., P.-D. G. SARTE, AND M. W. WATSON (2011): "Sectoral Versus Aggregate Shocks: A Structural Factor Analysis of Industrial Production," *Journal of Political Economy*, 119, 1–38.
- GAGLIARDINI, P., E. OSSOLA, AND O. SCAILLET (2016): "Time-Varying Risk Premium in Large Cross-Sectional Equity Data Sets," *Econometrica*, 84, 985–1046.
- GOBILLON, L. AND T. MAGNAC (2016): "Regional Policy Evaluation: Interactive Fixed Effects and Synthetic Controls," *Review of Economics and Statistics*, 98, 535–551.
- HAHN, J. AND G. KUERSTEINER (2002): "Asymptotically Unbiased Inference for a Dynamic Panel Model with Fixed Effects When Both N and T are Large," *Econometrica*, 70(4), 1639–1657.
- HAHN, J. AND H. R. MOON (2006): "Reducing Bias of MLE in A Dynamic Panel Model," *Econometric Theory*, 22, 499–512.
- HAHN, J. AND W. NEWEY (2004): "Jackknife and Analytical Bias Reduction for Nonlinear Panel Models," *Econometrica*, 72(4), 1295–1319.

- HALL, P. AND C. C. HEYDE (1980): *Martingale Limit Theory and Its Application*, Probability and Mathematical Statistics, Academic Press.
- HOLLY, S., M. H. PESARAN, AND T. YAMAGATA (2010): "A Spatio-temporal Model of House Prices in the USA," *Journal of Econometrics*, 158, 160 – 173, twenty Years of Cointegration.
- HSIAO, C. (2014): *Analysis of Panel Data*, Econometric Society Monographs, Cambridge University Press, 3 ed.
- HSIAO, C., M. H. PESARAN, AND A. PICK (2012a): "Diagnostic Tests of Cross-section Independence for Limited Dependent Variable Panel Data Models*," *Oxford Bulletin of Economics and Statistics*, 74, 253–277.
- HSIAO, C., H. STEVE CHING, AND S. KI WAN (2012b): "A Panel Data Approach For Program Evaluation: Measuring the Benefits of Political and Economic Integration of Hong Kong with Mainland China," *Journal of Applied Econometrics*, 27, 705–740.
- JUODIS, A. (2018): "Pseudo Panel Data Models with Cohort Interactive Effects," *Journal of Business & Economic Statistics*, 36, 47–61.
- JUODIS, A., H. KARABIYIK, AND J. WESTERLUND (2017): "On the Robustness of the Pooled CCE Estimator," Mimeo.
- KAPETANIOS, G., L. SERLENGA, AND Y. SHIN (2019): "Testing for Correlated Factor Loadings in Cross Sectionally Dependent Panels," SERIES Working papers N. 02/2019. Available at SSRN: <https://ssrn.com/abstract=3401745>
- KARABIYIK, H., S. REESE, AND J. WESTERLUND (2017): "On The Role of The Rank Condition in CCE Estimation of Factor-augmented Panel Regressions," *Journal of Econometrics*, 197, 60–64.
- LANCASTER, T. (2000): "The Incidental Parameter Problem since 1948," *Journal of Econometrics*, 95, 391–413.
- MAO, G. (2016): "Testing for Error Cross-sectional Independence using Pairwise Augmented Regressions," *The Econometrics Journal*, 19, 237–260.
- (2018): "Testing for sphericity in a two-way error components panel data model," *Econometric Reviews*, 37, 491–506.
- MASTROMARCO, C., L. SERLENGA, AND Y. SHIN (2016): "Modelling Technical Efficiency in Cross Sectionally Dependent Stochastic Frontier Panels," *Journal of Applied Econometrics*, 31, 281–297, jae.2439.

- MOSCONE, F. AND E. TOSETTI (2010): "Health Expenditure and Income in the United States," *Health Economics*, 19, 1385–1403.
- NEYMAN, J. AND E. L. SCOTT (1948): "Consistent Estimation from Partially Consistent Observations," *Econometrica*, 16, 1–32.
- PARKER, J. AND D. SUL (2016): "Identification of Unknown Common Factors: Leaders and Followers," *Journal of Business & Economic Statistics*, 34, 227–239.
- PESARAN, M. H. (2004): "General Diagnostic Tests for Cross Section Dependence in Panels," CESifo Working Paper No. 1229.
- (2006): "Estimation and Inference in Large Heterogeneous Panels with a Multifactor Error Structure," *Econometrica*, 74, 967–1012.
- (2015a): "Testing Weak Cross-Sectional Dependence in Large Panels," *Econometric Reviews*, 34, 1089–1117.
- (2015b): *Time Series and Panel Data Econometrics*, Oxford University Press.
- PESARAN, M. H., A. ULLAH, AND T. YAMAGATA (2008): "A Biasadjusted LM Test of Error Crosssection Independence," *The Econometrics Journal*, 11(1), 105–127.
- PHILLIPS, P. C. B. AND H. R. MOON (1999): "Linear Regression Limit Theory for Nonstationary Panel Data," *Econometrica*, 67, 1057–1111.
- REESE, S. AND J. WESTERLUND (2016): "Panicka: Panic on Cross-Section Averages," *Journal of Applied Econometrics*, 31, 961–981.
- ROBINSON, P. (2008): "Correlation Testing in Time Series, Spatial and Cross-sectional Data," *Journal of Econometrics*, 147, 5 – 16.
- SARAFIDIS, V., T. YAMAGATA, AND D. ROBERTSON (2009): "A Test of Cross Section Dependence for a Linear Dynamic Panel Model with Regressors," *Journal of Econometrics*, 148, 149–161.
- SU, L., Z. SHI, AND P. C. B. PHILLIPS (2016): "Identifying Latent Structures in Panel Data," *Econometrica*, 84, 2215–2264.
- WESTERLUND, J., H. KARABIYIK, AND P. NARAYAN (2017): "Testing for Predictability in Panels with General Predictors," *Journal of Applied Econometrics*, 32, 554–574.
- WESTERLUND, J. AND J.-P. URBAIN (2013): "On the Estimation and Inference in Factor-augmented Panel Regressions with Correlated Loadings," *Economics Letters*, 119, 247 – 250.

——— (2015): “Cross-sectional Averages Versus Principal Components,” *Journal of Econometrics*, 185, 372 – 377.

WHITE, H. (2001): *Asymptotic Theory for Econometricians*, Academic Press.

Online Supplement to

The Incidental Parameters Problem in Testing for Remaining Cross-section Correlation

by

Artūras Juodis and Simon Reese

October 2019

Table of Contents

| | |
|--|-----------|
| S.1 Additional theoretical contributions and discussion | 2 |
| S.1.1 An analytically bias-corrected CD test statistic | 2 |
| S.1.2 Group effects and principal components | 3 |
| S.1.3 Serial correlation | 4 |
| S.2 Proofs: CCE power | 6 |
| S.3 Auxiliary lemmas | 10 |
| S.4 Additional MC results | 25 |

S.1 Additional theoretical contributions and discussion

S.1.1 An analytically bias-corrected CD test statistic

Section 4.1 introduced parametric bias-correction as a feasible approach to re-establish asymptotically normal inference for the CD tests statistic under its null hypothesis. This section provides details on its implementation as a benchmark CD test statistic in the Monte Carlo experiments of Section 5. The analytically bias-corrected CD test statistic CD_{BC} is corrected with a plug-in estimate of the bias terms of Theorems 1 and 2. That is, whenever CD_{BC} is applied to 2WFE residuals, we construct it as

$$CD_{BC} = \hat{\Omega}_{FE}^{-1/2} \left(CD - \sqrt{T} \hat{\Xi} \right)$$

where

$$\hat{\Xi} = \sqrt{\frac{1}{2N(N-1)}} \left[\sum_{i=1}^N \left(1 - \sqrt{\hat{\sigma}_i^2} N^{-1} \sum_{\ell=1}^N (\hat{\sigma}_\ell^2)^{-1/2} \right)^2 - N \right],$$

$$\hat{\Omega}_{FE} = \frac{2(N-1)}{N} \left(\hat{\Xi} + 1 \right)^2$$

with $\hat{\sigma}_i^2 = T^{-1} \sum_{t=1}^T \hat{\varepsilon}_{i,t}^2$ for 2WFE residuals $\hat{\varepsilon}_{i,t}$. The expressions above arise from rearranging the terms of Theorem 1 so that their estimates can be obtained with minimal computational burden. In this regard, it is helpful to note that the asymptotic variance of Ω can be written as a function of the bias term Ξ . Likewise, for application of CD_{BC} to CCE residuals, we set

$$CD_{BC} = \hat{\Omega}_{CCE}^{-1/2} \left[CD - \sqrt{T} \left(\hat{\Phi}_1 - 2\hat{\Phi}_2 \right) \right],$$

where

$$\hat{\Phi}_1 = \sqrt{\frac{1}{2N(N-1)}} \left[N^{-1} \sum_{i=1}^N (\hat{\sigma}_i^2)^{-1/2} \right]^2 \bar{\lambda}' \hat{B}' \begin{bmatrix} \sum_{i=1}^N \hat{\sigma}_i^2; & \mathbf{0}'_m \\ \mathbf{0}_m; & \sum_{i=1}^N \hat{\Sigma}_i \end{bmatrix} \hat{B} \bar{\lambda},$$

$$\hat{\Phi}_2 = \sqrt{\frac{1}{2N(N-1)}} \left[N^{-1} \sum_{i=1}^N (\hat{\sigma}_i^2)^{-1/2} \right] \bar{\lambda}' \begin{bmatrix} \sum_{i=1}^N \hat{\sigma}_i^2 \\ \mathbf{0}_m \end{bmatrix},$$

$$\hat{\Omega}_{CCE} = \left[1 + \sqrt{\frac{2(N-1)}{N}} \left(\hat{\Phi}_1 - 2\hat{\Phi}_2 \right) \right]^2$$

with

$$\begin{aligned}\widehat{\mathbf{B}} &= \begin{bmatrix} 1 & \mathbf{0}'_m \\ \widehat{\boldsymbol{\beta}}^{\text{CCE}} & \mathbf{I}_m \end{bmatrix}, \\ \widehat{\boldsymbol{\beta}}^{\text{CCE}} &= \left((NT)^{-1} \sum_{i=1}^N \mathbf{X}'_i \mathbf{M}_{\widehat{\mathbf{F}}} \mathbf{X}_i \right)^{-1} (NT)^{-1} \sum_{i=1}^N \mathbf{X}'_i \mathbf{M}_{\widehat{\mathbf{F}}} \mathbf{y}_i, \\ \bar{\boldsymbol{\lambda}} &= N^{-1} \sum_{i=1}^N \widehat{\boldsymbol{\lambda}}_i, \\ \widehat{\boldsymbol{\lambda}}_i &= (\widehat{\mathbf{F}}' \widehat{\mathbf{F}})^{-1} \widehat{\mathbf{F}}' (\mathbf{y}_i - \mathbf{X}_i \widehat{\boldsymbol{\beta}}^{\text{CCE}}), \\ \widehat{\boldsymbol{\Sigma}}_i &= T^{-1} \mathbf{X}'_i \mathbf{M}_{\widehat{\mathbf{F}}} \mathbf{X}_i,\end{aligned}$$

as well as $\hat{\sigma}_i^2 = T^{-1} \hat{\boldsymbol{\varepsilon}}_i' \hat{\boldsymbol{\varepsilon}}_i$ and $\hat{\boldsymbol{\varepsilon}}_i = \mathbf{M}_{\widehat{\mathbf{F}}} \mathbf{y}_i - \mathbf{M}_{\widehat{\mathbf{F}}} \mathbf{X}_i \widehat{\boldsymbol{\beta}}^{\text{CCE}}$. The plug-in estimators $\widehat{\Phi}_1$ and $\widehat{\Phi}_2$ are calculated under the assumption of independence between factor loadings and error variances so that we can write $E[\boldsymbol{\lambda}_i \sigma_i^{-1}] = E[\boldsymbol{\lambda}_i] E[\sigma_i^{-1}]$. While this assumption allows making the estimates of both bias terms sufficiently precise in the setup of our Monte Carlo experiments, it may lead to size distortion in a DGP with dependence between factor loadings and error variances.

S.1.2 Group effects and principal components

Group effects

The idea put forward in Bonhomme and Manresa (2015) is that all individuals $i = 1, \dots, N$ are associated with an unobserved group membership variable $g_i \in \{1, \dots, G\}$. Regressors aside their model is given by:

$$y_{i,t} = \tau_{g_i,t} + \varepsilon_{i,t}, \quad (\text{S.1})$$

where $\tau_{g_i,t}$ are the group-specific unobservable effects. Following Theorem 2 in Bonhomme and Manresa (2015) we know that g_i can be estimated with high probability as $T \rightarrow \infty$, thus the estimation error from $\hat{g}_i - g_i$ can be ignored when evaluating the estimators of $\tau_{g_i,t}$. Note that if g_i is known then $\hat{\tau}_{g_i,t}$ is simply given by:

$$\hat{\tau}_{g_i,t} - \tau_{g_i,t} = \frac{1}{N} \sum_{i=1}^N \pi_g I(g_i = g) \varepsilon_{i,t}. \quad (\text{S.2})$$

where $\pi_g = N/N_g$ with $N_g = \sum_{i=1}^N I(g_i = g)$. The model in (S.1) admits an alternative representation as a factor model with G factors:

$$y_{i,t} = \boldsymbol{\lambda}'_i \mathbf{f}_t + \varepsilon_{i,t}, \quad (\text{S.3})$$

with $\boldsymbol{\lambda}_i = (I(g_i = 1), \dots, I(g_i = G))'$ and $\boldsymbol{f}_t = (\tau_{1,t}, \dots, \tau_{G,t})$. Thus the influence function $\boldsymbol{\psi}_{i,t}$ for $\hat{\boldsymbol{f}}_t$ is simply given by:

$$\boldsymbol{\psi}_{i,t} = \begin{bmatrix} \pi_1 I(g_i = 1) \varepsilon_{i,t} \\ \vdots \\ \pi_G I(g_i = G) \varepsilon_{i,t} \end{bmatrix}. \quad (\text{S.4})$$

Next consider the residuals in this model

$$\frac{1}{N} \sum_{i=1}^N (y_{i,t} - \boldsymbol{\lambda}'_i \hat{\boldsymbol{f}}_t) = \bar{\varepsilon}_t - \bar{\boldsymbol{\lambda}}' \bar{\boldsymbol{\psi}}_t = \bar{\varepsilon}_t - \frac{1}{N} \sum_{i=1}^N \sum_{g=1}^G I(g_i = 1) \varepsilon_{i,t} = 0, \quad (\text{S.5})$$

as $\sum_{g=1}^G I(g_i = 1) = 1$. This conclusion is analogous to the simple model with additive time effects only, i.e. $G = 1$.

Principal components

As in the main text we consider an exact factor model

$$y_{i,t} = \boldsymbol{\lambda}'_i \boldsymbol{f}_t + \varepsilon_{i,t}, \quad (\text{S.6})$$

with an unrestricted vector $\boldsymbol{\lambda}_i$. If $\boldsymbol{\lambda}_i$ are assumed to be known then \boldsymbol{f}_t can be estimated by least squares:

$$\hat{\boldsymbol{f}}_t - \boldsymbol{f}_t = \frac{1}{N} \sum_{i=1}^N \boldsymbol{\psi}_{i,t}, \quad \boldsymbol{\psi}_{i,t} = \left(N^{-1} \boldsymbol{\lambda}_i \boldsymbol{\lambda}'_i \right)^{-1} \boldsymbol{\lambda}_i \varepsilon_{i,t}. \quad (\text{S.7})$$

Estimation of $\boldsymbol{\lambda}_i$ implies that \boldsymbol{f}_t can only be introduced up to a rotation (see e.g. Bai (2003) Theorem 1) via principal components:

$$\boldsymbol{H}^{-1} \hat{\boldsymbol{f}}_t - \boldsymbol{f}_t = \frac{1}{N} \sum_{i=1}^N \boldsymbol{\psi}_{i,t} + \mathcal{O}_P(1), \quad \boldsymbol{\psi}_{i,t} = \boldsymbol{H}^{-1} \boldsymbol{V}^{-1} \boldsymbol{Q}^{-1} \boldsymbol{\lambda}_i \varepsilon_{i,t} \quad (\text{S.8})$$

where for precise definitions of $\boldsymbol{H}, \boldsymbol{V}, \boldsymbol{Q}$, as well as restrictions on N, T the interested reader is referred to Bai (2003, Theorem 1 and Proposition 1). Note that unlike all previous estimation techniques, the residuals obtained after PC estimation of factors are not zero on average for each t (in general).

S.1.3 Serial correlation

The original CD test of Pesaran (2004, 2015) assumes that error terms are serially uncorrelated, an assumption that can be difficult to justify in the case of economic data. While this problem is mostly ignored in practice, Baltagi et al. (2016) have very recently proposed a modification of the CD test statistic which ensures that the test statistic is asymptotically standard normal as

long as $\varepsilon_{i,t}$ is a stationary short memory process. In particular, under this type of assumption it can be shown that the relevant asymptotic variance for the result in Theorem 3 is given by:

$$\Omega = \sum_{s=-\infty}^{\infty} [\mathbb{E}[(w_i - \mathbb{E}[w_i])^2 \varepsilon_{i,t} \varepsilon_{i,t-s}]]^2. \quad (\text{S.9})$$

The above quantity is non-standard, as it not a function of the long-run variance of $(w_i - \mathbb{E}[w_i])\varepsilon_{i,t}$. In particular, one can use data which is over-differenced in the construction of the CD statistic.

In the context of our testing problem the natural plug-in estimator of Ω is given by

$$\hat{\Omega}_N = \frac{2}{TN(N-1)} \sum_{i=2}^N \sum_{j<i} (l'_i l_j)^2. \quad (\text{S.10})$$

Here $l_i = (w_i - \bar{w})\hat{\varepsilon}_i$ is constructed using residuals $\hat{\varepsilon}_i$. It can be expected that under reasonable regularity conditions on the memory properties of $\varepsilon_{i,t}$ as well as appropriate restrictions on N, T estimator $\hat{\Omega}_N$ is consistent in our setup. However, we do not attempt to prove this conjecture as it does not add to the main message of this paper.

Note that Baltagi et al. (2016) use mean adjustment variance estimate of Chen and Qin (2010), which is motivated by the need to obtain an unbiased (not only consistent) estimator of Ω . However, as in our setting x_i are correlated by construction, any theoretical justification for including $\bar{l}_{(i,j)}$ under null hypothesis is lost. Also note that a factor of 2 is missing in Baltagi et al. (2016).

S.2 Proofs: CCE power

Proof of Proposition 3.

Let $k_{N,T} = \sqrt{\frac{1}{2TN(N-1)}}$ and $\varpi_i = \left[\varsigma_{i,v}^{-1} \nu_i - \mathbf{D}_i \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \varsigma_{\ell,v}^{-1} \right) \right]$, where ν_i and $\varsigma_{v,i}^2$ are defined in the discussion following equation (31) and where

$$\mathbf{D}_i = \mathbf{F} \left[\boldsymbol{\lambda}_i^{(2)}, \mathbf{A}_i^{(2)} \right] + \left[\varepsilon_i, \mathbf{e}_i \right].$$

Using the same steps as in the Proof of Theorem 2, we can decompose the CD test statistic as

$$CD_{\mathbb{H}_1} = I - II$$

where

$$\begin{aligned} I &= 2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \varpi_i' \varpi_j + k_{N,T} \sum_{i=1}^N \varpi_i' \varpi_i - k_{N,T} \sum_{i,j}^N \varpi_i' \mathbf{P}_{\hat{\mathbf{F}}} \varpi_j \\ II &= k_{N,T} \sum_{i=1}^N \varsigma_{i,v}^{-2} \left(\nu_i' - \boldsymbol{\lambda}_i^{(1)'} \left(\overline{\mathbf{C}}^{(1)'} \right)^{-1} \overline{\mathbf{D}}' \right) \mathbf{M}_{\hat{\mathbf{F}}} \left(\nu_i - \overline{\mathbf{D}} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right). \end{aligned}$$

As shown in Lemmas 6 and 7, it holds that

$$k_{N,T} \sum_{i,j}^N \varpi_i' \mathbf{P}_{\hat{\mathbf{F}}} \varpi_j = \mathcal{O}_P \left[\max \left(N^{-1}, T^{-1} \right) \left(k_{N,T} \sum_{i,j}^N \varpi_i' \varpi_j \right) \right]$$

and

$$II = k_{N,T} \sum_{i=1}^N \varsigma_{i,v}^{-2} \nu_i' \nu_i + \mathcal{O}_P \left(N^{-1/2} \sqrt{T} \right) + \mathcal{O}_P \left(T^{-1/2} \right).$$

Furthermore, the expression $k_{N,T} \sum_{i=1}^N \varpi_i' \varpi_i$ is written out

$$\begin{aligned} k_{N,T} \sum_{i=1}^N \varpi_i' \varpi_i &= k_{N,T} \sum_{i=1}^N \varsigma_{i,v}^{-2} \nu_i' \nu_i \\ &\quad + \sqrt{\frac{TN}{(N-1)}} \left(N^{-1} \sum_{\ell=1}^N \varsigma_{\ell,v}^{-1} \boldsymbol{\lambda}_\ell^{(1)'} \right) \left(\overline{\mathbf{C}}^{(1)'} \right)^{-1} \left((NT)^{-1} \sum_{i=1}^N \mathbf{D}_i' \mathbf{D}_i \right) \\ &\quad \times \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \varsigma_{\ell,v}^{-1} \right) \\ &\quad - 2 \sqrt{\frac{TN}{(N-1)}} (NT)^{-1} \sum_{i=1}^N \varsigma_{i,v}^{-1} \nu_i' \mathbf{D}_i \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \varsigma_{\ell,v}^{-1} \right). \end{aligned}$$

The first term on the right-hand side above is canceled out by the leading term in II . The two remaining terms constitute expressions Φ_{1,\mathbb{H}_1} and Φ_{2,\mathbb{H}_1} , respectively. Next, consider the

properties of $2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \boldsymbol{\varpi}'_i \boldsymbol{\varpi}_j$, which we write as

$$2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \boldsymbol{\varpi}'_i \boldsymbol{\varpi}_j = 2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \zeta_{i,v}^{-1} \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \zeta_{j,v}^{-1} - 4k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \zeta_{i,v}^{-1} \boldsymbol{\nu}'_i \mathbf{D}_j \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,v}^{-1} \right) \quad (\text{S.11}) \\ + \left(N^{-1} \sum_{\ell=1}^N \zeta_{\ell,v}^{-1} \boldsymbol{\lambda}_\ell^{(1)'} \right) \left(\overline{\mathbf{C}}^{(1)'} \right)^{-1} \left(2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \mathbf{D}'_i \mathbf{D}_j \right) \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,v}^{-1} \right).$$

The order in probability of all three terms above is determined by the expected values of the elements within the double sum $\sum_{i=2}^N \sum_{j=1}^{i-1}$. Conditioning on the values taken by factor loadings and error variances, we have

$$\begin{aligned} \mathbb{E} \left[v_{i,t} v_{j,t} \mid \boldsymbol{\lambda}_i, \boldsymbol{\lambda}_j, \sigma_i^2, \sigma_j^2 \right] &= \boldsymbol{\lambda}_i^{(2)'} \boldsymbol{\Sigma}_{F,22} \boldsymbol{\lambda}_j^{(2)}, \\ \mathbb{E} \left[\mathbf{D}_{i,t} \mathbf{D}'_{j,t} \mid \boldsymbol{\lambda}_i, \boldsymbol{\lambda}_j, \mathbf{A}_i, \mathbf{A}_j, \sigma_i^2, \sigma_j^2 \right] &= \begin{bmatrix} \boldsymbol{\lambda}_i^{(2)'} \\ \mathbf{A}_i^{(2)'} \end{bmatrix} \boldsymbol{\Sigma}_{F,22} \begin{bmatrix} \boldsymbol{\lambda}_j^{(2)} \\ \mathbf{A}_j^{(2)} \end{bmatrix}, \\ \mathbb{E} \left[v_{i,t} \mathbf{D}'_{j,t} \mid \boldsymbol{\lambda}_i, \boldsymbol{\lambda}_j, \mathbf{A}_j, \sigma_i^2, \sigma_j^2 \right] &= \boldsymbol{\lambda}_i^{(2)'} \boldsymbol{\Sigma}_{F,22} \begin{bmatrix} \boldsymbol{\lambda}_j^{(2)} \\ \mathbf{A}_j^{(2)} \end{bmatrix}, \end{aligned}$$

for $i \neq j$. The zero mean property of $\boldsymbol{\lambda}_i^{(2)}$ and $\mathbf{A}_i^{(2)}$ implies that $\mathbb{E} \left[\mathbf{D}_{i,t} \mathbf{D}'_{j,t} \right] = \mathbf{0}$. This result carries over to $\mathbb{E} \left[\zeta_{i,v}^{-1} \boldsymbol{\nu}'_i \mathbf{D}_j \right]$ since independence of $\boldsymbol{\lambda}_i$ and σ_i^2 over cross-sections implies that we can write $\mathbb{E} \left[\zeta_{i,v}^{-1} \boldsymbol{\nu}'_i \mathbf{D}_j \right] = \mathbb{E} \left[\zeta_{i,v}^{-1} \boldsymbol{\lambda}_i^{(2)'} \right] \boldsymbol{\Sigma}_{F,22} \mathbb{E} \left\{ \begin{bmatrix} \boldsymbol{\lambda}_j^{(2)} \\ \mathbf{A}_j^{(2)} \end{bmatrix} \right\} = \mathbf{0}$. Consequently, the last two terms on the right-hand side of (S.11) are asymptotically centered around zero since their components $\overline{\mathbf{C}}^{(1)'}$ and $N^{-1} \sum_{\ell=1}^N \zeta_{\ell,v}^{-1} \boldsymbol{\lambda}_\ell^{(1)'}$ converge to nonstochastic limiting expressions. Lastly, by the LIE, we have

$$\mathbb{E} \left[\zeta_{i,v}^{-1} v_{i,t} v_{j,t} \zeta_{j,v}^{-1} \right] = \text{cov} \left[\zeta_{i,v}^{-1}, \boldsymbol{\lambda}_i^{(2)'} \right] \boldsymbol{\Sigma}_{F,22} \text{cov} \left[\boldsymbol{\lambda}_j^{(2)}, \zeta_{j,v}^{-1} \right],$$

and consequently

$$\mathbb{E} \left[2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \zeta_{i,v}^{-1} \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \zeta_{j,v}^{-1} \right] = N\sqrt{T} \text{cov} \left[\zeta_{1,v}^{-1}, \boldsymbol{\lambda}_1^{(2)'} \right] \boldsymbol{\Sigma}_{F,22} \text{cov} \left[\boldsymbol{\lambda}_1^{(2)}, \zeta_{1,v}^{-1} \right],$$

which parallels results obtained in the proof of Proposition 2. It will be shown below that variation around this expected value is of lower order in probability. Assuming this for now, we state that

$$\text{plim}_{N,T \rightarrow \infty} N^{-1} T^{-1/2} \sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \boldsymbol{\varpi}'_i \boldsymbol{\varpi}_j = \text{cov} \left[\zeta_{i,v}^{-1}, \boldsymbol{\lambda}_i^{(2)'} \right] \boldsymbol{\Sigma}_{F,22} \text{cov} \left[\boldsymbol{\lambda}_j^{(2)}, \zeta_{j,v}^{-1} \right].$$

A consequence of this result, as well as those on the second and third terms in equation (S.11) is that $2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \boldsymbol{\varpi}'_i \boldsymbol{\varpi}_j$ is asymptotically centered around 0 if $\zeta_{i,v}^{-1}$ and $\boldsymbol{\lambda}_i^{(2)'}$ are uncorrelated. In this special case, it is required to investigate the rate at which the variance of $2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \boldsymbol{\varpi}'_i \boldsymbol{\varpi}_j$ diverges. First, consider

$$\text{var} \left[2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \zeta_{i,v}^{-1} \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \zeta_{j,v}^{-1} \right] = \mathbb{E} \left[4k_{N,T}^2 \left(\sum_{i=2}^N \sum_{j=1}^{i-1} \zeta_{i,v}^{-1} \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \zeta_{j,v}^{-1} \right)^2 \right].$$

Apart from a superscript ⁽²⁾ on factors and factor loadings as well as the absence of the expected value of $\varsigma_{i,v}^{-1}$, the term $\varsigma_{i,v}^{-1} \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \varsigma_{j,v}^{-1}$ is identical to the one occurred in the proof of Proposition 2. Hence, we can follow the same reasoning to conclude that

$$\text{var} \left[\sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \varsigma_{i,v}^{-1} \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \varsigma_{j,v}^{-1} \right] = \mathcal{O}(T).$$

Next, consider the middle term of the third expression in equation (S.11). After vectorizing this matrix, we can write

$$\begin{aligned} \text{var} \left[\text{vec} \left(\sum_{i=2}^N \sum_{j=1}^{i-1} \mathbf{D}'_i \mathbf{D}_j \right) \right] &= \text{E} \left[\text{vec} \left(\sum_{i=2}^N \sum_{j=1}^{i-1} \mathbf{D}'_i \mathbf{D}_j \right) \text{vec} \left(\sum_{i'=2}^N \sum_{j'=1}^{i'-1} \mathbf{D}'_{i'} \mathbf{D}_{j'} \right)' \right] \\ &= \sum_{i=2}^N \sum_{j=1}^{i-1} \text{E} \left[\text{vec}(\mathbf{D}'_i \mathbf{D}_j) \text{vec}(\mathbf{D}'_i \mathbf{D}_j)' \right]. \end{aligned}$$

Recall here that

$$\begin{aligned} \mathbf{D}'_i \mathbf{D}_j &= \begin{bmatrix} \boldsymbol{\lambda}_i^{(2)} \\ \mathbf{A}_i^{(2)'} \end{bmatrix} \mathbf{F}^{(2)'} \mathbf{F}^{(2)} \begin{bmatrix} \boldsymbol{\lambda}_j^{(2)} \\ \mathbf{A}_j^{(2)} \end{bmatrix} + \begin{bmatrix} \boldsymbol{\varepsilon}'_i \\ \mathbf{E}'_i \end{bmatrix} [\boldsymbol{\varepsilon}_j, \mathbf{E}_j] \\ &\quad + \begin{bmatrix} \boldsymbol{\lambda}_i^{(2)} \\ \mathbf{A}_i^{(2)'} \end{bmatrix} \mathbf{F}^{(2)'} [\boldsymbol{\varepsilon}_j, \mathbf{E}_j] + \begin{bmatrix} \boldsymbol{\varepsilon}'_i \\ \mathbf{E}'_i \end{bmatrix} \mathbf{F}^{(2)} \begin{bmatrix} \boldsymbol{\lambda}_j^{(2)} \\ \mathbf{A}_j^{(2)} \end{bmatrix}, \end{aligned}$$

whose leading term is the first on the right-hand side of the first line and where we recall that $\mathbf{C}_i^{(2)} = \begin{bmatrix} \boldsymbol{\lambda}_i^{(2)} \\ \mathbf{A}_i^{(2)'} \end{bmatrix}$. Hence, the leading term of $\text{vec}(\mathbf{D}'_i \mathbf{D}_j) \text{vec}(\mathbf{D}'_i \mathbf{D}_j)'$ is written

$$\text{vec}(\mathbf{D}'_i \mathbf{D}_j) \text{vec}(\mathbf{D}'_i \mathbf{D}_j)' = \left(\mathbf{C}_i^{(2)} \otimes \mathbf{C}_j^{(2)} \right)' \text{vec}(\mathbf{F}^{(2)'} \mathbf{F}^{(2)}) \text{vec}(\mathbf{F}^{(2)'} \mathbf{F}^{(2)}) \left(\mathbf{C}_i^{(2)} \otimes \mathbf{C}_j^{(2)} \right),$$

which is nothing more than a multivariate version of the leading term in $\varsigma_{i,v}^{-1} \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \varsigma_{j,v}^{-1}$. The current case is even simpler due to the absence of the inverse error standard deviation $\varsigma_{i,v}^{-1}$. Consequently, finite fourth moments for factors and finite second moments for loadings together with the reasoning applied for the variance of $2k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \varsigma_{i,v}^{-1} \boldsymbol{\nu}'_i \boldsymbol{\nu}_j \varsigma_{j,v}^{-1}$ allow us to conclude that

$$\text{var} \left[\sqrt{\frac{2}{TN(N-1)}} \text{vec} \left(\sum_{i=2}^N \sum_{j=1}^{i-1} \mathbf{D}'_i \mathbf{D}_j \right) \right] = \mathcal{O}_P(T).$$

Since the outer terms $\left(N^{-1} \sum_{\ell=1}^N \varsigma_{\ell,v}^{-1} \boldsymbol{\lambda}_\ell^{(1)'} \right)$ and $\left(\overline{\mathbf{C}}^{(1)'} \right)^{-1}$ converge to finite, non-stochastic components, this rate of divergence carries over to the variance of the entire term $\left(N^{-1} \sum_{\ell=1}^N \varsigma_{\ell,v}^{-1} \boldsymbol{\lambda}_\ell^{(1)'} \right) \left(\overline{\mathbf{C}}^{(1)'} \right)^{-1}$

Lastly, consider the second term in (S.11), whose middle term is of particular interest. The variance of this term is given by

$$\begin{aligned} \text{var} \left[\sum_{i=2}^N \sum_{j=1}^{i-1} \varsigma_{i,v}^{-1} \boldsymbol{\nu}'_i \mathbf{D}_j \right] &= \mathbb{E} \left[\sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{i'=2}^N \sum_{j'=1}^{i'-1} \mathbf{D}'_j \boldsymbol{\nu}_i \boldsymbol{\nu}'_{i'} \mathbf{D}_{j'} \varsigma_{i,v}^{-1} \varsigma_{i',v}^{-1} \right] \\ &= \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t,t'}^T \mathbb{E} \left[\mathbf{D}_{j,t} \boldsymbol{\nu}_{i,t} \boldsymbol{\nu}'_{i,t'} \mathbf{D}'_{j,t'} \varsigma_{i,v}^{-2} \right]. \end{aligned}$$

Similar to the previous two terms, the leading expression in the term above is given by

$$\mathbb{E} \left[\mathbf{C}_j^{(2)'} \mathbf{F}_t^{(2)} \mathbf{F}_t^{(2)'} \boldsymbol{\lambda}_i^{(2)} \boldsymbol{\lambda}_i^{(2)'} \mathbf{F}_t^{(2)} \mathbf{F}_t^{(2)'} \mathbf{C}_j^{(2)} \varsigma_{i,v}^{-2} \right] \leq M \mathbb{E} \left[\mathbf{C}_j^{(2)'} \boldsymbol{\lambda}_i^{(2)} \varsigma_{i,v}^{-2} \boldsymbol{\lambda}_i^{(2)'} \mathbf{C}_j^{(2)} \right]$$

where the upper bound on the right-hand side above is established by the LIE and boundedness of the fourth-order moments of \mathbf{F}_t . Additionally assuming that $\mathbb{E} \left[\boldsymbol{\lambda}_i^{(2)} \varsigma_{i,v}^{-2} \boldsymbol{\lambda}_i^{(2)'} \right]$ is bounded and conditioning on $\mathbf{C}_j^{(2)}$, we arrive at

$$\sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t,t'}^T \mathbb{E} \left[\mathbf{D}_{j,t} \boldsymbol{\nu}_{i,t} \boldsymbol{\nu}'_{i,t'} \mathbf{D}'_{j,t'} \varsigma_{i,v}^{-2} \right] = \mathcal{O} \left((NT)^2 \right),$$

implying that

$$\text{var} \left[\sqrt{\frac{2}{TN(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \varsigma_{i,v}^{-1} \boldsymbol{\nu}'_i \mathbf{D}_j \right] = \mathcal{O}(T),$$

which carries over to the variance of $k_{N,T} \sum_{i=2}^N \sum_{j=1}^{i-1} \varsigma_{i,v}^{-1} \boldsymbol{\nu}'_i \mathbf{D}_j \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \varsigma_{\ell,v}^{-1} \right)$. \square

S.3 Auxiliary lemmas

Lemma 1. Let $\{a_{i,t}\}_{t=1}^T$ be a scalar and \mathbf{c}_i L -dimensional sequences of random variables for $i = 1, \dots, N$ such that:

- $a_{i,t}, a_{j,s}$ are independent for all $i \neq j$ and all s, t .
- $a_{i,t}, a_{i,s}$ are iid conditionally on \mathbf{c}_i for all s, t .
- $\mathbb{E}[|a_{i,t}|^8 | \mathbf{c}_i] < M < \infty$. \mathbf{c}_i are iid over i .
- $q_i^k \equiv \mathbb{E}[|a_{i,t}|^k | \mathbf{c}_i]$ and $\mathbb{E}[a_{i,t} | \mathbf{c}_i] = 0$ for $k \leq 8$.
- $\mathbb{E}[|q_i^k|] < \infty$ for $k \leq 8$.

Then:

$$U = \sqrt{\frac{2}{NT(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T a_{i,t} a_{j,t} \xrightarrow{d} N(0, \mathbb{E}[q_i^2]^2), \quad (\text{S.12})$$

jointly as $N, T \rightarrow \infty$.

Proof of Lemma 1.

To prove this lemma we use Theorem 3.2 of Hall and Heyde (1980). In particular, express U as:

$$U = \sum_{t=1}^T \zeta_{t,N,T}, \quad \zeta_{t,N,T} = \sqrt{\frac{2}{NT(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} a_{i,t} a_{j,t}. \quad (\text{S.13})$$

Denote by $\mathcal{C} = \sigma(\{\mathbf{c}_1, \dots, \mathbf{c}_N\})$ the σ -algebra generated by \mathbf{c}_i and let $\mathcal{F}_{t-1,N,T} = \sigma(\mathcal{C}, \zeta_{t-1,N,T}, \dots, \zeta_{1,N,T})$ be the σ -algebra generated by \mathcal{C} and $\zeta_{t-1,N,T}, \dots, \zeta_{1,N,T}$. It is easy to see that $\{\zeta_{t,N,T}, \mathcal{F}_{t-1,N,T} : t = 1, \dots, T\}$ is a Martingale Difference Array.

At first we establish the limiting variance of this MDS array. From Corollary 3.1 in Hall and Heyde (1980) the variance is determined from:

$$V_T = \sum_{t=1}^T \mathbb{E}[\zeta_{t,N,T}^2 | \mathcal{F}_{t-1,N,T}] \xrightarrow{p} \eta^2 \quad (\text{S.14})$$

By conditional independence of $a_{i,t}$, the above result simplifies:

$$\begin{aligned}
\sum_{t=1}^T \mathbb{E}[\zeta_{t,N,T}^2 | \mathcal{F}_{t-1,N,T}] &= \sum_{t=1}^T \mathbb{E}[\tilde{\zeta}_{t,N,T}^2 | \mathcal{C}] \\
&= \frac{2}{N(N-1)} \mathbb{E} \left[\left(\sum_{i=2}^N \sum_{j=1}^{i-1} a_{i,t} a_{j,t} \right)^2 \middle| \mathcal{C} \right] \\
&= \frac{2}{N(N-1)} \mathbb{E} \left[\left(\sum_{i=2}^N a_{i,t} A_{i-1,t} \right)^2 \middle| \mathcal{C} \right] \\
&= \frac{2}{N(N-1)} \mathbb{E} \left[\sum_{i=2}^N a_{i,t}^2 A_{i-1,t}^2 \middle| \mathcal{C} \right] \\
&= \frac{2}{N(N-1)} \sum_{i=2}^N q_i^2 \sum_{j=1}^{i-1} q_j^2,
\end{aligned}$$

where in the third line we defined the “integrated” variable $A_{i-1,t} = \sum_{j=1}^{i-1} a_{j,t}$, the fourth line uses the fact that $a_{i,t} A_{i-1,t}$ is a Martingale Difference Sequence. The last line uses the definition of q_i^2 . After re-arranging elements:

$$\begin{aligned}
V_T &= \frac{2}{N(N-1)} \sum_{i=2}^N q_i^2 \sum_{j=1}^{i-1} q_j^2 \\
&= \frac{1}{N(N-1)} \left(N^2 \left(\frac{1}{N} \sum_{i=1}^N q_i^2 \right)^2 - N \left(\frac{1}{N} \sum_{i=1}^N q_i^4 \right) \right) \\
&= \left(\frac{1}{N} \sum_{i=1}^N q_i^2 \right)^2 + \mathcal{O}_P(N^{-1}) \\
&= \mathbb{E}[q_i^2]^2 + \mathcal{O}_P(N^{-1}).
\end{aligned}$$

Here the third and the fourth lines follow from the application of the Kolmogorov’s SLLN to iid sequences q_i^2 and q_i^4 . Thus we can expect that

$$U = \sqrt{\frac{2}{NT(N-1)}} \sum_{i=2}^N \sum_{j=1}^{i-1} \sum_{t=1}^T a_{i,t} a_{j,t} \xrightarrow{d} N(0, \mathbb{E}[q_i^2]^2). \quad (\text{S.15})$$

It remains to prove that the sufficient condition for Corollary 3.1 in Hall and Heyde (1980) is satisfied. In particular, it is sufficient to show that $\zeta_{t,N,T}$ is a (conditionally) uniformly integrable sequence:

$$\sum_{t=1}^T \mathbb{E}[\zeta_{t,N,T}^2 I(|\zeta_{t,N,T}| > \varepsilon) | \mathcal{F}_{t-1,N,T}] \xrightarrow{p} 0. \quad (\text{S.16})$$

Instead of proving uniform integrability, we borrow the idea from Kao et al. (2012) and instead show that conditional Lyapunov's condition:

$$B_T = \sum_{t=1}^T \mathbb{E}[|\xi_{t,N,T}|^{2+\delta} | \mathcal{F}_{t-1,N,T}] \xrightarrow{p} 0, \quad \delta > 0, \quad (\text{S.17})$$

is satisfied under the maintained assumptions. In what follows we prove that the above condition is satisfied for $\delta = 2$. Observe how:

$$B_T = T \mathbb{E}[\zeta_{t,N,T}^4 | \mathcal{C}] = T^{-1} \frac{N}{4(N-1)} \mathbb{E} \left[\left(N\bar{a}^2 - \bar{a}^2 \right)^4 | \mathcal{C} \right] = T^{-1} \frac{N}{4(N-1)} \|N\bar{a}^2 - \bar{a}^2\|_4^4,$$

where $\bar{a} = N^{-1} \sum_{i=1}^N a_{i,t}$ and similarly for \bar{a}^2 . For the ease of exposition we drop the t subscript from the definition of the above averages as these quantities are identically distributed over t . The scaling in front of the expectation is of order $o(1)$ as long as $T \rightarrow \infty$. It remains to show that the second component is $\mathcal{O}_P(1)$. By Minkowski inequality with respect to the conditional probability measure:

$$\|N\bar{a}^2 - \bar{a}^2\|_4 \leq \|N\bar{a}^2\|_4 + \|\bar{a}^2\|_4. \quad (\text{S.18})$$

Similarly:

$$\begin{aligned} \|\bar{a}^2\|_4 &\leq \frac{1}{N} \sum_{i=1}^N \sqrt[4]{\mathbb{E}[a_{i,t}^8 | \mathcal{C}_i]} \\ &= \frac{1}{N} \sum_{i=1}^N \sqrt[4]{q_i^8} \\ &= \mathbb{E} \left[\sqrt[4]{q_i^8} \right] + \mathcal{O}_P(1). \end{aligned}$$

Here the first lines from the Minkowski inequality, second line from the definition of the $\|\cdot\|_4$ norm and the definition of q_i^8 . The final line follows from the application of the Kolmogorov's SLLN.

When it comes to $\|N\bar{a}^2\|_4$ observe that:

$$\|N\bar{a}^2\|_4^4 = \frac{1}{N^4} \sum_{i=1}^N \sum_{j=1}^N \sum_{k=1}^N \sum_{l=1}^N \sum_{m=1}^N \sum_{n=1}^N \sum_{p=1}^N \sum_{s=1}^N \mathbb{E}[a_{i,t} a_{j,t} a_{k,t} a_{l,t} a_{m,t} a_{n,t} a_{p,t} a_{s,t} | \mathcal{C}]. \quad (\text{S.19})$$

Because $a_{i,t}$ are conditionally independent, the expectation is nonzero only if the indices are pairwise equal. This leaves $\mathcal{O}(N^4)$ non-zero summands. When non-zero each expectation is a function of at most q_i^k , which we assume is integrable by assumption. Thus:

$$\begin{aligned} \|N\bar{a}^2\|_4^4 &\leq \left(\frac{1}{N} \sum_{i=1}^N q_i^2 \right)^4 + \mathcal{O}_P(1) \\ &= (\mathbb{E}[q_i^2])^4 + \mathcal{O}_P(1), \end{aligned}$$

where the last line follows from the Kolmogorov's SLLN and the Continuous Mapping Theorem. Combining all results we showed that:

$$\|N\bar{a}^2 - \bar{a}^2\|_4^4 = \mathcal{O}_P(1), \quad (\text{S.20})$$

and as a result:

$$B_T = T \mathbb{E}[z_{t,N,T}^4 | \mathcal{C}] = T^{-1} \frac{N}{4(N-1)} \|N\bar{a}^2 - \bar{a}^2\|_4^4 = \mathcal{O}_P(1).$$

As required. This completes the proof. \square

Lemma 2. *Let w_i denote any stochastic weights that are independent across i and satisfy $\mathbb{E}[w_i^2] < M$ for all i . Furthermore, let w_i be independent of \mathbf{f}_t , $\varepsilon_{i,t}$ and $\mathbf{e}_{i,t}$ for all i and t . Under Assumptions 2-6,*

$$\left(T^{-1} \hat{\mathbf{F}}' \hat{\mathbf{F}}\right)^{-1} = \mathcal{O}_P(1) \quad (\text{S.21})$$

$$\sqrt{NT} T^{-1/2} \left\| N^{-1} \sum_{i=1}^N w_i \mathbf{U}'_i \right\| = \mathcal{O}_P(1) \quad (\text{S.22})$$

$$\sqrt{NT} T^{-1/2} \left\| \left(N^{-1} \sum_{i=1}^N w_i \mathbf{U}'_i \right) \mathbf{F} \right\| = \mathcal{O}_P(1) \quad (\text{S.23})$$

Additionally, under Assumptions 2-6 and given weights w_i that satisfy Assumption 7, we have

$$\left\| (NT)^{-1} \sum_{i=1}^N \mathbf{U}'_i \varepsilon_i w_i \right\| = \mathcal{O}_P(N^{-1/2}) \quad (\text{S.24})$$

Proof of Lemma 2.

1. *Result (S.21):* By Pesaran (2006, equation (36)) it holds that

$$T^{-1} \hat{\mathbf{F}}' \hat{\mathbf{F}} = \bar{\mathbf{C}}' \left(T^{-1} \mathbf{F}' \mathbf{F} \right) \bar{\mathbf{C}} + \mathcal{O}_P((NT)^{-1/2}) + \mathcal{O}_P(N^{-1}),$$

where the function of true factors and true loadings itself converges to a positive definite matrix by assumptions 3 and 4. Hence, Theorem 1 in Karabiyik et al. (2017) applies and equation (S.21) follows.

2. *Result (S.22):* Taking the square of the expression of interest, we can write

$$NT^{-1} \left\| N^{-1} \sum_{i=1}^N w_i \mathbf{U}'_i \right\|^2 \leq (NT)^{-1} \sum_{i,j} \sum_{t=1}^T w_i w_j (\varepsilon_{i,t} \varepsilon_{j,t} + \mathbf{e}'_{i,t} \mathbf{e}_{j,t}) \|\mathbf{B}\|^2$$

where $(NT)^{-1} \sum_{i,j} \sum_{t=1}^T w_i w_j \varepsilon_{i,t} \varepsilon_{j,t} = \mathcal{O}_P(1)$ and $(NT)^{-1} \sum_{i,j} \sum_{t=1}^T w_i w_j \mathbf{e}'_{i,t} \mathbf{e}_{j,t} = \mathcal{O}_P(1)$ by Markov's inequality and zero correlation of idiosyncratic variation in both \mathbf{y}_i and \mathbf{X}_i across i . Result (S.22) follows accordingly.

3. *Result (S.23)*: Analogous to the proof of (S.22) we take the square of (S.23) and rearrange to arrive at

$$NT^{-1} \left\| N^{-1} \sum_{i=1}^N w_i \mathbf{U}'_i \mathbf{F} \right\|^2 \leq (NT)^{-1} \sum_{i,j}^N \sum_{t,t'}^T w_i w_j \mathbf{f}'_t \mathbf{f}_{t'} (\varepsilon_{i,t} \varepsilon_{j,t'} + \mathbf{e}'_{i,t} \mathbf{e}_{j,t'}) \|\mathbf{B}\|^2.$$

Now taking expectations of the non-negative expression $(NT)^{-1} \sum_{i,j}^N \sum_{t,t'}^T w_i w_j \mathbf{f}'_t \mathbf{f}_{t'} \varepsilon_{i,t} \varepsilon_{j,t'}$ and using uncorrelatedness of $\varepsilon_{i,t}$ across i and t as well as $E(w_i^2) \leq M$, we obtain

$$\begin{aligned} E \left[(NT)^{-1} \sum_{i,j}^N \sum_{t,t'}^T w_i w_j \mathbf{f}'_t \mathbf{f}_{t'} \varepsilon_{i,t} \varepsilon_{j,t'} \right] &\leq (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T M \operatorname{tr}(\boldsymbol{\Sigma}_{\mathbf{F}}) E[\varepsilon_{i,t}^2] \\ &= \mathcal{O}(1) \end{aligned}$$

An identical result is given for the term involving $\mathbf{f}'_t \mathbf{f}_{t'}$. *Result (S.23)* is then obtained via Markov's inequality.

4. *Result (S.24)*: Given the definition of \mathbf{U}_i , we can write

$$E \left[\left\| (NT)^{-1} \sum_{i=1}^N \mathbf{U}'_i \boldsymbol{\varepsilon}_i w_i \right\|^2 \right] \leq (NT)^{-2} \sum_{i=1}^N \sum_{t,t'}^T E[\varepsilon_{i,t} (\varepsilon_{i,t} \varepsilon_{i,t'} + \mathbf{e}'_{i,t} \mathbf{e}_{i,t'}) \varepsilon_{i,t'}] \operatorname{var}[w_i] \|\mathbf{B}\|^2$$

where we directly use independence of w_i across i as well as its zero expected value. Additionally,

$$\begin{aligned} (NT)^{-2} \sum_{i=1}^N \sum_{t,t'}^T E[\varepsilon_{i,t} \varepsilon_{i,t} \varepsilon_{i,t'} \varepsilon_{i,t'}] &= (NT)^{-2} \sum_{i=1}^N \sum_{t=1}^T \kappa_4[\varepsilon_{i,t}] + (NT)^{-2} \sum_{i=1}^N \sum_{t,t'}^T E[\varepsilon_{i,t}^2] E[\varepsilon_{i,t'}^2] \\ &= \mathcal{O}(N^{-1}) \end{aligned}$$

and

$$\begin{aligned} (NT)^{-2} \sum_{i=1}^N \sum_{t,t'}^T E[\varepsilon_{i,t} \mathbf{e}'_{i,t} \mathbf{e}_{i,t'} \varepsilon_{i,t'}] &= (NT)^{-2} \sum_{i=1}^N \sum_{t=1}^T E[\varepsilon_{i,t}^2] \operatorname{tr}(\boldsymbol{\Sigma}) \\ &= \mathcal{O}\left((NT)^{-1}\right), \end{aligned}$$

from which it follows that

$$\left\| (NT)^{-1} \sum_{i=1}^N \mathbf{U}'_i \boldsymbol{\varepsilon}_i w_i \right\|^2 = \mathcal{O}_P(N^{-1})$$

by Markov's inequality. Taking the square root leads to result (S.24). □

Lemma 3. Under Assumptions 2-6 and Assumption 7,

$$\begin{aligned} & \sqrt{\frac{1}{2TN(N-1)}} \sum_{ij}^N \left(\varepsilon_i - \bar{U} (\bar{\mathbf{C}}^{-1})' \boldsymbol{\lambda}_i \right)' \mathbf{P}_{\hat{\mathbf{F}}} \left(\varepsilon_j - \bar{U} (\bar{\mathbf{C}}^{-1})' \boldsymbol{\lambda}_j \right) w_i w_j \\ &= \mathcal{O}_P \left(T^{-1/2} \right) + \mathcal{O}_P \left(N^{-1/2} \right) + \mathcal{O}_P \left(N^{-1} \sqrt{T} \right) \end{aligned}$$

Proof of Lemma 3.

Let $\bar{\boldsymbol{\lambda}}_w = \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell w_\ell \right)$ and note that $\sum_{i=1}^N \left(\varepsilon_i - \bar{U} (\bar{\mathbf{C}}^{-1})' \boldsymbol{\lambda}_i \right) w_i = \sum_{i=1}^N \left(\varepsilon_i w_i - \mathbf{U}_i (\bar{\mathbf{C}}^{-1}) \bar{\boldsymbol{\lambda}}_w \right)$.

This allows us to write

$$\begin{aligned} & \sqrt{\frac{1}{2TN(N-1)}} \sum_{ij}^N \left(\varepsilon_i - \bar{U} (\bar{\mathbf{C}}^{-1})' \boldsymbol{\lambda}_i \right)' \mathbf{P}_{\hat{\mathbf{F}}} \left(\varepsilon_j - \bar{U} (\bar{\mathbf{C}}^{-1})' \boldsymbol{\lambda}_j \right) w_i w_j \\ & \leq \sqrt{\frac{1}{2TN(N-1)}} \left(\left\| \sum_{i=1}^N \hat{\mathbf{F}}' \varepsilon_i w_i \right\| + \left\| \sum_{i=1}^N \hat{\mathbf{F}}' \mathbf{U}_i \right\| \left\| \bar{\mathbf{C}}^{-1} \right\| \left\| \bar{\boldsymbol{\lambda}}_w \right\| \right)^2 \left\| (\hat{\mathbf{F}}' \hat{\mathbf{F}})^{-1} \right\|. \end{aligned}$$

First, note that

$$\left\| \bar{\boldsymbol{\lambda}}_w \right\|^2 = N^{-2} \sum_{ij}^N w_i w_j \boldsymbol{\lambda}_i' \boldsymbol{\lambda}_j = \mathcal{O}_P(1) \quad (\text{S.25})$$

since $\mathbb{E} \left[N^{-2} \sum_{ij}^N w_i w_j \boldsymbol{\lambda}_i' \boldsymbol{\lambda}_j \right] \leq \left(N^{-2} \sum_{ij}^N \mathbb{E} \left[w_i^2 w_j^2 \right] \right)^{1/2} \left(N^{-2} \sum_{ij}^N \mathbb{E} \left[\left\| \boldsymbol{\lambda}_i \right\|^2 \left\| \boldsymbol{\lambda}_j \right\|^2 \right] \right)^{1/2} = \mathcal{O}(1)$ by boundedness of the fourth moments of all stochastic components involved. Next, recall that $\hat{\mathbf{F}} = \mathbf{F} \bar{\mathbf{C}} + \bar{\mathbf{U}}$. Thus,

$$\begin{aligned} \left\| \sum_{i=1}^N \hat{\mathbf{F}}' \mathbf{U}_i \right\| & \leq \sqrt{NT} \left\| \bar{\mathbf{C}} \right\| \left\| \sqrt{N/T} \mathbf{F}' \bar{\mathbf{U}} \right\| + T \left\| \sqrt{N/T} \bar{\mathbf{U}} \right\|^2 \\ & = \mathcal{O}_P \left(\sqrt{NT} \right) + \mathcal{O}_P(T), \end{aligned}$$

where the last line is a consequence of results (S.22) and (S.23). Furthermore, since $\varepsilon_i = \mathbf{U}_i \mathbf{B}^{-1} [1; \mathbf{0}']'$, we can proceed analogously for $\left\| \sum_{i=1}^N \hat{\mathbf{F}}' \varepsilon_i w_i \right\| \leq \left\| \sum_{i=1}^N \hat{\mathbf{F}}' \mathbf{U}_i w_i \right\| \left\| \mathbf{B}^{-1} \right\|$ and obtain the same orders in probability as in the last equation above. Additionally using result (S.21), we arrive at

$$\begin{aligned} & \sqrt{\frac{1}{2TN(N-1)}} \sum_{ij}^N \left(\varepsilon_i - \bar{U} (\bar{\mathbf{C}}^{-1})' \boldsymbol{\lambda}_i \right)' \mathbf{P}_{\hat{\mathbf{F}}} \left(\varepsilon_j - \bar{U} (\bar{\mathbf{C}}^{-1})' \boldsymbol{\lambda}_j \right) w_i w_j \\ & = \sqrt{\frac{1}{2TN(N-1)}} \left[\mathcal{O}_P \left(\sqrt{NT} \right) + \mathcal{O}_P(T) \right]^2 \mathcal{O}_P \left(T^{-1} \right) \\ & = \mathcal{O}_P \left(T^{-1/2} \right) + \mathcal{O}_P \left(N^{-1/2} \right) + \mathcal{O}_P \left(N^{-1} \sqrt{T} \right), \end{aligned}$$

which concludes this proof. \square

Lemma 4. Under Assumptions 2-6 and Assumption 7,

$$\begin{aligned} & \sqrt{\frac{1}{2TN(N-1)}} \sum_{i=1}^N (\varepsilon_i - \bar{U} (\bar{\mathbf{C}}^{-1}) \lambda_i)' \mathbf{P}_{\hat{\mathbf{F}}} (\varepsilon_i - U (\bar{\mathbf{C}}^{-1}) \lambda_i) w_i^2 \\ & = \mathcal{O}_P(N^{-2}\sqrt{T}) + \mathcal{O}_P(N^{-3/2}) + \mathcal{O}_P(T^{-1/2}) \end{aligned}$$

Proof of Lemma 4.

Given that $\|\mathbf{A}_1 - \mathbf{A}_2\|^2 \leq 3\|\mathbf{A}_1\|^2 + 3\|\mathbf{A}_2\|^2$ for two arbitrary $m \times n$ matrices \mathbf{A}_1 and \mathbf{A}_2 , we have

$$\begin{aligned} & \sqrt{\frac{1}{2TN(N-1)}} \sum_{i=1}^N (\varepsilon_i - \bar{U} (\bar{\mathbf{C}}^{-1}) \lambda_i)' \mathbf{P}_{\hat{\mathbf{F}}} (\varepsilon_i - U (\bar{\mathbf{C}}^{-1}) \lambda_i) w_i^2 \\ & \leq 3\sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \varepsilon_i' \hat{\mathbf{F}} \right\|^2 \left\| (T^{-1} \hat{\mathbf{F}}' \hat{\mathbf{F}})^{-1} \right\| \\ & + 3\sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \lambda_i' (\bar{\mathbf{C}}^{-1})' \bar{U}' \hat{\mathbf{F}} \right\|^2 \left\| (T^{-1} \hat{\mathbf{F}}' \hat{\mathbf{F}})^{-1} \right\|. \end{aligned} \quad (\text{S.26})$$

Here,

$$\begin{aligned} & \sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \varepsilon_i' \hat{\mathbf{F}} \right\|^2 \\ & = 3\sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \varepsilon_i' \mathbf{F} \right\|^2 \|\bar{\mathbf{C}}\|^2 + 3\sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \varepsilon_i' \bar{U} \right\|^2. \end{aligned}$$

Concerning the first term on the right-hand side above, we can use Chebyshev's inequality together with

$$\begin{aligned} \mathbb{E} \left[\sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N T^{-2} w_i^2 \varepsilon_i' \mathbf{F} \mathbf{F}' \varepsilon_i \right] & = \sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \sum_{t,t'}^T T^{-2} \mathbb{E} [w_i^2] \mathbb{E} [\mathbb{E} [\varepsilon_{i,t} \varepsilon_{i,t'} | \sigma_i]] \mathbb{E} [\mathbf{f}_t' \mathbf{f}_{t'}] \\ & = \sqrt{\frac{1}{2TN(N-1)}} \sum_{i=1}^N \mathbb{E} [\sigma_i^2] \mathbb{E} [w_i^2] \text{tr}(\boldsymbol{\Sigma}_{\mathbf{F}}) \\ & = \mathcal{O}(T^{-1/2}) \end{aligned}$$

to arrive at $\sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \varepsilon_i' \mathbf{F} \right\|^2 \|\bar{\mathbf{C}}\|^2 = \mathcal{O}_P(T^{-1/2})$. Likewise, recalling that $\mathbf{u}_{i,t} =$

$\mathbf{B}' [\varepsilon_{i,t}; \mathbf{e}'_{i,t}]'$, we have

$$\begin{aligned}
& \mathbb{E} \left[\sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \varepsilon_i' \bar{\mathbf{U}} \right\|^2 \right] \\
& \leq \sqrt{\frac{T}{2N(N-1)}} \sum_{i,i',i''}^N \sum_{t,t'}^T (NT)^{-2} \mathbb{E} [w_i^2] \mathbb{E} [\varepsilon_{i,t} (\varepsilon_{i',t} \varepsilon_{i'',t} + \mathbf{e}'_{i',t} \mathbf{e}_{i'',t}) \varepsilon_{i,t}] \|\mathbf{B}\| \\
& = \sqrt{\frac{T}{2N(N-1)}} \left(\mathcal{O}(N^{-1}) + \mathcal{O}(T^{-1}) \right) \mathcal{O}(1) \\
& = \mathcal{O}(N^{-2} \sqrt{T}) + \mathcal{O}(N^{-1} T^{-1/2}).
\end{aligned}$$

This result is obtained by noting that

$$\begin{aligned}
(NT)^{-2} \sum_{i,i',i''}^N \sum_{t,t'}^T \mathbb{E} [\varepsilon_{i,t} \varepsilon_{i',t} \varepsilon_{i'',t}] &= (NT)^{-2} \sum_{i=1}^N \sum_{t=1}^T \kappa_4 [\varepsilon_{i,t}] + (NT)^{-2} \sum_{i=1}^N \sum_{t,t'}^T \mathbb{E} [\varepsilon_{i,t}^2] \mathbb{E} [\varepsilon_{i,t'}^2] \\
&+ (NT)^{-2} \sum_{i,i'}^N \sum_{t=1}^T \mathbb{E} [\varepsilon_{i,t}^2] \mathbb{E} [\varepsilon_{i',t}^2] \\
&= \mathcal{O}(N^{-1}) + \mathcal{O}_p(T^{-1})
\end{aligned}$$

and

$$\begin{aligned}
(NT)^{-2} \sum_{i,i',i''}^N \sum_{t,t'}^T \mathbb{E} [\varepsilon_{i,t} \mathbf{e}'_{i',t} \mathbf{e}_{i'',t} \varepsilon_{i,t}] &= (NT)^{-2} \sum_{i,i'}^N \sum_{t=1}^T \mathbb{E} [\varepsilon_{i,t}^2] \mathbb{E} [\mathbf{e}'_{i',t} \mathbf{e}_{i',t}] \\
&= \mathcal{O}(T^{-1}).
\end{aligned}$$

Combining the results obtained up to this point and additionally using (S.21), we obtain

$$\sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \varepsilon_i' \hat{\mathbf{F}} \right\|^2 \left\| (T^{-1} \hat{\mathbf{F}}' \hat{\mathbf{F}})^{-1} \right\| = \mathcal{O}_p(N^{-2} \sqrt{T}) + \mathcal{O}_p(T^{-1/2}). \quad (\text{S.27})$$

Next, consider

$$\sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \boldsymbol{\lambda}'_i (\bar{\mathbf{C}}^{-1})' \bar{\mathbf{U}} \hat{\mathbf{F}} \right\|^2 = \sqrt{\frac{NT}{2(N-1)}} \left(N^{-1} \sum_{i=1}^N w_i^2 \boldsymbol{\lambda}'_i \boldsymbol{\lambda}_i \right) \|\bar{\mathbf{C}}^{-1}\|^2 \left\| T^{-1} \bar{\mathbf{U}} \hat{\mathbf{F}} \right\|^2$$

where $N^{-1} \sum_{i=1}^N w_i^2 \boldsymbol{\lambda}'_i \boldsymbol{\lambda}_i = \mathcal{O}_p(1)$ since $\mathbb{E} \left[N^{-1} \sum_{i=1}^N w_i^2 \boldsymbol{\lambda}'_i \boldsymbol{\lambda}_i \right] \leq \left(N^{-1} \sum_{i=1}^N \mathbb{E} [w_i^4] \right)^{1/2} \left(N^{-1} \sum_{i=1}^N \mathbb{E} [\|\boldsymbol{\lambda}_i\|^4] \right)^{1/2}$

Additionally, we can write

$$\begin{aligned}
\left\| T^{-1} \bar{\mathbf{U}} \hat{\mathbf{F}} \right\|^2 &\leq \left(\left\| T^{-1} \mathbf{F}' \bar{\mathbf{U}} \right\| + \left\| T^{-1} \bar{\mathbf{U}}' \bar{\mathbf{U}} \right\| \right)^2 \\
&= \mathcal{O}_p(N^{-2}) + \mathcal{O}_p(N^{-3/2} T^{-1/2}) + \mathcal{O}_p((NT)^{-1})
\end{aligned}$$

with the last step following from (S.22) and (S.23). Consequently, it holds that

$$\begin{aligned} & \sqrt{\frac{T}{2N(N-1)}} \sum_{i=1}^N \left\| T^{-1} w_i \boldsymbol{\lambda}_i' (\overline{\mathbf{C}}^{-1})' \overline{\mathbf{U}} \hat{\mathbf{F}} \right\|^2 \left\| (T^{-1} \hat{\mathbf{F}}' \hat{\mathbf{F}})^{-1} \right\| \\ &= \mathcal{O}_P(N^{-2} \sqrt{T}) + \mathcal{O}_P(N^{-3/2}) + \mathcal{O}_P(N^{-1} T^{-1/2}). \end{aligned}$$

Summarizing the results derived so far, we have

$$\begin{aligned} & \sqrt{\frac{1}{2TN(N-1)}} \sum_{i=1}^N (\boldsymbol{\varepsilon}_i - \overline{\mathbf{U}} (\overline{\mathbf{C}}^{-1}) \boldsymbol{\lambda}_i)' \mathbf{P}_{\hat{\mathbf{F}}} (\boldsymbol{\varepsilon}_i - \mathbf{U} (\overline{\mathbf{C}}^{-1}) \boldsymbol{\lambda}_i) w_i^2 \\ &= \mathcal{O}_P(N^{-2} \sqrt{T}) + \mathcal{O}_P(N^{-3/2}) + \mathcal{O}_P(T^{-1/2}) \end{aligned}$$

which concludes the proof. \square

Lemma 5. *Suppose that the true model is (8) and that Assumptions 2–5 and 8 hold. Also, assume that the factor estimator $\hat{\mathbf{F}}$ is given by $\hat{\mathbf{F}} = [\overline{\mathbf{y}}, \overline{\mathbf{X}}]$. Then we can without loss of generality assume that the true factors and loadings can be partitioned into*

$$\mathbf{f}_t = \begin{bmatrix} \mathbf{f}_t^{(1)} \\ \mathbf{f}_t^{(2)} \end{bmatrix} \quad \text{and} \quad \boldsymbol{\lambda}_i = \begin{bmatrix} \boldsymbol{\lambda}_i^{(1)} \\ \boldsymbol{\lambda}_i^{(2)} \end{bmatrix} \quad \text{and} \quad \mathbf{A}_i = \begin{bmatrix} \mathbf{A}_i^{(1)} \\ \mathbf{A}_i^{(2)} \end{bmatrix}$$

such that $\mathbb{E} [\mathbf{f}_t^{(1)} \mathbf{f}_t^{(2)'}] = \mathbf{O}$, $\mathbb{E} [\boldsymbol{\lambda}_i^{(2)}] = \mathbf{0}$ and $\mathbb{E} [\mathbf{A}_i^{(2)}] = \mathbf{O}$. It holds that $\mathbf{f}^{(1)}$ and $\mathbf{f}^{(2)}$ are $(m+1) \times 1$ and $(r-m-1) \times 1$, respectively. $\boldsymbol{\lambda}_i^{(1)}$ and $\boldsymbol{\lambda}_i^{(2)}$ are $(m+1) \times 1$ and $(r-m-1) \times 1$. Lastly, $\mathbf{A}_i^{(1)}$ and $\mathbf{A}_i^{(2)}$ are $(m+1) \times m$ and $(r-m-1) \times m$.

Proof of Lemma 5.

Under a violation of the rank condition, cross-section averages of all variables will only identify the $m+1$ -dimensional linear combination $\mathbf{f}_t^{(1)} = \mathbf{C}_0' \mathbf{f}_t$ of the r true common factors. The $r \times (m+1)$ matrix \mathbf{C}_0 is given by

$$\mathbf{C}_0 = [\mathbb{E} [\boldsymbol{\lambda}_i], \mathbb{E} [\mathbf{A}_i]] \mathbf{B}.$$

Now define the $r \times (m+1)$ matrix $\boldsymbol{\Phi}_1 = \mathbb{E} [\mathbf{f}_t \mathbf{f}_t^{(1)'}] \left(\mathbb{E} [\mathbf{f}_t^{(1)} \mathbf{f}_t^{(1)'}] \right)^{-1}$ and the $r \times 1$ vector $\mathbf{f}_t^\perp = \mathbf{f}_t - \boldsymbol{\Phi}_1 \mathbf{f}_t^{(1)}$ and note that by construction it holds that $\mathbb{E} [\mathbf{f}_t^\perp \mathbf{f}_t^{(1)'}] = \mathbf{O}$. Using these two expressions, we can decompose

$$\boldsymbol{\lambda}_i' \mathbf{f}_t = \boldsymbol{\lambda}_i' \boldsymbol{\Phi}_1 \mathbf{f}_t^{(1)} + \boldsymbol{\lambda}_i' \mathbf{f}_t^\perp. \quad (\text{S.28})$$

The expression above decomposes the common component of a random variable that has a factor structure into the variation due to the $(m+1)$ -dimensional linear combination of the r true factors that is identified by a cross-section average-based factor estimator and residual

common variation that is not picked up. This residual source of common variation is bound to be part of the error term in a misspecified factor model.

Next, consider the product of C'_0 and \mathbf{f}_t^\perp

$$\begin{aligned} C'_0 \mathbf{f}_t^\perp &= C'_0 \mathbf{f}_t - C'_0 \mathbb{E} \left[\mathbf{f}_t \mathbf{f}_t^{(1)'} \right] \left(\mathbb{E} \left[\mathbf{f}_t^{(1)} \mathbf{f}_t^{(1)'} \right] \right)^{-1} \mathbf{f}_t^{(1)} \\ &= \mathbf{f}_t^{(1)} - \mathbb{E} \left[C'_0 \mathbf{f}_t \mathbf{f}_t^{(1)'} \right] \left(\mathbb{E} \left[\mathbf{f}_t^{(1)} \mathbf{f}_t^{(1)'} \right] \right)^{-1} \mathbf{f}_t^{(1)} \\ &= \mathbf{f}_t^{(1)} - \mathbf{f}_t^{(1)} \\ &= \mathbf{0}, \end{aligned}$$

In order to appreciate the implications of this result, let $\text{col}(C_0)$ denote the column space of C_0 and $\ker(C'_0)$ the kernel of C'_0 . The result reported in the last sequence of equations above states that $\mathbf{f}_t^\perp \in \ker(C'_0)$. Note now that $\dim(\text{col}(C_0)) + \dim(\ker(C'_0)) = r$ (see e.g. Abadir and Magnus 2005, Exercise 4.3(c)) and that $\dim(\text{col}(C_0)) = \dim(\text{col}(C'_0))$ (Abadir and Magnus 2005, Exercise 4.5(a)). As a result of the rank condition on C_0 , we have $\dim(\text{col}(C'_0)) = m + 1$ and hence $\dim(\ker(C'_0)) = r - m - 1$. Consequently, \mathbf{f}_t^\perp is an element in the $s = r - m - 1$ -dimensional vector space orthogonal to C_0 . This implies that, given an arbitrary basis for $\ker(C'_0)$, here denoted by the $r \times s$ -dimensional matrix Φ_2 , we can express \mathbf{f}_t^\perp as $\mathbf{f}_t^\perp = \Phi_2 \mathbf{f}_t^{(2)}$ where $\mathbf{f}_t^{(2)}$ is some $s \times 1$ vector of coordinates. Plugging this expression into (S.28) results in

$$\begin{aligned} \lambda'_i \mathbf{f}_t &= \lambda'_i \Phi_1 \mathbf{f}_t^{(1)} + \lambda'_i \Phi_2 \mathbf{f}_t^{(2)} \\ &= \lambda'_i \begin{bmatrix} \Phi_1 & \Phi_2 \end{bmatrix} \begin{bmatrix} \mathbf{f}_t^{(1)} \\ \mathbf{f}_t^{(2)} \end{bmatrix} \\ &= \begin{bmatrix} \lambda_i^{(1)'} & \lambda_i^{(2)'} \end{bmatrix} \begin{bmatrix} \mathbf{f}_t^{(1)} \\ \mathbf{f}_t^{(2)} \end{bmatrix} \end{aligned}$$

From the previously mentioned result $\mathbb{E} \left[\mathbf{f}_t^\perp \mathbf{f}_t^{(1)'} \right] = \mathbf{0}$ it follows that $\mathbb{E} \left[\mathbf{f}_t^{(2)} \mathbf{f}_t^{(1)'} \right] = \mathbf{0}$. Furthermore, given that $\mathbb{E}[\lambda_i] = C_0 B^{-1} [1, \mathbf{0}']'$ and that $\lambda_i^{(2)'} = \lambda'_i \Phi_2$ where Φ_2 is a basis for $\ker(C'_0)$,

$$\begin{aligned} \mathbb{E} \left[\lambda_i^{(2)} \right] &= \Phi_2' C_0 B^{-1} [1, \mathbf{0}']' \\ &= \mathbf{0}. \end{aligned}$$

Analogously, we can define $[\Lambda_i^{(1)'}, \Lambda_i^{(2)'}] = \Lambda'_i [\Phi_1, \Phi_2]$ where $\mathbb{E}[\Lambda_i^{(2)}] = \mathbf{0}$ by the same reasoning as above.

Consequently, whenever the factor estimates include cross-section averages of the dependent variable Y , we can rewrite the common component as being generated by a linear combination of the true factors and the true loadings such that the two moment conditions above are satisfied. The matrix $\Phi = (\Phi_1, \Phi_2)$ which links the loadings λ'_i (Λ'_i) and $[\lambda_i^{(1)'}, \lambda_i^{(2)'}]$

($[\Lambda_i^{(1)'}, \Lambda_i^{(2)'}]$) is of full rank by the rank assumptions on $E[\mathbf{f}_t \mathbf{f}_t']$ as well as C_0 . Hence, Φ is an invertible rotation and we can write $[\lambda_i^{(1)'}, \lambda_i^{(2)'}] [\mathbf{f}_t^{(1)'}, \mathbf{f}_t^{(2)'}]' = \lambda_i' \Phi \Phi^{-1} \mathbf{f}_t$. Now note that in factor models it is only the common component $\lambda_i' \mathbf{f}_t$ that is identified whereas the loadings λ_i and factors \mathbf{f}_t are identified only up to an invertible rotation. As argued above, the matrix Φ is invertible. Hence, irrespective of the true values of λ_i , Λ_i and \mathbf{f}_t (subject to the assumptions we make on them), we can assume that the data are generated by $\Phi' \lambda_i$, $\Phi' \Lambda_i$ and $\Phi^{-1} \mathbf{f}_t$ instead. \square

Lemma 6. *Under Assumptions 2-6,*

$$k_{N,T} \sum_{i,j} \boldsymbol{\varpi}_i' \mathbf{P}_{\hat{\mathbf{F}}} \boldsymbol{\varpi}_j = \mathcal{O}_P \left[\max(N^{-1}, T^{-1}) \left(k_{N,T} \sum_{i,j} \boldsymbol{\varpi}_i' \boldsymbol{\varpi}_j \right) \right] \quad (\text{S.29})$$

Proof of Lemma 6. Note that

$$\boldsymbol{\varpi}_i = \left[\zeta_{i,\nu}^{-1} \boldsymbol{\nu}_i - \mathbf{D}_i (\bar{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,\nu}^{-1} \right) \right]$$

where

$$\begin{aligned} \boldsymbol{\nu}_i &= \boldsymbol{\varepsilon}_i + \mathbf{F}^{(2)} \boldsymbol{\lambda}_i^{(2)}, \\ \mathbf{D}_i &= \mathbf{U}_i + \mathbf{F}^{(2)} \mathbf{C}_i^{(2)}. \end{aligned}$$

Then, we can rewrite

$$\begin{aligned} \boldsymbol{\varpi}_i &= \left[\boldsymbol{\varepsilon}_i \zeta_{i,\nu}^{-1} - \mathbf{U}_i (\bar{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,\nu}^{-1} \right) \right] \\ &\quad + \mathbf{F}^{(2)} \left[\boldsymbol{\lambda}_i^{(2)} \zeta_{i,\nu}^{-1} - \mathbf{C}_i^{(2)} (\bar{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,\nu}^{-1} \right) \right]. \end{aligned}$$

Consequently, we can write out

$$\begin{aligned} &k_{N,T} \sum_{i,j} \boldsymbol{\varpi}_i' \mathbf{P}_{\hat{\mathbf{F}}} \boldsymbol{\varpi}_j \\ &\leq 2 \sqrt{\frac{T}{2N(N-1)}} \left\| \left(T^{-1} \hat{\mathbf{F}}' \hat{\mathbf{F}} \right)^{-1} \right\| \left\| T^{-1} \hat{\mathbf{F}}' \mathbf{F}^{(2)} \right\|^2 \\ &\quad \times \left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(2)} \zeta_{i,\nu}^{-1} - \sum_{i=1}^N \mathbf{C}_i^{(2)} (\bar{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,\nu}^{-1} \right) \right\|^2 \\ &\quad + 2k_{N,T} \sum_{i,j} \left(\boldsymbol{\varepsilon}_i \zeta_{i,\nu}^{-1} - \mathbf{U}_i (\bar{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,\nu}^{-1} \right) \right)' \mathbf{P}_{\hat{\mathbf{F}}} \\ &\quad \times \mathbf{P}_{\hat{\mathbf{F}}} \left(\boldsymbol{\varepsilon}_j \zeta_{j,\nu}^{-1} - \mathbf{U}_j (\bar{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \zeta_{\ell,\nu}^{-1} \right) \right), \end{aligned} \quad (\text{S.30})$$

where the second line is of order $\mathcal{O}_P(T^{-1/2}) + \mathcal{O}_P(N^{-1/2}) + \mathcal{O}_P[N^{-1}\sqrt{T}]$ by Lemma 3 with $w_i = \varsigma_{i,\nu}^{-1}$. Concerning the first line, it holds that

$$\sqrt{\frac{T}{2N(N-1)}} \left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(2)} \varsigma_{i,\nu}^{-1} - \sum_{i=1}^N \mathbf{C}_i^{(2)} (\overline{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \varsigma_{\ell,\nu}^{-1} \right) \right\|^2 = \mathcal{O}_P \left(k_{N,T} \sum_{i,j}^N \boldsymbol{\varpi}'_i \boldsymbol{\varpi}_j \right). \quad (\text{S.31})$$

To appreciate this result, we write

$$\begin{aligned} & \left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(2)} \varsigma_{i,\nu}^{-1} - \sum_{i=1}^N \mathbf{C}_i^{(2)} (\overline{\mathbf{C}}^{(1)})^{-1} \left(N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \varsigma_{\ell,\nu}^{-1} \right) \right\|^2 \\ &= 3 \left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(2)} \varsigma_{i,\nu}^{-1} \right\|^2 + 3 \left\| \sum_{i=1}^N \mathbf{C}_i^{(2)} \right\|^2 \left\| (\overline{\mathbf{C}}^{(1)})^{-1} \right\|^2 \left\| N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \varsigma_{\ell,\nu}^{-1} \right\|^2 \end{aligned}$$

and note that $N^{-2} \left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(1)} \varsigma_{i,\nu}^{-1} \right\|^2 = \mathcal{O}_P(1)$ by the same reasoning leading to (S.25). Furthermore, $\left\| \sum_{i=1}^N \mathbf{C}_i^{(2)} \right\|^2 = \mathcal{O}_P(N)$ holds by the zero mean property of $\boldsymbol{\lambda}_i^{(2)}$ and $\boldsymbol{\lambda}_i^{(2)}$. Consequently,

$$\left\| \sum_{i=1}^N \mathbf{C}_i^{(2)} \right\|^2 \left\| (\overline{\mathbf{C}}^{(1)})^{-1} \right\|^2 \left\| N^{-1} \sum_{\ell=1}^N \boldsymbol{\lambda}_\ell^{(1)} \varsigma_{\ell,\nu}^{-1} \right\|^2 = \mathcal{O}_P(N),$$

which never has a higher order in probability than $\left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(2)} \varsigma_{i,\nu}^{-1} \right\|^2$. We prove this claim by noting that the expected value of $\left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(2)} \varsigma_{i,\nu}^{-1} \right\|^2$ is given by

$$E \left[\left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(2)} \varsigma_{i,\nu}^{-1} \right\|^2 \right] = \sum_{i=1}^N E \left[\varsigma_{i,\nu}^{-2} \boldsymbol{\lambda}_i^{(2)'} \boldsymbol{\lambda}_i^{(2)} \right] + \sum_{i \neq j}^N E \left[\varsigma_{i,\nu}^{-1} \boldsymbol{\lambda}_i^{(2)'} \right] E \left[\boldsymbol{\lambda}_j^{(2)} \varsigma_{j,\nu}^{-1} \right].$$

It follows straightforwardly from Markov's inequality that

$$\left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(2)} \varsigma_{i,\nu}^{-1} \right\|^2 = \begin{cases} \mathcal{O}_P(N) & \text{if } \text{cov} \left[\boldsymbol{\lambda}_j^{(2)}; \varsigma_{j,\nu}^{-1} \right] = \mathbf{0} \\ \mathcal{O}_P(N^2) & \text{otherwise} \end{cases}.$$

The reduction in the order in probability due to a zero correlation between $\boldsymbol{\lambda}_i^{(2)}$ and $\varsigma_{i,\nu}^{-1}$ is the same as observed in the case of $k_{N,T} \sum_{i,j}^N \boldsymbol{\varpi}'_i \boldsymbol{\varpi}_j$. Additionally using the scaling $\sqrt{\frac{T}{2N(N-1)}}$ and using our result on the order of $k_{N,T} \sum_{i,j}^N \boldsymbol{\varpi}'_i \boldsymbol{\varpi}_j$ in the proof of Proposition 3, we arrive at

$$\sqrt{\frac{T}{2N(N-1)}} \left\| \sum_{i=1}^N \boldsymbol{\lambda}_i^{(2)} \varsigma_{i,\nu}^{-1} \right\|^2 = \mathcal{O}_P \left(k_{N,T} \sum_{i,j}^N \boldsymbol{\varpi}'_i \boldsymbol{\varpi}_j \right),$$

which leads to result (S.31). Now returning to the first line of expression (S.30), we consider

$$\begin{aligned}
& \left\| T^{-1} \hat{\mathbf{F}}' \mathbf{F}^{(2)} \right\|^2 \\
& \leq 3 \left\| \bar{\mathbf{C}}^{(1)'} \right\|^2 \left\| T^{-1} \mathbf{F}^{(1)'} \mathbf{F}^{(2)} \right\|^2 + 9 \left\| \bar{\mathbf{C}}^{(2)'} \right\|^2 \left\| T^{-1} \mathbf{F}^{(2)'} \mathbf{F}^{(2)} \right\|^2 + k_{N,T} \sum_{i,j} \varpi_i' \varpi_j \left\| T^{-1} \bar{\mathbf{U}}' \mathbf{F}^{(2)} \right\|^2 \\
& = \mathcal{O}_P \left(T^{-1} \right) + \mathcal{O}_P \left(N^{-1} \right) + \mathcal{O}_P \left((NT)^{-1} \right), \tag{S.32}
\end{aligned}$$

which holds by zero correlation between $\mathbf{F}^{(1)}$ and $\mathbf{F}^{(2)}$, the zero expected value of $\bar{\mathbf{C}}^{(2)}$ and result (S.23). Consequently, an upper bound for the order in probability of (S.30) is given by

$$k_{N,T} \sum_{i,j} \varpi_i' \mathbf{P}_{\hat{\mathbf{F}}} \varpi_j = \mathcal{O}_P \left[\max \left[N^{-1}, T^{-1} \right] \left(k_{N,T} \sum_{i,j} \varpi_i' \varpi_j \right) \right].$$

□

Lemma 7. *Under Assumptions 2-6,*

$$\begin{aligned}
& k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \left(\boldsymbol{\nu}_i' - \boldsymbol{\lambda}_i^{(1)'} \left(\bar{\mathbf{C}}^{(1)'} \right)^{-1} \bar{\mathbf{D}}' \right) \mathbf{M}_{\hat{\mathbf{F}}} \left(\boldsymbol{\nu}_i' - \bar{\mathbf{D}} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right) \\
& = k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\nu}_i' \boldsymbol{\nu}_i + \mathcal{O}_P \left(N^{-1/2} \sqrt{T} \right) + \mathcal{O}_P \left(T^{-1/2} \right)
\end{aligned}$$

Proof of Lemma 7. The term that we are interested in can be expanded into

$$\begin{aligned}
& k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \left(\boldsymbol{\nu}_i' - \boldsymbol{\lambda}_i^{(1)'} \left(\bar{\mathbf{C}}^{(1)'} \right)^{-1} \bar{\mathbf{D}}' \right) \mathbf{M}_{\hat{\mathbf{F}}} \left(\boldsymbol{\nu}_i' - \bar{\mathbf{D}} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right) \\
& = k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\nu}_i' \boldsymbol{\nu}_i - 2k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\nu}_i' \bar{\mathbf{D}} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \\
& \quad + k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\lambda}_i^{(1)'} \left(\bar{\mathbf{C}}^{(1)'} \right)^{-1} \bar{\mathbf{D}}' \bar{\mathbf{D}} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \\
& \quad + k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \left(\boldsymbol{\nu}_i' - \boldsymbol{\lambda}_i^{(1)'} \left(\bar{\mathbf{C}}^{(1)'} \right)^{-1} \bar{\mathbf{D}}' \right) \mathbf{P}_{\hat{\mathbf{F}}} \left(\boldsymbol{\nu}_i' - \bar{\mathbf{D}} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right) \\
& = k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\nu}_i' \boldsymbol{\nu}_i - 2a_1 + a_2 + a_3,
\end{aligned}$$

where we recall that $\boldsymbol{\nu}_i = \boldsymbol{\varepsilon}_i + \mathbf{F}^{(2)} \boldsymbol{\lambda}_i^{(2)}$ and $\mathbf{D}_i = \mathbf{U}_i + \mathbf{F}^{(2)} \mathbf{C}_i^{(2)}$. Now consider a_3 . Similar to the proof of Lemma 6, we can write

$$\begin{aligned}
a_3 & \leq 2k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \left(\boldsymbol{\lambda}_i^{(2)} - \bar{\mathbf{C}}^{(2)} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right)' \mathbf{F}^{(2)'} \mathbf{P}_{\hat{\mathbf{F}}} \mathbf{F}^{(2)} \left(\boldsymbol{\lambda}_i^{(2)} - \bar{\mathbf{C}}^{(2)} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right) \\
& \quad + 2k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \left(\boldsymbol{\varepsilon}_i - \bar{\mathbf{U}} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right)' \mathbf{P}_{\hat{\mathbf{F}}} \left(\boldsymbol{\varepsilon}_i - \bar{\mathbf{U}} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right).
\end{aligned}$$

By Lemma 4, the last line in the expression above is $\mathcal{O}_P(N^{-2}\sqrt{T}) + \mathcal{O}_P(N^{-3/2}) + \mathcal{O}_P(T^{-1/2})$. Concerning the remaining term, we can write

$$\begin{aligned} & k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \left(\boldsymbol{\lambda}_i^{(2)} - \overline{\mathbf{C}}^{(2)} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right)' \mathbf{F}^{(2)'} \mathbf{P}_{\hat{\mathbf{F}}} \mathbf{F}^{(2)} \left(\boldsymbol{\lambda}_i^{(2)} - \overline{\mathbf{C}}^{(2)} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right) \\ & \leq T k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \left\| \boldsymbol{\lambda}_i^{(2)} - \overline{\mathbf{C}}^{(2)} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right\|^2 \left\| T^{-1} \hat{\mathbf{F}}' \mathbf{F}^{(2)} \right\|^2 \left\| \left(T^{-1} \hat{\mathbf{F}}' \hat{\mathbf{F}} \right)^{-1} \right\| \end{aligned}$$

where we focus on the term

$$\begin{aligned} & T k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \left\| \boldsymbol{\lambda}_i^{(2)} - \overline{\mathbf{C}}^{(2)} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right\|^2 \\ & \leq \left(T k_{N,T} 3 \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\lambda}_i^{(2)'} \boldsymbol{\lambda}_i^{(2)} \right) + \left(T k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\lambda}_i^{(1)'} \boldsymbol{\lambda}_i^{(1)} \right) \left\| \overline{\mathbf{C}}^{(2)} \right\|^2 \left\| \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \right\|^2 \\ & = \mathcal{O}_P(\sqrt{T}). \end{aligned}$$

This result follows from applying the reasoning given below equation (S.27) to both weighted sums over i in the expression above. Combining it with result (S.32), we arrive at

$$\begin{aligned} & k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \left(\boldsymbol{\lambda}_i^{(2)} - \overline{\mathbf{C}}^{(2)} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right)' \mathbf{F}^{(2)'} \mathbf{P}_{\hat{\mathbf{F}}} \mathbf{F}^{(2)} \left(\boldsymbol{\lambda}_i^{(2)} - \overline{\mathbf{C}}^{(2)} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right) \\ & = \mathcal{O}_P(N^{-1}\sqrt{T}) + \mathcal{O}_P(T^{-1/2}). \end{aligned}$$

which, together with Lemma 4, leads to

$$a_3 = \mathcal{O}_P(N^{-1}\sqrt{T}) + \mathcal{O}_P(T^{-1/2}). \quad (\text{S.33})$$

Next, we investigate a_2 and write

$$\begin{aligned} a_2 & \leq k_{N,T} \sum_{i=1}^N \left\| \varsigma_{i,\nu}^{-1} \boldsymbol{\lambda}_i^{(1)'} \right\|^2 \left\| \left(\overline{\mathbf{C}}^{(1)'} \right)^{-1} \right\|^2 3 \left(\left\| \overline{\mathbf{C}}^{(2)} \right\|^2 \left\| \mathbf{F} \right\|^2 + \left\| \overline{\mathbf{U}} \right\|^2 \right) \\ & = \mathcal{O}_P(N^{-1}\sqrt{T}), \end{aligned} \quad (\text{S.34})$$

where the last line follows from result (S.22), the zero mean of $\overline{\mathbf{C}}^{(2)}$ and the result on $\sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\lambda}_i^{(1)'} \boldsymbol{\lambda}_i^{(1)}$ referred to above. The last term to consider is given by

$$\begin{aligned} a_1 & = k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\varepsilon}_i' \overline{\mathbf{U}} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} + k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\lambda}_i^{(2)'} \mathbf{F}^{(2)'} \overline{\mathbf{U}} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \\ & \quad + k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\varepsilon}_i' \mathbf{F}^{(2)} \overline{\mathbf{C}}^{(2)} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \\ & \quad + k_{N,T} \sum_{i=1}^N \varsigma_{i,\nu}^{-2} \boldsymbol{\lambda}_i^{(2)'} \mathbf{F}^{(2)'} \mathbf{F}^{(2)} \overline{\mathbf{C}}^{(2)} \left(\overline{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \\ & = a_{11} + a_{12} + a_{13} + a_{14}. \end{aligned}$$

Here, $a_{11} = \mathcal{O}_p(N^{-1/2}) + \mathcal{O}_p(N^{-1}\sqrt{T})$ results from proceeding almost exactly as in the steps leading to equation (44) in the proof of Theorem 2. Concerning the second term, we can write

$$\begin{aligned} |a_{12}| &\leq k_{N,T} \sum_{i=1}^N \left\| \zeta_{i,\nu}^{-1} \boldsymbol{\lambda}_i^{(2)} \right\| \left\| \mathbf{F}^{(2)'} \bar{\mathbf{U}} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \right\| \left\| \zeta_{i,\nu}^{-1} \boldsymbol{\lambda}_i^{(1)} \right\| \\ &\leq k_{N,T} N \sqrt{\frac{T}{N}} \left(N^{-1} \sum_{i=1}^N \left\| \zeta_{i,\nu}^{-1} \boldsymbol{\lambda}_i^{(2)} \right\|^2 \right)^{1/2} \left(N^{-1} \sum_{i=1}^N \left\| \zeta_{i,\nu}^{-1} \boldsymbol{\lambda}_i^{(1)} \right\|^2 \right)^{1/2} \left\| \sqrt{\frac{N}{T}} \mathbf{F}^{(2)'} \bar{\mathbf{U}} \right\| \left\| \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \right\| \\ &= \mathcal{O}_p(N^{-1/2}), \end{aligned}$$

where we make use of (S.23) with $w_i = 1$. With regards to the third term, we have

$$\begin{aligned} |a_{13}| &\leq k_{N,T} \sqrt{NT} \left((NT)^{-1} \sum_{i=1}^N \left\| \boldsymbol{\varepsilon}_i' \mathbf{F}^{(2)} \right\|^2 \right)^{1/2} \left(N^{-1} \sum_{i=1}^N \left\| \boldsymbol{\lambda}_i^{(1)} \zeta_{i,\nu}^{-2} \right\|^2 \right)^{1/2} \left\| \sqrt{N} \bar{\mathbf{C}}^{(2)} \right\| \left\| \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \right\| \\ &= \mathcal{O}_p(N^{-1/2}). \end{aligned}$$

Here, we can use results from the proof of Lemma 4 to obtain $(NT)^{-1} \sum_{i=1}^N \left\| \boldsymbol{\varepsilon}_i' \mathbf{F}^{(2)} \right\|^2 = \mathcal{O}_p(1)$.

Lastly, we can apply Cauchy-Schwarz to $|a_{14}|$ in order to arrive at

$$\begin{aligned} |a_{14}| &= k_{N,T} \sqrt{NT} \left(N^{-1} \sum_{i=1}^N \left\| \zeta_{i,\nu}^{-1} \boldsymbol{\lambda}_i^{(2)} \right\|^2 \right)^{1/2} \left(N^{-1} \sum_{i=1}^N \left\| \zeta_{i,\nu}^{-1} \boldsymbol{\lambda}_i^{(1)} \right\|^2 \right)^{1/2} \\ &\quad \times \left\| T^{-1/2} \mathbf{F}^{(2)} \right\|^2 \left\| \sqrt{N} \bar{\mathbf{C}}^{(2)} \right\| \left\| \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \right\| \\ &= \mathcal{O}_p(N^{-1/2} \sqrt{T}). \end{aligned}$$

Using the last four intermediary results allows us to conclude that

$$a_1 = \mathcal{O}_p(N^{-1/2} \sqrt{T}).$$

This result, together with (S.33) and (S.34) establishes

$$\begin{aligned} &k_{N,T} \sum_{i=1}^N \zeta_{i,\nu}^{-2} \left(\boldsymbol{\nu}_i' - \boldsymbol{\lambda}_i^{(1)'} \left(\bar{\mathbf{C}}^{(1)'} \right)^{-1} \bar{\mathbf{D}}' \right) \mathbf{M}_{\hat{\mathbf{F}}} \left(\boldsymbol{\nu}_i' - \bar{\mathbf{D}} \left(\bar{\mathbf{C}}^{(1)} \right)^{-1} \boldsymbol{\lambda}_i^{(1)} \right) \\ &= k_{N,T} \sum_{i=1}^N \zeta_{i,\nu}^{-2} \boldsymbol{\nu}_i' \boldsymbol{\nu}_i + \mathcal{O}_p(N^{-1/2} \sqrt{T}) + \mathcal{O}_p(T^{-1/2}), \end{aligned}$$

which was to be shown. □

S.4 Additional MC results

This section reports additional results for the Monte Carlo experiments in Section 5 of the main paper. The five tables below are equivalents to Tables 1-5 and their underlying setup differs only in that the errors in $y_{i,t}$ and $x_{i,t}$ are drawn from a standardized $\chi^2(2)$ distribution with zero expected value and unit variance.

Table S.1: Sample moments of original CD test statistic for remaining CSD under \mathbb{H}_0

Part A: Application of CD to 2WFE residuals

| c_σ : | | 0.1 | 0.5 | 1 | 1.5 | 0.1 | 0.5 | 1 | 1.5 |
|--------------|-----|--------|--------|-------|-------|-----------------------|------|------|-------|
| N | T | Mean | | | | Variance $\times 100$ | | | |
| 25 | 25 | -3.35 | -3.31 | -3.20 | -2.97 | 1.28 | 1.77 | 3.94 | 9.92 |
| 25 | 50 | -4.91 | -4.87 | -4.72 | -4.43 | 0.54 | 0.89 | 2.70 | 8.46 |
| 25 | 100 | -7.08 | -7.02 | -6.82 | -6.43 | 0.23 | 0.60 | 2.71 | 9.99 |
| 25 | 200 | -10.11 | -10.02 | -9.75 | -9.21 | 0.10 | 0.49 | 3.35 | 14.57 |
| 50 | 25 | -3.30 | -3.26 | -3.14 | -2.91 | 1.03 | 1.38 | 2.91 | 7.33 |
| 50 | 50 | -4.86 | -4.80 | -4.65 | -4.33 | 0.38 | 0.65 | 1.99 | 6.37 |
| 50 | 100 | -7.00 | -6.94 | -6.73 | -6.32 | 0.14 | 0.36 | 1.69 | 6.41 |
| 50 | 200 | -10.00 | -9.91 | -9.63 | -9.06 | 0.06 | 0.28 | 1.94 | 8.66 |
| 100 | 25 | -3.28 | -3.24 | -3.11 | -2.86 | 0.88 | 1.16 | 2.35 | 5.87 |
| 100 | 50 | -4.83 | -4.78 | -4.62 | -4.30 | 0.26 | 0.45 | 1.38 | 4.50 |
| 100 | 100 | -6.97 | -6.90 | -6.68 | -6.26 | 0.09 | 0.23 | 1.06 | 4.22 |
| 100 | 200 | -9.95 | -9.86 | -9.57 | -8.99 | 0.03 | 0.16 | 1.07 | 4.77 |
| 200 | 25 | -3.27 | -3.23 | -3.10 | -2.85 | 0.73 | 0.96 | 1.95 | 4.90 |
| 200 | 50 | -4.82 | -4.76 | -4.60 | -4.28 | 0.21 | 0.35 | 1.01 | 3.24 |
| 200 | 100 | -6.95 | -6.88 | -6.66 | -6.23 | 0.07 | 0.17 | 0.75 | 2.95 |
| 200 | 200 | -9.92 | -9.83 | -9.54 | -8.94 | 0.02 | 0.11 | 0.71 | 3.21 |

Part B: Application of CD to CCE residuals

| c_σ : | | 0.1 | 0.5 | 1 | 1.5 | 0.1 | 0.5 | 1 | 1.5 |
|--------------|-----|--------|--------|-------|-------|-----------------------|------|------|-------|
| N | T | Mean | | | | Variance $\times 100$ | | | |
| 25 | 25 | -3.35 | -3.31 | -3.21 | -3.02 | 1.37 | 1.86 | 3.90 | 9.30 |
| 25 | 50 | -4.92 | -4.87 | -4.74 | -4.47 | 0.50 | 0.86 | 2.66 | 8.14 |
| 25 | 100 | -7.08 | -7.02 | -6.84 | -6.48 | 0.22 | 0.57 | 2.63 | 9.45 |
| 25 | 200 | -10.11 | -10.03 | -9.77 | -9.28 | 0.10 | 0.50 | 3.26 | 13.22 |
| 50 | 25 | -3.28 | -3.24 | -3.13 | -2.90 | 1.20 | 1.60 | 3.30 | 8.04 |
| 50 | 50 | -4.85 | -4.81 | -4.66 | -4.37 | 0.38 | 0.60 | 1.73 | 5.41 |
| 50 | 100 | -7.00 | -6.94 | -6.74 | -6.35 | 0.13 | 0.34 | 1.54 | 5.83 |
| 50 | 200 | -10.00 | -9.91 | -9.64 | -9.09 | 0.06 | 0.26 | 1.76 | 7.69 |
| 100 | 25 | -3.23 | -3.19 | -3.07 | -2.83 | 1.23 | 1.59 | 2.97 | 6.77 |
| 100 | 50 | -4.82 | -4.77 | -4.61 | -4.31 | 0.31 | 0.48 | 1.32 | 4.09 |
| 100 | 100 | -6.97 | -6.90 | -6.69 | -6.27 | 0.09 | 0.22 | 1.01 | 4.01 |
| 100 | 200 | -9.95 | -9.86 | -9.57 | -9.01 | 0.03 | 0.17 | 1.11 | 4.89 |
| 200 | 25 | -3.16 | -3.12 | -3.00 | -2.75 | 1.64 | 1.90 | 3.04 | 6.35 |
| 200 | 50 | -4.80 | -4.75 | -4.59 | -4.27 | 0.27 | 0.40 | 1.05 | 3.22 |
| 200 | 100 | -6.94 | -6.88 | -6.67 | -6.24 | 0.07 | 0.16 | 0.71 | 2.86 |
| 200 | 200 | -9.92 | -9.83 | -9.54 | -8.96 | 0.02 | 0.10 | 0.69 | 3.17 |

Notes. $\varepsilon_{i,t}$ and $e_{i,t}$ are standardized $\chi^2(2)$ with zero expected value and unit variance. $\sigma_{i,y}^2 = c_\sigma (\zeta_{i,y}^2 - 2) / 4 + 1$; $\zeta_{i,y}^2$ is $\chi^2(2)$. In part B, $r = 2$ and loadings λ_i are drawn from $U(0.5, 1.5)$. \mathbf{A}_i has first element from $U(0.5, 1.5)$ and second from $U(-0.5, 0.5)$. Factors \mathbf{f}_i drawn from $N(0, 1)$. In part A, we restrict $\mathbf{f}_i = [f_i^{(1)}, 1]'$, $\lambda_i = [1, \lambda_i^{(2)}]'$ and $\mathbf{A}_i = [1, \Lambda_i^{(2)}]'$.

Table S.2: Sample moments of original CD test statistic applied to 2WFE residuals under \mathbb{H}_1

| N | λ_i : σ_i^2 : T | Mean | | | | Variance | | | |
|-----|--------------------------------------|-------------------|----------------|-------------------|----------------|-------------------|----------------|-------------------|----------------|
| | | symmetric | | skewed | | symmetric | | skewed | |
| | | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ |
| 25 | 25 | -3.37 | -3.09 | -3.30 | -2.68 | 0.02 | 0.10 | 0.02 | 0.41 |
| 25 | 50 | -4.86 | -4.50 | -4.78 | -3.89 | 0.01 | 0.13 | 0.02 | 0.70 |
| 25 | 100 | -6.95 | -6.47 | -6.86 | -5.63 | 0.02 | 0.25 | 0.02 | 1.27 |
| 25 | 200 | -9.88 | -9.25 | -9.76 | -8.06 | 0.02 | 0.39 | 0.02 | 2.23 |
| 50 | 25 | -3.32 | -2.87 | -3.25 | -2.19 | 0.01 | 0.14 | 0.02 | 0.58 |
| 50 | 50 | -4.81 | -4.23 | -4.73 | -3.26 | 0.01 | 0.18 | 0.01 | 0.94 |
| 50 | 100 | -6.88 | -6.09 | -6.78 | -4.78 | 0.01 | 0.31 | 0.01 | 1.55 |
| 50 | 200 | -9.78 | -8.73 | -9.65 | -6.85 | 0.01 | 0.52 | 0.02 | 2.80 |
| 100 | 25 | -3.30 | -2.49 | -3.23 | -1.34 | 0.01 | 0.27 | 0.01 | 0.98 |
| 100 | 50 | -4.78 | -3.76 | -4.70 | -2.13 | 0.01 | 0.32 | 0.01 | 1.42 |
| 100 | 100 | -6.83 | -5.47 | -6.73 | -3.19 | 0.01 | 0.49 | 0.01 | 2.41 |
| 100 | 200 | -9.73 | -7.84 | -9.59 | -4.65 | 0.01 | 0.83 | 0.01 | 4.14 |
| 200 | 25 | -3.30 | -1.80 | -3.22 | 0.33 | 0.01 | 0.56 | 0.01 | 1.97 |
| 200 | 50 | -4.77 | -2.88 | -4.68 | 0.07 | 0.01 | 0.60 | 0.01 | 2.68 |
| 200 | 100 | -6.82 | -4.28 | -6.71 | -0.12 | 0.01 | 0.94 | 0.01 | 4.25 |
| 200 | 200 | -9.70 | -6.22 | -9.56 | -0.43 | 0.01 | 1.48 | 0.01 | 7.28 |

Notes. The model has a factor error structure with 3 factors. $\varepsilon_{i,t}$ and $e_{i,t}$ are standardized $\chi^2(2)$ with zero expected value and unit variance. " $\sigma_i^2: \perp \lambda_i$ " means that $\sigma_{i,y}^2 = (\zeta_{i,y}^2 - 2) / 4 + 1$ where $\zeta_{i,y}^2$ is $\chi^2(2)$. For " $\sigma_i^2: f(\lambda_i)$ ", we let $\sigma_{i,y}^2 = d_\sigma T^{-1} \sum_{t=1}^T (\lambda_i' \mathbf{f}_t)^2$. The case " λ_i : symmetric" corresponds to drawing λ_i from $U(0.5, 1.5)$. " λ_i : skewed" loadings are from a standardized $\chi^2(2)$ distribution with mean and variance equal to 1. Λ_i has first element from $U(0.5, 1.5)$ and all others from $U(-0.5, 0.5)$. Factors \mathbf{f}_t are drawn from $N(0, 1)$.

Table S.3: Sample moments of original CD test statistic applied to CCE residuals under \mathbb{H}_1

| N | λ_i : σ_i^2 : T | Mean | | | | Variance | | | |
|-----|--------------------------------------|-------------------|----------------|-------------------|----------------|-------------------|----------------|-------------------|----------------|
| | | symmetric | | skewed | | symmetric | | skewed | |
| | | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ | $\perp \lambda_i$ | $f(\lambda_i)$ |
| 25 | 25 | -3.27 | -3.21 | -3.25 | -3.01 | 0.03 | 0.04 | 0.03 | 0.19 |
| 25 | 50 | -4.78 | -4.72 | -4.76 | -4.42 | 0.02 | 0.03 | 0.02 | 0.25 |
| 25 | 100 | -6.87 | -6.80 | -6.86 | -6.40 | 0.02 | 0.04 | 0.02 | 0.41 |
| 25 | 200 | -9.80 | -9.70 | -9.78 | -9.16 | 0.03 | 0.06 | 0.02 | 0.68 |
| 50 | 25 | -3.21 | -3.14 | -3.19 | -2.93 | 0.02 | 0.04 | 0.02 | 0.15 |
| 50 | 50 | -4.71 | -4.65 | -4.70 | -4.37 | 0.02 | 0.02 | 0.01 | 0.17 |
| 50 | 100 | -6.78 | -6.70 | -6.78 | -6.32 | 0.01 | 0.03 | 0.01 | 0.26 |
| 50 | 200 | -9.68 | -9.58 | -9.67 | -9.04 | 0.02 | 0.04 | 0.01 | 0.42 |
| 100 | 25 | -3.16 | -3.10 | -3.14 | -2.90 | 0.02 | 0.03 | 0.02 | 0.10 |
| 100 | 50 | -4.67 | -4.61 | -4.66 | -4.34 | 0.01 | 0.02 | 0.01 | 0.11 |
| 100 | 100 | -6.73 | -6.66 | -6.73 | -6.28 | 0.01 | 0.02 | 0.01 | 0.16 |
| 100 | 200 | -9.62 | -9.52 | -9.61 | -8.99 | 0.01 | 0.03 | 0.01 | 0.27 |
| 200 | 25 | -3.11 | -3.05 | -3.09 | -2.84 | 0.02 | 0.03 | 0.02 | 0.09 |
| 200 | 50 | -4.66 | -4.58 | -4.64 | -4.31 | 0.01 | 0.02 | 0.01 | 0.09 |
| 200 | 100 | -6.71 | -6.64 | -6.71 | -6.26 | 0.01 | 0.02 | 0.01 | 0.11 |
| 200 | 200 | -9.59 | -9.48 | -9.58 | -8.96 | 0.01 | 0.02 | 0.01 | 0.21 |

Notes. See Table S.2.

Table S.4: Rejection rates for weighted CD test statistic when applied to 2WFE residuals

Part A: Size

| N | $\lambda_i:$ $\sigma_i^2:$ T | symmetric | | | | | | skewed | | | | | |
|-----|------------------------------------|-------------------|-----------|-----------|----------------|-----------|-----------|-------------------|-----------|-----------|----------------|-----------|-----------|
| | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | |
| | | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} |
| 25 | 25 | 4.9 | 5.4 | 2.5 | 5.1 | 5.6 | 2.9 | 5.0 | 5.9 | 2.5 | 6.1 | 6.6 | 2.7 |
| 25 | 50 | 6.5 | 7.0 | 3.9 | 6.8 | 7.4 | 3.9 | 5.7 | 6.5 | 3.5 | 6.2 | 6.8 | 3.3 |
| 25 | 100 | 8.8 | 9.0 | 5.0 | 8.0 | 8.2 | 4.6 | 8.4 | 8.8 | 5.1 | 8.0 | 8.1 | 3.9 |
| 25 | 200 | 10.8 | 10.9 | 5.7 | 9.7 | 9.8 | 6.1 | 11.0 | 11.0 | 7.0 | 10.9 | 11.1 | 4.3 |
| 50 | 25 | 5.1 | 5.9 | 2.8 | 5.7 | 6.2 | 3.6 | 5.7 | 6.7 | 2.3 | 5.1 | 5.7 | 2.9 |
| 50 | 50 | 5.0 | 6.6 | 4.3 | 4.6 | 5.9 | 4.0 | 6.5 | 7.6 | 3.4 | 5.4 | 6.8 | 3.1 |
| 50 | 100 | 6.6 | 7.1 | 3.9 | 5.9 | 6.7 | 3.2 | 6.6 | 7.0 | 4.6 | 5.9 | 6.5 | 3.5 |
| 50 | 200 | 6.6 | 6.8 | 4.7 | 5.7 | 5.9 | 4.4 | 5.8 | 6.2 | 4.4 | 6.1 | 6.3 | 3.5 |
| 100 | 25 | 5.3 | 5.8 | 2.8 | 5.3 | 5.5 | 3.1 | 4.6 | 4.7 | 3.5 | 5.0 | 5.4 | 3.7 |
| 100 | 50 | 5.9 | 8.7 | 5.1 | 4.5 | 6.7 | 4.1 | 4.6 | 7.4 | 3.9 | 4.8 | 7.4 | 4.0 |
| 100 | 100 | 5.6 | 6.6 | 4.5 | 5.2 | 7.0 | 4.3 | 4.9 | 6.1 | 5.0 | 4.2 | 5.9 | 2.8 |
| 100 | 200 | 5.8 | 6.3 | 4.0 | 5.5 | 6.2 | 4.8 | 5.5 | 5.9 | 3.1 | 5.9 | 6.3 | 4.1 |
| 200 | 25 | 4.4 | 4.5 | 2.9 | 4.8 | 4.8 | 3.5 | 5.0 | 5.1 | 3.2 | 4.8 | 4.9 | 3.1 |
| 200 | 50 | 4.6 | 7.4 | 3.6 | 4.8 | 8.6 | 3.8 | 4.5 | 8.0 | 3.7 | 5.8 | 8.7 | 4.0 |
| 200 | 100 | 4.7 | 8.3 | 4.2 | 5.5 | 9.7 | 4.7 | 5.2 | 8.6 | 4.1 | 5.7 | 9.4 | 4.1 |
| 200 | 200 | 4.2 | 5.4 | 5.0 | 5.4 | 7.4 | 5.0 | 5.9 | 6.8 | 5.2 | 4.9 | 6.6 | 3.6 |

Part B: Power

| N | $\lambda_i:$ $\sigma_i^2:$ T | symmetric | | | | | | skewed | | | | | |
|-----|------------------------------------|-------------------|-----------|-----------|----------------|-----------|-----------|-------------------|-----------|-----------|----------------|-----------|-----------|
| | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | |
| | | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} |
| 25 | 25 | 12.6 | 26.6 | 8.0 | 10.7 | 14.0 | 5.8 | 12.3 | 19.6 | 14.7 | 14.3 | 20.0 | 24.0 |
| 25 | 50 | 21.2 | 87.4 | 14.8 | 17.5 | 46.1 | 12.7 | 18.0 | 59.6 | 32.9 | 23.2 | 63.7 | 48.4 |
| 25 | 100 | 34.6 | 99.9 | 26.4 | 27.3 | 95.2 | 20.9 | 32.3 | 90.8 | 50.1 | 36.3 | 96.7 | 66.6 |
| 25 | 200 | 50.4 | 100 | 39.7 | 43.6 | 100 | 34.7 | 44.1 | 99.3 | 64.4 | 51.3 | 100 | 80.4 |
| 50 | 25 | 11.7 | 21.1 | 8.1 | 10.7 | 11.7 | 5.7 | 11.7 | 20.9 | 35.2 | 12.6 | 16.0 | 58.9 |
| 50 | 50 | 19.6 | 98.3 | 15.7 | 17.5 | 56.0 | 12.8 | 18.7 | 85.5 | 59.5 | 23.0 | 79.7 | 86.0 |
| 50 | 100 | 35.4 | 100 | 27.0 | 26.3 | 99.7 | 22.2 | 29.7 | 99.8 | 80.2 | 37.5 | 100 | 95.4 |
| 50 | 200 | 49.3 | 100 | 40.6 | 43.5 | 100 | 37.4 | 45.3 | 100 | 90.7 | 53.2 | 100 | 99.1 |
| 100 | 25 | 11.1 | 14.4 | 6.5 | 10.6 | 10.9 | 5.1 | 10.4 | 19.1 | 73.4 | 12.0 | 12.9 | 93.5 |
| 100 | 50 | 19.9 | 99.9 | 15.0 | 16.3 | 66.4 | 13.3 | 18.1 | 98.9 | 92.9 | 22.3 | 94.6 | 99.7 |
| 100 | 100 | 33.3 | 100 | 25.8 | 26.2 | 100 | 23.8 | 30.8 | 100 | 98.8 | 36.7 | 100 | 100 |
| 100 | 200 | 48.7 | 100 | 41.3 | 41.9 | 100 | 37.0 | 46.2 | 100 | 100 | 52.3 | 100 | 100 |
| 200 | 25 | 12.7 | 12.7 | 8.2 | 10.8 | 10.8 | 6.6 | 11.9 | 13.9 | 96.5 | 12.9 | 13.0 | 100 |
| 200 | 50 | 18.9 | 100 | 16.6 | 17.9 | 73.1 | 13.3 | 18.8 | 100 | 100 | 22.1 | 98.2 | 100 |
| 200 | 100 | 33.9 | 100 | 27.4 | 26.3 | 100 | 21.5 | 32.8 | 100 | 100 | 36.9 | 100 | 100 |
| 200 | 200 | 50.9 | 100 | 41.9 | 43.6 | 100 | 37.2 | 46.7 | 100 | 100 | 54.6 | 100 | 100 |

Notes. In Part A the model has two factors which are restricted as noted in Table 1. Part B corresponds to a model with factor error structure and 3 factors. For details on all other model parameters, see Table S.2.

CD_W is the weighted CD test statistic introduced in Theorem 3. CD_{W+} is its power-enhanced refinement. CD_{BC} is a CD test statistic with analytic bias correction.

Table S.5: Rejection rates for weighted CD test statistic when applied to CCE residuals

Part A: Size

| N | λ_i : σ_i^2 : T | symmetric | | | | | | skewed | | | | | |
|-----|--------------------------------------|-------------------|-----------|-----------|----------------|-----------|-----------|-------------------|-----------|-----------|----------------|-----------|-----------|
| | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | |
| | | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} |
| 25 | 25 | 6.2 | 6.8 | 4.6 | 5.7 | 6.4 | 4.4 | 5.6 | 6.1 | 4.7 | 5.6 | 6.6 | 5.9 |
| 25 | 50 | 7.4 | 8.1 | 5.7 | 5.8 | 6.6 | 5.9 | 7.0 | 7.9 | 6.1 | 6.7 | 7.3 | 8.9 |
| 25 | 100 | 8.3 | 8.5 | 6.6 | 8.0 | 7.9 | 7.8 | 9.0 | 9.7 | 6.4 | 8.2 | 8.8 | 14.5 |
| 25 | 200 | 9.2 | 10.1 | 7.4 | 11.3 | 11.6 | 8.8 | 10.9 | 12.2 | 9.5 | 13.0 | 14.0 | 24.2 |
| 50 | 25 | 5.1 | 5.8 | 4.1 | 4.4 | 5.0 | 4.8 | 5.9 | 6.6 | 4.9 | 5.5 | 6.7 | 8.0 |
| 50 | 50 | 5.1 | 6.5 | 5.1 | 6.1 | 7.0 | 4.2 | 5.5 | 7.0 | 5.8 | 7.2 | 8.0 | 8.1 |
| 50 | 100 | 5.5 | 6.0 | 4.7 | 5.2 | 5.5 | 6.3 | 6.3 | 6.6 | 4.7 | 6.6 | 7.2 | 11.8 |
| 50 | 200 | 6.4 | 6.5 | 5.9 | 6.9 | 6.9 | 6.7 | 6.0 | 6.6 | 6.9 | 7.4 | 7.9 | 21.9 |
| 100 | 25 | 5.2 | 5.4 | 5.9 | 5.5 | 5.7 | 6.7 | 5.4 | 5.8 | 6.6 | 5.5 | 5.7 | 8.9 |
| 100 | 50 | 5.0 | 7.4 | 4.3 | 5.3 | 8.3 | 5.9 | 5.3 | 7.0 | 4.5 | 5.1 | 6.9 | 8.0 |
| 100 | 100 | 5.5 | 6.6 | 5.1 | 5.1 | 7.0 | 5.2 | 5.8 | 7.0 | 5.4 | 5.1 | 6.9 | 11.3 |
| 100 | 200 | 5.0 | 5.5 | 4.8 | 6.0 | 6.2 | 5.1 | 5.2 | 5.9 | 5.7 | 5.8 | 6.1 | 19.2 |
| 200 | 25 | 5.8 | 5.8 | 11.7 | 5.1 | 5.1 | 12.7 | 6.3 | 6.3 | 11.7 | 4.9 | 5.0 | 16.3 |
| 200 | 50 | 6.3 | 9.4 | 4.5 | 5.6 | 9.7 | 5.3 | 5.0 | 9.1 | 5.0 | 5.6 | 9.8 | 9.5 |
| 200 | 100 | 5.4 | 8.5 | 5.5 | 5.1 | 8.3 | 5.8 | 4.9 | 8.3 | 4.9 | 5.1 | 8.4 | 10.8 |
| 200 | 200 | 5.9 | 7.0 | 5.3 | 4.8 | 5.9 | 5.9 | 4.8 | 6.7 | 4.7 | 5.0 | 7.1 | 15.8 |

Part B: Power

| N | λ_i : σ_i^2 : T | symmetric | | | | | | skewed | | | | | |
|-----|--------------------------------------|-------------------|-----------|-----------|----------------|-----------|-----------|-------------------|-----------|-----------|----------------|-----------|-----------|
| | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | | $\perp \lambda_i$ | | | $f(\lambda_i)$ | | |
| | | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} | CD_W | CD_{W+} | CD_{BC} |
| 25 | 25 | 10.4 | 16.3 | 5.7 | 9.2 | 10.7 | 5.6 | 8.5 | 13.4 | 6.2 | 9.7 | 12.8 | 7.1 |
| 25 | 50 | 15.0 | 47.6 | 9.9 | 10.2 | 19.8 | 8.9 | 12.5 | 37.4 | 11.8 | 15.2 | 31.4 | 14.5 |
| 25 | 100 | 22.6 | 83.9 | 14.0 | 17.6 | 50.6 | 13.2 | 20.5 | 64.4 | 18.7 | 23.3 | 67.0 | 24.8 |
| 25 | 200 | 33.6 | 97.6 | 22.5 | 26.0 | 85.7 | 22.5 | 28.2 | 86.6 | 28.8 | 34.4 | 91.1 | 42.4 |
| 50 | 25 | 9.7 | 12.6 | 5.7 | 7.8 | 8.5 | 4.9 | 8.7 | 15.0 | 5.9 | 8.6 | 10.3 | 7.4 |
| 50 | 50 | 11.5 | 67.0 | 8.6 | 11.5 | 23.5 | 8.2 | 10.9 | 58.8 | 9.4 | 13.5 | 42.7 | 14.9 |
| 50 | 100 | 20.6 | 98.4 | 14.7 | 14.7 | 68.5 | 13.9 | 18.1 | 91.6 | 21.6 | 21.2 | 90.4 | 28.1 |
| 50 | 200 | 28.0 | 100 | 23.5 | 24.3 | 98.6 | 22.9 | 26.6 | 99.0 | 32.1 | 33.2 | 99.6 | 43.0 |
| 100 | 25 | 9.3 | 9.9 | 4.8 | 7.5 | 7.8 | 5.5 | 9.0 | 13.0 | 6.2 | 9.0 | 9.3 | 8.5 |
| 100 | 50 | 12.6 | 80.4 | 6.4 | 9.3 | 25.4 | 7.6 | 12.2 | 82.9 | 11.9 | 13.1 | 54.9 | 15.9 |
| 100 | 100 | 18.2 | 100 | 12.5 | 15.2 | 85.8 | 12.8 | 17.6 | 99.7 | 21.8 | 22.2 | 98.9 | 29.5 |
| 100 | 200 | 29.4 | 100 | 21.9 | 23.6 | 99.9 | 24.3 | 28.6 | 100 | 36.2 | 29.7 | 100 | 46.7 |
| 200 | 25 | 9.4 | 9.7 | 7.9 | 8.1 | 8.1 | 6.6 | 7.5 | 8.1 | 7.6 | 10.7 | 10.8 | 8.6 |
| 200 | 50 | 12.1 | 90.8 | 9.4 | 10.9 | 33.4 | 7.2 | 11.9 | 95.2 | 11.2 | 12.8 | 64.5 | 17.2 |
| 200 | 100 | 20.0 | 100 | 13.0 | 14.8 | 96.3 | 13.0 | 16.6 | 100 | 21.8 | 21.1 | 99.8 | 29.8 |
| 200 | 200 | 27.2 | 100 | 22.6 | 23.5 | 100 | 21.7 | 26.6 | 100 | 42.4 | 31.0 | 100 | 49.0 |

Notes. In Part A the model has two factors which are restricted as noted in Table S.1. Part B corresponds to a model with factor error structure and 3 factors. For details on all other model parameters, see Table S.2.

CD_W is the weighted CD test statistic introduced in Theorem 3. CD_{W+} is its power-enhanced refinement. CD_{BC} is a CD test statistic with analytic bias correction.

References

- ABADIR, K. M. AND J. R. MAGNUS (2005): *Matrix algebra*, vol. 1 of *Econometric Exercises*, Cambridge University Press.
- BAI, J. (2003): "Inferential Theory for Factor Models of Large Dimensions," *Econometrica*, 71, 135–171.
- BONHOMME, S. AND E. MANRESA (2015): "Grouped Patterns of Heterogeneity in Panel Data," *Econometrica*, 83, 1147–1184.
- BALTAGI, B., C. KAO, AND B. PENG (2016): "Testing Cross-Sectional Correlation in Large Panel Data Models with Serial Correlation," *Econometrics*, 4, –.
- CHEN, S. X. AND Y.-L. QIN (2010): "A Two-sample Test for High-dimensional Data with Applications to Gene-set Testing," *The Annals of Statistics*, 38, 808–835.
- HALL, P. AND C. C. HEYDE (1980): *Martingale Limit Theory and Its Application*, Probability and Mathematical Statistics, Academic Press.
- JUODIS, A., H. KARABIYIK, AND J. WESTERLUND (2017): "On the Robustness of the Pooled CCE Estimator," Mimeo.
- KAO, C., L. TRAPANI, AND G. URGAS (2012): "Asymptotics for Panel Models with Common Shocks," *Econometric Reviews*, 31, 390–439.
- KARABIYIK, H., S. REESE, AND J. WESTERLUND (2017): "On The Role of The Rank Condition in CCE Estimation of Factor-augmented Panel Regressions," *Journal of Econometrics*, 197, 60–64.
- PESARAN, M. H. (2004): "General Diagnostic Tests for Cross Section Dependence in Panels," CESifo Working Paper No. 1229.
- (2006): "Estimation and Inference in Large Heterogeneous Panels with a Multifactor Error Structure," *Econometrica*, 74, 967–1012.
- (2015): "Testing Weak Cross-Sectional Dependence in Large Panels," *Econometric Reviews*, 34, 1089–1117.