

SOME RESULTS ON POLISH GROUPS

GIANLUCA PAOLINI AND SAHARON SHELAH

ABSTRACT. We prove that no quantifier-free formula in the language of group theory can define the \aleph_1 -half graph in a Polish group, thus generalising some results from [6]. We then pose some questions on the space of groups of automorphisms of a given Borel complete class, and observe that this space must contain at least one uncountable group. Finally, we prove some results on the structure of the group of automorphisms of a locally finite group: firstly, we prove that it is not the case that every group of automorphisms of a graph of power λ is the group of automorphism of a locally finite group of power λ ; secondly, we conjecture that the group of automorphisms of a locally finite group of power λ has a locally finite subgroup of power λ , and reduce the problem to a problem on p -groups, thus settling the conjecture in the case $\lambda = \aleph_0$.

1. INTRODUCTION

We collect some results (of different nature) on the theory of Polish groups.

Section 2. Definable \aleph_1 -Half Graphs in Polish Groups.

By the \aleph_1 -half graph $\Gamma(\aleph_1)$ we mean the graph on vertex set $\{a_\alpha : \alpha < \aleph_1\} \cup \{b_\beta : \beta < \aleph_1\}$ with edge relation $a_\alpha E_\Gamma b_\beta$ if and only if $\alpha < \beta$. In the process of characterization of the graph products of cyclic groups embeddable in a Polish group [6], we observed that the commutation relation $x^{-1}y^{-1}xy = e$ can never define the \aleph_1 -half graph in a Polish group G . Here we generalize this to:

Theorem 1.1. *No quantifier-free formula $\varphi(\bar{x}, \bar{y})$ in the language of group theory can define the \aleph_1 -half graph in a Polish group G .*

We actually prove a stronger result of independent interest, i.e. that Polish groups do not admit “polarized \aleph_1 -partitions”, see Theorem 2.1 for the detailed statement of this result. Finally, we would like to mention that Theorem 1.1 can be considered as a form of model-theoretic stability for Polish groups, in this direction see also [9].

Section 3. The Space of Automorphism Groups of a Borel Complete Class.

By a Borel complete class we mean a Borel class \mathbf{K} of structures with domain ω in a fixed language L such that the isomorphism relation on \mathbf{K} is as complicated as possible (equivalently, the countable graph isomorphism relation is reducible to it – cf. Definition 3.2). We wonder here: how complex can $\text{Aut}(\mathbf{K}) = \{\text{Aut}(A) : A \in \mathbf{K}\}$ be for a given Borel complete class? Can $\text{Aut}(\mathbf{K})$ contain only one isomorphism type, resp. finitely many, resp. countably many (cf. Problem 3.5)? In this direction:

Date: March 17, 2022.

Partially supported by European Research Council grant 338821. No. 1155 on Shelah’s publication list. The first author would like to thank Riccardo Camerlo for useful discussions related to the questions posed in Section 3.

Proposition 1.2. *Let \mathbf{K} be a Borel class of L -structures with domain \aleph_0 such that for every $G \in \text{Aut}(\mathbf{K})$ we have that $|G| \leq \aleph_0$. Then the isomorphism relation on \mathbf{K} is Borel, and so in particular \mathbf{K} is not Borel complete (cf. Definition 3.2).*

On questions affine to this topic see also the interesting recent work [5].

Section 4. Group of Automorphisms of Locally Finite Groups.

Notation 1.3. (1) We denote by \mathbf{K}_{lf} the class of locally finite groups.

(2) We denote by \mathbf{K}_{gf} the class of graphs.

(3) For \mathbf{K} a class of L -structures and λ an infinite cardinal, we let:

(3.1) $\mathbf{K}^\lambda = \{M \in \mathbf{K} : \text{the domain of } M \text{ is } \lambda\};$

(3.2) $\text{Aut}(\mathbf{K}) = \{\text{Aut}(M) : M \in \mathbf{K}\}.$

Fact 1.4 ([4]). *The class $\mathbf{K}_{\text{lf}}^{\aleph_0}$ is Borel complete (cf. Definition 3.2).*

In this section we deal with the following problem:

Problem 1.5. *Characterize $\text{Aut}(\mathbf{K}_{\text{lf}}^\lambda)$, for $\lambda \geq \aleph_0$.*

In this direction we first prove:

Theorem 1.6. *Let $\lambda \geq \aleph_0$, then:*

$$\text{Aut}(\mathbf{K}_{\text{gf}}^\lambda) \neq \text{Aut}(\mathbf{K}_{\text{lf}}^\lambda).$$

The proof of Theorem 1.6 leads to the following conjecture:

Conjecture 1.7. *If $G \in \mathbf{K}_{\text{lf}}^\lambda$, then $\text{Aut}(G)$ has a locally finite subgroup of power λ .*

In the case of \aleph_0 we prove that this is indeed the case:

Proposition 1.8. *If $G \in \mathbf{K}_{\text{lf}}^{\aleph_0}$, then $\text{Aut}(G)$ has a locally finite infinite subgroup.*

On the other hand, we do not settle here Conjecture 1.7 in general, but we prove:

Lemma 1.9. *To prove Conjecture 1.7 it suffices to prove Conjecture 1.10, where:*

Conjecture 1.10. *If $\lambda > \aleph_0$, $G \in \mathbf{K}_{\text{lf}}^\lambda$ is an abelian p -group, $H \leq G$ and $|H| < \lambda$, then $\text{Aut}_H(G)$ has a locally finite subgroup of power λ .*

Finally, we would like to mention that in [7] we give a close analysis of the group of automorphisms of Philip Hall's universal locally finite group.

2. DEFINABLE \aleph_1 -HALF GRAPHS IN POLISH GROUPS

Theorem 2.1. *Let G be a Polish group, and for $\ell < \ell(*) < \omega$ let:*

- (i) $\bar{g}^\ell = (\bar{g}_\alpha^\ell : \alpha \in A_\ell);$
- (ii) $A_\ell \in [\omega_1]^{\aleph_1};$
- (iii) $\bar{g}_\alpha^\ell \in G^{n(\ell)};$
- (iv) Δ be a finite set of q.f. formulas of the form $\varphi(\bar{x}_{n(0)}^0, \dots, \bar{x}_{n(\ell(*)-1)}^{\ell(*)-1})$ in the language of group theory such that $\text{lg}(\bar{x}_{n(\ell)}^\ell) = n(\ell).$

Then there are $B_\ell \in [A_\ell]^{\aleph_1}$, for $\ell < \ell(*)$, and truth value $t \in \{0, 1\}$ such that if $\alpha(\ell) \in B_\ell$, for $\ell < \ell(*)$, then:

$$G \models \varphi^t(\bar{g}_{\alpha(0)}^0, \dots, \bar{x}_{\alpha(\ell(*)-1)}^{\ell(*)-1}).$$

Proof. First of all notice that it suffices to prove the claim for:

$$\Delta = \{\sigma(\bar{x}_{n(0)}^0, \dots, \bar{x}_{n(\ell(*)-1)}^{\ell(*)-1}) = e\},$$

and $\sigma(\bar{x}_{n(0)}^0, \dots, \bar{x}_{n(\ell(*)-1)}^{\ell(*)-1})$ a term in the language of group theory $L = \{e, \cdot, ()^{-1}\}$. [Why? First of all, without loss of generality, we can assume that each $\varphi \in \Delta$ is a Boolean combination of formulas of the form $\sigma(\bar{x}_{n(0)}^0, \dots, \bar{x}_{n(\ell(*)-1)}^{\ell(*)-1}) = e$. So let $(\sigma_i(\bar{x}_{n(0)}^0, \dots, \bar{x}_{n(\ell(*)-1)}^{\ell(*)-1}) = e : i < i(*) < \omega)$ list them. Now choose $(B_{i,\ell} : \ell < \ell(*))$ by induction on $i \leq i*$ such that:

- (a) $B_{i,\ell} \in [\omega_1]^{\aleph_1}$;
- (b) $B_{0,\ell} = A_\ell$;
- (c) $B_{i+1,\ell} \subseteq B_{i,\ell}$;
- (d) $(B_{i+1,\ell} : \ell < \ell(*))$ satisfies the desired conclusion for:

$$\Delta = \{\sigma_i(\bar{x}_{n(0)}^0, \dots, \bar{x}_{n(\ell(*)-1)}^{\ell(*)-1}) = e\}.$$

Then $(B_{i(*),\ell} : \ell < \ell(*))$ is as wanted.]

Let (G, d) witness the Polishness of G . For $\ell < \ell(*)$ and $\alpha, \beta \in A_\ell$, let $d(\bar{g}_\alpha^\ell, \bar{g}_\beta^\ell) = \max\{d(g_\alpha^{\ell,i}, g_\beta^{\ell,i}) : i < n(\ell)\}$, and:

$$\mathcal{U}_\ell = \{\alpha < \omega_1 : \text{for some } \varepsilon \in (0, 1)_\mathbb{R} \text{ the set } \{\beta \in A_\ell : d(\bar{g}_\alpha^\ell, \bar{g}_\beta^\ell) < \varepsilon\} \text{ is countable}\}.$$

Since (G, d) is separable, for every $\ell < \ell(*)$, the set \mathcal{U}_ℓ is countable, and so we can find $\alpha(*) < \omega_1$ such that $\bigcup_{\ell < \ell(*)} \mathcal{U}_\ell \subseteq \alpha(*)$. Now, if $B_\ell = A_\ell - \alpha(*)$ is such that for every $\alpha(\ell) \in B_\ell$ we have that $G \models \sigma(\bar{g}_{\alpha(0)}^0, \dots, \bar{g}_{\alpha(\ell(*)-1)}^{\ell(*)-1}) = e$, then we are done. So suppose that this is not the case, then we can find $\alpha_\ell \in A_\ell - \alpha(*)$ such that:

$$\varepsilon = d(\sigma(\bar{g}_{\alpha(0)}^0, \dots, \bar{g}_{\alpha(\ell(*)-1)}^{\ell(*)-1}), e) \neq 0.$$

As G is Polish, there is $\xi \in (0, 1)_\mathbb{R}$ such that:

$$(1) \quad \text{if } \bar{a}_\ell \in G^{n(\ell)} \text{ and } d(\bar{a}_\ell, \bar{g}_{\alpha_\ell}^\ell), \text{ then } d(\sigma(\bar{a}_0^0, \dots, \bar{a}_{\ell(*)-1}^{\ell(*)-1}), e) > \varepsilon/2.$$

Now, for $\ell < \ell(*)$, let $B_\ell = \{\alpha \in A_\ell : d(\bar{g}_\alpha^\ell, \bar{g}_{\alpha_\ell}^\ell) < \xi\}$. Then $B_\ell \subseteq A_\ell$ and, as $\alpha(*) \leq \alpha_\ell$, clearly $|B_\ell| = \alpha_1$. Hence, by (1), $(B_\ell : \ell < \ell(*))$ is as required. ■

Theorem 1.1. *No quantifier-free formula $\varphi(\bar{x}, \bar{y})$ in the language of group theory can define the \aleph_1 -half graph in a Polish group G .*

Proof. Immediate from Theorem 2.1. ■

3. THE SPACE OF AUTOMORPHISM GROUPS OF A BOREL COMPLETE CLASS

For an overview (and careful explanation) of the descriptive set theoretic notions occurring in this section cf. e.g. [8, Chapter 11].

Notation 3.1. We denote by $\mathbf{K}_{\text{gf}}^{\aleph_0}$ the standard Borel space of graphs with domain \aleph_0 .

Definition 3.2. Let \mathbf{K} be a Borel class of L -structures with domain \aleph_0 . We say that \mathbf{K} is Borel complete if there exists a Borel map $f : \mathbf{K}_{\text{gf}}^{\aleph_0} \rightarrow \mathbf{K}$ such that for every $A, B \in \mathbf{K}_0$ we have $A \cong B$ if and only if $f(A) \cong f(B)$.

Notation 3.3. Let G and H be topological groups.

- (1) We write $G \cong H$ to mean that G and H are isomorphic as abstract groups.
- (2) We write $G \cong_t H$ to mean that G and H are isomorphic as topological groups.

Notation 3.4. (As in Notation 1.3) Given a class \mathbf{K} of L -structures, we let:

$$\text{Aut}(\mathbf{K}) = \{\text{Aut}(A) : A \in \mathbf{K}\}.$$

Proposition 1.2. Let \mathbf{K} be a Borel class of L -structures with domain \aleph_0 such that for every $G \in \text{Aut}(\mathbf{K})$ we have that $|G| \leq \aleph_0$. Then the isomorphism relation on \mathbf{K} is Borel, and so in particular \mathbf{K} is not Borel complete.

Proof. We show that for any such class \mathbf{K} the isomorphism relation \cong on \mathbf{K} is Borel. Notice that for $A, B \in \mathbf{K}$ we have that $A \cong B$ if and only if there are countably many $f \in S_\infty := \{f : \omega \rightarrow \omega : f \text{ is a bijection}\}$ such that $f : A \cong B$. Thus, the relation \cong on \mathbf{K} is the projection of a Borel relation R :

$$(\mathbf{K} \times \mathbf{K}) \times S_\infty \supseteq R = \{(A, B, f) : f : A \cong B\}$$

with countable sections $R_{(A,B)}$ (for $(A, B) \in \mathbf{K} \times \mathbf{K}$). Hence, by [3, Lemma 18.12], the relation \cong on \mathbf{K} is Borel, and so we are done. \blacksquare

We are interested in the following open problem:

Problem 3.5. Let \mathbf{K} be a Borel complete class.

- (1) Can $\text{Aut}(\mathbf{K})/\cong$ have size 1? Can $\text{Aut}(\mathbf{K})/\cong_t$ have size 1?
- (2) Can $\text{Aut}(\mathbf{K})/\cong$ be finite? Can $\text{Aut}(\mathbf{K})/\cong_t$ be finite?
- (3) Can $\text{Aut}(\mathbf{K})/\cong$ be countable? Can $\text{Aut}(\mathbf{K})/\cong_t$ be countable?

4. GROUPS OF AUTOMORPHISMS OF LOCALLY FINITE GROUPS

Definition 4.1. (1) We denote by P the set of prime numbers.

(2) For $p \in P$, we denote by $G_{(p,\infty)}^*$ the divisible abelian p -group of rank 1.

(3) For $p \in P$ and $\ell < \omega$ we denote by $G_{(p,\ell)}^*$ the finite cyclic group of order p^ℓ .

(4) For $p \in P$, we let $S_p = \{(p, \ell) : \ell < \omega\}$ and $S_p^+ = S_p \cup \{(p, \infty)\}$.

(5) For $s \in S_p^+$ and λ a cardinal, we let $G_{s,\lambda}^*$ be the direct sum of λ copies of G_s^* .

(6) For $p \in P$, we denote by J_p the group of p -adic integers.

(7) We say that an abelian group G is bounded if there exists $n < \omega$ such that for every $g \in G$ we have $ng = 0$.

(8) We say that G is unbounded if it is not bounded.

(9) We say that G is torsion if every element of G has finite order.

Fact 4.2 ([1][Theorem 17.2]). Let G be a bounded abelian group. Then G is a direct sum of cyclic groups.

Fact 4.3. If an abelian p -group G is bounded, then there exists $n < \omega$ such that:

$$G = \bigoplus_{\ell < n} G_{(p,\ell),\lambda_\ell}^*.$$

Proof. This is a consequence of Fact 4.2. \blacksquare

Fact 4.4 ([1][Theorem 8.4]). Let G be a torsion abelian group. Then:

$$G = \bigoplus_{p \in P} G_p,$$

with G_p a p -group, for every $p \in P$.

Remark 4.5 ([2][pg. 250]). *Let G be an abelian group and suppose that $G = \bigoplus_{i \in I} G_i$, then $\bigoplus_{i \in I} \text{Aut}(G_i)$ can be embedded into $\text{Aut}(G)$.*

Fact 4.6 ([2, Theorem 115.1]). *Let G be an unbounded abelian p -group, then there exists an embedding $f : J_p \rightarrow \text{Aut}(G)$.*

Lemma 4.7. *Let $G \in \mathbf{K}_{\text{lf}}^\lambda$ (cf. Definition 1.3). Then $\text{Aut}(G)$ has a non-trivial locally finite subgroup.*

Proof. We distinguish three cases:

- (i) G is not abelian.
- (ii) G is abelian and not bounded.
- (iii) G is abelian and bounded.

If (i), then $G/\text{Cent}(G) \in \mathbf{K}_{\text{lf}}$ is non-trivial and it can be embedded into $\text{Aut}(G)$, and so we are done. If (ii), then we are done by Facts 4.4 and 4.6, and Remark 4.5. Finally, if (iii), then by Facts 4.3 and 4.4, there exists a direct summand G_p of G such that $G_p = \bigoplus_{\ell < n} G_{(p, \ell), \lambda_\ell}^*$ and for some $0 < \ell(*) < n < \omega$ we have $\lambda_{\ell(*)} \geq \aleph_0$. Let $G_{p^\ell}(\alpha)$ be the α -th copy of $G_{(p, \ell), \lambda_\ell}^*$. If $p > 2$, then consider:

$$\{\pi \in \text{Aut}(G_p) : \pi \text{ maps } G_{p^\ell}(\alpha) \text{ onto itself, for every } \alpha < \lambda_\ell \text{ and } \ell < n\}.$$

If $p = 2$, then consider:

$$\{\pi \in \text{Aut}(G_p) : \pi \text{ maps } G_{p^{\ell(*)}}(2\alpha) \oplus G_{p^{\ell(*)}}(2\alpha + 1) \text{ onto itself, for every } \alpha < \lambda_{\ell(*)}\}.$$

Hence, by Remark 4.5, also (iii) is taken care of. \blacksquare

The following facts are folklore:

Fact 4.8. (1) *If M is a linear order, then $\text{Aut}(M)$ has no element of finite order.*
 (2) *For every infinite structure M there exists a graph Γ_M of the same cardinality of M such that $\text{Aut}(\Gamma_M) \cong \text{Aut}(M)$.*

Theorem 1.6. *Let $\lambda \geq \aleph_0$, then:*

$$\text{Aut}(\mathbf{K}_{\text{gf}}^\lambda) \neq \text{Aut}(\mathbf{K}_{\text{lf}}^\lambda).$$

Proof. By Lemma 4.7 and Fact 4.8. \blacksquare

We devote the rest of the section to the proof of Proposition 1.8 and Lemma 1.9.

Notation 4.9. *Given a group G and $H \leq G$ we let:*

$$\text{Aut}_H(G) = \{\pi \in \text{Aut}(G) : \pi \upharpoonright H = \text{id}_H\}.$$

Fact 4.10. *If $G \in \mathbf{K}_{\text{lf}}^\lambda$ and $|G/\text{Cent}(G)| < \lambda$, then there is $H \leq G$ such that $|H| < \lambda$ and $G = \langle H \cup \text{Cent}(G) \rangle_G$.*

Lemma 4.11. *If (A) then (B), where:*

- (A) (a) $G \in \mathbf{K}_{\text{lf}}^\lambda$;
 (b) $H \leq G$ and $|H| < \lambda$;
 (c) $G^* = \text{Cent}(G)$ and $G = \langle H \cup G^* \rangle_G$;
 (d) $G^* = \bigoplus_{p \in P} G_p$ and $H_p = H \cap G_p$;
- (B) (a) if $\pi \in \text{Aut}_H(G)$, then $\pi(p) := \pi \upharpoonright G_p \in \text{Aut}_{H_p}(G_p)$;
 (b) the mapping $\pi \mapsto (\pi(p) : p \in P)$ from $\text{Aut}_H(G)$ into $\prod_{p \in P} \text{Aut}_{H_p}(G_p)$ is an embedding;
 (c) the embedding in (b) is onto.

Proof. The non-trivial part is item (B)(c). To this extent, let $\pi_p \in \text{Aut}_{H_p}(G_p)$, for $p \in P$. It suffices to find $\pi \in \text{Aut}_H(G)$ such that $\pi(p) = \pi_p$, for every $p \in P$. We define π as follows. For $p_1 < \dots < p_n \in P$ an initial segment of P with the induced order, $y_{p_\ell} \in G_{p_\ell}$ and $y \in H$ we let:

$$\pi(yy_{p_1} \dots y_{p_n}) = y\pi_{p_1}(y_{p_1}) \dots \pi_{p_n}(y_{p_n}).$$

Now, by (A), every $g \in G$ has at least one representation of the form $g = yy_{p_1} \dots y_{p_n}$, and so, for every $g \in G$, $\pi(g)$ has at least one definition. We are then left to show that the choice of representation $g = yy_{p_1} \dots y_{p_n}$ does not matter. To this extent, let $g \in G$ and suppose that:

$$yy_{p_1} \dots y_{p_m} = g = y'y'_{p_1} \dots y'_{p_k},$$

By adding occurrences of e in the representations we can assume without loss of generality that:

$$yy_{p_1} \dots y_{p_n} = g = y'y'_{p_1} \dots y'_{p_n}.$$

Notice now that for $1 \leq \ell \leq n$ we have:

- (a) $y'_{p_\ell} \in y_{p_\ell}H_{p_\ell}$, say $y'_{p_\ell} = z_{p_\ell}y_{p_\ell}$ with $z_{p_\ell} \in H_{p_\ell}$;
- (b) $y'y'_{p_1} \dots y'_{p_n} = y'(z_{p_1}y_{p_1}) \dots (z_{p_n}y_{p_n}) = (y'(z_{p_1} \dots z_{p_n}))(y_{p_1} \dots y_{p_n})$;
- (c) $yy_{p_1} \dots y_{p_n} = (y'(z_{p_1} \dots z_{p_n}))(y_{p_1} \dots y_{p_n})$;
- (d) $y = y'(z_{p_1} \dots z_{p_n})$.

Hence, we have:

$$\begin{aligned} \pi(yy_{p_1} \dots y_{p_n}) &= y\pi_{p_1}(y_{p_1}) \dots \pi_{p_n}(y_{p_n}) \\ &= (y'(z_{p_1} \dots z_{p_n}))\pi_{p_1}(y_{p_1}), \dots, \pi_{p_n}(y_{p_n}) \\ &= y'z_{p_1}\pi_{p_1}(y_{p_1}) \dots z_{p_n}\pi_{p_n}(y_{p_n}) \\ &= y'\pi_{p_1}(z_{p_1}y_{p_1}) \dots \pi_{p_n}(z_{p_n}y_{p_n}) \\ &= y'\pi_{p_1}(y'_{p_1}) \dots \pi_{p_n}(y'_{p_n}) \\ &= \pi(y'y'_{p_1} \dots y'_{p_n}). \end{aligned}$$

■

Proposition 1.8. *If $G \in \mathbf{K}_{\text{lf}}^{\aleph_0}$, then $\text{Aut}(G)$ has a locally finite infinite subgroup.*

Proof. Let $G \in \mathbf{K}_{\text{lf}}^{\aleph_0}$. If $G/\text{Cent}(G)$ is infinite, then we are done, since we can embed $G/\text{Cent}(G)$ into $\text{Aut}(G)$. So suppose that $G/\text{Cent}(G)$ is finite and let G^* , G_p and H_p be as in Lemma 4.11. If for some $p \in P$ we have that G_p is infinite use Lemma 4.11 and Fact 4.6, unless G_p is bounded, in which case use Lemma 4.11 and Fact 4.3. If for every $p \in P$ we have that G_p is finite, then the set:

$$P^* = \{p \in P : G_p \neq H_p \text{ and } [G_p : H_p] > 2\}$$

is infinite, and so for every $p \in P^*$ we have that $\text{Aut}_{H_p}(G_p)$ is non-trivial. Hence, considering $\prod_{p \in P^*} \text{Aut}_{H_p}(G_p)$ and using Lemma 4.11 we are done. ■

Lemma 1.9. *To prove Conjecture 1.7 it suffices to prove Conjecture 1.10, where:*

Conjecture 1.10. *If $\lambda > \aleph_0$, $G \in \mathbf{K}_{\text{lf}}^\lambda$ is an abelian p -group, $H \leq G$ and $|H| < \lambda$, then $\text{Aut}_H(G)$ has a locally finite subgroup of power λ .*

Proof of Lemma 1.9. By Fact 4.10 and Lemma 4.11. ■

REFERENCES

- [1] László Fuchs. *Infinite Abelian Groups – Vol. I*. Pure and Applied Mathematics, Vol. 36. Academic Press, New York-London 1970.
- [2] László Fuchs. *Infinite Abelian Groups – Vol. II*. Pure and Applied Mathematics. Vol. 36-II. Academic Press, New York-London, 1973.
- [3] Alexander S. Kechris. *Classical Descriptive Set Theory*. Graduate Texts in Mathematics, 156. Springer-Verlag, New York, 1995.
- [4] Alan H. Mekler. *Stability of Nilpotent Groups of Class 2 and Prime Exponent*. J. Symbolic Logic **46** (1981), no. 4, 781-788.
- [5] Alexander S. Kechris, Andrés Nies, and Katrin Tent. *The Complexity of Topological Group Isomorphism*. J. Symbolic Logic **83** (2018), no. 3, 1190-1203.
- [6] Gianluca Paolini and Saharon Shelah. *Groups Metrics for Graph Products of Cyclic Groups*. Topology Appl. **232** (2017), 281-287.
- [7] Gianluca Paolini and Saharon Shelah. *The Automorphism Group of Hall’s Universal Group*. Proc. Amer. Math. Soc. **146** (2018), 1439-1445.
- [8] Su Gao. *Invariant Descriptive Set Theory*. Pure and Applied Mathematics (Boca Raton), 293. CRC Press, Boca Raton, FL, 2009.
- [9] Saharon Shelah. *Beginning of Stability Theory for Polish Spaces*. Israel J. Math. **214** (2016), no. 2, 507-537.

EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, ISRAEL

EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, ISRAEL AND
DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, U.S.A.