

Dynamic Assortment Optimization with Changing Contextual Information

Xi Chen ^{*1}, Yining Wang², and Yuan Zhou³

¹Stern School of Business, New York University

²Machine Learning Department, Carnegie Mellon University

³Computer Science Department, Indiana University at Bloomington

Abstract

In this paper, we study the dynamic assortment optimization problem under a finite selling season of length T . At each time period, the seller offers an arriving customer an assortment of substitutable products under a cardinality constraint, and the customer makes the purchase among offered products according to a discrete choice model. Most existing work associates each product with a real-valued fixed mean utility and assumes a multinomial logit choice (MNL) model. In many practical applications, feature/contextual information of products is readily available. In this paper, we incorporate the feature information by assuming a linear relationship between the mean utility and the feature. In addition, we allow the feature information of products to change over time so that the underlying choice model can also be non-stationary. To solve the dynamic assortment optimization under this changing contextual MNL model, we need to simultaneously learn the underlying unknown coefficient and make the decision on the assortment. To this end, we develop an upper confidence bound (UCB) based policy and establish the regret bound on the order of $\tilde{O}(d\sqrt{T})$, where d is the dimension of the feature and \tilde{O} suppresses logarithmic dependence. We further establish a lower bound $\Omega(d\sqrt{T}/K)$, where K is the cardinality constraint of an offered assortment, which is usually small. When K is a constant, our policy is optimal up to logarithmic factors. In the exploitation phase of the UCB algorithm, we need to solve a combinatorial optimization for assortment optimization based on the learned information. We further develop an approximation algorithm and an efficient greedy heuristic. The effectiveness of the proposed policy is further demonstrated by our numerical studies.

keywords: Dynamic assortment optimization, regret analysis, contextual information, bandit learning, upper confidence bounds.

1 Introduction

In operations, an important research problem facing a retailer is the selection of products/advertisements for display. For example, due to the limited shelf space, stocking restrictions, or available slots on a

* Author names listed in alphabetical order.

website, the retailer needs to carefully choose an assortment from the set of substitutable products. In an assortment optimization problem, choice model plays an important role since it characterizes a customer’s choice behavior. However, in many scenarios, customers’ choice behavior (e.g., mean utilities of products) is not given as *a priori* and cannot be easily estimated due to the insufficiency of historical data. This motivates the research of dynamic assortment optimization, which has attracted a lot of attentions from the revenue management community in recent years. A typical dynamic assortment optimization problem assumes a finite selling horizon of length T with a large T . At each time period, the seller offers an assortment of products (with the size upper bounded by K) to an arriving customer. The seller observes the customer’s purchase decision, which further provides useful information for learning utility parameters of the underlying choice model. The multinomial logit model (MNL) has been widely used in dynamic assortment optimization literature, see, e.g., [Caro & Gallien \(2007\)](#); [Rusmevichientong et al. \(2010\)](#); [Saure & Zeevi \(2013\)](#); [Agrawal et al. \(2017a,b\)](#); [Chen & Wang \(2018\)](#); [Wang et al. \(2018\)](#).

In the age of e-commerce, side information of products is widely available (e.g., brand, color, size, texture, popularity, historical selling information), which is important in characterizing customers’ preferences for products. Moreover, some features are not static and could change over time (e.g., popularity score or ratings). The feature/contextual information of products will facilitate accurate assortment decisions that are tailored to customers’ preferences. In particular, we assume at each time $t = 1, \dots, T$, each product j is associated with a d -dimensional feature vector $v_{tj} \in \mathbb{R}^d$. To incorporate the feature information, following the classical conditional logit model ([McFadden, 1973](#)), we assume that the mean utility of product j at time t (denoted by u_{tj}) obeys a linear model

$$u_{tj} = v_{tj}^\top \theta_0. \quad (1)$$

Here, $\theta_0 \in \mathbb{R}^d$ is the unknown coefficient to be learned. Based on this linear structure of the mean utility, we adopt the MNL model as the underlying choice model (see Section 2 and Eq. (3) for more details). As compared to the standard MNL, this *changing contextual MNL* model not only incorporates rich contextual information but also allows the utility to evolve over time. The changing utility is an attractive property as it captures the reality in many applications but also brings new technical challenges in learning and decision-making. For example, in existing works of ([Agrawal et al., 2017a](#)) for plain MNL choice models, upper confidence bands are constructed by providing the same assortment repetitively to incoming customers until a no-purchase activity is observed. Such an approach, however, can no longer be applied to MNL with changing contextual information as the utility parameters of products constantly evolve with time. To overcome such challenges, we propose a policy that performs optimization at every single time period, without repetitions of assortments in general.

Our model also allows the revenue for each product j to change over time. In particular, we associate the revenue parameter r_{tj} for the product j at time t .

This model generalizes the widely adopted (generalized) linear contextual bandit from machine learning literature (see, e.g., [Filippi et al. \(2010\)](#); [Chu et al. \(2011\)](#); [Abbasi-Yadkori et al. \(2011\)](#); [Agrawal & Goyal \(2013\)](#); [Li et al. \(2017\)](#) and references therein) in a non-trivial way since the MNL cannot be written in a generalized linear model form (when an assortment contains more than one product, see Section 1.1 for more details). It is also worthwhile noting that this model incorporates a personalized MNL model proposed by [Cheung & Simchi-Levi \(2017\)](#) as a special case, where each product j is associated with a fixed but unknown coefficient $\theta(j)$ and each arriving customer at time t with an observable feature vector x_t (see Section 1.1 for a more detailed discussion).

On the other hand, we choose to motivate our model from product contextual information since in practice, obtaining products’ features is usually easier (and less sensitive) than extracting customers’ preferences.

Given this contextual MNL choice model, the key challenge is how to design a policy that simultaneously learns the unknown coefficient θ_0 and sequentially makes the decision on offered assortment. The performance of a dynamic policy is usually measured by the *regret*, which is defined as the gap between the expected revenue generated by the policy and the oracle expected revenue when θ_0 (and thus the mean utilizes) is known as a priori.

The first contribution of the paper is the construction of an upper confidence bound (UCB) policy. Our UCB policy is based on the maximum likelihood estimator (MLE) and thus is named MLE-UCB. Although UCB has been a well-known technique for bandit problems, how to adopt this high-level idea to solve a problem with specific structures certainly requires technical innovations (e.g., how to build a confidence interval varies from one problem to another). In particular, our MLE-UCB contains two stages. The first stage is a *pure exploration stage* in which assortments are randomly offered and a “*pilot MLE*” is computed based on the observed purchase actions. As we will show in Lemma 1, this pilot estimator serves as a good initial estimator of θ_0 . After the exploration phase, the MLE-UCB enters the *simultaneous learning and decision-making phase*. We carefully construct an upper confidence bound of the expected revenue when offering an assortment. The added interval is based on the Fisher information matrix of the computed MLE from the previous step. Then we solve a combinatorial optimization problem to search the assortment that maximizes the upper confidence bound. By observing the customer’s purchase action based on the offered assortment, the policy updates the estimated MLE. In this update, we propose to compute a “*local MLE*”, which requires the solution to be close enough to our pilot estimator. The local MLE plays an important role in MLE-UCB policy since it guarantees that the obtained estimator at each time period is also close to the unknown true coefficient θ_0 .

Under some mild assumptions on features and coefficients, we are able to establish a regret bound $\tilde{O}(d\sqrt{T})$, where the \tilde{O} notation suppresses logarithmic dependence on T , K (cardinality constraint), and some other problem dependent parameters¹. One remarkable aspect of our regret bound is that our regret has no dependence on the total number of products N (not even in a logarithmic factor). This makes the result attractive to online applications where N is large (e.g., online advertisement). Moreover, it is also worthwhile noting the dependence of K is only through a logarithmic term.

Our second contribution is to establish the lower bound result $\Omega(d\sqrt{T}/K)$. When the maximum size of an assortment K is small (which usually holds in practice), this result shows that our policy is almost optimal.

Moreover, at each time period in the exploitation phase, our UCB policy needs to solve a combinatorial optimization problem, which searches for the best assortment (under the cardinality constraint) that minimizes the upper confidence bound of the expected revenue. Given the complicated structure of the upper confidence bound, there is no simple solution for this combinatorial problem. When K is small and N is not too large, one can directly search over all the possible sets with the size less than or equal to K . In addition to the solution of solving the combinatorial optimization exactly, the third contribution of the work is to provide an approximation algorithm based on dy-

¹For the ease of presentation in the introduction, we only present the dominating term under the common scenario that the selling horizon T is larger than the dimensionality d and the cardinality constraint K . Please refer to Theorem 1 for a more explicit expression of the obtained regret.

dynamic programming that runs in polynomial time with respect to N, K, T . Although the proposed approximation algorithm has a theoretical guarantee, it is still not efficient for dealing with large-scale applications. To this end, we further describe a computationally efficient greedy heuristic for solving this combinatorial optimization problem. The heuristic algorithm is based on the idea of local search by greedy swapping, with more details described in Sec. 5.2.

1.1 Related work

Due to the popularity of data-driven revenue management, dynamic assortment optimization, which adaptively learns unknown customers’ choice behavior, has received increasing attention in the past few years. Motivated by fast-fashion retailing, the work by Caro & Gallien (2007) first studied dynamic assortment optimization problem, but it makes a strong assumption that the demands for different product are independent. Recent works by Rusmevichientong et al. (2010); Saure & Zeevi (2013); Agrawal et al. (2017a,b); Chen & Wang (2018); Wang et al. (2018) incorporated MNL models into dynamic assortment optimization and formulated the problem into a online regret minimization problem. In particular, for capacitated MNL, Agrawal et al. (2017a) and Agrawal et al. (2017b) proposed UCB and Thompson sampling techniques and established the regret bound $\tilde{O}(\sqrt{NT})$ (when $T \gg N^2$). Chen & Wang (2018) further established a matching lower bound of $\Omega(\sqrt{NT})$. It is interesting to compare our regret to the bound for the standard MNL case. When the total number of products N is much larger than d (i.e., $N > d^2$), by incorporating the contextual information, the regret reduces from $\tilde{O}(\sqrt{NT})$ to $\tilde{O}(d\sqrt{T})$. The latter one only depends on d and is completely independent of the total number of products N , which also demonstrates the usefulness of the contextual information. Chen et al. (2018) further studied the dynamic assortment optimization under nested logit models. We also note that to highlight our key idea and focus on the balance between learning of θ_0 and revenue maximization, we study the stylized dynamic assortment optimization problems following the existing literature (Rusmevichientong et al., 2010; Saure & Zeevi, 2013; Agrawal et al., 2017a,b), which ignore operations considerations such as price decisions and inventory replenishment.

There is another line of recent research on investigating personalized assortment optimization. By incorporating the feature information of each arriving customer, both the static and dynamic assortment optimization problems are studied in Chen et al. (2015) and Cheung & Simchi-Levi (2017), respectively. It is worthwhile noting that although we do not approach our work from a personalized perspective², the *personalized MNL* considered in Cheung & Simchi-Levi (2017) can be viewed as a special of our model.

In particular, the personalized MNL assumes that each product j is associated with an unknown coefficient $\theta(j) \in \mathbb{R}^D$. When a customer arrives at time t with the *observed feature* x_t , the utility of product j at time t is $u_{tj} = x_t^\top \theta(j)$. Now we explain how to specialize our model to obtain the personalized MNL. Let us define $\theta_0 := \{\theta(1), \dots, \theta(N)\} \in \mathbb{R}^{DN}$ and the feature vector $v_{tj} := (0, \dots, x_t, \dots, 0) \in \mathbb{R}^{DN}$, which is a concatenation of N D -dimensional vectors with the j -th vector being x_t and all other vectors being 0. Then according to our linear model in Eq. (1), we have $u_{tj} = v_{tj}^\top \theta_0 = x_t^\top \theta(j)$, which recovers the personalized MNL model. Using our regret bound $\tilde{O}(d\sqrt{T})$ with $d = DN$ as the dimensionality of θ_0 , we directly obtain the regret $\tilde{O}(DN\sqrt{T})$ for the dynamic assortment optimization under the personalized MNL. As compared to the Bayesian regret

²This is because in some applications, the product features are easier to obtain by the seller as compared to customer features.

bound $\tilde{O}(DN\sqrt{KT})$ in Cheung & Simchi-Levi (2017) (see Theorem 3.3. therein), our approach still saves a factor of \sqrt{K} . We also remark that our results require a slightly stronger assumption on the contextual information vectors $\{v_{tj}\}$ compared to Cheung & Simchi-Levi (2017), which allows customer feature vectors $\{x_t\}$ to be adversarially chosen. More specifically, a stochastic assumption is imposed on $\{v_{tj}\}$ only during the *pure exploration* phase of our proposed policy. After this pure exploration phase, the feature vectors $\{v_{tj}\}$ can also be adversarially chosen. We refer the readers to Sec. 3.1 for further details.

In addition, the developed techniques in our work and Cheung & Simchi-Levi (2017) are different. Our policy is based on UCB, while the policy in Cheung & Simchi-Levi (2017) is based on Thompson sampling. Furthermore, other research studies personalized assortment optimization in an adversarial setting rather than stochastic setting. For example, Golrezaei et al. (2014); Chen et al. (2016) assumed that each customer’s choice behavior is known, but that the customers’ arriving sequence (or customers’ types) can be adversarially chosen and took the inventory level into consideration. Since the arriving sequence can be arbitrary, there is no learning component in the problem and both Golrezaei et al. (2014) and Chen et al. (2016) adopted the competitive ratio as the performance evaluation metric.

Another field of related research is the *contextual bandit* literature, in which the linear contextual bandit has been widely studied as a special case (see, e.g., Dani et al. (2008); Rusmevichientong & Tsitsiklis (2010); Chu et al. (2011); Abbasi-Yadkori et al. (2011); Agrawal & Goyal (2013) and references therein). Some recent work extends the linear contextual bandit to *generalized linear bandit* (Filippi et al., 2010; Li et al., 2017), which assumes a generalized linear reward structure. In particular, the reward r of pulling an arm given the observed feature vector of this arm x is modeled by

$$\mathbb{E}[r|x] = \sigma(x^\top \theta_0), \quad (2)$$

for an unknown linear model θ_0 and a known link function $\sigma : \mathbb{R} \rightarrow \mathbb{R}$. For example, for a linear contextual bandit, σ is the identity mapping, i.e., $\mathbb{E}[r|x] = \sigma(x^\top \theta_0)$. For the logistic contextual bandit, we have $r \in \{0, 1\}$ and $\Pr(r = 1|x) = \frac{\exp(x^\top \theta_0)}{1 + \exp(x^\top \theta_0)}$. In a standard generalized linear bandit problem (see, e.g., Li et al. (2017)) with N arms, it is assumed that a context vector v_{tj} is revealed at time t for each arm $j \in [N]$. Given a selected arm $i_t \in [N]$ at time t , the expected reward follows Eq. (2), i.e., $\mathbb{E}[r_t|v_{t,i_t}] = \sigma(v_{t,i_t}^\top \theta_0)$. At first glance, our contextual MNL model is a natural extension of the generalized linear bandit to the MNL choice model. However, when the size of an assortment $K \geq 2$, the contextual MNL *cannot* be written in the form of Eq. (2) and the denominator in the choice probability (see Eq. (3) in the next section) has a more complicated structure. Therefore, our problem is technically *not* a generalized linear model and is therefore more challenging. Moreover, in contextual bandit problems, only one arm is selected by the decision-maker at each time period. In contrast, each action in an assortment optimization problem involves a set of items, which makes the action space more complicated.

1.2 Notations and paper organization

Throughout the paper, we adopt the standard asymptotic notations. In particular, we use $f(\cdot) \lesssim g(\cdot)$ to denote that $f(\cdot) = O(g(\cdot))$. Similarly, by $f(\cdot) \gtrsim g(\cdot)$, we denote $f(\cdot) = \Omega(g(\cdot))$. We also use $f(\cdot) \asymp g(\cdot)$ for $f(\cdot) = \Theta(g(\cdot))$. Throughout this paper, we will use $C_0, C_1, C_2 \dots$ to denote universal constants. For a vector v and a matrix M , we will use $\|v\|_2$ and $\|M\|_{\text{op}}$ to denote the vector ℓ_2 -norm and the matrix spectral norm (i.e., the maximum singular value), respectively.

Moreover, for a real-valued symmetric matrix M , we denote the maximum eigenvalue and the minimum eigenvalue of M by $\lambda_{\max}(M)$ and $\lambda_{\min}(M)$, respectively, and define $\|v\|_M^2 := v^T M v$ for any given vector v . For a given integer N , we denote the set $\{1, \dots, N\}$ by $[N]$.

The rest of the paper is organized as follows. In Section 2, we introduce the mathematical formulation of our models and define the regret. In Section 3, we describe the proposed MLE-UCB policy and provide the regret analysis. The lower bound result is provided in Section 4. In Section 5, we investigate the combinatorial optimization problem in MLE-UCB and propose the approximation algorithm and greedy heuristic. The multivariate case of the approximation algorithm is relegated to the appendix. In Section 6, we provide the numerical studies. The conclusion and future directions are discussed in Section 7. Some technical proofs are provided in the online supplementary material.

2 The problem setup

There are N items, conveniently labeled as $1, 2, \dots, N$. At each time t , a set of time-sensitive “feature vectors” $v_{t1}, v_{t2}, \dots, v_{tN} \in \mathbb{R}^d$ and revenues $r_{t1}, \dots, r_{tN} \in [0, 1]$ are observed, reflecting time-varying changes of items’ revenues and customers’ preferences. A retailer, based on the features $\{v_{ti}\}_{i=1}^N$ and previous purchasing actions, picks an assortment $S_t \subseteq [N]$ under the cardinality constraint $|S_t| \leq K$ to present to an incoming customer; the retailer then observes a purchasing action $i_t \in S_t \cup \{0\}$ and collects the associated revenue r_{i_t} of the purchased item (if $i_t = 0$ then no item is purchased and zero revenue is collected).

We use an MNL model with features to characterize how a customer makes choices. Let $\theta_0 \in \mathbb{R}^d$ be an *unknown* time-invariant coefficient. For any $S \subseteq [N]$, the choice model $p_{\theta_0, t}(\cdot | S)$ is specified as (let $r_0 = 0$ and $v_{t0} = 0$)

$$p_{\theta_0, t}(j | S) = \frac{\exp\{v_{tj}^\top \theta_0\}}{1 + \sum_{k \in S} \exp\{v_{tk}^\top \theta_0\}} \quad \forall j \in S \cup \{0\}. \quad (3)$$

For simplicity, in the rest of the paper we use $p_{\theta, t}(\cdot | S)$ to denote the law of the purchased item i_t conditioned on given assortment S at time t , parameterized by the coefficient $\theta \in \mathbb{R}^d$. The expected revenue $R_t(S)$ of assortment $S \subseteq [N]$ at time t is then given by

$$R_t(S) := \mathbb{E}_{\theta_0, t}[r_{i_t} | S] = \frac{\sum_{j \in S} r_{tj} \exp\{v_{tj}^\top \theta_0\}}{1 + \sum_{j \in S} \exp\{v_{tj}^\top \theta_0\}}. \quad (4)$$

Note that throughout the paper, we use $\mathbb{E}_{\theta_0, t}[\cdot | S]$ to denote the expectation with respect to the choice probabilities $p_{\theta_0, t}(j | S)$ defined in Eq. (3).

Our objective is to design policy π such that the regret

$$\text{Regret}(\{S_t\}_{t=1}^T) = \mathbb{E}^\pi \sum_{t=1}^T R_t(S_t^*) - R_t(S_t) \quad \text{where } S_t^* = \arg \max_{S \subseteq [N], |S| \leq K} R_t(S) \quad (5)$$

is minimized. Here, S_t^* is an optimal assortment chosen when the full knowledge of choice probabilities is available (i.e., θ_0 is known).

Input: Number of pure explorations T_0 , constraint radius τ .

Output: Assortment selections $\{S_t\}_{t=1}^T \subseteq [N]$ satisfying $|S_t| \leq K$.

- 1 Pure exploration: for $t = 1, \dots, T_0$, pick $S_t = \{\ell_t\}$ for a single product ℓ_t sampled uniformly at random from $\{1, \dots, N\}$ and record purchasing actions (i_1, \dots, i_{T_0}) ;
- 2 Compute a pilot estimator using global MLE: $\theta^* \in \arg \max_{\theta \in \mathbb{R}^d} \sum_{t'=1}^{T_0} \log p_{\theta, t'}(i_{t'} | S_{t'})$;
- 3 **for** $t = T_0 + 1$ **to** T **do**
- 4 Observe revenue parameters $\{r_{tj}\}_{j=1}^N$ and preference features $\{v_{tj}\}_{j=1}^N$ at time t ;
- 5 Compute local MLE $\hat{\theta}_{t-1} \in \arg \max_{\|\theta - \theta^*\|_2 \leq \tau} \sum_{t'=1}^{t-1} \log p_{\theta, t'}(i_{t'} | S_{t'})$;
- 6 For every assortment $S \subseteq [N]$, $|S| \leq K$, compute its upper confidence bound

$$\bar{R}_t(S) := \mathbb{E}_{\hat{\theta}_{t-1, t}}[r_{tj} | S] + \min \left\{ 1, \omega \sqrt{\|\hat{I}_{t-1}^{-1/2}(\hat{\theta}_{t-1}) \widehat{M}_t(\hat{\theta}_{t-1} | S) \hat{I}_{t-1}^{-1/2}(\hat{\theta}_{t-1})\|_{\text{op}}} \right\};$$

$$\hat{I}_{t-1}(\theta) := \sum_{t'=1}^{t-1} \widehat{M}_{t'}(\theta | S_{t'}); \quad \widehat{M}_t(\theta | S) := \mathbb{E}_{\theta, t}[v_{tj} v_{tj}^\top | S] - \{\mathbb{E}_{\theta, t}[v_{tj} | S]\} \{\mathbb{E}_{\theta, t}[v_{tj} | S]\}^\top;$$

$$\omega \asymp \sqrt{d \log(\rho \nu T K)};$$

- 7 Pick $S_t \in \arg \max_{S \subseteq [N], |S| \leq K} \bar{R}_t(S)$ and observe purchasing action $i_t \in S_t \cup \{0\}$;
- 8 **end**
- 9 Remark: the expectations admit the following closed-form expressions:

$$\mathbb{E}_{\theta, t}[r_{tj} | S] = \sum_{j \in S} p_{\theta, t}(j | S) r_{tj} = \frac{\sum_{j \in S} r_{tj} \exp\{v_{tj}^\top \theta\}}{1 + \sum_{j \in S} \exp\{v_{tj}^\top \theta\}};$$

$$\mathbb{E}_{\theta, t}[v_{tj} | S] = \sum_{j \in S} p_{\theta, t}(j | S) v_{tj} = \frac{\sum_{j \in S} v_{tj} \exp\{v_{tj}^\top \theta\}}{1 + \sum_{j \in S} \exp\{v_{tj}^\top \theta\}};$$

$$\mathbb{E}_{\theta, t}[v_{tj} v_{tj}^\top | S] = \sum_{j \in S} p_{\theta, t}(j | S) v_{tj} v_{tj}^\top = \frac{\sum_{j \in S} v_{tj} v_{tj}^\top \exp\{v_{tj}^\top \theta\}}{1 + \sum_{j \in S} \exp\{v_{tj}^\top \theta\}}.$$

Algorithm 1: The MLE-UCB policy for dynamic assortment optimization with changing features

3 An MLE-UCB policy and its regret

We propose an MLE-UCB policy, described in Algorithm 1.

The policy can be roughly divided into two phases. In the first *pure exploration* phase, the policy selects assortments uniformly at random, consisting of only *one item*. The objective of the pure exploration is to establish a “pilot” estimator of the unknown coefficient θ_0 , i.e., a good initial estimator for θ_0 . For the simplicity of the analysis, we choose one item for each assortment in this phase, which facilitates us to adapt existing analysis in (Filippi et al., 2010; Li et al., 2017) as the MNL-logit choice model reduces to a *generalized linear model* when only one item is present in the assortment. In the second phase, we use a UCB-type approach that selects S_t as the assortment maximizing an *upper bound* $\bar{R}_t(S_t)$ of the expected revenue $R_t(S_t)$. Such upper bounds are built using a *local Maximum Likelihood Estimation (MLE)* of θ_0 . In particular, in Step 5, instead of computing an MLE, we compute a local MLE, where the estimator $\hat{\theta}_{t-1}$ lies in a ball centered at the pilot estimator θ^* with a radius τ . This localization also simplifies the technical analysis based

on Taylor expansion, which benefits from the constraint that $\widehat{\theta}_{t-1}$ is not too far away from θ^* .

To construct the confidence bound, we introduce the matrices $\widehat{M}_t(\widehat{\theta}_{t-1}|S)$ and $\widehat{I}_{t-1}(\widehat{\theta}_{t-1})$ in Step 6 of Algorithm 1, which are empirical estimates of the *Fisher's information* matrices $-\mathbb{E}[\nabla_{\theta}^2 \log p(\cdot|\theta)]$ corresponding to the MNL choice model $p(\cdot|S_t)$. The population version of the Fisher's information matrices are presented in Eq. (8) in Sec. 3.2.2. These quantities play an essential role in classical statistical analysis of maximum likelihood estimators (see, e.g., (Van der Vaart, 2000)).

The proposed MLE-UCB policy has three hyper-parameters: the coefficient $\omega > 0$ that controls the lengths of confidence intervals of $R_t(S)$, the number of pure exploration iterations T_0 , and the radius τ_0 in the local MLE formulation. While theoretical values of ω, T_0 and τ are given in Theorem 1, which potentially depend on several unknown problem parameters, in practice we recommend the usage of $T_0 = \max\{d \log T, T^{1/4}\}$, $\omega = \sqrt{d \log T}$ and $\tau = 1/K$.

In the rest of this section, we give a regret analysis that shows an $\widetilde{O}(d\sqrt{T})$ upper bound on the regret of the MLE-UCB policy. Additionally, we prove a lower bound of $\widetilde{O}(d\sqrt{T}/K)$ in Sec. 4 and show how the combinatorial optimization in Step 7 can be approximately computed efficiently in Sec. 5.

3.1 Regret analysis

To establish rigorous regret upper bounds on Algorithm 1, we impose the following assumptions:

- (A1) There exists a constant ν such that $\|v_{tj}\|_2 \leq \nu$ for all t and j . Moreover, for all $t \leq T_0$ and $j \in [N]$, v_{tj} are i.i.d. generated from an unknown distribution with the density μ satisfying that $\lambda_{\min}(\mathbb{E}_{\mu} v v^{\top}) \geq \lambda_0$ for some constant $\lambda_0 > 0$;
- (A2) There exists a constant $\rho < \infty$ such that for all $t \in [T]$ and $S \subseteq [N]$ with $|S| \leq K$, $\frac{p_{\theta_0,t}(j|S)}{p_{\theta_0,t}(j'|S)} \leq \rho$ for all $j, j' \in S \cup \{0\}$.

The item (A1) assumes that the contextual information vectors $\{v_{tj}\}$ in the pure-exploration phase with $t \leq T_0$ are *randomly generated* from a non-degenerate density. It also places a standard boundedness condition on $\{v_{tj}\}$ for all time periods t . Note that after the pure-exploration phase, we allow the contextual vectors $\{v_{tj}\}$ to be adversarially chosen, only subject to boundedness conditions. (A2) additionally assumes a bounded ratio between the probability of choosing any two different items in an arbitrary assortment set. We remark that if $\|\theta_0\|_2 \leq C$, then the boundedness assumption in (A1) implies (A2) with $\rho \leq e^{2 \max\{1, C\nu\}}$.

We are now ready to state our main result that upper bounds the worst-case accumulated regret of the proposed MLE-UCB policy in Algorithm 1.

Theorem 1. *Suppose that $T_0 \asymp \max\{\nu^2 d \log T / \lambda_0^2, \rho^2 (d + \log T) / (\tau^2 \lambda_0)\}$ and $\tau \asymp 1 / \sqrt{\rho^2 \nu^2 K^2}$ in Algorithm 1, then the regret of the MLE-UCB policy is upper bounded by*

$$C_1 \left[d\sqrt{T} \cdot \log(\lambda_0^{-1} \rho \nu T K) + d^2 \lambda_0^{-2} \rho^4 \nu^2 K^2 \log T \right] + C_2, \quad (6)$$

where $C_1, C_2 > 0$ are universal constants.

In addition to universal constants, the regret upper bound established in Theorem 1 has two terms. The first term, $d\sqrt{T} \cdot \log(\lambda_0^{-1} \rho \nu T K)$, is the main regret term that scales as $\widetilde{O}(d\sqrt{T})$ dropping logarithmic dependency. The second $d^2 \lambda_0^{-2} \rho^4 \nu^2 K^2 \log T$ term is a minor term, because it only

scales logarithmically with the time horizon T . One remarkable aspect of Theorem 1 is the fact that the regret upper bound has *no* dependency on the total number of items N (even in a logarithmic term). This is an attractive property of the proposed policy, which allows N to be very large, even exponentially large in d and K .

3.2 Proof sketch of Theorem 1

We provide a proof sketch of Theorem 1 in this section. The proofs of technical lemmas are relegated to the online supplement.

The proof is divided into four steps. In the first step, we analyze the pilot estimator θ^* obtained from the pure exploration phase of Algorithm 1, and show as a corollary that the true model θ_0 is feasible to all subsequent local MLE formulations with high probability (see Corollary 1). In the second step, we use an ε -net argument to analyze the estimation error of the local MLE. Afterwards, we show in the third step that an upper bound on the estimation error $\hat{\theta}_{t-1} - \theta_0$ implies an upper bound on the estimation error of the expected revenue $R_t(S)$, hence showing that $\bar{R}_t(S)$ are valid upper confidence bounds. Finally, we apply the *elliptical potential lemma*, which also plays a key role in linear stochastic bandit and its variants, to complete our proof.

3.2.1 Analysis of pure exploration and the pilot estimator

Our first step is to establish an upper bound on the estimation error $\|\theta^* - \theta_0\|_2$ of the pilot estimator θ^* , built using pure exploration data. It should be noted that in the pure exploration phase ($t \in \{1, \dots, T_0\}$), the assortments $\{S_t\}_{t=1}^{T_0}$ only consist of one item. Therefore the observation model reduces to a standard *generalized linear model* with the sigmoid function $\sigma(x) = 1/(1 + e^{-x}) = e^x/(1 + e^x)$ as the link function, which is essentially a logistic regression model of observing 1 if the customer makes a purchase.

Because the choice model in the pure exploration phase reduces to a generalized linear model, we can cite existing works to upper bound the error $\|\theta^* - \theta_0\|_2$. In particular, the following lemma is cited from (Li et al., 2017, Eq. (18)), adapted to our model and parameter settings. The details on how to adapt the result from (Li et al., 2017) provided in the supplementary material.

Lemma 1. *With probability $1 - \delta$ it holds that*

$$\|\theta^* - \theta_0\|_2 \leq \frac{2}{\kappa} \sqrt{\frac{d + \log(1/\delta)}{\lambda_{\min}(V)}} \quad \text{where } \kappa = \frac{1}{2e(1 + \rho)} \quad \text{and } V = \sum_{t=1}^{T_0} v_{t,it} v_{t,it}^\top. \quad (7)$$

The following corollary immediately follows Lemma 1, by lower bounding $\lambda_{\min}(V)$ using standard matrix concentration inequalities. Its proof is again deferred to the supplementary material.

Corollary 1. *There exists a universal constant $C_0 > 0$ such that for arbitrary $\tau \in (0, 1/2]$, if $T_0 \geq C_0 \max\{\nu^2 d \log T / \lambda_0^2, \rho^2 (d + \log T) / (\tau^2 \lambda_0)\}$ then with probability $1 - O(T^{-1})$, $\|\theta^* - \theta_0\|_2 \leq \tau$.*

The purpose of Corollary 1 is to establish a connection between the number of pure exploration iterations T_0 and the critical radius τ used in the local MLE formulation. It shows a lower bound on T_0 in order for the estimation error $\|\theta^* - \theta_0\|_2$ to be upper bounded by τ with high probability, which certifies that the true model θ_0 is also a *feasible* local estimator in our MLE-UCB policy. This is an important property for later analysis of local MLE solutions $\hat{\theta}_{t-1}$.

3.2.2 Analysis of the local MLE

The following lemma upper bounds a Mahalanobis distance between $\hat{\theta}_t$ and θ_0 . For convenience, we adopt the notation that $r_{t0} = 0$ and $v_{t0} = 0$ for all t throughout this section. We also define

$$I_t(\theta) := \sum_{t'=1}^t M_{t'}(\theta), \quad (8)$$

$$M_{t'}(\theta) := -\mathbb{E}_{\theta_0, t'}[\nabla_{\theta}^2 \log p_{\theta, t'}(j|S_{t'})]$$

$$= \mathbb{E}_{\theta_0, t'}[v_{t'j} v_{t'j}^{\top}] - \{\mathbb{E}_{\theta_0, t'} v_{t'j}\} \{\mathbb{E}_{\theta_0, t'} v_{t'j}\}^{\top} - \{\mathbb{E}_{\theta_0, t'} v_{t'j}\} \{\mathbb{E}_{\theta_0, t'} v_{t'j}\}^{\top} + \{\mathbb{E}_{\theta_0, t'} v_{t'j}\} \{\mathbb{E}_{\theta_0, t'} v_{t'j}\}^{\top}$$

where $\mathbb{E}_{\theta_0, t'}$ denotes the expectation evaluated under the law $j \sim p_{\theta_0, t'}(\cdot|S_{t'})$; that is, $p_{\theta_0, t'}(j|S_{t'}) = \exp\{v_{t'j}^{\top} \theta\} / (1 + \sum_{k \in S_{t'}} \exp\{v_{t'k}^{\top} \theta\})$ for $j \in S_{t'}$ and $p_{\theta_0, t'}(j|S_{t'}) = 0$ for $j \notin S_{t'}$.

Lemma 2. *Suppose $\tau \leq 1/\sqrt{8\rho\nu^2 K^2}$. Then there exists a universal constant $C > 0$ such that with probability $1 - O(T^{-1})$ the following holds uniformly over all $t = T_0, \dots, T-1$:*

$$(\hat{\theta}_t - \theta_0)^{\top} I_t(\theta_0) (\hat{\theta}_t - \theta_0) \leq C \cdot d \log(\rho\nu TK). \quad (9)$$

Remark 1. *For $\theta = \theta_0$, the expression of $M_{t'}(\theta)$ can be simplified as $M_{t'}(\theta_0) = \mathbb{E}_{\theta_0, t'}[v_{t'j} v_{t'j}^{\top}] - \{\mathbb{E}_{\theta_0, t'} v_{t'j}\} \{\mathbb{E}_{\theta_0, t'} v_{t'j}\}^{\top}$.*

The complete proof of Lemma 2 is given in the supplementary material, and here we provide some high-level ideas behind our proof.

Our proof is inspired by the classical convergence rate analysis of M-estimators (Van der Vaart, 2000, Sec. 5.8). The main technical challenge is to provide *finite-sample* analysis of several components in the proof of (Van der Vaart, 2000, Sec. 5.8).

In particular, for any $\theta \in \mathbb{R}^d$, consider

$$F_t(\theta) := \sum_{t' \leq t} f_{t'}(\theta) \quad \text{where} \quad f_{t'}(\theta) := \mathbb{E}_{\theta_0, t'} \left[\log \frac{p_{\theta, t'}(j|S_{t'})}{p_{\theta_0, t'}(j|S_{t'})} \right] = \sum_{j \in S_{t'} \cup \{0\}} p_{\theta_0, t'}(j|S_{t'}) \log \frac{p_{\theta, t'}(j|S_{t'})}{p_{\theta_0, t'}(j|S_{t'})}$$

and its ‘‘sample’’ version

$$\hat{F}_t(\theta) := \sum_{t' \leq t} \hat{f}_{t'}(\theta) \quad \text{where} \quad \hat{f}_{t'}(\theta) := \log \frac{p_{\theta, t'}(i_{t'}|S_{t'})}{p_{\theta_0, t'}(i_{t'}|S_{t'})}.$$

It is easy to verify by definition that $F_t(\hat{\theta}_t) \geq F_t(\theta_0) = 0$ and $\hat{F}_t(\hat{\theta}_t) \leq \hat{F}_t(\theta_0) = 0$, because $F_t(\cdot)$ is a Kullback-Leibler divergence, θ_0 is feasible to the local MLE formulation and $\hat{\theta}_{t-1}$ is the optimal solution. On the other hand, it can be proved that $|F_t(\theta) - \hat{F}_t(\theta)|$ is small for all θ with high probability, by using concentration inequalities for self-normalized empirical process (note that $\mathbb{E} \hat{f}_{t'}(\theta) = f_{t'}(\theta)$ for any θ). Moreover, by constructing a local quadratic approximation of $F_t(\cdot)$ around θ_0 , we can show that $F_t(\theta) - F_t(\theta_0)$ is large when θ is far away from θ_0 .

Following the above observations, we can use proof by contradiction to prove Lemma 2, which essentially claims that $\hat{\theta}_t$ and θ_0 are close under the quadratic distance $\|\cdot\|_{I_t(\theta_0)}$. Suppose by

contradiction that $\widehat{\theta}_t$ and θ_0 are far apart, which implies that $|F_t(\widehat{\theta}_t) - F_t(\theta_0)|$ is large. On the other hand, by the fact that $\widehat{F}_t(\widehat{\theta}_t) \leq 0 = F_t(\theta_0) \leq F_t(\widehat{\theta}_t)$, we have

$$|F_t(\widehat{\theta}_t) - F_t(\theta_0)| = |F_t(\widehat{\theta}_t)| \leq |F_t(\widehat{\theta}_t) - \widehat{F}_t(\widehat{\theta}_t)|.$$

By the established concentration result, we have $|F_t(\theta) - \widehat{F}_t(\theta)|$ is small for all θ with high probability (including $\theta = \widehat{\theta}_t$). This leads to the desired contradiction.

3.2.3 Analysis of upper confidence bounds

The following technical lemma shows that the upper confidence bounds constructed in Algorithm 1 are valid with high probability. Additionally, we establish an upper bound on the discrepancy between $\overline{R}_t(S)$ and the true value $R_t(S)$ defined in Eq. (4).

Lemma 3. *Suppose τ satisfies the condition in Lemma 2. With probability $1 - O(T^{-1})$ the following holds uniformly for all $t > T_0$ and $S \subseteq [N]$, $|S| \leq K$ such that*

1. $\overline{R}_t(S) \geq R_t(S)$;
2. $|\overline{R}_t(S) - R_t(S)| \lesssim \min\{1, \omega \sqrt{\|I_{t-1}^{-1/2}(\theta_0) M_t(\theta_0|S) I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}}\}$.

At a higher level, the proof of Lemma 3 can be regarded as a “finite-sample” version of the classical *Delta’s method*, which upper bounds estimation error of some functional φ of parameters, i.e., $|\varphi(\widehat{\theta}_{t-1}) - \varphi(\theta_0)|$ using the estimation error of the parameters themselves $\widehat{\theta}_{t-1} - \theta_0$. The complete proof is relegated to the supplementary material.

3.2.4 The elliptical potential lemma

Let S_t^* be the assortment that maximizes the expected revenue $R_t(\cdot)$ (defined in Eq. (4)) at time period t , and S_t be the assortment selected by Algorithm 1. Because $R_t(S) \leq \overline{R}_t(S)$ for all S (see Lemma 3), we have the following upper bound for each term in the regret (see Eq. (5)):

$$R_t(S_t^*) - R_t(S_t) \leq (\overline{R}_t(S_t^*) - \overline{R}_t(S_t)) + (\overline{R}_t(S_t) - R_t(S_t)) \leq \overline{R}_t(S_t) - R_t(S_t), \quad (10)$$

where the last inequality holds because $\overline{R}_t(S_t^*) - \overline{R}_t(S_t) \leq 0$ (note that S_t maximizes $\overline{R}_t(\cdot)$).

Subsequently, invoking Lemma 3 and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \sum_{t=T_0+1}^T R_t(S_t^*) - R_t(S_t) &\lesssim \sqrt{d \log(\rho \nu T K)} \cdot \sum_{t=T_0+1}^T \sqrt{\min\{1, \|I_{t-1}^{-1/2}(\theta_0) M_t(\theta_0|S_t) I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}\}} \\ &\lesssim \sqrt{dT \log(\rho \nu T K)} \cdot \sum_{t=T_0+1}^T \min\{1, \|I_{t-1}^{-1/2}(\theta_0) M_t(\theta_0|S_t) I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}^2\}. \end{aligned} \quad (11)$$

The following lemma is a key result that upper bounds $\sum_{t=T_0+1}^T \min\{1, \|I_{t-1}^{-1/2}(\theta_0) M_t(\theta_0|S_t) I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}^2\}$. It is usually referred to as the *elliptical potential lemma* and has found many applications in contextual bandit-type problems (see, e.g., Dani et al. (2008); Rusmevichientong et al. (2010); Filippi et al. (2010); Li et al. (2017)).

Lemma 4. *It holds that*

$$\sum_{t=T_0+1}^T \min\{1, \|I_{t-1}^{-1/2}(\theta_0)M_t(\theta_0|S_t)I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}^2\} \leq 4 \log \frac{\det I_T(\theta_0)}{\det I_{T_0}(\theta_0)} \lesssim d \log(\lambda_0^{-1} \rho \nu).$$

The proof of Lemma 4 is placed in the supplementary material. It is a routine proof following existing proofs of elliptical potential lemmas using matrix-determinant rank-1 updates.

We are now ready to give the final upper bound on $\text{Regret}(\{S_t\}_{t=1}^T)$ defined in Eq. (5). Note that the total regret incurred by the pure exploration phase is upper bounded by T_0 , because the revenue parameters r_{tj} are normalized so that they are upper bounded by 1. In addition, as the failure event of $\bar{R}_t(S) \leq R_t(S)$ for some S occurs with probability $1 - O(T^{-1})$, the total regret accumulated under the failure event is $O(T^{-1}) \cdot T = O(1)$. Further invoking Eq. (11) and Lemma 4, we have

$$\begin{aligned} \text{Regret}(\{S_t\}_{t=1}^T) &\leq T_0 + O(1) + \mathbb{E} \sum_{t=T_0+1}^T R_t(S_t^*) - R_t(S_t) \\ &\lesssim O(1) + \frac{\nu^2 d \log T}{\lambda_0^2} + \frac{\rho^2 (d + \log T)}{\tau^2 \lambda_0} + d\sqrt{T} \cdot \log(\lambda_0^{-1} \rho \nu T K) \\ &\lesssim O(1) + d^2 \lambda_0^{-2} \rho^4 \nu^2 K^2 \log T + d\sqrt{T} \cdot \log(\lambda_0^{-1} \rho \nu T K). \end{aligned} \quad (12)$$

4 Lower bound

To complement our regret analysis in Sec. 3.1, in this section we prove a *lower* bound for worst-case regret. Our lower bound is information theoretical, and therefore applies to *any* policy for dynamic assortment optimization with changing contextual features.

Theorem 2. *Suppose d is divisible by 4. There exists a universal constant $C_0 > 0$ such that for any sufficiently large T and policy π , there is a worst-case problem instance with $N \asymp K \cdot 2^d$ items and uniformly bounded feature and coefficient vector (i.e., $\|v_{ti}\|_2 \leq 1$ and $\|\theta_0\|_2 \leq 1$ for all $i \in [N]$, $t \in [T]$) such that the regret of π is lower bounded by $C_2 \cdot d\sqrt{T}/K$.*

Theorem 2 essentially implies that the $\tilde{O}(d\sqrt{T})$ regret upper bound established in Theorem 1 is tight (up to logarithmic factors) in T and d . Although there is an $O(K)$ gap between the upper and lower regret bounds, in practical applications K is usually small and can be generally regarded as a constant. It is an interesting technical open problem to close this gap of $O(K)$.

We also remark that an $\Omega(d\sqrt{T})$ lower bound was established in (Dani et al., 2008) for contextual linear bandit problems. However, in assortment selection, the reward function is *not* coordinate-wise decomposable, making techniques in Dani et al. (2008) not directly applicable. In the following subsection, we provide a high-level proof sketch of Theorem 2, with complete proofs of technical lemmas relegated to the supplementary material.

4.1 Proof sketch of Theorem 2

At a higher level, the proof of Theorem 2 can be divided into three steps (separated into three different sub-sections below). In the first step, we construct an *adversarial parameter* set and reduce

the task of lower bounding the *worst-case* regret of any policy to lower bounding the *Bayes risk* of the constructed parameter set. In the second step, we use a “counting argument” similar to the one developed in [Chen & Wang \(2018\)](#) to provide an explicit lower bound on the Bayes risk of the constructed adversarial parameter set, and finally we apply *Pinsker’s inequality* (see, e.g., [Tsybakov \(2009\)](#)) to derive a complete lower bound.

4.1.1 Adversarial construction and the Bayes risk

Let $\epsilon \in (0, 1/d\sqrt{d})$ be a small positive parameter to be specified later. For every subset $W \subseteq [d]$, define the corresponding parameter $\theta_W \in \mathbb{R}^d$ as $[\theta_W]_i = \epsilon$ for all $i \in W$, and $[\theta_W]_i = 0$ for all $i \notin W$. The parameter set we consider is

$$\theta \in \Theta := \{\theta_W : W \in \mathcal{W}_{d/4}\} := \{\theta_W : W \subseteq [d], |W| = d/4\}. \quad (13)$$

Note that $d/4$ is a positive integer because d is divisible by 4, as assumed in [Theorem 2](#). Also, to simplify notation, we use \mathcal{W}_k to denote the class of all subsets of $[d]$ whose size is k .

The feature vectors $\{v_{ti}\}$ are constructed to be invariant across time iterations t . For each t and $U \in \mathcal{W}_{d/4}$, K identical feature vectors v_U are constructed as (recall that K is the maximum allowed assortment capacity)

$$[v_U]_i = 1/\sqrt{d} \quad \text{for } i \in U; \quad [v_U]_i = 0 \quad \text{for } i \notin U. \quad (14)$$

It is easy to check that with the condition $\epsilon \in (0, 1/\sqrt{d})$, $\|\theta_W\|_2 \leq 1$ and $\|v_U\|_2 \leq 1$ for all $W, U \in \mathcal{W}_{d/4}$. Hence the worst-case regret of any policy π can be lower bounded by the worst-case regret of parameters belonging to Θ , which can be further lower bounded by the “average” regret over a uniform prior over Θ :

$$\begin{aligned} \sup_{v, \theta} \mathbb{E}_{v, \theta}^\pi \sum_{t=1}^T R(S_\theta^*) - R(S_t) &\geq \max_{\theta_W \in \Theta} \mathbb{E}_{v, \theta_W}^\pi \sum_{t=1}^T R(S_{\theta_W}^*) - R(S_t) \\ &\geq \frac{1}{|\mathcal{W}_{d/4}|} \sum_{W \in \mathcal{W}_{d/4}} \mathbb{E}_{v, \theta_W}^\pi \sum_{t=1}^T R(S_{\theta_W}^*) - R(S_t). \end{aligned} \quad (15)$$

Here S_θ^* is the optimal assortment of size at most K that maximizes (expected) revenue under parameterization θ . By construction, it is easy to verify that $S_{\theta_W}^*$ consists of all K items corresponding to feature v_W . We also employ constant revenue parameters $r_{ti} \equiv 1$ for all $t \in [T]$, $i \in [N]$.

4.1.2 The counting argument

In this section we drive an explicit lower bound on the Bayes risk in [Eq. \(15\)](#). For any sequences $\{S_t\}_{t=1}^T$ produced by the policy π , we first describe an alternative sequence $\{\tilde{S}_t\}_{t=1}^T$ that provably enjoys less regret under parameterization θ_W , while simplifying our analysis.

Let v_{U_1}, \dots, v_{U_L} be the distinct feature vectors contained in assortment S_t (if $S_t = \emptyset$ then one may choose an arbitrary feature v_U) with $U_1, \dots, U_L \in \mathcal{W}_{d/4}$. Let U^* be the subset among U_1, \dots, U_L that maximizes $\langle v_{U^*}, \theta_W \rangle$, where θ_W is the underlying parameter. Let \tilde{S}_t be the assortment consisting of all K items corresponding to feature v_{U^*} . We then have the following observation:

Proposition 1. $R(S_t) \leq R(\tilde{S}_t)$ under θ_W .

Proof. Because $r_{tj} \equiv 1$ in our construction, we have $R(S_t) = (\sum_{j \in S_t} u_j) / (1 + \sum_{j \in S_t} u_j)$ where $u_j = \exp\{v_j^\top \theta_W\}$ under θ_W . Clearly $R(S)$ is a monotonically non-decreasing function in u_j . By replacing all $v_j \in S_t$ with $v_{U^*} \in \tilde{S}_t$, the u_j values do not decrease and therefore the Proposition holds true. \square

To simplify notation we also use \tilde{U}_t to denote the unique $U^* \in \mathcal{W}_{d/4}$ in \tilde{S}_t . We also use \mathbb{E}_W and \mathbb{P}_W to denote the law parameterized by θ_W and policy π . The following lemma gives a lower bound on $R(\tilde{S}_t) - R(S_{\theta_W}^*)$ by comparing it with W , which is also proved in the supplementary material.

Lemma 5. Suppose $\epsilon \in (0, 1/d\sqrt{d})$ and define $\delta := d/4 - |\tilde{U}_t \cap W|$. Then

$$R(S_{\theta_W}^*) - R(\tilde{S}_t) \geq \frac{\delta \epsilon}{4K\sqrt{d}}.$$

Define random variables $\tilde{N}_i := \sum_{t=1}^T \mathbf{1}\{i \in \tilde{U}_t\}$. Lemma 5 immediately implies

$$\mathbb{E}_W \sum_{t=1}^T R(S_{\theta_W}^*) - R(\tilde{S}_t) \geq \frac{\epsilon}{4K\sqrt{d}} \left(\frac{dT}{4} - \sum_{i \in W} \mathbb{E}_W[\tilde{N}_i] \right), \quad \forall W \in \mathcal{W}_{d/4}. \quad (16)$$

Denote $\mathcal{W}_{d/4}^{(i)} := \{W \in \mathcal{W}_{d/4} : i \in W\}$ and $\mathcal{W}_{d/4-1} := \{W \subseteq [d] : |W| = d/4 - 1\}$. Averaging both sides of Eq. (16) with respect to all $W \in \mathcal{W}_{d/4}$ and swapping the summation order, we have

$$\begin{aligned} \frac{1}{|\mathcal{W}_{d/4}|} \sum_{W \in \mathcal{W}_{d/4}} \mathbb{E}_W \sum_{t=1}^T R(S_{\theta_W}^*) - R(S_t) &\geq \frac{\epsilon}{4K\sqrt{d}} \frac{1}{|\mathcal{W}_{d/4}|} \sum_{W \in \mathcal{W}_{d/4}} \left(\frac{dT}{4} - \sum_{i \in W} \mathbb{E}_W[\tilde{N}_i] \right) \\ &= \frac{\epsilon}{4K\sqrt{d}} \left(\frac{dT}{4} - \frac{1}{|\mathcal{W}_{d/4}|} \sum_{i=1}^d \sum_{W \in \mathcal{W}_{d/4}^{(i)}} \mathbb{E}_W[\tilde{N}_i] \right) \\ &= \frac{\epsilon}{4K\sqrt{d}} \left(\frac{dT}{4} - \frac{1}{|\mathcal{W}_{d/4}|} \sum_{W \in \mathcal{W}_{d/4-1}} \sum_{i \notin W} \mathbb{E}_{W \cup \{i\}}[\tilde{N}_i] \right) \\ &\geq \frac{\epsilon}{4K\sqrt{d}} \left(\frac{dT}{4} - \frac{|\mathcal{W}_{d/4-1}|}{|\mathcal{W}_{d/4}|} \max_{W \in \mathcal{W}_{d/4-1}} \sum_{i \notin W} \mathbb{E}_{W \cup \{i\}}[\tilde{N}_i] \right) \\ &= \frac{\epsilon}{4K\sqrt{d}} \left(\frac{dT}{4} - \frac{|\mathcal{W}_{d/4-1}|}{|\mathcal{W}_{d/4}|} \max_{W \in \mathcal{W}_{d/4-1}} \sum_{i \notin W} \mathbb{E}_W[\tilde{N}_i] + \mathbb{E}_{W \cup \{i\}}[\tilde{N}_i] - \mathbb{E}_W[\tilde{N}_i] \right). \end{aligned}$$

Note that for any fixed W , $\sum_{i \notin W} \mathbb{E}_W[\tilde{N}_i] \leq \sum_{i=1}^d \mathbb{E}_W[\tilde{N}_i] \leq dT/4$. Also, $|\mathcal{W}_{d/4-1}|/|\mathcal{W}_{d/4}| =$

$\binom{d}{d/4-1}/\binom{d}{d/4} = \frac{d/4}{3d/4+1} \leq 1/3$. Subsequently,

$$\frac{1}{|\mathcal{W}_{d/4}|} \sum_{W \in \mathcal{W}_{d/4}} \mathbb{E}_W \sum_{t=1}^T R(S_{\theta_W}^*) - R(S_t) \geq \frac{\epsilon}{4K\sqrt{d}} \left(\frac{dT}{6} - \max_{W \in \mathcal{W}_{d/4-1}} \sum_{i \notin W} |\mathbb{E}_{W \cup \{i\}}[\tilde{N}_i] - \mathbb{E}_W[\tilde{N}_i]| \right). \quad (17)$$

4.1.3 Pinsker's inequality

In this section we concentrate on upper bounding $|\mathbb{E}_{W \cup \{i\}}[\tilde{N}_i] - \mathbb{E}_W[\tilde{N}_i]|$ for any $W \in \mathcal{W}_{d/4-1}$. Let $P = \mathbb{P}_W$ and $Q = \mathbb{P}_{W \cup \{i\}}$ denote the laws under θ_W and $\theta_{W \cup \{i\}}$, respectively. Then

$$\begin{aligned} |\mathbb{E}_P[\tilde{N}_i] - \mathbb{E}_Q[\tilde{N}_i]| &\leq \sum_{j=0}^T j \cdot |P[\tilde{N}_i = j] - Q[\tilde{N}_i = j]| \\ &\leq T \cdot \sum_{j=0}^T |P[\tilde{N}_i = j] - Q[\tilde{N}_i = j]| \\ &\leq T \cdot \|P - Q\|_{\text{TV}} \leq T \cdot \sqrt{\frac{1}{2} \text{KL}(P\|Q)}, \end{aligned}$$

where $\|P - Q\|_{\text{TV}} = \sup_A |P(A) - Q(A)|$ is the total variation distance between P, Q , $\text{KL}(P\|Q) = \int (\log dP/dQ) dP$ is the Kullback-Leibler (KL) divergence between P, Q , and the inequality $\|P - Q\|_{\text{TV}} \leq \sqrt{\frac{1}{2} \text{KL}(P\|Q)}$ is the celebrated Pinsker's inequality.

For every $i \in [d]$ define random variables $N_i := \sum_{t=1}^T \frac{1}{K} \sum_{v_U \in S_t} \mathbf{1}\{i \in U\}$. The next lemma upper bound the KL divergence, which is proved in the supplementary material.

Lemma 6. *For any $W \in \mathcal{W}_{d/4-1}$ and $i \in [d]$, $\text{KL}(P_W\|P_{W \cup \{i\}}) \leq C_{\text{KL}} \cdot \mathbb{E}_W[N_i] \cdot \epsilon^2/d$ for some universal constant $C_{\text{KL}} > 0$.*

Combining Lemma 6 and Eq. (17), we have

$$\frac{1}{|\mathcal{W}_{d/4}|} \sum_{W \in \mathcal{W}_{d/4}} \mathbb{E}_W \sum_{t=1}^T R(S_{\theta_W}^*) - R(S_t) \geq \frac{\epsilon}{4K\sqrt{d}} \left(\frac{dT}{6} - T \sum_{i=1}^d \sqrt{C_{\text{KL}} \mathbb{E}_W[N_i] \epsilon^2/d} \right).$$

Further using Cauchy-Schwartz inequality, we have

$$\sum_{i=1}^d \sqrt{C_{\text{KL}} \mathbb{E}_W[N_i] \epsilon^2/d} \leq \sqrt{d} \cdot \sqrt{\sum_{i=1}^d C_{\text{KL}} \mathbb{E}_W[N_i] \epsilon^2/d},$$

which is further upper bounded by $\sqrt{d} \cdot \sqrt{C_{\text{KL}} T \epsilon^2/4}$ because $\sum_{i=1}^d \mathbb{E}_W[N_i] \leq dT/4$. Subsequently,

$$\frac{1}{|\mathcal{W}_{d/4}|} \sum_{W \in \mathcal{W}_{d/4}} \mathbb{E}_W \sum_{t=1}^T R(S_{\theta_W}^*) - R(S_t) \geq \frac{\epsilon}{4K\sqrt{d}} \left(\frac{dT}{6} - T \sqrt{C'_{\text{KL}} dT \epsilon^2} \right), \quad (18)$$

where $C'_{\text{KL}} = C_{\text{KL}}/4$. Setting $\epsilon = \sqrt{d/144C'_{\text{KL}}T}$ we complete the proof of Theorem 2.

5 The combinatorial optimization subproblem

The major computational bottleneck of our algorithm is its Step 6, which involves solving a *combinatorial* optimization problem. For notational simplicity, we equivalently reformulate this problem as follows:

$$\max_{S \subseteq [N], |S| \leq K} \text{ESTR}(S) + \min\{1, \omega \cdot \text{CI}(S)\} \quad \text{where } \text{ESTR}(S) := \frac{\sum_{j \in S} r_{tj} \hat{u}_{tj}}{1 + \sum_{j \in S} \hat{u}_{tj}} \quad \text{and} \quad (19)$$

$$\text{CI}(S) := \sqrt{\left\| \left\| \frac{\sum_{j \in S} \hat{u}_{tj} x_{tj} x_{tj}^\top}{1 + \sum_{j \in S} \hat{u}_{tj}} - \left(\frac{\sum_{j \in S} \hat{u}_{tj} x_{tj}}{1 + \sum_{j \in S} \hat{u}_{tj}} \right) \left(\frac{\sum_{j \in S} \hat{u}_{tj} x_{tj}}{1 + \sum_{j \in S} \hat{u}_{tj}} \right)^\top \right\|_{\text{op}} \right\|}.$$

Here $\hat{u}_{tj} := \exp\{v_{tj}^\top \hat{\theta}_{t-1}\}$ and $x_{tj} := \hat{I}_{t-1}^{-1/2}(\hat{\theta}_{t-1})v_{tj}$, both of which can be pre-computed before solving Eq. (19).

A brute-force way to compute Eq. (19) is to enumerate all subsets $S \subseteq [N]$, $|S| \leq K$ and select the one with the largest objective value. Such an approach is *not* an efficient (polynomial-time) algorithm and is therefore not scalable.

In this section we provide two alternative methods for (approximately) solving the combinatorial optimization problem in Eq. (19). Our first algorithm is based on discretized dynamic programming and enjoys rigorous approximation guarantees. The second algorithm is a computationally efficient greedy heuristic. Although the greedy heuristic does not have rigorous guarantees, our numerical result suggests it works reasonably well (see Sec. 6).

5.1 Approximation algorithms for assortment optimization

In this section we introduce algorithms with polynomial running times and rigorous approximation guarantees for the optimization task described in Eq. (19). We first formally introduce the concept of $(\alpha, \varepsilon, \delta)$ -approximation to characterize the approximation performance, and show that such approximation guarantees imply certain upper bounds on the final regret.

Definition 1 ($(\alpha, \varepsilon, \delta)$ -approximation). *Fix $\alpha \geq 1$, $\varepsilon \geq 0$ and $\delta \in [0, 1)$. An algorithm is an $(\alpha, \varepsilon, \delta)$ -approximation algorithm if it produces $\hat{S} \subseteq [N]$, $|\hat{S}| \leq K$ such that with probability at least $1 - \delta$,*

$$\text{ESTR}(\hat{S}) + \min\{1, \alpha\omega \cdot \text{CI}(\hat{S})\} + \varepsilon \geq \text{ESTR}(S^*) + \min\{1, \omega \cdot \text{CI}(S^*)\}, \quad (20)$$

where S^* is the assortment set maximizing the actual objective in Eq. (19)³.

The following lemma shows how $(\alpha, \varepsilon, \delta)$ -approximation algorithms imply an upper bound on the accumulated. It is proved using standard analysis of UCB type algorithms, with the complete proof given in the supplementary material.

³We slightly abuse the notation S^* here following the optimization convention that S^* denotes the optimal solution. Note that S^* is different from S_t^* in (5), where the latter means the assortment that maximizes the expected revenue at time t .

Lemma 7. Suppose an $(\alpha, \varepsilon, \delta)$ -approximation algorithm is used instead of exact optimization in the MLE-UCB policy at each time period t . Then its regret can be upper bounded by

$$\alpha \cdot \text{Regret}^* + \varepsilon T + \delta T^2 + O(1),$$

where Regret^* is the regret upper bound shown by Theorem 1 for Algorithm 1 with exact optimization in Step 6.

In the rest of this section we introduce our proposed approximation algorithm and the approximation guarantee. To highlight the main idea of the approximation algorithm, we only describe how the algorithm operates in the *univariate* ($d = 1$) case, while leaving the general multivariate ($d > 1$) case to the appendix.

Our approximation algorithm can be roughly divided into three steps. In the first step, we use a “discretization” trick to approximate the objective function using “rounded” parameter values. Such rounding motivates the second step, in which we define “reachable states” and present a simple yet computationally expensive brute-force method to enumerate all reachable states, and establish approximation guarantees for such methods. This brute-force method is only presented for illustration purposes and will be replaced by a dynamic programming algorithm proposed in the third step. In particular, a *dynamic programming* algorithm is developed to compute which states are “reachable” in polynomial time.

5.1.1 The discretization trick

In the univariate case, $\{x_{tj}\}$ are scalars and therefore $x_{tj}x_{tj}^\top$ is simply x_{tj}^2 . Let $\Delta > 0$ be a small positive discretization parameter to be specified later. For all $i \in [N]$, define

$$\mu_i := \left\lceil \frac{\widehat{u}_{ti}}{\Delta} \right\rceil \Delta, \quad \alpha_i := \left\lceil \frac{\widehat{u}_{ti}x_{ti}}{\Delta} \right\rceil \Delta, \quad \beta_i := \left\lceil \frac{\widehat{u}_{ti}x_{ti}^2}{\Delta} \right\rceil \Delta, \quad \gamma_i := \left\lceil \frac{\widehat{u}_{ti}r_{ti}}{\Delta} \right\rceil \Delta, \quad (21)$$

where $\lceil a \rceil$ denotes the nearest integer a real number a is rounded into. Intuitively, μ_i is the real number closest to \widehat{u}_{ti} that is an *integer* multiple of the discretization parameter Δ , and similarly for $\alpha_i, \beta_i, \gamma_i$.

The motivation for the definitions of $\{\mu_i, \alpha_i, \beta_i, \gamma_i\}$ is their *sufficiency* in computing the objective function $\text{ESTR}(S) + \min\{1, \omega \cdot \text{CI}(S)\}$. Indeed, for any $S \subseteq [n]$, $|S| \leq K$, define $\mu = \sum_{j \in S} \mu_j$, $\alpha = \sum_{j \in S} \alpha_j$, $\beta = \sum_{j \in S} \beta_j$, $\gamma = \sum_{j \in S} \gamma_j$ and

$$\widehat{\text{ESTR}}(S) := \frac{\gamma}{1 + \mu}, \quad \widehat{\text{CI}}(S) := \max \left\{ 0, \sqrt{\frac{\beta}{1 + \mu} - \left(\frac{\alpha}{1 + \mu} \right)^2} \right\}.$$

Following the definition of $\text{ESTR}(S)$ and $\text{CI}(S)$, it is easy to see that $\widehat{\text{ESTR}}(S) \rightarrow \text{ESTR}(S)$ and $\widehat{\text{CI}}(S) \rightarrow \text{CI}(S)$ as $\Delta \rightarrow 0^+$. The following lemma gives a more precise control of the error between $\widehat{\text{ESTR}}(S)$, $\widehat{\text{CI}}(S)$ and $\text{ESTR}(S)$, $\text{CI}(S)$ using the values of Δ and the maximum utility parameter in S .

Lemma 8. For any $S \subseteq [N]$, $|S| \leq K$, suppose $U = \max_{j \in S} \{1, \widehat{u}_{tj}\}$ and $\Delta = \varepsilon_0 U / K$ for some $\varepsilon_0 > 0$. Suppose also $|x_{tj}| \leq \nu$ for all t, j . Then

$$|\text{ESTR}(S) - \widehat{\text{ESTR}}(S)| \leq 6\varepsilon_0 \quad \text{and} \quad |\text{CI}(S) - \widehat{\text{CI}}(S)| \leq \sqrt{24\varepsilon_0}(1 + \nu), \quad (22)$$

The complete proof of Lemma 8 is relegated to the supplementary material.

5.1.2 Reachable states and a brute-force algorithm

To apply the estimation error bounds in Lemma 8 one needs to first enumerate $q \in [N]$ giving rise to the item in S with the largest utility parameter \hat{u}_{tq} . After such an element q is enumerated, the discretization parameter $\Delta = \epsilon_0 U / K = \epsilon_0 \max\{1, \hat{u}_{tq}\} / K$ can be determined and discretized values $\mu_i, \alpha_i, \beta_i, \gamma_i$ can be computed for all $i \in [N] \setminus \{q\}$. It is also easy to verify that there are at most $O(K/\epsilon)$ possible values of μ_i, γ_i , $O(K\nu/\epsilon)$ possible values of α_i and $O(K\nu^2/\epsilon)$ possible values of β_i (recall that ν is the upper bound of $|x_{tj}|$ for all t and j).

For any $i \in [N] \cup \{0\}$, $k \in [K] \cup \{0\}$ and $\mu, \alpha, \beta, \gamma \geq 0$ being *integer* multiples of Δ , we use a tuple $\zeta_i^k(\mu, \alpha, \beta, \gamma)$ to denote a *state*. Here the indices i and k mean that the assortment $S \subseteq \{1, 2, \dots, i\}$ and $|S| = k$. Clearly there are at most $O(NK^5\nu^3/\epsilon^4)$ different types of states. A state $\zeta_i^k(\mu, \alpha, \beta, \gamma)$ can be either *reachable* or *non-reachable*, as defined below:

Definition 2. Let $q \in [N]$ be the enumerated item with maximal utility parameter and $U = \max\{1, \hat{u}_{tq}\}$, $\Delta = \epsilon_0 U / K$. A state $\zeta_i^k(\mu, \alpha, \beta, \gamma)$ is *reachable* if there exists $S \subseteq [N]$ satisfying the following:

1. $S \subseteq \{1, 2, \dots, i\}$ and $|S| = k$;
2. $\hat{u}_{tj} \leq \hat{u}_{tq}$ for all $j \in S$;
3. if $i \geq q$ then $q \in S$;
4. $\mu = \sum_{j \in S} \mu_j$, $\alpha = \sum_{j \in S} \alpha_j$, $\beta = \sum_{j \in S} \beta_j$ and $\gamma = \sum_{j \in S} \gamma_j$.

On the other hand, a state $\zeta_i^k(\mu, \alpha, \beta, \gamma)$ is *non-reachable* if at least one condition above is violated.

A simple way to find all reachable states is to enumerate all $S \subseteq [N]$, $|S| \leq K$ and verify the three conditions in Definition 2. While such a procedure is clearly computationally intractable, in the next section we will present a dynamic programming approach to compute all reachable states in polynomial time. After all reachable states are computed, enumerate over every $q \in [N]$ and reachable $\zeta_N^k(\cdot, \cdot, \cdot, \cdot)$ for $k \in [K]$ and find \hat{S} that maximizes $\widehat{\text{ESTR}}(\hat{S}) + \min\{1, \omega \cdot \widehat{\text{CI}}(\hat{S})\}$. The following corollary establishes the approximation guarantee for \hat{S} , following Lemma 8.

Corollary 2. Let $\hat{S} \subseteq [N]$, $|\hat{S}| \leq K$ be a subset corresponding to a reachable state $\zeta_N^k(\cdot, \cdot, \cdot, \cdot)$ for some $k \in [K]$, $q \in [N]$, that maximizes $\widehat{\text{ESTR}}(\hat{S}) + \min\{1, \omega \cdot \widehat{\text{CI}}(\hat{S})\}$. Then

$$\widehat{\text{ESTR}}(\hat{S}) + \min\{1, \omega \cdot \widehat{\text{CI}}(\hat{S})\} \geq \max_{S \subseteq [N], |S| \leq K} \text{ESTR}(S) + \min\{1, \omega \cdot \text{CI}(S)\} - (6\epsilon_0 + \omega(1 + \nu)\sqrt{24\epsilon_0}).$$

Corollary 2 follows easily by plugging in the upper bounds of estimation error in Lemma 8. By setting $\epsilon_0 = \min\{\epsilon/12, \epsilon^2/96\omega^2(1 + \nu)^2\}$, the algorithm that produces \hat{S} satisfies $(1, \epsilon, 0)$ -approximation as defined in Definition 1.

5.1.3 A dynamic programming method for computation of reachable states

In this section we describe a *dynamic programming* algorithm to compute reachable states in polynomial time. The dynamic programming algorithm is exact and deterministic, therefore approximation guarantees in Corollary 2 remain valid.

The first step is again to enumerate $q \in [N]$ corresponding to the item in S with the largest utility parameter \widehat{u}_{tq} , and calculating the discretization parameter $\Delta = \epsilon \max\{1, \widehat{u}_{tq}\}/K$. Afterwards, reachable states are computed in an iterative manner, from $i = 0, 1, \dots$ until $i = N$. The initialization is that $\varsigma_0^0(0, 0, 0, 0)$ is reachable. Once a state $\varsigma_i^k(\mu, \alpha, \beta, \gamma)$ is determined to be reachable, the following two states are *potentially* reachable:

$$\varsigma_{i+1}^k(\mu, \alpha, \beta, \gamma) \quad \text{and} \quad \varsigma_{i+1}^{k+1}(\mu + \mu_{i+1}, \alpha + \alpha_{i+1}, \beta + \beta_{i+1}, \gamma + \gamma_{i+1}).$$

The first future state $\varsigma_{i+1}^k(\mu, \alpha, \beta, \gamma)$ corresponds to the case of $i + 1 \notin S$. To determine when such a state is reachable, we review the conditions in Definition 2 and observe that whenever $i + 1 \neq q$, the decision $i + 1 \notin S$ is legal because q must belong to S whenever $i \geq q$ (note that q is the item in S with the largest estimated utility). The second future state $\varsigma_{i+1}^{k+1}(\mu + \mu_{i+1}, \alpha + \alpha_{i+1}, \beta + \beta_{i+1}, \gamma + \gamma_{i+1})$ corresponds to the case of $i + 1 \in S$. Reviewing again the conditions listed in Definition 2, such a state is reachable if $k + 1 \leq K$ (meaning that there is still room to include a new item in S) and $\widehat{u}_{t,i+1} \leq \widehat{u}_{tq}$ (meaning that the new item ($i + 1$) to be included has an estimated utility smaller than \widehat{u}_{tq}). Combining both cases, we arrive at the following updated rule of reachability:

1. If $i + 1 \neq q$, then $\varsigma_{i+1}^k(\mu, \alpha, \beta, \gamma)$ is reachable;
2. If $k < K$ and $\widehat{u}_{t,i+1} \leq \widehat{u}_{tq}$, then $\varsigma_{i+1}^{k+1}(\mu + \mu_{i+1}, \alpha + \alpha_{i+1}, \beta + \beta_{i+1}, \gamma + \gamma_{i+1})$ is reachable.

Algorithms 3 and 2 give pseudo-codes for the proposed dynamic programming approach of computing reachable states and an approximate optimizer of $\widehat{\text{ESTR}}(S) + \min\{1, \omega \cdot \widehat{\text{CI}}(S)\}$.

Finally, we remark on the time complexity of the proposed algorithm. Because the items j we consider in the assortment satisfy $|\widehat{u}_{tj}| \leq U$, $|r_{tj}| \leq 1$, and $|x_{tj}| \leq \nu$, and all $\mu_i, \alpha_i, \beta_i, \gamma_i$ are integral multiples of Δ , we have (1) μ_i and γ_i take at most $O(K\epsilon_0^{-1})$ possible values; (2) α_i takes at most $(K\nu\epsilon_0^{-1})$ possible values; and (3) β_i takes at most $(K\nu^2\epsilon_0^{-1})$ values. Therefore, the total number of states $\varsigma_i^k(\cdot, \cdot, \cdot, \cdot)$ for fixed $i \in [N] \cup \{0\}$, $k \in [K]$ can be upper bounded by $O(K^8\nu^3\epsilon_0^{-4})$. The time complexity of Algorithm 3 is thus upper bounded by $O(K^9N\nu^3\epsilon_0^{-4})$. Alternatively, to achieve $(1, \epsilon, 0)$ -approximation, one may set $\epsilon_0 = \min\{\epsilon/12, \epsilon^2/(96(1+\nu)^2\omega^2)\}$ as suggested by Corollary 2, resulting in a time complexity of $O(K^9N\nu^3 \max\{\epsilon^{-4}, (1+\nu)^8\omega^8\epsilon^{-8}\})$.

This dynamic programming based approximation algorithm can be extended to multivariate feature vector with $d > 1$. The details are presented in Appendix A.

5.2 Greedy swapping heuristics

While the proposed approximation has rigorous approximation guarantees and runs in polynomial time, the large time complexity still prohibits its application to moderately large scale problem instances. In this subsection, we consider a practically efficient greedy swapping heuristic to approximately solve the combinatorial optimization problem in Eq. (19).

At a higher level, the heuristic algorithm is a “local search” method similar to the Lloyd’s algorithm for K-means clustering (Lloyd, 1982), which continuously tries to improve an assortment

Input: $\{\hat{u}_{ti}, r_{ti}, x_{ti}\}_{i=1}^N$, the designated maximum utility item q , and approximation parameter ϵ .

Output: An approximate maximizer \hat{S} of $\text{ESTR}(\hat{S}) + \min\{1, \omega \cdot \text{CI}(\hat{S})\}$

- 1 Initialization: compute $\mu_i, \alpha_i, \beta_i, \gamma_i$ for all $i \in [N]$ as in Eq. (21);
- 2 Declare $\varsigma_1^0(0, 0, 0, 0)$ as reachable;
- 3 **for** $i = 0, 1, \dots, N - 1$ **do**
- 4 **for all reachable states** $\varsigma_{i-1}^k(\mu, \alpha, \beta, \gamma)$ **do**
- 5 **if** $i + 1 \neq q$ **then**
- 6 Declare $\varsigma_{i+1}^k(\mu, \alpha, \beta, \gamma)$ as reachable;
- 7 **end**
- 8 **if** $\hat{u}_{t,i+1} \leq \hat{u}_{tq}$ and $k + 1 \leq K$ **then**
- 9 Declare $\varsigma_{i+1}^{k+1}(\mu + \mu_{i+1}, \alpha + \alpha_{i+1}, \beta + \beta_{i+1}, \gamma + \gamma_{i+1})$ as reachable;
- 10 **end**
- 11 **end**
- 12 **end**
- 13 For all reachable states $\varsigma_N^k(\mu, \alpha, \beta, \gamma)$, trace back the actual assortment $S \subseteq [N]$, $|S| \leq K$ and select the one with the largest $\widehat{\text{ESTR}}(S) + \min\{1, \omega \cdot \widehat{\text{CI}}(S)\}$ as the output \hat{S} .

Algorithm 2: Approximate combinatorial optimization, the univariate ($d = 1$) case, and with the designated maximum utility item.

Input: $\{\hat{u}_{ti}, r_{ti}, x_{ti}\}_{i=1}^N$ and additive approximation parameter ϵ .

Output: An approximate maximizer \hat{S} of $\text{ESTR}(\hat{S}) + \min\{1, \omega \cdot \text{CI}(\hat{S})\}$.

- 1 **for** $i = 1, 2, \dots, N$ **do**
- 2 Invoke Algorithm 2 with parameters $q = i$ and ϵ and denote the returned assortment by \hat{S}_i .
- 3 **end**
- 4 Among $\hat{S}_1, \dots, \hat{S}_N$, select the one with the largest $\widehat{\text{ESTR}}(S) + \min\{1, \omega \cdot \widehat{\text{CI}}(S)\}$ as the output \hat{S} .

Algorithm 3: Approximate combinatorial optimization, the univariate ($d = 1$) case.

solution by considering local swapping/addition/deletions until no further improvements are possible. A pseudo-code description of our heuristic method is given in Algorithm 4.

While the greedy heuristic does not have rigorous guarantees in general, we would like to mention a special case of $\omega = 0$, in which Algorithm 4 does converge to the optimal assortment S maximizing $\text{ESTR}(S) + \min\{1, \omega \cdot \text{CI}(S)\}$ in polynomial time. More specifically, we have the following proposition which is proved in the supplementary material.

Proposition 2. *If $\omega = 0$, then Algorithm 4 terminates in $O(N^4)$ iterations and produces an output S that maximizes $\text{ESTR}(S)$.*

Input: problem parameters $\{\widehat{u}_{ti}, r_{ti}, x_{ti}\}_{i=1}^N$.

Output: approximate maximizer \widehat{S} of $\text{ESTR}(\widehat{S}) + \min\{1, \omega \cdot \text{CI}(\widehat{S})\}$.

- 1 Initialization: select $S \subseteq [N]$, $|S| = K$ uniformly at random;
- 2 **while** $\text{ESTR}(S) + \min\{1, \omega \cdot \text{CI}(S)\}$ *can be improved* **do**
- 3 For every $i \notin S$ and $j \in S$, consider new candidate assortments $S' = S \cup \{i\} \setminus \{j\}$
 (swapping), $S' = S \cup \{i\}$ if $|S| < K$ (addition) and $S' = S \setminus \{j\}$ if $|S| > 1$ (deletion);
- 4 let S'_* be the considered assortments with the largest $\text{ESTR}(S'_*) + \min\{1, \omega \cdot \text{CI}(S'_*)\}$;
- 5 If S can be improved update $S \leftarrow S'_*$;
- 6 **end**

Algorithm 4: A greedy heuristic for combinatorial assortment optimization

6 Numerical studies

In this section, we present numerical results of our proposed MLE-UCB algorithm. We use the greedy swapping heuristics (Algorithm 4) as the subroutine to solve the combinatorial optimization problem in Eq. (19). We will also study the quality of the solution of the greedy swapping heuristics.

Experiment setup. The unknown model parameter $\theta_0 \in \mathbb{R}^d$ is generated as a uniformly random unit d -dimensional vector. The revenue parameters $\{r_{tj}\}$ for $j \in [N]$ are independently and identically generated from the uniform distribution $[0.5, 0.8]$. For the feature vectors $\{v_{tj}\}$, each of them is independently generated as a uniform random vector v such that $\|v\| = 2$ and $v^\top \theta_0 < -0.6$. Here we set an upper bound of -0.6 for the inner product so that the utility parameters $u_{tj} = \exp\{v_{tj}^\top \theta_0\}$ are upper bounded by $\exp(-0.6) \approx 0.55$. We set such an upper bound because if the utility parameters are uniformly large, the optimal assortment is likely to pick very few items, leading to degenerated problem instances. In the implementation of our MLE-UCB algorithm, we simply set $T_0 = \lfloor \sqrt{T} \rfloor$ and $\omega = \sqrt{d \ln(TK)}$.

The greedy swapping heuristics. We first numerically evaluate the solution quality of the greedy swapping heuristic algorithm by focusing on the optimization problem in Eq. (19). We compare the obtained objective values in Eq. (19) to the proposed greedy heuristic and the optimal solution (obtained by brute-force search). Instead of generating purely random instances, we consider more realistic instances generated from a dynamic assortment planning process. In particular, for a given T , we generate a dynamic assortment optimization problem with parameters $N = 10$, $K = 4$ and $d = 5$, and run the MLE-UCB algorithm till the T -th time period. Now the combinatorial optimization problem in Eq. (19) to be solved at the T -th time is kept as one testing instance for the greedy swapping algorithm.

For each $T \in \{50, 200, 800\}$, we generate 1000 such test instances, and compare the solution of the greedy swapping heuristics with the optimal solution obtained by brute-force search in terms of the objective value in Eq. (19). Table 1 shows the relative differences between the two solutions at several percentiles, and the mean relative differences. We can see that the approximation quality of the greedy swapping algorithm has already been desirable when $T = 50$, and becomes even better as T grows.

T	percentile rank					mean relative difference in objective value
	94th	96th	98th	99th	99.5th	
50	0	0.0159	0.0293	0.0393	0.0687	0.00207
200	0	0.0001	0.0040	0.0080	0.0123	0.00024
800	0	0	0	0.0014	0.0037	0.00004

Table 1: relative differences in terms of objective value in Eq. (19) between the greedy swapping algorithm and the optimal solution.




Figure 1: Illustration of the performance of MLE-UCB.

Performance of the MLE-UCB algorithm. In Figure 1a we plot the average regret (i.e. regret/T) of MLE-UCB algorithm with $N = 1000$, $K = 10$, $d = 5$ for the first $T = 10000$ time periods. For each experiment (in both Figure 1a and other figures), we repeat the experiment for 100 times and report the average value. In Figure 1b we compare our algorithm with the UCB algorithm for multinomial logit bandit (MNL-UCB) from Agrawal et al. (2017a) without utilizing the feature information. Since the MNL-UCB algorithm assumes fixed item utilities that do not change over time, in this experiment we randomly generate one feature vector for each of the $N = 1000$ items and this feature vector will be fixed for the entire time span. We can observe that our MLE-UCB algorithm performs much better than MNL-UCB, which suggests the importance of taking the advantage of the contextual information.

Impact of the dimension size d . We study how the dimension of the feature vector impacts the performance of our MLE-UCB algorithm. We fix $N = 1000$ and $K = 10$ and test our algorithm for dimension sizes in $5, 7, 9, 11, \dots, 25$. In Figure 2, we report the average regret at times $T \in \{4000, 6000, 8000, 10000\}$. We can see that the average regret increases approximately linearly with d . This phenomenon matches the linear dependency on d of the main term of the regret Eq. (6) of the MLE-UCB.




Figure 2: Average regret of MLE-UCB for various d 's.




Figure 3: Average regret of MLE-UCB for various N 's.

Impact of the number of items N . We compare the performance of our MLE-UCB algorithm for the varying number of items N . We fix $K = 10$ and $d = 5$, and test MLE-UCB for $N \in \{1000, 2000, 3000, 4000\}$. In Figure 3, we report the average regret for the first $T = 10000$ time periods. We observe that the regret of the algorithm is almost not affected by a bigger N . This confirms the fact that the regret Eq. (6) of MLE-UCB is totally independent of N .

7 Conclusion and future directions

In this paper, we study the dynamic assortment planning problem under a contextual MNL model, which incorporates rich feature information into choice modeling. We propose an upper confidence bound (UCB) algorithm based on the local MLE that simultaneously learns the underlying coefficient and makes the decision on the assortment selection. We establish both the upper and lower bounds of the regret. Moreover, we develop an approximation algorithm and a greedy heuristic for solving the key optimization problem in our UCB algorithm.

There are a few possibilities for future work. Technically, there is still a gap of $1/K$ between our upper and lower bounds on regret. Although the cardinality constraint of an assortment K is usually small in practice, it is still a technically interesting question to close this gap. Second, introducing contextual information into choice model is a natural idea for many online applications. This paper explores the standard MNL model, and it would be interesting to extend this work to contextual nested logit and other popular choice models. Finally, it is interesting to incorporate other operational considerations into the model, such as prices or inventory constraints.

A Multivariate approximation algorithm

In this appendix we describe an approximation algorithm for the combinatorial optimization problem studied in Sec. 5.1 for the general multivariate ($d > 1$) case. The multivariate case is dealt with by *randomized* reductions to several univariate problem instances.

More specifically, for any $y \in \mathbb{R}^d$, $\|y\|_2 = 1$, a univariate problem instance can be constructed

Input: $\{\hat{u}_{ti}, r_{ti}, x_{ti}\}_{i=1}^N$, multiplicative approximation parameter α , additive approximation parameter ϵ , repetition $L \in \mathbb{N}$.

Output: An approximate maximizer \hat{S} of $\text{ESTR}(\hat{S}) + \min\{1, \omega \cdot \text{CI}(\hat{S})\}$.

- 1 Generalize L vectors $y^{(1)}, \dots, y^{(L)} \in \mathbb{R}^d$ independently and uniformly from the unit sphere;
- 2 **for** $\ell = 1, 2, \dots, L$ **do**
- 3 Replace each x_{ti} with $\langle x_{ti}, y^{(\ell)} \rangle$;
- 4 Invoke Algorithm 3 on the reduced univariate problem instance, and let $\hat{S}^{(\ell)}$ be the output;
- 5 **end**
- 6 Output $\hat{S}^{(\ell)}$ that maximizes $\text{ESTR}(\hat{S}^{(\ell)}) + \min\{1, \alpha\omega \cdot \text{CI}(\hat{S}^{(\ell)})\}$.

Algorithm 5: Approximate combinatorial optimization, the multivariate ($d > 1$) case

by replacing every occurrences of x_{ti} with $x_{ti}^\top y$. The univariate approximation Algorithm 3 is then invoked on L independent univariate problem instances, each corresponding to a y vector sampled uniformly at random from the d -dimensional unit sphere. The L output maximizers \hat{S} of Algorithm 3 are then compared against each other and the one leading to the largest value of $\text{ESTR}(\hat{S}) + \min\{1, \alpha\omega \cdot \text{CI}(\hat{S})\}$ is selected, where α is the preset multiplicative approximation parameter. A pseudo-code description is given in Algorithm 5.

A.1 Approximation guarantees

The approximation performance of Algorithm 5 can be analyzed based on the following observation: if y is close to y^* , the leading eigenvector of

$$\frac{\sum_{j \in S^*} \hat{u}_{tj} x_{tj} x_{tj}^\top}{1 + \sum_{j \in S^*} \hat{u}_{tj}} - \left(\frac{\sum_{j \in S^*} \hat{u}_{tj} x_{tj}}{1 + \sum_{j \in S^*} \hat{u}_{tj}} \right) \left(\frac{\sum_{j \in S^*} \hat{u}_{tj} x_{tj}}{1 + \sum_{j \in S^*} \hat{u}_{tj}} \right)^\top,$$

where S^* is the exact maximizer of Eq. (19), then the reduction to a univariate problem instance $x_{tj} \mapsto x_{tj}^\top y$ does not lose much accuracy. More specifically, we have the following lemma:

Lemma 9. *Suppose there exists $\ell \in [L]$ such that $\langle y^{(\ell)}, y^* \rangle \geq 1/\alpha$ for some $\alpha \geq 1$ in Algorithm 5, then $\text{ESTR}(\hat{S}^{(\ell)}) + \min\{1, \alpha\omega \cdot \text{CI}(\hat{S}^{(\ell)})\} + \epsilon \geq \text{ESTR}(S^*) + \min\{1, \omega \cdot \text{CI}(S^*)\}$, where $\epsilon > 0$ is the approximation parameter of the univariate problem instances.*

Lemma 9 is proved in the supplementary material using elementary linear algebra. At a higher level, Lemma 9 shows that when the sampled vector $y^{(\ell)}$ is close to the underlying leading eigenvector y^* (in the sense that the inner product between $y^{(\ell)}$ and y^* is large), the produced subset $\hat{S}^{(\ell)}$ will have good performance in maximizing the objective function $\text{ESTR}(S) + \min\{1, \omega \cdot \text{CI}(S)\}$.

The following proposition additionally gives the proximity between a random y and y^* .

Proposition 3. *Assume that $d \geq 2$. Let $y^* \in \mathbb{R}^d$, $\|y^*\|_2 = 1$ be fixed and y be sampled uniformly at random from the unit d -dimensional sphere. Then*

$$\Pr[\langle y, y^* \rangle \geq 1/\sqrt{d}] = \Omega(1) \quad \text{and} \quad \Pr[\langle y, y^* \rangle \geq 1/2] = \exp\{-O(d)\}.$$

Proposition 3 is again proved in the supplementary material, using isotropy of y and classical concentration inequalities.

Combining Lemma 9 and Proposition 3 we can give some recommendations on the choice of L in Algorithm 5, which is the number of random $y^{(\ell)}$ vectors sampled. First, if $L \asymp \log(1/\delta)$ initializations are taken, then with probability $1 - \delta$ Lemma 9 is satisfied with $\alpha = \sqrt{d}$, yielding a $(\sqrt{d}, \epsilon, \delta)$ -approximation. Additionally, if $L \asymp e^{O(d)} \log(1/\delta)$ initializations are taken, then with probability $1 - \delta$ Lemma 9 is satisfied with $\alpha = 2$, yielding a $(2, \epsilon, \delta)$ -approximation.

A.2 Time complexity analysis

To achieve a $(\sqrt{d}, \epsilon, \delta)$ -approximation L is set to $L \asymp \log(1/\delta)$ and the overall running time of Algorithm 5 is $O(K^9 N \nu^3 \max\{\epsilon^{-4}, (1+\nu)^8 \omega^8 \epsilon^{-8}\} \log \delta^{-1})$. To achieve a $(2, \epsilon, \delta)$ -approximation L is set to $L \asymp e^{O(d)} \log(1/\delta)$ and the overall running time of Algorithm 5 is $e^{O(d)} K^9 N \nu^3 \max\{\epsilon^{-4}, (1+\nu)^8 \omega^8 \epsilon^{-8}\}$.

Now we use Algorithm 5 to solve the combinatorial optimization problem in Step 6 of Algorithm 1 and examine the cumulative regret. If we let Algorithm 5 achieve to $(\sqrt{d}, \epsilon, \delta)$ -approximation guarantee with $\epsilon = T^{-1/2}$ and $\delta = T^{-2}$, the computational time complexity at each time slot will be $\tilde{O}(K^9 N \nu^3 (1+\nu)^8 d^4 T^4)$,⁴ and the cumulative regret will be upper bounded by $O(\sqrt{d}) \cdot \text{Regret}^*$. If we let Algorithm 5 to achieve $(1/2, \epsilon, \delta)$ -approximation guarantee with $\epsilon = T^{-1/2}$ and $\delta = T^{-2}$, the computational time complexity at each time slot will be $e^{O(d)} \cdot \tilde{O}(K^9 N \nu^3 (1+\nu)^8 d^4 T^4)$, and the cumulative regret will be upper bounded by $O(1) \cdot \text{Regret}^*$.

Acknowledgement : The authors would like to thank Vineet Goyal for helpful discussions, and Zikai Xiong for helping with the numerical studies.

References

- Abbasi-Yadkori, Y., Pál, D., & Szepesvári, C. (2011). Improved algorithms for linear stochastic bandits. In *Proceedings of the Advances in Neural Information Processing Systems (NIPS)*.
- Agrawal, S., Avadhanula, V., Goyal, V., & Zeevi, A. (2017a). MNL-bandit: A dynamic learning approach to assortment selection. *arXiv preprint arXiv:1706.03880*.
- Agrawal, S., Avadhanula, V., Goyal, V., & Zeevi, A. (2017b). Thompson sampling for MNL-bandit. In *Proceedings of the Conference on Learning Theory (COLT)*.
- Agrawal, S., & Goyal, N. (2013). Thompson sampling for contextual bandits with linear payoffs. In *Proceedings of the International Conference on Machine Learning (ICML)*.
- Caro, F., & Gallien, J. (2007). Dynamic Assortment with Demand Learning for Seasonal Consumer Goods. *Management Science*, 53(2), 276–292.
- Chen, X., Ma, W., Simchi-Levi, D., & Xin, L. (2016). Dynamic recommendation at checkout under inventory constraint. Available at SSRN: <https://www.ssrn.com/abstract=2853093>.

⁴A polylogarithmic factor dependent on $T, K, \delta^{-1}, \nu, \rho$ is hidden in the $\tilde{O}(\cdot)$ notation.

- Chen, X., Owen, Z., Pixton, C., & Simchi-Levi, D. (2015). A statistical learning approach to personalization in revenue management. Available at SSRN: <http://ssrn.com/abstract=2579462>.
- Chen, X., & Wang, Y. (2018). A note on tight lower bound for mnl-bandit assortment selection models. *Operations Research Letters*, 46(5), 534–537.
- Chen, X., Wang, Y., & Zhou, Y. (2018). Dynamic assortment selection under nested logit models. *arXiv preprint arXiv:1806.10410*.
- Cheung, W. C., & Simchi-Levi, D. (2017). Thompson sampling for online personalized assortment optimization problems with multinomial logit choice models. Available at SSRN: https://papers.ssrn.com/?abstract_id=3075658.
- Chu, W., Li, L., Reyzin, L., & Schapire, R. (2011). Contextual bandits with linear payoff functions. In *Proceedings of the International Conference on Artificial Intelligence and Statistics (AISTATS)*.
- Dani, V., Hayes, T. P., & Kakade, S. M. (2008). Stochastic Linear Optimization under Bandit Feedback. In *Proceedings of the Annual Conference on Learning Theory (COLT)*.
- Fan, X., Grama, I., & Liu, Q. (2015). Exponential inequalities for martingales with applications. *Electronic Journal of Probability*, 20(1), 1–22.
- Filippi, S., Cappe, O., Garivier, A., & Szepesvári, C. (2010). Parametric bandits: The generalized linear case. In *Proceedings of the Advances in Neural Information Processing Systems (NIPS)*.
- Freedman, D. A. (1975). On tail probabilities for martingales. *The Annals of Probability*, 3(1), 100–118.
- Golrezaei, N., Nazerzadeh, H., & Rusmevichientong, P. (2014). Real-time optimization of personalized assortments. *Management Science*, 60(6), 1532–1551.
- Li, L., Lu, Y., & Zhou, D. (2017). Provably optimal algorithms for generalized linear contextual bandits. In *Proceedings of International Conference on Machine Learning (ICML)*.
- Lloyd, S. (1982). Least squares quantization in pcm. *IEEE Transactions on Information Theory*, 28(2), 129–137.
- McFadden, D. (1973). Conditional logit analysis of qualitative choice behaviour. In P. Zarembka (Ed.) *Frontiers in Econometrics*, (pp. 105–142). New York, NY, USA: Academic Press New York.
- Rusmevichientong, P., Shen, Z.-J., & Shmoys, D. (2010). Dynamic assortment optimization with a multinomial logit choice model and capacity constraint. *Operations Research*, 58(6), 1666–1680.
- Rusmevichientong, P., & Tsitsiklis, J. N. (2010). Linearly parameterized bandits. *Mathematics of Operations Research*, 35(2), 395–411.
- Saure, D., & Zeevi, A. (2013). Optimal dynamic assortment planning with demand learning. *Manufacturing & Service Operations Management*, 15(3), 387–404.

- Tsybakov, A. B. (2009). *Introduction to nonparametric estimation*. Springer Series in Statistics. Springer, New York.
- van de Geer, S. A. (2000). *Empirical Processes in M-Estimation*. Cambridge University Press.
- Van der Vaart, A. W. (2000). *Asymptotic statistics*. Cambridge university press.
- Vershynin, R. (2012). How close is the sample covariance matrix to the actual covariance matrix? *Journal of Theoretical Probability*, 25(3), 655–686.
- Wang, Y., Chen, X., & Zhou, Y. (2018). Near-optimal policies for dynamic multinomial logit assortment selection models. In *Advances in Neural Information Processing Systems (NeurIPS)*.

Supplementary Material for: Dynamic Assortment Optimization with Changing Contextual Information

Xi Chen ^{¶1}, Yining Wang², and Yuan Zhou³

¹Stern School of Business, New York University

²Machine Learning Department, Carnegie Mellon University

³Computer Science Department, Indiana University at Bloomington

This supplementary material provides detailed proofs for technical lemmas whose proofs are omitted in the main text.

A Proofs of technical lemmas for Theorem 1 (upper bound)

A.1 Proof of Lemma 1

Lemma 1 (restated). *With probability $1 - \delta$ it holds that*

$$\|\theta^* - \theta_0\|_2 \leq \frac{2}{\kappa} \sqrt{\frac{d + \log(1/\delta)}{\lambda_{\min}(V)}} \quad \text{where } \kappa = \frac{1}{2e(1 + \rho)} \quad \text{and } V = \sum_{t=1}^{T_0} v_{t,i_t} v_{t,i_t}^\top. \quad (\text{S1})$$

Proof. Because the noise in a logistic regression model is clearly centered and sub-Gaussian with parameter at most $1/4$, it only remains to check (Li et al., 2017, Assumption 1), that $\inf_{\|x\|_2 \leq 1, \|\theta - \theta_0\|_2 \leq 1} \sigma'(x^\top \theta) \geq \kappa = 2e(1 + \rho)$ where $\sigma(x) = 1/(1 + e^{-x})$ is the sigmoid link function. Because $\sigma'(x) = \sigma(x)(1 - \sigma(x))$, we have $\sigma'(x^\top \theta) = \wp_\theta(1 - \wp_\theta) \geq 0.5\wp_\theta$ where $\wp_\theta = \min\{p_\theta(1), 1 - p_\theta(1)\}$ and $p_\theta(1) = \sigma(x^\top \theta) = 1/(1 + \exp\{-x^\top \theta\})$. By (A2), we know that $\wp_{\theta_0} \geq 1/(1 + \rho)$. Subsequently, for any $\|x\|_2 \leq 1$ and $\|\theta - \theta_0\|_2 \leq 1$, we have

$$\wp_\theta = \frac{1}{1 + \exp\{-x^\top \theta\}} = \frac{1}{1 + \exp\{-x^\top(\theta - \theta_0)\} \exp\{-x^\top \theta_0\}} \geq \frac{1}{e} \frac{1}{1 + \exp\{x^\top \theta_0\}} \geq \frac{1}{e(1 + \rho)}.$$

Lemma 1 is then an immediate consequence of (Li et al., 2017, Eq. (18)). \square

* Author names listed in alphabetical order.

¶ Author names listed in alphabetical order.

A.2 Proof of Corollary 1

Corollary 1 (restated). *There exists a universal constant $C_0 > 0$ such that for arbitrary $\tau \in (0, 1/2]$, if $T_0 \geq C_0 \max\{\nu^2 d \log T / \lambda_0^2, \rho^2(d + \log T) / (\tau^2 \lambda_0)\}$ then with probability $1 - O(T^{-1})$, $\|\theta^* - \theta_0\|_2 \leq \tau$.*

Proof. Denote $\Lambda := \mathbb{E}_\mu x x^\top$ and $\widehat{\Lambda} := V/T_0 = \frac{1}{T_0} \sum_{t=1}^{T_0} x_{t,i_t} x_{t,i_t}^\top$. Clearly $\mathbb{E}\widehat{\Lambda} = \Lambda$. In addition, because $\|v_{tj}\|_2 \leq \nu$ almost surely, v_{tj} are sub-Gaussian random variables with parameter ν^2 . By standard concentration inequalities (see, e.g., (Vershynin, 2012, Proposition 2.1)), we have with probability $1 - O(T^{-2})$ that $\|\widehat{\Lambda} - \Lambda\|_{\text{op}} \lesssim \nu \sqrt{\frac{d \log T}{T_0}}$. Hence, if $T_0 \geq C_0 \nu^2 d \log T / \lambda_0^2$ for some sufficiently large universal constant C_0 , we have $\|\widehat{\Lambda} - \Lambda\|_{\text{op}} \leq 0.5\lambda_0 = \lambda_{\min}(\Lambda)$ and therefore $\lambda_{\min}(V) = T_0 \lambda_{\min}(\widehat{\Lambda}) \geq 0.5T_0 \lambda_0$. The corollary then immediately follows Lemma 1. \square

A.3 Proof of Lemma 2

Lemma 2 (restated). *Suppose $\tau \leq 1/\sqrt{8\rho\nu^2 K^2}$. Then there exists a universal constant $C > 0$ such that with probability $1 - O(T^{-1})$ the following holds uniformly over all $t = T_0, \dots, T-1$:*

$$(\widehat{\theta}_t - \theta_0)^\top I_t(\theta_0)(\widehat{\theta}_t - \theta_0) \leq C \cdot d \log(\rho\nu TK). \quad (\text{S2})$$

Proof. For any $\theta \in \mathbb{R}^d$ define

$$f_{t'}(\theta) := \mathbb{E}_{\theta_0, t'} \left[\log \frac{p_{\theta, t'}(j|S_{t'})}{p_{\theta_0, t'}(j|S_{t'})} \right] = \sum_{j \in S_{t'} \cup \{0\}} p_{\theta_0, t'}(j|S_{t'}) \log \frac{p_{\theta, t'}(j|S_{t'})}{p_{\theta_0, t'}(j|S_{t'})}.$$

By simple algebra calculations, the first and second order derivatives of $f_{t'}$ with respect to θ can be computed as

$$\nabla_\theta f_{t'}(\theta) = \mathbb{E}_{\theta_0, t'}[v_{t'j}] - \mathbb{E}_{\theta, t'}[v_{t'j}]; \quad (\text{S3})$$

$$\begin{aligned} \nabla_\theta^2 f_{t'}(\theta) &= -\mathbb{E}_{\theta_0, t'}[v_{t'j} v_{t'j}^\top] + \{\mathbb{E}_{\theta_0, t'} v_{t'j}\} \{\mathbb{E}_{\theta, t'} v_{t'j}\}^\top \\ &\quad + \{\mathbb{E}_{\theta, t'} v_{t'j}\} \{\mathbb{E}_{\theta_0, t'} v_{t'j}\}^\top - \{\mathbb{E}_{\theta, t'} v_{t'j}\} \{\mathbb{E}_{\theta, t'} v_{t'j}\}^\top. \end{aligned} \quad (\text{S4})$$

In the rest of the section we drop the subscript in ∇_θ , ∇_θ^2 , and the ∇ , ∇^2 notations should always be understood as with respect to θ .

Define $F_t(\theta) := \sum_{t'=1}^t f_{t'}(\theta)$. It is easy to verify that $-F_t(\theta)$ is the Kullback-Leibler divergence between the conditional distribution of (i_1, \dots, i_t) parameterized by θ and θ_0 , respectively. Therefore, $F_t(\theta)$ is always non-positive. Note also that $F_t(\theta_0) = 0$, $\nabla F_t(\theta_0) = 0$, $\nabla^2 f_{t'}(\theta) = -M_{t'}(\theta)$ and $\nabla^2 F_t(\theta) \equiv -I_t(\theta)$. By Taylor expansion with Lagrangian remainder, there exists $\bar{\theta}_t = \alpha\theta_0 + (1 - \alpha)\widehat{\theta}_t$ for some $\alpha \in (0, 1)$ such that

$$F_t(\widehat{\theta}_t) = -\frac{1}{2}(\widehat{\theta}_t - \theta_0)^\top I_t(\bar{\theta}_t)(\widehat{\theta}_t - \theta_0). \quad (\text{S5})$$

Our next lemma shows that, if $\bar{\theta}_t$ is close to θ_0 (guaranteed by the constraint that $\|\widehat{\theta}_t - \theta^*\|_2 \leq \tau$), then $I_t(\bar{\theta}_t)$ can be *spectrally* lower bounded by $I_t(\theta_0)$. It is proved in the supplementary material.

Lemma 12. Suppose $\tau \leq 1/\sqrt{8\rho\nu^2K^2}$. Then $I_t(\bar{\theta}_t) \succeq \frac{1}{2}I_t(\theta_0)$ for all t .

As a corollary of Lemma 12, we have

$$F_t(\hat{\theta}_t) \leq -\frac{1}{4}(\hat{\theta}_t - \theta_0)^\top I_t(\theta_0)(\hat{\theta}_t - \theta_0). \quad (\text{S6})$$

On the other hand, consider the ‘‘empirical’’ version $\hat{F}_t(\theta) := \sum_{t'=1}^t \hat{f}_{t'}(\theta)$, where

$$\hat{f}_{t'}(\theta) := \log \frac{p_{\theta, t'}(i_{t'} | S_{t'})}{p_{\theta_0, t'}(i_{t'} | S_{t'})}. \quad (\text{S7})$$

It is easy to verify that $\hat{F}_t(\theta_0) = 0$ remains true; in addition, for any fixed $\theta \in \mathbb{R}^d$, $\{\hat{F}_t(\theta)\}_t$ forms a martingale¹ and satisfies $\mathbb{E}\hat{F}_t(\theta) = F_t(\theta)$ for all t . This leads to our following lemma, which upper bounds the uniform convergence of $\hat{F}_t(\theta)$ towards $F_t(\theta)$ for all $\|\theta - \theta_0\| \leq 2\tau$.

Lemma 13. Suppose $\tau \leq 1/\sqrt{8\rho^2\nu^2K^2}$. Then there exists a universal constant $C > 0$ such that with probability $1 - O(T^{-1})$ the following holds uniformly for all $t \in \{T_0 + 1, \dots, T\}$ and $\|\theta - \theta_0\|_2 \leq 2\tau$:

$$|\hat{F}_t(\theta) - F_t(\theta)| \leq C \left[d \log(\rho\nu TK) + \sqrt{|F_t(\theta)| d \log(\rho\nu TK)} \right]. \quad (\text{S8})$$

Lemma 13 can be proved by using a standard ε -net argument. Since the complete proof is quite involved, we defer it to the supplementary material.

We are now ready to prove Lemma 2. By Eq. (S8) and the fact that $\hat{F}_t(\hat{\theta}_t) \leq 0 \leq F_t(\hat{\theta}_t)$, we have

$$|F_t(\hat{\theta}_t)| \leq |\hat{F}_t(\hat{\theta}_t) - F_t(\hat{\theta}_t)| \lesssim d \log(\rho\nu TK) + \sqrt{|F_t(\hat{\theta}_t)| d \log(\rho\nu TK)}. \quad (\text{S9})$$

Subsequently,

$$|F_t(\hat{\theta}_t)| \lesssim d \log(\rho\nu NT). \quad (\text{S10})$$

In addition, because $F_t(\hat{\theta}_t) \leq 0$, by Eq. (S6) we have

$$-\frac{1}{2}(\hat{\theta}_t - \theta_0)^\top I_t(\theta_0)(\hat{\theta}_t - \theta_0) \geq F_t(\hat{\theta}_t) \geq d \log(\rho\nu TK). \quad (\text{S11})$$

Lemma 2 is thus proved. \square

Proof of Lemma 12

Lemma 12 (restated). Suppose $\tau \leq 1/\sqrt{8\rho\nu^2K^2}$. Then $I_t(\bar{\theta}_t) \succeq \frac{1}{2}I_t(\theta_0)$ for all t .

Proof. Because $\hat{\theta}_t$ is a feasible solution of the local MLE, we know $\|\hat{\theta}_t - \theta^*\|_2 \leq \tau$. Also by Corollary 1 we know that $\|\theta^* - \theta_0\|_2 \leq \tau$ with high probability. By triangle inequality and the definition of $\bar{\theta}_t$ we have that $\|\bar{\theta}_t - \theta_0\|_2 \leq 2\tau$.

¹ $\{X_k\}_k$ forms a martingale if $\mathbb{E}[X_{k+1} | X_1, \dots, X_k] = X_k$ for all k .

To prove $I_t(\bar{\theta}_t) \succeq \frac{1}{2}I_t(\theta_0)$ we only need to show that $M_{t'}(\bar{\theta}_t) - M_{t'}(\theta_0) \preceq \frac{1}{2}M_{t'}(\theta_0)$ for all $1 \leq t' \leq t$. This reduces to proving

$$\{\mathbb{E}_{\bar{\theta}_t, t'} v_{t'j} - \mathbb{E}_{\theta_0, t'} v_{t'j}\} \{\mathbb{E}_{\bar{\theta}_t, t'} v_{t'j} - \mathbb{E}_{\theta_0, t'} v_{t'j}\}^\top \preceq \frac{1}{2} \mathbb{E}_{\theta_0, t'} \left[(v_{t'j} - \mathbb{E}_{\theta_0, t'} v_{t'j})(v_{t'j} - \mathbb{E}_{\theta_0, t'} v_{t'j})^\top \right]. \quad (\text{S12})$$

Fix arbitrary $S_{t'} \subseteq [N]$, $|S_{t'}| = J \leq K$ and for convenience denote $x_1, \dots, x_J \in \mathbb{R}^d$ as the feature vectors of items in $S_{t'}$ (i.e., $\{v_{t'j}\}_{j \in S_{t'}}$). Let also $p_{\theta_0}(j)$ and $p_{\bar{\theta}_t}(j)$ be the probability of choosing action $j \in [J]$ corresponding to x_j parameterized by θ_0 or $\bar{\theta}_t$. Define $\bar{x} := \sum_{j=1}^J p_{\theta_0}(j)x_j$, $w_j := x_j - \bar{x}$ and $\delta_j := p_{\bar{\theta}_t}(j) - p_{\theta_0}(j)$. Recall also that $x_0 = 0$ and $w_0 = -\bar{x}$. Eq. (S12) is then equivalent to

$$\left\{ \sum_{j=0}^J \delta_j w_j \right\} \left\{ \sum_{j=0}^J \delta_j w_j \right\}^\top \preceq \frac{1}{2} \sum_{j=0}^J p_{\theta_0}(j) w_j w_j^\top. \quad (\text{S13})$$

Let $L = \text{span}\{w_j\}_{j=0}^J$ and $H \in \mathbb{R}^{L \times d}$ be a whitening matrix such that $H(\sum_j p_{\theta_0}(j)w_j w_j^\top)H^\top = I_{L \times L}$, where $I_{L \times L}$ is the identity matrix of size L . Denote $\tilde{w}_j := Hw_j$. We then have $\sum_{j=0}^J p_{\theta_0}(j)\tilde{w}_j \tilde{w}_j^\top = I_{L \times L}$. Eq. (S13) is then equivalent to

$$\left\| \sum_{j=0}^J \delta_j \tilde{w}_j \right\|_2^2 \leq \frac{1}{2}. \quad (\text{S14})$$

On the other hand, by (A2) we know that $p_{\theta_0}(j) \geq 1/\rho K$ for all j and therefore $\|\tilde{w}_j\|_2 \leq \sqrt{\rho K}$ for all j . Subsequently, we have

$$\left\| \sum_{j=0}^J \delta_j \tilde{w}_j \right\|_2^2 \leq \left(\max_j |\delta_j| \cdot \sum_{j=0}^J \|\tilde{w}_j\|_2 \right)^2 \leq \max_j |\delta_j|^2 \cdot \rho K^2. \quad (\text{S15})$$

Recall that $\delta_i = p_{\bar{\theta}_t}(i) - p_{\theta_0}(i)$ where $p_\theta(i) = \exp\{x_i^\top \theta\} / (1 + \sum_{j \in S_{t'}} \exp\{x_j^\top \theta\})$. Simple algebra yields that $\nabla_\theta p_\theta(i) = p_\theta(i)[x_i - \mathbb{E}_\theta x_j]$, where $\mathbb{E}_\theta x_j = \sum_{j \in S_{t'}} p_\theta(j)x_j$. Using the mean-value theorem, there exists $\tilde{\theta}_t = \tilde{\alpha}\bar{\theta}_t + (1 - \tilde{\alpha})\theta_0$ for some $\tilde{\alpha} \in (0, 1)$ such that

$$\delta_i = \langle \nabla_\theta p_{\tilde{\theta}_t}(i), \hat{\theta}_t - \theta_0 \rangle = p_{\tilde{\theta}_t}(i) \langle x_i - \mathbb{E}_{\tilde{\theta}_t} x_j, \bar{\theta}_t - \theta_0 \rangle. \quad (\text{S16})$$

Because $\|x_{ti}\|_2 \leq \nu$ almost surely for all $t \in [T]$ and $i \in [N]$, we have

$$\max_j |\delta_j|^2 \cdot \rho K^2 \leq 4 \cdot \max_i \|x_i\|_2^2 \cdot \|\bar{\theta}_t - \theta_0\|_2^2 \cdot \rho K^2 \leq 4\rho\nu^2 K^2 \cdot \tau^2. \quad (\text{S17})$$

The lemma is then proved by plugging in the condition on τ . \square

Proof of Lemma 13

Lemma 13 (restated). *Suppose $\tau \leq 1/\sqrt{8d\rho^2\nu^2K^2}$. Then there exists a universal constant $C > 0$ such that with probability $1 - O(T^{-1})$ the following holds uniformly for all $t \in \{T_0 + 1, \dots, T\}$ and $\|\theta - \theta_0\|_2 \leq 2\tau$:*

$$|\hat{F}_t(\theta) - F_t(\theta)| \leq C \left[d \log(\rho\nu TK) + \sqrt{|F_t(\theta)| d \log(\rho\nu TK)} \right]. \quad (\text{S18})$$

Proof. We first consider a fixed $\theta \in \mathbb{R}^d$, $\|\theta - \theta_0\|_2 \leq 2\tau$. Define

$$\mathcal{M} := \max_{t' \leq t} |\widehat{f}_{t'}(\theta)| \quad \text{and} \quad \mathcal{V}^2 := \sum_{t'=1}^t \mathbb{E}_{j \sim \theta_0, t'} \left| \log \frac{p_{\theta, t'}(j|S_{t'})}{p_{\theta_0, t'}(j|S_{t'})} \right|^2. \quad (\text{S19})$$

Using an Azuma-Bernstein type inequality (see, for example, (Fan et al., 2015, Theorem A), (Freedman, 1975, Theorem (1.6))), we have

$$|\widehat{F}_t(\theta) - F_t(\theta)| \lesssim \mathcal{M} \log(1/\delta) + \sqrt{\mathcal{V}^2 \log(1/\delta)} \quad \text{with probability } 1 - \delta. \quad (\text{S20})$$

The following lemma upper bounds \mathcal{M} and \mathcal{V}^2 using $F_t(\theta)$ and the fact that θ is close to θ_0 . It will be proved right after this proof.

Lemma 16. *If $\tau \leq 1/\sqrt{8\rho^2\nu^2K^2}$ then $\mathcal{M} \leq 1$ and $\mathcal{V}^2 \leq 8|F_t(\theta)|$.*

Corollary 4. *Suppose τ satisfies the condition in Lemma 16. Then for any $\|\theta - \theta_0\|_2 \leq 2\tau$,*

$$|\widehat{F}_t(\theta) - F_t(\theta)| \lesssim \log(1/\delta) + \sqrt{|F_t(\theta)| \log(1/\delta)} \quad \text{with probability } 1 - \delta. \quad (\text{S21})$$

Our next step is to construct an ϵ -net over $\{\theta \in \mathbb{R}^d : \|\theta - \theta_0\|_2 \leq 2\tau\}$ and apply union bound on the constructed ϵ -net. This together with a deterministic perturbation argument delivers *uniform* concentration of $\widehat{F}_t(\theta)$ towards $F_t(\theta)$.

For any $\epsilon > 0$, let $\mathcal{H}(\epsilon)$ be a finite covering of $\{\theta \in \mathbb{R}^d : \|\theta - \theta_0\|_2 \leq 2\tau\}$ in $\|\cdot\|_2$ up to precision ϵ . That is, $\sup_{\|\theta - \theta_0\|_2 \leq 2\tau} \min_{\theta' \in \mathcal{H}(\epsilon)} \|\theta - \theta'\|_2 \leq \epsilon$. By standard covering number arguments (e.g., (van de Geer, 2000)), such a finite covering set $\mathcal{H}(\epsilon)$ exists whose size can be upper bounded by $\log |\mathcal{H}(\epsilon)| \lesssim d \log(\tau/\epsilon)$. Subsequently, by Corollary 4 and the union bound, we have with probability $1 - O(T^{-1})$ that

$$|\widehat{F}_t(\theta) - F_t(\theta)| \lesssim d \log(T/\epsilon) + \sqrt{|F_t(\theta)| d \log(T/\epsilon)} \quad \forall T_0 < t \leq T, \theta \in \mathcal{H}(\epsilon). \quad (\text{S22})$$

On the other hand, with probability $1 - O(T^{-1})$ such that Eq. (S16) holds, we have for arbitrary $\|\theta - \theta'\|_2 \leq \epsilon$ that

$$\begin{aligned} |\widehat{F}_t(\theta) - \widehat{F}_t(\theta')| &\leq t \cdot \sup_{t' \leq t, j \in S_{t'} \cup \{0\}} \left| \log \frac{p_{\theta, t'}(j|S_{t'})}{p_{\theta', t'}(j|S_{t'})} \right| \\ &\leq t \cdot \sup_{t' \leq t, j \in S_{t'} \cup \{0\}} \frac{|p_{\theta, t'}(j|S_{t'}) - p_{\theta', t'}(j|S_{t'})|}{p_{\theta', t'}(j|S_{t'})} \end{aligned} \quad (\text{S23})$$

$$\leq 2\rho TK \cdot \sup_{t' \leq t, j \in S_{t'} \cup \{0\}} |p_{\theta, t'}(j|S_{t'}) - p_{\theta', t'}(j|S_{t'})| \quad (\text{S24})$$

$$\leq 2\rho TK \cdot \sup_{t' \leq t, j \in [N]} 4\|v_{t'j}\|_2^2 \cdot \|\theta - \theta'\|_2$$

$$\lesssim \rho TK \cdot \nu^2 \cdot \epsilon. \quad (\text{S25})$$

Here Eq. (S23) holds because $\log(1+x) \leq x$; Eq. (S24) holds because $p_{\theta', t'}(j|S_{t'}) \geq p_{\theta_0, t'}(j|S_{t'}) - |p_{\theta', t'}(j|S_{t'}) - p_{\theta_0, t'}(j|S_{t'})| \geq 1/2\rho K$ thanks to (A2) and Eq. (S17).

Combining Eqs. (S22,S25) and setting $\epsilon \asymp 1/(\rho\nu^2TK)$ we have with probability $1 - O(T^{-1})$ that

$$|\widehat{F}_t(\theta) - F_t(\theta)| \lesssim d \log(\rho\nu TK) + \sqrt{|F_t(\theta)|d \log(\rho\nu TK)} \quad \forall T_0 < t \leq T, \|\theta - \theta_0\|_2 \leq 2\tau, \quad (\text{S26})$$

which is to be demonstrated in Lemma 13. \square

Proof of Lemma 16

Lemma 16 (restated). *If $\tau \leq 1/\sqrt{8\rho^2\nu^2K^2}$ then $\mathcal{M} \leq 1$ and $\mathcal{V}^2 \leq 8|F_t(\theta)|$.*

Proof. We first derive an upper bound for M . By (A2), we know that $p_{\theta_0,t'}(j|S_{t'}) \geq 1/\rho K$ for all j . Also, Eqs. (S16,S17) shows that $|p_{\theta,t'}(j|S_{t'}) - p_{\theta_0,t'}(j|S_{t'})| \leq 4\nu^2 \cdot \tau^2$. If $\tau^2 \leq 1/\sqrt{8\rho\nu^2K}$ we have $|p_{\theta,t'}(j|S_{t'}) - p_{\theta_0,t'}(j|S_{t'})| \leq 0.5p_{\theta_0,t'}(j|S_{t'})$ and therefore $|\widehat{f}_{t'}(\theta)| \leq \log^2 2 \leq 1$.

We next give upper bounds on \mathcal{V}^2 . Fix arbitrary t' , and for notational simplicity let $p_j = p_{\theta_0,t'}(j|S_{t'})$ and $q_j = p_{\theta,t'}(j|S_{t'})$. Because $\log(1+x) \leq x$ for all $x \in (-1, \infty)$, we have

$$\mathbb{E}_{j \sim \theta_0,t'} \left| \log \frac{p_{\theta,t'}(j|S_{t'})}{p_{\theta_0,t'}(j|S_{t'})} \right|^2 = \sum_{j \in S_{t'} \cup \{0\}} p_j \log^2 \left(1 + \frac{q_j - p_j}{p_j} \right) \leq \sum_{j \in S_{t'} \cup \{0\}} \frac{(q_j - p_j)^2}{p_j}. \quad (\text{S27})$$

On the other hand, by Taylor expansion we know that for any $x \in (-1, \infty)$, there exists $\bar{x} \in (0, x)$ such that $\log(1+x) = x - x^2/2(1+\bar{x})^2$. Subsequently,

$$-f_{t'}(\theta) = -\mathbb{E}_{j \sim \theta_0,t'} \left[\log \frac{p_{\theta,t'}(j|S_{t'})}{p_{\theta_0,t'}(j|S_{t'})} \right] = - \sum_{j \in S_{t'} \cup \{0\}} p_j \log \left(1 + \frac{q_j - p_j}{p_j} \right) \quad (\text{S28})$$

$$= - \sum_{j \in S_{t'} \cup \{0\}} p_j \left(\frac{q_j - p_j}{p_j} - \frac{1}{2(1+\bar{\delta}_j)^2} \frac{|q_j - p_j|^2}{p_j^2} \right) \quad (\text{S29})$$

$$\geq \frac{1}{2(1 + \max_j |p_j - q_j|/p_j)^2} \cdot \sum_{j \in S_{t'} \cup \{0\}} \frac{(q_j - p_j)^2}{p_j}. \quad (\text{S30})$$

Here $\bar{\delta}_j \in (0, (q_j - p_j)/p_j)$ and the last inequality holds because $\sum_j p_j = \sum_j q_j = 1$.

By Eqs. (S16) and (S17), we have that $|q_j - p_j|^2 \leq 4\nu^2 \cdot \tau^2$. In addition, (A2) implies that $p_j \geq 1/\rho K$ for all j . Therefore, if $\tau \leq 1/\sqrt{4\rho^2\nu^2K^2}$ we have $|p_j - q_j|/p_j \leq 1$ for all j and hence

$$\mathbb{E}_{j \sim \theta_0,t'} \left| \log \frac{p_{\theta,t'}(j|S_{t'})}{p_{\theta_0,t'}(j|S_{t'})} \right|^2 \leq \sum_{j \in S_{t'} \cup \{0\}} \frac{(q_j - p_j)^2}{p_j} \leq 8|f_{t'}(\theta)|. \quad (\text{S31})$$

Summing over all $t' = 1, \dots, t$ and noting that $f_{t'}(\theta)$ is always non-positive, we complete the proof of Lemma 16. \square

A.4 Proof of Lemma 3

Lemma 3 (restated). *Suppose τ satisfies the condition in Lemma 2. With probability $1 - O(T^{-1})$ the following holds uniformly for all $t > T_0$ and $S \subseteq [N]$, $|S| \leq K$ such that*

1. $\bar{R}_t(S) \geq R_t(S)$;
2. $|\bar{R}_t(S) - R_t(S)| \lesssim \min\{1, \omega \sqrt{\|I_{t-1}^{-1/2}(\theta_0)M_t(\theta_0|S)I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}}\}$.

Proof. Without explicit clarification, all statements are conditioned on the success event in Lemma 2, which occurs with probability $1 - O(T^{-1})$ if τ is sufficiently large and satisfies the condition in Lemma 2.

We present below a key technical lemma in the proof of Lemma 3, which is an upper bound on the absolute value difference between $R_t(S) := \mathbb{E}_{\theta_0,t}[r_{tj}|S]$ and $\hat{R}_t(S) := \mathbb{E}_{\hat{\theta}_{t-1,t}}[r_{tj}|S]$ using $I_{t-1}(\theta_0)$ and $M_t(\theta_0|S)$, where $I_{t-1}(\theta) = \sum_{t'=1}^{t-1} M_{t'}(\theta)$ and $M_{t'}(\theta) = \mathbb{E}_{\theta_0,t'}[v_{t'j}v_{t'j}^\top] - \{\mathbb{E}_{\theta_0,t'}v_{t'j}\}\{\mathbb{E}_{\theta_0,t'}v_{t'j}\}^\top - \{\mathbb{E}_{\theta_0,t'}v_{t'j}\}\{\mathbb{E}_{\theta_0,t'}v_{t'j}\}^\top + \{\mathbb{E}_{\theta_0,t'}v_{t'j}\}\{\mathbb{E}_{\theta_0,t'}v_{t'j}\}^\top$. This key lemma can be regarded as a finite sample version of the celebrated *Delta's method* (e.g., (Van der Vaart, 2000)) used widely in classical statistics to estimate and/or infer a functional of unknown quantities.

Lemma 19. *For all $t > T_0$ and $S \subseteq [N]$, $|S| \leq K$, it holds that $|\hat{R}_t(S) - R_t(S)| \lesssim \sqrt{d \log(\rho\nu TK)} \cdot \sqrt{\|I_{t-1}^{-1/2}(\theta_0)M_t(\theta_0|S)I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}}$, where in \lesssim notation we only hide numerical constants.*

Below we state our proof of Lemma 19, while deferring the proof of some detailed technical lemmas to the supplementary material. Fix $S \subseteq [N]$. We use $\mathfrak{R}_t(\theta) = \mathbb{E}_{\theta,t}[r_{tj}] = [\sum_{j \in S} r_{tj} \exp\{v_{tj}^\top \theta\}] / [1 + \sum_{j \in S} \exp\{v_{tj}^\top \theta\}]$ to denote the expected revenue of assortment S at time t , evaluated using a specific model $\theta \in \mathbb{R}$. Then

$$\begin{aligned} \nabla_{\theta} \mathfrak{R}_t(\theta) &= \frac{\sum_{j \in S} r_{tj} \exp\{v_{tj}^\top \theta\} (1 + \sum_{j \in S} \exp\{v_{tj}^\top \theta\})^2 - (\sum_{j \in S} r_{tj} \exp\{v_{tj}^\top \theta\}) (\sum_{j \in S} \exp\{v_{tj}^\top \theta\})}{(1 + \sum_{j \in S} \exp\{v_{tj}^\top \theta\})^2} \\ &= \mathbb{E}_{\theta,t}[r_{tj}v_{tj}] - \{\mathbb{E}_{\theta,t}r_{tj}\}\{\mathbb{E}_{\theta,t}v_{tj}\}. \end{aligned} \quad (\text{S32})$$

By the mean value theorem, there exists $\tilde{\theta}_{t-1} = \theta_0 + \xi(\hat{\theta}_{t-1} - \theta_0)$ for some $\xi \in (0, 1)$ such that

$$\begin{aligned} |\hat{R}_t(S) - R_t(S)| &= |\mathfrak{R}_t(\hat{\theta}_{t-1}) - \mathfrak{R}_t(\theta_0)| = |\langle \nabla \mathfrak{R}_t(\tilde{\theta}_{t-1}), \hat{\theta}_{t-1} - \theta_0 \rangle| \\ &= \sqrt{(\hat{\theta}_{t-1} - \theta_0)^\top [\nabla \mathfrak{R}_t(\tilde{\theta}_{t-1}) \nabla \mathfrak{R}_t(\tilde{\theta}_{t-1})^\top] (\hat{\theta}_{t-1} - \theta_0)}. \end{aligned} \quad (\text{S33})$$

Recall that $\nabla \mathfrak{R}_t(\tilde{\theta}_{t-1}) = \mathbb{E}_{\tilde{\theta}_{t-1,t}}[r_{tj}v_{tj}] - \{\mathbb{E}_{\tilde{\theta}_{t-1,t}}r_{tj}\}\{\mathbb{E}_{\tilde{\theta}_{t-1,t}}v_{tj}\} = \mathbb{E}_{\tilde{\theta}_{t-1,t}}[(r_{tj} - \mathbb{E}_{\tilde{\theta}_{t-1,t}}r_{tj})(v_{tj} - \mathbb{E}_{\tilde{\theta}_{t-1,t}}v_{tj})]$. Subsequently, by Jensen's inequality and the fact that $r_{tj} \in [0, 1]$ almost surely,

$$\begin{aligned} \nabla \mathfrak{R}_t(\tilde{\theta}_{t-1}) \nabla \mathfrak{R}_t(\tilde{\theta}_{t-1})^\top &\preceq \mathbb{E}_{\tilde{\theta}_{t-1,t}} \left[(r_{tj} - \mathbb{E}_{\tilde{\theta}_{t-1,t}}r_{tj})^2 (v_{tj} - \mathbb{E}_{\tilde{\theta}_{t-1,t}}v_{tj})(v_{tj} - \mathbb{E}_{\tilde{\theta}_{t-1,t}}v_{tj})^\top \right] \\ &\preceq \mathbb{E}_{\tilde{\theta}_{t-1,t}} \left[(v_{tj} - \mathbb{E}_{\tilde{\theta}_{t-1,t}}v_{tj})(v_{tj} - \mathbb{E}_{\tilde{\theta}_{t-1,t}}v_{tj})^\top \right] = \widehat{M}_t(\tilde{\theta}_{t-1}|S). \end{aligned} \quad (\text{S34})$$

Define $\widehat{M}_t(\theta|S) := \mathbb{E}_{\theta,t}[(v_{tj} - \mathbb{E}_{\theta,t}v_{tj})(v_{tj} - \mathbb{E}_{\theta,t}v_{tj})^\top]$, where $S \subseteq [N]$ is the assortment supplied at iteration t . Combining Eqs. (S33, S34) with Lemma 2, we have

$$|\hat{R}_t(S) - R_t(S)| \lesssim \sqrt{d \log(\rho\nu NT)} \cdot \sqrt{\|I_{t-1}(\theta_0)^{-1/2} \widehat{M}_t(\tilde{\theta}_{t-1}|S) I_{t-1}(\theta_0)^{-1/2}\|_{\text{op}}}. \quad (\text{S35})$$

It remains to show that $\widehat{M}_t(\tilde{\theta}_{t-1}|S)$ and $M_t(\theta_0|S)$ are close, for which we first recall the definitions of both quantities:

$$\begin{aligned}\widehat{M}_t(\tilde{\theta}_{t-1}|S) &= \mathbb{E}_{\tilde{\theta}_{t-1,t}} \left[(v_{tj} - \mathbb{E}_{\tilde{\theta}_{t-1,t}} v_{tj})(v_{tj} - \mathbb{E}_{\tilde{\theta}_{t-1,t}} v_{tj})^\top \right]; \\ M_t(\theta_0|S) &= \mathbb{E}_{\theta_0,t} [v_{tj} v_{tj}^\top] - \{\mathbb{E}_{\theta_0,t} v_{tj}\} \{\mathbb{E}_{\theta_0,t} v_{tj}\}^\top = \widehat{M}_t(\theta_0|S).\end{aligned}$$

The next lemma shows that under suitable conditions $\widehat{M}_t(\tilde{\theta}_{t-1}|S)$ is close to $\widehat{M}_t(\theta_0|S) = M_t(\theta_0|S)$, implying that $\frac{1}{4}M_t(\theta_0|S) \preceq \widehat{M}_t(\tilde{\theta}_{t-1}|S) \preceq 4M_t(\theta_0|S)$. It is proved in the supplementary material.

Lemma 20. *Suppose $\tau \leq 1/\sqrt{8\rho^2\nu^2K^2}$. Then $\frac{1}{4}M_t(\theta_0|S) \preceq \widehat{M}_t(\tilde{\theta}_{t-1}|S) \preceq 4M_t(\theta_0|S)$ for all t , S and θ .*

As a consequence of Lemma 20, the right-hand side of Eq. (S35) can be upper bounded by

$$\sqrt{d \log(\rho\nu TK)} \cdot \sqrt{4\|I_{t-1}(\theta_0)^{-1/2}M_t(\theta_0|S)I_{t-1}(\theta_0)^{-1/2}\|_{\text{op}}}.$$

Lemma 19 is thus proved. We are now ready to prove Lemma 3. By Lemma 19, we know that with high probability

$$|\widehat{R}_t(S) - R_t(S)| \lesssim \sqrt{d \log(\rho\nu TK)} \cdot \sqrt{\|I_{t-1}(\theta_0)^{-1/2}M_t(\theta_0|S)I_{t-1}(\theta_0)^{-1/2}\|_{\text{op}}} \quad (\text{S36})$$

In addition, by Lemma 20 and the fact that $\|\widehat{\theta}_{t-1} - \theta_0\|_2 \leq \tau$ thanks to the local MLE formulation, we have $\frac{1}{4}M_t(\theta_0|S) \preceq \widehat{M}_t(\widehat{\theta}_{t-1}|S) \preceq 4M_t(\theta_0|S)$ and subsequently $\frac{1}{4}I_{t-1}(\theta_0) \preceq \widehat{I}_{t-1}(\widehat{\theta}_{t-1}) \preceq 4I_{t-1}(\theta_0)$ because $I_{t-1}(\cdot)$ and $\widehat{I}_{t-1}(\cdot)$ are summations of $M_{t'}(\cdot)$ and $\widehat{M}_{t'}(\cdot)$ terms. Setting $\omega \gtrsim \sqrt{d \log(\rho\nu TK)}$ we proved that $\widehat{R}_t(S) \geq R_t(S)$. The second property of Lemma 3 can be proved similarly, by invoking the spectral similarities between $I_{t-1}(\cdot)$, $M_{t'}(\cdot)$ and $\widehat{I}_{t-1}(\cdot)$, $\widehat{M}_{t'}(\cdot)$. \square

Proof of Lemma 20

Lemma 20 (restated). *Suppose $\tau \leq 1/\sqrt{8\rho^2\nu^2K^2}$. Then $\frac{1}{4}M_t(\theta_0|S) \preceq \widehat{M}_t(\tilde{\theta}_{t-1}|S) \preceq 4M_t(\theta_0|S)$ for all t , S and θ .*

Proof. Define $\overline{M}_t(\theta|S) := \mathbb{E}_{\theta_0,t}[(v_{tj} - \mathbb{E}_{\theta_0,t} v_{tj})(v_{tj} - \mathbb{E}_{\theta_0,t} v_{tj})^\top]$, where only the outermost expectation is replaced by taking with respect to the probability law under θ_0 . Denote also $\tilde{w}_j := v_{tj} - \mathbb{E}_{\theta_0,t} v_{tj}$. Then $\overline{M}_t(\theta|S) = \sum_j p_{\theta_0,t}(j) \tilde{w}_j \tilde{w}_j^\top$ and $\overline{M}_t(\theta|S) - \widehat{M}_t(\theta|S) = \sum_j \delta_j \tilde{w}_j \tilde{w}_j^\top$, where $\delta_j = p_{\theta_0,t}(j) - p_{\theta,t}(j)$. By Eq. (S16) and the fact that $\|v_{ti}\|_2 \leq \nu$, $\|\theta - \theta_0\|_2 \leq \tau$, we have

$$\max_j |\delta_j| \leq \sqrt{4\nu^2 \cdot \tau}. \quad (\text{S37})$$

On the other hand, by (A2) we know that $\min_j p_{\theta_0,t}(j) \geq 1/\rho K$ and therefore

$$\overline{M}_t(\theta|S) = \sum_j p_{\theta_0,t}(j) \tilde{w}_j \tilde{w}_j^\top \succeq \frac{1}{\rho K} \sum_j \tilde{w}_j \tilde{w}_j^\top. \quad (\text{S38})$$

Combining Eqs. (S37,S38) and the fact that $\overline{M}_t(\theta|S) - \widehat{M}_t(\theta|S) = \sum_j \delta_j \widetilde{w}_j \widetilde{w}_j^\top$, we have $\overline{M}_t(\theta|S) - \widehat{M}_t(\theta|S) \preceq \overline{M}_t(\theta|S)/2$ and $\widehat{M}_t(\theta|S) - \overline{M}_t(\theta|S) \preceq \overline{M}_t(\theta|S)/2$, provided that $\tau \leq 1/\sqrt{8\rho^2\nu^2K^2}$. This also implies $\frac{1}{2}\overline{M}_t(\theta|S) \preceq \widehat{M}_t(\theta|S) \preceq 2\overline{M}_t(\theta|S)$.

We next prove that $\frac{1}{2}M_t(\theta_0|S) \preceq \overline{M}_t(\theta|S) \preceq 2M_t(\theta_0|S)$ which, together with $\frac{1}{2}\overline{M}_t(\theta|S) \preceq \widehat{M}_t(\theta|S) \preceq 2\overline{M}_t(\theta|S)$ established in the previous section, implies Lemma 20. Recall the definitions that

$$\begin{aligned} M_t(\theta_0|S) &= \mathbb{E}_{\theta_0,t} \left[(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj})(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj})^\top \right]; \\ \overline{M}_t(\theta|S) &= \mathbb{E}_{\theta_0,t} \left[(v_{tj} - \mathbb{E}_{\theta,t}v_{tj})(v_{tj} - \mathbb{E}_{\theta,t}v_{tj})^\top \right]. \end{aligned}$$

Adding and subtracting $\mathbb{E}_{\theta,t}v_{tj}, \mathbb{E}_{\theta_0,t}v_{tj}$ terms, we have

$$\begin{aligned} &\overline{M}_t(\theta|S) - M_t(\theta_0|S) \\ &= \mathbb{E}_{\theta_0,t} \left[(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj} + \mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj} + \mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})^\top \right] \\ &\quad - \mathbb{E}_{\theta_0,t} \left[(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj})(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj})^\top \right] \\ &= \mathbb{E}_{\theta_0,t} \left[(\mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj})^\top \right] + \mathbb{E}_{\theta_0,t} \left[(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj})(\mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})^\top \right] \\ &\quad + (\mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})(\mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})^\top \\ &= (\mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})(\mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})^\top. \end{aligned}$$

By Eq. (S12) in the proof of Lemma 12, we have that

$$(\mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})(\mathbb{E}_{\theta_0,t}v_{tj} - \mathbb{E}_{\theta,t}v_{tj})^\top \preceq \frac{1}{2}\mathbb{E}_{\theta_0,t}[(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj})(v_{tj} - \mathbb{E}_{\theta_0,t}v_{tj})^\top] = \frac{1}{2}M_t(\theta_0|S)$$

provided that $\tau \leq 1/\sqrt{8\rho^2\nu^2K^2}$, thus implying $\frac{1}{2}M_t(\theta_0|S) \preceq \overline{M}_t(\theta|S) \preceq 2M_t(\theta_0|S)$. \square

A.5 Proof of Lemma 4

Lemma 4 (restated). *It holds that*

$$\sum_{t=T_0+1}^T \min\{1, \|I_{t-1}^{-1/2}(\theta_0)M_t(\theta_0|S_t)I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}^2\} \leq 4 \log \frac{\det I_T(\theta_0)}{\det I_{T_0}(\theta_0)} \lesssim d \log(\lambda_0^{-1}\rho\nu).$$

Proof. Denote $A_t := I_{t-1}^{-1/2}(\theta_0)M_t(\theta_0|S_t)I_{t-1}^{-1/2}(\theta_0)$ as d -dimensional positive semi-definite matrices with eigenvalues sorted as $\sigma_1(A_t) \geq \dots \geq \sigma_d(A_t) \geq 0$. By simple algebra,

$$\begin{aligned} \sum_{t=T_0+1}^T \min\{1, \|I_{t-1}^{-1/2}(\theta_0)M_t(\theta_0|S_t)I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}^2\} &= \sum_{t=T_0+1}^T \min\{1, \sigma_1(A_t)^2\} \\ &\leq \sum_{t=T_0+1}^T 2 \log(1 + \sigma_1(A_t)^2) \leq \sum_{t=T_0+1}^T 4 \log(1 + \sigma_1(A_t)). \end{aligned} \tag{S39}$$

On the other hand, note that $I_t(\theta_0) = I_{t-1}(\theta_0) + M_t(\theta_0|S_t) = I_{t-1}(\theta_0)^{1/2}[I_{d \times d} + A_t]I_{t-1}(\theta_0)^{1/2}$. Hence,

$$\log \det I_t(\theta_0) = \log \det I_{t-1}(\theta_0) + \sum_{j=1}^d \log(1 + \sigma_j(A_t)). \quad (\text{S40})$$

Comparing Eqs. (S39) and (S40), we have

$$\sum_{t=T_0+1}^T \min\{1, \|I_{t-1}^{-1/2}(\theta_0)M_t(\theta_0|S_t)I_{t-1}^{-1/2}(\theta_0)\|_{\text{op}}^2\} \leq 4 \log \frac{\det I_T(\theta_0)}{\det I_{T_0}(\theta_0)}, \quad (\text{S41})$$

which proves the first inequality in Lemma 4.

We next prove the second inequality in Lemma 4. Because assortments have size 1 throughout the pure exploration phase ($t \leq T_0$), we have

$$I_{T_0}(\theta_0) = \sum_{t=1}^{T_0} p_{\theta_0,t}(j_t)(1 - p_{\theta_0,t}(j_t))^2 v_{t,j_t} v_{t,j_t}^\top \geq \frac{1}{(1 + \rho)^3} \cdot \sum_{t=1}^{T_0} v_{t,j_t} v_{t,j_t}^\top, \quad (\text{S42})$$

where the last inequality holds thanks to assumption (A2), which implies $p_{\theta_0,t}(j_t) \in [1/(1 + \rho), \rho/(1 + \rho)]$. In addition, by the proof of Corollary 1, with high probability $\lambda_{\min}(\sum_{t=1}^{T_0} v_{t,j_t} v_{t,j_t}^\top) \geq 0.5T_0\lambda_0$, where $\lambda_0 > 0$ is a parameter specified in assumption (A1). Therefore,

$$\det I_{T_0}(\theta_0) \gtrsim [T_0\lambda_0/\rho^3]^d. \quad (\text{S43})$$

On the other hand, because $\max_{t,j} \|v_{t,j}\|_2 \leq \nu$ we have $I_T(\theta_0) \lesssim T \cdot \nu^2$ and subsequently

$$\det I_T(\theta_0) \lesssim [\nu^2 T]^d. \quad (\text{S44})$$

Combining Eqs. (S43) and (S44) we proved the second inequality in Lemma 4. \square

B Proofs of technical lemmas for Theorem 2 (lower bound)

B.1 Proof of Lemma 5

Lemma 5 (restated). *Suppose $\epsilon \in (0, 1/d\sqrt{d})$ and define $\delta := d/4 - |\tilde{U}_t \cap W|$. Then*

$$R(S_{\theta_W}^*) - R(\tilde{S}_t) \geq \frac{\delta\epsilon}{4K\sqrt{d}}.$$

Proof. Let $v = v_W$ and $\hat{v} = v_{\tilde{U}_t}$ be the corresponding feature vectors. Then

$$\begin{aligned} R(S_{\theta_W}^*) - R(\tilde{S}_t) &= \frac{K \exp\{v^\top \theta_W\}}{1 + K \exp\{v^\top \theta_W\}} - \frac{K \exp\{\hat{v}^\top \theta_W\}}{1 + K \exp\{\hat{v}^\top \theta_W\}} \\ &= \frac{K[\exp\{v^\top \theta_W\} - \exp\{\hat{v}^\top \theta_W\}]}{(1 + K \exp\{v^\top \theta_W\})(1 + K \exp\{\hat{v}^\top \theta_W\})} \\ &\geq \frac{\exp\{v^\top \theta_W\} - \exp\{\hat{v}^\top \theta_W\}}{2Ke}. \end{aligned}$$

Here the last inequality holds because $\max(\exp\{v^\top \theta_W\}, \exp\{\hat{v}^\top \theta_W\}) \leq e$. In addition, by Taylor expansion we know that $1 + x \leq e^x \leq 1 + x + x^2/2$ for all $x \in [0, 1]$. Subsequently,

$$R(S_{\theta_W}^*) - R(\tilde{S}_t) \geq \frac{(v - \hat{v})^\top \theta_W - (\hat{v}^\top \theta_W)^2/2}{2Ke} \geq \frac{\delta\epsilon/\sqrt{d} - (\sqrt{d}\epsilon)^2/2}{2Ke}.$$

Finally, noting that $d\epsilon^2/2 \leq \delta\epsilon/2\sqrt{d}$ provided that $\epsilon \in (0, 1/d\sqrt{d})$, we finish the proof of Lemma 5. \square

B.2 Proof of Lemma 6

Lemma 6 (restated). *For any $W \in \mathcal{W}_{d/4-1}$ and $i \in [d]$, $\text{KL}(P_W \| P_{W \cup \{i\}}) \leq C_{\text{KL}} \cdot \mathbb{E}_W[N_i] \cdot \epsilon^2/d$ for some universal constant $C_{\text{KL}} > 0$.*

Proof. Fix a time t with policy's assortment choice S_t , and define $n_i(S_t) := \sum_{v_U \in S_t} \mathbf{1}\{i \in U\}/K$. Let $\{p_j\}_{j \in S_t \cup \{0\}}$ and $\{q_j\}_{j \in S_t \cup \{0\}}$ be the probabilities of purchasing item j under parameterization θ_W and $\theta_{W \cup \{i\}}$, respectively. Then

$$\text{KL}(P_W(\cdot|S_t) \| P_{W \cup \{i\}}(\cdot|S_t)) = \sum_{j \in S_t \cup \{0\}} p_j \log \frac{q_j}{p_j} \leq \sum_j p_j \frac{p_j - q_j}{q_j} \leq \sum_j \frac{|p_j - q_j|^2}{q_j}, \quad (\text{S45})$$

where the only inequality holds because $\log(1+x) \leq x$ for all $x > -1$. Because $q_j \geq e^{-1}/(1+Ke) \geq 1/(2Ke^2)$ for all $j \in S_t \cup \{0\}$, Eq. (S45) is reduced to

$$\text{KL}(P_W(\cdot|S_t) \| P_{W \cup \{i\}}(\cdot|S_t)) \leq 2e^2 K \cdot \sum_{j \in S_t \cup \{0\}} |p_j - q_j|^2. \quad (\text{S46})$$

We next upper bound $|p_j - q_j|$ separately. First consider $j = 0$. We have

$$\begin{aligned} |p_j - q_j| &= \left| \frac{1}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_W\}} - \frac{1}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_{W \cup \{i\}}\}} \right| \\ &\leq \frac{1}{(1 + K/e)^2} \cdot 2 \sum_{j \in S_t} |v_j^\top (\theta_W - \theta_{W \cup \{i\}})| \\ &\leq \frac{2Kn_i(S_t)\epsilon/\sqrt{d}}{(1 + K/e)^2} \leq \frac{8e^2 n_i(S_t)\epsilon}{K\sqrt{d}}. \end{aligned}$$

Here the first inequality holds because $e^x \leq 1 + 2x$ for all $x \in [0, 1]$.

For $j > 0$ corresponding to $v_j = v_U$ where $i \notin U$, we have

$$\begin{aligned} |p_j - q_j| &= \left| \frac{\exp\{v_U^\top \theta_W\}}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_W\}} - \frac{\exp\{v_U^\top \theta_{W \cup \{i\}}\}}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_{W \cup \{i\}}\}} \right| \\ &\leq \left| \frac{1}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_W\}} - \frac{1}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_{W \cup \{i\}}\}} \right| \\ &\leq \frac{8e^2 n_i(S_t)\epsilon}{K\sqrt{d}}. \end{aligned}$$

Here the first inequality holds because $\exp\{v_U^\top \theta_W\} = \exp\{v_U^\top \theta_{W \cup \{i\}}\} \leq 1$, since $i \notin U$. For $j > 0$ corresponding to $v_j = v_U$ and $i \in U$, we have

$$\begin{aligned} |p_j - q_j| &= \left| \frac{\exp\{v_U^\top \theta_W\}}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_W\}} - \frac{\exp\{v_U^\top \theta_{W \cup \{i\}}\}}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_{W \cup \{i\}}\}} \right| \\ &\leq \exp\{v_U^\top \theta_{W \cup \{i\}}\} \cdot \left| \frac{1}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_W\}} - \frac{1}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_{W \cup \{i\}}\}} \right| \\ &\quad + \left| \exp\{v_U^\top \theta_W\} - \exp\{v_U^\top \theta_{W \cup \{i\}}\} \right| \cdot \left| \frac{1}{1 + \sum_{j \in S_t} \exp\{v_j^\top \theta_W\}} \right| \\ &\leq \frac{8e^2 n_i(S_t) \epsilon}{K\sqrt{d}} + \frac{\epsilon}{\sqrt{d}} \cdot \frac{1}{1 + K/e} \leq \frac{8e^2 n_i(S_t) \epsilon}{K\sqrt{d}} + \frac{2e\epsilon}{K\sqrt{d}}. \end{aligned}$$

Combining all upper bounds on $|p_j - q_j|$ and Eq. (S46), we have

$$\begin{aligned} \text{KL}(P_W(\cdot|S_t) \| P_{W \cup \{i\}}(\cdot|S_t)) &\leq 2e^2 K \cdot \left[\frac{128e^4 n_i(S_t)^2 \epsilon^2}{K^2 d} (1 + K) + K n_i(S_t) \cdot \frac{8e^4 \epsilon^2}{K^2 d} \right] \\ &\lesssim n_i(S_t) \epsilon^2 / d. \end{aligned}$$

Here the last inequality holds because $n_i(S_t) \leq 1$. Note also that $N_i = \sum_{t=1}^T n_i(S_t)$ by definition, and subsequently summing over all $t = 1$ to T we have

$$\text{KL}(P_W \| P_{W \cup \{i\}}) \lesssim \mathbb{E}_W[N_i] \cdot \epsilon^2 / d,$$

which is to be demonstrated. \square

C Proofs of approximation algorithms

C.1 Proof of Lemma 7

Lemma 7 (restated). *Suppose an $(\alpha, \epsilon, \delta)$ -approximation algorithm is used instead of exact optimization in the MLE-UCB policy at each time period t . Then its regret can be upper bounded by*

$$\alpha \cdot \text{Regret}^* + \epsilon T + \delta T^2 + O(1),$$

where Regret^* is the regret upper bound shown by Theorem 1 for Algorithm 1 with exact optimization in Step 6.

Proof. By union bound, we know the approximation guarantee in Eq. (20) for all t with probability at least $1 - \delta T$. In the event of failure, the accumulated regret is upper bounded by T almost surely, because the regret incurred by each time period t is at most 1. This gives rise to the δT^2 term in Lemma 7, and in the rest of the proof we shall assume Eq. (20) holds for all t .

Let S_t^* be the solution to the exact optimization problem in Step 6 of Algorithm 1, $S_t^\#$ be the assortment with the optimal revenue the same step, and \hat{S}_t be the solution by an $(\alpha, \epsilon, \delta)$ -approximation algorithm.

For each $t > T_0$, we bound the expected regret incurred at time t by

$$\begin{aligned}
& \text{ESTR}(S_t^\#) - \text{ESTR}(S_t) \\
&= \left(\text{ESTR}(S_t^\#) + \min\{1 + \omega \cdot \text{CI}(S_t^\#)\} \right) - \left(\text{ESTR}(S_t^*) + \min\{1 + \omega \cdot \text{CI}(S_t^*)\} \right) \\
&\quad + \left(\text{ESTR}(S_t^*) + \min\{1 + \omega \cdot \text{CI}(S_t^*)\} \right) - \left(\text{ESTR}(S_t) + \min\{1 + \alpha\omega \cdot \text{CI}(S_t)\} - \varepsilon \right) \\
&\quad + \left(\varepsilon + \min\{1 + \alpha\omega \cdot \text{CI}(S_t)\} - \min\{1 + \omega \cdot \text{CI}(S_t^\#)\} \right) \\
&\leq \varepsilon + \min\{1 + \alpha\omega \cdot \text{CI}(S_t)\} - \min\{1 + \omega \cdot \text{CI}(S_t^\#)\} \leq \varepsilon + \min\{1 + \alpha\omega \cdot \text{CI}(S_t)\}.
\end{aligned}$$

Therefore, the total expected regret is bounded by

$$T_0 + \sum_{t=T_0+1}^T (\varepsilon + \min\{1 + \alpha\omega \cdot \text{CI}(S_t)\}) \leq \varepsilon T + T_0 + \alpha \sum_{t=T_0+1}^T \min\{1 + \omega \cdot \text{CI}(S_t)\},$$

which, by the same analysis in Section 3.2.4, can be bounded by $\alpha \cdot \text{Regret}^* + \varepsilon T$. \square

C.2 Proof of Lemma 8

Lemma 8 (restated). *For any $S \subseteq [N]$, $|S| \leq K$, suppose $U = \max_{j \in S} \{1, \hat{u}_{tj}\}$ and $\Delta = \varepsilon_0 U / K$ for some $\varepsilon_0 > 0$. Suppose also $|x_{tj}| \leq \nu$ for all t, j . Then*

$$|\text{ESTR}(S) - \widehat{\text{ESTR}}(S)| \leq 6\varepsilon_0 \quad \text{and} \quad |\text{CI}(S) - \widehat{\text{CI}}(S)| \leq \sqrt{24\varepsilon_0}(1 + \nu), \quad (\text{S47})$$

Proof. We first prove the upper bound on $|\text{ESTR}(S) - \widehat{\text{ESTR}}(S)|$, which is

$$\left| \frac{\sum_{i \in S} \hat{u}_{ti} r_{ti}}{1 + \sum_{i \in S} \hat{u}_{ti}} - \frac{\sum_{i \in S} \gamma_i}{1 + \sum_{i \in S} \mu_i} \right| \leq 6\varepsilon_0, \quad (\text{S48})$$

where $\mu_i = [\hat{u}_{ti} / \Delta] \cdot \Delta$, $\gamma_i = [\hat{u}_{ti} r_{ti} / \Delta] \cdot \Delta$.

Denote $A := \sum_{i \in S} \hat{u}_{ti} r_{ti}$ and $B := 1 + \sum_{i \in S} \hat{u}_{ti}$. Because $r_{ti} \leq 1$, we have $A \leq B$. Let also $\tau_1 := \sum_{i \in S} \gamma_i - A$ and $\tau_2 := 1 + \sum_{i \in S} \mu_i - B$. Because $\max\{|\gamma_i - \hat{u}_{ti} r_{ti}|, |\mu_i - \hat{u}_{ti}|\} \leq \Delta$, we have $\max\{|\tau_1|, |\tau_2|\} \leq \Delta \cdot K$. Subsequently,

$$\begin{aligned}
\left| \frac{\sum_{i \in S} \hat{u}_{ti} r_{ti}}{1 + \sum_{i \in S} \hat{u}_{ti}} - \frac{\sum_{i \in S} \gamma_i}{1 + \sum_{i \in S} \mu_i} \right| &= \left| \frac{A}{B} - \frac{A + \tau_1}{B + \tau_2} \right| = \left| \frac{A\tau_2 - B\tau_1}{B(B + \tau_2)} \right| = \left| \frac{A\tau_2 - B\tau_2 + B\tau_2 - B\tau_1}{B(B + \tau_2)} \right| \\
&\leq \left| \frac{(A - B)\tau_2}{B(B + \tau_2)} \right| + \frac{|\tau_1| + |\tau_2|}{B - |\tau_2|} \leq \frac{|\tau_1| + 2|\tau_2|}{B - |\tau_2|},
\end{aligned}$$

where the last inequality holds because $A \leq B$. Using $B = 1 + \sum_{i \in S} \hat{u}_{ti} \geq 1 + \hat{u}_{tq} \geq U$ (since $q \in S$ and $U = \max\{1, u_{tq}\}$, and $\max\{|\tau_1|, |\tau_2|\} \leq \Delta \cdot K = \varepsilon_0 U$, we have

$$\frac{|\tau_1| + 2|\tau_2|}{B - |\tau_2|} \leq \frac{3\varepsilon_0 U}{U - \varepsilon_0 U} \leq 6\varepsilon_0,$$

provided that $\varepsilon_0 \in (0, 1/2]$. Eq. (S48) is thus proved.

We next prove the upper bound on $|\text{CI}(S) - \widehat{\text{CI}}(S)|$, which is

$$\left| \sqrt{\frac{\sum_{i \in S} \widehat{u}_{ti} x_{ti}^2}{1 + \sum_{i \in S} \widehat{u}_{ti}} - \left(\frac{\sum_{i \in S} \widehat{u}_{ti} x_{ti}}{1 + \sum_{i \in S} \widehat{u}_{ti}} \right)^2} - \sqrt{\frac{\sum_{i \in S} \beta_i}{1 + \sum_{i \in S} \mu_i} - \left(\frac{\sum_{i \in S} \alpha_i}{1 + \sum_{i \in S} u_i} \right)^2} \right| \leq \sqrt{24\epsilon_0}(1 + \nu), \quad (\text{S49})$$

where $\sqrt{\cdot} = \sqrt{\max\{0, \cdot\}}$, $\mu_i = [\widehat{u}_{ti}/\Delta] \cdot \Delta$, $\alpha_i = [\widehat{u}_{ti} x_{ti}/\Delta] \cdot \Delta$, $\beta_i = [\widehat{u}_{ti} x_{ti}^2/\Delta] \cdot \Delta$.

Denote $C := \frac{\sum_{i \in S} \widehat{u}_{ti} x_{ti}}{1 + \sum_{i \in S} \widehat{u}_{ti}}$ and $D := \frac{\sum_{i \in S} \widehat{u}_{ti} x_{ti}^2}{1 + \sum_{i \in S} \widehat{u}_{ti}}$. Because $|x_{ti}| \leq \nu$ for all t and i , we have $C \in [-\nu, \nu]$ and $D \in [0, \nu^2]$. Denote also $\tau_3 := \frac{\sum_{i \in S} \alpha_i}{1 + \sum_{i \in S} \mu_i} - C$ and $\tau_4 := \frac{\sum_{i \in S} \beta_i}{1 + \sum_{i \in S} \mu_i} - D$. Using the same analysis as in the proof of Eq. (S48), we have $|\tau_3| \leq 6\epsilon_0(1 + \nu)$ and $|\tau_4| \leq 6\epsilon_0(1 + \nu^2)$.

With the definitions of C , D , τ_3 and τ_4 , the left-hand side of Eq. (S49) can be re-written as

$$\left| \sqrt{D - C^2} - \sqrt{(D + \tau_4) - (C + \tau_3)^2} \right|. \quad (\text{S50})$$

Case 1: $D - C^2 > -(\tau_4 - 2\tau_3 C - \tau_3^2)$. In this case, we have

$$\begin{aligned} \text{Eq. (S50)} &= \frac{|\tau_4 - 2\tau_3 C - \tau_3^2|}{\sqrt{D - C^2} + \sqrt{D - C^2 + (\tau_4 - 2\tau_3 C - \tau_3^2)}} \leq \sqrt{|\tau_4 - 2\tau_3 C - \tau_3^2|} \\ &\leq \sqrt{6\epsilon_0(1 + \nu^2) + 2 \cdot 6\epsilon_0(1 + \nu)^2 + 6\epsilon_0(1 + \nu)} \leq \sqrt{24\epsilon_0}(1 + \nu). \end{aligned}$$

Case 2: $D - C^2 \leq -(\tau_4 - 2\tau_3 C - \tau_3^2)$. In this case, we have $(D + \tau_4) - (C + \tau_3)^2 \leq 0$ and subsequently

$$\text{Eq. (S50)} = \sqrt{D - C^2} \leq \sqrt{|\tau_4 - 2\tau_3 C - \tau_3^2|} \leq \sqrt{24\epsilon_0}(1 + \nu).$$

Combining both cases we prove Eq. (S49). \square

C.3 Proof of Lemma 9

Lemma 9 (restated). *Suppose there exists $\ell \in [L]$ such that $\langle y^{(\ell)}, y^* \rangle \geq 1/\alpha$ for some $\alpha \geq 1$ in Algorithm 5, then $\text{ESTR}(\widehat{S}^{(\ell)}) + \min\{1, \alpha\omega \cdot \text{CI}(\widehat{S}^{(\ell)})\} + \varepsilon \geq \text{ESTR}(S^*) + \min\{1, \omega \cdot \text{CI}(S^*)\}$, where $\varepsilon > 0$ is the approximation parameter of the univariate problem instances.*

Proof. For each assortment S , define $\text{CI}^{(\ell)}(S)$ by

$$\text{CI}^{(\ell)}(S) := y^{(\ell)\top} \left(\frac{\sum_{j \in S} \widehat{u}_{tj} x_{tj} x_{tj}^\top}{1 + \sum_{j \in S} \widehat{u}_{tj}} - \left(\frac{\sum_{j \in S} \widehat{u}_{tj} x_{tj}}{1 + \sum_{j \in S} \widehat{u}_{tj}} \right) \left(\frac{\sum_{j \in S} \widehat{u}_{tj} x_{tj}}{1 + \sum_{j \in S} \widehat{u}_{tj}} \right)^\top \right) y^{(\ell)},$$

Since $\text{CI}^{(\ell)}(S) \leq \text{CI}(S)$, we have

$$\text{ESTR}(\widehat{S}^{(\ell)}) + \min\{1, \alpha\omega \cdot \text{CI}(\widehat{S}^{(\ell)})\} + \varepsilon \geq \text{ESTR}(\widehat{S}^{(\ell)}) + \min\{1, \alpha\omega \cdot \text{CI}^{(\ell)}(\widehat{S}^{(\ell)})\} + \varepsilon. \quad (\text{S51})$$

By the approximation guarantee of Algorithm 3, we have

$$\text{ESTR}(\widehat{S}^{(\ell)}) + \min\{1, \alpha\omega \cdot \text{CI}^{(\ell)}(\widehat{S}^{(\ell)})\} + \varepsilon \geq \text{ESTR}(S^*) + \min\{1, \alpha\omega \cdot \text{CI}^{(\ell)}(S^*)\}. \quad (\text{S52})$$

Since $\langle y^{(\ell)}, y^* \rangle \geq 1/\alpha$, we have $\text{CI}^{(\ell)}(S^*) \geq (1/\alpha) \cdot \text{CI}(S^*)$. Therefore,

$$\text{ESTR}(S^*) + \min\{1, \alpha\omega \cdot \text{CI}^{(\ell)}(S^*)\} \geq \text{ESTR}(S^*) + \min\{1, \omega \cdot \text{CI}(S^*)\}. \quad (\text{S53})$$

The lemma is proved by combining Eq. (S51), Eq. (S52), and Eq. (S53). \square

C.4 Proof of Proposition 2

Proposition 2 (restated). *If $\omega = 0$, then Algorithm 4 terminates in $O(N^4)$ iterations and produces an output S that maximizes $\text{ESTR}(S)$.*

Proof. We first show that when the algorithm terminates with $\text{ESTR}(S) = r$, S is one of the optimal assortments. Suppose S is not an optimal assortment, i.e. there exists $S^\#$ such that $\text{ESTR}(S^\#) > r$, we show that the algorithm will not terminate. By the definition of $\text{ESTR}(\cdot)$ we have $\sum_{i \in S^\#} \widehat{u}_{ti}(r_{ti} - r) > r$ and $\sum_{i \in S} \widehat{u}_{ti}(r_{ti} - r) = r$. By comparing $S^\#$ and S , one can find a new candidate assortment S' via swapping, adding, or deleting an item from/to S such that $\sum_{i \in S'} \widehat{u}_{ti}(r_{ti} - r) > r$. Therefore, $\text{ESTR}(S') > r$ and the algorithm will not terminate.

It remains to show that the algorithm terminates in $O(N^4)$ iterations.

For each $r \in [0, 1]$, we define a total order \geq_r on $[N] \cup \{\perp\}$, where $[N]$ corresponds to the N items and \perp is a special element with the definition $\widehat{u}_{t\perp} = r_{t\perp} = 0$ for convenience, as follows: $i \geq_r j$ if and only if $\widehat{u}_{ti}(r_{ti} - r) \geq \widehat{u}_{tj}(r_{tj} - r)$ (and consequently $i >_r j$ if and only if $\widehat{u}_{ti}(r_{ti} - r) > \widehat{u}_{tj}(r_{tj} - r)$). It is straightforward to verify that there exists $O(N^2)$ section points $\theta_0 < \theta_1 < \theta_2 < \dots < \theta_{L-1} < \theta_L = 1$ so that for any two r_1, r_2 that sandwiched by the same pair of neighboring section points (i.e. $\exists \ell \in [L] : r_1, r_2 \in (\theta_{\ell-1}, \theta_\ell)$), we have $\geq_{r_1} \equiv \geq_{r_2}$. Indeed, one can set the section points to be the solutions to the equalities $\widehat{u}_{ti}(r_{ti} - r) = \widehat{u}_{tj}(r_{tj} - r)$ for every pair of $i, j \in [N] \cup \{\perp\}$.

We will show that if $\text{ESTR}(S) \in (\theta_{\ell-1}, \theta_\ell)$ for some $\ell \in [L]$, after at most $O(N^2)$ iterations, either the algorithm terminates or $\text{ESTR}(S) \geq \theta_\ell$. This directly leads to an $O(N^4)$ upper bound on the total number of iterations that the algorithm performs. We pick an arbitrary $r \in (\theta_{\ell-1}, \theta_\ell)$ and define the following two potential functions: $I(S) = |\{(i, j) : i \in S, j \in [N] \setminus S, i <_r j\}|$, and $J(S) = |\{i \in S : i <_r \perp\}|$. We have the following observations:

- When a swapping operation is performed on S , $I(S)$ strictly decreases and $J(S)$ does not increase.
- When a deletion operation is performed on S , $I(S)$ increases by at most N and $J(S)$ strictly decreases.
- When an addition operation is performed on S , $I(S)$ increases by at most N and $J(S)$ does not increase.

We let $F(S) = I(S) + (2N + 1) \cdot J(S) \in [0, O(N^2)]$. Suppose there are a swapping operations, b deletion operations, and c addition operations done in total, S is the assortment that the algorithm begins with and T is the last assortment satisfying $\text{ESTR}(T) < \theta_\ell$. Observe that there are at most $c \leq b + N$ addition operations. Together with the three observations above, we have

$$\begin{aligned} 0 \leq F(T) &\leq F(S) - a + Nb - (2N + 1)b + Nc \leq F(S) - a + Nb - (2N + 1)b + N(b + N) \\ &\leq F(S) + N^2 - a - b \leq O(N^2) - a - b. \end{aligned}$$

In total, we have $a + b \leq O(N^2)$. Therefore, the total number of iterations where $\text{ESTR}(S) \in (\theta_{\ell-1}, \theta_\ell)$ is $a + b + c \leq a + 2b + N \leq O(N^2)$. \square

C.5 Proof of Proposition 3

Proposition 3 (restated). *Assume that $d \geq 2$. Let $y^* \in \mathbb{R}^d$, $\|y^*\|_2 = 1$ be fixed and y be sampled uniformly at random from the unit d -dimensional sphere. Then*

$$\Pr[\langle y, y^* \rangle \geq 1/\sqrt{d}] = \Omega(1) \quad \text{and} \quad \Pr[\langle y, y^* \rangle \geq 1/2] = \exp\{-O(d)\}.$$

Proof. Assume without loss of generality that $y^* = (1, 0, 0, \dots, 0)$, and let y be sampled as follows. Sample $z_i \sim N(0, 1)$ independently for each $i \in [d]$, and let $y = z/\|z\|_2$. Now, $\langle y, y^* \rangle = z_1/\|z\|_2$.

We first prove $\Pr[\langle y, y^* \rangle \geq 1/\sqrt{d}] = \Pr[z_1/\|z\|_2 \geq 1/\sqrt{d}] = \Omega(1)$. Note that when $z_1 \geq 10$ and $\sqrt{z_2^2 + \dots + z_d^2} \leq 5\sqrt{d}$, we have $z_1/\|z\|_2 = 1/\sqrt{1 + (z_2^2 + \dots + z_d^2)/z_1^2} \geq 1/\sqrt{1 + (5\sqrt{d})^2/10^2} \geq 1/\sqrt{d}$, where the last inequality holds for $d \geq 2$. Therefore,

$$\begin{aligned} \Pr[z_1/\|z\|_2 \geq 1/\sqrt{d}] &\geq \Pr\left[z_1 \geq 10 \wedge \sqrt{z_2^2 + \dots + z_d^2} \leq 5\sqrt{d}\right] \\ &= \Pr[z_1 \geq 10] \cdot \Pr\left[\sqrt{z_2^2 + \dots + z_d^2} \leq 5\sqrt{d}\right] = \Omega(1). \end{aligned}$$

Now we prove $\Pr[\langle y, y^* \rangle \geq 1/2] = \Pr[z_1/\|z\|_2 \geq 1/2] = \exp\{-O(d)\}$. Similarly, when $z_1 \geq 5\sqrt{d}$ and $\sqrt{z_2^2 + \dots + z_d^2} \leq 5\sqrt{d}$, we have $z_1/\|z\|_2 = 1/\sqrt{1 + (z_2^2 + \dots + z_d^2)/z_1^2} \geq 1/\sqrt{1 + 1} > 1/2$. Therefore,

$$\begin{aligned} \Pr[z_1/\|z\|_2 \geq 1/2] &\geq \Pr\left[z_1 \geq 5\sqrt{d} \wedge \sqrt{z_2^2 + \dots + z_d^2} \leq 5\sqrt{d}\right] \\ &= \Pr[z_1 \geq 5\sqrt{d}] \cdot \Pr\left[\sqrt{z_2^2 + \dots + z_d^2} \leq 5\sqrt{d}\right] \\ &= \exp\{-O(d)\} \cdot \Omega(1) = \exp\{-O(d)\}. \end{aligned}$$

\square