

Comments on "Light deflection by Damour-Solodukhin wormholes and Gauss-Bonnet theorem, Phys. Rev. D 98, 044033 (2018) by Ali Övgün"

Amrita Bhattacharya*

Department of Mathematics, Kidderpore College, 2,
Pitamber Sircar Lane, Kolkata 700023, WB, India

Ramis Kh. Karimov†

Zel'dovich International Center for Astrophysics, Bashkir State Pedagogical University,
3A, October Revolution Street, Ufa 450008, RB, Russia

(Dated: March 15, 2022)

In the article cited above, Övgün [1] calculated the leading order deflection angle of light by the static Damour-Solodukhin wormhole [2] using its redefined static and Kerr-like forms proposed by Bueno *et al* [3]. Because of the importance of wormholes, especially with regard to strong field lensing [4,5] and gravitational waves [2, 3, 6–10], we decided to take a closer look at the work in [1].

The purpose of this Comment is to point out that there are conceptual and mathematical flaws in the analysis in [1]. First, the coefficients \bar{a} and \bar{b} derived by the author are incorrect as \bar{a} in Eqn.(2.31) has a wrong sign (probably a typo) but \bar{b} in Eqn.(2.32) is in error. Second, the deflection angle in Eqn.(3.10) in the Kerr-like case is plainly incorrect.

Övgün takes the redefined form of the metric given in [3], viz.,

$$ds^2 = - \left(1 - \frac{2M}{r}\right) c^2 dt^2 + \left[1 - \frac{2M(1 + \lambda^2)}{r}\right]^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2, \quad (1)$$

where λ is a constant parameter. The strong field deflection angle in the static case due to Bozza is [11]

$$\hat{\alpha}(\theta) = -\bar{a} \log \left(\frac{u}{u_m} - 1 \right) + \bar{b}. \quad (2)$$

The author obtains the Bozza coefficients \bar{a} and \bar{b} of the strong field deflection angle for the static case as follows (Eqs.(2.31)-(2.32))

$$\begin{aligned} \bar{a} &= -\frac{1}{\sqrt{1 - 2\lambda^2}}, \\ \bar{b} &= \frac{\log(6) \sqrt{-2\lambda^2 + 1}}{\log(10)2\lambda^2 - 1} + (r_m - \pi). \end{aligned}$$

The correct expressions of the coefficients for the metric (1) however work out to

$$\bar{a} = \frac{1}{\sqrt{1 - 2\lambda^2}}, \quad (3)$$

$$\bar{b} = \frac{\log(6)}{\sqrt{1 - 2\lambda^2}} + (b_R - \pi), \quad (4)$$

$$b_R = \int_0^1 g(z, r_m) dz, \quad (5)$$

$$g(z, r_m) = 2 \left[\frac{\sqrt{3 + 6z}}{z \sqrt{(3 - 2z)\{1 + 2z - 2(1 - z)\lambda^2\}}} - \frac{1}{z \sqrt{1 - 2\lambda^2}} \right]. \quad (6)$$

*Electronic address: amrita'852003@yahoo.co.in

†Electronic address: karimov'ramis'1992@mail.ru

Observe that the first part in \bar{b} in Eqn.(2.32) is different from the corresponding part in the expression in (4). The second part $(r_m - \pi)$ in Eqn.(2.32) depending on M via $r_m = 3M$ becomes dimensionful differing from $(b_R - \pi)$ in the expression in (4). However, note that the Bozza coefficients are dimensionless, which is achieved in (6) since at $r_m = 3M$ the expression for $g(z, r_m)$ becomes independent of M , and consequently $(b_R - \pi)$ becomes dimensionless, as required. The illustrative table below gives a glimpse of the variation of coefficients with λ allowing a comparison with the known Schwarzschild values:

λ	\bar{a}	\bar{b}
0 (Sch)	1.0000	-0.4002
0.1	1.0101	-0.4131
0.2	1.0425	-0.4573
0.3	1.1043	-0.5536

TABLE I: Strong field lensing coefficients for Damour-Solodukhin wormhole.

Next, Övgün [1] obtained the leading order deflection in the Kerr-like case as (Eqn.(3.10))

$$\hat{\alpha} = \frac{2M(\lambda^2 + 2)}{b} \pm \frac{4Ma}{b^2},$$

meaning that the term in $1/b^2$ is symmetric on both sides of the lens. This result is incorrect. Conceptually, the deflection produced by the presence of spin a (inducing frame dragging) explicitly depends on whether the light motion takes place in the direction of spin or opposite to it [12] (see also [13]). Compared to the static case, the deflection is greater for direct and smaller for retrograde orbits resulting in the *loss* of symmetry about the spin axis. The higher the spin, the tighter is the winding of the direct orbit and more unwinding of the retrograde orbits. Mathematically, the exact bending angle for the Kerr case ($\lambda = 0$) has been worked out by Iyer and Hansen [12] and in the weak field it expands as

$$\hat{\alpha} = \frac{4M}{b} + \left[\frac{15\pi}{4} - 4s \left(\frac{a}{M} \right) \right] \left(\frac{M}{b} \right)^2 + \left[\frac{128}{3} - 10\pi s \left(\frac{a}{M} \right) + 4 \left(\frac{a}{M} \right)^2 \right] \left(\frac{M}{b} \right)^3 + \dots \quad (7)$$

where $s = +1$ for direct orbits and -1 for retrograde orbits. The loss of symmetry in the coefficient of $\left(\frac{M}{b} \right)^2$ (as well as in higher order terms) is quite evident. The limiting Kerr value $\lambda = 0$ in Eqn.(3.10) does not reproduce the second term in Eqn.(7).

-
- [1] A. Övgün, Phys. Rev. D **98**, 044033 (2018).
 - [2] T. Damour and S.N. Solodukhin, Phys. Rev. D **76**, 024016 (2007).
 - [3] P. Bueno, P.A. Cano, F. Goelen, T. Hertog and B. Vercnocke, Phys. Rev. D **97**, 024040 (2018).
 - [4] K.K. Nandi, Y.-Z. Zhang and A.V. Zakharov, Phys. Rev. D **74**, 024020 (2006).
 - [5] K.K. Nandi, R.N. Izmailov, E.R. Zhdanov and A. Bhattacharya, JCAP 07 (2018) 027.
 - [6] V. Cardoso, E. Franzin and P. Pani, Phys. Rev. Lett. **116**, 171101 (2016); **117**, 089902(E) (2016).
 - [7] V. Cardoso and P. Pani, Nature Astronomy **1**, 586 (2017).
 - [8] R.A. Konoplya and A. Zhidenko, JCAP 12 (2016) 043.
 - [9] K.K. Nandi, R.N. Izmailov, A.A. Yanbekov and A.A. Shayakhmetov, Phys. Rev. D **95**, 104011 (2017).
 - [10] S.H. Völkel and K.D. Kokkotas, Class. Quantum Grav. **35**, 105018 (2018).
 - [11] V. Bozza, Phys. Rev. D **66**, 103001 (2002).
 - [12] S.V. Iyer and E.C. Hansen, Phys. Rev. D **80**, 124023 (2009).
 - [13] S.V. Iyer, *Frame-dragging induced asymmetry in the bending of light near a Kerr black hole* [arXiv:1808.06630].