

Entropy functions of BPS black holes in AdS_4 and AdS_6

Sunjin Choi¹, Chiung Hwang^{2,3}, Seok Kim¹ and June Nahmgoong¹

¹*Department of Physics and Astronomy & Center for Theoretical Physics,
Seoul National University, Seoul 08826, Korea.*

²*School of Physics, Korea Institute for Advanced Study, Seoul 02455, Korea.*

³*Dipartimento di Fisica, Università di Milano-Bicocca & INFN,
Sezione di Milano-Bicocca, I-20126 Milano, Italy.*

E-mails: csj37100@snu.ac.kr, chwang@kias.re.kr,
skim@phya.snu.ac.kr, earendil25@snu.ac.kr

Abstract

We find the entropy functions of supersymmetric black holes in AdS_4 and AdS_6 with electric charges and angular momenta. Extremizing these functions, one obtains the entropies and the chemical potentials of known analytic black hole solutions.

Contents

1	Introduction	1
2	AdS₄ black holes	2
2.1	Black hole solutions	2
2.2	Entropy function	5
3	AdS₆ black holes	8
3.1	Black hole solutions	8
3.2	Entropy function	10
4	Concluding remarks	11

1 Introduction

Understanding black holes [1, 2, 3] is an important subject in AdS/CFT [4]. In models with supersymmetry, one expects that quantitative analysis at strong coupling would be easier in the BPS sectors of SCFTs. Supersymmetric AdS black holes correspond to thermal ensembles of BPS states, carrying angular momenta and also internal charges (electric charges in AdS). In AdS_d with $d > 3$, supersymmetric black holes have very complicated structures. First of all, it is known that there are no BPS black holes with electric charges only, at zero angular momenta. This is because in the dual field theory, the local BPS operators will reduce to chiral rings which do not have enough numbers of microstates to form black holes: e.g. see [5] for the case with $d = 5$. With nonzero angular momenta, the solutions appear very involved. See, e.g. [6, 7] for $d = 4$, [8, 9, 10, 11] for $d = 5$, [12] for $d = 6$, and [13, 14] for $d = 7$.

In [15, 16], simple underlying structures of BPS black holes were discovered in AdS₅ and AdS₇. Firstly, one can obtain the entropies and the chemical potentials of known BPS black holes [8, 9, 10, 11] in AdS₅ \times S^5 by extremizing the following entropy function [15],

$$S(\Delta_I, \omega_i) = \frac{N^2 \Delta_1 \Delta_2 \Delta_3}{2\omega_1 \omega_2} + \sum_{I=1}^3 \Delta_I Q_I + \sum_{i=1}^2 \omega_i J_i, \quad (1.1)$$

subject to the constraint $\Delta_1 + \Delta_2 + \Delta_3 - \omega_1 - \omega_2 = 2\pi i$. Here Q_I and J_i are $U(1)^3 \subset SO(6)$ electric charges and $U(1)^2 \subset SO(4)$ angular momenta, respectively. The Bekenstein-Hawking entropy of the black hole is the extremal value of $\text{Re}(S)$, at one of the extremum solutions for Δ_I, ω_i [15]. The black hole chemical potentials are the extremal values of $\text{Re}(\Delta_I)$, $\text{Re}(\omega_i)$ [17].

Similarly, the properties of known BPS black holes [13, 14] in $\text{AdS}_7 \times S^4$ can be understood by extremizing the following entropy function [16],

$$S(\Delta_I, \omega_i) = -\frac{N^3(\Delta_1\Delta_2)^2}{24\omega_1\omega_2\omega_3} + \sum_{I=1}^2 \Delta_I Q_I + \sum_{i=1}^3 \omega_i J_i, \quad (1.2)$$

subject to the constraint $\Delta_1 + \Delta_2 - \omega_1 - \omega_2 - \omega_3 = 2\pi i$. Q_I and J_i are $U(1)^2 \subset SO(5)$ electric charges and $U(1)^3 \subset SO(6)$ angular momenta. These extremely simple formulae encode apparently complicated properties of supersymmetric AdS black holes. They triggered interesting follow-up works aiming at microscopic accounts for these black holes [18, 17, 19]. In particular, [17] derived (1.1) for large AdS_5 black holes, from the index of 4d $\mathcal{N} = 4$ Yang-Mills theory. ([17] also found a generalization of [15] in a different charge sector.) [17, 20] also provided an anomaly-based microscopic discussion which leads to (1.2) for large AdS_7 black holes. Therefore, the simple functions (1.1) and (1.2) provided very useful inspirations for microscopic studies.

We thus find that it will be very helpful to have entropy function formalisms for supersymmetric AdS black holes in other dimensions. In this note, we provide such functions in AdS_4 and AdS_6 , simplifying the apparently complicated structures of known black hole solutions. These led to better microscopic understandings based on CFT_3 [21] and CFT_5 duals [22]. In particular, [21] and [22] statistically accounted for large BPS black holes in AdS_4 and AdS_6 , from the indices of CFT_3 and CFT_5 duals.

The rest of this note is organized as follows. Section 2 summarizes the properties of known supersymmetric black holes in $\text{AdS}_4 \times S^7$, and show that an entropy function we suggest encodes these properties. Section 3 makes similar studies with supersymmetric AdS_6 black holes. Section 4 concludes with remarks.

2 AdS_4 black holes

2.1 Black hole solutions

We study the supersymmetric black holes in $\text{AdS}_4 \times S^7$ of [7]. These are obtained by taking supersymmetric limits of [23], also demanding the existence of smooth horizons.

Black holes in $\text{AdS}_4 \times S^7$ can carry six kinds of conserved quantities: mass (or energy) E , angular momentum J on S^2 of global AdS_4 , and four Cartan charges Q_I ($I = 1, 2, 3, 4$) of $SO(8)$ symmetry on S^7 . The last four conserved quantities Q_I appear in 4d gravity as $U(1)^4$ electric charges. The convention of [7] for Q_I is to take four angular momenta acting on the orthogonal 2-planes of \mathbb{R}^8 related to S^7 . The most general black holes known to date have

pairwise equal electric charges, $Q_1 = Q_3$, $Q_2 = Q_4$. With the last charge restrictions, the four conserved quantities E, J, Q_1, Q_2 are labeled by four parameters m, a, δ_1, δ_2 as [7]

$$\begin{aligned} E &= \frac{m}{2G\Xi^2}(\cosh 2\delta_1 + \cosh 2\delta_2) \quad , \quad J = \frac{ma}{2G\Xi^2}(\cosh 2\delta_1 + \cosh 2\delta_2) \quad , \\ Q_1 = Q_3 &= \frac{m}{4G\Xi} \sinh 2\delta_1 \quad , \quad Q_2 = Q_4 = \frac{m}{4G\Xi} \sinh 2\delta_2 \quad , \end{aligned} \quad (2.1)$$

where $\Xi = 1 - a^2 g^2$. The entropy is given by

$$S = \frac{\pi(r_1 r_2 + a^2)}{G\Xi} \quad , \quad (2.2)$$

where $r_i = r_+ + 2m \sinh^2 \delta_i$. $r = r_+$ is the location of the event horizon. G is the 4d Newton constant, which will be replaced by microscopic parameters later. (In [7], all charges and entropy are computed omitting the overall $\frac{1}{G}$ factor, or at $G = 1$. E.g. the entropy is computed by dividing the horizon area by 4, rather than $S = \frac{A}{4G}$.) g is a parameter of the 4d gauged supergravity, and is related to the radius ℓ of AdS_4 as $g = \ell^{-1}$.

The BPS limit of these black holes is given by

$$e^{2\delta_1 + 2\delta_2} = 1 + \frac{2}{ag} \quad , \quad (2.3)$$

which corrects a typo of [7]. Only after this correction, the BPS relation

$$E = gJ + \sum_{I=1}^4 Q_I = gJ + 2Q_1 + 2Q_2 \quad (2.4)$$

is met. A further condition to have a regular horizon is $\Delta_r = 0$ having a double root at $r = r_+$. (See [7] for the definition of the function Δ_r .) This happens only after a further tuning of m . After the tuning, the horizon location $r = r_+$ is given by

$$r_+ = \frac{2m \sinh \delta_1 \sinh \delta_2}{\cosh(\delta_1 + \delta_2)} \quad , \quad (2.5)$$

when m satisfies

$$(mg)^2 = \frac{\cosh^2(\delta_1 + \delta_2)}{e^{\delta_1 + \delta_2} \sinh^3(\delta_1 + \delta_2) \sinh(2\delta_1) \sinh(2\delta_2)} \quad . \quad (2.6)$$

This again corrects the formula $mg = \frac{\cosh(\delta_1 + \delta_2)}{e^{\frac{\delta_1 + \delta_2}{2}} \sinh^2(\delta_1 + \delta_2) \sinh(2\delta_1) \sinh(2\delta_2)}$ of [7]. The typos found in this paragraph are also reported in [24].

Taking the BPS limit, the entropy of the supersymmetric black hole is given by

$$S = \frac{2\pi}{g^2 G (e^{2\delta_1 + 2\delta_2} - 3)} \quad . \quad (2.7)$$

The two conditions (2.3), (2.6) leave two independent parameters among m, a, δ_1, δ_2 . Even after restricting E as (2.4) due to the BPS condition, the remaining charges Q_1, Q_2, J satisfy a

relation. Together with S , we find the following two relations after taking the BPS limit:

$$\begin{aligned} \left(\frac{2Q_1}{g} + \frac{2Q_2}{g} \right) S &= \frac{\pi}{g^2 G} J, \\ S^2 + \frac{\pi}{g^2 G} S - 4\pi^2 \frac{2Q_1}{g} \frac{2Q_2}{g} &= 0. \end{aligned} \quad (2.8)$$

Since these equations determine S twice, one will get a charge relation between Q_1, Q_2, J from the compatibility of two equations. Explicitly, we insert the solution of the first equation to the second equation, demanding two equations have the same solution for S . Then, taking the unique positive solution assuming $Q_1, Q_2, J > 0$, one obtains

$$\begin{aligned} S &= \frac{\pi}{g^2 G} \frac{J}{\left(\frac{2Q_1}{g} + \frac{2Q_2}{g} \right)}, \\ J &= \frac{1}{2} \left(\frac{2Q_1}{g} + \frac{2Q_2}{g} \right) \left(-1 + \sqrt{1 + 16g^4 G^2 \frac{2Q_1}{g} \frac{2Q_2}{g}} \right). \end{aligned} \quad (2.9)$$

Thus, we have explicitly found the charge relation between Q_1, Q_2, J .

The black hole chemical potentials and the free energy F satisfy

$$S = -T^{-1}F(T) + T^{-1}E - T^{-1}\Omega J - T^{-1} \sum_{I=1}^4 \Phi_I Q_I, \quad (2.10)$$

where T is the temperature, Ω is the angular velocity, and Φ_i 's are the electrostatic potentials. The chemical potentials are evaluated on the horizon. In the BPS limit we are interested in,

$$T = \frac{\Delta'_r}{4\pi(r_1 r_2 + a^2)} \rightarrow 0 \quad (2.11)$$

because Δ_r has a double root at the horizon. On the other hand, as one inserts the value of the variables in the BPS limit, $a = \frac{2}{g(e^{2\delta_1+2\delta_2}-1)}$, mg given by (2.6), and then the horizon location $r \rightarrow r_+$ (2.5), one finds

$$\Omega = \frac{a(1+g^2 r_1 r_2)}{r_1 r_2 + a^2} \rightarrow g, \quad \Phi_1 = \frac{m r_2 \sinh(2\delta_1)}{r_1 r_2 + a^2} \rightarrow 1, \quad \Phi_2 = \frac{m r_1 \sinh(2\delta_2)}{r_1 r_2 + a^2} \rightarrow 1. \quad (2.12)$$

Defining ΔE by $E = \Delta E + 2Q_1 + 2Q_2 + gJ$, one finds that

$$S = -T^{-1}F(T) + T^{-1}\Delta E - T^{-1}(\Omega - g)J - T^{-1} \sum_{I=1}^4 (\Phi_I - 1)Q_I. \quad (2.13)$$

The BPS limit satisfies $T \rightarrow 0$, $\Delta E \rightarrow 0$. One first finds that

$$\omega = -\lim_{T \rightarrow 0} (T^{-1}(\Omega - g)) , \quad \Delta_I = -\lim_{T \rightarrow 0} (T^{-1}(\Phi_I - 1)) \quad (2.14)$$

are well defined in the BPS limit, by explicitly computing them (although the expressions are very complicated). Since S is also finite in this limit, the ‘BPS free energy’ $F_{\text{BPS}} \equiv \lim_{T \rightarrow 0}(T^{-1}(F - \Delta E))$ should also be well defined. So one finds

$$S = -F_{\text{BPS}} + \omega J + \sum_{I=1}^4 \Delta_I Q_I \quad (2.15)$$

in the BPS limit. $-F_{\text{BPS}}$ is to be interpreted as $\log Z$, where Z is the BPS partition function of this system. We again stress that the BPS limit is taken by first inserting $ag \rightarrow \frac{2}{e^{2\delta_1 + 2\delta_2} - 1}$, $mg \rightarrow \sqrt{\frac{(\coth(\delta_1 + \delta_2) - 1) \coth^2(\delta_1 + \delta_2)}{\sinh(2\delta_1) \sinh(2\delta_2)}}$ and then $r \rightarrow \frac{2m \sinh \delta_1 \sinh \delta_2}{\cosh(\delta_1 + \delta_2)}$. This results in quite complicated expressions for ω, Δ_i . After taking the BPS limit, one can show that they satisfy

$$\Delta_1 + \Delta_2 = \frac{1}{g}\omega \quad \Rightarrow \quad \sum_{I=1}^4 \frac{g}{2} \Delta_I - \omega = 0 . \quad (2.16)$$

This is an alternative statement of the charge relation between Q_1, Q_2, J .

2.2 Entropy function

We now present an entropy function, whose suitable Legendre transformation in Δ_I, ω yields the entropy $S(Q_I, J)$ and the BPS chemical potentials of the supersymmetric black holes. Our entropy function $S(\Delta_I, \omega; Q_I, J)$ is given by

$$S(\Delta_I, \omega; Q_I, J) = -i \frac{4\sqrt{2}N^{\frac{3}{2}}}{3} \frac{\sqrt{\Delta_1 \Delta_2 \Delta_3 \Delta_4}}{\omega} + \omega J + \sum_{I=1}^4 \Delta_I Q_I . \quad (2.17)$$

We extremize S in Δ_I, ω with the constraint

$$\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4 - \omega = 2\pi i . \quad (2.18)$$

A microscopic derivation of the entropy function (2.17) from the CFT_3 dual was studied in [21], in the Cardy limit $\omega \rightarrow 0$. Just like AdS_5 , AdS_7 black holes analyzed in [17], the constraint (2.18) is given an interpretation in [21]. Here the number of M2-branes N is related to the 4d Newton constant G as follows:

$$G_{11} = 16\pi^7 \ell_{\text{P}}^9, \ell_{S^7} = 2\ell = \ell_{\text{P}}(2^5 \pi^2 N)^{1/6} \Rightarrow \frac{1}{g^2 G} = \frac{\text{vol}(S^7)}{g^2 G_{11}} = \frac{2\sqrt{2}}{3} \frac{N^{3/2}}{g^2 \ell^2} = \frac{2\sqrt{2}}{3} N^{3/2} . \quad (2.19)$$

ℓ_{P} is the 11d Planck scale, ℓ_{S^7} is the radius of S^7 , and ℓ is the AdS_4 radius as defined in the previous subsection. We claim that the resulting extremal value of $\text{Re}(S)$ is the entropy of supersymmetric black holes. We shall check this against the known solutions summarized in the previous subsection, at $Q_1 = Q_3, Q_2 = Q_4$ (which is equivalent to $\Delta_1 = \Delta_3, \Delta_2 = \Delta_4$). Here, note that the chemical potentials Δ_I, ω are all complexified. With complex Δ_I , the square

root $\sqrt{\Delta_1\Delta_2\Delta_3\Delta_4}$ in (2.17) should be understood as to take the argument of $\Delta_1\Delta_2\Delta_3\Delta_4$ in the principal branch $(-\pi, \pi)$ [21].

We show our claim by extremizing S , subject to the constraint (2.18). We introduce the Lagrange multiplier λ and extremize

$$S = -i \frac{4\sqrt{2}N^{\frac{3}{2}}}{3} \frac{\sqrt{\Delta_1\Delta_2\Delta_3\Delta_4}}{\omega} + \omega J + \sum_{I=1}^4 \Delta_I Q_I + \lambda \left(\sum_{I=1}^4 \Delta_I - \omega - 2\pi i \right). \quad (2.20)$$

The extremum conditions are given by

$$\begin{aligned} \lambda + Q_I &= i \frac{4\sqrt{2}N^{\frac{3}{2}}}{3\omega} \frac{\sqrt{\Delta_1\Delta_2\Delta_3\Delta_4}}{2\Delta_I} \quad (I = 1, \dots, 4), \\ \lambda - J &= i \frac{4\sqrt{2}N^{\frac{3}{2}}}{3\omega^2} \sqrt{\Delta_1\Delta_2\Delta_3\Delta_4}. \end{aligned} \quad (2.21)$$

Inserting these charges into (2.20), to eliminate the appearances of Q_I, J , one obtains

$$S = -2\pi i \lambda. \quad (2.22)$$

Multiplying the four equations on the first line of (2.21), one finds

$$(\lambda + Q_1)(\lambda + Q_2)(\lambda + Q_3)(\lambda + Q_4) = \frac{64N^6}{81\omega^4} \Delta_1\Delta_2\Delta_3\Delta_4 = -\frac{2N^3}{9}(\lambda - J)^2. \quad (2.23)$$

So one obtains a very useful expression,

$$\left(\frac{S}{2\pi i} - Q_1 \right) \left(\frac{S}{2\pi i} - Q_2 \right) \left(\frac{S}{2\pi i} - Q_3 \right) \left(\frac{S}{2\pi i} - Q_4 \right) = -\frac{2N^3}{9} \left(\frac{S}{2\pi i} + J \right)^2. \quad (2.24)$$

One needs care to treat the above expression. While the above is the quartic equation in S , only the half of them are the true solutions to (2.21) satisfying the constraint (2.18). The other halves are the extraneous solutions. Hence, after solving the above equation, one should check whether the resulting solution is a true one.

After extremizing the entropy function, one would generally obtain complex solutions for S by solving (2.24). Along the spirit of [17], we shall generally regard $\text{Re}(S)$ as the entropy at the extremum. See [17, 21] for the interpretation of the imaginary part. However, we are primarily interested in comparing our results against the known black hole solutions of section 2.1. Therefore, we impose the charge relation of these black holes and compare the thermodynamic quantities on that surface only. Somewhat remarkably, the charge relation of known black holes will turn out to be $\text{Im}(S) = 0$ at the extremum of our entropy function. So from now on, we demand the existence of a real solution for S in (2.24), and compare the results with the known black holes. Demanding real S for real charges Q_1, Q_2, Q_3, Q_4, J , the complex equation (2.24) is separated into two real equations as follows:

$$\begin{aligned} \frac{1}{16\pi^4} S^4 - \frac{\sum_{I<J} Q_I Q_J}{4\pi^2} S^2 + Q_1 Q_2 Q_3 Q_4 &= \frac{N^3}{18\pi^2} S^2 - \frac{2N^3 J^2}{9}, \\ -\frac{\sum_I Q_I}{8\pi^3} S^3 + \frac{\sum_{I<J<K} Q_I Q_J Q_K}{2\pi} S &= \frac{2N^3 J}{9\pi} S. \end{aligned} \quad (2.25)$$

These equations determine S twice as functions of charges. From the compatibility of two equations, one will get a relation of Q_I, J . Explicitly, one may take the unique positive solution of the second equation and insert it to the first equation, to obtain the charge relation. One can check that this solution is a true solution satisfying (2.21) and (2.18).

To compare with known black holes summarized in section 2.1, we set $Q_1 = Q_3, Q_2 = Q_4$. Then, taking the unique positive solution assuming $Q_1, Q_2, J > 0$, one obtains

$$\begin{aligned} S &= \frac{2\pi}{3} \sqrt{\frac{9Q_1Q_2(Q_1+Q_2) - 2N^3J}{Q_1+Q_2}}, \\ 0 &= 2N^3J^2 + 2N^3(Q_1+Q_2)J - 9Q_1Q_2(Q_1+Q_2)^2. \end{aligned} \quad (2.26)$$

These can be rearranged as

$$\begin{aligned} S &= \frac{2\sqrt{2}\pi N^{\frac{3}{2}}}{3} \frac{J}{Q_1+Q_2} = \frac{\pi}{g^2G} \frac{J}{Q_1+Q_2}, \\ J &= \frac{1}{2}(Q_1+Q_2) \left(-1 + \sqrt{1 + \frac{18}{N^3}Q_1Q_2} \right) = \frac{1}{2}(Q_1+Q_2) \left(-1 + \sqrt{1 + 16g^4G^2Q_1Q_2} \right). \end{aligned} \quad (2.27)$$

One can easily check that this solution indeed satisfies (2.21) and (2.18), i.e. it is not an extraneous solution. The above expressions are exactly the same as (2.9), which we obtained from the supersymmetric black holes. Note that the charges and chemical potentials of the entropy function (2.20) are related to those of supersymmetric black holes as

$$\begin{aligned} S_{\text{BH}} &= S, \quad J_{\text{BH}} = J, \quad \frac{2}{g}Q_{I,\text{BH}} = Q_I, \\ \omega_{\text{BH}} &= \text{Re}(\omega), \quad \frac{g}{2}\Delta_{I,\text{BH}} = \text{Re}(\Delta_I). \end{aligned} \quad (2.28)$$

Here, the subscripts ‘BH’ denote the black hole quantities, while the others are the quantities used in the entropy function. The second line can be shown by a rather straightforward but tedious calculus. One also finds that the relation between the chemical potentials in the entropy function (2.18) is equivalent to that of the supersymmetric black holes (2.16).

To summarize, our entropy function (2.20) indeed reproduces the Bekenstein-Hawking entropy of the supersymmetric AdS_4 black holes (2.7) and the corresponding charge/chemical potential relations (2.9), (2.16), at $Q_1 = Q_3, Q_2 = Q_4$ where solutions are known. Recently, 4 parameter BPS black hole solutions with all different Q_I ’s were discovered in [25], whose physics is successfully described by our entropy function (2.20).

3 AdS₆ black holes

3.1 Black hole solutions

In this section, we study the supersymmetric AdS₆ black holes, and find an entropy function which accounts for their physics. We construct an entropy function for the solution of [12]. The solution may be regarded as describing BPS states of any large N 5d SCFT dual. For instance, as our favorite example, results in this section may be understood in the context of massive type IIA string theory on warped AdS₆ \times S^4/\mathbb{Z}_2 product background. This system is dual to 5d $\mathcal{N} = 1$ SCFT living on N D4-branes probing the O8-D8 system [26]. The 5d SCFT dual has a gauge theory description, with $Sp(N)$ gauge group, rank 2 antisymmetric hypermultiplet, and $N_f \leq 7$ fundamental hypermultiplets. However, we expect that our general analysis can be embedded to AdS₆ black holes in the backgrounds of [27, 28, 29].

The 6d $\mathcal{N} = (1, 0)$ $SU(2)$ gauged supergravity was obtained by a consistent Kaluza-Klein truncation of massive type IIA supergravity on S^4/\mathbb{Z}_2 [30]. In [12], the charged rotating AdS₆ black hole solution in this gauged supergravity was obtained. It has four kinds of conserved quantities: mass E , two angular momenta J_1, J_2 , which describe the orthogonal 2-plane rotations on S^4 in global AdS₆, and one $U(1) \subset SU(2)$ electric charge Q . They are given in terms of four parameters m, a, b, δ of the solution as [12]

$$\begin{aligned} E &= \frac{2\pi m}{3G\Xi_a\Xi_b} \left[\frac{1}{\Xi_a} + \frac{1}{\Xi_b} + \sinh^2 \delta \left(1 + \frac{\Xi_a}{\Xi_b} + \frac{\Xi_b}{\Xi_a} \right) \right], \quad Q = \frac{\pi m}{G\Xi_a\Xi_b} \sinh 2\delta, \\ J_1 &= \frac{2\pi m a}{3G\Xi_a^2\Xi_b} (1 + \Xi_b \sinh^2 \delta), \quad J_2 = \frac{2\pi m b}{3G\Xi_a\Xi_b^2} (1 + \Xi_a \sinh^2 \delta), \end{aligned} \quad (3.1)$$

where $\Xi_a = 1 - a^2 g^2$ and $\Xi_b = 1 - b^2 g^2$. The entropy is given by

$$S = \frac{2\pi^2 [(r_+^2 + a^2)(r_+^2 + b^2) + 2mr_+ \sinh^2 \delta]}{3G\Xi_a\Xi_b}. \quad (3.2)$$

The event horizon is located at $r = r_+$. Here, G is the 6d Newton constant. (In [12], the unit $G = 1$ is used.) g is a gauge coupling constant in 6d gravity, setting the inverse-radius of AdS₆.

This black hole solution admits the supersymmetric limit without naked closed timelike curves. The BPS condition

$$E = gJ_1 + gJ_2 + Q \quad (3.3)$$

is satisfied if

$$e^{2\delta} = 1 + \frac{2}{(a+b)g}. \quad (3.4)$$

In addition, a smooth horizon exists only if

$$m = \frac{(a+b)^2(1+ag)(1+bg)(2+ag+bg)}{2(1+ag+bg)} \sqrt{\frac{ab}{1+ag+bg}} \quad (3.5)$$

is satisfied. The horizon is located at

$$r_+ = \sqrt{\frac{ab}{1+ag+bg}} . \quad (3.6)$$

Taking the BPS limit, the entropy of the supersymmetric black hole is given by

$$S = \frac{2\pi^2 ab(a+b)}{3gG(1-ag)(1-bg)(1+ag+bg)} . \quad (3.7)$$

The two conditions (3.4), (3.5) leave two independent parameters among m, a, b, δ . Even after restricting E as (3.3) from the BPS condition, the remaining charges J_1, J_2, Q carried by the supersymmetric black holes will satisfy a charge relation. Equivalently, together with S , we find the following two relations:

$$\begin{aligned} S^3 - \frac{2\pi^2}{3g^4G} S^2 - 12\pi^2 \left(\frac{Q}{3g}\right)^2 S + \frac{8\pi^4}{3g^4G} J_1 J_2 &= 0 , \\ \frac{Q}{3g} S^2 + \frac{2\pi^2}{9g^4G} (J_1 + J_2) S - \frac{4\pi^2}{3} \left(\frac{Q}{3g}\right)^3 &= 0 . \end{aligned} \quad (3.8)$$

Since these equations determine S twice, one will get a charge relation between J_1, J_2, Q from the compatibility of two equations. Explicitly, one may take the unique positive solution of the second equation and insert it to the first equation, to get the charge relation.

The black hole chemical potentials and the free energy F satisfy

$$S = -T^{-1}F + T^{-1}E - T^{-1}\Omega_1 J_1 - T^{-1}\Omega_2 J_2 - T^{-1}\Phi Q , \quad (3.9)$$

where T is the temperature, Ω_1, Ω_2 are the angular velocities, and Φ is the electrostatic potential. The temperature of the supersymmetric black hole is zero in the BPS smooth horizon limit,

$$T = \frac{2r_+^2(1+g^2r_+^2)(2r_+^2+a^2+b^2) - (1-g^2r_+^2)(r_+^2+a^2)(r_+^2+b^2) + 8mg^2r_+^3 \sinh^2 \delta - 4m^2g^2 \sinh^4 \delta}{4\pi r_+[(r_+^2+a^2)(r_+^2+b^2) + 2mr_+ \sinh^2 \delta]} \rightarrow 0 . \quad (3.10)$$

The other chemical potentials in the BPS limit are given by

$$\begin{aligned} \Omega_1 &= a \frac{(1+g^2r_+^2)(r_+^2+b^2) + 2mg^2r_+ \sinh^2 \delta}{(r_+^2+a^2)(r_+^2+b^2) + 2mr_+ \sinh^2 \delta} \rightarrow g , \\ \Omega_2 &= b \frac{(1+g^2r_+^2)(r_+^2+a^2) + 2mg^2r_+ \sinh^2 \delta}{(r_+^2+a^2)(r_+^2+b^2) + 2mr_+ \sinh^2 \delta} \rightarrow g , \\ \Phi &= \frac{mr_+ \sinh 2\delta}{(r_+^2+a^2)(r_+^2+b^2) + 2mr_+ \sinh^2 \delta} \rightarrow 1 . \end{aligned} \quad (3.11)$$

Similar to the analysis in section 2.1, the following limits exist,

$$F_{\text{BPS}} = \lim_{T \rightarrow 0} (T^{-1}(F - \Delta E)) , \quad \omega_i = - \lim_{T \rightarrow 0} (T^{-1}(\Omega_i - g)) , \quad \Delta = - \lim_{T \rightarrow 0} (T^{-1}(\Phi - 1)) , \quad (3.12)$$

where $\Delta E \equiv E - Q - gJ_1 - gJ_2$. Then, in the zero temperature BPS limit, one obtains

$$S = -F_{\text{BPS}} + \omega_1 J_1 + \omega_2 J_2 + \Delta Q . \quad (3.13)$$

Using the computed expressions for ω_i, Δ , one finds that

$$\omega_1 + \omega_2 = 3g\Delta . \quad (3.14)$$

Again, this is the alternative statement of the charge relation of J_1, J_2, Q .

3.2 Entropy function

We now present an entropy function which encodes the physics of the BPS black holes presented in the previous subsection. The entropy function is given by

$$S = -i \frac{\pi}{81g^4 G} \frac{\Delta^3}{\omega_1 \omega_2} + \Delta Q + \omega_1 J_1 + \omega_2 J_2 + \lambda \left(\Delta - \omega_1 - \omega_2 - 2\pi i \right) , \quad (3.15)$$

where G is the 6d Newton constant as before. Having in mind the concrete example of massive IIA supergravity on warped $\text{AdS}_6 \times S^4/\mathbb{Z}_2$ background, one would find $\frac{1}{g^4 G} = \frac{27\sqrt{2}}{5\pi} \frac{N^{\frac{5}{2}}}{\sqrt{8-N_f}}$ [22]. In that case, a microscopic derivation of the entropy function (3.15) from the CFT_5 dual was studied in [22], in the Cardy limit $\omega_{1,2} \rightarrow 0$. Here, we introduced the Lagrange multiplier λ to extremize S in $\Delta, \omega_1, \omega_2$ subject to the constraint

$$\Delta - \omega_1 - \omega_2 = 2\pi i . \quad (3.16)$$

Differentiating with respect to the chemical potentials, one obtains

$$\lambda + Q = i \frac{\pi}{27g^4 G} \frac{\Delta^2}{\omega_1 \omega_2} , \quad \lambda - J_1 = i \frac{\pi}{81g^4 G} \frac{\Delta^3}{\omega_1^2 \omega_2} , \quad \lambda - J_2 = i \frac{\pi}{81g^4 G} \frac{\Delta^3}{\omega_1 \omega_2^2} . \quad (3.17)$$

Inserting these back to the original entropy function formula, one obtains

$$S = -2\pi i \lambda . \quad (3.18)$$

Multiplying the last two equations of (3.17), one obtains

$$(\lambda - J_1)(\lambda - J_2) = - \left(\frac{\pi}{81g^4 G} \right)^2 \frac{\Delta^6}{\omega_1^3 \omega_2^3} = -i \frac{3g^4 G}{\pi} (\lambda + Q)^3 . \quad (3.19)$$

Hence, one obtains

$$\left(\frac{S}{2\pi i} + J_1 \right) \left(\frac{S}{2\pi i} + J_2 \right) = i \frac{3g^4 G}{\pi} \left(\frac{S}{2\pi i} - Q \right)^3 . \quad (3.20)$$

As in our section 2 and [17], we dismiss $\text{Im}(S)$, focussing on $\text{Re}(S)$ as our entropy. However, again note that all known supersymmetric AdS_6 black holes have a charge relation. This charge

relation will coincide with the condition $\text{Im}(S) = 0$ at the saddle point. So we demand real S for real charges Q, J_1, J_2 . Then, (3.20) is separated into two real equations as follows:

$$\begin{aligned} S^3 - \frac{2\pi^2}{3g^4G}S^2 - 12\pi^2Q^2S + \frac{8\pi^4}{3g^4G}J_1J_2 &= 0, \\ QS^2 + \frac{2\pi^2}{9g^4G}(J_1 + J_2)S - \frac{4\pi^2}{3}Q^3 &= 0. \end{aligned} \quad (3.21)$$

These equations determine S twice as functions of charges. Therefore, from the compatibility of two equations, one obtains a charge relation of Q, J_1, J_2 . These two equations of S, Q, J_1, J_2 (3.21), derived from the entropy function (3.15), are exactly the same as those from the supersymmetric black holes (3.8). Note that the charges and chemical potentials of the entropy function (3.15) are related to those of the black holes as

$$\begin{aligned} S_{\text{BH}} &= S, \quad J_{i,\text{BH}} = J_i, \quad \frac{1}{3g}Q_{\text{BH}} = Q, \\ \omega_{i,\text{BH}} &= \text{Re}(\omega_i), \quad 3g\Delta_{\text{BH}} = \text{Re}(\Delta). \end{aligned} \quad (3.22)$$

The subscripts ‘BH’ denote the black hole quantities, while the others are the quantities used in the entropy function. One can also realize that the relation between the chemical potentials in the entropy function (3.16) is equivalent to that of the supersymmetric black holes (3.14).

Thus, our entropy function (3.15) indeed reproduces the Bekenstein-Hawking entropy of the supersymmetric AdS_6 black holes (3.7), and also their chemical potentials.

4 Concluding remarks

In this note, we presented the entropy functions of supersymmetric black holes in AdS_4 and AdS_6 . Complicated black hole quantities can be very concisely understood from simple extremization principles of these entropy functions. Considering the inspirations given by the similar entropy functions in AdS_5 [15] or AdS_7 [16] to their microscopic studies, we expect that our entropy functions will also play similar roles.

The entropy function in AdS_4 , presented in section 2, was actually motivated by our microscopic study [21] of BPS states in the M2-brane QFT (see, e.g. [32, 33, 34]) from its index [31]. In particular, [21] derived the entropy function (2.17) for large AdS_4 black holes, from the index of the radially quantized SCFT on M2-branes, where the condensation of the magnetic monopole operators gives rise to the novel deconfined $N^{\frac{3}{2}}$ degrees of freedom.

One can also derive our results on AdS_6 black holes from 5d SCFT duals, for instance, the strong coupling limits of 5d gauge theories on the D4-D8-O8 system [22]. The indices of such SCFTs on $S^4 \times \mathbb{R}$ were explored in [35], where the problem reduced to studies of the 5d

instanton partition functions: see, e.g. [36, 37, 38]. Later, [22] derived the entropy function (3.15) for large AdS_6 black holes, from the indices of such 5d SCFTs and their orbifold theories, where the instanton solitons play subtle roles to realize deconfined $N^{\frac{5}{2}}$ degrees of freedom. Furthermore, while AdS_6 black hole solution known to date has only one electric charge dual to R-charge, [22] obtained a more general form of the entropy function, which describes AdS_6 black holes carrying various electric charges, dual to R-charge, mesonic charge and baryonic charges, yet to be discovered. For example, when the black hole has one more electric charge dual to the mesonic charge, the numerator $\sim \Delta^3$ of our entropy function (3.15) is refined to $[(\Delta + \hat{m})(\Delta - \hat{m})]^{\frac{3}{2}}$, where $m \equiv \hat{m} + 2\pi i$ is the chemical potential conjugate to the mesonic charge.

One may think of generalizations of our results on $\text{AdS}_4 \times S^7$, to more general 4d $\mathcal{N} = 2$ gauged supergravity models arising from string or M-theory. To see a natural possibility of generalization, note that the numerator $\sim \sqrt{\Delta_1 \Delta_2 \Delta_3 \Delta_4}$ of our entropy function (2.17) is the homogeneous degree 2 prepotential of the $U(1)^4$ supergravity [39]. The prepotential is the square root of a degree 4 polynomial. See, e.g. [40, 41] for such structures in other backgrounds. We conjecture that, for BPS black holes in 4d $\mathcal{N} = 2$ gauged supergravity, an entropy function like (2.17) can be constructed by replacing the numerator by the prepotential of the theory. Recently, such an entropy function was found in [25], and also microscopically studied in [42] from the indices of CFT₃ duals in the Cardy limit $\omega \rightarrow 0$.

Here, note that similar prepotentials appeared in the entropy functions of magnetic/dyonic AdS_4 black holes [43] (see also [44]). The unifying underlying structures for all these entropy functions were microscopically studied in [42] from CFT₃ duals. We finally note that the entropy functions of electric AdS_6 black holes in our paper also appear to have some similarities with magnetized black holes in AdS_6 , with boundaries replaced by more general 4-manifolds [45, 46].

Acknowledgements

We thank Dongmin Gang, Hee-Cheol Kim, Joonho Kim, Kimyeong Lee and Jaemo Park for helpful discussions. This work is supported in part by the National Research Foundation of Korea (NRF) Grant 2018R1A2B6004914 (SC, SK, JN), NRF-2017-Global Ph.D. Fellowship Program (SC), the ERC-STG grant 637844-HBQFTNCER (CH), the INFN (CH), and Hyundai Motor Chung Mong-Koo Foundation (JN).

References

- [1] S. W. Hawking and D. N. Page, Commun. Math. Phys. **87**, 577 (1983).
doi:10.1007/BF01208266

- [2] E. Witten, Adv. Theor. Math. Phys. **2**, 505 (1998) doi:10.4310/ATMP.1998.v2.n3.a3 [hep-th/9803131].
- [3] O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas and M. Van Raamsdonk, Adv. Theor. Math. Phys. **8**, 603 (2004) doi:10.4310/ATMP.2004.v8.n4.a1 [hep-th/0310285].
- [4] J. M. Maldacena, Int. J. Theor. Phys. **38**, 1113 (1999) [Adv. Theor. Math. Phys. **2**, 231 (1998)] doi:10.1023/A:1026654312961, 10.4310/ATMP.1998.v2.n2.a1 [hep-th/9711200].
- [5] J. Kinney, J. M. Maldacena, S. Minwalla and S. Raju, Commun. Math. Phys. **275**, 209 (2007) doi:10.1007/s00220-007-0258-7 [hep-th/0510251].
- [6] V. A. Kostelecky and M. J. Perry, Phys. Lett. B **371**, 191 (1996) doi:10.1016/0370-2693(95)01607-4 [hep-th/9512222].
- [7] M. Cvetič, G. W. Gibbons, H. Lu and C. N. Pope, hep-th/0504080.
- [8] J. B. Gutowski and H. S. Reall, JHEP **0402**, 006 (2004) doi:10.1088/1126-6708/2004/02/006 [hep-th/0401042].
- [9] J. B. Gutowski and H. S. Reall, JHEP **0404**, 048 (2004) doi:10.1088/1126-6708/2004/04/048 [hep-th/0401129].
- [10] Z. W. Chong, M. Cvetič, H. Lu and C. N. Pope, Phys. Rev. D **72**, 041901 (2005) doi:10.1103/PhysRevD.72.041901 [hep-th/0505112].
- [11] H. K. Kunduri, J. Lucietti and H. S. Reall, JHEP **0604**, 036 (2006) doi:10.1088/1126-6708/2006/04/036 [hep-th/0601156].
- [12] D. D. K. Chow, Class. Quant. Grav. **27**, 065004 (2010) doi:10.1088/0264-9381/27/6/065004 [arXiv:0808.2728 [hep-th]].
- [13] Z.-W. Chong, M. Cvetič, H. Lu and C. N. Pope, Phys. Lett. B **626**, 215 (2005) doi:10.1016/j.physletb.2005.07.054 [hep-th/0412094].
- [14] D. D. K. Chow, Class. Quant. Grav. **25**, 175010 (2008) doi:10.1088/0264-9381/25/17/175010 [arXiv:0711.1975 [hep-th]].
- [15] S. M. Hosseini, K. Hristov and A. Zaffaroni, JHEP **1707**, 106 (2017) [arXiv:1705.05383 [hep-th]].
- [16] S. M. Hosseini, K. Hristov and A. Zaffaroni, JHEP **1805**, 121 (2018) [arXiv:1803.07568 [hep-th]].
- [17] S. Choi, J. Kim, S. Kim and J. Nahmgoong, arXiv:1810.12067 [hep-th].

- [18] A. Cabo-Bizet, D. Cassani, D. Martelli and S. Murthy, arXiv:1810.11442 [hep-th].
- [19] S. Choi, J. Kim, S. Kim and J. Nahmgoong, arXiv:1811.08646 [hep-th].
- [20] J. Nahmgoong, arXiv:1907.12582 [hep-th].
- [21] S. Choi, C. Hwang and S. Kim, arXiv:1908.02470 [hep-th].
- [22] S. Choi and S. Kim, arXiv:1904.01164 [hep-th].
- [23] Z.-W. Chong, M. Cvetič, H. Lu and C. N. Pope, Nucl. Phys. B **717**, 246 (2005) doi:10.1016/j.nuclphysb.2005.03.034 [hep-th/0411045].
- [24] D. D. K. Chow and G. Compère, Phys. Rev. D **89**, no. 6, 065003 (2014) doi:10.1103/PhysRevD.89.065003 [arXiv:1311.1204 [hep-th]].
- [25] K. Hristov, S. Katmadas and C. Toldo, Phys. Rev. D **100**, no. 6, 066016 (2019) doi:10.1103/PhysRevD.100.066016 [arXiv:1907.05192 [hep-th]].
- [26] N. Seiberg, Phys. Lett. B **388**, 753 (1996) doi:10.1016/S0370-2693(96)01215-4 [hep-th/9608111].
- [27] E. D'Hoker, M. Gutperle, A. Karch and C. F. Uhlemann, JHEP **1608**, 046 (2016) doi:10.1007/JHEP08(2016)046 [arXiv:1606.01254 [hep-th]]; E. D'Hoker, M. Gutperle and C. F. Uhlemann, JHEP **1705**, 131 (2017) doi:10.1007/JHEP05(2017)131 [arXiv:1703.08186 [hep-th]]; E. D'Hoker, M. Gutperle and C. F. Uhlemann, JHEP **1711**, 200 (2017) doi:10.1007/JHEP11(2017)200 [arXiv:1706.00433 [hep-th]].
- [28] F. Apruzzi, M. Fazzi, A. Passias, D. Rosa and A. Tomasiello, JHEP **1411**, 099 (2014) Erratum: [JHEP **1505**, 012 (2015)] doi:10.1007/JHEP11(2014)099, 10.1007/JHEP05(2015)012 [arXiv:1406.0852 [hep-th]].
- [29] H. Kim, N. Kim and M. Suh, Eur. Phys. J. C **75**, no. 10, 484 (2015) doi:10.1140/epjc/s10052-015-3705-1 [arXiv:1506.05480 [hep-th]].
- [30] M. Cvetič, H. Lu and C. N. Pope, Phys. Rev. Lett. **83**, 5226 (1999) [arXiv:hep-th/9906221].
- [31] S. Kim, Nucl. Phys. B **821**, 241 (2009) Erratum: [Nucl. Phys. B **864**, 884 (2012)] doi:10.1016/j.nuclphysb.2012.07.015, 10.1016/j.nuclphysb.2009.06.025 [arXiv:0903.4172 [hep-th]].
- [32] J. Bagger and N. Lambert, Phys. Rev. D **75**, 045020 (2007) doi:10.1103/PhysRevD.75.045020 [hep-th/0611108]; J. Bagger and N. Lambert, Phys. Rev. D **77**, 065008 (2008) doi:10.1103/PhysRevD.77.065008 [arXiv:0711.0955 [hep-th]]; J. Bagger and N. Lambert, JHEP **0802**, 105 (2008) doi:10.1088/1126-6708/2008/02/105 [arXiv:0712.3738 [hep-th]].

- [33] A. Gustavsson, Nucl. Phys. B **811**, 66 (2009) doi:10.1016/j.nuclphysb.2008.11.014 [arXiv:0709.1260 [hep-th]].
- [34] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, JHEP **0810**, 091 (2008) doi:10.1088/1126-6708/2008/10/091 [arXiv:0806.1218 [hep-th]].
- [35] H. C. Kim, S. S. Kim and K. Lee, JHEP **1210**, 142 (2012) doi:10.1007/JHEP10(2012)142 [arXiv:1206.6781 [hep-th]].
- [36] N. A. Nekrasov, Adv. Theor. Math. Phys. **7**, no. 5, 831 (2003) doi:10.4310/ATMP.2003.v7.n5.a4 [hep-th/0206161].
- [37] N. Nekrasov and S. Shadchin, Commun. Math. Phys. **252**, 359 (2004) doi:10.1007/s00220-004-1189-1 [hep-th/0404225].
- [38] C. Hwang, J. Kim, S. Kim and J. Park, JHEP **1507**, 063 (2015) Addendum: [JHEP **1604**, 094 (2016)] doi:10.1007/JHEP07(2015)063, 10.1007/JHEP04(2016)094 [arXiv:1406.6793 [hep-th]].
- [39] M. Cvetič *et al.*, Nucl. Phys. B **558**, 96 (1999) doi:10.1016/S0550-3213(99)00419-8 [hep-th/9903214].
- [40] J. P. Gauntlett, S. Kim, O. Varela and D. Waldram, JHEP **0904**, 102 (2009) doi:10.1088/1126-6708/2009/04/102 [arXiv:0901.0676 [hep-th]].
- [41] S. Lee and D. Yokoyama, JHEP **1503**, 103 (2015) doi:10.1007/JHEP03(2015)103 [arXiv:1412.8703 [hep-th]].
- [42] S. Choi and C. Hwang, arXiv:1911.01448 [hep-th].
- [43] F. Benini, K. Hristov and A. Zaffaroni, JHEP **1605**, 054 (2016) doi:10.1007/JHEP05(2016)054 [arXiv:1511.04085 [hep-th]]; F. Benini, K. Hristov and A. Zaffaroni, Phys. Lett. B **771**, 462 (2017) doi:10.1016/j.physletb.2017.05.076 [arXiv:1608.07294 [hep-th]].
- [44] S. M. Hosseini and A. Zaffaroni, JHEP **1608**, 064 (2016) doi:10.1007/JHEP08(2016)064 [arXiv:1604.03122 [hep-th]]; S. M. Hosseini and N. Mekareeya, JHEP **1608**, 089 (2016) doi:10.1007/JHEP08(2016)089 [arXiv:1604.03397 [hep-th]].
- [45] S. M. Hosseini, I. Yaakov and A. Zaffaroni, arXiv:1808.06626 [hep-th]; S. M. Hosseini, K. Hristov, A. Passias and A. Zaffaroni, arXiv:1809.10685 [hep-th].
- [46] M. Suh, JHEP **1901**, 035 (2019) doi:10.1007/JHEP01(2019)035 [arXiv:1809.03517 [hep-th]]; M. Suh, JHEP **1902**, 108 (2019) doi:10.1007/JHEP02(2019)108 [arXiv:1810.00675 [hep-th]]; M. Suh, arXiv:1812.10491 [hep-th].