

LEBESGUE'S DENSITY THEOREM AND DEFINABLE SELECTORS FOR IDEALS

SANDRA MÜLLER, PHILIPP SCHLICHT, DAVID SCHRITTESSER, AND THILO WEINERT

ABSTRACT. We introduce a notion of density point and prove results analogous to Lebesgue's density theorem for various well-known ideals on Cantor space and Baire space. In fact, we isolate a class of ideals for which our results hold.

In contrast to these results, we show that there is no reasonably definable selector that chooses representatives for the equivalence relation on the Borel sets of having countable symmetric difference. In other words, there is no notion of density which makes the ideal of countable sets satisfy an analogue to the density theorem.

The proofs of the positive results use only elementary combinatorics of trees, while the negative results rely on forcing arguments.

1. INTRODUCTION

Given a σ -ideal I on a Polish space X , consider the equivalence relation $=_I$ on $\mathcal{P}(X)$ defined by $A =_I B \iff A \Delta B \in I$. A map D with $\text{dom}(D), \text{ran}(D) \subseteq \mathcal{P}(X)$ is called a *selector for $=_I$* if $D(A) =_I A$ (i.e., D chooses a representative from the $=_I$ -equivalence class of A) and $A =_I B \Rightarrow D(A) = D(B)$ (i.e., D is invariant with respect to $=_I$) for all $A, B \in \text{dom}(D)$. While a selector for $=_I$ on $\mathcal{P}(X)$ can always be found using the axiom of choice, it is natural to ask: For which σ -ideals is there a selector D with domain the Borel subsets of X that is *definable* in some given sense? For instance, we are interested in Borel measurable and projective selectors.

To make this more precise, let us consider the ideal of Lebesgue null sets on \mathbb{R} and the meager ideal on any Polish space as examples. First we consider the equivalence relation $=_\mu$ on the collection MEAS of Lebesgue measurable subsets of \mathbb{R} , given by equality up to a Lebesgue null set. By Lebesgue's density theorem, we obtain a selector by assigning to each measurable set $A \subseteq \mathbb{R}$ the set $D_\mu(A)$ of points of density 1, as follows: Recall that

$$d_A^\mu(x) = \liminf_{\epsilon \rightarrow 0} \frac{\mu(B_\epsilon(x) \cap A)}{\mu(B_\epsilon(x))}$$

and $D_\mu(A) = \{x \in \mathbb{R} \mid d_A^\mu(x) = 1\}$ for $A \in \text{MEAS}$.¹ Note that in measure theory $D_\mu(A)$ is usually denoted by $\Phi(A)$.

Theorem 1.1 (Lebesgue's density theorem). *For any $A \in \text{MEAS}$, $A =_\mu D_\mu(A)$.*

It follows that $A \mapsto D_\mu(A)$ is a selector with domain MEAS. To gauge the definability of this selector, take recourse to a standard coding of Borel sets (see e.g. [Kec95, 35.B]): Fix a Π_1^1 set $\mathcal{B} \subseteq \mathcal{P}(\omega)$ of Borel codes and Σ_1^1 sets $P, Q \subseteq \mathcal{P}(\omega) \times \mathbb{R}$ such that $\{y \in \mathbb{R} \mid P(x, y)\} = \{y \in \mathbb{R} \mid \neg Q(x, y)\}$ for all $x \in \mathcal{B}$ and every Borel set is of this form. Write B_x for this set, the Borel set *coded by x* . With such a coding, standard arguments on the complexity of measure (see [Kec73, Theorem 2.2.3]) show that D_μ restricted to the Borel sets is induced by a definable map on the Borel codes as follows: There is a map $D: \mathcal{B} \rightarrow \mathcal{B}$ such that

$$B_{D(x)} = D_\mu(B_x)$$

Date: March 17, 2022.

2010 Mathematics Subject Classification. 03E15, 28A05.

Key words and phrases. Density points, ideals, selectors, forcing.

¹See also [ACC19] and [AC19] for recent results on the descriptive set theory of Lebesgue's density theorem.

for all Borel codes $x \in \mathcal{B}$, and the graph of D is a Boolean combination of Σ_1^1 sets. In particular, D is universally Baire measurable.²

The situation is entirely analogous for the Lebesgue measure (i.e. coin-tossing measure) on Cantor space ${}^\omega 2$ (see e.g. [AC13, Section 8] or [Mil08, Proposition 2.10]). It further follows from the isomorphism theorem for measures [Kec95, Theorem 17.41] that a definable selector exists for the null ideal with respect to any Borel probability measure on a standard Borel space.

Similarly, a definable selector (with domain the Borel sets) exists for the *meager ideal* on any Polish space. This is the smallest σ -ideal containing all nowhere dense sets (i.e., sets whose closure has empty interior). When I is the meager ideal we can define a selector by letting

$$D_I(A) = \bigcup \{U \mid U \text{ open, } A \cap U \text{ comeager in } U\}$$

for any set A with the Baire property (i.e., any set which has meager symmetric difference with some Borel set).³

In this article we isolate a class of ideals, the *strongly linked tree ideals* for which likewise there exists a definable selector (cf. Definition 4.15). For this result, we utilize the connection between σ -ideals and *forcing*.⁴ To make our paper as accessible as possible we give a short but self-contained review of this connection in Section 4.

We shall restrict our attention to spaces X of the form ${}^\omega \Omega$ with the product topology, where $\Omega = \{0, 1\}$ or $\Omega = \omega$, i.e., Cantor and Baire space.⁵ A *tree forcing* is a preorder \mathbb{P} , with respect to inclusion, where \mathbb{P} is a collection of perfect subtrees of ${}^{<\omega} \Omega$. A standard construction associates to each tree forcing \mathbb{P} an ideal $I_{\mathbb{P}}$ on ${}^\omega \Omega$ —we call ideals of this form *tree ideals*. Similarly, a family $\text{MEAS}({}^\omega \Omega, \mathbb{P})$ of \mathbb{P} -measurable subsets of ${}^\omega \Omega$ is associated to each tree forcing. For details, see Sections 4.1–4.3.

Recall that \sqsubseteq means *initial segment* and the *stem* of a tree T , denoted by $\text{stem}(T)$, is the maximal node in T that is \sqsubseteq -comparable with all other nodes in T .

If I is a tree-ideal associated to \mathbb{P} define the set of *I -shift density points* (or short, just *I -density points*) of $A \subseteq {}^\omega \Omega$ as follows:

$$D_I(A) = \{x \in {}^\omega \Omega \mid \exists m \in \omega \forall n \geq m \forall T \in \mathbb{P} [\text{stem}(T) = x \restriction n \Rightarrow [T] \cap A \notin I]\}. \quad (1)$$

In Section 4 we prove the following theorem (as Corollary 4.18); this section should be readable without prior knowledge of forcing.

Theorem 1.2. *Let I be a strongly linked tree ideal (cf. Definition 4.15). Then for any $A, B \in \text{MEAS}({}^\omega \Omega, \mathbb{P})$*

- (A) $A =_I B \Rightarrow D_I(A) = D_I(B)$, and
- (B) $D_I(A) =_I A$.

We show in Section 4.6 that D_I has reasonable complexity under the additional assumption that \mathbb{P} is Suslin. Then the map $D_I \restriction \text{BOREL}({}^\omega \Omega)$ is induced by an absolutely- Δ_2^1 and therefore universally Baire measurable map $D: \mathcal{B} \rightarrow \mathcal{B}$ (where $\text{BOREL}({}^\omega \Omega)$ denotes the collection of Borel subsets of ${}^\omega \Omega$ and \mathcal{B} the set of their codes, as described above for \mathbb{R}).

It follows immediately from Theorem 1.2 that D_I is a useful notion of density point for the ideals associated to Cohen forcing, Hechler forcing, eventually different forcing, Laver forcing with a filter, and Mathias forcing with a shift invariant filter (cf. Section 6). Although one can define D_I as in (1) for arbitrary tree ideals, we verify in Section 5.1 that the statement of Theorem 1.2 fails for \mathbb{P} equal to Sacks, Miller, Mathias, Laver, or Silver forcing.

²We refer the reader to [Kec95] for standard definitions in descriptive set theory throughout this paper. We also review the definition of universally Baire in Section 4.6.

³For a different definition of density point which generalizes the standard notion and is at the same time meaningful for the meager ideal, see [PWBW85] (we discuss this in Section 5.3).

⁴This connection has been explored for instance in [BKW18, GRSS95, KSZ13, Ike10, SS18, Zap08].

⁵Note that for any σ -ideal I that contains all singletons on a Polish space X , there is a set $A \in I$ such that $X \setminus A$ is homeomorphic to ${}^\omega \omega$; so this is not a serious restriction.

While the null ideal is *not* a strongly linked tree ideal (see Remark 4.16) our methods also yield a variant of density point for the null ideal. Namely, when $I = I_\mu$ is the ideal of null sets with respect to the Lebesgue measure μ on ${}^\omega 2$ and \mathbb{P} is random forcing (see Definition 6.1(1)) let, for $A \subseteq {}^\omega 2$,

$$D_{I_\mu}(A) = \left\{ x \in {}^\omega \Omega \mid \exists m \in \omega \forall n \geq m \forall T \in \mathbb{P} \left[\frac{\mu([T] \cap N_{x \upharpoonright n})}{\mu(N_{x \upharpoonright n})} > \frac{1}{2} \Rightarrow [T] \cap A \notin I_\mu \right] \right\}. \quad (2)$$

We show in Section 3 that $D_{I_\mu}(A) =_\mu A$ for any μ -measurable set $A \subseteq {}^\omega 2$.

One expects that for many ideals, no definable selector exists; in fact it turns out that this is the case for the smallest non-trivial σ -ideal, the ideal of countable sets. We prove the following result in Theorem 5.5:

Theorem 1.3. *Let I denote the ideal of countable subsets of ${}^\omega \Omega$. There is no selector $\text{BOREL}({}^\omega \Omega) \rightarrow \text{BOREL}({}^\omega \Omega)$ for $=_I$ which is induced by a universally Baire measurable function on Borel codes.*

We extend this result in Lemma 5.15 and Theorem 5.18 to selectors with projective values, assuming the Axiom of Projective Determinacy (PD). To state the next result, let $\text{PROJ}({}^\omega \Omega)$ denote the collection of projective subsets of ${}^\omega \Omega$ (we shall discuss the notion of codes for projective sets in Section 5.2).

Theorem 1.4. *Assuming PD, there is no selector $\text{BOREL}({}^\omega \Omega) \rightarrow \text{PROJ}({}^\omega \Omega)$ for $=_I$ which is induced by a universally Baire measurable function on the codes.*

We further prove in Theorem 5.19 that it is consistent with ZFC that there is no selector which is definable by a first order formula in the language of set theory (with parameter a sequence of ordinals); and that it is consistent with ZF that there is no selector for I at all. In fact there is no such selector in Solovay's model.

We would like to point out two interesting recent results that came to our attention after this paper was submitted. Firstly, there is a definable selector on Δ_2^0 sets for $=_I$, where I denotes the ideal of countable sets [Kan19]. Secondly, the existence of a (not necessarily definable) *lower density operator* (essentially, a selector preserving \cap) on Borel sets is equivalent to the Continuum Hypothesis [BG20].

What we aim for. Our aim is to find dividing lines between ideals with and without a good notion of “density point”. The results show that for all tree forcings listed in Section 6, the following three conditions (a)-(c) are equivalent. Moreover for strongly linked tree forcings, random and Sacks forcing, all four conditions (a)-(d) are equivalent.

- (a) \mathbb{P} is σ -linked.
- (b) \mathbb{P} satisfies the countable chain condition (ccc).
- (c) The analogue of Theorem 1.2 holds for $I = I_\mathbb{P}$ and all $A, B \in \text{BOREL}({}^\omega \Omega)$, with D_I as in Definition 2.3.
- (d) There is a selector $\text{BOREL}({}^\omega \Omega) \rightarrow \text{BOREL}({}^\omega \Omega)$ for $=_{I_\mathbb{P}}$ that is induced by a universally Baire measurable function on the codes.

Whether (d) holds for other well-known forcings is an important question left open in this paper (see Section 8).

Structure of the paper. We introduce density points for ideals in Section 2 and study them for the null ideal in Section 3. In Sections 4.1 to 4.3, we introduce tree ideals and study some of their properties. For instance, we show in Section 4.2 that for any tree forcing \mathbb{P} with the ω_1 -covering property, all Borel sets are \mathbb{P} -measurable. This improves a result of Ikegami [Ike10, Lemma 3.5]. We then prove the main result on strongly linked tree ideals (Theorem 1.2) in Sections 4.4 and 4.5. In Section 4.6, we compute a bound on the complexity of density operators for strongly linked tree ideals and show that they are universally Baire measurable. Section 5.1 contains counterexamples for the remaining tree forcings listed in Section 6. In Section 5.2, we prove that there are no selectors for the ideal of countable

sets which are induced by universally Baire measurable functions (Theorems 1.3 and 1.4). Moreover, we show that it is consistent with ZF that there is no selector at all for the ideal of countable sets. Section 6 contains a list of the tree forcings which we consider in this article. As an additional result of independent interest, we show in Section 7 that one can effectively construct density points of a closed set from a sequence of weights attached to basic open sets. We end with some open questions in Section 8.

Acknowledgments. The authors would like to thank Alessandro Andretta, Martin Goldstern, Daisuke Ikegami and Luca Motto Ros for discussions, Jindra Zapletal for the suggestion to study eventually different forcing and Vladimir Kanovei for reading a previous version of this article and sending us his version of the proof of Theorem 1.3 (see [KL20]). We further thank the anonymous referee for their work and many useful suggestions for improvement.

This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 794020 (IMIC) for the second-listed author and No 706219 (REGPROP) for the first, third, and last-listed authors. The first-listed author, formerly known as Sandra Uhlenbrock, was partially supported by FWF grant number P28157, the second-listed author was partially supported by FWF grant number I4039, the third-listed author by the DNRf Niels Bohr Professorship of Lars Hesselholt as well as FWF grant numbers F29999 and Y1012, and the last-listed author by FWF grants numbers I3081 and Y1012-N35. The first and second-listed authors further wish to thank the Erwin Schrödinger International Institute for Mathematics and Physics (ESI), Vienna for support during the workshop “Current Trends in Descriptive Set Theory” in December 2016.

2. TREES AND DENSITY POINTS

In this section we review some notation and introduce some terminology, including our notion of density point, in a way that allows us to treat *both* D_I , as defined in (1) for strongly linked tree ideals, *and* D_{I_μ} , our variant of the density operator for the null ideal from (2), simultaneously.

Recall that we write Ω to mean either 2 (i.e., $\{0, 1\}$) or ω (i.e., \mathbb{N}). We consider subtrees T of ${}^{<\omega}\Omega$ and write

$$[T] = \{x \in {}^\omega\Omega \mid (\forall n \in \omega) x \restriction n \in T\}$$

for the set of branches through T . A tree T is *perfect* if it has no end nodes and some splitting node above each node. Let

$$s^\frown T = \{s^\frown t \mid t \in T\}$$

$$T_u = \{t \in T \mid u \subseteq t \vee t \subseteq u\}$$

for $s \in {}^{<\omega}\Omega$ and $u \in T$. For $s \in {}^{<\omega}\Omega$ and $U \subseteq {}^{<\omega}\Omega$ write

$$U/s = \{t \in {}^{<\omega}\Omega \mid s^\frown t \in U\}.$$

This is frequently used as T/s when $T \subseteq {}^{<\omega}\Omega$ is a perfect tree or as A/s when $A \subseteq {}^\omega\Omega$. In the latter case, A/s is also called the *detail* or *localization* of A at s and is often denoted by $A_{[s]}$ (see, e.g., [AC13]).

We write

$$\sigma_s : {}^{<\omega}\Omega \rightarrow {}^{<\omega}\Omega, \sigma_s(x) = s^\frown x$$

denote the shift by $s \in {}^{<\omega}\Omega$. Thus $\sigma_s^{-1}[T] = [T/s] = \{t \mid s^\frown t \in [T]\}$. For $s, t \in {}^{<\omega}\Omega$, we write $s \sqsubseteq t$ if s is an initial segment of t . The longest $s \in T$ such that $s \sqsubseteq t$ or $t \sqsubseteq s$ for all nodes $t \in T$ is called the *stem* of T , denoted (as mentioned previously) by $\text{stem}(T)$. The set of *splitting nodes* of T (those with at least two direct successors in T) is denoted split_T . Let $|t|$ denote the *length* of a finite sequence $t \in {}^{<\omega}2$ (note that $|t| = \text{dom}(t)$). Moreover, let $s \wedge t$ denote the longest common initial segment of s and t .

Of course an ideal I on a set X is a collection $I \subseteq \mathcal{P}(X)$ such that $A, B \in I$ implies $A \cup B \in I$ and for any $C \subseteq B \in I$, $C \in I$. A σ -ideal is an ideal which is closed not just under finite, but under countable unions. If I is an ideal on ${}^\omega\Omega$ and A and B are subsets of ${}^\omega\Omega$, recall that we write $A =_I B$ for $A \triangle B \in I$. We also write $A \subseteq_I B$ for $A \setminus B \in I$ and $A \perp_I B$

for $A \cap B \in I$. An ideal on ${}^\omega\Omega$ is called *shift invariant* if it is closed under pointwise images and preimages under σ_t for all $t \in {}^{<\omega}\Omega$. For an ideal I , a set is called *I -positive* if it is not in I ; recall that the set of I -positive sets (the co-ideal of I) is denoted I^+ .

The central notion for this article is the *density property*:

Definition 2.1. If I is an ideal and D is a map into $\mathcal{P}({}^\omega\Omega)$ with $\text{BOREL}({}^\omega\Omega) \subseteq \text{dom}(D) \subseteq \mathcal{P}({}^\omega\Omega)$ and $A =_I B \Rightarrow D(A) = D(B)$, we say that the *density property* holds (with respect to D and I) if $D(A) =_I A$ for all $A \in \text{dom}(D)$.

Ideally, we would like to define a notion of density points relative to an arbitrary shift invariant ideal I on ${}^\omega\Omega$. For our definition we find it necessary to fix a collection \mathbf{L}_I of sets which we consider “large”; but for strongly linked ideals and for the null ideal there is a natural choice of \mathbf{L}_I —namely when $I = I_\mu$, \mathbf{L}_I is defined as the set of perfect sets of measure at least $\frac{1}{2}$ and when $I = I_\mathbb{P}$ and \mathbb{P} is strongly linked, \mathbf{L}_I is defined to be the set $\{[T] \in \mathbb{P} \mid \text{stem}(T) = \emptyset\}$.

Since we want to speak about arbitrary tree ideals in some of our results below, we make the following convention.

Convention 2.2. Let I be a tree ideal, and fix \mathbb{P} such that $I = I_\mathbb{P}$. If $I = I_\mu$ we shall assume that \mathbb{P} is random forcing, i.e. the collection of trees $T \subseteq {}^{<\omega}2$ such that for all $s \in T$, $\mu([T] \cap N_s) > 0$; further, we let

$$\mathbf{L}_I = \{[T] \mid T \in \mathbb{P}, \mu([T]) > \frac{1}{2}\}.$$

If \mathbb{P} is any other tree forcing then we let

$$\mathbf{L}_I = \{[T] \mid T \in \mathbb{P}, \text{stem}(T) = \emptyset\}.$$

We say that elements of \mathbf{L}_I are *large with respect to I* .

Definition 2.3. Suppose that A is a subset of ${}^\omega\Omega$.

- (a) An element x of ${}^\omega\Omega$ is an *I -shift density point* of A if there is some n_x such that for all $B \in \mathbf{L}_I$ and $n \geq n_x$

$$\sigma_{x \upharpoonright n}(B) \cap A \notin I.$$

- (b) $D_I(A)$ denotes the set of I -shift density points of A .

We further say that the *I -shift density property* holds if $D_I(A) \triangle A \in I$ for all Borel sets A .

For simplicity, we sometimes just write *I -density point* and *I -density property*. It is clear that by Convention 2.2, Definition 2.3 just repeats the definition of D_I given in (1) for strongly linked tree ideals as well as the one in (2) for the null ideal. Note that $D_I(A)$ is Σ_2^0 for any subset A of ${}^\omega\Omega$. To see this, let

$$S = \{s \in {}^{<\omega}\Omega \mid \forall B \in \mathbf{L}_I \sigma_s(B) \cap A \notin I\}$$

and observe that $x \in D_I(A) \iff \exists m \forall n \geq m \ x \upharpoonright n \in S$; thus $D_I(A)$ is $\Sigma_2^0(S)$.

Finally, note that in those cases where we verify the *I -shift density property*, we obtain that $D_I(A) \triangle A \in I$ for all \mathbb{P} -measurable⁶ subsets A of ${}^\omega\Omega$ rather than just for Borel sets, since in all these cases \mathbb{P} is ccc (cf. Remark 4.19).

Remark 2.4. Definition 2.3 can be rephrased in the following fashion. We call a subset A of ${}^\omega\Omega$ *I -full* if for all $B \in \mathbf{L}_I$, the set $B \cap A$ is I -positive (this is analogous to the definition of stationary sets from club sets). Then x is a density point of A if and only if $A/(x \upharpoonright n)$ —which by definition is the same as $\sigma_{x \upharpoonright n}^{-1}(A)$ —is eventually I -full as n increases.⁷

⁶ \mathbb{P} -measurability is defined in Section 4.2.

⁷As mentioned above, the localization A/s of A at s is often denoted by $A_{[s]}$ (see e.g. [AC13]).

Remark 2.5. We notice that Definition 2.3 can also be rephrased via the following notion of convergence. We say that a sequence $\vec{f} = \langle f_n \mid n \in \omega \rangle$ of functions $f_n: {}^\omega\Omega \rightarrow \mathbb{R}$ *converges in* I to a function $f: {}^\omega\Omega \rightarrow \mathbb{R}$ if the following condition holds:⁸ For all $\epsilon > 0$, there is some n_0 such that for all $B \in \mathbf{L}_I$ and $n \geq n_0$,

$$B \setminus \{x \in {}^\omega\Omega \mid |f_n(x) - f(x)| \geq \epsilon\} \notin I.$$

By shift invariance, the condition $\sigma_{x \upharpoonright n}(B) \cap A \notin I$ in Definition 2.3 (a) is equivalent to $B \cap \sigma_{x \upharpoonright n}^{-1}(A) \notin I$. Moreover, $B \cap \sigma_{x \upharpoonright n}^{-1}(A) = B \setminus \{x \in {}^\omega\Omega \mid |1_{\sigma_{x \upharpoonright n}^{-1}(A)}(x) - 1| \geq \epsilon\}$ for any ϵ with $0 < \epsilon < 1$. Therefore x is an I -shift density point of A if and only if the sequence $\langle 1_{\sigma_{x \upharpoonright n}^{-1}(A)} \mid n \in \omega \rangle$ of characteristic functions of $\sigma_{x \upharpoonright n}^{-1}(A)$ converges in I to the constant function with value 1.

3. THE NULL IDEAL

We now outline the situation in the special case of the σ -ideal of Lebesgue null subsets of ${}^\omega 2$ to illustrate some ideas used in this paper.

Recall that μ denotes Lebesgue measure on ${}^\omega 2$ and I_μ the σ -ideal of μ -null sets. The next lemma implies that the I_μ -shift density property holds.

Lemma 3.1. *Let x be an element and A a Borel subset of ${}^\omega 2$.*

- (1) *If $d_A^\mu(x) = 1$ and $\epsilon > 0$, then there is some $n_{x,\epsilon}$ such that for all $n \geq n_{x,\epsilon}$ and all Borel sets B with $\mu(B) \geq \epsilon$, $\mu(\sigma_{x \upharpoonright n}(B) \cap A) > 0$.*
- (2) *If $d_A^\mu(x) = 0$ and $\epsilon > 0$, then there is a Borel set B with*

$$\exists^\infty n \ (\sigma_{x \upharpoonright n}(B) \cap A = \emptyset)$$

and $\mu(B) \geq 1 - \epsilon$.

Proof. For the first claim, note that there is some $n_{x,\epsilon}$ with $\frac{\mu(A \cap N_{x \upharpoonright n})}{\mu(N_{x \upharpoonright n})} > 1 - \epsilon$ for all $n \geq n_{x,\epsilon}$, since

$$d_A^\mu(x) = \liminf_n \frac{\mu(A \cap N_{x \upharpoonright n})}{\mu(N_{x \upharpoonright n})} = 1.$$

If B is any I_μ -positive Borel set of size at least ϵ , then $\mu(\sigma_{x \upharpoonright n}(B) \cap A) > 0$ for all $n \geq n_{x,\epsilon}$.

For the second claim, let $\vec{\epsilon} = \langle \epsilon_i \mid i < \omega \rangle$ be a sequence in \mathbb{R}^+ with $\sum_i \epsilon_i \leq \epsilon$. Since

$$d_A^\mu(x) = \liminf_n \frac{\mu(A \cap N_{x \upharpoonright n})}{\mu(N_{x \upharpoonright n})} = 0,$$

there is a strictly increasing sequence $\vec{n} = \langle n_i \mid i \in \omega \rangle$ with $\frac{\mu(A \cap N_{x \upharpoonright n_i})}{\mu(N_{x \upharpoonright n_i})} < \epsilon_i$ for all $i \in \omega$. Let $B_i = \sigma_{x \upharpoonright n_i}^{-1}(A)$ for $i \in \omega$ and $B = \bigcup_{i \in \omega} B_i$. Then

$$\mu(B) \leq \sum_i \mu(B_i) \leq \sum_i \frac{\mu(A \cap N_{x \upharpoonright n_i})}{\mu(N_{x \upharpoonright n_i})} \leq \sum_i \epsilon_i \leq \epsilon.$$

Let C be an I_μ -positive set with $\mu(C) \geq 1 - \epsilon$ that is disjoint from B . Then $C \cap B_i = \emptyset$ for all $i \in \omega$. Since $B_i = \sigma_{x \upharpoonright n_i}^{-1}(A)$, it follows that $\sigma_{x \upharpoonright n_i}(C) \cap A = \sigma_{x \upharpoonright n_i}(C) \cap \sigma_{x \upharpoonright n_i}(B_i) = \sigma_{x \upharpoonright n_i}(C \cap B_i) = \emptyset$. \square

For $\epsilon = \frac{1}{2}$, we obtain that $d_A^\mu(x) = 1$ implies that x is an I_μ -shift density point and $d_A^\mu(x) = 0$ implies that this fails. Using Lebesgue's density theorem for μ and ${}^\omega 2$ (see [AC13, Section 8] or [Mil08, Proposition 2.10]) this yields the I_μ -shift density property (cf. Definition 2.3).

Corollary 3.2. *For every Lebesgue measurable subset A of ${}^\omega 2$, $D_{I_\mu}(A) = {}_\mu D_\mu(A)$, recalling that by $D_\mu(A)$ we denote the points of A of density 1 (see p. 1). In particular, the I_μ -shift density property holds.*

⁸Compare this with convergence in measure as in Lemma 5.21.

We shall give another proof of Lebesgue's density theorem in Section 7 below, thus making the previous argument for the I_μ -shift density property self-contained. The I_μ -density property also follows as a special case from the results in Section 4.5.

In the next lemma, we give two examples which show that if $d_A^\mu(x) \in (0, 1)$, then x can but does not have to be an I_μ -shift density point of A .

Lemma 3.3. *Each of the following statements is satisfied by some Borel subset A of ${}^\omega 2$ and some $x \in {}^\omega 2$ with $d_A^\mu(x) \in (0, 1)$.*

- (a) x is an I_μ -shift density point of A .
- (b) x is not an I_μ -shift density point of A .

Proof. Let $A = \{0^n \smallfrown 1^3 \smallfrown x \in {}^\omega 2 \mid n \in \omega, x \in {}^\omega 2\}$ and B its complement.

For (a) note that $d_B^\mu(0^\omega) \in (0, 1)$ since $\frac{\mu(B \cap N_{0^n})}{\mu(N_{0^n})} = \frac{3}{4}$ for all $n \in \omega$. Thus $\mu(\sigma_{0^n}(C) \cap B) > 0$ for any Borel set C with $\mu(C) \geq \frac{1}{2}$. So 0^ω is an I_μ -shift density point of B .

For (b) we have $d_A^\mu(0^\omega) = 1 - d_B^\mu(0^\omega) \in (0, 1)$. Since $\mu(B) \geq \frac{1}{2}$ and $\sigma_{0^n}(B) \cap A = \emptyset$ for all $n \in \omega$, 0^ω is not an I_μ -shift density point of A . \square

4. TREE IDEALS

In this section, we study ideals induced by collections of trees. We introduce the class of strongly linked tree ideals and show that the shift density property holds for this class. Recall again that we work in the Polish space ${}^\omega \Omega$ where Ω is either 2 (i.e., $\{0, 1\}$) or ω (i.e., \mathbb{N}).

4.1. What is a tree ideal? A tree ideal on ${}^\omega \Omega$ is induced by a collection \mathbb{P} of perfect subtrees of ${}^{<\omega} \Omega$ that contains ${}^{<\omega} \Omega$ and T_s for all $T \in \mathbb{P}$ and $s \in T$. We will always assume this condition for any collection of trees.

Any such collection of trees carries the partial order $S \leq T : \iff S \subseteq T \iff [S] \subseteq [T]$. Such partial orders are also called *tree forcings*,⁹ some well-known examples are listed in Section 6. For instance, the null ideal is induced by the collection of random trees, given as follows:

Example 4.1. A subtree T of ${}^{<\omega} 2$ is *random* if $\mu([T_s]) > 0$ for all $s \in T$ with $\text{stem}(T) \sqsubseteq s$.

We next associate an ideal to any collection of trees (we follow [Ike10, Definition 2.6]). The underlying idea is based on the special case that the sets $[T]$ for $T \in \mathbb{P}$ form a base for a topology. In this case, \mathbb{P} is called *topological* and its topology is denoted $\tau_\mathbb{P}$. For instance, the collection of Hechler trees (see Definition 6.1 (5)) is topological. In fact, all strongly linked collections of trees (as defined in Section 4.5) have this property.¹⁰

Usually, one defines nowhere dense sets relative to a given topology, or equivalently, to a base of that topology. Moreover, meager sets are defined as countable unions of these sets. In the next definition, these notions are generalized by replacing a base by an arbitrary collection of trees.

Definition 4.2. [Ike10, Definition 2.6] Let \mathbb{P} be a collection of trees.

- (a) A set A is \mathbb{P} -*nowhere dense* if for all $T \in \mathbb{P}$ there is some $S \leq T$ with $[S] \cap A = \emptyset$.
Moreover, $N_\mathbb{P}$ is the ideal of \mathbb{P} -nowhere dense sets.
- (b) $I_\mathbb{P}$ is the σ -ideal of \mathbb{P} -*meager sets* generated by $N_\mathbb{P}$.

Tree ideals are those of the form $I_\mathbb{P}$ for a collection \mathbb{P} of trees.¹¹ This presentation allows for uniform proofs of results for various ideals. Moreover, many well-known ideals are of this form; for instance, for Cohen forcing¹² $\tau_\mathbb{P}$ is the standard topology, so $N_\mathbb{P}$ is the collection

⁹These forcings, but without the condition that ${}^\omega \Omega$ is in \mathbb{P} , are called *strongly arboreal* in [Ike10, Definition 2.4].

¹⁰Topological does not imply the density property. For instance, assuming CH one can construct a dense topological (shift-invariant) subforcing of Sacks forcing, while we show in Proposition 5.1 and Theorem 5.5 that the density property fails for the ideal associated to Sacks forcing.

¹¹In [BL99, Section 2] and various other papers, tree ideals mean the ideals $N_\mathbb{P}$ instead of $I_\mathbb{P}$.

¹²See Section 6 for this and the following forcings.

of nowhere dense sets and $I_{\mathbb{P}}$ that of meager sets. For random forcing, $N_{\mathbb{P}}$ and $I_{\mathbb{P}}$ equal the σ -ideal of null sets. Sacks forcing is the collection of all perfect trees; here both ideals equal the *Marczewski ideal* (see [Szp35, 3.1] or for a more modern presentation, [Buk11, p. 306]). Its restriction to Borel sets equals the ideal of countable sets by the perfect set property for Borel sets. For Mathias forcing, $\tau_{\mathbb{P}}$ is the Ellentuck topology, and $N_{\mathbb{P}}$ and $I_{\mathbb{P}}$ are equal to the ideal of nowhere Ramsey sets (see [BKW18]). The ideal associated to Silver forcing consists of the completely doughnut null sets (see [Hal03]).

4.2. Measurability for tree ideals. Let \mathbb{P} be a collection of subtrees of ${}^{<\omega}\Omega$. The next definition introduces a form of indivisibility¹³ of \mathbb{P} with respect to A : If $T \in \mathbb{P}$ and $[T]$ is split into the two pieces $[T] \cap A$ and $[T] \setminus A$, then at least one of these pieces contains a set of the form $[S]$ for some $S \in \mathbb{P}$, up to some \mathbb{P} -meager set.

Definition 4.3. [Ike10, Definition 2.8]¹⁴ A subset A of ${}^\omega\Omega$ is called \mathbb{P} -*measurable* if for every $T \in \mathbb{P}$, there is some $S \leq T$ with at least one of the properties (a) $[S] \subseteq_{I_{\mathbb{P}}} A$ and (b) $[S] \perp_{I_{\mathbb{P}}} A$. The collection of \mathbb{P} -measurable sets is denoted $\text{MEAS}({}^\omega\Omega, \mathbb{P})$.

Note that the properties (a) and (b) are mutually exclusive (see Lemma 4.10 below).

The main motivation for introducing this notion is the fact that it formalizes various well-known properties. For instance, we will see in Lemma 4.8 that \mathbb{P} -measurability for random forcing means Lebesgue measurability. For Sacks forcing it is equivalent to the Bernstein property for collections of sets closed under continuous preimages and intersections with closed sets [BL99, Lemma 2.1]. Moreover, for Mathias forcing it is equivalent to being completely Ramsey.

Our next goal is to show that for a very large class of forcings, all Borel sets are \mathbb{P} -measurable. This will be important in the proofs of the following sections.

[Ike10, Lemma 3.5] shows that for proper tree forcings \mathbb{P} , all Borel sets are \mathbb{P} -measurable.¹⁵ We will show a slightly more general version of this result. To state this, recall that a forcing \mathbb{P} has the ω_1 -*covering property* if for any \mathbb{P} -generic filter G over V , any countable set of ordinals in $V[G]$ is covered by (i.e., is a subset of) a set in V that is countable in V . For instance, this statement holds for all proper and thus for all Axiom A forcings.¹⁶ In particular, it holds for all forcings considered in this paper.

We will need the following characterization of the ω_1 -covering property. To state it, we introduce the following notation: For $D \subseteq \mathbb{P}$ and $p \in \mathbb{P}$ let us write

$$D^{\parallel p} = \{q \in D \mid p \parallel q\},$$

where $p \parallel q$ denotes that there is an $r \in \mathbb{P}$ such that $r \leq p$ and $r \leq q$.

Lemma 4.4. *The following conditions are equivalent for any forcing \mathbb{P} :*

- (a) \mathbb{P} has the ω_1 -covering property.
- (b) *For any condition $p \in \mathbb{P}$ and any sequence $\vec{D} = \langle D_n \mid n \in \omega \rangle$ of antichains in \mathbb{P} , there is some $q \leq p$ such that for any $n \in \omega$, the set $D_n^{\parallel q}$ is countable.*

Proof. We first assume (a). Let $p \in \mathbb{P}$ and let $\vec{D} = \langle D_n \mid n \in \omega \rangle$ be as in (b). Take a \mathbb{P} -generic filter G over V . Moreover, let $f(n)$ be an element of $D_n \cap G$ for each $n \in \omega$. By the ω_1 -covering property, there is a countable subset $C \in V$ of \mathbb{P} such that $f(n) \in C$ for all $n \in \omega$. Let \dot{f} be a name for f such that $q \leq p$ forces $\dot{f}(n) \in \check{C}$ for all $n \in \omega$. It follows that $D_n^{\parallel q} \subseteq C$, since for any \mathbb{P} -generic filter H over V that contains both q and r we have $r = \dot{f}^H(n) \in C$.

For the converse implication, assume (b) and suppose that G is \mathbb{P} -generic over V and C is a countable set of ordinals in $V[G]$. Moreover, let f be an enumeration of C and \dot{f} a name with $\dot{f}^G = f$. Then there is a condition $p \in G$ which forces that $\dot{f}: \omega \rightarrow \text{Ord}$ is a function. For

¹³See e.g. [LNVTPS11] for the concept of indivisibility in combinatorics.

¹⁴This is a variant of a definition in [BHL05, Section 0].

¹⁵The proof of this and several other results in [Ike10] can also be done from the weaker assumption that \mathbb{P} has the ω_1 -covering property. We give a more direct proof.

¹⁶See [Jec03, Definition 31.10].

each $n \in \omega$, let D_n be a maximal antichain of conditions deciding $\dot{f}(n)$. By our assumption, there are densely many conditions $q \leq p$ as in (b). Hence there is some $q \in G$ as in (b). Since $D_n^{\parallel q}$ is countable for all $n \in \omega$, $C_n = \{\alpha \mid r \Vdash \dot{f}(n) = \alpha \text{ for some } r \leq q, q' \text{ for a } q' \in D_n^{\parallel q}\}$ is countable and hence $C = \bigcup_{n \in \omega} C_n$ is a countable cover of $\text{ran}(f)$. \square

To show that all Borel sets are \mathbb{P} -measurable if \mathbb{P} has the ω_1 -covering property, we need the next two easy lemmas. We will use the following notation. If A is a subset of ${}^\omega\Omega$, let $\mathbb{P}_A = \{T \in \mathbb{P} \mid [T] \subseteq_{I_{\mathbb{P}}} A\}$. We further say that a subset of \mathbb{P} is *A-good* if it is contained in $\mathbb{P}_{(A)} = \mathbb{P}_A \cup \mathbb{P}_{\omega\Omega \setminus A}$.

Lemma 4.5. *A subset A of ${}^\omega\Omega$ is \mathbb{P} -measurable if and only if there is an A -good maximal antichain in \mathbb{P} .*

Proof. If A is \mathbb{P} -measurable, then $\mathbb{P}_{(A)}$ is a dense subset of \mathbb{P} . Then there is a maximal antichain in \mathbb{P} contained in $\mathbb{P}_{(A)}$ and hence an A -good maximal antichain. Conversely, if D is a maximal A -good antichain in \mathbb{P} and $S \in \mathbb{P}$, let $T \in D^{\parallel S}$ and $R \leq S, T$. Then $[R] \subseteq_{I_{\mathbb{P}}} A$ or $[R] \subseteq_{I_{\mathbb{P}}} \omega\Omega \setminus A$. \square

If $D \subseteq \mathbb{P}$, write

$$\bigcup^\square D = \bigcup_{T \in D} [T].$$

Lemma 4.6. *If D is a maximal antichain in \mathbb{P} and $T \in \mathbb{P}$, then $[T] \setminus \bigcup^\square D^{\parallel T} \in \mathbb{N}_{\mathbb{P}}$.*

Proof. Let $S \in \mathbb{P}$, $R \in D^{\parallel S}$ and $Q \leq R, S$. If $Q \perp T$, then there is some $P \leq Q$ with $[P] \cap [T] = \emptyset$ by the closure property of \mathbb{P} defined in the beginning of Section 4.1. If $Q \parallel T$, let $P \leq Q, T$. Since $P \in D^{\parallel T}$, $[P]$ is disjoint from $[T] \setminus \bigcup^\square D^{\parallel T}$, as required. \square

The next result shows that Borel sets are \mathbb{P} -measurable in all relevant cases.

Lemma 4.7. *If \mathbb{P} has the ω_1 -covering property, then the \mathbb{P} -measurable sets form a σ -algebra. In particular, all Borel sets are \mathbb{P} -measurable.*

Proof. For the first claim, it suffices to show that the class of \mathbb{P} -measurable sets is closed under forming countable unions. To this end, let $\vec{A} = \langle A_n \mid n \in \omega \rangle$ be a sequence of \mathbb{P} -measurable subsets of ${}^\omega\Omega$. Furthermore, let D_n be an A_n -good maximal antichain for each $n \in \omega$ by Lemma 4.5. We will show that $A = \bigcup_{n \in \omega} A_n$ is \mathbb{P} -measurable.

Fix any $T \in \mathbb{P}$. Since \mathbb{P} has the ω_1 -covering property, there is some $S \leq T$ such that the sets $E_n = D_n^{\parallel S}$ are countable for all $n \in \omega$ by Lemma 4.4. First assume that for some $n \in \omega$, there is a tree $R \in E_n$ with $[R] \subseteq_{I_{\mathbb{P}}} A_n$. Then there is some $Q \leq S$ with $[Q] \subseteq_{I_{\mathbb{P}}} A_n \subseteq A$ as required. So we can assume that the previous assumption fails. We claim that then $[S] \perp_{I_{\mathbb{P}}} A$. It suffices to show that $[S] \cap A_n \in I_{\mathbb{P}}$ for each $n \in \omega$, since $I_{\mathbb{P}}$ is a σ -ideal. To see this, note that $[R] \cap A_n \in I_{\mathbb{P}}$ for all $R \in E_n$ by our case assumption. Hence $\bigcup^\square E_n \cap A_n \in I_{\mathbb{P}}$. Moreover, $[S] \setminus \bigcup^\square E_n \in \mathbb{N}_{\mathbb{P}}$ by Lemma 4.6 and therefore $[S] \cap A_n \in I_{\mathbb{P}}$.

Since it is easy to see that closed sets are \mathbb{P} -measurable, it follows that all Borel sets are \mathbb{P} -measurable. \square

Next is the observation that \mathbb{P} -measurability for random forcing means Lebesgue measurability. We recall the argument from [Ike10] for the benefit of the reader.

Lemma 4.8. [Ike10, Proposition 2.9] *If \mathbb{P} is a ccc tree forcing and A is any subset of ${}^\omega\Omega$, then the following conditions are equivalent.*

- (a) A is \mathbb{P} -measurable.
- (b) There is a Borel set B with $A \triangle B \in I_{\mathbb{P}}$.

Proof. If A is \mathbb{P} -measurable, then $\mathbb{P}_{(A)}$ is dense in \mathbb{P} . Let $D \subseteq \mathbb{P}_{(A)}$ be a maximal antichain in \mathbb{P} . Since D is countable, the sets $B_0 = \bigcup^\square (D \cap \mathbb{P}_A) \subseteq_{I_{\mathbb{P}}} A$ and $B_1 = \bigcup^\square (D \setminus \mathbb{P}_A) \subseteq_{I_{\mathbb{P}}} \omega\Omega \setminus A$ are Borel. Since D is maximal, $\omega\Omega \setminus \bigcup^\square D \in \mathbb{N}_{\mathbb{P}}$ by Lemma 4.6. Thus $A \triangle B_0 \in I_{\mathbb{P}}$.

The reverse implication follows from the fact that all Borel sets are \mathbb{P} -measurable by Lemma 4.7. \square

4.3. Positive Borel sets. The following characterization of positive Borel sets via trees will be important below. It uses an auxiliary ideal from [Ike10].

Definition 4.9. [Ike10, Definitions 2.11] $A \in I_{\mathbb{P}}^*$ if for all $T \in \mathbb{P}$, there is some $S \leq T$ with $[S] \cap A \in I_{\mathbb{P}}$.

It is clear that $I_{\mathbb{P}} \subseteq I_{\mathbb{P}}^*$, but it is open whether equality holds for all proper tree forcings.¹⁷ Note that equality holds for ccc forcings. To see this, assume that $A \in I_{\mathbb{P}}^*$. Then $\mathbb{P}_{\omega\Omega \setminus A}$ (as defined before Lemma 4.5) is dense in \mathbb{P} and therefore contains a (countable) maximal antichain D . We have $A \cap \bigcup^\square D \in I_{\mathbb{P}}$.¹⁸ Since $\omega\Omega \setminus \bigcup^\square D \in N_{\mathbb{P}}$ by Lemma 4.6, we have $A \in I_{\mathbb{P}}$. Moreover, equality holds for fusion forcings (then in fact $N_{\mathbb{P}} = I_{\mathbb{P}}$)¹⁹ and for topological tree forcings as in the proof of [FKK16, Lemma 3.8].

The next lemma characterizes $I_{\mathbb{P}}^*$ -positive sets.

Lemma 4.10. *Suppose that \mathbb{P} is a tree forcing.*

- (1) *For any $T \in \mathbb{P}$, $[T] \notin I_{\mathbb{P}}^*$.*
- (2) *A \mathbb{P} -measurable subset A of $\omega\Omega$ is $I_{\mathbb{P}}^*$ -positive if and only if there is some $T \in \mathbb{P}$ with $[T] \subseteq_{I_{\mathbb{P}}^*} A$.*

Proof. We show the first claim. If $[T] \in I_{\mathbb{P}}^*$, then $[S] \in I_{\mathbb{P}}$ for some $S \leq T$. Let $\vec{A} = \langle A_n \mid n \in \omega \rangle$ be a sequence of sets in $N_{\mathbb{P}}$ with $[S] \subseteq \bigcup_{n \in \omega} A_n$. We can then recursively construct a sequence $\vec{S} = \langle S_n \mid n \in \omega \rangle$ in \mathbb{P} such that $S_0 = S$, $S_{n+1} \subseteq S_n$, $[S_n] \cap A_n = \emptyset$ for all $n \in \omega$ and the sequence $\vec{s} = \langle \text{stem}(S_n) \mid n \in \omega \rangle$ of stems is strictly increasing. Then $x = \bigcup_{n \in \omega} \text{stem}(S_n) \in \bigcap_{n \in \omega} [S_n]$. Since $[S_n] \cap A_n = \emptyset$ for all $n \in \omega$ and $[S] \subseteq \bigcup_{n \in \omega} A_n$, we have $x \notin [S]$. But this contradicts the fact that $x \in [S]$.

We now show the second claim. By the first part, it is sufficient to take any \mathbb{P} -measurable $I_{\mathbb{P}}^*$ -positive set A and find some $S \in \mathbb{P}$ with $[S] \subseteq_{I_{\mathbb{P}}^*} A$. Assume that there is no such tree. Since A is \mathbb{P} -measurable, we have that for any $T \in \mathbb{P}$ there is some $S \leq T$ with $[S] \perp_{I_{\mathbb{P}}} A$. Hence $A \in I_{\mathbb{P}}$ by the definition of $I_{\mathbb{P}}^*$. \square

For ideals I of the form $I_{\mathbb{P}}^*$ such that \mathbb{P} has the ω_1 -covering property, the definition of I -shift density points (Definition 2.3) of a Borel set A can now be formulated in the following way: An element x of $\omega\Omega$ is an I -shift density point of A if there is some n_x such that for all $B \in \mathbf{L}_I$ and $n \geq n_x$, there is some $T \in \mathbb{P}$ with

$$[T] \subseteq_I \sigma_{x \upharpoonright n}(B) \cap A.$$

Note that it is easy to see that \mathbb{P} -measurability remains equivalent if $I_{\mathbb{P}}$ is replaced with $I_{\mathbb{P}}^*$. Moreover, the ideals $N_{\mathbb{P}}$, $I_{\mathbb{P}}$ and $I_{\mathbb{P}}^*$ remain the same if \mathbb{P} is replaced by a dense subset by the next remark.

Remark 4.11. \mathbb{P} is dense in \mathbb{Q} if for every $T \in \mathbb{P}$, there is some $S \leq T$ in \mathbb{Q} . We define \mathbb{P} and \mathbb{Q} to be *mutually dense* if \mathbb{P} is dense in \mathbb{Q} and conversely.

We claim that $N_{\mathbb{P}} = N_{\mathbb{Q}}$, $I_{\mathbb{P}} = I_{\mathbb{Q}}$ and $I_{\mathbb{P}}^* = I_{\mathbb{Q}}^*$ if \mathbb{P} and \mathbb{Q} are mutually dense. To see that $N_{\mathbb{P}} \subseteq N_{\mathbb{Q}}$, take any $A \in N_{\mathbb{P}}$ and $T \in \mathbb{Q}$. As \mathbb{P} is dense in \mathbb{Q} , there is some $T' \leq T$ in \mathbb{P} . Since $A \in N_{\mathbb{P}}$, there is some $S \leq T'$ in \mathbb{P} with $[S] \cap A = \emptyset$. As \mathbb{Q} is dense in \mathbb{P} , there is some $S' \leq S$ with $S' \in \mathbb{Q}$ such that $[S'] \cap A \subseteq [S] \cap A = \emptyset$. Thus $N_{\mathbb{P}} = N_{\mathbb{Q}}$ and $I_{\mathbb{P}} = I_{\mathbb{Q}}$. A similar argument shows $I_{\mathbb{P}}^* = I_{\mathbb{Q}}^*$.

¹⁷To our knowledge, every known proper tree forcing satisfies either the ccc or fusion and thus equality holds.

¹⁸This notation is defined before Lemma 4.6.

¹⁹For a tree forcing \mathbb{P} , we define *fusion* as the existence of a sequence $\vec{\leq} = \langle \leq_n \mid n \in \omega \rangle$ of partial orders on \mathbb{P} with $\leq_0 = \leq$ which satisfy the following conditions:

- (a) (*decreasing*) If $S \leq_n T$ and $m \leq n$, then $S \leq_m T$.
- (b) (*limit*) If $\vec{T} = \langle T_n \mid n \in \omega \rangle$ is a sequence in \mathbb{P} with $T_{n+1} \leq_n T_n$ for all $n \in \omega$, then there is some $S \in \mathbb{P}$ with $S \leq_n T_n$ for all $n \in \omega$.
- (c) (*covering*) If $T \in \mathbb{P}$, $n \in \omega$ and D is dense below T , then there is some $S \leq_n T$ with $[S] \subseteq \bigcup^\square D$.

Conversely, any two collections of trees \mathbb{P} and \mathbb{Q} with the ω_1 -covering property and $I_{\mathbb{P}}^* = I_{\mathbb{Q}}^*$ are mutually dense by Lemmas 4.7 and 4.10.

4.4. An equivalence to the density property. Let I be an ideal on ${}^\omega\Omega$. We say that a function $D: \text{BOREL}({}^\omega\Omega) \rightarrow \text{BOREL}({}^\omega\Omega)$ is *lifted from* $\text{BOREL}({}^\omega\Omega)/I$ if for any $A, B \in \text{BOREL}({}^\omega\Omega)$, $A =_I B \implies D(A) = D(B)$.

For instance, D_I as introduced in Definition 2.3 is lifted from $\text{BOREL}({}^\omega\Omega)/I$ if I is shift invariant (see p. 5). Then in fact $A \subseteq_I B \implies D_I(A) \subseteq D_I(B)$ by the definition of D_I .

Recall that the density property holds for D and I if $D(A) =_I A$ for all Borel sets A (see Definition 2.1). We now give a useful condition for proving this for specific ideals. To state this condition, we say that D is *I -compatible* if $A \subseteq_I B \implies D(A) \subseteq_I D(B)$ and $A \perp_I B \implies D(A) \perp_I D(B)$ for all Borel sets A and B . We further say that D is *I -positive* if $D(A) \cap A$ is I -positive for all I -positive Borel sets A .

Proposition 4.12. *If $D: \text{BOREL}({}^\omega\Omega) \rightarrow \text{BOREL}({}^\omega\Omega)$ is a function that is lifted from $\text{BOREL}({}^\omega\Omega)/I$, then the following statements are equivalent.*

- (a) *D is I -compatible and I -positive.*
- (b) *The density property holds for D and I .*

Proof. It is clear that the I -density property implies that D is I -positive and I -compatible. To see that these conditions imply the density property, take any Borel set A . We aim to show that $D(A) =_I A$.

We first show that $B_0 = A \setminus D(A)$ is in I . Towards a contradiction, assume that B_0 is I -positive. Then $D(B_0) \setminus D(A)$ is also I -positive, since it contains $D(B_0) \cap B_0$ as a subset, and the latter is I -positive since D is I -positive. On the other hand, we have $D(B_0) \setminus D(A) \in I$ since $B_0 \subseteq A$ and D is I -compatible, contradiction.

It remains to show that $B_1 = D(A) \setminus A$ is in I . Assume that B_1 is I -positive, so that in particular $D(A)$ is I -positive. The set $C = D(B_1) \cap B_1$ is I -positive, since D is I -positive. We thus obtain $C \perp_I D(A)$, as $C \subseteq D(B_1)$ and we have $D(B_1) \perp_I D(A)$ since $B_1 \perp_I A$ (in fact B_1 and A are disjoint) and D is I -compatible. However, this contradicts the fact that C is I -positive and $C \subseteq B_1 \subseteq D(A)$. \square

4.5. The density property for strongly linked tree ideals. To obtain the density property for $D_{I_{\mathbb{P}}}$, we will make two modest assumptions on \mathbb{P} . Let \mathbf{K}_I denote a fixed subset of \mathbb{P} coding \mathbf{L}_I (from Convention 2.2) in the sense that $\mathbf{L}_I = \{[T] \mid T \in \mathbf{K}_I\}$.

Definition 4.13. We say $(\mathbb{P}, \mathbf{K}_I)$ has the *stem property* if for all $T \in \mathbb{P}$ and $I_{\mathbb{P}}^*$ -almost all $x \in [T]$, there are infinitely many $n \in \omega$ such that there is some $S \leq T$ with $x \in [S]$ and $S/(x \upharpoonright n) \in \mathbf{K}_I$. Since \mathbf{K}_I is fixed for each \mathbb{P} , we may also just say \mathbb{P} has the stem property.

The condition is trivially true for all $x \in [T]$ when $\mathbf{K}_I = \{T \in \mathbb{P} \mid \text{stem}(T) = \emptyset\}$ and we only introduce it to deal with the case of random forcing. For random forcing, recall $\mathbf{K}_I = \{T \in \mathbb{P} \mid \mu([T]) > \frac{1}{2}\}$. Then the stem property follows from Theorem 7.1 or Lebesgue's density theorem.

The next lemma shows that D_I is I -compatible for $I = I_{\mathbb{P}}^*$ provided that D_I is I -positive and \mathbb{P} has the stem property.

Lemma 4.14. *Assume that \mathbb{P} is a collection of trees with the stem property, all Borel sets are \mathbb{P} -measurable, $I = I_{\mathbb{P}}^*$ and $D_I([T]) \cap [T] \notin I$ for all $T \in \mathbb{P}$. Then $D_I(A) \cap D_I(B) =_I D_I(A \cap B)$ for all Borel sets A, B . In particular, D_I is I -compatible.*

Proof. It is easy to see that $D_I(A \cap B) \subseteq D_I(A) \cap D_I(B)$, so suppose towards a contradiction that there are Borel sets A, B with $C = (D_I(A) \cap D_I(B)) \setminus D_I(A \cap B) \notin I$. Since C is Borel (it is a Boolean combination of Σ_2^0 sets) and hence \mathbb{P} -measurable, by Lemma 4.10 there is some $S_0 \in \mathbb{P}$ with $[S_0] \subseteq_I C$.

Since $[S_0] \subseteq_I D_I(A)$ we can pick $x \in [S_0] \cap D_I(A)$ witnessing the stem property for S_0 . Let $n_x \in \omega$ witness that $x \in D_I(A)$. By the choice of x , there is some $m \geq n_x$ and $S_1 \leq S_0$

with $S_1/(x \upharpoonright m) \in \mathbf{K}_I$. Since $m \geq n_x$, $[S_1] \cap A \notin I$. By Lemma 4.10, there is some $S_2 \in \mathbb{P}$ with $[S_2] \subseteq_I [S_1] \cap A$.

Thus $[S_2] \subseteq_I C \subseteq D_I(B)$. Repeating the previous argument with A replaced by B yields some $S_3 \in \mathbb{P}$ with $[S_3] \subseteq_I [S_2] \cap B$.

Since $[S_3] \subseteq_I A \cap B$, we have $D_I([S_3]) \subseteq D_I(A \cap B)$. Since $[S_3] \subseteq_I C$ it follows that $D_I([S_3]) \cap [S_3] \subseteq_I C \cap D_I(A \cap B) = \emptyset$. This contradicts the assumption that $D_I([S_3]) \cap [S_3] \notin I$. \square

To obtain I -positivity we assume the following property (already mentioned in Section 1).

Definition 4.15. A collection of trees \mathbb{P} is called *strongly linked* if any $S, T \in \mathbb{P}$ with $\text{stem}(S) \subseteq \text{stem}(T)$ and $\text{stem}(T) \in S$ are compatible in \mathbb{P} .

This condition holds for all ccc tree forcings that we study in this paper except random forcing. For instance for Hechler forcing, eventually different forcing, Laver forcing \mathbb{L}_F with a filter and Mathias forcing \mathbb{R}_F with a shift invariant filter. Clearly the condition implies that \mathbb{P} is σ -linked and ccc. Thus $I_{\mathbb{P}} = I_{\mathbb{P}}^*$ by Section 4.3.

Remark 4.16. The null ideal I_{μ} is not a strongly linked tree ideal. For suppose that \mathbb{P} is a strongly linked tree forcing with $I_{\mathbb{P}} = I_{\mu}$. Fix a nowhere dense closed set C of positive measure and find $S \in \mathbb{P}$ with $[S] \subseteq_{\mu} C$ by Lemma 4.10. We write $A \subseteq_{\mu} B$ if $\mu(A \setminus B) = 0$. Find some $t \in S$ with $\text{stem}(S) \subseteq t$ and $\frac{\mu([S] \cap N_t)}{\mu(N_t)} < 1$; such a $t \in S$ exists since C is nowhere dense and $[S] \subseteq_{\mu} C$. Let further $A = N_t \setminus [S]$. Since $\frac{\mu(A \cap N_t)}{\mu(N_t)} > 0$, there is some $T \in \mathbb{P}$ with $[T] \subseteq_{\mu} A$ (again by Lemma 4.10). Then $\text{stem}(S) \subseteq t \subseteq \text{stem}(T)$, but S and T are incompatible in \mathbb{P} .

Lemma 4.17. Assume that \mathbb{P} is a strongly linked tree forcing—whence by Convention 2.2 $\mathbf{L}_{I_{\mathbb{P}}} = \{[T] \in \mathbb{P} \mid \text{stem}(T) = \emptyset\}$. Let $I = I_{\mathbb{P}}$. Then D_I is I -positive. In fact for any $T \in \mathbb{P}$, $D_I([T]) = [T]$.

Proof. It follows from the definition that $D_I([T]) \subseteq [T]$. To see the other inclusion, let $x \in [T]$ be given; we show that x is an I -density point of $[T]$. Let $m \in \omega$ be such that $\text{stem}(T) = x \upharpoonright m$. Suppose we are given $S \in \mathbb{P}$ such that $\text{stem}(S) = x \upharpoonright n$ for some $n \geq m$. Since the stems of S and T are compatible, $\text{stem}(S) \in T$, and \mathbb{P} is strongly linked, S and T are compatible. Therefore $[S] \cap [T]$ is an I -positive set as required. \square

Since $I_{\mathbb{P}} = I_{\mathbb{P}}^*$ when \mathbb{P} is ccc, Proposition 4.12 together with Lemmas 4.14 and 4.17 imply the following version of Lebesgue's density theorem, viz. Theorem 1.2:

Corollary 4.18. For any strongly linked tree forcing \mathbb{P} , the $I_{\mathbb{P}}$ -shift density property (cf. Definition 2.3) holds. In particular, the $I_{\mathbb{P}}$ -shift density property holds when \mathbb{P} is Cohen forcing \mathbb{C} , Hechler forcing \mathbb{H} , eventually different forcing \mathbb{E} , Laver forcing \mathbb{L}_F with a filter and Mathias forcing \mathbb{R}_F with a shift invariant filter (cf. Section 6).

For random forcing, the I_{μ} -shift density property follows from Lemma 3.1 and Lebesgue's density theorem. A self-contained proof is obtained from the fact that D_I is I -positive by Theorem 7.1 together with Proposition 4.12 and Lemma 4.14.

Note that if \mathbb{P} is topological and $I_{\mathbb{P}}$ has the density property, then one can describe $D_{I_{\mathbb{P}}}(A)$ as follows using $\tau_{\mathbb{P}}$. First note that $I_{\mathbb{P}} = I_{\mathbb{P}}^*$ by the proof of [FKK16, Lemma 3.8]. Moreover, for any Borel set A there is some $\tau_{\mathbb{P}}$ -open set U with $A \triangle U$ $\tau_{\mathbb{P}}$ -meager by the $\tau_{\mathbb{P}}$ -Baire property. Since $I_{\mathbb{P}}$ equals the set of $\tau_{\mathbb{P}}$ -meager sets, $D_{I_{\mathbb{P}}}(A)$ is almost equal to a $\tau_{\mathbb{P}}$ -open set. However, note that even for ccc collections it is not clear how to find such a set in a simply definable way.

We conclude this section with some observations about other variants of the Definition 2.3 of density points and the shift density property.

Remark 4.19. If \mathbb{P} is ccc and the density property holds for all Borel sets, then it already holds for all \mathbb{P} -measurable sets. This is the case because any \mathbb{P} -measurable set equals $A \triangle B$ for some Borel set A and some $B \in I_{\mathbb{P}}$ by the ccc and Lemma 4.8.

Remark 4.20.

- (1) If we let $\mathbf{L}_I = I^+$ be the I -positive sets for Hechler forcing \mathbb{H} and $I = I_{\mathbb{H}}$, then the density property fails. To see this, let T be the Hechler tree with empty stem given by the constant function with value 1. With this variant $[T]$ does not satisfy the density property, since no $x \in [T]$ is an I -shift density point of $[T]$, as witnessed by $N_{\langle 0 \rangle} \in \mathbf{L}_I$.
- (2) If n_0 depends on $B \in \mathbf{L}_I$ in Definition 2.3, then the density property fails for $I_{\mathbb{H}}$ as well. To see this, let x be the function with $x(n) = n + 1$ for all $n \in \omega$ and let $T_{\emptyset, x}$ be the tree with empty stem given by x . Let further $A = \bigcup_{t \in {}^\omega \omega} [T_{t \restriction \langle |t| \rangle, t \restriction \langle |t| \rangle \smallfrown 0^\omega}]$; this contains all $y \in {}^\omega \omega$ with $y(n) = n$ for some $n \in \omega$. It is sufficient to show $[T_{\emptyset, x}] \subseteq D(A) \setminus A$ by Lemma 4.10.

It is easy to see that $[T_{\emptyset, x}] \cap A = \emptyset$, since any $y \in [T_{\emptyset, x}]$ satisfies $y(n) \geq n + 1$ for all $n \in \omega$.

We now claim that $y \in D(A)$ for all $y \in {}^\omega \omega$. So take any $I_{\mathbb{H}}$ -positive Borel set B . It is sufficient to assume that $B = [T_{s, u}]$ for some $s \in {}^{<\omega} \omega$ and $u \in {}^\omega \omega$ by Lemma 4.10. Then for any $n \geq u(|s|)$, $\sigma_{y \restriction n}^{-1}([T_{s, u/s}]) \cap A = [(y \restriction n) \smallfrown (T_{s, u/s})] \cap A$ contains $[T_{t, t \restriction v}]$ for $t = y \restriction n \smallfrown \langle n \rangle$ and $v \in {}^\omega \omega$ with $v(i) = u(|s| + i + 1)$ for all $i \in \omega$. Hence it is $I_{\mathbb{H}}$ -positive and thus y is a density point of A .

4.6. Complexity of the density operator. In this section, we give an upper bound for the complexity of the operator D_I for relevant cases of tree ideals I of the form $I_{\mathbb{P}}$. \mathbf{K}_I is fixed as in the previous section.

Recall that a subset A of ${}^\omega \omega$ is called *universally Baire* if for any topological space Y and any continuous function $f: Y \rightarrow {}^\omega \omega$, $f^{-1}(A)$ has the property of Baire. A function $f: A \rightarrow B$ between subsets of ${}^\omega \omega$ is called *universally Baire measurable* if $f^{-1}(U)$ is universally Baire for every relatively open subset U of B . Note that all universally Baire sets have the Baire property, are Lebesgue measurable and Ramsey [FMW92, Theorem 2.2]. A definable forcing \mathbb{P} is called *absolutely ccc* if the ccc holds in all generic extensions. A Δ_2^1 predicate is *absolutely* Δ_2^1 if its Σ_2^1 and Π_2^1 definitions are equivalent in all generic extensions.

Lemma 4.21. *Suppose that $I = I_{\mathbb{P}}$, where \mathbb{P} , $\leq_{\mathbb{P}}$ and $\perp_{\mathbb{P}}$ are Σ_1^1 , \mathbf{K}_I is a Σ_1^1 subset of \mathbb{P} and \mathbb{P} is absolutely ccc. Then*

$$D_I: \text{BOREL}({}^\omega \Omega) \rightarrow \text{BOREL}({}^\omega \Omega)$$

is induced by a Δ_2^1 function from Borel codes to Borel codes. Moreover, this function is universally Baire measurable.

Proof. Recall that we write \mathcal{B} for the Π_1^1 set of Borel codes and B_x the set coded by a Borel code x . Let $\varphi(x)$ denote the formula $x \in \mathcal{B} \ \& \ \forall T \in \mathbf{K}_I \ B_x \cap [T] \notin I$. Let $\psi(x)$ denote the following statement: $x \in \mathcal{B}$ and there is some $m \leq \omega$ and a sequence $\vec{T} = \langle T_i \mid i < m \rangle$ from \mathbb{P} with $\forall i < m \ [T_i] \subseteq_I B_x$ and $\forall T \in L_I \ \exists i < m \ T \parallel T_i$.

Claim 4.22. $\forall x \ \varphi(x) \iff \psi(x)$.

Proof. If $\varphi(x)$ holds, inductively define an antichain $\vec{T} = \langle T_\xi \mid \xi < \theta \rangle$ from \mathbb{P} with $[T_\xi] \subseteq_{I_{\mathbb{P}}} B_x$ for all $\xi < \theta$. Suppose that $\vec{T}^{(\alpha)} = \langle T_\xi \mid \xi < \alpha \rangle$ is already defined. If there is $T \in L_I$ which is incompatible with each element of $\vec{T}^{(\alpha)}$, we may find $T_\alpha \in \mathbb{P}$ with $T_\alpha \subseteq T$ and $[T_\alpha] \subseteq B_x$ since $\varphi(x)$ holds. We can thus extend the antichain by adding T_α . By the ccc we must reach some $\theta < \omega_1$ such that each $T \in L_I$ is compatible to an element of $\vec{T} = \langle T_\xi \mid \xi < \theta \rangle$. Enumerate \vec{T} in order type $m \leq \omega$ to obtain a witness to $\psi(x)$.

If conversely $\psi(x)$ holds, then for any $T \in \mathbf{K}_I$ we may find some $i \in \omega$ with $T_i \parallel T$ and $[T_i] \subseteq_I B_x$, so one can infer $B_x \cap [T] \notin I$ from Lemma 4.10. Thus $\varphi(x)$ holds. \square

We now check that $\varphi(x)$ is a Π_2^1 and $\psi(x)$ a Σ_2^1 formula. First note that the statement $B_x \in \mathbf{N}_{\mathbb{P}}$ is Σ_2^1 , since this holds if and only if there is a (countable) maximal antichain $\vec{S} = \langle S_i \mid i < m \rangle$ in \mathbb{P} with $B_x \cap \bigcup_{i < m} [S_i] = \emptyset$. Since $I = I_{\mathbb{P}}$ is the σ -ideal generated by $\mathbf{N}_{\mathbb{P}}$, the statements $B_x \in I$ and $[T] \subseteq_I B_x$ are Σ_2^1 as well. Thus $\varphi(x)$ and $\psi(x)$ are indeed of said complexity.

Since \mathbb{P} is absolutely ccc, the argument above shows that $\forall x \Phi(x) \iff \Psi(x)$ is absolute to generic extensions. The above easily shows that $S = \{(x, s) \in \mathcal{B} \times {}^{<\omega}\Omega \mid \forall T \in \mathbf{K}_I \sigma_s([T]) \cap B_x \neq \emptyset\}$ is absolutely Δ_2^1 , where \mathcal{B} denotes the set of Borel codes. Thus the function \hat{D} which sends each Borel code x to a Borel code $\hat{D}(x)$ for $D_I(B_x)$ has a Σ_2^1 graph. Finally for each $s \in {}^{<\omega}\Omega$, $\{x \mid (x, s) \in S\}$ is absolutely Δ_2^1 and hence universally Baire by [FMW92, Theorem 2.1]. It follows easily that \hat{D} is universally Baire measurable. \square

We want to point out that the previous lemma remains true with virtually the same proof if we replace *absolutely ccc* and *absolutely Δ_2^1* by *provably ccc* and *provably Δ_2^1* .

We now show that for all strongly linked tree forcings listed in Section 6, the density operator $D_{I_{\mathbb{P}}}$ is induced by a universally Baire measurable function. Recall that a forcing is called *Suslin* if \mathbb{P} , $\leq_{\mathbb{P}}$ and $\perp_{\mathbb{P}}$ are Σ_1^1 .

Proposition 4.23. *For any strongly linked Suslin tree forcing \mathbb{P} , $D_{I_{\mathbb{P}}}$ is induced by a universally Baire measurable function.*

Proof. If \mathbb{P} is strongly linked, then $S \parallel T$ if and only if $\text{stem}(S) \in T$, $\text{stem}(T) \in S$, and $\text{stem}(S)$ and $\text{stem}(T)$ are compatible, so $\perp_{\mathbb{P}}$ is arithmetical and hence Σ_1^1 . Moreover, the fact that a Suslin tree forcing is strongly linked is Π_2^1 and hence absolute. Thus $D_{I_{\mathbb{P}}}$ is induced by a Σ_2^1 , universally Baire measurable function on the Borel codes by Lemma 4.21. \square

5. IDEALS WITHOUT DENSITY

In this section, we study various counterexamples to density properties.

5.1. Counterexamples. We first give counterexamples to the density property in Definition 2.3 for several non-ccc tree forcings.

Proposition 5.1. *Let \mathbb{R} , \mathbb{V} , \mathbb{S} denote Mathias, Silver and Sacks forcing. Then $I_{\mathbb{R}}$, $I_{\mathbb{V}}$ and $I_{\mathbb{S}}$ do not have the shift density property.*

Proof. To see that $I_{\mathbb{R}}$ does not have the $I_{\mathbb{R}}$ -shift density property, let $A = \{f \in {}^\omega 2 \mid f(2n+1) = 1 \text{ for all } n \in \omega\}$. Note that $A = [S]$ for some $S \in \mathbb{R}$ and hence $A \notin I_{\mathbb{R}}$. We aim to show that no $x \in A$ is an $I_{\mathbb{R}}$ -density point of A , i.e. $A \cap D_{\mathbb{R}}(A) = \emptyset$. Then in particular $A \triangle D_{\mathbb{R}}(A) = A \cup D_{\mathbb{R}}(A) \notin I_{\mathbb{R}}$.

Let $x \in A$ be arbitrary and let $T \in \mathbb{R}$ be a perfect tree such that $\text{split}(T) = 2\mathbb{N}$ and $t \smallfrown \langle i \rangle \smallfrown \langle j \rangle \in T$ iff $j = 0$ for all $t \in \text{split}(T)$ and $i, j \in \{0, 1\}$. In particular $\text{stem}(T) = \emptyset$. Let $n_0 \in \omega$ be arbitrary and let $n \geq n_0$ be even. Then $f_{x \upharpoonright n}[T] \cap A = \emptyset$ and thus x is not an $I_{\mathbb{R}}$ -density point of A .

As $\mathbb{R} \subseteq \mathbb{V} \subseteq \mathbb{S}$, the claim also holds for $I_{\mathbb{V}}$ and $I_{\mathbb{S}}$. \square

The following is a similar counterexample for Laver and Miller forcing.

Proposition 5.2. *Let \mathbb{L} , \mathbb{M} denote Laver and Miller forcing. Then $I_{\mathbb{L}}$ and $I_{\mathbb{M}}$ do not have the shift density property.*

Proof. Let $A = \{f \in {}^\omega \omega \mid f(n) \text{ is even for all } n \in \omega\}$. Then $A = [S]$ for some $S \in \mathbb{L}$ so in particular $A \notin I_{\mathbb{L}}$. We aim to show that no $x \in A$ is an $I_{\mathbb{L}}$ -density point of A , i.e. $A \cap D_{\mathbb{L}}(A) = \emptyset$. As above this implies $A \triangle D_{\mathbb{L}}(A) \notin I_{\mathbb{L}}$.

Let $x \in A$ be arbitrary and let $T \in \mathbb{L}$ be a perfect tree such that $\text{stem}(T) = \emptyset$ and $[T] = \{g \in {}^\omega \omega \mid g(n) \text{ is odd for all } n \in \omega\}$. Let $n_0 \in \omega$ be arbitrary and let $n \geq n_0$. Then $f_{x \upharpoonright n}[T] \cap A = \emptyset$ and thus x is not an $I_{\mathbb{L}}$ -density point of A .

Since $\mathbb{L} \subseteq \mathbb{M}$, the claim for $I_{\mathbb{M}}$ follows. \square

5.2. Selectors for the ideal of countable sets. We now study the ideal I of countable sets. In contrast to the previous results, we will show that there is no Baire measurable selector with Borel values for the equivalence relation of having countable symmetric difference on the set of Borel subsets. This implies that the density property fails for I for any reasonable notion of density point.

To state the result formally, we need the following notions. A *selector* for an equivalence relation E is a function that chooses an element from each equivalence class. We generalize this notion by replacing equality with a subequivalence relation E' of E .

Definition 5.3. Suppose that $E' \subseteq E$ are equivalence relations on a set B and $A \subseteq B$. A *selector for E/E' on A* is an (E, E') -homomorphism $A \rightarrow B$ that uniformizes E .

Equivalently, the induced map on B/E' is a selector for the equivalence relation on B/E' induced by E .

In the following, E' will be equality of decoded sets, E the equivalence relation of having countable symmetric difference, A will be the set of F_σ -codes, denoted also by \mathcal{B}_{F_σ} , and $B = \mathcal{B}$, the set of Borel codes. More precisely, an F_σ -code is a sequence $\vec{T} = \langle T_n \mid n \in \omega \rangle$, where T_n is a subtree of ${}^{<\omega}2$ for each $n \in \omega$. Note that \mathcal{B}_{F_σ} is therefore a Polish space. We can assume that $\mathcal{B}_{F_\sigma} \subseteq \mathcal{B}$.

Equality of Borel sets and equality modulo I induces the following equivalence relations on \mathcal{B} . As before, let B_x denote the Borel set coded by $x \in \mathcal{B}$. Let $E_ =$ denote the equivalence relation on \mathcal{B} of equality of decoded sets, i.e. $(x, y) \in E_ = \iff B_x = B_y$. Let further $(x, y) \in E_I \iff B_x \triangle B_y \in I$ for $x, y \in \mathcal{B}$.

Definition 5.4. A *selector for I with Borel values* is a selector $\mathcal{B}_{F_\sigma} \rightarrow \mathcal{B}$ for $E_I/E_ =$.

The restriction to F_σ -codes is purely for a technical reason: the proof of Theorem 5.5 will show that there is no reasonably definable selector for I on \mathcal{B}_{F_σ} . It follows that there is no such selector on the set of all Borel codes.

The motivation for this definition is as follows. Consider any notion of density points for I with Borel values, i.e. such that for any Borel set A , the set of density points of A is Borel. If the density property holds for this notion, then the density operator induces a selector for I on the set of Borel codes.

Theorem 5.5. *There is no Baire measurable selector for I , the ideal of countable sets, with Borel values.*

It follows that there is no selector $\text{BOREL}({}^\omega\Omega) \rightarrow \text{BOREL}({}^\omega\Omega)$ for $=_I$ that is induced by a universally Baire measurable function (on the codes).

Note that this result is analogous to the fact that E_0 does not have a Baire measurable selector. However, the rather short proof of the latter (see e.g. [Hjo10, Example 1.6.2]) is of no use here.

The current proof simplifies that of a previous version, and resembles Kanovei's version [KL20] in some aspects, but was arrived at independently.

Before we begin the long proof of Theorem 5.5 let us sketch the main ideas. Suppose that $F: \mathcal{B}_{F_\sigma} \rightarrow \mathcal{B}$ is a selector for $E_I/E_ =$. Thus, given a code C for an F_σ set, $B_{F(C)}$ is a representative for the equivalence class of B_C modulo I .

We first construct a sequence of names for perfect trees $\langle \dot{T}_n \mid n \in \omega \rangle$ added by Cohen forcing, \mathbb{C} . Thus it is forced that $\bigcup_{n \in \omega} [\dot{T}_n]$ is an F_σ set; let \dot{C} be a name for its code (i.e., for the sequence of trees). We will fix a large enough countable elementary submodel M of H_{ω_1} and carefully construct a name $\sigma \in M$ for an element of the representative $B_{F(\dot{C})}$ of $\bigcup_{n \in \omega} [\dot{T}_n]$.

Now let $g \times h$ be any \mathbb{C}^2 -generic over M . We will find a different $g' \in M[g][h]$ which is also Cohen-generic over M such that $\bigcup h = \sigma^{g'}$ and

$$\bigcup_{n \in \omega} [\dot{T}_n^g] = \left(\bigcup_{n \in \omega} [\dot{T}_n^{g'}] \right) \setminus \{\sigma^{g'}\}.$$

Thus letting $C = \langle \dot{T}_n^g \mid n \in \omega \rangle$, a code for $\bigcup_{n \in \omega} [\dot{T}_n^g]$, and $C' = \langle \dot{T}_n^{g'} \mid n \in \omega \rangle$, a code for $\bigcup_{n \in \omega} [\dot{T}_n^{g'}]$, it follows that $B_{F(C)} = B_{F(C')}$, i.e., the two sets have the same representative. A careful choice of the name σ and of the model M is crucial for the construction of g' . Since $\bigcup h = \sigma^{g'} \in B_{F(C')} = B_{F(C)}$, we will be able to conclude that any Cohen real over $M[g]$ is an element of $B_{F(C)} \setminus \bigcup_{n \in \omega} [\dot{T}_n^g]$. Since there are uncountably many Cohen reals over $M[g]$ this contradicts the assumption that F is a selector.

We now introduce some notation to aid us in the proof of Theorem 5.5. To obtain our sequence $\langle \dot{T}_n \mid n < \omega \rangle$ we will use a particular presentation of Cohen forcing.

Notation 5.6. Let \mathbb{T} be the forcing whose conditions are finite subtrees t of ${}^{<\omega}2$, ordered by end extension. By this we mean that one can only extend a tree at its maximal nodes. For any \mathbb{T} -generic filter G we refer to the tree $\bigcup G$ as the tree added by \mathbb{T} .

Let $\mathbb{T}^\omega = \prod_{n \in \omega} \mathbb{T}_n$ denote the finite support product of ω copies of \mathbb{T} and \dot{T}_n a \mathbb{T}_n -name for the tree added by \mathbb{T}_n . We can identify each \dot{T}_n canonically with a \mathbb{T}^ω -name.

Since \mathbb{T}^ω is countable and atomless, indeed it is a presentation of Cohen forcing.

We are now ready to prove Theorem 5.5.

Proof of Theorem 5.5. Suppose that there is a selector F on \mathcal{B}_{F_σ} as in the statement of the theorem. Since F is Baire measurable, there is a comeager G_δ subset A of the Polish space \mathcal{B}_{F_σ} such that $F \upharpoonright A$ is continuous. Let x_A be a real in which both a Borel code for A and a code for $F \upharpoonright A$ are computable. Moreover, let $M < H_{\omega_1}$ be countable with $x_A \in M$.

Work in M . Note that $\mathbb{T}^\omega \in M$; the same holds for all other objects defined in 5.6. Let $\dot{C} \in M$ be a \mathbb{T}^ω -name for the canonical F_σ -code for $\bigcup_{n \in \omega} [\dot{T}_n]$ and $\dot{F} \in M$ a \mathbb{T}^ω -name for $F \upharpoonright A$.

Working in V again, $\dot{F}^g(x) = F(x)$ for all $x \in A \cap M[g]$ and all \mathbb{T}^ω -generic filters $g \in V$ over M .

Claim 5.7. For all $n \in \omega$, $1 \Vdash_{\mathbb{T}^\omega} [\dot{T}_n] \setminus B_{\dot{F}(\dot{C})}$ is countable.

Proof. Assume towards a contradiction that $p \Vdash_{\mathbb{T}^\omega} [\dot{T}_n] \setminus B_{\dot{F}(\dot{C})}$ is uncountable for some $p \in \mathbb{T}^\omega$ and some $n \in \omega$. Since A is comeager, any \mathbb{T}^ω -generic filter g over M with $p \in g$ will satisfy $\dot{C}^g \in A$. Let $C = \dot{C}^g$. Then $\dot{F}^g(\dot{C}^g) = F(C)$. Since $p \in g$, $M[g] \models [\dot{T}_n^g] \setminus B_{F(C)}$ is uncountable.

Work in $M[g]$. By the perfect set property for Borel sets, there is a perfect tree S whose branches are all elements of $[\dot{T}_n^g] \setminus B_{F(C)}$.

By Π_1^1 -absoluteness between $M[g]$ and V , all branches of S in V are elements of $[\dot{T}_n^g] \setminus B_{F(C)}$. But this contradicts the assumption that F is a selector. \square

We now resume working in M . For the construction of σ we need some more notation:

Notation 5.8.

- (1) If S is a subtree of ${}^{<\omega}2$, let $\text{split}(S)$ denote the set of splitting nodes of S , i.e. those with at least two direct successors and let $\text{term}(S)$ denote the set of terminal (i.e. maximal) nodes in S .
- (2) If S is an arbitrary subtree of ${}^{<\omega}2$, let π_S denote the unique order isomorphism from a subtree of ${}^{<\omega}2$ onto $\text{split}(S) \cup \text{term}(S)$ such that every node in $\text{dom}(\pi_S)$ is either splitting or maximal in $\text{dom}(\pi_S)$. (If S is perfect, then π_S is the unique order isomorphism from ${}^{<\omega}2$ to $\text{split}(S)$.)
- (3) For each $x \in {}^\omega 2$, let σ_x be a \mathbb{T}^ω -name for $\bigcup_{n \in \omega} \pi_{\dot{T}_0}(x \upharpoonright n)$.

Using the previous claim, elementarity, the fact that \mathbb{T}^ω preserves cardinality and $({}^\omega 2)^M$ is uncountable in M , there is $x \in ({}^\omega 2)^M$ and $p \in \mathbb{T}^\omega$ with $p \Vdash_{\mathbb{T}^\omega} \sigma_x \in [\dot{T}_0] \cap B_{\dot{F}(\dot{C})}$.

Notation 5.9. Until the end of the proof of Theorem 5.5 and Claim 5.10, let us fix $x \in ({}^\omega 2)^M$ and $p_0 \in \mathbb{T}^\omega$ with $p_0 \Vdash_{\mathbb{T}^\omega} \sigma_x \in [\dot{T}_0] \cap B_{\dot{F}(\dot{C})}$. Moreover, we assume $p_0 = \emptyset$ for notational convenience. We shall also write σ for σ_x and $\text{stem}(T)$ for $\text{stem}(T)$.

Our next goal is to demonstrate that for any \mathbb{T}^ω -generic g over M , $B_{F(\dot{C}^g)} \setminus B_{\dot{C}^g}$ must contain every real which is Cohen over $M[g]$ (and hence must be uncountable, leading to a contradiction). The next claim will be crucial.

Claim 5.10. In every $\mathbb{T}^\omega \times \mathbb{C}$ -generic extension $M[g \times h]$ we can find g' which is \mathbb{T}^ω -generic over M such that

$$\sigma^{g'} = \bigcup h, \quad (3)$$

$$\left(\bigcup_{n \in \omega} [\dot{T}_n^{g'}] \right) \setminus \{\sigma^{g'}\} = \bigcup_{n \in \omega} [\dot{T}_n^g]. \quad (4)$$

In fact, there exist in M a dense subset D of $\mathbb{T}^\omega \times \mathbb{C}$ and a projection

$$\pi: D \rightarrow \mathbb{T}^\omega$$

so that the above holds with $g' = \pi[(g \times h) \cap D]$. (See [Cum10, Definition 5.2] for the definition of projections.)

Assuming Claim 5.10 for the moment, we can easily finish the proof of Theorem 5.5 as follows. Fix any \mathbb{T}^ω -generic filter g over M in V . Since A is comeager and is coded in M , we have $C = \dot{C}^g \in A$.

Claim 5.11. $B_{F(C)} \setminus B_C$ contains every Cohen real over $M[g]$ in V .

Proof. Let h be any Cohen generic filter over $M[g]$ in V and let $c = \bigcup h$. One easily obtains $c \notin B_C$ by a density argument.

It remains to show that $c \in B_{F(C)}$. By the previous claim we can find $g' \in M[g \times h]$ which is \mathbb{T}^ω -generic over M and so that $c = \sigma^{g'}$ and $B_{\dot{C}^{g'}} = B_C \cup \{\sigma^{g'}\}$. Moreover $\sigma^{g'} \in B_{F(\dot{C}^{g'})}$ by the choice of σ and Borel absoluteness between $M[g']$ and V . Since F is a selector, $B_{F(\dot{C}^{g'})} = B_{F(C)}$. We thus have $c = \sigma^{g'} \in B_{F(\dot{C}^{g'})} = B_{F(C)}$. \square

By the previous claim, $B_{F(C)} \setminus B_C$ is uncountable. But this contradicts our assumption that F is a selector.

To finish the proof of Theorem 5.5, it remains to prove Claim 5.10. We need some more notation:

Notation 5.12. For $s, t \in {}^{<\omega}2$ write $s \wedge t$ for the longest common initial segment of s and t .

Proof of Claim 5.10. Before we define π , we define a map $\hat{\pi}$ on a larger domain. The projection π will be the restriction of $\hat{\pi}$ to a dense subset D of $\mathbb{T}^\omega \times \mathbb{C}$. This approach is convenient since $\hat{\pi}$ captures both the action of π on conditions in $\mathbb{T}^\omega \times \mathbb{C}$, as well the “continuous extension” of this action to generic objects for $\mathbb{T}^\omega \times \mathbb{C}$.

The domain of $\hat{\pi}$ is the set of pairs $(\langle t_n \mid n < l \rangle, p)$ such that $l \in \omega + 1$, $p \in {}^{<\omega}2$, and each t_n is a subtree of ${}^{<\omega}2$. Suppose we are given $q = (\langle t_n \mid n < l \rangle, p) \in \text{dom}(\hat{\pi})$. We shall let

$$\hat{\pi}(q) = \langle t'_j \mid j < l' \rangle,$$

where $\langle t'_j \mid j < l' \rangle$ is a sequence of subtrees of ${}^{<\omega}2$ with $l' \in \omega + 1$ which is constructed as follows.

We first construct a sequence $\langle n(i) \mid i < m \rangle$ of natural numbers $n(i) < l$, where $m \in \omega + 1$. Let $k \in \omega$ and suppose we have already defined $\langle n(i) \mid i < k \rangle$ (if $k = 0$, this is the empty sequence). Let $n(k)$ be least number, if it exists, that satisfies the following conditions:

Conditions 5.13.

- (a) If $k > 0$, then
 - (i) $n(k) > n(k-1)$,
 - (ii) $p \wedge \text{stem}(t_{n(k-1)}) \subsetneq p \wedge \text{stem}(t_{n(k)})$ and
 - (iii) $\text{stem}(t_{n(k)}) \perp p$.
- (b) Letting $t = \{p \restriction j \mid j \in \omega\} \cup \bigcup_{i \leq k} t_{n(i)}$, we have

- (i) $x \upharpoonright (k+1) \in \text{dom}(\pi_t)$ and²⁰
- (ii) $\pi_t(x \upharpoonright (k+1)) \subseteq p$.

If such $n(k)$ does not exist, let $m = k$ and declare the construction of $\langle n(i) \mid i < m \rangle$ complete. If $n(k)$ is defined for each $k < \omega$ we let $m = \omega$. This completes the construction of $\langle n(i) \mid i < m \rangle$.

We now define $\langle t'_j \mid j < l' \rangle$ and thus $\hat{\pi}$. First, let

$$t'_0 = \{p \upharpoonright j \mid j \in \omega\} \cup \bigcup_{i < m} t_{n(i)}. \quad (5)$$

Note that $m < \omega$ and $\pi_{t'_0}(x \upharpoonright m) = p$ if $p \in \mathbb{C}$ and in general, $\bigcup_{k < m} \pi_{t'_0}(x \upharpoonright (k+1)) = p$. Further, let $\langle t'_j \mid 0 < j < l' \rangle$ be the increasing re-enumeration of what remains of the sequence $\langle t_n \mid n < l \rangle$ after removing the subsequence $\langle t_{n(i)} \mid i < m \rangle$. In other words, letting $\langle \bar{n}(i) \mid i < \bar{l} \rangle$ be the increasing enumeration of $l \setminus \{n(i) \mid i < m\}$, we define $l' = 1 + \bar{l}$ and $t'_{j+1} = t_{\bar{n}(j)}$ for $j < \bar{l}$. (Note that $l' = l - m + 1$ if l is finite.) If $(\langle t_n \mid n < l \rangle, p) \in \mathbb{T}^\omega \times \mathbb{C}$, then clearly $\langle t'_j \mid j < l' \rangle \in \mathbb{T}^\omega$.

To define D , we introduce following notation. Suppose that in the context of the above construction of $\langle n(i) \mid i < m \rangle$, we have $(\langle t_n \mid n < l \rangle, p) \in \mathbb{T}^\omega \times \mathbb{C}$ (whence $m < \omega$) and furthermore, $m > 0$. Then we denote the last initial segment $p \wedge \text{stem}(t_n)$ of p picked up in this construction as

$$s(\langle t_n \mid n < l \rangle, p) := p \wedge \text{stem}(t_{n(m-1)}). \quad (6)$$

When this notation is used in the definition of D below, it is implied that $0 < m < \omega$, so that $t_{n(m-1)}$ is defined.

Note that $\pi_t(x \upharpoonright (k+1)) \perp \text{stem}(t_{n(k)})$ and $\pi_t(x \upharpoonright k) \subseteq \text{stem}(t_{n(k)})$ for each $k < m$ and $p \wedge \text{stem}(t_{n(k)}) \subsetneq p \wedge \text{stem}(t_{n(k)}) \subsetneq p$ for each $k' < k < m$. Let

$$D = \{(\langle t_n \mid n < l \rangle, p) \in \mathbb{T}^\omega \times \mathbb{C} \mid |s(\langle t_n \mid n < l \rangle, p)| = |p| - 1 \text{ and} \\ \forall n < l \text{ split}(t_n) \neq \emptyset \wedge |p \wedge \text{stem}(t_n)| < |p|\}, \quad (7)$$

where $s(\langle t_n \mid n < l \rangle, p)$ is as defined in (6) above. Clearly D is dense in $\mathbb{T}^\omega \times \mathbb{C}$. The reason for restricting the domain of $\hat{\pi}$ in this way is that for $q \in D$, it is guaranteed that the sequence $\langle n(i) \mid i < m \rangle$ grows in the intended manner when extending the \mathbb{T}^ω -component of q (see below).

Subclaim 5.14. The map $\pi = \hat{\pi} \upharpoonright D$ is a projection to \mathbb{T}^ω .

Proof. To see that π is order-preserving, suppose we are given $q = (\langle t_j \mid j < l \rangle, p)$ and $\tilde{q} = (\langle \tilde{t}_j \mid j < \tilde{l} \rangle, \tilde{p})$ from D with $\tilde{q} \leq q$. Write $\pi(q) = \langle t'_j \mid j < l' \rangle$ and $\pi(\tilde{q}) = \langle \tilde{t}'_j \mid j < \tilde{l}' \rangle$. We must show that $l' \leq \tilde{l}'$ and \tilde{t}'_j end-extends t'_j for each $j < l'$.

Write $\langle n(i) \mid i < m \rangle$ for the sequence constructed from q as above (in the definition of $\hat{\pi}$) and write $\langle \tilde{n}(i) \mid i < \tilde{m} \rangle$ for the analogous sequence constructed from \tilde{q} . By the requirement in (7) that all trees have at least one splitting node, and since trees can only be extended at their maximal nodes, we have $\text{stem}(t_j) = \text{stem}(\tilde{t}_j)$ for all $j < l$. By the requirement in (7) that $p \wedge \text{stem}(t_j)$ is strictly shorter than p for all $j < l$, $\langle n(i) \mid i < m \rangle$ is an initial segment of $\langle \tilde{n}(i) \mid i < \tilde{m} \rangle$. Hence the increasing enumeration of $l \setminus \{n(i) \mid i < m\}$ is an initial segment of the increasing enumeration of $\tilde{l} \setminus \{\tilde{n}(i) \mid i < \tilde{m}\}$.

It follows that $l' \leq \tilde{l}'$ and \tilde{t}'_i end-extends t'_i for $0 < i < l$. Since $q \in D$ and thus $|p \wedge \text{stem}(t_{n(m-1)})| = |p| - 1$ by (7), it follows that $p \subseteq \text{stem}(t_{\bar{n}(j)})$ whenever $m < j < \bar{m}$. Thus \tilde{t}'_0 end-extends t'_0 by the definition in (5).

To see that π is a projection, let $q = (\langle t_j \mid j < l \rangle, p) \in D$ be given; write $q' = \langle t'_n \mid n < l' \rangle$ for $\pi(q)$. Further suppose we are given $q'' = \langle t''_n \mid n < l'' \rangle$ in \mathbb{T}^ω with $q'' \leq q' = \pi(q)$. We aim to find $\tilde{q} = (\langle \tilde{t}_j \mid j < \tilde{l} \rangle, \tilde{p})$ in D with $\tilde{q} \leq q$ and $\pi(\tilde{q}) \leq q''$.

²⁰See Notation 5.8(2) for the definition of the isomorphism π_t .

Let \tilde{m} be maximal so that $\pi_{t_0''}(x \upharpoonright \tilde{m})$ is defined and let

$$\tilde{p} = \pi_{t_0''}(x \upharpoonright \tilde{m}).$$

Write $\langle s_i \mid i < \tilde{m} \rangle$ for the enumeration in order of increasing length of the set of $s \in t_0''$ such that $s \perp \tilde{p}$ but each proper initial segment of s is an initial segment of \tilde{p} .

We now take some precautions to ensure that \tilde{q} as defined below will be an element of D and that $\pi(\tilde{q}) = q''$.

Without loss of generality, by replacing each tree t_i'' for $i < l''$ by an end-extension if necessary, we can assume that each tree $(t_0'')_{s_i}$ for $i < \tilde{m}$ as well as each tree t_j'' for $0 < j < l''$ has at least one splitting node; this is automatically true only for $i < m$ and $j < l'$. (Recall here that $(t_0'')_{s_i} = \{s \in t_0'' \mid s \subseteq s_i \vee s_i \subseteq s\}$ as in Section 2, p. 2.)

Likewise, replacing t_0'' by an end-extension if necessary, we can assume that \tilde{p} has greater length than $\tilde{p} \wedge \text{stem}(t_j'')$ for each $j < l''$.

Finally, again replacing t_0'' by an end-extension and increasing \tilde{m} if necessary, we can assume that $|s_{\tilde{m}-1}| = |\tilde{p}|$, or equivalently,

$$|\tilde{p} \wedge s_{\tilde{m}-1}| = |\tilde{p}| - 1. \quad (8)$$

Making these assumptions will suffice to see that \tilde{q} as defined below is in D .

Note that p is an initial segment of \tilde{p} , since $q'' \leq q' = \pi(q)$ (whence t_0'' end-extends t_0') and by the definition of $\hat{\pi}$. We further have $p \wedge \text{stem}(t_{n(i)}) \subseteq s_i \subseteq \text{stem}(t_{n(i)})$ for each $i < m$ by the definition of $\langle n(i) \mid i < m \rangle$ from q as above. Moreover $p \subseteq s_j$ for any j with $m \leq j < \tilde{m}$, since $|s_{m-1}| = |p|$ or equivalently, $|p \wedge \text{stem}(t_{n(m-1)})| = |p| - 1$. The latter holds since $q \in D$.

We now define \tilde{q} . With $\langle \tilde{n}(i) \mid i < l - m \rangle$ defined from q as above, let for $j < l$

$$\tilde{t}_j = \begin{cases} (t_0'')_{s_i} & \text{if } j = n(i) \text{ for some } i < m, \\ t_{i+1}'' & \text{if } j = \tilde{n}(i) \text{ for some } i < l - m. \end{cases}$$

Let $\tilde{l} = l'' + \tilde{m} - 1$ and define

$$\tilde{q} = (\langle \tilde{t}_0, \dots, \tilde{t}_{l-1}, \overbrace{(t_0'')_{s_m}, \dots, (t_0'')_{s_{\tilde{m}-1}}}^{\text{if } \tilde{m} > m}, \overbrace{t_{l'}'', \dots, t_{l''-1}''}^{\text{if } l'' > l'}, \tilde{p} \rangle.$$

$\begin{array}{ccccccc} & & & \uparrow & & \uparrow & \uparrow \\ \text{\scriptsize } l\text{th position} & \longrightarrow & & \text{\scriptsize } (l+\tilde{m}-m)\text{th position} & \longrightarrow & & \text{\scriptsize } (\tilde{l}-1)\text{th position} \end{array}$

The precautions taken above ensure that $\tilde{q} \in D$. To see that $\pi(\tilde{q}) \leq q''$, observe that by the definition of $\hat{\pi}$ and (8), the sequence of $\tilde{n}(i)$ associated with \tilde{q} as above has length \tilde{m} ; we denote it by $\langle \tilde{n}(i) \mid i < \tilde{m} \rangle$. Moreover

$$\tilde{n}(i) = \begin{cases} n(i) & \text{if } i < m, \\ l - m + i & \text{if } m \leq i < \tilde{m}. \end{cases}$$

It is then a matter of straightforward computation to check that $\pi(\tilde{q}) = q''$. \square

It remains to verify (3) and (4). Suppose $g \times h$ is $\mathbb{T}^\omega \times \mathbb{C}$ -generic. Let $T_n = \dot{T}_n^g$ and let $c = \bigcup h$. By the definition of $\hat{\pi}$, letting $g' = \hat{\pi}(g \times h)$ and writing $\langle T'_n \mid n < \omega \rangle$ for the sequence of trees given as

$$\langle T'_n \mid n < \omega \rangle = \hat{\pi}(\langle T_n \mid n < \omega \rangle, c),$$

we have $T'_n = \dot{T}_n^{g'}$ for each $n < \omega$. Also by the definition of $\hat{\pi}$, in particular by (5),

$$c = \bigcup_{k \in \omega} \pi_{T'_0}(x \upharpoonright k) = \sigma_x^{g'} = \sigma^{g'}$$

proving (3). Moreover

$$T'_0 = \bigcup_{i < \omega} T_{n(i)}$$

with $n(i)$ as above (in the definition of $\hat{\pi}$) and therefore by construction

$$[T'_0] \setminus \{c\} = \bigcup_{i < \omega} [T_{n(i)}],$$

proving (4). \square

This completes the proof of the theorem, as we argued after Claim 5.10. \square

The next results use the following principle. Let *internal projective Cohen absoluteness* ($\text{IA}_{\text{proj}}^{\text{C}}$) denote the statement that $H_{\omega_1}^{M[g]} < H_{\omega_1}$ holds for all sufficiently large regular cardinals θ , countable elementary submodels $M < H_\theta$ and Cohen generic filters g over M in V .

We will only need the first part of the next lemma; the second part is an easy observation. Recall that PD denotes the axiom of projective determinacy.²¹

Lemma 5.15.

- (1) PD implies $\text{IA}_{\text{proj}}^{\text{C}}$.
- (2) If $\text{IA}_{\text{proj}}^{\text{C}}$ holds, then all projective set have the property of Baire.

Proof. (1) Take a Σ_{2n}^1 -universal Σ_{2n}^1 subset A_n of $(\omega 2)^2$ for each $n \geq 1$. If the Σ_{2n}^1 -formula $\varphi_n(x, y)$ defining A_n is chosen in a reasonable way, then for any Σ_{2n}^1 -formula $\psi(x)$ with (hidden) real parameters, there is some $y_\psi \in \omega 2$ with $\forall x (\psi(x) \iff \varphi_n(x, y_\psi))$. Moreover, this holds in all transitive models of ZFC^- and the map $\psi \mapsto y_\psi$ is absolute between such models. We assume that in any transitive model of ZFC^- , A_n denotes the set defined by $\varphi_n(x, y)$.

Let $M < H_\theta$ be a countable elementary submodel for some large enough regular cardinal θ . By [Mos09, Corollary 6C.4] and PD, there is a Σ_{2n}^1 -scale $\sigma_n = \langle \leq_n^m \mid m \in \omega \rangle$ on A_n for each $n \geq 1$. Let $r_n^m(x)$ denote the rank of $x \in A_n$ with respect to \leq_n^m . Moreover, recall that the *tree of the scale* σ_n ²² is defined as

$$T_{\sigma_n} = \{(x \restriction m, (r_n^0(x), \dots, r_n^{m-1}(x))) \mid x \in A_n, m \in \omega\}$$

and $p[T_{\sigma_n}] = A_n$. Since $\text{ZFC}^- + \text{PD}$ is sufficient to prove the existence of such scales, let T_{σ_n} denote the tree defined via $\varphi_n(x, y)$ in any transitive model of this theory containing $\mathcal{P}(\omega)$ as an element. Let $g \in V$ be Cohen generic over M .

Claim 5.16. $M[g] \models A_n = p[T_{\sigma_n}]$.

Proof. $p[T_{\sigma_n}]$ has a projective definition via the above definitions of T_{σ_n} and A_n . Thus the claim holds by 1-step projective Cohen absoluteness in M [Woo82, Lemma 2]. \square

Claim 5.17. $T_{\sigma_n}^M = T_{\sigma_n}^{M[g]}$.

Proof. By PD any projective pre-wellorder E is *thin*, i.e. there is no perfect set of reals that are pairwise inequivalent with respect to E . By 1-step projective Cohen absoluteness [Woo82, Lemma 2], [Sch14, Lemma 3.18] and PD, Cohen forcing does not add new equivalence classes to E . Equality now follows from [Sch14, Theorem 5.15]. The inclusion $T_{\sigma_n}^M \subseteq T_{\sigma_n}^{M[g]}$ holds since the rank function r_n is upwards absolute from M to $M[g]$ by the previous statement for $E = \leq_n$. The converse inclusion is proved in [Sch14, Claim 5.16] from PD. \square

We now show that $\text{IA}_{\text{proj}}^{\text{C}}$ holds. Assume that $\psi(x)$ is a Σ_{2n}^1 -formula and $x \in M[g]$. Note that the equivalences $\psi(x) \iff \varphi_n(x, y_\psi) \iff (x, y_\psi) \in A_n \iff (x, y_\psi) \in p[T_{\sigma_n}]$ hold in V and in $M[g]$ by the first claim. It remains to show that $M[g] \models (x, y_\psi) \in p[T_{\sigma_n}]$ if and only if $(x, y_\psi) \in p[T_{\sigma_n}]$ holds in V .

We claim that $T_{\sigma_n}^{M[g]} = T_{\sigma_n} \cap M[g]$. To see this, note that $T_{\sigma_n}^M = T_{\sigma_n} \cap M$ since $M < H_\theta$. Using the second claim and the fact that $\text{Ord}^M = \text{Ord}^{M[g]}$, we obtain $T_{\sigma_n}^{M[g]} = T_{\sigma_n}^M =$

²¹See [Jec03, Chapter 33].

²²The tree of a scale is defined in the discussion before [Mos09, Theorem 8G.10].

$T_{\sigma_n} \cap M = T_{\sigma_n} \cap M[g]$. By absoluteness of wellfoundedness, we have $M[g] \models (x, y_\psi) \in p[T_{\sigma_n}] \iff (x, y_\psi) \in p[T_{\sigma_n}^{M[g]}] \iff (x, y_\psi) \in p[T_{\sigma_n}]$.

(2) Since M is countable, the set of Cohen reals over M in V is comeager. Therefore, this claim holds by the argument for the Baire property for definable sets in Solovay's model. \square

The above argument for the non-existence of a selector with Borel values can now be used for the following results. We fix codes for Σ_n^1 sets via Σ_n^1 -universal Σ_n^1 sets.

Theorem 5.18. *Assuming $\text{IA}_{\text{proj}}^{\mathbb{C}}$, there is no Baire measurable selector for I with projective values.*

Proof. This is proved similarly to Theorem 5.5. We only indicate the two necessary changes. In the proof of the first claim, $\text{IA}_{\text{proj}}^{\mathbb{C}}$ implies that the (projective) statement $[\dot{T}_n^g] \cap B_{F(\dot{C}^g)} = [\dot{T}_n^g] \cap B_{\dot{F}^g(\dot{C}^g)} = \emptyset$ is absolute between $M[g]$ and V . The second change is at the very end of the proof. Here projective absoluteness between $M[g \times h]$ and V by $\text{IA}_{\text{proj}}^{\mathbb{C}}$ guarantees that $x = \sigma^G \in B_{F(\dot{C}^g)}$ and thus $B_{F(\dot{C}^g)} \setminus B_{\dot{C}^g}$ contains every Cohen real over $M[g]$ in V . As before, this contradicts the fact that F is a selector for I . \square

Note that by Lemma 5.15, PD is sufficient for the previous result. Thus PD implies that there is no projective selector for I . In particular, there is no selector $\text{BOREL}(\omega\Omega) \rightarrow \text{PROJ}(\omega\Omega)$ for $=_I$ that is induced by a universally Baire measurable function (assuming PD).

Note that some assumption beyond ZFC is necessary to prove this statement. For instance, in L there are projective selectors for all projectively coded ideals I , since there is a projective, in fact a Σ_2^1 , wellorder of the reals.

The previous arguments can also be used to show that it is consistent with ZF that there is no selector at all for I .

Suppose that κ is an uncountable cardinal and G is a \mathbb{P} -generic filter over V for $\mathbb{P} = \text{Add}(\omega, \kappa)$ or $\mathbb{P} = \text{Col}(\omega, <\kappa)$. We call $V^* = \bigcup_{\alpha < \kappa} V[G \restriction \alpha]$ a *symmetric extension* for \mathbb{P} . Note that $V^* = \text{HOD}_{V^*}^{V[G]}$ (the class of sets which are hereditarily ordinal definable in $V[G]$ with parameters from V^*) by homogeneity and for cardinals κ of uncountable cofinality, $\mathbb{R}^{V^*} = \mathbb{R}^{V[G]}$.

Theorem 5.19. *There are no selectors for I in the following models of ZF.*

- (a) *Symmetric extensions V^* for $\text{Add}(\omega, \kappa)$ and $\text{Col}(\omega, <\kappa)$, where κ is any uncountable cardinal.*
- (b) *Solovay's model.*
- (c) *$L(\mathbb{R})$ assuming $\text{AD}^{L(\mathbb{R})}$.*

Proof. The proof of the first claim is similar to that of Theorem 5.5.

Suppose that F is a selector for I in V^* . It is definable from an element x_0 of $V[G \restriction \alpha]$ for some $\alpha < \kappa$. We can further assume that $x_0 \in V$. Hence F is continuous on the set of Cohen reals over V in V^* ; let \dot{F} be a name for this function.

We follow the proof of Theorem 5.5 but work with V instead of M . The first claim in the proof of Theorem 5.5 is replaced by the following claim.

We will write V^* for a \mathbb{P} -name for V^* to keep the notation simple.

Claim 5.20. $1 \Vdash_{\mathbb{T}^\omega \times \mathbb{P}}^V V^* \models [\dot{T}_n] \cap B_{\dot{F}(\dot{C})} \neq \emptyset$ for all $n \in \omega$.

Proof. Assume towards a contradiction that $(p, q) \Vdash_{\mathbb{T}^\omega \times \mathbb{P}}^V V^* \models [\dot{T}_n] \cap B_{\dot{F}(\dot{C})} = \emptyset$ for some $p \in \mathbb{T}^\omega$, $q \in \mathbb{P}$ and $n \in \omega$. We can assume $q = 1$ by homogeneity. Let $g \times h$ be a $\mathbb{T}^\omega \times \mathbb{P}$ -generic filter over V with $p \in g$ whose symmetric model is V^* . Then $[\dot{T}_n^g] \cap B_{F(\dot{C}^g)} = [\dot{T}_n^g] \cap B_{\dot{F}^g(\dot{C}^g)} = \emptyset$. But this contradicts the assumption that F is a selector with respect to I . \square

There is a \mathbb{T}^ω -name σ in V that is forced to be an element of $[\dot{T}_n] \cap B_{\dot{F}(\dot{C})} \cap V^*$ by the previous claim. The next steps of the proof are as before.

In the end of the proof, we rearrange $V[g \times h]$ as a $\mathbb{T}^\omega \times \text{Add}(\omega, 1)$ -generic extension $V[G \times H]$ with $\sigma^G = x$ as before. Then $\sigma^G \in B_{F(\dot{C}^g)}$ by the choice of σ . Thus $B_{F(\dot{C}^g)} \setminus B_{\dot{C}^g}$

contains every Cohen real over $V[g]$ in V^* . Since there are uncountably many Cohen reals over $V[g]$ in V^* , this contradicts the assumption that F is a selector.

The second claim holds since Solovay's model is obtained via a symmetric extension for $\text{Col}(\omega, <\kappa)$. Note that this model is also a $\text{Add}(\omega, \omega_1)$ -generic extension of an intermediate model.

The last claim follows from [SS06, Theorem 0.1] and the first claim. By this theorem and $\text{AD}^{L(\mathbb{R})}$, $L(\mathbb{R})^V = L(\mathbb{R})^{V^*}$ for some symmetric extension V^* for $\text{Col}(\omega, <\kappa)$ for some κ over some ground model N which is an inner model of some generic extension of V . \square

5.3. Density points via convergence. In this section we discuss the notion of density point introduced in [PWBW85]. We show that this notion does not satisfy the analogue of Lebesgue's density theorem for any of the tree forcings listed in the next section, except Cohen and random forcing.

Lebesgue's density theorem was generalized to the σ -ideals of meager sets on Polish metric spaces in [PWBW85, Theorem 2]. To this end, a notion of density points for ideals was introduced. This notion is based on the following well-known measure theoretic lemma.

Lemma 5.21. *Suppose that (X, d, μ) is a metric measure space, f is a Borel-measurable function and $\vec{f} = \langle f_n \mid n \in \omega \rangle$ is a sequence of Borel-measurable functions from (X, d) to \mathbb{R} . The following statements are equivalent:*

- (a) $f_n \rightarrow f$ converges in measure, i.e. for all $\epsilon > 0$ we have

$$\lim_{n \rightarrow \infty} \mu(\{x \in X \mid |f_n(x) - f(x)| \geq \epsilon\}) = 0.$$

- (b) Every subsequence of \vec{f} has a further subsequence that converges pointwise μ -almost everywhere.

Condition (b) is suitable for generalizations to other ideals, since it only mentions the ideal of null sets, but not the measure itself.

Definition 5.22. Let I be a σ -ideal on ${}^\omega\Omega$ and $A \subseteq {}^\omega\Omega$.

- (1) Given $x \in {}^\omega\Omega$, let f_n denote the characteristic function of $\sigma_{x \upharpoonright n}^{-1}(A \cap N_{x \upharpoonright n})$ for each $n \in \omega$. Define x to be an I -convergence density point of A if every subsequence of $\vec{f} = \langle f_n \mid n \in \omega \rangle$ has a further subsequence that converges I -almost everywhere (i.e. on a set A with ${}^\omega\Omega \setminus A \in I$) to the constant function on ${}^\omega\Omega$ with value 1.
- (2) By the I -convergence density property we mean the statement that for any $B \in \text{BOREL}({}^\omega\Omega)$ and the set C of I -convergence density points of B , we have $B \triangle C \in I$.

We first consider tree forcings on ${}^\omega\omega$. For $s \in {}^{<\omega}\omega$ and $f_0, \dots, f_m \in {}^\omega\omega$, let

$$T_{s, f_0, \dots, f_m} = \{t \in C_s \mid \forall i \leq m \forall n \geq |s| \ t(n) \neq f_i(n)\}$$

(cf. Definition 6.1 (6) below) and let $f \in {}^\omega\omega$ denote the constant function with value 0.

Lemma 5.23. *If $N_{\langle 0 \rangle}$ and $[T_{\emptyset, f}]$ are I -positive, then the I -convergence density property fails. In particular, this holds for $I_{\mathbb{H}}$, $I_{\mathbb{E}}$, $I_{\mathbb{M}}$ and $I_{\mathbb{L}_A}$ if A contains all cofinite sets.*

Proof. We claim that no $x \in {}^\omega\omega$ is an I -convergence density point of $[T_{\emptyset, f}]$. We have $\sigma_t^{-1}([T_{\emptyset, f}]) = [T_{\emptyset, f}]$ if $t \in T_{\emptyset, f}$ and $\sigma_t^{-1}([T_{\emptyset, f}]) = \emptyset$ otherwise. Hence $\sigma_t^{-1}([T_{\emptyset, f}]) \cap N_{\langle 0 \rangle} = \emptyset$ for all $t \in {}^{<\omega}\omega$. Thus x is not an I -convergence density point of $[T_{\emptyset, f}]$. \square

We now turn to tree forcings on ${}^\omega 2$. For $s \in {}^{<\omega} 2$ and $N \in [\omega]^\omega$ let

$$T_{s, N} = \{t \in {}^{<\omega} 2 \mid t \in C_s \ \& \ \forall n \ t(n) = 1 \Rightarrow n \in N\}$$

(cf. Definition 6.1 (8) below).

Lemma 5.24. *Suppose that $N_{\langle 1 \rangle}$ and $[T_{\emptyset, N}]$ are I -positive for some infinite set N . Then the I -convergence density property fails. In particular, this holds for \mathbb{R}_A and \mathbb{V}_A for any subset A of $\mathcal{P}(\omega)$ that contains at least one infinite set.*

Proof. It is sufficient to show that no $x \in [T_{\emptyset, N}]$ is an I -convergence density point of $[T_{\emptyset, N}]$. We have $\sigma_{x \upharpoonright n}^{-1}([T_{\emptyset, N}]) \cap N_{\langle 1 \rangle} = \emptyset$ for all $n \notin N$. Since $N_{\langle 1 \rangle}$ is I -positive, any infinite strictly increasing sequence in N witnesses that x is not an I -convergence density point. \square

The next lemma takes care of the remaining tree forcings.

Lemma 5.25. *Suppose that I is an ideal on ${}^\omega 2$ with $I_{\mathbb{S}} \subseteq I \subseteq I_{\mathbb{R}}$. Then the I -convergence density property fails.*

Proof. Let

$$T = \{t \in {}^{<\omega} 2 \mid \forall i \in \omega \ (t(i) = 1 \Rightarrow \exists j \in \omega \ (i = 2^j + 1))\}.$$

It is sufficient to show that no $x \in {}^\omega 2$ is an I -convergence density point of $[T]$. Otherwise there is a strictly increasing sequence $\vec{n} = \langle n_i \mid i < \omega \rangle$ such that

$$A = \{y \in {}^\omega 2 \mid \exists i \ \forall j \geq i \ y \in \sigma_{x \upharpoonright n_j}^{-1}([T])\}$$

is co-countable. Let

$$A_{i,j} = \sigma_{x \upharpoonright n_i}^{-1}([T]) \cap \sigma_{x \upharpoonright n_j}^{-1}([T])$$

for $i < j$ in ω . Since $A \subseteq \bigcup_{i,j \in \omega, i < j} A_{i,j}$ is uncountable, $A_{i,j}$ is uncountable for some $i < j$. Let $a <_{\text{lex}} b <_{\text{lex}} c$ be elements of $A_{i,j}$. We denote the longest common initial segment of $d, e \in {}^{<\omega} 2$ by $d \wedge e$. We can assume without loss of generality that $a \wedge b = s$, $a \wedge c = b \wedge c = t$ and $s \subsetneq t$. Then s and t are splitting nodes in $\sigma_{x \upharpoonright n_i}^{-1}(T)$ and $\sigma_{x \upharpoonright n_j}^{-1}(T)$. Hence $s + n_i$, $s + n_j$, $t + n_i$, $t + n_j$ are of the form $2^k + 1$. Since $s \neq t$, $n_j - n_i$ can be written in the form $2^k - 2^l$ in two different ways. But this contradicts the easy fact that k, l are uniquely determined by $2^k - 2^l$. \square

Let \mathbb{P}_{E_0} denote E_0 -forcing [Zap08, Section 4.7.1] and \mathbb{W} an appropriate representation of *Willoutree forcing* [Bre95, Section 1.1]. Since $\mathbb{R} \subseteq \mathbb{V} \subseteq \mathbb{W} \subseteq \mathbb{P}_{E_0} \subseteq \mathbb{S}$, the previous result holds for the ideals $I_{\mathbb{P}}$ associated to these forcings as well.

6. A LIST OF TREE FORCINGS

We review definitions of some tree forcings for the reader's convenience. If $N \subseteq \omega$ and $m \in \omega$, we write $m + N = \{m + n \mid n \in N\}$. Moreover, a subset A of $\mathcal{P}(\omega)$ is called *shift invariant* if $N \in A \iff m + N \in A$ for all $m \in \omega$.

Definition 6.1. Assume that A is a subset of $\mathcal{P}(\omega)$.

- (1) *Random forcing* is the collection of perfect subtrees T of ${}^{<\omega} 2$ with $\mu([T_s]) > 0$ for all $s \in T$ with $\text{stem}(T) \sqsubseteq s$.
- (2) *Cohen forcing* \mathbb{C} is collection of *cones* $C_s = \{t \in {}^{<\omega} \omega \mid s \sqsubseteq t \text{ or } t \sqsubseteq s\}$ for $s \in {}^{<\omega} \omega$.
- (3) *Sacks forcing* \mathbb{S} is the collection of all perfect subtrees of ${}^{<\omega} 2$.
- (4) *Miller forcing* \mathbb{M} is the collection of *superperfect* subtrees T of ${}^{<\omega} \omega$. This means that above every node in T there is some infinitely splitting node t in T , i.e. t has infinitely many direct successors.
- (5) *Hechler forcing* \mathbb{H} is the collection of trees

$$T_{s,f} = \{t \in C_s \mid \forall n \geq |s| \ t(n) \geq f(n)\}$$

for $s \in {}^{<\omega} \omega$ and $f \in {}^\omega \omega$.

- (6) *Eventually different forcing* \mathbb{E} is the collection of trees

$$T_{s,f_0,\dots,f_m} = \{t \in C_s \mid \forall i \leq m \ \forall n \geq |s| \ t(n) \neq f_i(n)\}$$

for $s \in {}^{<\omega} \omega$ and $f_0, \dots, f_m \in {}^\omega \omega$.

- (7) *A-Laver forcing* \mathbb{L}_A is the collection of subtrees T of ${}^{<\omega} \omega$ such that for every $t \in T$ with $\text{stem}(T) \sqsubseteq t$, the set of immediate successors of t in T is an element of A . *Laver forcing* is \mathbb{L}_F for the Fréchet filter F of cofinite sets.

(8) *A-Mathias forcing* \mathbb{R}_A is the collection of trees

$$T_{s,N} = \{t \in {}^{<\omega}2 \mid t \in C_s \text{ \& } \forall n \ t(n) = 1 \Rightarrow n \in N\}$$

for $s \in {}^{<\omega}2$ and $N \in A$. *Mathias forcing* \mathbb{R} is \mathbb{R}_A for $A = [\omega]^\omega$.

(9) *A-Silver forcing* \mathbb{V}_A is the collection of trees

$$T_f = \{t \in {}^{<\omega}2 \mid \forall n \in \text{dom}(t) \cap \text{dom}(f) \ f(n) \leq t(n)\},$$

where $\text{dom}(f) = \omega \setminus N$ for some $N \in A$. Silver forcing \mathbb{V} is \mathbb{V}_A for $A = [\omega]^\omega$.

All of these satisfy the above condition for collections of trees \mathbb{P} that $T_s \in \mathbb{P}$ for all $T \in \mathbb{P}$ and $s \in T$. Moreover, random forcing, Cohen forcing, Sacks forcing, Hechler forcing, eventually different forcing, and A -Laver forcing are *shift invariant* in the sense that for all T and $s \in {}^{<\omega}2$, $T \in \mathbb{P} \iff \sigma_s(T) \in \mathbb{P}$. If A is shift invariant, then \mathbb{R}_A and \mathbb{V}_A are also shift invariant.

Note that Cohen forcing, Hechler forcing, eventually different forcing, Laver forcing, and Silver forcing are topological, while random forcing, Sacks forcing, and Miller forcing are not.

7. AN EXPLICIT CONSTRUCTION OF DENSITY POINTS

In this section, we show how to explicitly construct density points of a closed set C of positive measure.

In fact, by the next result there is an algorithm that takes as input a list of data from C and outputs a perfect tree (level by level) all of whose branches are density points. (By choosing e.g. the leftmost branch, we can approximate a single density point with arbitrary precision.) The input is a tree T together with weights $w_t = \frac{\mu([T] \cap N_t)}{\mu(N_t)} > 0$ for all $t \in T$; we call this a *weighted tree*. The weights are given as inputs with arbitrary precision.

Theorem 7.1. *There is a partial computable function that takes as input any pair (T, q) , where T is a weighted tree and $q \in \mathbb{Q} \cap (0, 1)$, and produces a perfect tree $S \subseteq T$ with*

- (a) $[S] \subseteq D_\mu([T])$ and
- (b) $\mu([S]) \geq q\mu([T])$.

Since $\mu([T_s])$ is right-c.e.²³ in the oracle T for all $s \in T$, one has the following immediate consequence.

Corollary 7.2. *For any tree T with $\mu([T]) > 0$ and $q \in \mathbb{Q} \cap (0, 1)$, there is a $\Delta_2^0(T)$ -definable perfect tree S such that (a) and (b) of Theorem 7.1 hold. Moreover, there is a $\Delta_2^0(T)$ -coded F_σ set $A \subseteq D_\mu([T])$ with $\mu(A) = \mu([T])$.*

Note that for all strongly linked collections of trees \mathbb{P} (see Definition 4.15) listed in Section 6 and all $T \in \mathbb{P}$, $S = T$ already satisfies the conditions in Theorem 7.1. For these collections, any $T \in \mathbb{P}$ has the property that for all $x \in [T]$, there are infinitely many $n \in \omega$ such that there is some $S \leq T$ with $x \in [S]$ and $\text{stem}(S) = x \restriction n$. Hence all elements of $[T]$ are density points of $[T]$ by the proof of Lemma 4.17 below.

Theorem 7.1 will follow from the next lemmas. To state them, we fix the following notation: Let $C = [T]$, $L_{t,i} = \text{Lev}_{|t|+i}(T_t)$ be the level of T_t at height $|t| + i$ and write

$$w_{t,i} = \frac{|L_{t,i}|}{2^i} = \mu(N_t)^{-1} \frac{|L_{t,i}|}{2^{|t|+i}}$$

for all $t \in {}^{<\omega}2$ and $i \in \omega$. This is the *relative size* of levels of T above t . The next result shows that these values converge to the relative measure at t .

Lemma 7.3. $\lim_{i \rightarrow \infty} w_{t,i} = w_t$ for all $t \in {}^{<\omega}2$.

Proof. We have $w_t \leq \lim_{i \rightarrow \infty} w_{t,i}$, since $C \cap N_t \subseteq \bigcup_{u \in L_{t,i}} N_u$ and hence $w_t \leq w_{t,i}$ for all $i \in \omega$. To prove that $\lim_{i \rightarrow \infty} w_{t,i} \leq w_t$, suppose that $\epsilon > 0$ is given. Let U be an open set with $C \cap N_t \subseteq U$ and $\mu(U) < \mu(C \cap N_t) + \epsilon \cdot \mu(N_t)$. By compactness of C , we can

²³See [Nie09, Definition 1.8.14].

assume that U is a finite union of basic open sets. We can thus write $U = \bigcup_{j \leq n} N_{s_j}$ for some $\vec{s} = \langle s_j \mid j \leq n \rangle$ that consists of pairwise incompatible sequences s_j of the same length $|t| + i$. Since $C \cap N_t \subseteq U$, we have

$$w_{t,i} = \mu(N_t)^{-1} \frac{|L_{t,i}|}{2^{|t|+i}} \leq \frac{\mu(\bigcup_{j \leq n} N_{s_j})}{\mu(N_t)} = \frac{\mu(U)}{\mu(N_t)}.$$

Hence $w_{t,i} - w_t \leq \frac{\mu(U) - \mu(C \cap N_t)}{\mu(N_t)} < \epsilon$ by the previous inequality and the definition of w_t . \square

For any $t \in {}^{<\omega}2$ and $i \in \omega$, let

$$r_{t,i} = \inf \{c \in (0, 1) \mid \frac{|\{u \in L_{t,i} \mid w_u \geq c\}|}{|L_{t,i}|} \geq c\}$$

denote the ratio of nodes on level $|t| + i$ above t with large weight.

Lemma 7.4. $\liminf_{i \rightarrow \infty} r_{t,i} = 1$ for all $t \in {}^{<\omega}2$.

Proof. Let $b = w_t$ and assume that $c \in (0, 1)$ is given. Since $b > bc + b(1 - c)c$, there is some $\epsilon > 0$ with $b > (b + \epsilon)c + (b + \epsilon)(1 - c)c$. By Lemma 7.3, we can take $i \in \omega$ to be sufficiently large such that $w_{t,i} \leq b + \epsilon$. Moreover, let α denote the fraction of nodes $u \in L_{t,i}$ with weight $w_u \geq c$. Then

$$b = w_t = 2^{-i} \sum_{u \in L_{t,i}} w_u \leq w_{t,i} \alpha + w_{t,i} (1 - \alpha) c.$$

We claim that $\alpha \geq c$. Otherwise $\alpha < c$ and $b \leq w_{t,i} \alpha + w_{t,i} (1 - \alpha) c \leq w_{t,i} c + w_{t,i} (1 - c) c$. Since $w_{t,i} \leq b + \epsilon$, we obtain $b \leq (b + \epsilon)c + (b + \epsilon)(1 - c)c$, contradicting the definition of ϵ . \square

We need the following notion to ensure that weights converge to 1 along branches of the tree constructed below. We say that $v \sqsupseteq t$ is (t, a) -good if $w_u \geq a$ for all u with $t \sqsubseteq u \sqsubseteq v$; otherwise it is called (t, a) -bad. Let

$$s_{t,a,i} = \frac{|\{u \in L_{t,i} \mid u \text{ is } (t, a)\text{-good}\}|}{|L_{t,i}|}$$

denote the fraction of (t, a) -good nodes on level $|t| + i$ above t .

We fix a computable function $f: \mathbb{Q} \cap (0, 1) \rightarrow \mathbb{Q} \cap (0, 1)$ such that $\frac{1-b}{b(b-a)} < \frac{1-a}{a}$ holds for all $a, b \in \mathbb{Q} \cap (0, 1)$ with $b > f(a)$.

Lemma 7.5. If $a \in \mathbb{Q} \cap (0, 1)$ and $w_t = b > f(a)$, then $\liminf_{i \rightarrow \infty} s_{t,a,i} \geq a$.

Proof. Since $b > f(a)$, there is some $c \in (0, 1)$ with $1 - b < \frac{1-a}{a} b(b-a)c$. Let i be sufficiently large such that the fraction of $v \in L_{t,i}$ with $w_v \geq c$ is at least b by Lemma 7.4. Then the fraction of nodes $v \in L_{t,i}$ with $w_v < c$ is at most $1 - b$ and their number at most $2^i w_{t,i} (1 - b)$.

Let A the set of (t, a) -bad nodes in $L_{t,i}$, $U = \bigcup_{v \in A} N_v$ and $\alpha = \frac{|A|}{|L_{t,i}|}$. We aim to show that $\alpha \leq 1 - a$. The number of (t, a) -bad nodes in $L_{t,i}$ is $2^i w_{t,i} \alpha$. Since all of these except at most $2^i w_{t,i} (1 - b)$ have weight at least c , we have

$$2^{|t|+i} \mu(C \cap U) = \sum_{v \in A} w_v \geq (2^i w_{t,i} \alpha - 2^i w_{t,i} (1 - b)) c.$$

Claim 7.6. $1 - b \geq \frac{1-a}{a} w_{t,i} (\alpha - (1 - b)) c$.

Proof. For each $v \in A$, take some u_v with $t \sqsubseteq u_v \sqsubseteq v$ and $w_{u_v} < a$ by the definition of (t, a) -bad. In particular $\frac{1-w_{u_v}}{w_{u_v}} > \frac{1-a}{a}$.

Let $B = \{u_v \mid v \in A\}$ and B^* the set of \sqsubseteq -minimal elements of B . For each $v \in A$ and $u = u_v$

$$\mu(N_u \setminus C) = 2^{-|u|} (1 - w_u) > 2^{-|u|} w_u \frac{1-a}{a} = \frac{1-a}{a} \mu(C \cap N_u).$$

Since the sets N_u for $u \in B^*$ are pairwise disjoint, the previous inequality implies

$$\mu(U \setminus C) = \sum_{u \in B^*} \mu(N_u \setminus C) > \frac{1-a}{a} \sum_{u \in B^*} \mu(C \cap N_u) = \frac{1-a}{a} \mu(C \cap U).$$

By this inequality and the one before the claim, we have $\mu(N_t \setminus C) \geq \mu(U \setminus C) > \frac{1-a}{a} \mu(C \cap U) \geq 2^{-|t|} \frac{1-a}{a} (w_{t,i} \alpha - w_{t,i} (1-b))c$. Since $\frac{\mu(N_t \setminus C)}{2^{-|t|}} = \frac{\mu(N_t \setminus C)}{\mu(N_t)} = 1 - w_t = 1 - b$ and $\mu(N_t) = 2^{-|t|}$, the claim follows. \square

It is sufficient to show that $\alpha \leq 1 - a$. Otherwise by the previous claim and since $b = w_t \leq w_{t,i}$

$$\begin{aligned} 1 - b &\geq \frac{1-a}{a} w_{t,i} (\alpha - (1-b))c \\ &> \frac{1-a}{a} w_{t,i} ((1-a) - (1-b))c \\ &= \frac{1-a}{a} w_{t,i} (b-a)c \\ &\geq \frac{1-a}{a} b(b-a)c. \end{aligned}$$

But this contradicts the choice of c . \square

Proof of Theorem 7.1. Let $\vec{a} = \langle a_i \mid i \in \omega \rangle$ be a computable sequence in $\mathbb{Q} \cap (0, 1)$ with $\prod_{i \in \omega} a_i^2 > q$. Using Lemmas 7.4 and 7.5, we will inductively construct a strictly increasing sequence $\vec{n} = \langle n_i \mid i \in \omega \rangle$ and sets $S_i \subseteq \text{Lev}_{n_i}(T)$ with $n_0 = 0$ and $S_0 = \{\emptyset\}$ by induction on $i \in \omega$. The sets S_i are compatible levels of a tree in the sense that each $t \in S_i$ has an extension $u \in S_{i+1}$ and conversely, each $u \in S_{i+1}$ extends some $t \in S_i$. We further let $T^{(i)} = \{t \in T \mid \exists u \in S_i (t \sqsubseteq u \vee u \sqsubseteq t)\}$ denote the subtree of T induced by S_i .

We will maintain during the induction that (a) u is (t, a_i) -good for all $t \in S_{i+1}$ and $u \in S_{i+2}$ and (b) $\frac{\mu([T^{(i+1)}])}{\mu([T^{(i)}])} \geq a_i^2$.

Description. We now describe the construction. We simultaneously construct auxiliary numbers $a'_i, a''_i, b_i \in \mathbb{Q}$ and $j_i \in \omega$ with $a_i < a'_i < a''_i < b_i < 1$ and $b_i > f(a''_i)$ for all $i \geq 1$. It is not hard to see that all steps are effective.

Let $n_0 = 0$ and $S_0 = \text{Lev}_{n_0}(T) = \text{Lev}_0(T)$.

For $i = 1$, we first choose some $a'_1 \in \mathbb{Q}$ with $a_1 < a'_1 < 1$ and $b_1 \in \mathbb{Q} \cap (0, 1)$ with $b_1 > f(a'_1)$. By Lemma 7.4 applied to $t = \emptyset$, there is some j_1 with $r_{\emptyset, j_1} > b_1$. Let $n_1 = j_1$ and S_1 a subset of $\text{Lev}_{n_1}(T)$ with $w_u > b_1$ for all $u \in S_1$ and $\frac{|S_1|}{|\text{Lev}_{n_1}(T)|} > b_1$. We can further take b_1

to be sufficiently large such that $\frac{\mu([T^{(1)}])}{\mu([T^{(0)}])} > a_0^2$ by $r_{\emptyset, j_1} > b_1$ and the definition of r_{\emptyset, j_1} .

Fix $i \geq 1$ and assume that step i is completed. First take some $a'_{i+1}, a''_{i+1} \in \mathbb{Q}$ with $a_{i+1} < a'_{i+1} < a''_{i+1} < 1$ and $1 - a'_{i+1} < a'_i - a_i$. Let $b_{i+1} \in \mathbb{Q} \cap (0, 1)$ with $b_{i+1} > f(a''_{i+1})$, a''_{i+1}, a_i . By Lemma 7.4, there is some j_{i+1} such that $r_{t, j_{i+1}} > b_{i+1}$ for all $t \in S_i$. Since $b_i > f(a''_i)$ by the inductive hypothesis and $a'_i < a''_i$, we can take j_{i+1} to satisfy $s_{t, a''_i, j_{i+1}} > a'_i$ for all $t \in S_i$ by Lemma 7.5. Let $n_{i+1} = n_i + j_{i+1}$. By definition of j_{i+1} , there is a subset S_{i+1} of $\text{Lev}_{n_{i+1}}(T)$ such that for all $u \in S_{i+1}$, we have $w_u > b_{i+1}$, there is a (unique) $t \in S_i$ with $t \sqsubseteq u$, u is (t, a'_i) -good and $\frac{|L_{t, j_{i+1}} \setminus S_{i+1}|}{|L_{t, j_{i+1}}|} < (1 - a'_i) + (1 - b_{i+1})$.

Verification. We show that the algorithm computes the required tree. Clearly condition (a) is maintained in the construction. The next claim shows (b).

Claim 7.7. $\frac{\mu([T^{(i+1)}])}{\mu([T^{(i)}])} \geq a_i^2$ for all $i \in \omega$.

Proof. This is clear for $i = 0$. Let $i \geq 1$ and fix any $t \in S_i$. Since $a'_{i+1} < b_{i+1}$ and by the definition of S_{i+1} , we have $\frac{|L_{t, j_{i+1}} \setminus S_{i+1}|}{|L_{t, j_{i+1}}|} \leq (1 - a'_i) + (1 - b_{i+1}) \leq (1 - a'_i) + (1 - a'_{i+1}) \leq 1 - a_i$. Hence $\frac{|L_{t, j_{i+1}} \cap S_{i+1}|}{|L_{t, j_{i+1}}|} \geq a_i$.

Since each $u \in S_{i+1}$ has weight at least b_{i+1} , we have $c := \frac{\mu([T^{(i+1)}] \cap N_t)}{\mu([T^{(i)}] \cap N_t)} \geq c' := a_i b_{i+1}$. By the definition of $T^{(i+1)}$ from S_{i+1} , $d := \frac{\mu([T^{(i)}] \cap N_t) \setminus [T^{(i+1)}])}{\mu([T^{(i)}] \cap N_t)} \leq d' := 1 - a_i$. Moreover $c + d = 1$. Since $c \geq c'$, $\frac{c+d'}{c} = 1 + \frac{d'}{c} \leq 1 + \frac{d'}{c'} = \frac{c'+d'}{c'}$. Since $d \leq d'$ and by the last inequality

$\frac{c}{c+d} \geq \frac{c}{c+d'} \geq \frac{c'}{c'+d'}$. Therefore $c = \frac{c}{c+d} \geq \frac{c'}{c'+d'} = \frac{a_i b_{i+1}}{a_i b_{i+1} + (1-a_i)} \geq a_i b_{i+1} \geq a_i^2$. Since this inequality holds for all $t \in S_i$, we have $\frac{\mu([T^{(i+1)}])}{\mu([T^{(i)}])} \geq a_i^2$. \square

To see that conditions (a) and (b) are sufficient, let S be the unique perfect subtree of ${}^{<\omega}2$ with $\text{Lev}_{n_i}(S) = S_i$ for all $i \in \omega$. This tree can be computed level by level via the algorithm above. We have $\lim_{i \rightarrow \infty} a_i = 1$ by the definition of \vec{a} . Thus (a) implies that all elements of $[S]$ are density points of $[T]$. Moreover, $\mu([S]) = \inf_{i \in \omega} \mu([T^{(i)}]) \geq \prod_{i \in \omega} a_i^2 \mu([T]) < q\mu([T])$ by (b) as required. \square

The previous result provides a finitized proof of Lebesgue's density theorem for Lebesgue measure on Cantor space, since any Borel set can be approximated in measure by closed subsets. To see this, note that trivially $D(A) \cap D({}^\omega\Omega \setminus A) = \emptyset$ for any subset A of ${}^\omega\Omega$. Thus it is sufficient to show that for any Borel set A and any $\epsilon > 0$, there is a closed subset C of A with $\mu(A \setminus C) < \epsilon$ consisting of density points of A ; the density property for A follows by applying this property to both A and its complement. To see that this property holds, take a closed subset B of A with $\mu(A \setminus B) < \frac{\epsilon}{2}$. By Theorem 7.1, there is a closed subset C of B with $\mu(B \setminus C) < \frac{\epsilon}{2}$ that consists of density points of B and therefore also of A . Since $\mu(A \setminus C) < \epsilon$, C is as required.

Note that the algorithm also produces lower bounds for weights along branches of S .

8. OPEN PROBLEMS

We end with some open problems. Our main goal is to prove the equivalence of the properties discussed in Section 1 for many other ideals.

Question 8.1. *Are the shift density property, the existence of a simply definable selector and the ccc equivalent for all simply definable σ -ideals?*

This is of interest for the ideals studied in [Zap08], for instance for the K_σ -ideal. For the latter, we suggest to generalize the proof idea of Theorem 5.5 to answer the next question.

Question 8.2. *Is there a Baire measurable selector with Borel values for the K_σ -ideal?*

A promising related problem is whether there is a relation between the shift density property and the condition that the collection of Borel sets modulo I carries a natural Polish metric.

Our proof of the density property for strongly linked tree ideals is based on the fact from Section 4.2 that \mathbb{P} -measurable sets form a σ -ideal if \mathbb{P} has the ω_1 -covering property. Can the latter assumption be omitted? In this case, the next question would also have a positive answer.

Question 8.3. *Are all Borel sets \mathbb{P} -measurable for all tree forcings \mathbb{P} ?*

Note that any counterexample \mathbb{P} collapses ω_1 if we assume CH; in this case \mathbb{P} preserves ω_1 if and only if it has the ω_1 -covering property by a similar argument as in the proof of Lemma 4.4.

In Section 5.2, we proved from PD that there is no selector $\text{BOREL}({}^\omega\Omega) \rightarrow \text{PROJ}({}^\omega\Omega)$ for $=_I$ that is induced by a universally Baire measurable function on the codes (Theorem 1.4). To this end, we introduced the principle $\text{IA}_{\text{proj}}^{\mathbb{C}}$ of generic absoluteness. This follows from PD by Lemma 5.15. Our results leave its precise consistency strength open.

Question 8.4. *What is the consistency strength of $\text{IA}_{\text{proj}}^{\mathbb{C}}$?*

Recent unpublished work of the first and second-listed authors answers this by showing that the consistency strength of $\text{IA}_{\text{proj}}^{\mathbb{C}}$ is that of ZFC (see [MS]).

REFERENCES

- [AC13] Alessandro Andretta and Riccardo Camerlo. The descriptive set theory of the Lebesgue density theorem. *Adv. Math.*, 234:1–42, 2013.
- [AC19] Alessandro Andretta and Riccardo Camerlo. Analytic sets of reals and the density function in the Cantor space. *Eur. J. Math.*, 5(1):49–80, 2019.
- [ACC13] Alessandro Andretta, Riccardo Camerlo, and Camillo Costantini. The density point property. Slides of a talk given by Alessandro Andretta at the 4th European Set Theory Conference, Mon St. Benet, Catalunya, 2013.
- [ACC19] Alessandro Andretta, Riccardo Camerlo, and Camillo Costantini. Lebesgue density and exceptional points. *Proc. Lond. Math. Soc. (3)*, 118(1):103–142, 2019.
- [BG20] Marek Balcerzak and Szymon Głąb. A lower density operator for the borel algebra. *Results in Mathematics*, 75(2):50, 2020.
- [BHL05] Jörg Brendle, Lorenz Halbeisen, and Benedikt Löwe. Silver measurability and its relation to other regularity properties. *Math. Proc. Cambridge Philos. Soc.*, 138(1):135–149, 2005.
- [BJ95] Tomek Bartoszyński and Haim Judah. *Set theory*. A K Peters, Ltd., Wellesley, MA, 1995. On the structure of the real line.
- [BKW18] Jörg Brendle, Yurii Khomskii, and Wolfgang Wohofsky. Cofinalities of Marczewski-like ideals. Preprint, 2018.
- [BL99] Jörg Brendle and Benedikt Löwe. Solovay-type characterizations for forcing-algebras. *The Journal of Symbolic Logic*, 64(3):1307–1323, 1999.
- [Bre95] Jörg Brendle. Strolling through paradise. *Fund. Math.*, 148(1):1–25, 1995.
- [Buk11] Lev Bukovský. *The structure of the real line*, volume 71 of *Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series)*. Birkhäuser/Springer Basel AG, Basel, 2011.
- [Cum10] James Cummings. Iterated forcing and elementary embeddings. In *Handbook of set theory. Vols. 1, 2, 3*, pages 775–883. Springer, Dordrecht, 2010.
- [FKK16] Sy Friedman, Yurii Khomskii, and Vadim Kulikov. Regularity properties on the generalized reals. *Ann. Pure Appl. Logic*, 167(4):408–430, 2016.
- [FMW92] Qi Feng, Menachem Magidor, and Hugh Woodin. Universally Baire sets of reals. In *Set theory of the continuum (Berkeley, CA, 1989)*, volume 26 of *Math. Sci. Res. Inst. Publ.*, pages 203–242. Springer, New York, 1992.
- [GRSS95] Martin Goldstern, Miroslav Repický, Saharon Shelah, and Otmar Spinas. On tree ideals. *Proceedings of the American Mathematical Society*, 123(5):1573–1581, 1995.
- [Hal03] Lorenz Halbeisen. Making doughnuts of Cohen reals. *Mathematical Logic Quarterly*, 49(2):173–178, 2003.
- [Hjo10] Greg Hjorth. Borel equivalence relations. In *Handbook of set theory. Vols. 1, 2, 3*, pages 297–332. Springer, Dordrecht, 2010.
- [Ike10] Daisuke Ikegami. Forcing absoluteness and regularity properties. *Ann. Pure Appl. Logic*, 161(7):879–894, 2010.
- [Jec03] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded.
- [JR95] Haim Judah and Miroslav Repický. No random reals in countable support iterations. *Israel J. Math.*, 92(1-3):349–359, 1995.
- [Kan09] Akihiro Kanamori. *The higher infinite*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, second edition, 2009. Large cardinals in set theory from their beginnings, Paperback reprint of the 2003 edition.
- [Kan19] V. Kanovei. Definable selectors for Δ_2^0 sets modulo countable. Preprint at [arXiv:1510.04193](https://arxiv.org/abs/1510.04193), October 2019.
- [Kec95] A. Kechris. *Classical Descriptive Set Theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag New York, 1995.
- [Kec73] Alexander S. Kechris. Measure and category in effective descriptive set theory. *Ann. Math. Logic*, 5:337–384, 1972/73.
- [KL20] V. G. Kanovei and V. A. Lyubetsky. On the equality relation modulo a countable set. *Mathematical Notes*, 108(4):615–616, 2020.
- [KPW08] Piotr Kalembe, Szymon Plewik, and Anna Wojciechowska. On the ideal (v^0) . *Cent. Eur. J. Math.*, 6(2):218–227, 2008.
- [KRS16] Antti Käenmäki, Tapio Rajala, and Ville Suomala. Local homogeneity and dimensions of measures. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 16(4):1315–1351, 2016.
- [KSZ13] Vladimir Kanovei, Marcin Sabok, and Jindřich Zapletal. *Canonical Ramsey theory on Polish spaces*, volume 202. Cambridge University Press, 2013.
- [LNVTPS11] C. Laflamme, L. Nguyen Van Thé, M. Pouzet, and N. Sauer. Partitions and indivisibility properties of countable dimensional vector spaces. *J. Combin. Theory Ser. A*, 118(1):67–77, 2011.

- [Mat95] Pertti Mattila. Geometry of sets and measures in Euclidean spaces, fractals and rectifiability, volume 44 of *Cambridge Studies in Advanced Mathematics*. *Cambridge University Press, Cambridge*, 7:11, 1995.
- [Mil08] Benjamin D. Miller. The existence of quasi-invariant measures of a given cocycle, I: Atomless, ergodic sigma-finite measures. *Ergodic Theory and Dynamical Systems*, 28(5):1599–1613, 2008.
- [Mos09] Yiannis N. Moschovakis. *Descriptive set theory*, volume 155 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2009.
- [MS] Sandra Müller and Philipp Schlicht. Preprint on generic absoluteness. In preparation.
- [Nie09] André Nies. *Computability and randomness*, volume 51 of *Oxford Logic Guides*. Oxford University Press, Oxford, 2009.
- [PWBW85] Wiesława Poreda, Elżbieta Wagner-Bojakowska, and Władysław Wilczyński. A category analogue of the density topology. *Fundamenta Mathematicae*, 125:167–173, 1985.
- [Sch14] Philipp Schlicht. Thin equivalence relations and inner models. *Ann. Pure Appl. Logic*, 165(10):1577–1625, 2014.
- [Sik48] Roman Sikorski. On the representation of Boolean algebras as fields of sets. *Fund. Math.*, 35:247–258, 1948.
- [SS06] Ralf Schindler and John Steel. The strength of AD. Unpublished note, available at <https://ivv5hpp.uni-muenster.de/u/rds/>, 2006.
- [SS18] Saharon Shelah and Otmar Spinas. Different cofinalities of tree ideals. Preprint, 2018.
- [Szp35] Edward Szpilrajn. Sur une classe de fonctions de M. Sierpiński et la classe correspondante d'ensembles. *Fundamenta Mathematicae*, 24(1):17–34, 1935.
- [Woo82] W. Hugh Woodin. On the consistency strength of projective uniformization. In *Proceedings of the Herbrand symposium (Marseilles, 1981)*, volume 107 of *Stud. Logic Found. Math.*, pages 365–384. North-Holland, Amsterdam, 1982.
- [Zap04] Jindřich Zapletal. Descriptive set theory and definable forcing. *Mem. Amer. Math. Soc.*, 167(793), 2004.
- [Zap08] Jindřich Zapletal. *Forcing idealized*, volume 174 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2008.

SANDRA MÜLLER, INSTITUTE OF MATHEMATICS, UNIVERSITY OF VIENNA, UZA 1—GEBÄUDE 2, AUGASSE 2–6, 1090 WIEN, AUSTRIA

Email address: `mueller.sandra@univie.ac.at`

PHILIPP SCHLICHT, SCHOOL OF MATHEMATICS, UNIVERSITY OF BRISTOL, FRY BUILDING, WOODLAND ROAD, BRISTOL, BS8 1UG, UK

Email address: `philipp.schlicht@bristol.ac.uk`

DAVID SCHRITTESSER, INSTITUTE OF MATHEMATICS, UNIVERSITY OF VIENNA, UZA 1—GEBÄUDE 2, AUGASSE 2–6, 1090 WIEN, AUSTRIA

Email address: `david.schrittesser@univie.ac.at`

THILO WEINERT, DEPARTMENT OF MATHEMATICS NATIONAL UNIVERSITY OF SINGAPORE BLK S17 LEVEL 4 10 LOWER KENT RIDGE ROAD 119076 SINGAPORE

Email address: `thilo.weinert@univie.ac.at`