

THE OPERATOR-VALUED PARALLELISM AND NORM-PARALLELISM IN MATRICES

M. MOHAMMADI GOHARI AND M. AMYARI*

ABSTRACT. Let \mathcal{H} be a Hilbert space, and let $K(\mathcal{H})$ be the C^* -algebra of compact operators on \mathcal{H} . In this paper, we present some characterizations of the norm-parallelism for elements of a Hilbert $K(\mathcal{H})$ -module by employing the Birkhoff–James orthogonality. Among other things, we present a characterization of transitive relation of the norm-parallelism for elements in a certain Hilbert $K(\mathcal{H})$ -module. We also give some characterizations of the Schatten p -norms and the operator norm-parallelism for matrices.

1. INTRODUCTION AND PRELIMINARIES

Throughout the paper, let $(\mathcal{H}, [\cdot, \cdot])$ be a Hilbert space. Let $B(\mathcal{H})$ and $K(\mathcal{H})$ denote the C^* -algebras of all bounded linear operators and compact linear operators on \mathcal{H} , respectively. The numerical range of an operator $T \in B(\mathcal{H})$ is defined by $W(T) = \{[T\xi, \xi] : \xi \in \mathcal{H}, \|\xi\| = 1\}$. Using the Gelfand–Naimark theorem, we identify any C^* -algebra \mathcal{A} as a C^* -subalgebra of $B(\mathcal{H})$ for some Hilbert space \mathcal{H} . An (left) inner product \mathcal{A} -module is a left \mathcal{A} -module \mathcal{E} equipped with an \mathcal{A} -valued inner product $\langle \cdot, \cdot \rangle$, which is linear in the first variable and conjugate linear in the second and satisfies (i) $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0$ if and only if $x = 0$, (ii) $\langle ax, y \rangle = a\langle x, y \rangle$, (iii) $\langle x, y \rangle^* = \langle y, x \rangle$ for all $x, y \in \mathcal{E}$, $a \in \mathcal{A}$. If \mathcal{E} is complete with respect to the norm defined by $\|x\| = \|\langle x, x \rangle\|^{\frac{1}{2}}$, then \mathcal{E} is called a Hilbert \mathcal{A} -module. Obviously, every Hilbert space is a Hilbert \mathbb{C} -module. Further, every C^* -algebra \mathcal{A} can be regarded as a Hilbert C^* -module over itself under the inner product $\langle a, b \rangle = ab^*$. More details on the theory of Hilbert C^* -modules can be found, for example, in [11].

A projection $E \in B(\mathcal{H})$ is minimal if $EB(\mathcal{H})E = \mathbb{C}E$. For $\xi, \eta \in \mathcal{H}$, let $\xi \otimes \eta$ denote the corresponding elementary operator $(\xi \otimes \eta)(\zeta) = [\zeta, \eta]\xi$. It is easy to see that $\|\xi \otimes \eta\| = \|\xi\|\|\eta\|$. If $\xi \in \mathcal{H}$ is a unit vector, then $\xi \otimes \xi$ is a minimal projection. Conversely, any minimal

2010 *Mathematics Subject Classification.* Primary 46L08, Secondary 47A30, 47B10, 47B47.

Key words and phrases. Hilbert C^* -module, orthonormal basis, parallelism, Schatten p -norm.

*Corresponding author.

projection is of this form. The set of all minimal projections in $K(\mathcal{H})$ is denoted by P_m . A system $\{x_\alpha\}_{\alpha \in \Lambda}$ in a Hilbert $K(\mathcal{H})$ -module \mathcal{E} is orthonormal if (i) $\|x_\alpha\| = 1$ for all $\alpha \in \Lambda$, (ii) $\langle x_\alpha, x_\beta \rangle = 0$ for all $\alpha \neq \beta$, (iii) $\langle x_\alpha, x_\alpha \rangle$ is a minimal projection for all $\alpha \in \Lambda$. An orthonormal system is said to be an orthonormal basis for \mathcal{E} if it generates a dense submodule of \mathcal{E} . The orthogonal dimension of \mathcal{E} is defined as the cardinal number of any of its orthonormal bases; see [2, 6, 13].

Let $(X, \|\cdot\|)$ be a complex normed space, and let $x, y \in X$. We say that x is orthogonal to y in the Birkhoff–James sense, denoted by $x \perp_B y$, if

$$\|x\| \leq \|x + \alpha y\| \quad \text{for all } \alpha \in \mathbb{C}.$$

It is easy to see that the Birkhoff–James orthogonality is equivalent to the usual orthogonality in the case when X is an inner product space. For other definitions and more properties of the Birkhoff–James orthogonality in Hilbert C^* -modules; see [1, 3, 4, 5, 7, 8, 9, 15].

Let $x, y \in X$, we say that x is norm-parallel to y , denoted by $x \| y$, if

$$\|x + \lambda y\| = \|x\| + \|y\| \quad \text{for some } \lambda \in \mathbb{T} = \{\alpha \in \mathbb{C} : |\alpha| = 1\}.$$

For more details one can see [14, 18].

Clearly, two elements of a Hilbert space are norm-parallel if they are linearly dependent. In the case of normed linear spaces, two linearly dependent vectors are norm-parallel, but the converse is false in general. Notice that the norm parallelism is symmetric and \mathbb{R} -homogenous, but not transitive (i.e., $x \| y$ and $y \| z$ do not imply $x \| z$ in general; see [19, Example 2.7], unless X is smooth at y ; see [16, Theorem 3.1]). Some characterizations of the norm-parallelism for operators and elements of Hilbert C^* -modules were given in [16, 17, 18, 19].

Some characterizations of the norm-parallelism for elements of a Hilbert $K(\mathcal{H})$ -module are given in Section 2. Let \mathcal{F} be a Hilbert C^* -module, and let $x, y \in \mathcal{F}$. Zamani and Moslehian [19, Theorem 2.3] proved that $x \| y$ if and only if there exist a sequence of unit vectors ξ_n in a Hilbert space \mathcal{H} and $\lambda \in \mathbb{T}$ such that

$$\lim_{n \rightarrow \infty} \operatorname{Re}[\langle x, \lambda y \rangle \xi_n, \xi_n] = \|x\| \|y\|.$$

Suppose that \mathcal{E} is a Hilbert $K(\mathcal{H})$ -module, that $x, y \in \mathcal{E}$, and that $\langle x, x \rangle$ is a minimal projection in P_m . We show that $x \parallel y$ if and only if there exists a unit vector $\xi \in \mathcal{H}$ such that

$$|[\langle x, y \rangle \xi, \xi]| = \|y\|.$$

Bakić and Guljaš [2] proved that every Hilbert $K(\mathcal{H})$ -module \mathcal{E} has an orthonormal basis. Moreover there exists an orthonormal basis $\{x_\alpha\}_{\alpha \in \Lambda}$ for \mathcal{E} such that $\langle x_\alpha, x_\alpha \rangle = E$ for all $\alpha \in \Lambda$ and for some minimal projection $E \in P_m$; see [2, Remark 4]. In the rest of paper, $\{x_\alpha\}_{\alpha \in \Lambda}^\xi$ denotes the orthonormal basis of \mathcal{E} such that $\langle x_\alpha, x_\alpha \rangle = E = \xi \otimes \xi$ for all $\alpha \in \Lambda$ and for some unit vector $\xi \in \mathcal{H}$. We show that there is no element $0 \neq x \in \mathcal{E}$ such that $x \parallel x_\alpha$ for all $\alpha \in \Lambda$.

In Section 3, we use some techniques of [12] to characterize matrix pairs which are norm-parallelism with respect to the Schatten p -norms. Let $\|\cdot\|_\nu$ be an operator norm on $\mathcal{M}_{n \times n}$ induced by the vector norm ν on $\mathcal{M}_{n \times n}$. In [3], Bhatia, and Šemrl conjectured that for $A, B \in \mathcal{M}_{n \times n}$ if $A \perp_B B$ in $\|\cdot\|_\nu$, then $Ay \perp_B By$ for some unit vector $y \in \mathcal{M}_{n \times 1}$ with $\nu(y) = 1$ such that $\nu(Ay) = \|A\|_\nu$. Li and Schneider in [12] proved that above conclusion is not true in general.

We show that if $Ay \parallel By$ in $\nu(\cdot)$ for some unit vector $y \in \mathcal{M}_{n \times 1}$ with $\nu(y) = 1$ such that $\nu(Ay) = \|A\|_\nu$, then $A \parallel B$ in $\|\cdot\|_\nu$, but by [12, Proposition 4.2 and Example 4.3], the converse is not true in general.

2. NORM-PARALLELISM IN HILBERT $\mathbb{K}(\mathcal{H})$ -MODULES

Throughout, \mathcal{E} denotes a Hilbert $K(\mathcal{H})$ -module. We begin with a characterization of the norm-parallelism for elements of \mathcal{E} . Let \mathcal{F} be a Hilbert C^* -module and let $x, y \in \mathcal{F}$. Zamani and Moslehian [19, Theorem 2.3] proved that $x \parallel y$ if and only if there exist a sequence of unit vectors ξ_n in a Hilbert space \mathcal{H} and $\lambda \in \mathbb{T}$ such that

$$\lim_{n \rightarrow \infty} \operatorname{Re}[\langle x, \lambda y \rangle \xi_n, \xi_n] = \|x\| \|y\|.$$

Now suppose that $x, y \in \mathcal{E}$. Following we show that $x \parallel y$ if and only if there exists a unit vector $\xi \in \mathcal{H}$ such that $|[\langle x, x \rangle \langle x, y \rangle \xi, \xi]| = \|x\|^3 \|y\|$.

Theorem 2.1. *Let $x, y \in \mathcal{E}$. Then the following statements are equivalent:*

- (i) $x \parallel y$.

(ii) *There exists a unit vector $\xi \in \mathcal{H}$ such that*

$$|[\langle x, x \rangle \langle x, y \rangle \xi, \xi]| = \|x\|^3 \|y\|.$$

Proof. (i) \Rightarrow (ii): Let $x \parallel y$. We may assume that $x \neq 0$. From [18, Theorem 4.7] $\langle x, x \rangle \parallel \langle x, y \rangle$ and $\|\langle x, y \rangle\| = \|x\| \|y\|$. Theorem 2.4 of [19] states that there exists $\lambda \in \mathbb{T}$ such that

$$\langle x, x \rangle \perp_B \|\langle x, y \rangle\| \langle x, x \rangle + \lambda \|x\|^2 \langle x, y \rangle.$$

By the proof of [17, Theorem 2.10], there exists a unit vector $\xi \in \mathcal{H}$ such that $\|\langle x, x \rangle \xi\| = \|\langle x, x \rangle\|$ and

$$\left[\left(\|\langle x, y \rangle\| \langle x, x \rangle + \lambda \|x\|^2 \langle x, y \rangle \right) \xi, \langle x, x \rangle \xi \right] = 0.$$

Thus $|[\langle x, x \rangle \langle x, y \rangle \xi, \xi]| = \|x\|^2 \|\langle x, y \rangle\| = \|x\|^3 \|y\|$.

(ii) \Rightarrow (i): Suppose that (ii) holds. Then there exists a unit vector $\xi \in \mathcal{H}$ such that

$$\|x\|^3 \|y\| = |[\langle x, x \rangle \langle x, y \rangle \xi, \xi]| \leq \|x\|^2 \|\langle x, y \rangle\| \leq \|x\|^2 \|x\| \|y\|.$$

Hence $\|\langle x, y \rangle\| = \|x\| \|y\|$ and

$$|[\langle x, x \rangle \langle x, y \rangle \xi, \xi]| = \|x\|^2 \|\langle x, y \rangle\| = \|\langle x, x \rangle\| \|\langle x, y \rangle\|.$$

By [17, Theorem 2.10], we have $\langle x, x \rangle \parallel \langle x, y \rangle$. Then [18, Theorem 4.7] implies that $x \parallel y$. \square

Lemma 2.2. [19, Corollary 2.5] *Let $x, y \in \mathcal{E}$. If $x \parallel y$, then there exists $\lambda \in \mathbb{T}$ such that $x \perp_B (\|y\| \langle x, x \rangle x + \lambda \|x\| \langle y, x \rangle x)$ and $y \perp_B (\|x\| \langle y, y \rangle y + \lambda \|y\| \langle y, x \rangle y)$.*

Now, we apply Lemma 2.2 to obtain the following characterization of the norm-parallelism for elements of a Hilbert $K(\mathcal{H})$ -module.

Corollary 2.3. *Let $x, y \in \mathcal{E}$ and $\langle x, x \rangle$ be idempotent. Then the following statements are equivalent:*

(i) $x \parallel y$;

(ii) $x \perp_B (\|y\| \langle x, x \rangle x + \lambda \|x\| \langle y, x \rangle x)$ and $y \perp_B (\|x\| \langle y, y \rangle y + \lambda \|y\| \langle y, x \rangle y)$

for some $\lambda \in \mathbb{T}$.

Proof. (i) \Rightarrow (ii): This implication follows immediately from Lemma 2.2.

(ii) \Rightarrow (i): Suppose that (ii) holds. By [1, Lemma 2.1 part(5)], we have

$$\langle x, x \rangle \perp_B \left(\|y\| \langle x, x \rangle^2 + \bar{\lambda} \|x\| \langle x, x \rangle \langle x, y \rangle \right).$$

By [17, Theorem 2.10], there exists a vector ξ in \mathcal{H} such that

$$\left[\left(\|y\| \langle x, x \rangle^2 + \bar{\lambda} \|x\| \langle x, x \rangle \langle x, y \rangle \right) \xi, \langle x, x \rangle \xi \right] = 0$$

and

$$\|\langle x, x \rangle \xi\| = \|x\|^2.$$

Since $\langle x, x \rangle$ is idempotent, we reach

$$|[\langle x, x \rangle \langle x, y \rangle \xi, \xi]| = \|x\|^3 \|y\|.$$

From (ii) \Rightarrow (i) of Theorem 2.1, we obtain $x\|y$. □

Recall that if x is an element of \mathcal{E} such that $\langle x, x \rangle$ is a minimal projection, then $\langle x, x \rangle x = x$; see [2]. Let $x, y \in \mathcal{E}$. If there exist a unit vector $\xi \in \mathcal{H}$ and $\lambda \in \mathbb{T}$ such that $\operatorname{Re}[\langle x, \lambda y \rangle \xi, \xi] = \|x\| \|y\|$, then [19, Theorem 2.3] implies that $x\|y$. The question is under which conditions the converse is true. When $\langle x, x \rangle$ is a minimal projection, by using Theorem 2.1, we get $x\|y$, if and only if there exists a unit vector $\xi \in \mathcal{H}$ such that $|\langle \langle x, y \rangle \xi, \xi \rangle| = \|y\|$. Now we prove it by a different approach.

Theorem 2.4. *Let $x, y \in \mathcal{E}$. If $\langle x, x \rangle$ is a minimal projection, then $x\|y$ if and only if there exists a unit vector $\xi \in \mathcal{H}$ such that*

$$|\langle \langle x, y \rangle \xi, \xi \rangle| = \|y\|.$$

Proof. Let $x\|y$. By Lemma 2.2, there exists a scalar μ in \mathbb{T} such that

$$x \perp_B \left(\|y\| \langle x, x \rangle x + \mu \|x\| \langle y, x \rangle x \right).$$

Since $\langle x, x \rangle$ is a minimal projection, by [1, Proposition 2.4], we have

$$\langle x, x \rangle \left\langle \left(\|y\| \langle x, x \rangle x + \mu \langle y, x \rangle x \right), x \right\rangle = 0,$$

and so

$$\|y\| \langle x, x \rangle + \mu \langle x, x \rangle \langle y, x \rangle \langle x, x \rangle = 0.$$

Hence

$$\langle x, x \rangle \langle x, y \rangle \langle x, x \rangle = -\bar{\mu} \|y\| \langle x, x \rangle. \quad (2.1)$$

Now, by knowing the fact that $\langle x, x \rangle = \xi \otimes \xi$ for some unit vector $\xi \in \mathcal{H}$, since $\xi \otimes \xi \langle x, y \rangle \xi \otimes \xi = [\langle x, y \rangle \xi, \xi] \xi \otimes \xi$, we obtain

$$\left| [\langle x, y \rangle \xi, \xi] \right| \|\xi \otimes \xi\| = \|y\| \|\xi \otimes \xi\|,$$

whence

$$\left| [\langle x, y \rangle \xi, \xi] \right| = \|y\|.$$

(ii) \Rightarrow (i): Let $\left| [\langle x, y \rangle \xi, \xi] \right| = \|y\|$. Then there exists $\lambda \in \mathbb{T}$ such that

$$\operatorname{Re}[\langle x, \lambda y \rangle \xi, \xi] = \|x\| \|y\|.$$

From [19, Theorem 2.3], we get $x\|y$. □

In the following theorem we use some ideas of [19].

Theorem 2.5. *Let $\{x_\alpha\}_{\alpha \in \Lambda}^\xi$ be an orthonormal basis for \mathcal{E} such that as a Hilbert module is not of dimension one. Then there is no nonzero element $x \in \mathcal{E}$ such that $x\|x_\alpha$ for all $\alpha \in \Lambda$.*

Proof. Let there be a nonzero element $x \in \mathcal{E}$ such that $x\|x_\alpha$ for all $\alpha \in \Lambda$. By [19, Corollary 2.5], there exists a sequence $\{\lambda_\alpha\}$ in \mathbb{T} such that

$$x_\alpha \perp_B \left(\|x\| \langle x_\alpha, x_\alpha \rangle x_\alpha + \lambda_\alpha \langle x_\alpha, x \rangle x_\alpha \right) \quad (\alpha \in \Lambda). \quad (2.2)$$

Taking $E = \langle x_\alpha, x_\alpha \rangle$, since $E x_\alpha = E$, equation (2.1) implies that

$$\langle x_\alpha, x \rangle E = E \langle x_\alpha, x \rangle E = -\overline{\lambda_\alpha} \|x\| E \quad (\alpha \in \Lambda).$$

Hence

$$(\langle x_\alpha, x \rangle E)^* (\langle x_\alpha, x \rangle E) = \lambda_\alpha \overline{\lambda_\alpha} \|x\|^2 E^2 = \|x\|^2 E \quad (2.3)$$

or equivalently

$$E \langle x, x_\alpha \rangle \langle x_\alpha, x \rangle E = \|x\|^2 E \quad (\alpha \in \Lambda).$$

Therefore $[\langle x, x_\alpha \rangle \langle x_\alpha, x \rangle \xi, \xi] = \|x\|^2$ and

$$[\langle x, x \rangle \xi, \xi] = \left[\sum_{\alpha \in \Lambda} \langle x, x_\alpha \rangle \langle x_\alpha, x \rangle \xi, \xi \right] = \sum_{\alpha \in \Lambda} \|x\|^2.$$

Thus

$$\sum_{\alpha \in \Lambda} \|x\|^2 = [\langle x, x \rangle \xi, \xi] = \left| [\langle x, x \rangle \xi, \xi] \right| \leq \|[\langle x, x \rangle \xi, \xi]\| \leq \|x\|^2,$$

which implies that $x = 0$, that is a contradiction. \square

Now we have the following characterization of the transitive relation for elements of a certain Hilbert $K(\mathcal{H})$ -module.

Corollary 2.6. *Let $x, y, z \in \mathcal{E}$ be such that \mathcal{E} as a Hilbert module is of dimension one with orthonormal basis $\{y\}^\xi$. If $x\|y$ and $y\|z$, then $x\|z$.*

Proof. Let $x\|y$ and $y\|z$. By [19, Corollary 2.5], there exist $\lambda, \mu \in \mathbb{T}$ such that $y \perp_B (\|x\| \langle y, y \rangle y + \lambda \|y\| \langle y, x \rangle y)$ and $y \perp_B (\|z\| \langle y, y \rangle y + \mu \|y\| \langle z, y \rangle y)$. Taking $E = \langle y, y \rangle$. Equation (2.1) ensures that $E \langle x, y \rangle = -\bar{\lambda} \|x\| E$ and $\langle y, z \rangle E = -\bar{\mu} \|z\| E$. Then $E \langle x, y \rangle \langle y, z \rangle E = \bar{\lambda} \bar{\mu} \|x\| \|z\| E$. Therefore

$$\left| [\langle x, z \rangle \xi, \xi] \right| = \left| [\langle x, y \rangle \langle y, z \rangle \xi, \xi] \right| = \|x\| \|z\|.$$

That is $x\|z$. \square

Example 2.7. Suppose that \mathcal{H} is an infinite dimensional Hilbert space. Then \mathcal{H} equipped with $\langle x, y \rangle = x \otimes y$ is a Hilbert $K(\mathcal{H})$ -module and $\|x\|^2 = \|x \otimes x\|$ is the original norm on \mathcal{H} . We can assume that each unit vector $y \in \mathcal{H}$ gives an orthonormal basis $\{y\}^y$ for \mathcal{H} . Therefore the dimension of H as a Hilbert module is equal to one. Finally, from Corollary 2.6, we see that if $x\|y$ and $y\|z$, for some unit vector y , then

$$\left| [\langle x, z \rangle y, y] \right| = \|x\| \|z\|.$$

Therefore, $x\|z$.

3. NORM-PARALLELISM IN MATRICES

Let $\mathcal{M}_{n \times n}$ be the space of $n \times n$ complex matrices, let $A, B \in \mathcal{M}_{n \times n}$, and let $\|\cdot\|$ be any norm on $\mathcal{M}_{n \times n}$. We say that A is norm-parallel to B if

$$\|A + \lambda B\| = \|A\| + \|B\|$$

for some $\lambda \in \mathbb{T}$. We also consider the inner product $\langle A, B \rangle = \text{tr}(AB^*)$. The dual norm of $\|\cdot\|$ is defined by

$$\|A\|^D = \max\{|\langle A, B \rangle| : \|B\| \leq 1\}.$$

The singular values of $A \in \mathcal{M}_{n \times n}$, which are the square roots of the eigenvalues of the matrix A^*A , are denoted by $s_1(A) \geq \cdots \geq s_n(A)$ and the Schatten p -norm of A is defined by

$$\|A\|_p = \left\{ \sum_{i=1}^n s_i(A)^p \right\}^{\frac{1}{p}},$$

if $1 \leq p < \infty$ and if $p = \infty$, $\|A\|_p = \max\{s_j(A) : 1 \leq j \leq n\}$; see [12]. If $A \in \mathcal{M}_{n \times n}$, then for a positive integer $1 \leq k \leq n$, the Ky-Fan k -norm is defined by

$$\|A\|_{(k)} = \sum_{j=1}^k s_j(A),$$

where $s_1(A) \geq \cdots \geq s_n(A) \geq 0$, are the singular values of A . The cases when $k = 1$ and $k = n$ correspond to the operator norm $\|\cdot\|_\infty$ and the trace norm $\|\cdot\|_1$, respectively. we have the following result.

Proposition 3.1. *Let $A, B \in \mathcal{M}_{n \times n}$. Then the following statement are equivalent:*

- (i) $A \| B$, in $\|\cdot\|_\infty$.
- (ii) There exist unit vectors $x \in \mathcal{M}_{n \times 1}$ and $y \in \mathcal{M}_{n \times 1}$ such that $\|A\|_\infty = x^* A y$ and $|x^* B y| = \|B\|_\infty$.
- (iii) For any $U \in \mathcal{M}_{n \times n}$ with orthonormal columns that form a basis for the eigenspace of AA^* corresponding to the largest eigenvalue and $V = \frac{1}{\|A\|_\infty} A^* U \in \mathcal{M}_{n \times n}$, it holds that

$$-\|B\|_\infty \in W(\lambda U^* B V)$$

for some $\lambda \in \mathbb{T}$.

Proof. (i) \Rightarrow (ii): Let $A \parallel B$. Thus

$$A \perp_B (\|B\|_\infty A + \lambda \|A\|_\infty B)$$

for some $\lambda \in \mathbb{T}$. By [12, Theorem 3.1], there exist unit vectors $x \in \mathcal{M}_{n \times 1}$ and $y \in \mathcal{M}_{n \times 1}$ such that $\|A\|_\infty = x^* A y$ and $x^* (\|B\|_\infty A + \lambda \|A\|_\infty B) y = 0$. Thus we get $|x^* B y| = \|B\|_\infty$.

(ii) \Rightarrow (i): It is obvious.

(i) \Rightarrow (iii): If $A \parallel B$, then

$$A \perp_B (\|B\|_\infty A + \lambda \|A\|_\infty B)$$

for some $\lambda \in \mathbb{T}$. By [12, Theorem 3.1], for any $U \in \mathcal{M}_{n \times n}$ with orthonormal columns that form a basis for the eigenspace of AA^* corresponding to the largest eigenvalue and $V = \frac{1}{\|A\|_\infty} A^* U \in \mathcal{M}_{n \times n}$, we have

$$\begin{aligned} 0 &\in W(U^* (\|B\|_\infty A + \lambda \|A\|_\infty B) V) = W(\|B\|_\infty \|A\|_\infty V^* V + \lambda \|A\|_\infty U^* B V) \\ &= W(\|B\|_\infty \|A\|_\infty I + \lambda \|A\|_\infty U^* B V). \end{aligned}$$

Thus there exists a unit vector $\zeta \in \mathcal{M}_{n \times 1}$ such that $\langle (\|B\|_\infty I + \lambda \|A\|_\infty U^* B V) \zeta, \zeta \rangle = 0$ or $\langle \lambda U^* B V \zeta, \zeta \rangle = -\frac{\|B\|_\infty}{\|A\|_\infty}$. Hence

$$-\|B\|_\infty \in W(\lambda U^* B V)$$

for some $\lambda \in \mathbb{T}$.

(iii) \Rightarrow (i): It is obvious. □

Remark 3.1. By the equivalence (i) \Leftrightarrow (ii) of Proposition 3.1, $A \parallel B$, in $\|\cdot\|_\infty$ if and only if there exist unit vectors $x \in \mathcal{M}_{m \times 1}$ and $y \in \mathcal{M}_{n \times 1}$ such that $\|A\|_\infty = x^* A y$ and $|x^* B y| = \|B\|_\infty$, this condition is equivalent to the fact that there is a unit vector $y \in \mathcal{M}_{n \times 1}$ such that $\|A\|_\infty = \langle A y, A y \rangle^{\frac{1}{2}}$ and $|\langle A y, B y \rangle| = \|A\|_\infty \|B\|_\infty$.

Remark 3.2. Suppose that $A, B \in \mathcal{M}_{n \times n}$. If $A \parallel B$ in $\|\cdot\|_{(k)}$, then $A \perp_B (\|B\|_{(k)} A + \lambda \|A\|_{(k)} B)$. So, by [10, Remark 1], there exists a matrix $F \in \mathcal{M}_{n \times n}$ such that $\|F\|_\infty \leq 1$, $\|F\|_1 \leq k$, $\text{tr}(F^* A) = \|A\|_{(k)}$ and $\text{tr}\left(F^* (\|B\|_{(k)} A + \lambda \|A\|_{(k)} B)\right) = 0$. Hence $A \parallel B$ in $\|\cdot\|_{(k)}$ if and only if there exists a matrix $F \in \mathcal{M}_{n \times n}$ such that $\|F\|_\infty \leq 1$, $\|F\|_1 \leq k$, $\text{tr}(F^* A) = \|A\|_{(k)}$ and $|\text{tr}(F^* B)| = \|A\|_{(k)}$.

Now by [12, Theorem 3.2], we obtain the following result.

Corollary 3.2. *Let $1 < p < \infty$ and $A, B \in \mathcal{M}_{n \times n}$, where $A = DC$ for some positive matrices $D \in \mathcal{M}_{m \times m}$ and $C \in \mathcal{M}_{m \times n}$ with $CC^* = I_n$. Then $A \parallel B$, in $\|\cdot\|_p$ if and only if $|\operatorname{tr}(D^{p-1}CB^*)| = \frac{\|B\|_p \|D\|_p^p}{\|A\|_p}$.*

Proof. Let $A \parallel B$. Thus

$$A \perp_B (\|B\|_p A + \lambda \|A\|_p B)$$

for some $\lambda \in \mathbb{T}$. By [12, Theorem 3.2], $\operatorname{tr}(D^{p-1}C(\|B\|_p A + \lambda \|A\|_p B)^*) = 0$.

Thus $|\operatorname{tr}(D^{p-1}CB^*)| = \frac{\|B\|_p \|D\|_p^p}{\|A\|_p}$. The converse is obvious. \square

As a consequence of [12, Theorem 3.3], we have the following result.

Remark 3.3. Let $A, B \in \mathcal{M}_{n \times n}$ and $A \parallel B$ in $\|\cdot\|_1$. By a similar computation as previous theorem, we deduce that there exists $F \in \mathcal{M}_{n \times n}$ such that $\|F\|_\infty \leq 1$, $\operatorname{tr}(AF^*) = \|A\|_1$ and $|\operatorname{tr}(BF^*)| = \|B\|_1$.

Let ν be a norm on $\mathcal{M}_{n \times 1}$, and let $\|\cdot\|_\nu$ be the operator norm on $\mathcal{M}_{n \times n}$ induced by ν defined by

$$\|A\|_\nu = \max\{\nu(Ax) : x \in \mathcal{M}_{n \times 1}, \nu(x) \leq 1\}.$$

The dual norm of ν is defined by

$$\nu^D(x) = \max\{|\langle x, y \rangle| : y \in \mathcal{M}_{n \times 1}, \nu(y) \leq 1\}$$

and the dual norm of $\|\cdot\|_\nu$ is defined as

$$\|A\|_\nu^D = \max\{|\langle A, B \rangle| : B \in \mathcal{M}_{n \times n}, \|B\|_\nu \leq 1\}.$$

We say $F \in \mathcal{M}_{n \times n}$ is an extreme point of the unit norm ball of $\mathcal{M}_{n \times n}$ if only if

- (i) $p = 1$ and $F = xy^*$ for some unit vectors $x \in \mathcal{M}_{n \times 1}$ and $y \in \mathcal{N}_{n \times 1}$,
- (ii) $1 < p < \infty$ and $\|F\|_p = 1$,
- (iii) $p = \infty$ and $FF^* = I_n$.

We have the following characterization of the norm parallelism for operator norms in $\mathcal{M}_{n \times n}$.

Proposition 3.3. *Let $\|\cdot\|_\nu$ be an operator norm on $\mathcal{M}_{n \times n}$ induced by the vector norm ν on $\mathcal{M}_{n \times n}$. Let $A, B \in \mathcal{M}_{n \times n}$. Denote by \mathcal{E} and \mathcal{E}^D the set of extreme points of the unit norm balls of ν and ν^D , respectively. If*

$$V(A) = \{xy^* : x \in \mathcal{E}^D, y \in \mathcal{E}, \langle A, xy^* \rangle = \|A\|_\nu\},$$

then $A \parallel B$ in $\|\cdot\|_\nu$ if and only if there exist k extreme points $x_1y_1^, \dots, x_ky_k^*$ with $k \leq 3$ in the complex case and $k \leq 2$ in the real case, and positive numbers t_1, \dots, t_k with $t_1 + \dots + t_k = 1$ such that*

$$|\sum_{j=1}^k t_j \langle B, x_jy_j^* \rangle| = \|B\|_\nu.$$

Proof. Let $A \parallel B$. Then

$$A \perp_B (\|B\|_\nu A + \lambda \|A\|_\nu B)$$

for some $\lambda \in \mathbb{T}$. So by [12, Proposition 4.2], there exist k extreme points $x_1y_1^*, \dots, x_ky_k^*$ with $k \leq 3$ in the complex case and $k \leq 2$ in the real case, and positive numbers t_1, \dots, t_k with $t_1 + \dots + t_k = 1$ such that

$$\sum_{j=1}^k t_j \langle (\|B\|_\nu A + \lambda \|A\|_\nu B), x_jy_j^* \rangle = 0.$$

Thus $|\sum_{j=1}^k t_j \langle B, x_jy_j^* \rangle| = \|B\|_\nu$. The converse is obvious. \square

Remark 3.4. Suppose that ν is a norm on $\mathcal{M}_{n \times 1}$ and that $\|\cdot\|_\nu$ is the corresponding operator norm on $\mathcal{M}_{n \times n}$. Given $A, B \in \mathcal{M}_{n \times n}$. If there exists a vector $y \in \mathcal{M}_{n \times 1}$ with $\nu(y) = 1$ such that $\nu(Ay) = \|A\|_\nu$ and $\nu\left(Ay + \mu(\nu(By)Ay + \lambda\nu(Ay)By)\right) \geq \nu(Ay)$ for all $\mu \in \mathbb{C}$, that is, $Ay \parallel By$. Then

$$\|A + \mu(\|B\|_\nu A + \lambda \|A\|_\nu B)\|_\nu \geq \nu\left(Ay + \mu(\nu(By)Ay + \lambda\nu(Ay)By)\right) \geq \nu(Ay) = \|A\|_\nu$$

for all $\mu \in \mathbb{C}$; hence $A \parallel B$. On the other hand [12, Proposition 4.2 and Example 4.3] show that the converse of the above result is not true, in general.

REFERENCES

- [1] Lj. Arambašić and R. Rajić, *On symmetry of the (strong) Birkhoff-James orthogonality in Hilbert C^* -modules*, Ann. Funct. Anal. **7** (2016), no. 1, 17–23.
- [2] D. Bakić, and B. Guljaš, *Hilbert C^* -modules over C^* -algebras of compact operators*, Acta Sci. Math. (Szeged) **68** (2002), no. 1-2, 249–269.
- [3] R. Bhatia, and P. Šemrl, *Orthogonality of matrices and some distance problems*, Linear Algebra Appl. **287** (1999), no. 1-3, 77–85.
- [4] T. Bhattacharyya, and P. Grover, *Characterization of Birkhoff-James orthogonality*, J. Math. Anal. Appl. **407** (2013), no. 2, 350–358.
- [5] T. Bottazzi, C. Conde, M. S. Moslehian, P. Wójcik and A. Zamani, *Orthogonality and parallelism of operators on various Banach spaces*, J. Aust. Math. Soc. (to appear), arXiv:1609.06615.
- [6] M. Cabrera, J. Martínez, and A. Rodríguez, *Hilbert modules revisited: Orthonormal bases and Hilbert-Schmidt operators*, Glasgow Math. J. **37** (1995), 45–54.
- [7] J. Chmieliński, *Remarks on orthogonality preserving mappings in normed spaces and some stability problems*, Banach J. Math. Anal. **1** (2007), no. 1, 117–124.
- [8] J. Chmieliński, R. Lukasik, and P. Wójcik, *On the stability of the orthogonality equation and the orthogonality-preserving property with two unknown functions*, Banach J. Math. Anal. **10** (2016), no. 4, 828–847.
- [9] P. Ghosh, D. Sain, and K. Paul, *On symmetry of Birkhoff-James orthogonality of linear operators*, Adv. Oper. Theory **2** (2017), no. 4, 428–434.
- [10] P. Grover, *Orthogonality of matrices in the Ky Fan k -norms*, Linear Multilinear Algebra **65** (2017), no. 3, 496–509.
- [11] E. C. Lance, *Hilbert C^* -Modules. A Toolkit for Operator Algebraists*, in: London Mathematical Society Lecture Note Series, vol. **210**, Cambridge University Press, Cambridge, 1995.
- [12] C. K. Li, and H. Schneider, *Orthogonality of matrices*, Linear Algebra App. **81** (2003), no. 2, 175–181.
- [13] L. Molnár, *Modular bases in a Hilbert A -module*, Czechoslovak Math. J. **42** (1992), no. 4, 649–656.
- [14] A. Seddik, *Rank one operators and norm of elementary operators*, Linear Algebra Appl. **424** (2007), no. 1, 177–183.
- [15] A. Turnšek, *On operators preserving James' orthogonality*, Univ. Beograd. Publ. Elektrotehn. Linear Algebra Appl. **407** (2005), 189–195.
- [16] P. Wójcik, *Norm-parallelism in classical M -ideals*, Indag. Math. **28** (2017), no. 2, 287–293.
- [17] A. Zamani, *The operator-valued parallelism*, Linear Algebra Appl. (N.S.). **505** (2016), 282–295.
- [18] A. Zamani, and M. S. Moslehian, *Exact and approximate operator parallelism*, Canad. Math. Bull. **58** (2015), no. 1, 207–224.
- [19] A. Zamani, and M. S. Moslehian, *Norm-parallelism in the geometry of Hilbert C^* -modules*, Indag. Math. (N.S.). **27** (2016), no. 1, 266–281.

DEPARTMENT OF MATHEMATICS, MASHHAD BRANCH, ISLAMIC AZAD UNIVERSITY, MASHHAD, IRAN.

E-mail address: mahdi.mohammadigohari@gmail.com

E-mail address: maryam_amyari@yahoo.com and amyari@mshdiau.ac.ir