

THE STRUCTURE OF NORMAL HAUSDORFF OPERATORS ON LEBESGUE SPACES

A. R. MIROTIN

amirotin@yandex.ru

ABSTRACT. We consider a generalization of Hausdorff operator and introduce the notion of the symbol of such an operator. Using this notion we describe the structure and investigate important properties (such as invertibility, spectrum, norm, and compactness) of normal generalized Hausdorff operators on Lebesgue spaces over \mathbb{R}^n . The examples of Cesàro operators are considered. ¹ ²

INTRODUCTION. Hausdorff operators on Lebesgue spaces originated from some classical summation methods. They were introduced by Hardy [7, Chapter XI] on the unit interval and by C. Georgakis and independently by E. Liflyand and F. Moricz on the whole real line [6], [12]. Its natural multidimensional extension is the operator

$$(\mathcal{H}f)(x) = \int_{\mathbb{R}^m} \Phi(u)f(A(u)x)du, \quad x \in \mathbb{R}^n$$

where $A(u)$ is a family of $n \times n$ matrices satisfying $\det A(u) \neq 0$ almost everywhere in the support of Φ . It was introduced by Brown and Moricz [2] on the Lebesgue space and by Lerner and Liflyand [9] on the real Hardy space. This class of operators has attracted considerable attention in recent decades. It includes some important examples, such as the Hardy operator, the adjoint Hardy operator, the Cesàro operator and their multidimensional analogs. The survey articles [11], [4] contain main results on Hausdorff operators and bibliography up to 2014. As far as the author is aware all known results refer to the boundedness of general Hausdorff operators in various settings only. But this question, being solved positively, naturally entails such questions as whether the operator is invertible and what is its inverse, what is the spectrum of the operator, and a number of others. To deepen the investigations it is natural to begin with Hausdorff operators on the Hilbert space. This work is mainly devoted to this case. We shall consider a generalization of Hausdorff operator and introduce the notion of the symbol of this operator. This notion is crucial for our considerations. Our aim is to describe

¹Key words: Hausdorff operator, Cesàro operator, symbol of an operator, Lebesgue space, normal operator, spectrum, compact operator.

²AMS Mathematics Subject Classification: 47B38, 47B15, 46E30.

the structure and to investigate important properties (such as invertibility, spectrum, norm, and compactness) of generalized normal Hausdorff operators on Lebesgue spaces over \mathbb{R}^n . Using the Mellin transform we prove under some extra condition that a normal Hausdorff operator in $L^2(\mathbb{R}^n)$ is unitary equivalent to the operator of coordinate-wise multiplication by the symbol in the space $L^2(\mathbb{R}^n, \mathbb{C}^{2^n})$. As examples, discrete Hausdorff operators and Cesàro operators are considered. The study of the one-dimensional Cesàro operator on L^2 was pioneered by Brown, Halmos, and Shields [1], the L^p case was considered in [2].

Definition 1 [13]. Let (Ω, μ) be some σ -compact locally compact topological space endowed with positive regular Borel measure μ , Φ a locally integrable function on Ω , and $(A(u))_{u \in \Omega}$ a μ -measurable family of $n \times n$ matrices defined almost everywhere in the support of Φ and satisfying $\det A(u) \neq 0$. We define the *Hausdorff operator* with the kernel Φ by

$$(\mathcal{H}f)(x) = (\mathcal{H}_{\Phi, A}f)(x) = \int_{\Omega} \Phi(u) f(A(u)x) d\mu(u)$$

($x \in \mathbb{R}^n$ is a column vector).

As was mentioned by Hardy in the case $n = 1$, $\Omega = [0, 1]$ [7, Theorem 217] the Hausdorff operator possesses some regularity property. The multi-dimensional version of his result looks as follows.

Proposition 1. *Under the conditions of definition 1 let the matrices $A(u)$ be positive definite for almost all $u \in \Omega$. In order that the transformation $\mathcal{H}_{\Phi, A}$ should be regular, i.e. that $f \in C(\mathbb{R}^n)$, $f(x) \rightarrow l$ when $x \rightarrow \infty$ should imply $\mathcal{H}_{\Phi, A}f(x) \rightarrow l$, it is necessary and sufficient that $\int_{\Omega} \Phi(u) d\mu(u) = 1$.*

Proof. If $f(x) = 1$ then $\mathcal{H}f(x) = \int_{\Omega} \Phi(u) d\mu(u)$. Thus, $\int_{\Omega} \Phi(u) d\mu(u) = 1$ is a necessary condition.

To prove the sufficiency, first note that if $A(u)$ is positive definite then $x \rightarrow \infty$ implies $A(u)x \rightarrow \infty$ (this follows from the representation $A(u) = C^{-1}A'C$ where C is an orthogonal matrix and A' is a diagonal one with positive eigenvalues). But if, in addition, $f \in C(\mathbb{R}^n)$ and $f(x) \rightarrow l$ then f is bounded and therefore $\mathcal{H}f(x) \rightarrow l$ ($x \rightarrow \infty$) by the Lebesgue Theorem.

Proposition 1 motivates our work with positive definite matrices in the sequel.

1. THE STRUCTURE OF NORMAL HAUSDORFF OPERATORS IN L^2 .

We need the following lemmas to prove our main result.

Lemma 1 [13] (cf. [7, (11.18.4)], [2]). *Let $(\det A(u))^{-1/p}\Phi(u) \in L^1(\Omega)$. Then the operator $\mathcal{H}_{\Phi,A}$ is bounded in $L^p(\mathbb{R}^n)$ ($1 \leq p < \infty$) and*

$$\|\mathcal{H}_{\Phi,A}\| \leq \int_{\Omega} |\Phi(u)| |\det A(u)|^{-1/p} d\mu(u).$$

This estimate is sharp (see corollary 2 below).

In the next corollary we as usual consider $L^p(\mathbb{R}_+^n)$ as a subspace of $L^p(\mathbb{R}^n)$.

Corollary 1. *If, in addition, every $A(u)$ maps \mathbb{R}_+^n into itself the operator $\mathcal{H}_{\Phi,A}$ is bounded in $L^p(\mathbb{R}_+^n)$ ($1 \leq p < \infty$).*

Lemma 2 (cf. [2]). *The adjoint for the Hausdorff operator in $L^p(\mathbb{R}^n)$ has the form*

$$(\mathcal{H}^* f)(x) = (\mathcal{H}_{\Phi,A}^* f)(x) = \int_{\Omega} \overline{\Phi(v)} |\det A(v)|^{-1} f(A(v)^{-1}x) d\mu(v).$$

Thus, the adjoint for a Hausdorff operator is also Hausdorff.

Lemma 3. *Under the conditions of definition 1 the Hausdorff operator $\mathcal{H}_{\Phi,A}$ is normal in $L^2(\mathbb{R}^n)$ if $n \times n$ -matrices $A(u)$ (defined μ -a. e. on the support of Φ) form a pairwise commuting family.*

Proof. Indeed, this follows from the equalities

$$(\mathcal{H}\mathcal{H}^* f)(x) = \int_{\Omega} \int_{\Omega} \Phi(u) \overline{\Phi(v)} |\det A(v)|^{-1} f(A(u)A(v)^{-1}x) d\mu(v) d\mu(u),$$

$$(\mathcal{H}^*\mathcal{H} f)(x) = \int_{\Omega} \int_{\Omega} \Phi(u) \overline{\Phi(v)} |\det A(v)|^{-1} f(A(v)^{-1}A(u)x) d\mu(u) d\mu(v)$$

and the Fubini Theorem.

In the sequel for $x \in \mathbb{R}^n$, $\alpha \in \mathbb{C}^n$, and $r \in \mathbb{R}$ we put $dx = dx_1 \dots dx_n$, $|x|^{r+\alpha} := |x_1|^{r+\alpha_1} \dots |x_n|^{r+\alpha_n}$.

The following notion is crucial for our investigations.

Definition 2. Let the conditions of definition 1 are fulfilled. Let $a(u) := (a_1(u), \dots, a_n(u))$ be the family of eigenvalues (with their multiplicities) of the matrix $A(u)$ and $|\det A(u)|^{-1/p}\Phi(u) \in L^1(\Omega)$. Then we call the function ($s \in \mathbb{R}^n$)

$$\varphi(s) := \int_{\Omega} \Phi(u) (\det A(u))^{-1/p} a(u)^{-is} d\mu(u)$$

the symbol of the Hausdorff operator $\mathcal{H}_{\Phi,A}$ in $L^p(\mathbb{R}^n)$ ($1 \leq p < \infty$).

In the previous definition we assume that for every $j = 1, \dots, n$ a μ -measurable branch of $\log a_j(u)$ in Ω exists and is fixed, and $a(u)^z := \prod_{j=1}^n a_j(u)^{z_j}$ where $a_j(u)^{z_j} := \exp(z_j \log a_j(u))$ ($z \in \mathbb{C}^n$). (Otherwise, φ should be considered as a multi-valued function.)

It is clear that φ is bounded and continuous on \mathbb{R}^n .

Note that if $A(u)$ is a pairwise commuting family of real selfadjoint $n \times n$ -matrices, there is such an orthogonal $n \times n$ -matrix C and a family of real diagonal matrices $A'(u) = \text{diag}[a_1(u), \dots, a_n(u)]$, that $A'(u) = C^{-1}A(u)C$ for almost all $u \in \Omega$. If each $A(u)$ is positive definite the corresponding Hausdorff operator $\mathcal{H}_{\Phi,A'}$ in $L^p(\mathbb{R}^n)$ is bounded provided $(\det A(u))^{-1/p}\Phi(u) = a(u)^{-1/p}\Phi(u) \in L^1(\Omega)$.

Example 1 (cf. [2]). Consider the n -dimensional Cesàro operator:

$$(\mathcal{C}_n f)(x_1, \dots, x_n) = \frac{1}{x_1 \dots x_n} \int_0^{x_1} \dots \int_0^{x_n} f(t_1, \dots, t_n) dt_1 \dots dt_n.$$

This is a bounded Hausdorff operator in $L^p(\mathbb{R}^n)$ (and in $L^p(\mathbb{R}_+^n)$) for $1 < p < \infty$ where $\Omega = [0, 1]^n$ is endowed with the Lebesgue measure, $\Phi = 1$, and $A(u) = \text{diag}[u_1, \dots, u_n]$. Its symbol is $(s = (s_j) \in \mathbb{R}^n)$

$$\varphi(s) = \int_0^1 \dots \int_0^1 \prod_{j=1}^n u_j^{-\frac{1}{2} - is_j} du_1 \dots du_n = \prod_{j=1}^n (1/2 - is_j)^{-1}.$$

Example 2. (Discrete Hausdorff operators.) Let $\Omega = \mathbb{Z}_+$, and μ be a counting measure. Then the definition 1 takes the form ($f \in L^p(\mathbb{R}^n)$)

$$(\mathcal{H}_{\Phi,A} f)(x) = \sum_{k=0}^{\infty} \Phi(k) f(A(k)x).$$

Assume that $\sum_{k=0}^{\infty} |\Phi(k)| |\det A(k)|^{-1/p} < \infty$. Then $\mathcal{H}_{\Phi,A}$ is bounded on $L^p(\mathbb{R}^n)$ and if $a_j(k) > 0$ for all j and k the symbol of $\mathcal{H}_{\Phi,A}$ is

$$\varphi(s) = \sum_{k=0}^{\infty} \Phi(k) (\det A(k))^{-1/p} a(k)^{-is}$$

where $s = (s_j) \in \mathbb{R}^n$ and the principal values of the exponents are considered. Since this series converges on \mathbb{R}^n absolutely (and therefore uniformly), φ is uniformly almost periodic.

The next theorem describes under some additional assumptions the structure of normal Hausdorff operator in $L^2(\mathbb{R}^n)$.

Theorem 1. *Let $A(u)$ be a pairwise commuting family of real positive definite $n \times n$ -matrices (defined μ -a. e. on the support of Φ), and $(\det A(u))^{-1/2}\Phi(u) \in L^1(\Omega)$. Then the Hausdorff operator $\mathcal{H}_{\Phi,A}$ in $L^2(\mathbb{R}^n)$ with symbol φ is normal and unitary equivalent to the operator M_φ of coordinate-wise multiplication by φ in the space $L^2(\mathbb{R}^n, \mathbb{C}^{2^n})$ of \mathbb{C}^{2^n} -valued functions. More precisely, $\mathcal{H}_{\Phi,A} = \mathcal{V}^{-1}M_\varphi\mathcal{V}$ where \mathcal{V} is a unitary operator from $L^2(\mathbb{R}^n)$ to $L^2(\mathbb{R}^n, \mathbb{C}^{2^n})$ which does not depend on Φ .*

Proof. Since matrices $A(u)$ pairwise commutes and are positive definite, there are an orthogonal $n \times n$ -matrix C and a family of diagonal positive definite matrices $A'(u)$ such that $A'(u) = C^{-1}A(u)C$ for all $u \in \Omega$. If $A'(u) = \text{diag}[a_1(u), \dots, a_n(u)]$, then all functions $a_j(u)$ are μ -measurable and positive and $\det A(u) = a_1(u) \dots a_n(u)$. Consider the unitary operator $\hat{C}f(x) := f(Cx)$ in $L^2(\mathbb{R}^n)$. It is easy to verify that

$$\mathcal{H}_{\Phi,A} = \hat{C}^{-1}\mathcal{H}_{\Phi,A'}\hat{C}.$$

Thus, the operator $\mathcal{H}_{\Phi,A}$ is unitary equivalent to $\mathcal{H}_{\Phi,A'}$.

Let U_1, \dots, U_{2^n} be open hyperoctants in \mathbb{R}^n . Then $L^2(\mathbb{R}^n) = \bigoplus_{j=1}^{2^n} L^2(U_j)$, the orthogonal sum of subspaces, and each $L^2(U_j)$ is $\mathcal{H}_{\Phi,A'}$ -invariant. For every j consider the operator \mathcal{H}'_j on $L^2(\mathbb{R}^n)$ which on $L^2(U_j)$ coincides with the restriction of $\mathcal{H}_{\Phi,A'}$ to this space and equals to zero on the orthogonal complement of $L^2(U_j)$. Then

$$\mathcal{H}_{\Phi,A'} = \bigoplus_{j=1}^{2^n} \mathcal{H}'_j,$$

the orthogonal sum of operators.

Consider the modified n -dimensional Mellin transform for the n -hyperoctant U_j in the form ($j = 1, \dots, 2^n$; $s \in \mathbb{R}^n$)

$$(\mathcal{M}_j f)(s) := \frac{1}{(2\pi)^{n/2}} \int_{U_j} |x|^{-\frac{1}{2}+is} f(x) dx.$$

Then \mathcal{M}_j is a unitary operator from $L^2(U_j)$ to $L^2(\mathbb{R}^n)$. This can be easily obtained from the Plancherel theorem for the n -dimensional Fourier transform by using an exponential change of variables (see [3]). Moreover, if we

assume that $|y|^{-1/2}f(y) \in L^1(U)$ then making use of Fubini's theorem, and integrating by substitution $x = A'(u)^{-1}y$ yield the following ($s \in \mathbb{R}^n$):

$$\begin{aligned} (\mathcal{M}_j \mathcal{H}'_j f)(s) &= \frac{1}{(2\pi)^{n/2}} \int_{U_j} |x|^{-\frac{1}{2}+is} dx \int_{\Omega} \Phi(u) f(A'(u)x) d\mu(u) = \\ &= \frac{1}{(2\pi)^{n/2}} \int_{\Omega} \Phi(u) d\mu(u) \int_{U_j} |x|^{-\frac{1}{2}+is} f(A'(u)x) dx = \\ &= \int_{\Omega} \Phi(u) a(u)^{-\frac{1}{p}-is} d\mu(u) \frac{1}{(2\pi)^{n/2}} \int_U |y|^{-\frac{1}{2}+is} f(y) dy = \varphi(s) (\mathcal{M}_j f)(s). \end{aligned}$$

By continuity we get for all $f \in L^2(U_j)$ that

$$\mathcal{M}_j \mathcal{H}'_j f = \varphi \mathcal{M}_j f.$$

So, \mathcal{H}'_j is unitary equivalent to the operator M'_φ in $L^2(\mathbb{R}^n)$ of multiplication by φ and

$$\mathcal{H}_{\Phi, A'} = \bigoplus_{j=1}^{2^n} \mathcal{M}_j^{-1} M'_\varphi \mathcal{M}_j.$$

If we define the operator \mathcal{V}' from $L^2(\mathbb{R}^n)$ to the space $L^2(\mathbb{R}^n) \oplus \dots \oplus L^2(\mathbb{R}^n)$ (2^n direct summands) which is isomorphic to the Hilbert space $L^2(\mathbb{R}^n, \mathbb{C}^{2^n})$ by the equalities $\mathcal{V}'|_{L^2(U_j)} := \mathcal{M}_j$ ($j = 1, \dots, 2^n$) then \mathcal{V}' is unitary and it is easy to verify that

$$\mathcal{V}'^{-1} M_\varphi \mathcal{V}' = \bigoplus_{j=1}^{2^n} \mathcal{M}_j^{-1} M'_\varphi \mathcal{M}_j.$$

Thus, $\mathcal{H}_{\Phi, A'} = \mathcal{V}'^{-1} M'_\varphi \mathcal{V}'$ and the proof is complete.

The following corollary shows that properties of a Hausdorff operator are closely related to the properties of its symbol.

Corollary 2. (i) *The operator $\mathcal{H}_{\Phi, A}$ is invertible if and only if $\inf |\varphi| > 0$; in this case the inverse $\mathcal{H}_{\Phi, A}^{-1}$ is unitary equivalent to the operator $M_{1/\varphi}$ in $L^2(\mathbb{R}^n, \mathbb{C}^{2^n})$;*

(ii) *Let $\mathcal{H}_{\Phi, A}$ and $\mathcal{H}_{\Psi, B}$ be two Hausdorff operators over the same measure space (Ω, μ) such that $(A(u), B(v))$ is a pairwise commuting family of real positive definite $n \times n$ -matrices (u and v run over the support of Φ and Ψ respectively), and $(\det A(u))^{-1/2} \Phi(u), (\det B(v))^{-1/2} \Psi(v) \in L^1(\Omega)$. Then the product $\mathcal{H}_{\Phi, A} \mathcal{H}_{\Psi, B}$ is unitary equivalent to the operator $M_{\varphi\psi}$ in $L^2(\mathbb{R}^n, \mathbb{C}^{2^n})$ (ψ denotes the symbol of $\mathcal{H}_{\Psi, B}$).*

- (iii) The spectrum $\sigma(\mathcal{H}_{\Phi,A})$, the point spectrum $\sigma_p(\mathcal{H}_{\Phi,A})$, and the continuous spectrum of $\mathcal{H}_{\Phi,A}$ equal to the spectrum (i. e. to the closure of the range of the symbol φ), to the point spectrum, and to the continuous spectrum of M'_φ in $L^2(\mathbb{R}^n)$ respectively, the residual spectrum $\sigma_r(\mathcal{H}_{\Phi,A})$ is empty;
- (iv) The symbol of the adjoint $\mathcal{H}_{\Phi,A}^*$ is $\overline{\varphi}$.
- (v) $\|\mathcal{H}_{\Phi,A}\| = \sup |\varphi|$. If, in addition, $\Phi(u) \geq 0$ for μ -almost all u then

$$\|\mathcal{H}_{\Phi,A}\| = \int_{\Omega} \Phi(u)(\det A(u))^{-1/2} d\mu(u).$$

Proof. (i) Evidently the operator $\mathcal{H}_{\Phi,A}$ is invertible if and only if M_φ is invertible, i. e. if $\inf |\varphi| > 0$, and in this case $\mathcal{H}_{\Phi,A}^{-1}$ is unitary equivalent to $M_{1/\varphi}$.

To prove (ii) first note that the orthogonal matrix C exists such that both $A'(u) = C^{-1}A(u)C$ and $B'(v) := C^{-1}B(v)C$ are diagonal. Then the proof of theorem 1 shows that $\mathcal{V}\mathcal{H}_{\Phi,A}\mathcal{V}^{-1} = M_\varphi$, $\mathcal{V}\mathcal{H}_{\Psi,B}\mathcal{V}^{-1} = M_\psi$ for some unitary operator \mathcal{V} from $L^2(\mathbb{R}^n)$ to $L^2(\mathbb{R}^n, \mathbb{C}^{2^n})$ and (ii) follows.

(iii) For $\lambda \in \mathbb{C}$ we have

$$M_\varphi - \lambda = \bigoplus_1^{2^n} (M'_\varphi - \lambda) := (M'_\varphi - \lambda) \oplus \cdots \oplus (M'_\varphi - \lambda).$$

Since a finite orthogonal sum of operators is invertible if and only if each summand is invertible (see, e.g., [8, p. 439]), $\sigma(\mathcal{H}) = \sigma(M_\varphi) = \sigma(M'_\varphi)$.

Moreover,

$$\sigma_p(\mathcal{H}) = \sigma_p(M_\varphi) = \sigma_p\left(\bigoplus_1^{2^n} M'_\varphi\right) = \bigcup_1^{2^n} \sigma_p(M'_\varphi) = \sigma_p(M'_\varphi).$$

Next, by theorem 1 $\sigma_r(\mathcal{H}) = \sigma_r(M_\varphi)$. Let $\lambda \notin \sigma_p(M_\varphi)$. Since $\sigma_r(M'_\varphi) = \emptyset$, we get (in the following $\text{cl}(S)$ denotes the closure of the subset $S \subset L^2(\mathbb{R}^n)$ and $\text{Im}(T)$ denotes the image of the operator T)

$$\text{cl}(\text{Im}(M_\varphi - \lambda)) = \bigoplus_1^{2^n} \text{cl}(\text{Im}(M'_\varphi - \lambda)) = \bigoplus_1^{2^n} L^2(\mathbb{R}^n) = L^2(\mathbb{R}^n, \mathbb{C}^{2^n}).$$

So, $\sigma_r(M_\varphi) = \emptyset$, and (iii) follows.

(iv) Since the adjoint of $\mathcal{H}_{\Phi,A}$ is also of Hausdorff type (with $A(u)^{-1}$ instead of $A(u)$), the this statement follows from the definition 2 and lemma 2.

Finally, the equality $\|\mathcal{H}_{\Phi,A}\| = \sup |\varphi|$ follows from (iii) and the normality of $\mathcal{H}_{\Phi,A}$. If, in addition, $\Phi(u) \geq 0$ for μ -almost all u we have $\sup |\varphi| = \varphi(0)$ which completes the proof.

Corollary 3. *Under the assumptions of theorem 1 the Hausdorff operator $\mathcal{H}_{\Phi,A}$ is self-adjoint (positive, unitary) in $L^2(\mathbb{R}^n)$ if and only if its symbol φ is real-valued (respectively, nonnegative, $|\varphi| = 1$).*

Proof. Indeed, the spectral decomposition of a normal operator implies that the operator $\mathcal{H}_{\Phi,A}$ is self-adjoint (positive, unitary) if and only if $\sigma(\mathcal{H}_{\Phi,A}) = \text{cl}(\varphi(\mathbb{R}^n))$ is contained in \mathbb{R} (respectively, in \mathbb{R}_+ , \mathbb{T}).

Corollary 4. *Under the assumptions and notation of theorem 1 (ii) the operator $\mathcal{H}_{\Psi,B}$ is the inverse of $\mathcal{H}_{\Phi,A}$ if and only if $\varphi\psi = 1$.*

Corollary 5. *Under the assumptions of theorem 1 the Hausdorff operator $\mathcal{H}_{\Phi,A}$ in $L^2(\mathbb{R}_+^n)$ is unitary equivalent to the operator of multiplication by φ in $L^2(\mathbb{R}_+^n)$.*

Proof. This was established in the process of proving theorem 1.

Example 3. Consider the Cesàro operator in $L^2(\mathbb{R}^n)$ (see example 1). It is normal and its symbol is $\varphi(s) = \prod_{j=1}^n (1/2 - is_j)^{-1}$.

The statement (iv) of corollary 2 implies $\|\mathcal{C}_n\| = 2^n$ (and by the corollary 5 the same is true in $L^2(\mathbb{R}_+^n)$).

According to the statement (iii) of corollary 2 $\sigma_r(\mathcal{C}_n) = \emptyset$, $\sigma_p(\mathcal{C}_n) = \emptyset$, and $\sigma(\mathcal{C}_n)$ equals to the closure of the range of φ . Consider the circle $S := \{1/(1/2 - it) : t \in \mathbb{R}\} = \{z \in \mathbb{C} : |z - 1| = 1\}$. Then is $\sigma(\mathcal{C}_n) = \{z_1 \dots z_n : z_j \in S, j = 1, \dots, n\}$. Let $z_j \in S$, $z_j = 1 + e^{i\theta_j}$ ($\theta_j \in [-\pi, \pi]$). Then $|z_j| = 2 \cos(\theta_j/2)$, and $\arg(z_j) = \theta_j/2$ ($j = 1, \dots, n$). Let $z = z_1 \dots z_n$, $r := |z|$, $\theta := \arg(z)$. Then $\theta = (\theta_1 + \dots + \theta_n)/2$, and $r = 2^n \prod_{j=1}^n \cos(\theta_j/2)$. Using the identity

$$\prod_{j=1}^n \cos(\alpha_j) = \frac{1}{2^{n-1}} \sum_{\varepsilon \in \{-1,1\}^n} \cos(\varepsilon \cdot \alpha) = \frac{1}{2^{n-1}} \sum_{\varepsilon' \in \{-1,1\}^{n-1}} \cos(\alpha_1 + \varepsilon' \cdot \alpha'),$$

where $\alpha = (\alpha_j)_{j=1}^n = (\theta_j/2)_{j=1}^n$, $\alpha' = (\theta_j/2)_{j=2}^n$, and \cdot denotes the dot product we conclude that

$$\sigma(\mathcal{C}_n) = \{re^{i\theta} : r \leq 2(2^{n-1} - 1 + \cos \theta), \theta \in [-\pi, \pi]\}.$$

By the corollary 5 the same is true for the Cesàro operator in $L^2(\mathbb{R}_+^n)$.

In particular, for the bivariate Cesàro operator we have

$$\sigma(\mathcal{C}_2) = \{re^{i\theta} : r \leq 2(1 + \cos \theta), \theta \in [-\pi, \pi]\},$$

the region in \mathbb{C} bounded by cardioid (cf. [15, p. 95]).

Results of the example 3 are consistent with classical results from [1] where the case $n = 1$ and the space $L^2(\mathbb{R}_+)$ were considered (see p. 137 therein; in [1] the Cesàro operator in $L^2(\mathbb{R}_+)$ was denoted by \mathcal{C}_∞).

Example 4. Consider the (C, k) mean of a function $f \in L^2(\mathbb{R})$ (see, e.g., [7, p. 276])

$$(C, k)f(x) = \frac{k}{x^k} \int_0^x (x-t)^{k-1} f(t) dt \quad (k > 0).$$

This is an operator of Hausdorff type where $n = 1$, $\Omega = [0, 1]$ endowed with the Lebesgue measure, $\Phi(u) = k(1-u)^{k-1}$, and $A(u) = u$. Its symbol is

$$\varphi(s) = \int_0^1 k(1-u)^{k-1} u^{-1/2-is} du = \frac{\Gamma(k+1)\Gamma(1/2-is)}{\Gamma(k+1/2-is)}.$$

Using [17, Section 12.13, Example 1] we can rewrite this as follows

$$\varphi(s) = \prod_{l=1}^{\infty} \frac{l(k+l-1/2-is)}{(k+l)(l-1/2-is)}.$$

So, the spectrum $\sigma((C, k))$ is the closure of the curve with complex equation

$$z = \prod_{l=1}^{\infty} \frac{l(k+l-1/2-is)}{(k+l)(l-1/2-is)}, \quad s \in \mathbb{R}.$$

Moreover, the last representation of φ implies also that $\max_{\mathbb{R}} |\varphi| = \varphi(0)$. It follows in view of corollary 2 that

$$\|(C, k)\| = \sqrt{\pi} \frac{\Gamma(k+1)}{\Gamma(k+1/2)}$$

(since $\Phi \geq 0$, this follows also from the last equality in corollary 2 (v)).

If, in addition, $k \in \mathbb{N}$, we have $\|(C, k)\| = k!2^k/1 \cdot 3 \dots (2k-1)$ and

$$\varphi(s) = k! \prod_{j=0}^{k-1} (j+1/2-is)^{-1}.$$

Example 5. Let $\mathcal{H}_{\Phi, A}$ and φ be as in the example 2. If $p = 2$ corollary 2 implies that $\sigma(\mathcal{H}_{\Phi, A}) = \text{cl}(\varphi(\mathbb{R}^n))$. Assume, in addition, that $A(k) = A^k$ where $A \neq E$ be an $n \times n$ -matrix with positive eigenvalues $\lambda_1, \dots, \lambda_n$ (taking into account their multiplicities). Consider the function ($z \in \mathbb{D} := \{|z| \leq 1\}$)

$$F(z) := \sum_{k=0}^{\infty} \Phi(k) (\det A)^{-k/2} z^k.$$

Then F belongs to the commutative Banach algebra \mathcal{A}^+ of functions on $\overline{\mathbb{D}}$ with absolutely convergent Taylor series and $\varphi(s) = F(\exp(-i \sum_j s_j \log \lambda_j))$. Now theorem 1 implies that $\sigma(\mathcal{H}_{\Phi,A}) = F(\mathbb{T})$ where $\mathbb{T} = \{|z| = 1\}$. It follows also that $\|\mathcal{H}_{\Phi,A}\| = \sup_{\mathbb{T}} |F|$, and $\mathcal{H}_{\Phi,A}$ is invertible if and only if $\inf_{\mathbb{T}} |F| > 0$. Since the last condition is equivalent to $\inf_{\overline{\mathbb{D}}} |F| > 0$, under this condition the function $G := 1/F$ belongs to the algebra \mathcal{A}^+ , as well (it follows from the Gelfand theory). Let $G(z) = \sum_{k=0}^{\infty} b(k)z^k$ be the Taylor expansion for G ($z \in \overline{\mathbb{D}}$). Then the inverse of $\mathcal{H}_{\Phi,A}$ is the Hausdorff operator

$$\mathcal{H}_{\Phi,A}^{-1}f(x) := \sum_{k=0}^{\infty} b(k)(\det A)^{k/2}f(A^k x).$$

Indeed, in this case all the conditions of corollary 2 (ii) for the pair $\mathcal{H}_{\Phi,A}$, $\mathcal{H}_{\Phi,A}^{-1}$ are fulfilled and the symbol of the last operator is

$$\psi(s) = \sum_{k=0}^{\infty} b(k)a(k)^{-is} =$$

$$G(\exp(-i \sum_j s_j \log \lambda_j)) = \frac{1}{F(\exp(-i \sum_j s_j \log \lambda_j))} = \frac{1}{\varphi(s)}.$$

Thus, the result follows from corollary 4.

2. THE NONCOMPACTNESS OF HAUSDORFF OPERATORS IN L^p . The problem of compactness of (nontrivial) Hausdorff operators was posed by Liflyand in [10] (see also [11]). Below we shall show that such an operator is noncompact in $L^p(\mathbb{R}^n)$ ($1 \leq p < \infty$) if $(A(u))_{u \in \Omega}$ is a pairwise commuting family of positive definite matrices. The case $p = 2$ was considered in [13].

Theorem 2. *Let $A(u)$ be a pairwise commuting family of positive definite $n \times n$ -matrices (u runs over the support of Φ) and $(\det A(u))^{-1/p}\Phi(u) \in L^1(\Omega)$. Then the Hausdorff operator $\mathcal{H}_{\Phi,A}$ is noncompact in $L^p(\mathbb{R}^n)$ ($1 \leq p < \infty$) provided it is non-zero.*

Proof. Assume the contrary. As in the proof of theorem 1 consider the (bounded and invertible) operator $\hat{C}f(x) := f(Cx)$ in $L^p(\mathbb{R}^n)$. Since $\hat{C}\mathcal{H}_{\Phi,A}\hat{C}^{-1} = \mathcal{H}_{\Phi,A'}$, the operator $\mathcal{H} := \mathcal{H}_{\Phi,A'}$ is non-zero and compact, too. Note that every open n -hyperoctant in \mathbb{R}^n is \mathcal{H} -invariant. There is an open n -hyperoctant U such that the restriction $\mathcal{K} := \mathcal{H}|_{L^p(U)}$ is non-zero. Then \mathcal{K} is compact as well.

We shall use the modified n -dimensional Mellin transform for the n -hyperoctant U in the form ($s \in \mathbb{R}^n$)

$$(\mathcal{M}f)(s) := \frac{1}{(2\pi)^{n/2}} \int_U |x|^{-\frac{1}{q}+is} f(x) dx$$

to get a contradiction. The map \mathcal{M} is a bounded operator from $L^p(U)$ to $L^q(\mathbb{R}^n)$ if $1 \leq p \leq 2$ ($1/p + 1/q = 1$) and a unitary operator from $L^2(U)$ to $L^2(\mathbb{R}^n)$. This can be easily obtained from the Hausdorff–Young inequality (and from the Plancherel theorem if $p = 2$) for the n -dimensional Fourier transform by using an exponential change of variables (see [3]). Let $f \in L^p(U)$. First assume that $|y|^{-1/q}f(y) \in L^1(U)$. Then as in the proof of theorem 1 making use of Fubini’s theorem, and integrating by substitution $x = A(u)^{-1}y$ yield the following ($s \in \mathbb{R}^n$):

$$\begin{aligned} (\mathcal{MK}f)(s) &= \frac{1}{(2\pi)^{n/2}} \int_U |x|^{-\frac{1}{q}+is} dx \int_{\Omega} \Phi(u) f(A(u)x) d\mu(u) = \\ &= \frac{1}{(2\pi)^{n/2}} \int_{\Omega} \Phi(u) d\mu(u) \int_U |x|^{-\frac{1}{q}+is} f(A(u)x) dx = \\ &= \int_{\Omega} \Phi(u) a(u)^{-\frac{1}{p}-is} d\mu(u) \frac{1}{(2\pi)^{n/2}} \int_U |y|^{-\frac{1}{q}+is} f(y) dy = \varphi(s)(\mathcal{M}f)(s), \end{aligned}$$

where the symbol φ is continuous and bounded on \mathbb{R}^n . Thus, if $|y|^{-1/q}f(y) \in L^1(U)$ then

$$\mathcal{MK}f = \varphi \mathcal{M}f. \quad (1)$$

By continuity the last equality is valid for all $f \in L^p(U)$.

If $p = 2$ it follows that \mathcal{K} is unitary equivalent to the operator M_{φ} of multiplication by φ in $L^2(\mathbb{R}^n)$. Since \mathcal{K} is non-zero, the continuous function φ is nonzero, too. So, there is such a constant $c > 0$ that the set $\{s \in U : |\varphi(s)| > c\}$ contains some open ball S . Then the restriction of M_{φ} to the space $L^2(S)$ is an invertible and compact operator, which is contrary to the fact that $L^2(S)$ is infinite dimensional.

Now let $1 \leq p < 2$. It follows from (1) then $M_{\psi} P_{L^q(S)} \mathcal{MK} = P_{L^q(S)} \mathcal{M}$ where $\psi = (1/\varphi)|_S$ and $P_{L^q(S)} f := f|_{L^q(S)}$ a projection. Let $T = P_{L^q(S)} \mathcal{M}$. Passing to conjugates gives $\mathcal{K}^* T^* M_{\psi}^* = T^*$. By the theorem 1 from [5] this implies that the operator $T^* = \mathcal{M}^* P_{L^q(S)}^*$ has finite rank. Note that the conjugate $P_{L^q(S)}^*$ is the operator of natural embedding $L^p(S) \subset L^p(\mathbb{R}^n)$. We clame also that for $g \in L^p(\mathbb{R}^n), x \in U$

$$(\mathcal{M}^*g)(x) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} |x|^{-\frac{1}{q}+is} g(s) ds.$$

First note that the right-hand side here is a bounded operator from $L^p(\mathbb{R}^n)$ to $L^q(U)$. As above this can be obtained from the Hausdorff–Young inequality by using an exponential change of variables. Then (and by lemma 1) both sides of the equality ($f \in L^p(U), g \in L^q(\mathbb{R}^n)$)

$$\int_{\mathbb{R}^n} (\mathcal{M}f)(s) g(s) ds = \int_U f(x) (\mathcal{M}^*g)(x) dx$$

that we want to prove exist as Lebesgue integrals. So we may apply Fubini's theorem as follows:

$$\begin{aligned} \int_U f(x)(\mathcal{M}^*g)(x)dx &= \int_U f(x)\frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} |x|^{-\frac{1}{q}+is} g(s)dsdx = \\ &= \int_{\mathbb{R}^n} \frac{1}{(2\pi)^{n/2}} \int_U f(x)|x|^{-\frac{1}{q}+is} dxg(s)ds = \int_{\mathbb{R}^n} (\mathcal{M}f)(s)g(s)ds. \end{aligned}$$

It was shown above that the restriction of the operator \mathcal{M}^* to $L^p(S)$ has finite rank. Since \mathcal{M}^* can be easily reduced to the Fourier transform, this contradicts to the Paley-Wiener Theorem about the image of $L^2(S)$ ($L^2(S) \subset L^p(S)$) under the Fourier transform (see, e. g., [16, Theorem III.4.9]).

Finally, if $2 < p < \infty$ one can use duality arguments. Indeed, by lemma 2 the adjoint operator $\mathcal{H}^* = \mathcal{H}_{\Phi, A'}^*$ (as an operator in $L^q(\mathbb{R}^n)$) is also of Hausdorff type. More precisely, it equals to $\mathcal{H}_{\Psi, B}$, where $\Psi(u) = \Phi(u)|\det A(u)'^{-1}| = \Phi(u)/a(u)$, $B(u) = A(u)'^{-1} = \text{diag}[1/a_1(u), \dots, 1/a_n(u)]$. It is easy to verify that $\mathcal{H}_{\Psi, B}$ satisfies all the conditions of theorem 2 (with q , Ψ , and B in place of p , Φ , and A respectively). Since $1 < q < 2$, the operator $\mathcal{H}_{\Psi, B}$ is noncompact in $L^q(\mathbb{R}^n)$ and the result follows.

Corollary 5. *Let $\forall j a_j(u) = b(u)$, where $b(u)$ is μ -measurable and positive and $b(u)^{-n/p}\Phi(u) \in L^1(\Omega)$, $1 \leq p < \infty$. Then the corresponding Hausdorff operator \mathcal{H}_Φ is noncompact in $L^p(\mathbb{R}^n)$ provided it is non-zero.*

References

- [1] A. Brown, P.R. Halmos, A.L. Shields, Cesàro operators, Acta Sci. Math. (Szeged) 26, 125–137 (1965)
- [2] G. Brown, F. Móricz, Multivariate Hausdorff operators on the spaces $L^p(\mathbb{R}^n)$, J. Math. Anal. Appl., 271, 443–454 (2002)
- [3] Yu.A. Brychkov, H.-J. Glaeske, A.P. Prudnikov, and Vu Kim Tuan, Multidimensional Integral Transformations. Gordon and Breach, New York - Philadelphia - London - Paris - Montreux - Tokyo - Melbourne - Singapore (1992)
- [4] J. Chen, D. Fan, S. Wang, Hausdorff operators on Euclidean space (a survey article), Appl. Math. J. Chinese Univ. Ser. B (4) 28, 548–564 (2014)

- [5] C. K. Fong, E. A. Nordgren, M. Radjabalipour, H. Radjavi, P. Rosenthal, Extensions of Lomonosov's invariant subspace theorem. *Acta Sci. Math. (Szeged)*, 41, 55 – 62 (1979)
- [6] C. Georgakis, The Hausdorff mean of a Fourier-Stieltjes transform, *Proc. Am. Math. Soc.* 116, 465–471 (1992)
- [7] G.H. Hardy, *Divergent Series*. Clarendon Press, Oxford (1949)
- [8] C.S. Kubrusly, *Elements of Operator Theory*, Birkhauser, Boston, 2001.
- [9] A. Lerner and E. Liflyand, Multidimensional Hausdorff operators on the real Hardy space, *J. Austr. Math. Soc.*, 83, 79–86 (2007)
- [10] E. Liflyand, Open problems on Hausdorff operators, In: *Complex Analysis and Potential Theory*, Proc. Conf. Satellite to ICM 2006, Gebze, Turkey, 8-14 Sept. 2006; Eds. T. Aliyev Azeroglu and P.M. Tamrazov; World Sci., 280–285 (2007)
- [11] E. Liflyand, Hausdorff operators on Hardy spaces, *Eurasian Mathematical Journal*, 4, 101 – 141 (2013)
- [12] E. Liflyand, F. Móricz, The Hausdorff operator is bounded on the real Hardy space $H^1(\mathbb{R})$, *Proc. Am. Math. Soc.*, 128, 1391–1396 (2000).
- [13] A. R. Mirotin, Boundedness of Hausdorff operators on Hardy spaces H^1 over locally compact groups. To appear in the *Journal of Mathematical Analysis and Applications*, Preprint. arXiv:1808.08257v2 [math.FA] 1 Sep 2018.
- [14] A. R. Mirotin, Hilbert transform in context of locally compact abelian groups, *Int. J. Pure and Appl. Math.* 51, 463 – 474 (2009)
- [15] B. E. Rhoades, Spectra of some Hausdorff operators, *Acta. Sci. Math. (Szeged)* 32, 91 – 100 (1972)
- [16] E. Stein, G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton University Press, Princeton (1971)
- [17] E. T. Whittaker and G. N. Watson. *A Course of Modern Analysis. Part II. The Transcendental Functions*, Cambridge University Press; 4th edition, Cambridge (1927)

Adolf R. Mirotin,
 DEPARTMENT OF MATHEMATICS AND PROGRAMMING TECHNOLOGIES,
 F. SKORINA GOMEL STATE UNIVERSITY, 246019, Sovietskaya, 104, Gomel,
 Belarus