

Realizing data features by deep nets

Zheng-Chu Guo, Lei Shi, and Shao-Bo Lin

Abstract—This paper considers the power of deep neural networks (deep nets for short) in realizing data features. Based on refined covering number estimates, we find that, to realize some complex data features, deep nets can improve the performances of shallow neural networks (shallow nets for short) without requiring additional capacity costs. This verifies the advantage of deep nets in realizing complex features. On the other hand, to realize some simple data feature like the smoothness, we prove that, up to a logarithmic factor, the approximation rate of deep nets is asymptotically identical to that of shallow nets, provided that the depth is fixed. This exhibits a limitation of deep nets in realizing simple features.

Index Terms—Neural networks, Approximation rates, Deep nets, Covering numbers, Data feature.

I. INTRODUCTION

Deep learning [12] is recognized to be a state-of-the-art scheme in artificial intelligence and machine learning and has recently triggered enormous research activities. Deep neural networks (deep nets for short) is believed to be capable of discovering deep features of data which are important but are impossible to be found by shallow neural networks (shallow nets for short). It, however, simultaneously produces a series of challenges such as the efficient computation, algorithmic solvability, robustness, interpretability and so on. A direct consequence of these challenges is that users hesitate to utilize deep learning in learning tasks with high risk such as the clinical diagnosis and financial investment, since it is not clear whether deep nets perform essentially better than the scheme in hand. Thus, it is urgent and crucial to provide the theoretical guidance on “when do deep nets perform better than shallow nets?”

Generally speaking, there are three steps to study the above problem. The first step is to correspond specific real-world applications to some data features. For example, figures are assumed to be local similarity [47]; earthquake forecasting is related to rotation-invariant features [1]; and computer vision requires the sparseness of activated neurons on the receptive field [48]. The second step is to connect these data features with a-priori information which can be mathematically reflected by specific properties of functions. In particular, local similarity usually corresponds to piece-wise smooth functions [38]; rotation-invariance generally corresponds to radial functions [6] and sparseness on the receptive field frequently corresponds to sparseness in the spacial domain [25]. The last step is to pursue the outperformance of deep nets in approximating or learning these application-related

functions. In fact, the outperformance of deep nets has been rigorously verified in approximating piece-wise smooth functions [38], rotation-invariant functions [6] and sparse functions [25], which coincides with the empirical evidences on image classification [20], earthquake prediction [46] and computer vision [21].

With the rapid development in deep nets approximation theory, there are numerous features that are proved to be realizable by deep nets [5], [25], [33], [36], [38], [45] with much less neurons than shallow nets. Different from these encouraging results, studies in learning theory showed that, however, to realize these features, capacities of deep nets are much larger than those of shallow nets with comparable number of free parameters. In particular, under some specified capacity measurements such as the number of linear regions [37], Betti numbers [2], number of monomials [7], it was proved that the capacity of deep nets increases exponentially with respect to the depth but polynomially with respect to the width. An extreme case is that there exist deep nets with two hidden layers whose capacity measured by the pseudo-dimension is infinite [28], [29]. The large capacity of deep nets inevitably makes the deep nets learner sensitive to noise and requires a large amount of computations to find a good estimator.

In a nutshell, previous studies on advantages of deep nets showed that deep nets are capable of realizing various application-related data features, but it requires additional capacity costs. The first purpose of our study is to figure out whether the large capacity of deep nets to realize data features is necessary. Our study is based on two interesting observations from the literature [3], [25], [33], [36], [38], [45], [49]. One is that the number of layers of deep nets to realize various data features is small, the order of which is at most the logarithm of the number of free parameters. The other is that the magnitude of free parameters is relatively small, which is at most a polynomial with respect to the number of free parameters. With these two findings, we adopt the well known covering number [51], [52] to measure the capacity of deep nets with controllable number of layers and magnitude of weights and present a refined estimate of the covering number of deep nets. In particular, we prove that the covering number of deep nets with controllable depth and magnitude of weights is similar as that of shallow nets with comparable free parameters. This finding together with existing results in approximation theory shows that, to realize various features such as sparseness, hierarchy, rotation-invariance and manifold structures, deep nets improve the performance of shallow nets without bringing additional capacity costs.

As is well known, advantages of deep nets in realizing some special features do not mean that deep nets are always better than shallow nets. Our second purpose is to demon-

Z. C. Guo is with School of Mathematical Sciences, Zhejiang University, Hangzhou, China. L. Shi is with Shanghai Key Laboratory for Contemporary Applied Mathematics, School of Mathematical Sciences, Fudan University, Shanghai, China. S. B. Lin is with Department of Mathematics, Wenzhou University, Wenzhou, China. Z. C. Guo and L. Shi are the co-first author. The corresponding author is S. B. Lin (email: sblin1983@gmail.com).

strate the necessity of deepening networks in realizing some simple data features. After building a close relation between approximation rates and covering number estimates, we prove that if only the smoothness feature is explored, then up to a logarithmic factor, approximation rates of shallow nets and deep nets with controllable depth and magnitude of weights are asymptotically identical. Combining the above two statements, we indeed present rigorous theoretical verifications to support that deep nets are necessary in a large number of applications corresponding to complex data features, in the sense that deep nets realize data features without any additional capacity costs, but not all.

The rest of paper is organized as follows. In the next section, after reviewing some advantages of deep nets in approximation, we present a refined covering number estimate for deep nets. In Section III, we give a lower bound for deep nets approximation to show the limitation for deep nets in realizing simple features. In the last section, we draw a simple conclusion of this paper.

II. ADVANTAGES OF DEEP NETS IN REALIZING FEATURE

In this section, we study advantages of deep nets in approximating classes of functions with complex features. After introducing some mathematical concepts associated with deep nets, we review some important results in approximation theory which show that deep nets can realize some application-related features that cannot be approximated by shallow nets with comparable free parameters. Then, we present a refined covering number estimate for deep nets to show that deepening networks in some special way does not enlarge the capacity of shallow nets.

A. Deep nets with fixed structures

Great progress of deep learning is built on deepening neural networks with structures. Deep nets with different structures have been proved to be universal, i.e., [53], [54] for deep convolutional nets, [14] for deep nets with tree structures and [10] for deep fully-connected neural networks.

Let $\mathbb{I} := [-1, 1]$ and $x = (x^{(1)}, \dots, x^{(d)}) \in \mathbb{I}^d = [-1, 1]^d$. Let $L \in \mathbb{N}$ and $d_0, d_1, \dots, d_L \in \mathbb{N}$ with $d_0 = d$. Assume $\sigma_k : \mathbb{R} \rightarrow \mathbb{R}$, $k = 1, \dots, L$, be univariate nonlinear functions. For $\vec{h} = (h^{(1)}, \dots, h^{(d_k)})^T \in \mathbb{R}^{d_k}$, define $\vec{\sigma}_k(\vec{h}) = (\sigma_k(h^{(1)}), \dots, \sigma_k(h^{(d_k)}))^T$. Deep nets with depth L and width d_j in the j -th hidden layer can be mathematically represented as


$$h_{\{d_0, \dots, d_L, \sigma\}}(x) = \vec{a} \cdot \vec{h}_L(x), \quad (1)$$

where

$$\vec{h}_k(x) = \vec{\sigma}_k(W_k \cdot \vec{h}_{k-1}(x) + \vec{b}_k), \quad k = 1, 2, \dots, L, \quad (2)$$

$\vec{h}_0(x) = x$, $\vec{a} \in \mathbb{R}^{d_L}$, $\vec{b}_k \in \mathbb{R}^{d_k}$, and $W_k = (W_k^{i,j})_{i=1, j=1}^{d_k, d_{k-1}}$ be a $d_k \times d_{k-1}$ matrix. Denote by $\mathcal{H}_{\{d_0, \dots, d_L, \sigma\}}$ the set of all these deep nets. When $L = 1$, the function defined by (1) is the classical shallow net.

The structure of deep nets can be reflected by structures of the weight matrices W_k and parameter vectors \vec{b}_k and \vec{a} , $k = 1, 2, \dots, L$. For examples, deep convolutional neural networks corresponds to Toeplitz-type weight matrices [54] and



(a) Deep fully-connected nets (b) Deep nets with tree structure

Fig. 1. Structures for deep nets

deep nets with tree structures usually correspond extremely sparse weight matrices [36]. Throughout this paper, a deep net with specific structures refers to a deep nets with specific structures of all W_k, \vec{b}_k , $k = 1, \dots, L$ and \vec{a} . Figure 1 shows two structures for deep nets.

Although deep fully-connected neural networks possess better approximation ability than other networks, the number of free parameters of this type networks is

$$\mathcal{A}_L = d_L + \sum_{k=1}^L (d_{k-1}d_k + d_k), \quad (3)$$

which is huge when the width and depth are large. A recent focus in deep nets approximation is to pursue the approximation ability of deep nets with fixed structures. Up till now, numerous theoretical results [39], [54], [6], [38] showed that the approximation ability of deep fully-connected neural networks can be maintained by deep nets with some special structures with much less free parameters.

In this paper, we are interested in deep nets with structures. For $k = 1, \dots, L$, we assume that the structure of deep nets is fixed and there are $\mathcal{F}_{k,w}$ free parameters in W_k , $\mathcal{F}_{k,b}$ free thresholds in \vec{b}_k and $\mathcal{F}_{L,a}$ free parameters in \vec{a} . Then, there are totally

$$n := \sum_{k=1}^L (\mathcal{F}_{k,w} + \mathcal{F}_{k,b}) + \mathcal{F}_{L,a} \quad (4)$$

free parameters in the deep nets. We assume further $n \ll \mathcal{A}_L$. Throughout the paper, we say there are $\mathcal{F}_{k,w}$ free parameters in W_k , if the weight matrix W_k is generated through the following three ways. The first way is that the matrix has $\mathcal{F}_{k,w}$ entries that can be determined freely, while the reminder $d_k d_{k-1} - \mathcal{F}_{k,w}$ entries are fixed, e.g., the weight matrix in deep nets with tree structures. The second way is that the weight matrix W_k is exactly generated by $\mathcal{F}_{k,w}$ free parameters, e.g., the Toeplitz-type weight matrix in deep convolutional neural networks. The third way is that the weight matrix is generated jointly by both way above, that is, part of the weight matrix is fixed, while the remaining part are totally generated by $\mathcal{F}_{k,w}$ free parameters. Denote by $\mathcal{H}_{\{n, L, \sigma\}}$ the set of all these deep nets with L hidden layers, fixed structure and n free parameters. Denote further

$$\begin{aligned} \mathcal{H}_{\{n, L, \sigma, \mathcal{R}\}} &= \{h_{n, L, \sigma} \in \mathcal{H}_{\{n, L, \sigma\}} : \\ &|w_k^{i,j}|, |b_k^i|, |a_i| \leq \mathcal{R}, 1 \leq i \leq d_k, 1 \leq j \leq d_{k-1}, \\ &1 \leq k \leq L\} \end{aligned} \quad (5)$$

the set of deep nets whose weights and thresholds are uniformly bounded by \mathcal{R} , where \mathcal{R} is some positive number which may depend on $n, d_k, k = 0, 1, \dots, L$ and L . We aim at studying the approximation ability and capacity of $\mathcal{H}_{\{n, L, \sigma, \mathcal{R}\}}$.

It should be mentioned that the boundedness assumption in (5) is necessary. In fact, without such an assumption, [28], [13] proved that for arbitrary $\varepsilon > 0$ and arbitrary continuous function f , a deep net with two hidden layers and finitely many free parameters is fully able to generate an approximation H_f , such that

$$\|f - H_f\|_{L_p(\mathbb{I}^d)} \leq \varepsilon. \quad (6)$$

This implies that the capacity of deep nets with two hidden layers and finitely many free parameters is comparable with that of $L_p(\mathbb{I}^d)$, showing its extremely large capacity. Therefore, to further control the capacity of deep nets, the boundedness assumption has been employed in large literature [14], [25], [38].

B. A fast review for realizing data features by deep nets

In approximation and learning theory, data features are usually formulated by a-priori information for corresponding functions, like the target function [3] for approximation, regression function [25] for regression and Bayes decision function [23] for classification. Studying advantages of deep nets in approximating functions with different a-priori information is a classical topic. It can date back to 1994, when [3] deduced the localized approximation property of deep nets which is far beyond the capability of shallow nets.

The localized approximation of a neural network shows that if the target function is modified only on a small subset of the Euclidean space, then only a few neurons, rather than the entire network, need to be retrained. We refer to [3, Def.2.1] for a formal definition of localized approximation. Since the localized approximation is an important step-stone in approximating piecewise smooth functions [38] and sparse functions in spacial domains [25], deep nets perform much better than shallow nets in related applications such as image processing and computer vision [8]. The following proposition, which can be found in [3, Theorem 2.3] (see also [25]), shows the localized approximation property of deep nets.

Proposition 1. Suppose that $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ is a bounded measurable function with the sigmoidal property

$$\lim_{t \rightarrow -\infty} \sigma(t) = 0, \quad \lim_{t \rightarrow +\infty} \sigma(t) = 1. \quad (7)$$

Then there exists a deep net with two hidden layers, $2d + 1$ neurons and activation function σ provides localized approximation.

Rotation-invariance, is another popular data feature, which abounds in statistical physics [22], earthquake early warning [42] and image rendering [34]. Mathematically, rotation-invariant property corresponds to a radial function which is by definition a function whose value at each point depends only on the distance between that point and the origin. In the nice papers [15], [16], shallow nets were proved to be incapable of embodying rotation-invariance features. To show the power

of depth in approximating radial functions, we present the definition of smooth radial function as follows.

Definition 1. Let $\mathbb{A} \subset \mathbb{R}$, $c_0 > 0$ and $r = s + v$ with $s \in \mathbb{N}_0 := \{0\} \cup \mathbb{N}$ and $0 < v \leq 1$. We say a univariate function $g : \mathbb{A} \rightarrow \mathbb{R}$ is (r, c_0) -Lipschitz continuous if g is s -times differentiable and its s -th derivative satisfies the Lipschitz condition

$$|g^{(s)}(t) - g^{(s)}(t_0)| \leq c_0 |t - t_0|^v, \quad \forall t, t_0 \in \mathbb{A}. \quad (8)$$

Denote by $Lip_{\mathbb{A}}^{(r, c_0)}$ the set of all (r, c_0) -Lipschitz continuous functions defined on \mathbb{A} . Denote also by $Lip^{(\diamond, r, c_0)}$ the set of radial functions $f = g(\|x\|_2^2)$ with $g \in Lip_{[0, 1]}^{(r, c_0)}$.

The following proposition, which can be found in [6], shows that deep nets can realize rotation-invariance and smoothness features of target functions, simultaneously.

Proposition 2. Let $d \geq 2$ and $1 \leq p \leq \infty$. If σ is the logistic function, i.e. $\sigma(t) = \frac{1}{1+e^{-t}}$, then for arbitrary $f \in Lip^{(\diamond, r, c_0)}$, there is an $h \in \mathcal{H}_{\{n, 3, \sigma, \mathcal{R}\}}$ such that

$$\|f - h\|_{L_p(\mathbb{I}^d)} \leq C_1 n^{-r}. \quad (9)$$

Furthermore, for arbitrary $h' \in \mathcal{H}_{\{n, 1, \sigma, \mathcal{R}\}}$, there always exists a function $f_0 \in Lip^{(\diamond, r, c_0)}$ satisfying

$$\|f_0 - h'\|_{L_\infty(\mathbb{I}^d)} \geq C_2 n^{-r/(d-1)}, \quad (10)$$

where C_1, C_2 are constants independent of d_0, d_1, \dots, d_L or n .

Numerous learning problems [26] in computer vision, gene analysis and speech processing involve high dimensional data. These data are often governed by many fewer variables, producing manifold-structure features in a high dimensional ambient space. A large number of theoretical studies [5], [45], [50] have revealed that shallow nets are difficult to realize smooth and manifold-structure features simultaneously. Conversely, deep nets, as studied in [45], [5], is capable of reflecting these features, which is shown by the following proposition [45] (see also [5]).

Proposition 3. Let $\Gamma \subset \mathbb{I}^d$ be a smooth d' -dimensional compact manifold (without boundary) with $d' \ll d$. If σ is the ReLU activation function, i.e. $\sigma(t) = \max\{t, 0\}$, and f is defined on Γ and twice differentiable, then there exists a $g \in \mathcal{H}_{n, 4, \sigma, \mathcal{R}}$ such that

$$\|f - g\|_{L^2(\mathbb{I}^d)} \leq C_3 n^{-\frac{2}{d'}}. \quad (11)$$

where C_3 is a constant independent of d_0, d_1, \dots, d_L or n .

The previous studies showed that, compared with shallow nets, deep nets equipped with fewer parameters are enough to approximate functions with complex features to the same accuracy. In the following Table I, we list some literature on studying the advantages of realizing data futures.

C. Covering number estimates

In the above subsection, we have reviewed some results on the advantages of deep nets in realizing data features. However, it does not mean that deep nets are better than shallow nets, since we do not know what price is paid for

References	Features	σ	L
[3], [5]	Localized approximation	Sigmoidal	2
[25]	Sparse+Smooth	Sigmoidal	2
[45]	Smooth+Manifold	ReLU	4
[35], [14]	Hierarchical+Smooth	Sigmoidal	Hierarchical
[38]	piecewise smooth	ReLU	Finite
[41]	ℓ_1 radial+smooth	ReLU	$\log(\varepsilon^{-1})$
[22], [40]	Sparse (frequency)	Analytic	$\log(\varepsilon^{-1})$

TABLE I
POWERS OF DEEP NETS IN APPROXIMATION (WITHIN ACCURACY ε)

such advantages in approximation. In this subsection, we use the covering number, which is widely used in learning theory [23], [43], [44], [51], [52], to measure the capacity of $\mathcal{H}_{n,L,\sigma,\mathcal{R}}$ and then unify the comparison within the same framework to show the outperformance of deep nets.

Let \mathbb{B} be a Banach space and V be a subset of \mathbb{B} . Denote by $\mathcal{N}(\varepsilon, V, \mathbb{B})$ the ε -covering number of V under the metric of \mathbb{B} , which is the minimal number of elements in an ε -net of V . If $\mathbb{B} = L_1(\mathbb{I}^d)$, we denote $\mathcal{N}(\varepsilon, V) := \mathcal{N}(\varepsilon, V, L_1(\mathbb{I}^d))$ for brevity. Our purpose is a tight bound for covering numbers of $\mathcal{H}_{n,L,\sigma,\mathcal{R}}$. To this end, we need the following assumption.

Assumption 1. For arbitrary $t \in \mathbb{R}$ and every $k \in \{1, \dots, L\}$, assume

$$|\sigma_k(t) - \sigma_k(t')| \leq c_1 |t - t'| \quad (12)$$

and

$$|\sigma_k(t)| \leq c(|t| + 1) \quad (13)$$

for some $c, c_1 \geq 1$.

To be detailed, (12) shows the Lipschitz continuous property of σ_k and (13) exhibits the linear increasing condition of σ_k . These assumptions have been utilized in [32], [17], [25] to quantify covering numbers of neural networks with different structures. We can see that almost all widely used activation functions such as the logistic function, hyperbolic tangent sigmoidal function $\sigma(t) = \frac{1}{2}(\tanh(t) + 1)$ with $\tanh(t) = (e^{2t} - 1)/(e^{2t} + 1)$, arctan sigmoidal function $\sigma(t) = \frac{1}{\pi} \arctan(t) + \frac{1}{2}$, Gompertz function $\sigma(t) = e^{-ae^{-bt}}$ with $a, b > 0$, ReLU $\sigma(t) = \max\{t, 0\}$, and Gaussian function $\sigma(t) = e^{-t^2}$ satisfy Assumption 1. With this assumption, we present our first main result in the following theorem, whose proof can be found in Appendix A.

Theorem 1. Let $\mathcal{H}_{n,L,\sigma,\mathcal{R}}$ be defined by (5). Under Assumption 1, there holds

$$\mathcal{N}(\varepsilon, \mathcal{H}_{n,L,\sigma,\mathcal{R}}) \leq (c_2 \mathcal{R} D_{\max})^{3(L+1)^2 n} \varepsilon^{-n},$$

where $D_{\max} := \max_{0 \leq \ell \leq L} d_\ell$ and $c_2 \geq 1$ is a constant depending only on c, c_1 and d .

For σ satisfying Assumption 1, it was deduced in [31], [9] that

$$\log \mathcal{N}(\varepsilon, \mathcal{H}_{n,1,\sigma,\mathcal{R}}) = \mathcal{O}\left(n \log \frac{C_4 \mathcal{R}}{\varepsilon}\right), \quad (14)$$

where C_4 is a constant independent of ε or n . From Theorem 1, we can derive

$$\log \mathcal{N}(\varepsilon, \mathcal{H}_{n,L,\sigma,\mathcal{R}}) = \mathcal{O}\left(L^2 n \log \frac{C_5 \mathcal{R}}{\varepsilon}\right) \quad (15)$$

for some C_5 independent of $\varepsilon, L, d_0, d_1, \dots, d_L$ or n . Comparing (15) with (14), we find that, up to a logarithmic factor, deep nets do not essentially enlarge the capacity of shallow nets, provided that they have same number of free parameters and the depth of deep nets is at most $\log n$. Noting that the depths of deep nets in Table I all satisfy this constraint, Theorem 1 shows that to realize various data features presented in Table I, deep nets can improve the performance of shallow nets without imposing additional capacity costs. Therefore, Theorem 1 together with Table I yields the reason why deep nets perform much better than shallow nets in some complex learning tasks such as image processing and computer vision.

Recently, [11] presented a tight VC-dimension bounds for piecewise linear neural networks. In particular, they proved that

$$VCDim(\text{sgn}(\mathcal{H}_{\{d_0, \dots, d_L, \sigma\}})) = \mathcal{O}(Ln \log n), \quad (16)$$

where $VCDim(V)$ denotes the VC-dimension of the set V and $\text{sgn}(V) := \{x \rightarrow \text{sgn}(f(x)) : f \in V\}$, where $\text{sgn}(f(x)) = 1$ if $f(x) \geq 0$ and $\text{sgn}(f(x)) = -1$ otherwise. Using the standard approach in [9, Chap.9], we can derive

$$\log \mathcal{N}(\varepsilon, \mathcal{H}_{n,L,\sigma,\mathcal{R}}) = \mathcal{O}\left(Ln \log \frac{C_5 \mathcal{R}}{\varepsilon}\right) \quad (17)$$

provided that $\sigma_1 = \sigma_2 = \dots = \sigma_L$ are piecewise linear, where C_6 is a constant independent of $\varepsilon, L, d_0, \dots, d_L$ or n . Comparing (17), there is an additional L in our analysis. The reason is that we focus on all activation functions satisfying (1) rather than piecewise activation functions. It should be also mentioned that similar covering number estimates for deep nets with tree structures has been studied in [6], [14], [25]. We highlight that different structures yield essentially non-trivial approaches. In fact, due to tree structures, the approach in [6], [14], [25] is just to decouple layers by using the boundedness and Lipschitz property of activation functions. However, in estimating covering number of deep nets with arbitrarily fixed structure, we need a novel matrix-vector transformation technique, as presented in Appendix A.

III. NECESSITY OF THE DEPTH

Previous studies showed that, to realize some complex data features, deep nets can improve the performance of shallow nets without additional capacity costs. In this section, we study in a different direction to prove that, to realize some simple data features, deep nets are not essentially better than shallow nets.

A. Limitations of deep nets approximation

Smoothness or regularity is a widely used feature that has been adopted in a vast literature [3], [4], [15], [16], [28], [29], [49]. To present the approximation result, we at first introduce the following definition.

Definition 2. Let $c_0 > 0$ and $r = s + v$ with $s \in \mathbb{N}_0 := \{0\} \cup \mathbb{N}$ and $0 < v \leq 1$. We say a function $f : \mathbb{I}^d \rightarrow \mathbb{R}$ is (r, c_0) -smooth if f is s -times differentiable and for every $\alpha_j \in \mathbb{N}_0$, $j = 1, \dots, d$ with $\alpha_1 + \dots + \alpha_d = s$, its s -th partial derivative satisfies the Lipschitz condition

$$\left| \frac{\partial^s f}{\partial x_1^{\alpha_1} \dots \partial x_d^{\alpha_d}}(x) - \frac{\partial^s f}{\partial x_1^{\alpha_1} \dots \partial x_d^{\alpha_d}}(x') \right| \leq c_0 \|x - x'\|^v, \quad (18)$$

where $x, x' \in \mathbb{I}^d$ and $\|x\|$ denotes the Euclidean norm of $x \in \mathbb{R}^d$. Denote by $\text{Lip}^{(r, c_0)}$ the set of all (r, c_0) -smooth functions defined on \mathbb{I}^d .

Approximating smooth functions is a classical topic in neural networks approximation. It is well known that the approximation rate can be as fast as $O(n^{-r/d})$ for neural networks with n free parameters. In particular, the Jackson-type error estimate

$$\text{dist}(\text{Lip}^{(r, c_0)}, \mathcal{H}_{\{n, 1, \sigma, \mathcal{R}\}}, L_p(\mathbb{I}^d)) \leq C'_1 n^{-\frac{r}{d}} \quad (19)$$

has been established [30] for shallow nets with analytic activation functions, where

$$\begin{aligned} \text{dist}(U, V, L_p(\mathbb{I}^d)) &:= \sup_{f \in U} \text{dist}(f, V, L_p(\mathbb{I}^d)) \\ &:= \sup_{f \in U} \inf_{g \in V} \|f - g\|_{L_p(\mathbb{I}^d)} \end{aligned}$$

denotes the deviations of U from V in $L_p(\mathbb{I}^d)$ for $U, V \subseteq L_p(\mathbb{I}^d)$. Similar results has been derived in [3] with deep nets with two hidden layers and a sigmoidal activation function. Recently, [49] derived an error estimate taking the form of

$$\text{dist}(\text{Lip}^{(r, c_0)}, \mathcal{H}_{n, L, \sigma, \mathcal{R}}, L_p(\mathbb{I}^d)) \leq C'_2 n^{-\frac{r}{d}} \log n \quad (20)$$

for deep nets with $L = \log n$ and ReLU activation functions. We would like to point out that, for shallow nets with ReLU activation functions, estimates (20) holds only for $0 < r \leq 1$, which is also considered as the approximation bottleneck of shallow nets. The paper [49] showed that deepening the networks can overcome this bottleneck for shallow nets. However, it should be mentioned from (19) that for other activation functions except the ReLU activation functions, such a bottleneck does not exist. Thus, the paper [49] indeed conduct a nice analysis on the necessity of deepening ReLU nets. However, their established results can not illustrate the necessity of depth.

In the following theorem that will be proved in Appendix C, we show that deep nets cannot be essentially better than shallow nets in realizing the smoothness feature.

Theorem 2. Let $1 \leq p \leq \infty$, $L \in \mathbb{N}$. Then


$$\begin{aligned} &\text{dist}(\text{Lip}^{(r, c_0)}, \mathcal{H}_{n, L, \sigma, \mathcal{R}}, L_1(\mathbb{I}^d)) \\ &\geq C [L^2 n \log_2 n \log_2(\mathcal{R} D_{\max})]^{-\frac{r}{d}}, \end{aligned} \quad (21)$$

where C is a constant depending only on c, c_0, c_1, d and r .

Combining the estimates (21) and (19), and noting

$$\|f\|_{L^1(\mathbb{I}^d)} \leq C_{d,p} \|f\|_{L^p(\mathbb{I}^d)}$$

with $C_{d,p}$ a constant depending only on d and p , we see that, when L is not too large, deep nets cannot essentially improve



(a) Approximation by shallow nets (b) Approximation by DFCNs

Fig. 2. Comparison between deep and shallow nets

the approximation rate if one only considers the smoothness feature. When L is too large, it follows from Theorem 1 that we will need additional capacity cost for deep nets to improve the approximation ability of shallow nets. In other words, the smoothness feature is not sufficient to judge whether the depth of neural networks is necessary.

B. Remarks and discussions

Limitations of the approximation capabilities of shallow nets were firstly studied in [4] in terms of providing lower bounds of approximation of smooth functions in the minimax sense. Recently, [24] highlighted that there exists a probabilistic measure, under which, all smooth functions cannot be approximated by shallow nets very well with high confidence. In another two interesting papers [18], [19], limitations of shallow nets were presented in terms of establishing lower bound of approximating functions with some variation restrictions. However, due to these results, it is still not clear whether the depth of neural networks is necessary, if only the smoothness information is given.

Theorem 2 goes further along this direction and presents a negative answer. In Theorem 2, to realize smoothness features, deep nets perform almost the same as shallow nets. This result verifies the common consensus that deep learning outperforms shallow learning in some “difficult” learning tasks [8], but not always. Moreover, our result also implies that whether deep nets can help to improve the performance of the existing learning schemes depends on what features for data we are exploring. Combining our work with [35], [36], [14], [38], [33], [5], [6], [25], we can illustrate the comparison between shallow and deep nets in Figure 2.

We declare that Theorem 2 only presents limitations of deep nets in realizing smooth features. As shown in Figure 2, if more features are explored, we believe that the approximation rate of deep nets can break through the lower bound presented in (21).

IV. CONCLUSION

In this paper, we study the advantages and limitations of deep nets in realizing different data features. Our results showed that, in realizing some complex data features such as the rotation-invariance, manifold structure, hierarchical structure, sparseness, deep nets can improve the performance of shallow nets without additional capacity costs. We also exhibit that for some simple data features like the smoothness, deep nets performs essentially similar as shallow nets.

APPENDIX A: PROOF OF THEOREM 1

For $1 \leq \ell \leq L$, let $\mathcal{W}_{\mathcal{F}_{\ell,w}}^*$ be the set of $d_\ell \times d_{\ell-1}$ matrices with fixed structures and total $\mathcal{F}_{\ell,w}$ free parameters and $\vec{B}_{\mathcal{F}_{\ell,b}}^*$ be the set of $\mathcal{F}_{\ell,b}$ -dimensional vectors with fixed structures and total $\mathcal{F}_{\ell,b}$ free parameters. Denote

$$\mathcal{W}_{\mathcal{F}_{\ell,w}} := \{W \in \mathcal{W}_{\mathcal{F}_{\ell,w}}^* : |W^{i,j}| \leq \mathcal{R}\}$$

and

$$\vec{\mathcal{B}}_{\mathcal{F}_{\ell,b}} := \{\vec{b} \in \vec{B}_{\mathcal{F}_{\ell,b}}^* : |b_i| \leq \mathcal{R}\}.$$

For $x \in \mathbb{I}^d$, let $\vec{\mathcal{H}}_0 = \{x\}$ and define iteratively

$$\begin{aligned} \vec{\mathcal{H}}_\ell &= \{\vec{h}_\ell(x) = \vec{\sigma}_\ell(W_\ell \vec{h}_{\ell-1}(x) + \vec{b}_\ell) : \\ \vec{h}_{\ell-1} &\in \vec{\mathcal{H}}_{\ell-1}, W_\ell \in \mathcal{W}_{\mathcal{F}_{\ell,w}}, \vec{b}_\ell \in \vec{\mathcal{B}}_{\mathcal{F}_{\ell,b}}\}, \quad \ell = 1, 2, \dots, L. \end{aligned} \quad (22)$$

For each $\vec{h}_\ell = (h_\ell^1, \dots, h_\ell^{d_\ell}) \in \vec{\mathcal{H}}_\ell$, define $\|\vec{h}_\ell\|_{*,d_\ell} := \max_{1 \leq i \leq d_\ell} \|h_\ell^i\|_{L^1(\mathbb{I}^d)}$. The following lemma devotes to the uniform bound of functional vectors in $\vec{\mathcal{H}}_\ell$. In our analysis, we always assume that the activation functions satisfy Assumption 1 with uniform constants c and c_1 . Moreover, we also suppose that $\mathcal{R} \geq 1$ and $c \geq 1$.

Lemma 1. *For each $\ell = 1, 2, \dots, L$ and $\vec{h}_\ell \in \vec{\mathcal{H}}_\ell$, if σ_ℓ satisfies (13), then there holds*

$$\|\vec{h}_\ell\|_{*,d_\ell} \leq (c(1 + 2^{d+1})\mathcal{R})^\ell d_{\ell-1} \cdots d_0. \quad (23)$$

Proof: For arbitrary $\ell = 1, \dots, L$, it follows from (13) that

$$\begin{aligned} \|\vec{h}_\ell\|_{*,d_\ell} &= \left\| \vec{\sigma}_\ell(W_\ell \vec{h}_{\ell-1}(x) + \vec{b}_\ell) \right\|_{*,d_\ell} \\ &= \max_{1 \leq i \leq d_\ell} \int_{\mathbb{I}^d} \left| \sigma_\ell(W_\ell^i \cdot \vec{h}_{\ell-1}(x) + b_\ell^i) \right| dx \\ &\leq c \max_{1 \leq i \leq d_\ell} \int_{\mathbb{I}^d} \left(\left| \sum_{j=1}^{d_{\ell-1}} W_\ell^{i,j} h_{\ell-1}^j(x) + b_\ell^i \right| + 1 \right) dx \\ &\leq c \max_{1 \leq i \leq d_\ell} \left\{ \sum_{j=1}^{d_{\ell-1}} |W_\ell^{i,j}| \int_{\mathbb{I}^d} |h_{\ell-1}^j(x)| dx + |b_\ell^i| 2^d \right\} + c 2^d \\ &\leq (cd_{\ell-1} \|\vec{h}_{\ell-1}\|_{*,d_{\ell-1}} + c 2^d \mathcal{R}) + c 2^d, \end{aligned}$$

where W_ℓ^i denotes the i -row of the matrix W_ℓ , $W_\ell^{i,j}$ denotes the (i, j) -element of W_ℓ , $\vec{b}_\ell = (b_\ell^1, \dots, b_\ell^{d_\ell})$ and $\vec{h}_{\ell-1} \in \vec{\mathcal{H}}_{\ell-1}$. Noting $\|\vec{h}_0\|_{*,d_0} = \max_{1 \leq i \leq d_0} \int_{\mathbb{I}} |x^i| dx = 1$, we then have

$$\|\vec{h}_\ell\|_{*,d_\ell} \leq (c(1 + 2^{d+1})\mathcal{R})^\ell d_{\ell-1} \cdots d_0.$$

This finishes the proof of Lemma 1. \blacksquare

Our second lemma aims at deriving covering number of some matrix and vector with fixed free parameters.

Lemma 2. *For arbitrary $\varepsilon > 0$ and $1 \leq \ell \leq L$, we have*

$$\mathcal{N}(\varepsilon, \mathcal{W}_{\mathcal{F}_{\ell,w}}, \|\cdot\|_1) \leq \left(\frac{2d_\ell d_{\ell-1} \mathcal{R}}{\varepsilon} \right)^{\mathcal{F}_{\ell,w}},$$

and

$$\mathcal{N}(\varepsilon, \vec{\mathcal{B}}_{\mathcal{F}_{\ell,b}}, \ell_\infty^m) \leq \left(\frac{2\mathcal{R}}{\varepsilon} \right)^{\mathcal{F}_{\ell,b}},$$

where $\|W_\ell\|_1 := \sum_{i=1}^{d_\ell} \sum_{j=1}^{d_{\ell-1}} |W_\ell^{i,j}|$ denotes the 1-norm of the matrix W_ℓ .

Proof: For arbitrary $d_\ell \times d_{\ell-1}$ matrix, we can rewrite it as a $d_\ell \times d_{\ell-1}$ -dimensional vector as $\{w_1, \dots, w_{d_\ell \times d_{\ell-1}}\}$. Without loss of generality, we assume that the first $\mathcal{F}_{\ell,w}$ elements of the $d_\ell \times d_{\ell-1}$ -dimensional vector are free parameters. Let \mathcal{E}_{F_i} be the ε -cover nets of $\{w_i : |w_i| \leq \mathcal{R}\}$, that is, for each $|w_i| \leq \mathcal{R}$, there is a $w'_i \in \mathcal{E}_{F_i}$ such that

$$|w_i - w'_i| \leq \varepsilon, \quad \forall i = 1, \dots, \mathcal{F}_{\ell,w}.$$

Then, for arbitrary $W, W' \in \mathcal{W}_{\mathcal{F}_{\ell,w}}$ with W, W' the matrices corresponding to the vector $(w_1, w_2, \dots, w_{\mathcal{F}_{\ell,w}}, \dots)$ and $(w'_1, w'_2, \dots, w'_{\mathcal{F}_{\ell,w}}, \dots)$ respectively, there holds

$$\begin{aligned} \|W - W'\|_1 &= \sum_{i=1}^{d_\ell} \sum_{j=1}^{d_{\ell-1}} |W^{i,j} - W'^{i,j}| \\ &= \sum_{i=1}^{\mathcal{F}_{\ell,w}} |w_i - w'_i| + \sum_{i=\mathcal{F}_{\ell,w}+1}^{d_\ell d_{\ell-1}} |w_i - w'_i|. \end{aligned}$$

If the reminder $d_\ell d_{\ell-1} - \mathcal{F}_{\ell,w}$ are fixed constants, we have $\sum_{i=\mathcal{F}_{\ell,w}+1}^{d_\ell d_{\ell-1}} |w_i - w'_i| = 0$. If the weight matrix is generated by the other two ways, which implies some elements in the remainder $d_\ell - \mathcal{F}_{\ell,w}$ terms sharing the same values as some elements in the first $\mathcal{F}_{\ell,w}$ terms, then we have

$$\sum_{i=\mathcal{F}_{\ell,w}+1}^{d_\ell d_{\ell-1}} |w_i - w'_i| \leq (d_\ell d_{\ell-1} - \mathcal{F}_{\ell,w}) \max_{1 \leq i \leq \mathcal{F}_{\ell,w}} |w_i - w'_i|.$$

Both cases yield

$$\|W - W'\|_1 \leq d_\ell d_{\ell-1} \varepsilon.$$

Hence $\mathcal{F}_{\ell,w}$ ε -covers for sets $\{w_i : |w_i| \leq \mathcal{R}\}$ with $i = 1, \dots, \mathcal{F}_{\ell,w}$ constitute a $d_\ell d_{\ell-1} \varepsilon$ -cover for $\mathcal{W}_{\mathcal{F}_{\ell,w}}$, which together with $|\mathcal{E}_{F_i}| \leq \frac{2\mathcal{R}}{\varepsilon}$, $i = 1, 2, \dots, \mathcal{F}_{\ell,w}$ implies

$$\mathcal{N}(\varepsilon, \mathcal{W}_{\mathcal{F}_{\ell,w}}, \|\cdot\|_1) \leq \left(\frac{2d_\ell d_{\ell-1} \mathcal{R}}{\varepsilon} \right)^{\mathcal{F}_{\ell,w}},$$

where $|\mathcal{E}|$ denotes the cardinality of the set \mathcal{E} . This completes the first estimate. The second estimates can be derived by using the similar approach. With these, we completes the proof of Lemma 2. \blacksquare

Based on the previous lemmas, we can derive the following iterative estimates for the covering number associated with the affine mapping $\vec{\sigma}(W\vec{h} + \vec{b})$.

Lemma 3. *If σ_ℓ satisfies Assumption 1 for each $\ell = 1, 2, \dots, L$, then*

$$\begin{aligned} \mathcal{N}(\varepsilon, \vec{\mathcal{H}}_\ell, \|\cdot\|_{*,d_\ell}) &\leq (c'_1 \mathcal{R})^{\ell \mathcal{F}_\ell} D_\ell^{2\mathcal{F}_\ell \varepsilon - \mathcal{F}_\ell} \\ &\times \mathcal{N}\left(\frac{\varepsilon}{(c'_1 \mathcal{R})^{\ell-1} D_\ell}, \vec{\mathcal{H}}_{\ell-1}, \|\cdot\|_{*,d_{\ell-1}}\right), \end{aligned}$$

holds for $\ell = 2, \dots, L$ and

$$\mathcal{N}(\varepsilon, \vec{\mathcal{H}}_1, \|\cdot\|_{*,d_1}) \leq \left(\frac{c'_1 \mathcal{R} D_1}{\varepsilon} \right)^{\mathcal{F}_1}.$$

where $D_\ell = d_\ell \cdots d_0$, $\mathcal{F}_\ell = \mathcal{F}_{\ell,w} + \mathcal{F}_{\ell,b}$ and $c'_1 = 6c_1c(1 + 2^{d+1})$.

Proof: For each $\ell = 1, 2, \dots, L$, let $\mathcal{E}_{\ell,w}$ and $\mathcal{E}_{\ell,b}$ be ε -cover nets of $\mathcal{W}_{\mathcal{F}_{\ell,w}}$ and $\mathcal{B}_{\mathcal{F}_{\ell,b}}$ respectively. For $\ell = 2, \dots, L$, let $\mathcal{E}_{\ell,h}$ be the ε -cover nets for $\vec{\mathcal{H}}_{\ell-1}$. Therefore, for each $\vec{h}_{\ell-1} \in \vec{\mathcal{H}}_{\ell-1}$, $W_\ell \in \mathcal{W}_{\mathcal{F}_{\ell,w}}$ and $\vec{b}_\ell \in \mathcal{B}_{\mathcal{F}_{\ell,b}}$, there exist $\vec{h}'_{\ell-1} \in \mathcal{E}_{\ell,h}$, $W'_\ell \in \mathcal{E}_{\ell,w}$, $\vec{b}'_\ell \in \mathcal{E}_{\ell,b}$ such that

$$\|\vec{h}_{\ell-1} - \vec{h}'_{\ell-1}\|_{*,d_{\ell-1}} \leq \varepsilon, \quad \ell = 2, \dots, L \quad (24)$$

and

$$\|W_\ell - W'_\ell\|_1 \leq \varepsilon, \quad \|\vec{b}_\ell - \vec{b}'_\ell\|_{\ell_\infty^{d_\ell}} \leq \varepsilon, \quad \ell = 1, \dots, L \quad (25)$$

Then, for arbitrary $\vec{h}_\ell \in \vec{\mathcal{H}}_\ell$ and $\ell = 2, 3, \dots, L$, there holds

$$\begin{aligned} & \|\vec{h}_\ell - \vec{\sigma}_\ell(W'_\ell \vec{h}'_{\ell-1} + \vec{b}'_\ell)\|_{*,d_\ell} \\ & \leq \|\vec{\sigma}_\ell(W_\ell \vec{h}_{\ell-1} + \vec{b}_\ell) - \vec{\sigma}_\ell(W'_\ell \vec{h}_{\ell-1} + \vec{b}_\ell)\|_{*,d_\ell} \\ & + \|\vec{\sigma}_\ell(W'_\ell \vec{h}_{\ell-1} + \vec{b}_\ell) - \vec{\sigma}_\ell(W'_\ell \vec{h}'_{\ell-1} + \vec{b}_\ell)\|_{*,d_\ell} \\ & + \|\vec{\sigma}_\ell(W'_\ell \vec{h}'_{\ell-1} + \vec{b}_\ell) - \vec{\sigma}_\ell(W'_\ell \vec{h}'_{\ell-1} + \vec{b}'_\ell)\|_{*,d_\ell}. \end{aligned} \quad (26)$$

Due to (12), we get from Lemma 1 that

$$\begin{aligned} & \|\vec{\sigma}_\ell(W_\ell \vec{h}_{\ell-1} + \vec{b}_\ell) - \vec{\sigma}_\ell(W'_\ell \vec{h}_{\ell-1} + \vec{b}_\ell)\|_{*,d_\ell} \\ & \leq \max_{1 \leq i \leq d_\ell} \int_{\mathbb{I}^d} |\sigma_\ell(W_\ell^i \cdot \vec{h}_{\ell-1}(x) + b_\ell^i) \\ & - \sigma_\ell(W'_\ell^i \cdot \vec{h}_{\ell-1}(x) + b_\ell^i)| dx \\ & \leq c_1 \sum_{i=1}^{d_\ell} \int_{\mathbb{I}^d} |(W_\ell^i - W'_\ell^i) \cdot \vec{h}_{\ell-1}(x)| dx \\ & \leq c_1 \|\vec{h}_{\ell-1}\|_{*,d_{\ell-1}} \sum_{i=1}^{d_\ell} \sum_{j=1}^{d_{\ell-1}} |W_\ell^{ij} - W'_\ell^{ij}| \\ & \leq c_1 (c(1 + 2^{d+1})\mathcal{R})^{\ell-1} d_{\ell-2} \cdots d_0 \|W_\ell - W'_\ell\|_1. \end{aligned}$$

For $\ell = 1, 2, \dots, L$, we have from (12) that

$$\begin{aligned} & \|\vec{\sigma}_\ell(W'_\ell \vec{h}_{\ell-1} + \vec{b}_\ell) - \vec{\sigma}_\ell(W'_\ell \vec{h}'_{\ell-1} + \vec{b}_\ell)\|_{*,d_\ell} \\ & \leq \max_{1 \leq i \leq d_\ell} \int_{\mathbb{I}^d} |\sigma_\ell(W'_\ell^i \cdot \vec{h}_{\ell-1}(x) + b_\ell^i) \\ & - \sigma_\ell(W'_\ell^i \cdot \vec{h}'_{\ell-1}(x) + b_\ell^i)| dx \\ & \leq c_1 \max_{1 \leq i \leq d_\ell} \int_{\mathbb{I}^d} |W'_\ell^i \cdot (\vec{h}_{\ell-1}(x) - \vec{h}'_{\ell-1}(x))| dx \\ & \leq c_1 \sum_{i=1}^{d_\ell} \sum_{j=1}^{d_{\ell-1}} |W'_\ell^{ij}| \|\vec{h}_{\ell-1} - \vec{h}'_{\ell-1}\|_{*,d_{\ell-1}} \\ & \leq c_1 d_\ell d_{\ell-1} \mathcal{R} \|\vec{h}_{\ell-1} - \vec{h}'_{\ell-1}\|_{*,d_{\ell-1}} \end{aligned}$$

and

$$\begin{aligned} & \|\vec{\sigma}_\ell(W'_\ell \vec{h}'_{\ell-1} + \vec{b}_\ell) - \vec{\sigma}_\ell(W'_\ell \vec{h}'_{\ell-1} + \vec{b}'_\ell)\|_{*,d_\ell} \\ & = \max_{1 \leq i \leq d_\ell} \int_{\mathbb{I}^d} |\sigma_\ell(W'_\ell^i \cdot \vec{h}'_{\ell-1}(x) + b_\ell^i) \\ & - \sigma_\ell(W'_\ell^i \cdot \vec{h}'_{\ell-1}(x) + b_\ell'^i)| dx \\ & \leq c_1 \max_{1 \leq i \leq d_\ell} \int_{\mathbb{I}^d} |b_\ell^i - b_\ell'^i| dx \leq 2^d c_1 \|\vec{b}_\ell - \vec{b}'_\ell\|_{\ell_\infty^{d_\ell}}. \end{aligned}$$

For $\ell = 2, \dots, L$, plugging the above three estimates into (26), we then get from (24) and (25) that

$$\begin{aligned} & \|\vec{h}_\ell - \vec{\sigma}_\ell(W'_\ell \vec{h}'_{\ell-1} + \vec{b}'_\ell)\|_{*,d_\ell} \\ & \leq 3c_1 (c(1 + 2^{d+1})\mathcal{R})^{\ell-1} d_\ell \cdots d_0 \varepsilon. \end{aligned}$$

This implies that

$$\{\vec{\sigma}_\ell(W'_\ell \vec{h}'_{\ell-1} + \vec{b}'_\ell) : W'_\ell \in \mathcal{E}_{\ell,w}, \vec{h}'_{\ell-1} \in \mathcal{E}_{\ell,h}, \vec{b}'_\ell \in \mathcal{E}_{\ell,b}\}$$

is a $3c_1 (2c(1 + 2^{d+1})\mathcal{R})^{\ell-1} d_\ell \cdots d_0 \varepsilon$ -net of $\vec{\mathcal{H}}_\ell$. This together with Lemma 2 implies

$$\begin{aligned} & \mathcal{N}\left(3c_1 (c(1 + 2^{d+1})\mathcal{R})^{\ell-1} d_\ell \cdots d_0 \varepsilon, \vec{\mathcal{H}}_\ell, \|\cdot\|_{*,d_\ell}\right) \\ & \leq \left(\frac{2d_\ell d_{\ell-1} \mathcal{R}}{\varepsilon}\right)^{\mathcal{F}_{\ell,w} + \mathcal{F}_{\ell,b}} \mathcal{N}\left(\varepsilon, \vec{\mathcal{H}}_{\ell-1}, \|\cdot\|_{*,d_{\ell-1}}\right). \end{aligned}$$

Scaling ε to $\frac{\varepsilon}{3c_1 (c(1 + 2^{d+1})\mathcal{R})^{\ell-1} d_\ell \cdots d_0}$, we then have

$$\begin{aligned} & \mathcal{N}(\varepsilon, \vec{\mathcal{H}}_\ell, \|\cdot\|_{*,d_\ell}) \leq ((c'_1 \mathcal{R}))^{\ell \mathcal{F}_\ell} D_\ell^{2\mathcal{F}_\ell} \varepsilon^{-\mathcal{F}_\ell} \\ & \times \mathcal{N}\left(\frac{\varepsilon}{(c'_1 \mathcal{R})^{\ell-1} D_\ell}, \vec{\mathcal{H}}_{\ell-1}, \|\cdot\|_{*,d_{\ell-1}}\right), \end{aligned}$$

where $c'_1 = 6c_1c(1 + 2^{d+1})$ and $D_\ell := d_\ell \cdots d_0$. This proves Lemma 3 for $\ell = 2, \dots, L$. If $\ell = 1$, then for arbitrary $\vec{h}_1 \in \vec{\mathcal{H}}_1$, we have

$$\begin{aligned} & \|\vec{h}_1 - \vec{\sigma}_1(W'_1 x + \vec{b}'_1)\|_{*,d_1} \\ & \leq \|\vec{\sigma}_1(W_1 x + \vec{b}_1) - \vec{\sigma}_1(W'_1 x + \vec{b}_1)\|_{*,d_1} \\ & + \|\vec{\sigma}_1(W'_1 x + \vec{b}_1) - \vec{\sigma}_1(W'_1 x + \vec{b}'_1)\|_{*,d_1}. \end{aligned} \quad (27)$$

The same approach as above yields that

$$\{\vec{\sigma}_1(W'_1 x + \vec{b}'_1) : W'_1 \in \mathcal{E}_{1,w}, \vec{b}'_1 \in \mathcal{E}_{1,b}\}$$

is a $c_1 2^{d+1} \varepsilon$ -net of $\vec{\mathcal{H}}_1$. Using Lemma 2 again, we obtain

$$\mathcal{N}(c_1 2^{d+1} \varepsilon, \vec{\mathcal{H}}_1, \|\cdot\|_{*,d_1}) \leq \left(\frac{2D_1 \mathcal{R}}{\varepsilon}\right)^{\mathcal{F}_{1,w} + \mathcal{F}_{1,b}}.$$

Scaling ε to $\varepsilon/c_1 2^{d+1}$, we get

$$\mathcal{N}(\varepsilon, \vec{\mathcal{H}}_1, \|\cdot\|_{*,d_1}) \leq \left(\frac{c'_1 \mathcal{R} D_1}{\varepsilon}\right)^{\mathcal{F}_1}.$$

This completes the proof of Lemma 3. \blacksquare

With the help of the above two lemmas, we are in a position to prove Theorem 1.

Proof of Theorem 1: Let $\vec{\mathcal{A}}_{\mathcal{F}_{L,a}}^*$ be the set of d_L -dimensional vectors with fixed structures and totally $\mathcal{F}_{L,a}$ free parameters. Denote $\vec{\mathcal{A}}_{\mathcal{F}_{L,a}} := \{\vec{a} \in \vec{\mathcal{A}}_{\mathcal{F}_{L,a}}^* : |a_i| \leq \mathcal{R}\}$. Assume that $\mathcal{E}_{L,a}$ is an ε -cover of the set $\vec{\mathcal{A}}_{\mathcal{F}_{L,a}}^*$ under the metric of $\ell_1^{d_L}$. Then, for arbitrary $\vec{a} \in \mathbb{R}^{d_L}$ and $\vec{h}_L \in \vec{\mathcal{H}}_L$ there is a $\vec{a}^* \in \mathcal{E}_{L,a}$ and $\vec{h}_L^* \in \mathcal{E}_{L,h}$ such that

$$\|\vec{a} - \vec{a}^*\|_{\ell_1^{d_L}} \leq \varepsilon, \quad \text{and} \quad \|\vec{h}_L - \vec{h}_L^*\|_{*,d_L} \leq \varepsilon.$$

Note that

$$\begin{aligned} & \|\vec{a} \cdot \vec{h}_L - \vec{a}^* \cdot \vec{h}_L^*\|_{L^1(\mathbb{I}^d)} \leq \|\vec{a} \cdot \vec{h}_L - \vec{a}^* \cdot \vec{h}_L\|_{L^1(\mathbb{I}^d)} \\ & + \|\vec{a}^* \cdot \vec{h}_L - \vec{a}^* \cdot \vec{h}_L^*\|_{L^1(\mathbb{I}^d)}. \end{aligned} \quad (28)$$

Moreover, Lemma 1 shows

$$\begin{aligned} & \|\vec{a} \cdot \vec{h}_L - \vec{a}^* \cdot \vec{h}_L\|_{L^1(\mathbb{I}^d)} \leq \|\vec{a} - \vec{a}^*\|_{\ell_1^{d_L}} \|\vec{h}_L\|_{*,d_L} \\ & \leq (c(1+2^{d+1})\mathcal{R})^L d_{L-1} \cdots d_0 \|\vec{a} - \vec{a}^*\|_{\ell_1^{d_L}} \\ & \leq (c(1+2^{d+1})\mathcal{R})^L d_{L-1} \cdots d_0 \varepsilon \end{aligned}$$

and

$$\begin{aligned} & \|\vec{a}^* \cdot \vec{h}_L - \vec{a}^* \cdot \vec{h}_L^*\|_{L^1(\mathbb{I}^d)} \leq d_L \mathcal{R} \|\vec{h}_L - \vec{h}_L^*\|_{*,d_L} \\ & \leq d_L \mathcal{R} \varepsilon. \end{aligned}$$

Plugging the above estimates into (28), we have

$$\|\vec{a} \cdot \vec{h}_L - \vec{a}^* \cdot \vec{h}_L^*\|_{L^1(\mathbb{I}^d)} \leq (c'_2 \mathcal{R})^L D_L \varepsilon,$$

where $c'_2 = 2c(1+2^{d+1})$. Since Lemma 2 implies

$$\begin{aligned} & \mathcal{N}\left(\frac{\varepsilon}{(c'_2 \mathcal{R})^L D_L}, \vec{\mathcal{A}}_{\mathcal{F}_L, a}, \ell_1^{d_L}\right) \\ & \leq (c'_2 \mathcal{R})^{(L+1)\mathcal{F}_L, a} D_L^{2\mathcal{F}_L, a} \varepsilon^{-\mathcal{F}_L, a}, \end{aligned}$$

there holds

$$\begin{aligned} & \mathcal{N}(\varepsilon, \mathcal{H}_{n, L, \sigma, \mathcal{R}}, L_1(\mathbb{I}^d)) \\ & \leq (c'_2 \mathcal{R})^{(L+1)\mathcal{F}_L, a} D_L^{2\mathcal{F}_L, a} \varepsilon^{-\mathcal{F}_L, a} \\ & \times \mathcal{N}\left(\frac{\varepsilon}{(c'_2 \mathcal{R})^L D_L}, \vec{\mathcal{H}}_L, \|\cdot\|_{*,d_L}\right). \end{aligned} \quad (29)$$

We then use Lemma 3 to estimate the second part of the above term. Let

$$\begin{aligned} B_\ell &:= 2(\max\{c'_1, c'_2\}\mathcal{R})^\ell D_\ell^2 D_{\ell+1}, \ell = 1, 2, \dots, L-1, \\ B_L &:= 2(\max\{c'_1, c'_2\}\mathcal{R})^L D_L^2, \\ B_{L+1} &:= 2(\max\{c'_1, c'_2\}\mathcal{R})^{L+1} D_L^2. \end{aligned} \quad (30)$$

Then, the first estimate of Lemma 3 shows

$$\begin{aligned} & \mathcal{N}(\varepsilon, \vec{\mathcal{H}}_\ell, \|\cdot\|_{*,d_\ell}) \leq B_\ell^{\mathcal{F}_\ell} \varepsilon^{-\mathcal{F}_\ell} \\ & \times \mathcal{N}\left(\frac{\varepsilon}{B_{\ell-1}}, \vec{\mathcal{H}}_{\ell-1}, \|\cdot\|_{*,d_{\ell-1}}\right), \quad \ell = L, \dots, 2. \end{aligned}$$

Using the above inequality iteratively with $\ell = L, L-1, \dots, 2$, we obtain

$$\begin{aligned} & \mathcal{N}\left(\frac{\varepsilon}{(c'_2 \mathcal{R})^L D_L}, \vec{\mathcal{H}}_L, \|\cdot\|_{*,d_L}\right) \\ & \leq \left(\prod_{\ell=2}^{L-1} B_\ell^{\mathcal{F}_\ell}\right) \left(\varepsilon^{-\sum_{\ell=2}^L \mathcal{F}_\ell}\right) \mathcal{N}\left(\frac{\varepsilon}{\prod_{\ell=1}^L B_\ell}, \vec{\mathcal{H}}_1, \|\cdot\|_{*,d_1}\right) \\ & \times B_L^{\mathcal{F}_L} (B_L B_{L-1})^{\mathcal{F}_{L-1}} \cdots (B_L \cdots B_2)^{\mathcal{F}_2} \\ & = \left(\prod_{\ell=2}^{L-1} B_\ell^{\mathcal{F}_\ell}\right) \left(\prod_{\ell=2}^L B_\ell^{\sum_{j=2}^{\ell} \mathcal{F}_j}\right) \varepsilon^{-\sum_{\ell=2}^L \mathcal{F}_\ell} \\ & \times \mathcal{N}\left(\frac{\varepsilon}{\prod_{\ell=1}^L B_\ell}, \vec{\mathcal{H}}_1, \|\cdot\|_{*,d_1}\right) \end{aligned}$$

But the second estimate in Lemma 3 and the definition of B_ℓ yield

$$\mathcal{N}\left(\frac{\varepsilon}{\prod_{\ell=1}^L B_\ell}, \vec{\mathcal{H}}_1, \|\cdot\|_{*,d_1}\right) \leq \left(B_1 \prod_{\ell=1}^L B_\ell\right)^{\mathcal{F}_1} \varepsilon^{-\mathcal{F}_1}.$$

Then,

$$\begin{aligned} & \mathcal{N}\left(\frac{\varepsilon}{(c'_2 \mathcal{R})^L D_L}, \vec{\mathcal{H}}_L, \|\cdot\|_{*,d_L}\right) \\ & \leq \left(\prod_{\ell=1}^{L-1} B_\ell^{\mathcal{F}_\ell}\right) \left(\prod_{\ell=1}^L B_\ell^{\sum_{j=1}^{\ell} \mathcal{F}_j}\right) \varepsilon^{-\sum_{\ell=1}^L \mathcal{F}_\ell} \end{aligned} \quad (31)$$

Inserting the above estimate into (29), we have

$$\begin{aligned} & \mathcal{N}(\varepsilon, \mathcal{H}_{n, L, \sigma, \mathcal{R}}, L_1(\mathbb{I}^d)) \\ & \leq B_{L+1}^{\mathcal{F}_L, a} \prod_{\ell=1}^L B_\ell^{\mathcal{F}_\ell + \sum_{j=1}^{\ell} \mathcal{F}_j} \varepsilon^{-\sum_{\ell=1}^L \mathcal{F}_\ell - \mathcal{F}_L, a}. \end{aligned}$$

It follows from (30) that

$$\max_{1 \leq \ell \leq L+1} B_\ell \leq B_{L+1} D_L.$$

Then,

$$\begin{aligned} & \mathcal{N}(\varepsilon, \mathcal{H}_{n, L, \sigma, \mathcal{R}}, L_1(\mathbb{I}^d)) \\ & \leq (B_{L+1} D_L)^{\mathcal{F}_L, a + (L+1) \sum_{\ell=1}^L \mathcal{F}_\ell} \varepsilon^{-n} \\ & \leq (B_{L+1} D_L)^{(L+1)n} \varepsilon^{-n}. \end{aligned}$$

This together with (30) yields

$$\mathcal{N}(\varepsilon, \mathcal{H}_{n, L, \sigma, \mathcal{R}}, L_1(\mathbb{I}^d)) \leq ((c_3 \mathcal{R})^{L+1} D_L^3)^{(L+1)n} \varepsilon^{-n},$$

where $c_3 = 2 \max\{c'_1, c'_2\}$. This completes the proof of Theorem 1 by noting $D_L \leq D_{\max}^{L+1}$. \blacksquare

APPENDIX B: COVERING NUMBERS AND APPROXIMATION

The main tool in our analysis is a relation between covering numbers and lower bounds of approximation, which is presented in the following theorem.

Theorem 3. *Let $n \in \mathbb{N}$ and $V \subseteq L_1(\mathbb{I}^d)$. For arbitrary $\varepsilon > 0$, if*

$$\mathcal{N}(\varepsilon, V) \leq \tilde{C}_1 \left(\frac{\tilde{C}_2 n^\beta}{\varepsilon}\right)^n \quad (32)$$

with $\beta, \tilde{C}_1, \tilde{C}_2 > 0$, then

$$\text{dist}(Lip^{(r, c_0)}, V, L_1(\mathbb{I}^d)) \geq C'(n \log_2(n+1))^{-r/d}, \quad (33)$$

where

$$\begin{aligned} C' &:= \frac{1}{4} d^{-d/2} \left[32(1 + \beta + 3r/d) \left(\log_2(2\tilde{C}_1) \right. \right. \\ & \quad \left. \left. + 8d^{d/2}(1 + \beta + 3r/d + \tilde{C}_2) \right) + 1 \right]^{-\frac{r}{d}}. \end{aligned}$$

We postpone the proof of Theorem 3 to the end of this section. Theorem 3 shows that to approximate functions in $Lip^{(r, c_0)}$, the capacity of the approximations, measured by the covering number, plays a crucial role. To present the limitations of deep nets, Theorem 3 implies that we only need to estimate their covering numbers. We highlight that Theorem 3 is motivated by [29], in which a relation between the so-called pseudo-dimensions and lower bounds of approximation is established. However, estimating pseudo-dimensions of classes of functions is not so easy, even for shallow nets [31].

To prove Theorem 3, we need the following four technical lemmas. At first, we introduce the definition of the ε -packing number (see [51], [52]) by

$$\mathcal{M}(\varepsilon, V, B) = \max\{m : \exists f_1, \dots, f_m \in B, \|f_i - f_j\|_B \geq \varepsilon, \forall i \neq j\}.$$

We also denote $\mathcal{M}(\varepsilon, V) := \mathcal{M}(\varepsilon, V, L_1(\mathbb{I}^d))$. The following lemma which was proved in [9, Lemma 9.2] establishes a relation between $\mathcal{N}(\varepsilon, V)$ and $\mathcal{M}(\varepsilon, V)$.

Lemma 4. *For arbitrary $\varepsilon > 0$ and $V \subseteq L_1(\mathbb{I}^d)$, there holds*

$$\mathcal{M}(2\varepsilon, V) \leq \mathcal{N}(\varepsilon, V) \leq \mathcal{M}(\varepsilon, V).$$

For arbitrary $N^* \in \mathbb{N}$, denote $E^{(N^*)^d} := \{\epsilon = (\epsilon_1, \dots, \epsilon_{(N^*)^d}) : \epsilon_i \in \{-1, 1\}, 1 \leq i \leq (N^*)^d\}$. The following lemma can be found in [27, P.489] (see also [29, Claim 1]).

Lemma 5. *For arbitrary $N^* \in \mathbb{N}$, there exists a set $G^{(N^*)^d} \subseteq E^{(N^*)^d}$ with $|G^{(N^*)^d}| \geq 2^{(N^*)^d/16}$ such that for any $v, v' \in G^{(N^*)^d}$ with $v \neq v'$, there holds $\|v - v'\|_{\ell_1} \geq (N^*)^d/2$, where $\|v\|_{\ell_1} = \sum_{i=1}^{(N^*)^d} |v_i|$ for $v = (v_1, \dots, v_{(N^*)^d})$ and $|G^{(N^*)^d}|$ denotes the cardinality of $G^{(N^*)^d}$.*

Define $g : \mathbb{R} \rightarrow \mathbb{R}$ such that $\text{supp}(g) \subseteq [-1/\sqrt{d}, 1/\sqrt{d}]^d$, $g(x) = 1$ for $x \in [-1/(2\sqrt{d}), 1/(2\sqrt{d})]^d$ and $g \in \text{Lip}^{(r, c_0 2^{v-1})}$, where $\text{supp}(g)$ denotes the support of g . Partition \mathbb{I}^d by $(N^*)^d$ sub-cubes $\{A_k\}_{k=1}^{(N^*)^d}$ of side length $1/N^*$ and with centers $\{\xi_k\}_{k=1}^{(N^*)^d}$. For arbitrary $x \in \mathbb{I}^d$, define

$$g_k(x) := (N^*)^{-r} g(N^*(x - \xi_k)) \quad (34)$$

and

$$\mathcal{F}_{G^{(N^*)^d}} := \left\{ \sum_{k=1}^{(N^*)^d} \epsilon_k g_k(x) : \epsilon = (\epsilon_1, \dots, \epsilon_{(N^*)^d}) \in G^{(N^*)^d} \right\}. \quad (35)$$

The following lemma shows that $\mathcal{F}_{G^{(N^*)^d}} \subset \text{Lip}^{(r, c_0)}$

Lemma 6. *For arbitrary $N^* \in \mathbb{N}$, we have*

$$\mathcal{F}_{G^{(N^*)^d}} \subset \text{Lip}^{(r, c_0)},$$

where $G^{(N^*)^d}$ is defined in Lemma 5.

Proof: Let $\vec{\alpha} = (\alpha_1, \dots, \alpha_d)$. Denote by

$$f^{(\vec{\alpha})}(x) = \frac{\partial^s f}{\partial x_1^{\alpha_1} \dots \partial x_d^{\alpha_d}}(x)$$

for every $\alpha_j \in \mathbb{N}_0$, $j = 1, \dots, d$ with $\alpha_1 + \dots + \alpha_d = s$. Since

$$\|N^*(x - \xi_k) - N^*(x - \xi_{k'})\| = N^* \|\xi_k - \xi_{k'}\| \geq 1, \quad \forall k \neq k', \quad (36)$$

$N^*(x - \xi_k)$ and $N^*(x - \xi_{k'})$ do not belong to the set $(-1/\sqrt{d}, 1/\sqrt{d})^d$ simultaneously. Then it follows from $\text{supp}(g) \subseteq [-1/\sqrt{d}, 1/\sqrt{d}]^d$ that for arbitrary $x \in \mathbb{I}^d$, there is at most one $k \in \{1, 2, \dots, (N^*)^d\}$ such that $g_k(x) \neq 0$, $g_k^{(\vec{\alpha})}(x) \neq 0$, that is,

$$g_k(x) = 0, \quad g_k^{(\vec{\alpha})}(x) = 0, \quad \text{if } x \in A_{k'} \text{ with } k' \neq k. \quad (37)$$

If $x, x' \in A_{k_0}$ for some $k_0 \in \{1, \dots, (N^*)^d\}$, then $g_k(x) = 0$ for $k \neq k_0$. So, for each $f \in \mathcal{F}_{G^{(N^*)^d}}$, we get from $|\epsilon_k| = 1$, (34) and $g \in \text{Lip}^{(r, c_0 2^{v-1})}$ with $r = s + v$ and $0 < v \leq 1$ that

$$\begin{aligned} |f^{(\vec{\alpha})}(x) - f^{(\vec{\alpha})}(x')| &= \left| \sum_{k=1}^{(N^*)^d} \epsilon_k [g_k^{(\vec{\alpha})}(x) - g_k^{(\vec{\alpha})}(x')] \right| \\ &= |g_{k_0}^{(\vec{\alpha})}(x) - g_{k_0}^{(\vec{\alpha})}(x')| \\ &= (N^*)^{-r+s} \left| [g^{(\vec{\alpha})}(N^*(x - \xi_{k_0})) - g^{(\vec{\alpha})}(N^*(x' - \xi_{k_0}))] \right| \\ &\leq c_0 2^{v-1} \|x - x'\|^v \leq c_0 \|x - x'\|^v. \end{aligned}$$

If $x \in A_{k_1}$ but $x' \in A_{k_2}$ for some $k_1, k_2 \in \{1, \dots, (N^*)^d\}$ with $k_1 \neq k_2$, we can choose $z \in \partial A_{k_1}$ and $z' \in \partial A_{k_2}$ such that z, z' are on the segment between x and x' , where ∂A denotes the boundary of the sub-cube A . Then

$$\|x - z\| + \|x' - z'\| \leq \|x - x'\|.$$

Due to the fact that $\text{supp}(g) \subseteq [-1/\sqrt{d}, 1/\sqrt{d}]^d$, g is smooth on \mathbb{R}^d and (34), we get

$$g_{k_1}^{(\vec{\alpha})}(z) = g_{k_2}^{(\vec{\alpha})}(z') = 0. \quad (38)$$

So, $g \in \text{Lip}^{(r, c_0 2^{v-1})}$ with $0 < v \leq 1$ and Jensen's inequality yield

$$\begin{aligned} |f^{(\vec{\alpha})}(x) - f^{(\vec{\alpha})}(x')| &= \left| \sum_{k=1}^{(N^*)^d} \epsilon_k [g_k^{(\vec{\alpha})}(x) - g_k^{(\vec{\alpha})}(x')] \right| \\ &\leq |g_{k_1}^{(\vec{\alpha})}(x)| + |g_{k_2}^{(\vec{\alpha})}(x')| \\ &= |g_{k_1}^{(\vec{\alpha})}(x) - g_{k_1}^{(\vec{\alpha})}(z)| + |g_{k_2}^{(\vec{\alpha})}(x') - g_{k_2}^{(\vec{\alpha})}(z')| \\ &\leq (N^*)^{d-r} \left[|g^{(\vec{\alpha})}(N^*(x - \xi_{k_1})) - g^{(\vec{\alpha})}(N^*(z - \xi_{k_1}))| \right. \\ &\quad \left. + |g^{(\vec{\alpha})}(N^*(x' - \xi_{k_2})) - g^{(\vec{\alpha})}(N^*(z' - \xi_{k_2}))| \right] \\ &\leq c_0 2^v \left[\frac{\|x - z\|^v}{2} + \frac{\|x' - z'\|^v}{2} \right] \\ &\leq c_0 2^v \left[\frac{\|x - z\|}{2} + \frac{\|x' - z'\|}{2} \right]^v \leq c_0 \|x - x'\|^v. \end{aligned}$$

Both assertions yield $f \in \text{Lip}^{(r, c_0)}$ and proves Lemma 6 ■

The last lemma describes the geometry of $\mathcal{F}_{G^{(N^*)^d}}$.

Lemma 7. *Let $N^* \in \mathbb{N}$ and $G^{(N^*)^d}$ be defined in Lemma 5. For any $f \neq f_1 \in \mathcal{F}_{G^{(N^*)^d}}$, there holds*

$$\|f - f_1\|_{L_1(\mathbb{I}^d)} \geq \frac{1}{2} d^{-d/2} (N^*)^{-r}. \quad (39)$$

Proof: For arbitrary $f, f_1 \in \mathcal{F}_{G^{(N^*)^d}}$ with $f \neq f_1$, it follows from (35) that there exist $\epsilon, \epsilon' \in G^{(N^*)^d}$ with $\epsilon \neq \epsilon'$ such that

$$\|f - f_1\|_{L_1(\mathbb{I}^d)} = \int_{\mathbb{I}^d} \left| \sum_{k=1}^{(N^*)^d} (\epsilon_k - \epsilon'_k) g_k(x) \right| dx. \quad (40)$$

Since $g_k(x) = 0$ for $x \in \partial A_k$, $k = 1, 2, \dots, (N^*)^d$, we get from (40), (34), (37) and $g \in Lip^{(r, c_0 2^{v-1})}$ that

$$\begin{aligned} \|f - f_1\|_{L_1(\mathbb{I}^d)} &= \sum_{k'=1}^{(N^*)^d} \int_{A_{k'}} \left| \sum_{k=1}^{(N^*)^d} (\epsilon_k - \epsilon'_k) g_k(x) \right| dx \\ &= \sum_{k'=1}^{(N^*)^d} \int_{A_{k'}} |(\epsilon_{k'} - \epsilon'_{k'}) g_{k'}(x)| dx \\ &= (N^*)^{-r} \sum_{k'=1}^{(N^*)^d} |\epsilon_{k'} - \epsilon'_{k'}| \int_{A_{k'}} |g(N^*(x - \xi_{k'}))| dx. \end{aligned} \quad (41)$$

For each $k' = 1, 2, \dots, (N^*)^d$, when x runs over $A_{k'}$, $N^*(x - \xi_{k'})$ runs over a cube S centered at $\xi_{k'}$ and with side-length 1. Then, $g(x) = 1$ for $x \in [-1/(2\sqrt{d}), 1/(2\sqrt{d})]^d$ yields

$$\begin{aligned} &\int_{A_{k'}} |g(N^*(x - \xi_{k'}))| dx \\ &= \int_{A_{k'}} |g(N^*(x - \xi_{k'}))| d(x - \xi_{k'}) \\ &\geq (N^*)^{-d} \int_{[-1/(2\sqrt{d}), 1/(2\sqrt{d})]^d} |g(x)| dx \geq (\sqrt{d} N^*)^{-d}. \end{aligned} \quad (42)$$

But Lemma 5 with $\epsilon, \epsilon' \in G^{(N^*)^d}$ shows

$$\sum_{k'=1}^{(N^*)^d} |\epsilon_{k'} - \epsilon'_{k'}| \geq (N^*)^d / 2. \quad (43)$$

Hence, for arbitrary $f, f_1 \in \mathcal{F}_{G^{(N^*)^d}}$, inserting (42) and (43) into (41), we obtain

$$\|f - f_1\|_{L_1(\mathbb{I}^d)} \geq (N^*)^{-r} (\sqrt{d} N^*)^{-d} (N^*)^d / 2 \geq \frac{1}{2} d^{-d/2} (N^*)^{-r}. \quad (44)$$

This completes the proof of Lemma 7. \blacksquare

By the help of the above four lemmas, we are in a position to prove Theorem 3.

Proof of Theorem 3: For arbitrary $\nu > 0$, denote

$$\delta = \text{dist}(\mathcal{F}_{G^{(N^*)^d}}, V, L_1(\mathbb{I}^d)) + \nu. \quad (45)$$

For any $f \in \mathcal{F}_{G^{(N^*)^d}}$, define a function $Pf \in V$ such that

$$\|f - Pf\|_{L_1(\mathbb{I}^d)} \leq \delta. \quad (46)$$

Due to (45), there are more than one Pf satisfying (46). Define $\mathcal{T}_{G^{(N^*)^d}} := \{Pf : f \in \mathcal{F}_{G^{(N^*)^d}}\} \subseteq V$. For arbitrary $f, f_1 \in \mathcal{F}_{G^{(N^*)^d}}$ with $f \neq f_1$, write $f^* = Pf$ and $f_1^* = Pf_1$. Then

$$\begin{aligned} \|f^* - f_1^*\|_{L_1(\mathbb{I}^d)} &= \|Pf - Pf_1\|_{L_1(\mathbb{I}^d)} \\ &= \|Pf - f + f - f_1 + f_1 - Pf_1\|_{L_1(\mathbb{I}^d)} \\ &\geq \|f - f_1\|_{L_1(\mathbb{I}^d)} - \|Pf - f\|_{L_1(\mathbb{I}^d)} - \|Pf_1 - f_1\|_{L_1(\mathbb{I}^d)}, \end{aligned}$$

which together with (39) and (45) shows

$$\|f^* - f_1^*\|_{L_1(\mathbb{I}^d)} \geq \frac{1}{2} d^{-d/2} (N^*)^{-r} - 2\delta. \quad (47)$$

We now claim $\delta > \frac{1}{8} d^{-d/2} (N^*)^{-r}$ for N^* satisfying

$$\begin{aligned} &(N^*)^d = \lceil 32(1 + \beta + 3r/d)n \\ &\times \log_2(2\tilde{C}_1 + 8d^{d/2}(1 + \beta + 3r/d + \tilde{C}_2) + n) \rceil \end{aligned} \quad (48)$$

with $\lceil a \rceil$ denoting the smallest integer not smaller than the positive number a . To prove the claim, suppose, to the contrary, that

$$\delta \leq \frac{1}{8} d^{-d/2} (N^*)^{-r}, \quad (49)$$

then (47) implies

$$\|f^* - f_1^*\|_{L_1(\mathbb{I}^d)} \geq \frac{1}{4} d^{-d/2} (N^*)^{-r}.$$

This shows that $f \neq f_1$ implies $f^* \neq f_1^*$. So it follows from Lemma 5 that

$$|\mathcal{T}_{G^{(N^*)^d}}| = |\mathcal{F}_{G^{(N^*)^d}}| = |G^{(N^*)^d}| \geq 2^{(N^*)^d/16}.$$

Fixing $\varepsilon_0 = \frac{1}{4} d^{-d/2} (N^*)^{-r}$, we then have

$$\mathcal{M}(\varepsilon_0, V) \geq 2^{(N^*)^d/16}.$$

On the other hand, since $\mathcal{T}_{G^{(N^*)^d}} \subseteq V$, it follows from (32) and Lemma 4 that

$$\begin{aligned} \mathcal{M}(\varepsilon_0, V) &\leq \mathcal{N}(\varepsilon_0/2, V) \leq \tilde{C}_1 \left(\frac{2\tilde{C}_2 n^\beta}{\varepsilon_0} \right)^n \\ &= \tilde{C}_1 \left(2\tilde{C}_2 n^\beta 4d^{d/2} (N^*)^r \right)^n. \end{aligned}$$

Combining the above two inequalities, we get

$$2^{(N^*)^d/16} \leq \tilde{C}_1 \left(2\tilde{C}_2 n^\beta 4d^{d/2} (N^*)^r \right)^n. \quad (50)$$

This together with (48) shows

$$\begin{aligned} &2(1 + \beta + 3r/d)n \\ &\times \log_2(2\tilde{C}_1 + 8d^{d/2}(1 + \beta + 3r/d + \tilde{C}_2) + n) \\ &< \log_2(\tilde{C}_1) + n \log_2(\tilde{C}_2 4d^{d/2}) \\ &+ \beta n \log n + \frac{rn}{d} \log_2(32(\beta + 1 + 3r/d)) + \frac{rn}{d} \log_2 n \\ &+ \frac{rn}{d} \log_2 \log_2(2\tilde{C}_1 + 8d^{d/2}(1 + \beta + 3r/d + \tilde{C}_2) + n). \end{aligned} \quad (51)$$

Since the righthand of the above inequality is smaller than

$$(2 + \beta + 3r/d)n \log_2(2\tilde{C}_1 + 8d^{d/2}(1 + \beta + 3r/d + \tilde{C}_2) + n),$$

it leads to a contradiction. This proves the claim $\delta > \frac{1}{8} d^{-d/2} (N^*)^{-r}$ for N^* satisfying (48). Noting for arbitrary $u \geq 2$,

$$\log_2(n + u) \leq \log_2 u + \log_2(n + 1) \leq (\log_2 u + 1) \log_2(n + 1),$$

we have

$$\delta > \frac{1}{8} d^{-d/2} (N^*)^{-r} \geq 2C'(n \log_2(n + 1))^{-r/d}.$$

But (45) with $\nu = \delta/2$ shows

$$\text{dist}(\mathcal{F}_{G^{(N^*)^d}}, V, L_1(\mathbb{I}^d)) = \frac{\delta}{2} > C'(n \log_2(n + 1))^{-r/d}.$$

Therefore, it follows from Lemma 6 that

$$\begin{aligned} \text{dist}(Lip^{(r, c_0)}, V, L_1(\mathbb{I}^d)) &\geq \text{dist}(\mathcal{F}_{G^{(N^*)^d}}, V, L_1(\mathbb{I}^d)) \\ &\geq C'(n \log_2(n + 1))^{-r/d}. \end{aligned}$$

This completes the proof of Theorem 3. \blacksquare

APPENDIX C: PROOF OF THEOREM 2

Combining Theorem 1 and Theorem 3, we can prove Theorem 2 as follows.

Proof of Theorem 2: It suffices to prove Theorem 2 for $p = 1$, since $2^d \|f\|_{L_p(\mathbb{I}^d)} \geq \|f\|_{L_1(\mathbb{I}^d)}$ for arbitrary $f \in L_p(\mathbb{I}^d)$ and $p \geq 1$. Due to Theorem 1, (32) in Theorem 3 is satisfied with $V = \mathcal{H}_{n,L,\gamma,\mathcal{R}}$, $\tilde{C}_1 = 1$, $\beta = 0$ and $\tilde{C}_2 = (c_3 \mathcal{R} D_{\max})^{2(L+1)L}$. Hence, it follows from Theorem 3 that

$$\begin{aligned} & \text{dist}(Lip^{(r,c_0)}, \mathcal{H}_{n,L,\sigma,\mathcal{R}}, L_1(\mathbb{I}^d)) \\ & \geq C' [2\mathcal{A}_L \log_2(2\mathcal{A}_L + 1)]^{-\frac{r}{d}}. \end{aligned}$$

where

$$\begin{aligned} C' &= \frac{1}{4} d^{-d/2} \left[32(1 + 3r/d) \left(\log_2(2 + 8d^{d/2}(1 + 3r/d) \right. \right. \\ & \left. \left. + (c_3 \mathcal{R} D_{\max})^{2(L+1)L} + 1) \right) \right]^{-\frac{r}{d}}. \end{aligned}$$

Since

$$\begin{aligned} & 2 + 8d^{d/2}(1 + 3r/d + (c_3 \mathcal{R} D_{\max})^{2(L+1)L}) \\ & \leq (48d^{d/2} c_3 \mathcal{R} D_{\max})^{2(L+1)L}, \end{aligned}$$

and

$$\log_2(48d^{d/2} c_3 \mathcal{R} D_{\max}) \leq (\log_2(48d^{d/2} c_3) + 1) \log_2(\mathcal{R} D_{\max}),$$

we have

$$C' \geq \bar{C}'_1 [L^2 \log_2(\mathcal{R} D_{\max})]^{-r/d}$$

where $\bar{C}'_1 := \frac{1}{2} [128(1 + 3r/d)(\log_2(48d^{d/2} c_3 + 1))]^{-\frac{r}{d}}$. Therefore,

$$\begin{aligned} & \text{dist}(Lip^{(r,c_0)}, \mathcal{H}_{n,L,\sigma,\mathcal{R}}, L_1(\mathbb{I}^d)) \\ & \geq \bar{C}'_1 [L^2 \log_2(\mathcal{R} D_{\max})]^{-\frac{r}{d}} [2n \log_2(2n + 1)]^{-\frac{r}{d}} \\ & \geq C [L^2 n \log_2 n \log_2(\mathcal{R} D_{\max})]^{-\frac{r}{d}} \end{aligned}$$

with $C = 3^{-r/d} \bar{C}'_1$. This completes the proof of Theorem 2. \blacksquare

ACKNOWLEDGEMENT

The research was supported by the National Natural Science Foundation of China [Grant Nos. 11531013, 61816133, 11571078, 11631015]. Lei Shi is also supported by the Program of Shanghai Subject Chief Scientist (Project No.18XD1400700).

REFERENCES

- [1] H. Adeli and A. Panakktat. A probabilistic neural network for earthquake magnitude prediction. *Neural Networks*, vol. 22, pp. 1018-1024, 2009.
- [2] M. Bianchini and F. Scarselli. "On the complexity of neural network classifiers: a comparison between shallow and deep architectures", *IEEE Trans. Neural Netw. & Learn. Sys.*, vol. 25, pp. 1553-1565, 2014.
- [3] C. K. Chui, X. Li and H. N. Mhaskar. Neural networks for localized approximation. *Math. Comput.*, vol. 63, pp. 607-623, 1994.
- [4] C. K. Chui, X. Li and H. N. Mhaskar. Limitations of the approximation capabilities of neural networks with one hidden layer. *Adv. Comput. Math.*, vol. 5, pp. 233-243, 1996.
- [5] C. K. Chui, S. B. Lin and D. X. Zhou. Construction of neural networks for realization of localized deep learning. *Front. Appl. Math. Stat.*, vol. 4, no. 14, 2018.
- [6] C. K. Chui, S. B. Lin and D. X. Zhou. Deep Neural Networks for Rotation-Invariance Approximation and Learning. Submitted to *Analysis and Applications*.
- [7] O. Delalleau and Y. Bengio. "Shallow vs. deep sum-product networks", *NIPS*, 666-674, 2011.
- [8] I. Goodfellow, Y. Bengio and A. Courville. *Deep Learning*. MIT Press, 2016.
- [9] L. Györfy, M. Kohler, A. Krzyzak and H. Walk. *A Distribution-Free Theory of Nonparametric Regression*. Springer, Berlin, 2002.
- [10] B. Hanin. Universal function approximation by deep neural nets with bounded width and relu activations. *arXiv preprint arXiv:1708.02691*, 2017.
- [11] N. Harvey, C. Liaw and A. Mehrabian. Nearly-tight VC-dimension bounds for piecewise linear neural networks. *Conference on Learning Theory*, 2017: 1064-1068.
- [12] G. E. Hinton, S. Osindero and Y. W. Teh. A fast learning algorithm for deep belief networks. *Neural Comput.*, vol. 18, pp. 1527-1554, 2006.
- [13] V. E. Ismailov. On the approximation by neural networks with bounded number of neurons in hidden layers. *J. Math. Anal. Appl.*, vol. 417, pp. 963-969, 2014.
- [14] M. Kohler and A. Krzyzak. Nonparametric regression based on hierarchical interaction models. *IEEE Trans. Inf. Theory*, vol. 63, pp. 1620-1630, 2017.
- [15] V. N. Konovalov, D. Leviatan and V. E. Maiorov. Approximation by polynomials and ridge functions of classes of s-monotone radial functions. *J. Approx. Theory*, vol. 152, pp. 20-51, 2008.
- [16] V. N. Konovalov, D. Leviatan and V. E. Maiorov. Approximation of Sobolev classes by polynomials and ridge functions. *J. Approx. Theory*, vol. 159, pp. 97-108, 2009.
- [17] V. Kůrková and M. Sanguinetti. Estimates of covering numbers of convex sets with slowly decaying orthogonal subsets. *Discrete Appl. Math.*, vol. 155, pp. 1930-1942, 2007.
- [18] V. Kůrková and M. Sanguinetti. Probabilistic lower bounds for approximation by shallow perceptron networks. *Neural Networks*, vol. 91, pp. 34-41, 2017.
- [19] V. Kůrková. Constructive lower bounds on model complexity of shallow perceptron networks. *Neural Comput. Appl.*, Doi: 10.1007/s00521-017-2965-0.
- [20] A. Krizhevsky, I. Sutskever and G. E. Hinton. Imagenet classification with deep convolutional neural networks. *NIPS*, 2097-1105, 2012.
- [21] Y. LeCun, Y. Bengio and G. Hinton. Deep learning. *Nature*, vol. 521, no. 7553, pp. 436-444, 2015.
- [22] H. W. Lin, M. Tegmark and D. Rolnick. Why does deep and cheap learning works so well? *J. Stat. Phys.*, vol. 168, pp. 1223-1247, 2017.
- [23] S. B. Lin, J. Zeng and X. Chang. Learning rates for classification with Gaussian kernels. *Neural Comput.*, vol. 29, pp. 3353-3380, 2017.
- [24] S. B. Lin. Limitations of shallow nets approximation. *Neural Networks*, vol. 94, pp. 96-102, 2017.
- [25] S. B. Lin. Generalization and expressivity for deep nets. *IEEE Trans. Neural Netw. Learn. Syst.*, In Press.
- [26] T. Lin and H. Zha. Riemannian manifold learning. *IEEE Trans. Pattern Anal. Mach. Intel.*, vol. 30, pp. 796-809, 2008.
- [27] G. G. Lorentz, M. Z. Golitschek and Y. Makovoz. *Constructive Approximation, Advanced Problems*. Springer-Verlag, New York, 1996.
- [28] V. Maiorov and A. Pinkus. Lower bounds for approximation by MLP neural networks. *Neurocomputing*, vol. 25, pp. 81-91, 1999.
- [29] V. Maiorov and J. Ratsaby. On the degree of approximation by manifolds of finite pseudo-dimension. *Constr. Approx.*, vol. 15, pp. 291-300, 1999.
- [30] V. Maiorov and R. Meir. On the near optimality of the stochastic approximation of smooth functions by neural networks. *Adv. Comput. Math.*, vol. 13, pp. 79-103, 2000.
- [31] V. Maiorov. Pseudo-dimension and entropy of manifolds formed by affine invariant dictionary. *Adv. Comput. Math.*, vol. 25, pp. 435-450, 2006.
- [32] Y. Makovoz. Random approximants and neural networks. *J. Approx. Theory*, vol. 85, pp. 98-109, 1996.
- [33] B. McCane and L. Szymanski. Deep radial kernel networks: approximating radially symmetric functions with deep networks. *arXiv preprint arXiv:1703.03470*, 2017.
- [34] L. Meylan and S. Susstrunk. High dynamic range image rendering with a retinex-based adaptive filter. *IEEE Trans. Image Proc.*, vol. 15, pp. 2820-2830, 2006.
- [35] H. Mhaskar, Q. Liao and T. Poggio. Learning real and Boolean functions: When is deep better than shallow. *arXiv preprint arXiv:1603.00988*, 2016.
- [36] H. N. Mhaskar and T. Poggio. Deep vs. shallow networks: An approximation theory perspective, *Anal. Appl.*, vol. 14, pp. 829-848, 2016.

- [37] G. Montúfar, R. pascanu, K. Cho and Y. Bengio. On the number of linear regions of deep neural networks. *Nips*, 2014, 2924-2932.
- [38] P. Petersen and F. Voigtlaender. Optimal approximation of piecewise smooth functions using deep ReLU neural networks, *Neural Networks*, vol. 108, pp. 296-330, 2018.
- [39] T. Poggio, H. Mhaskar, L. Rosasco, B. Miranda and Q. Liao. Why and when can deep-but not shallow-networks avoid the curse of dimensionality: A review. *Intern. J. Auto. Comput.*, DOI: 10.1007/s11633-017-1054-2, 2017.
- [40] D. Rolnick and M. Tegmark. The power of deeper networks for expressing natural functions. *arXiv:1705.05502v1*, 2017.
- [41] I. Safran and O. Shamir. Depth-width tradeoffs in approximating natural functions with neural networks. *arXiv reprint arXiv:1610.09887v2*, 2016.
- [42] C. Satriano, Y. M. Wu, A. Zollo and H. Kanamori. Earthquake early warning: Concepts, methods and physical grounds. *Soil Dynamics Earth. Engineer.*, vol. 31, pp. 106-118, 2011.
- [43] L. Shi, Y. L. Feng and D. X. Zhou. Concentration estimates for learning with l_1 -regularizer and data dependent hypothesis spaces. *Appl. Comput. Harmon. Anal.*, vol. 31, pp. 286-302, 2011.
- [44] L. Shi. Learning theory estimates for coefficient-based regularized regression. *Appl. Comput. Harmon. Anal.*, vol. 34, pp. 252-265, 2013.
- [45] U. Shaham, A. Cloninger and R. R. Coifman, Provable approximation properties for deep neural networks. *Appl. Comput. Harmon. Anal.*, To appear.
- [46] K. Vikraman. A deep neural network to identify foreshocks in real time, *arXiv preprint arXiv:1611.08655*, 2016.
- [47] Z. Wang, A. C. Bovik, H. R. Sheikh and E. P. Simoncelli. Image quality assessment: from error visibility to structural similarity. *IEEE Trans. Image Process.*, vol. 13, pp. 600-612, 2004.
- [48] J. Wright, Y. Ma, J. Mairal, G. Sapiro, T. S. Huang and S. Yan. Sparse representation for computer vision and pattern recognition. *Proc. IEEE*, vol. 98, pp. 1031-1044, 2010.
- [49] D. Yarotsky. Error bounds for approximations with deep ReLU networks. *Neural Networks*, vol. 94, pp. 103-114, 2017.
- [50] G. B. Ye and D. X. Zhou. Learning and approximation by Gaussians on Riemannian manifolds. *Adv. Comput. Math.*, vol. 29, pp. 291-310, 2008.
- [51] D. X. Zhou. The covering number in learning theory. *J. Complex.*, vol. 18, pp. 739-767, 2002.
- [52] D. X. Zhou. Capacity of reproducing kernel spaces in learning theory. *IEEE Trans. Inf. Theory*, vol. 49, pp. 1743-1752, 2003.
- [53] D. X. Zhou. Deep distributed convolutional neural networks: Universality. *Anal. Appl.*, In Press, 2018.
- [54] D. X. Zhou. Universality of Deep Convolutional Neural Networks. *arXiv:1805.10769*, 2018.