Preprint typeset in JHEP style - HYPER VERSION

arXiv:1901.00134v3 [hep-th] 8 Mar 2020

Supersymmetric S-matrices from the worldsheet in 10 & 11d

Yvonne Geyer,

Department of Physics, Faculty of Science, Chulalongkorn University, Thanon Phayathai, Pathumwan, Bangkok 10330, Thailand,
& Institute for Advanced Study, 1 Einstein Drive, 08540 Princeton NJ, USA

Lionel Mason,

Mathematical Institute, University of Oxford, Woodstock Road, Oxford OX2 6GG, UK

Abstract: We obtain compact formulae for tree super-amplitudes for 10 and 11-dimensional supergravity and 10-dimensional supersymmetric Yang-Mills and Born-Infeld. These are based on the *polarised scattering equations*. These incorporate polarization data into a spinor field on the Riemann sphere and arise from a twistorial representation of ambitwistor strings in 10 and 11 dimensions. They naturally extend amplitude formulae to manifest maximal supersymmetry. The framework is the natural generalization of twistorial ambitwistor string formulae found previously in four and six dimensions and is informally motivated from a vertex operator prescription for a family of supersymmetric worldsheet ambitwistor string models.

Contents

1. Introduction **1**

2. 11d supergravity **2**

3. 10d superamplitudes **7**

4. Discussion **12**

1. Introduction

M-theory is approximated by 11d supergravity and is often characterised as the theory that provides the natural geometric backgrounds for supersymmetric membranes. One might therefore expect that supermembranes should be needed to construct amplitudes for 11d supergravity [1] rather than superstrings, whose backgrounds are naturally described by supergravity theories in 10d. However, in this paper we propose formulae for the massless tree-level S-matrix of 11d supergravity based on string theories in ambitwistor space, the space of complex null geodesics. We also explain the analogous framework for 10d superamplitudes.

Ambitwistor strings [2, 3, 4] provide novel formulations of massless quantum field theories that naturally generalize the 4d twistor-strings [5, 6, 7, 8, 9]. They directly yield the remarkable formulae of Cachazo, He and Yuan (CHY), that express tree-level amplitudes as integrals over the moduli space of marked Riemann spheres, that localize on solutions to the *scattering equations* [10, 11]. However, the CHY formulae do not naturally manifest supersymmetry. Fermionic amplitudes are accessible from the Ramond sector of the ambitwistor string [3] and the pure spinor ambitwistor string [12, 13] manifests supersymmetry, but it remains difficult to generate explicit closed-form formulae beyond four points.

In 4d [14][1] and 6d [15], this was remedied by working in a twistorial representation of the model. This naturally manifests supersymmetry giving rise to compact formulae for superamplitudes, manifesting supersymmetry, now localizing on the *polarized scattering equations* that extend the scattering equations to incorporate polarization data.

Here we give the natural extension of these ideas to 10 and 11 dimensions, and present the full, manifestly supersymmetric S-matrix for 11d supergravity and a variety of theories in 10d. Some ingredients have already been presented in the literature: the *tiny group* that

[1] For the 4d case, see also [16] for analagous formulae arising from the original twistor-string; they have twice as many delta functions and more moduli, but are shown to be equivalent to the 4d ambitwistor formulation in §5.2.2. of [17].

leads to the definition of supermomenta in [18] and its links to ambitwistor-strings in twistor coordinates in [19]. The formulae are again localized on the polarized scattering equations. We give the basic structure of ambitwistor string vertex operators in these coordinates and show how they lead to polarized scattering equations in 10 and 11 dimensions. Although we do not give a complete quantization of these models, the structures we obtain provide the necessary ingredients for supersymmetric amplitude formulae. We first set out the 11 dimensional framework for M-theory amplitudes, then the corresponding formulae in 10 dimensions, and explain how to reduce to four dimensions to make contact with [14] providing a proof at least for the lower lying formulae.

2. 11d supergravity

Little groups and tiny groups. In d-dimensions, the little group is $\mathrm{SO}(d-2) \subset \mathrm{SO}(d)$ inside the stabilizer of a null momentum vector, k_μ, $\mu = 0, \ldots, d-1$. Polarization states for massless particles are representations of this little group. Let Γ_μ denote the Clifford matrices, then the null condition gives $(k \cdot \Gamma)^2 = 0$. It is a standard result that the kernel of $k \cdot \Gamma$ is half the dimension of the spin space and can be identified with the spin space of the little group. These little group spinors give, for example, the polarization states for the massless chiral Dirac equation of momentum k.

In 11d, the spin space is 32 dimensional indexed by $\mathfrak{a}, \mathfrak{b} = 1, \ldots 32$, and spinor indices can be raised and lowered with a skew form $\varepsilon_{\mathfrak{a}\mathfrak{b}}$. The kernel of $k \cdot \Gamma$, the spin space for the little group, can be indexed by $\alpha, \beta = 1 \ldots 16$ that can be raised and lowered with a symmetric form $\delta^{\alpha\beta}$. We introduce the basis $\kappa_{\mathfrak{a}\alpha}$ of the kernel of $k \cdot \Gamma$ normalized by [2]

$$\kappa_{\mathfrak{a}\alpha}\kappa^{\alpha}_{\mathfrak{b}} = \Gamma^{\mu}_{\mathfrak{a}\mathfrak{b}} k_\mu \,, \quad \Gamma^{\mathfrak{a}\mathfrak{b}}_{\mu}\kappa_{\mathfrak{a}\alpha}\kappa_{\mathfrak{b}\beta} = -2k_\mu \delta_{\alpha\beta}. \tag{2.1}$$

We take gluon polarization data to be null vectors e_μ with $k \cdot e = 0$. With respect to such a choice, the *tiny group* [18] is the (now complex) $\mathrm{SO}(d-4)$ inside the stabilizer of both k_μ and e_μ. In such a situation we will have a common kernel to $k \cdot \Gamma$ and $e \cdot \Gamma$ as

$$\{e \cdot \Gamma, k \cdot \Gamma\} = k \cdot e \mathbb{I} = 0. \tag{2.2}$$

This joint kernel can be identified with the 8 dimensional spin space of the tiny group, indexed by $a = 1, \ldots, 8$, and we represent its basis by $\epsilon_{\mathfrak{a}a} = \kappa_{\mathfrak{a}\alpha}\epsilon^{\alpha}_{a}$. When e and k are linearly independent, these satisfy the important (semi-) purity relations

$$\Gamma^{\mathfrak{a}\mathfrak{b}}_{\mu}\epsilon_{\mathfrak{a}a}\epsilon_{\mathfrak{b}b} = 2k_\mu \epsilon^{\alpha}_{a}\epsilon_{\alpha b} = 0 \,. \tag{2.3}$$

This follows from using (2.1) and its analogue for e_μ to see that $\Gamma^{\mathfrak{a}\mathfrak{b}}_{\mu}\epsilon_{\mathfrak{a}a}\epsilon_{\mathfrak{b}b}$ is proportional to both k_μ and e_μ, and so must vanish. We also impose the normalizations

$$\epsilon_{\mathfrak{a}a}\epsilon^{a}_{\mathfrak{b}} = \Gamma^{2\,\mu\nu}_{\mathfrak{a}\mathfrak{b}} e_\mu k_\nu, \quad \Gamma^{2\,\mathfrak{a}\mathfrak{b}}_{\mu\nu}\epsilon_{\mathfrak{a}a}\epsilon_{\mathfrak{b}b} = -8\delta_{ab} e_{[\mu}k_{\nu]}, \tag{2.4}$$

where $\Gamma^2_{\mu\nu} = \Gamma_{[\mu}\Gamma_{\nu]}$ etc. as usual.

[2] We follow the conventions of Penrose & Rindler [20] for spinors in higher dimensions.

The polarized scattering equations. We take gravity polarization data to be metric perturbations of the form $\delta g_{\mu\nu} = e_\mu e_\nu \mathrm{e}^{ik\cdot x}$ where e_μ is null, or equivalently $\epsilon_{\mathfrak{a}a}$ or ϵ_a^α satisfying (2.3) and (2.4).

The scattering equations associate n points σ_i on the Riemann sphere to n null momenta $k_{i\mu} \in \mathbb{R}^d$, $i = 1, \dots n$, subject to momentum conservation $\sum_i k_i = 0$. First introduce the meromorphic, Möbius-invariant one-form

$$P_\mu(\sigma) = \sum_i \frac{k_{i\mu}}{\sigma - \sigma_i} \mathrm{d}\sigma \,. \tag{2.5}$$

The scattering equations are n equations on the σ_i, encoding $P^2 = 0$ for all σ:

$$\operatorname*{Res}_{\sigma_i} \frac{1}{2} P^2(\sigma) = k_i \cdot P(\sigma_i) = \sum_j \frac{k_i \cdot k_j}{\sigma_{ij}} = 0 \,. \tag{2.6}$$

Since P is null, we can hope to find $\lambda_\alpha^{\mathfrak{a}}(\sigma)$, satisfying analogues of (2.1),

$$\lambda_{\alpha\mathfrak{a}} \lambda_{\mathfrak{b}}^\alpha = \Gamma^\mu_{\mathfrak{a}\mathfrak{b}} P_\mu \,, \quad \Gamma^{\mathfrak{a}\mathfrak{b}}_\mu \lambda^\alpha_{\mathfrak{a}} \lambda^\beta_{\mathfrak{b}} = -2 P_\mu \delta^{\alpha\beta} \,. \tag{2.7}$$

Since $k \cdot P = 0$, we can again apply the tiny group argument now to k_i and $P(\sigma)$ near σ_i, leading to a joint 8-dimensional kernel of $k \cdot \Gamma$ and $P \cdot \Gamma$. This kernel is spanned by a pair of 8×16 matrices $(u_{a\alpha}, v_{a\alpha})$ subject to the *polarized scattering equations*

$$u_{ia\alpha} \lambda^\alpha_{\mathfrak{a}}(\sigma_i) = v_{ia\alpha} \kappa^\alpha_{i\mathfrak{a}} \,. \tag{2.8}$$

Note that the α-indices on the $v_{ia\alpha}$s are those for the little group for $k_{i\mu}$ whereas that on the $u_{ia\alpha}$ are global little group indices associated to P_μ. The variables $(u_{a\alpha}, v_{a\alpha})$ are defined up to a GL(8)-transformation of the a-indices, and satisfy

$$u_{a\alpha} u_{b\beta} \delta^{\alpha\beta} = 0 \,, \quad v_{a\alpha} v_{b\beta} \delta^{\alpha\beta} = 0 \,, \tag{2.9}$$

so that these subspaces are (semi-)pure. We have the freedom to further normalize against $\epsilon_{\alpha a}$ by

$$\epsilon_{\alpha a} v_b^\alpha = \delta_{ab} \,. \tag{2.10}$$

This reduces the GL(8) freedom on the a-index down to SO(8).[3]

Equation (2.7) implies that $\lambda_{\alpha\mathfrak{a}}$ is a worldsheet spinor. Motivated by the ambitwistor-string model introduced later, we make the Ansatz

$$\lambda_{\mathfrak{a}\alpha}(\sigma) = \sum_{i=1}^n \frac{u_{i\alpha a} \epsilon^a_{i\mathfrak{a}}}{\sigma - \sigma_i} \sqrt{d\sigma} \,, \tag{2.11}$$

where $\epsilon^a_{i\mathfrak{a}}$ is the polarization data for the ith particle.

It is a key fact that for each solution to the scattering equations $k_i \cdot P(\sigma_i) = 0$, with momenta and polarization data in general position, there exists a unique $\lambda_{\mathfrak{a}\alpha}$ satisfying

[3]Further normalization can be done to reduce to the Spin(7) tiny group but we wont use that in the following.

(2.8) and (2.11), [21]. Briefly, this follows from a degree count of the subbundle $E \subset \mathbb{S}_{\mathfrak{a}}$ where $\mathbb{S}_{\mathfrak{a}}$ is the trivial bundle of spinors over $\mathbb{CP}^1$ and E the subbundle that is annihilated by $P^m \gamma_m^{\mathfrak{ab}}$. For each index α, $\lambda_\alpha^{\mathfrak{a}}$ is a section of $E \otimes \mathcal{O}(-1)$. It follows from the defining exact sequences that $E \otimes \mathcal{O}(n-1)$, the bundle in which $\lambda_\alpha^{\mathfrak{a}} \prod_{i=1}^n (\sigma - \sigma_i)$ takes its values, has degree $8n$. However, the ansatz (2.11) imposes 8 conditions per marked point thus reducing the degree to zero. Thus this 16 dimensional bundle is generically trivial with 16 sections. These can then be normalized to satisfy (2.7).

Supersymmetry and the tiny group. The tiny group was introduced in [18] to define supermomenta in higher dimensions, and it was argued that there are natural choices for the ambitwistor-string in [19]. Although there the proposed reduction arises from the null $P(\sigma)$ at σ_i, but in our context this is a pole with residue k_i, which is not independent of k_i, and so does not work directly. We can nevertheless use a variant to introduce supermomenta in the context of our polarized scattering equations (2.8) as follows. On a momentum eigenstate, the supersymmetry generators satisfy $\{Q_{\mathfrak{a}}, Q_{\mathfrak{b}}\} = \Gamma^\mu_{\mathfrak{ab}} k_\mu$. This allows us to define (little-group) Q_α via

$$Q_{\mathfrak{a}} = \kappa_{\mathfrak{a}}^\alpha Q_\alpha \,, \qquad \text{satisfying} \qquad \{Q_\alpha, Q_\beta\} = \delta_{\alpha\beta} \,.$$

The introduction of supermomenta requires the choice of an anticommuting 8 dimensional subspace of the 16 Q_α's. For us, a natural choice arises from the polarization data and solution to the polarized scattering equations $(\epsilon_{i\alpha}^a, v_{i\alpha}^a)$ as these satisfy $v_{i\alpha a} v_{ib}^\alpha = 0 = \epsilon_{i\alpha a}\epsilon_{ib}^\alpha, \epsilon_{i\alpha a} v_{ib}^\alpha = \delta_{ab}$. This however would lead to a supersymmetry representation that depends on the solution to the polarized scattering equations via v_i. Instead, we choose one additional basis spinor ξ_{ia}^α for each particle, such that $(\epsilon_{i\alpha}^a, \xi_{i\alpha}^a)$ satisfy $\xi_{i\alpha a}\xi_{ib}^\alpha = 0$ and $\epsilon_{i\alpha a}\xi_{ib}^\alpha = \delta_{ab}$. We can then define fermionic supermomenta q_i^a by the relations

$$Q_{i\alpha} = \xi_{i\alpha a} q_i^a + \epsilon_{i\alpha}^a \frac{\partial}{\partial q_i^a} \,. \tag{2.12}$$

The 11d supergravity massless multiplet consists of the triplet $(h_{\mu\nu}, C_{\mu\nu\rho}, \psi_\mu^{\mathfrak{b}})$, containing a metric, 3-form potential and Rarita-Schwinger field. The full supermultiplet is then generated from the pure graviton state $(e_\mu e_\nu, 0, 0)$ at $q_i^a = 0$. At $O(q^a)$ we see the 8 components of the Rarita-Schwinger field $\psi_\mu^{\mathfrak{a}} = e_\mu \epsilon_a^{\mathfrak{a}} q^a$ and at $O(q^a q^b)$ the 3-form $C_{\mu\nu\rho} = \Gamma^{\mathfrak{ab}}_{\mu\nu\rho}\epsilon_{\mathfrak{a}}^a \epsilon_{\mathfrak{b}}^b q_a q_b$ and so-on (see §3 for full details of the 10d analogues).

We define the total supersymmetry generator for n particles by

$$Q_{\mathfrak{a}} = \sum_i \kappa_{i\mathfrak{a}}^\alpha Q_{i\alpha} = \sum_i \kappa_{i\mathfrak{a}}^\alpha \left(\xi_{i\alpha a} q_i^a + \epsilon_{i\alpha}^a \frac{\partial}{\partial q_i^a} \right) . \tag{2.13}$$

A clear consistency requirement on supergravity amplitudes is that they must be annihilated by $Q_{\mathfrak{a}}$. We will see that the total dependence of the supergravity superamplitude on the supermomenta in this representation should take the form of an exponential factor e^F, with

$$F = \sum_{i<j}^n \frac{u_{ia\alpha} u_{jb}^\alpha}{\sigma_{ij}} q_i^a q_j^b - \frac{1}{2} \sum_{i=1}^n \xi_{ia\alpha} v_{ib}^\alpha \; q_i^a q_i^b \,. \tag{2.14}$$

We discuss the origin of this factor from a worldsheet model in the next section. Supermomentum conservation is then easily verified,

$$Q_{\mathfrak{a}}\mathrm{e}^F = \left(\sum_i \kappa_{i\mathfrak{a}}^{\alpha} v_{i\alpha a} q_i^a - \sum_j \lambda_{\mathfrak{a}\alpha}(\sigma_j) u_{jb}^{\alpha} q_j^b\right) \mathrm{e}^F = 0\,,$$

with the second equality following from the polarized scattering equations (2.8). This guarantees invariance under supersymmetry provided the q-dependence is encoded in the exponential e^F.

11d SUGRA amplitudes. Our amplitude formulae take the form

$$\mathcal{M}_n = \int_{\mathfrak{M}_{0,n}} \mathrm{d}\mu_{\mathrm{CHY}}\, \mathcal{I}_n\,, \tag{2.15}$$

where the CHY measure on the moduli space $\mathfrak{M}_{0,n}$ of n points σ_i on $\mathbb{CP}^1$ is given by

$$\mathrm{d}\mu_{\mathrm{CHY}} := \frac{\prod_{i=1}^n \bar{\delta}(k_i \cdot P(\sigma_i)) d\sigma_i}{\mathrm{vol}(\mathrm{SL}(2) \times \mathbb{C}^3)}\,, \tag{2.16}$$

with the Möbius transformation quotient defined via the usual Faddeev-Popov methods and the $\mathbb{C}^3$ quotient leading to the removal of three $\bar{\delta}$-functions and a further Faddeev-Popov factor [2]. For 11d supergravity our formula arises simply from

$$\mathcal{I}_n = \det{}' M\, \mathrm{e}^F\,, \tag{2.17}$$

where $M = \begin{pmatrix} A & C \\ -C^t & B \end{pmatrix}$ is the $2n \times 2n$ CHY matrix constructed from our polarization data,

$$A_{ij} = \frac{k_i \cdot k_j}{\sigma_{ij}},\ B_{ij} = \frac{e_i \cdot e_j}{\sigma_{ij}},\ C_{ij} = \begin{cases} \frac{e_i \cdot k_j}{\sigma_{ij}}, & i \neq j \\ -e_i \cdot P(\sigma_i), & i = j\,, \end{cases} \tag{2.18}$$

with $\sigma_{ij} = \sigma_i - \sigma_j$. The reduced determinant is defined as $\det{}' M = \det M_{[ij]}/\sigma_{ij}^2$, where $M_{[ij]}$ is M with rows and columns i, j removed.

For $q_{ia} = 0$, it is clear that our formula reduces to the standard CHY formula for gravity amplitudes. Thus our work provides a natural supersymmetric extension to provide the full 11d supergravity multiplet.

Worldsheet model and vertex operators. To motivate the polarized scattering quations and supersymmetry factors, we introduce here a twistorial ambitwistor string model. A full quantum description of the model is beyond the scope of this letter.

Let us work in an 11d superspace with coordinates $(x^\mu, \theta^{\mathfrak{a}})$. The Green-Schwartz ambitwistor-string for supergravity [2] has the worldsheet action

$$S = \int_\Sigma P_\mu(\bar{\partial} X^\mu - i\theta^{\mathfrak{a}} \bar{\partial} \theta^{\mathfrak{b}} \Gamma^\mu_{\mathfrak{ab}}) + \frac{e}{2} P^2\,, \tag{2.19}$$

Following [19], we solve the $P^2 = 0$ constraint by $2P_\mu \delta^{\alpha\beta} = \lambda^\alpha_{\mathfrak{a}} \lambda^\beta_{\mathfrak{b}} \Gamma^{\mathfrak{a}\mathfrak{b}}_\mu$. We introduce twistors $Z^{\mathfrak{A}} = (\lambda_{\mathfrak{a}}, \mu^{\mathfrak{b}}, \eta)$, $\mathfrak{A} = 1, \ldots, 64|0$ as a supersymmetric extension of spinors for the conformal group SO(13) [20], with a skew inner product

$$\varepsilon_{\mathfrak{A}\mathfrak{B}} Z_1^{\mathfrak{A}} Z_2^{\mathfrak{B}} = \lambda_{1\mathfrak{a}} \mu_2^{\mathfrak{a}} - \mu_1^{\mathfrak{a}} \lambda_{2\mathfrak{a}} + \eta_1 \eta_2 \,, \tag{2.20}$$

that can be used to raise and lower indices. The spinorial representation of the ambitwistor string is then formulated using 16 such twistors $Z^{\mathfrak{A}}_\alpha$, related to space-time via the incidence relations

$$(\mu^{\mathfrak{a}}_\alpha, \eta_\alpha) = \left(\frac{1}{32} \lambda_{\alpha\mathfrak{b}} \left(X^\mu \Gamma^{\mathfrak{a}\mathfrak{b}}_\mu - 16 i \theta^{\mathfrak{a}} \theta^{\mathfrak{b}} \right), \lambda_{\alpha\mathfrak{a}} \theta^{\mathfrak{a}} \right).$$

Using $32 P_\mu = \lambda^\alpha_{\mathfrak{a}} \lambda_{\mathfrak{b}\alpha} \Gamma^{\mathfrak{a}\mathfrak{b}}_\mu$, the Green Schwarz action transforms in twistor coordinates to

$$S = \int_\Sigma Z^{\mathfrak{A}}_\alpha \bar\partial Z^\alpha_{\mathfrak{A}} + A^{\alpha\beta}_M \Gamma^M_{\mathfrak{A}\mathfrak{B}} Z^{\mathfrak{A}}_\alpha Z^{\mathfrak{B}}_\beta \tag{2.21}$$

Here the $Z^{\mathfrak{A}}_\alpha$ are worldsheet spinors, $A^{\alpha\beta}_M$ are $(0,1)$-form gauge fields on the worldsheet, and $\Gamma^M_{\mathfrak{A}\mathfrak{B}}$ are the SO(13) gamma matrices. The $A^{\alpha\beta}_M$ are Lagrange multipliers for the 13d semi-purity constraints $\Gamma^M_{\mathfrak{A}\mathfrak{B}} Z^{\mathfrak{A}}_\alpha Z^{\mathfrak{B}}_\beta = 0$ that follow from the existence of $(x^\mu, \theta^{\mathfrak{a}}, P_\mu)$ such that the incidence relations (2.21) hold [21].

The vertex operators for this model need to reduce to $\bar\delta(k \cdot P)\, \mathrm{e}^{ik\cdot x}$ in the bosonic case. Including supermomenta q^a, we propose[4]

$$V = \int \delta(k \cdot P)\, w \exp\left(\mu^{\mathfrak{a}}_\alpha \epsilon^a_{\mathfrak{a}} u^\alpha_a + \eta_\alpha u^\alpha_a q^a - \frac{1}{2} \xi_{a\alpha} v^\alpha_b \; q^a q^b \right).$$

Here w is an additional worldsheet operator depending on the polarization data whose correlators provide the determinant $\det' M$ as in [2, 4]. This reduces correctly to the bosonic case: using the unique solution (u, v) to the polarized scattering equations and the incidence relations together with (2.23), the argument of the exponential becomes $k \cdot x + \theta^{\mathfrak{a}} \theta^{\mathfrak{b}} k_{\mathfrak{a}\mathfrak{b}} + \theta^{\mathfrak{a}} \kappa^\alpha_{\mathfrak{a}} \xi_{\alpha a} q^a$ as appropriate for a supermomentum eigenstate.

Consider now a path-integral with n vertex operators. The exponentials in the vertex operators can then be taken into the action, providing sources $(\epsilon^a_{i\mathfrak{a}} u^\alpha_{ia}, u^\alpha_{ia} q^a_i)$ in the equations of motion for $(\lambda^\alpha_{\mathfrak{a}}, \eta^\alpha)$

$$\bar\partial(\lambda^\alpha_{\mathfrak{a}}, 2\eta^\alpha) = \sum_i (\epsilon^a_{i\mathfrak{a}} u^\alpha_{ia}, u^\alpha_{ia} q^a_i) \bar\delta(\sigma - \sigma_i) \,, \tag{2.22}$$

The path integral then localizes onto the classical solution

$$\big(\lambda^\alpha_{\mathfrak{a}}(\sigma), \eta^\alpha(\sigma)\big) = \sum_i \left(\frac{\epsilon^a_{i\mathfrak{a}} u^\alpha_{ia}}{\sigma - \sigma_i}, \frac{u^\alpha_{ia} q^a_i}{2(\sigma - \sigma_i)} \right), \tag{2.23}$$

yielding (2.11) as promised. Furthermore, localising on these classical solution with $\mu^{\mathfrak{a}}_\alpha = 0$ leads to the exponential factor in the fermions

$$\exp\left(\sum_i \eta_\alpha(\sigma_i) u^\alpha_{ia} q^a_i - \frac{1}{2} \xi_{ia\alpha} v^\alpha_{ib} \; q^a_i q^b_i \right) = e^F \,,$$

giving the exponential supermomentum factor introduced earlier.

[4]Note here that the weight of u^α_a as a worldsheet co-spinor cancels that of $(\mu^{\mathfrak{a}}_a, \eta_\alpha)$.

3. 10d superamplitudes

Much of the analysis in 11d extends straightforwardly to 10d, both by analogy and dimensional reduction. We redefine the space-time and little-group indices to $\mu = 1, \ldots 10$ and $m = 1, \ldots, 8$, but maintain our spinor conventions. Note that there is no metric on the 10d spin space $\alpha, \beta = 1, \ldots, 16$. The little group is now SO(8) with two types of chiral spinor indices $a = 1, \ldots, 8$ and $\dot{a} = \dot{1}, \ldots, \dot{8}$. The Clifford matrices Γ decompose into chiral Pauli matrices $\gamma_{\mu\alpha\beta}$, $\gamma_\mu^{\alpha\beta}$.

Little and tiny groups in 10d. Denote the basis of the kernel of $k \cdot \gamma_{\alpha\beta}$ by κ_a^α, normalized by

$$\kappa_{a\alpha}\kappa_\beta^a = \gamma_{\alpha\beta}^\mu k_\mu \,, \quad \gamma_\mu^{\alpha\beta}\kappa_\alpha^a\kappa_\beta^b = -2k_\mu\delta^{ab} \,, \tag{3.1}$$

with similar dotted versions for $\kappa_{\dot{a}}^\alpha$. For null polarization vectors e_μ, the joint kernel of $k\cdot\gamma$ and $e\cdot\gamma$ is now 4-dimensional in each chiral spin space,

$$\epsilon_\alpha^A = \kappa_\alpha^a\epsilon_a^A \,, \quad \epsilon_A^\alpha = \kappa^{\dot{a}\alpha}\epsilon_{\dot{a}A} \,, \quad A = 1, \ldots 4 \tag{3.2}$$

where A is a spinor index for the SO(6) tiny group. As in 11d, we impose the normalizations

$$\epsilon_\alpha^A\epsilon_A^\beta = (\gamma^{\mu\nu})_\alpha^\beta\, e_\mu k_\nu \,, \quad \epsilon_\alpha^A\epsilon_A^\beta\,(\gamma_{\mu\nu})_\beta^\alpha = -8\,\delta_A^B\, e_{[\mu}k_{\nu]} \,. \tag{3.3}$$

We now have full purity conditions

$$\epsilon_a^A\epsilon^{aB} = 0, \qquad \epsilon_A^{\dot{a}}\epsilon_{\dot{a}B} = 0, \tag{3.4}$$

following as before because $\gamma_\mu^{\alpha\beta}\epsilon_\alpha^A\epsilon_\beta^B$ is proportional to both k_μ and e_μ and so must vanish.

The polarized scattering equations. On the scattering equations, we decompose $P(\sigma)$ again into spinors λ_α^a via

$$\lambda_{a\alpha}\lambda_\beta^a = \gamma_{\alpha\beta}^\mu P_\mu \,, \qquad \gamma_\mu^{\alpha\beta}\lambda_\alpha^a\lambda_\beta^b = -2P_\mu\delta^{ab} \,, \tag{3.5}$$

together with a similarly normalized $\lambda_{\dot{a}}^\alpha$. Since this is again a worlsheet spinor, we take

$$\lambda_{a\alpha}(\sigma) = \sum_{i=1}^n \frac{u_{iaA}\epsilon_{i\alpha}^A}{\sigma - \sigma_i} \,. \tag{3.6}$$

where $\epsilon_{i\alpha}^A$ is the polarization data for the ith particle. As before, the scattering equations $k \cdot P = 0$ ensure that $k\cdot\gamma$ and $P\cdot\gamma$ share a 4-dimensional kernel, parametrized by a pair of 4×8 matrices (u_{aA}, v_{aA}). These are again subject to the *polarized scattering equations*,

$$u_{iaA}\lambda_\alpha^a(\sigma_i) = v_{iaA}\kappa_{i\alpha}^a \,, \tag{3.7}$$

and similarly $u_i^{\dot{a}A}\lambda_{\dot{a}}^\alpha(\sigma_i) = v_i^{\dot{a}A}\kappa_{i\dot{a}}^\alpha$ for the opposite chirality. The purity conditions

$$u_{aA}u_{bB}\delta^{ab} = 0 \,, \qquad v_{aA}v_{bB}\delta^{ab} = 0 \,, \tag{3.8}$$

ensure that these subspaces are totally null. Moreover, they are dual to the 4-space defined by the polarization data due to the normalization $v_A^a\epsilon_a^B = \delta_A^B$, giving a unique tiny group for each particle. As in 11d, there exists a unique solution (u_{aA}, v_{aA}) for each solution $\{\sigma_i\}$ to the scattering equations.

Supermomenta for Yang-Mills theory. For super Yang-Mills theory, the supersymmetry generators Q_α act on the supermultiplet by

$$Q_\beta(e_\mu, \zeta^\alpha) = \left(\frac{1}{2}\gamma_{\alpha\beta\mu}\zeta^\alpha, \gamma_\beta^{\mu\nu\alpha} e_{[\mu} k_{\nu]}\right). \tag{3.9}$$

These reduce to the little group data $(e_m, \zeta^{\dot{a}} = \kappa_{\dot{a}}^\alpha \zeta^{\dot{a}})$ by $Q_\alpha = \kappa_\alpha^a Q_a$ where $\{Q_a, Q_b\} = \delta_{ab}$ acting by

$$Q_a(e_m, \zeta^{\dot{a}}) = \left(-\frac{1}{2}\gamma_{ma\dot{a}}\zeta^{\dot{a}}, \gamma_a^{m\dot{a}} e_m\right), \tag{3.10}$$

where $\gamma_{ma\dot{a}}$ are 8d gamma matrices that relate the polarization data e_m to ϵ_a^A, $\epsilon_A^{\dot{a}}$ by

$$e_m \gamma_{a\dot{a}}^m = \epsilon_{aA}\epsilon_{\dot{a}}^A, \quad \gamma_{ma\dot{a}}\epsilon_A^a \epsilon^{\dot{a}B} = -2e_m \delta_A^B, \tag{3.11}$$

To construct the supersymmetry representation, we again introduce additional little group spinors ξ_{iA}^a and $\xi_{i\dot{a}}^A$ for each particle, such that

$$\xi_{aA}\xi_{bB}\delta^{ab} = 0, \qquad \xi_A^a \epsilon_a^B = \delta_A^B, \tag{3.12}$$

and similarly for $\xi_{\dot{a}}^A$. The 8d vector ξ_m relates to ξ_A^a and $\xi_{\dot{a}}^A$ via the analogous relations to (3.11), and the ξ_A^a and $\epsilon_A^{\dot{a}}$ further determine 6d γ-matrices by

$$\gamma_{mAB}^{(6)} := \gamma_{ma\dot{a}}\epsilon_{[A}^{\dot{a}}\xi_{B]}^a = \frac{1}{2}\varepsilon_{ABCD}\gamma_{ma\dot{a}}\xi^{\dot{a}C}\epsilon^{aD}. \tag{3.13}$$

We use the polarization data and the solutions to the scattering equations to parametrize the super Yang-Mills multiplet,

$$(e_m, \xi^{\dot{a}}) = (q^4 \xi_m + 2\gamma_{mAB}^6 q^A q^B + e_m, \epsilon_A^{\dot{a}} q^A + \xi^{\dot{a}A} q_A^3). \tag{3.14}$$

Here, q^A are fermionic supermomenta, with $q^4 = \frac{1}{4!}\varepsilon_{ABCD}q^A \dots q^D$ and $q_A^3 = \frac{1}{3!}\varepsilon_{ABCD}q^B q^C q^D$. On these representatives, the supersymmetry generators take the now-familiar form

$$Q_{ia} = \xi_{iaA} q_i^A + \epsilon_{ia}^A \frac{\partial}{\partial q_i^A}. \tag{3.15}$$

The full supermultiplet $(e_{[\mu}k_{\nu]}, \zeta^\alpha)$ is then given by

$$\left(\gamma_{\mu\nu\beta}^\alpha(\xi_{\alpha A}\xi^{\beta A} q^4 + 2\xi_{\alpha A}\epsilon_B^\beta q^A q^B + \epsilon_\alpha^A \epsilon_A^\beta),\ \epsilon_A^\alpha q^A + \xi^{\alpha A} q_A^3\right), \tag{3.16}$$

where $(\xi_{\alpha A}, \xi^{\alpha A}) = (\xi_A^a \kappa_{a\alpha}, \xi^{\dot{a}A}\kappa_{\dot{a}}^\alpha)$, with supersymmetry generators

$$Q_{i\alpha} = \xi_{i\alpha A} q_i^A + \epsilon_{i\alpha}^A \frac{\partial}{\partial q_i^A}. \tag{3.17}$$

For n superparticles, the total supersymmetry generator is again given by $Q_\alpha = \sum_i Q_{i\alpha}$. Motivated by the ambitwistor string model, we define $\eta_a(\sigma)$ in analogy to (2.23). Super

Yang-Mills amplitudes then only depend on the supermomenta q^A via an exponential factor e^{F_1}, where

$$\begin{aligned} F_1 &:= \sum_{i=1}^{n} \eta_a(\sigma_i) u_{iA}^a q_i^A - \frac{1}{2}\sum_{i=1}^{n} \xi_{iaA} v_{iB}^a \; q_i^A q_j^B \\ &= \sum_{i<j} \frac{u_{iA}^a u_{jBa}}{\sigma_{ij}} \, q_i^A q_j^B - \frac{1}{2}\sum_{i=1}^{n} \xi_{iaA} v_{iB}^a \; q_i^A q_j^B \,. \end{aligned} \tag{3.18}$$

In 10 dimensions, we can extend the supersymmetry to $\mathcal{N} = 2$ with an SO(2) R-symmetry, indexed by $I = 1, 2$ with a symmetric metric δ^{IJ}. This doubles the number of supermomenta to q_{iI}^A, and superamplitudes now carry factors of e^{F_2} with

$$F_2 := \sum_{i<j} \frac{u_{iA}^a u_{jBa}}{\sigma_{ij}} \, q_{iI}^A q_j^{BI} - \frac{1}{2}\sum_{i=1}^{n} \xi_{iaA} v_{iB}^a \; q_{iI}^A q_j^{IB} \,. \tag{3.19}$$

Alternatively, we can extend the supersymmetry in a parity invariant way by introducing supersymmetry generators Q^α of the opposite chirality, leading to supermomenta q_{iA} in the conjugate representation of the tiny group. This leads to exponential supersymmetry factors $\exp \tilde{F}_1$, now built out of conjugate $\tilde{u}_{i\dot{a}}^A$s and $\tilde{q}_{iA}$s.

The supersymmetry factors e^{F_1}, $\mathrm{e}^{\tilde{F}_1}$ and e^{F_2} are supersymmetric under Q_α by an identical calculation to the 11d case. Thus, any formula will be supersymmetric if there is no q-dependence in the rest of the integrand.

10d formulae. We can now introduce 10d formulae that are supersymmetric extensions of the CHY formulae of [10, 11]. In these, gravity amplitudes arise as a double copy of Yang-Mills amplitudes. Thus our integrands $\mathcal{I}_n$ in (2.15) are constructed from reduced Pfaffians or determinants of the CHY matrix M in (2.18) or of the submatrix A, as well as the supersymmetric factors e^F,

Super Yang-Mills:	$\mathrm{PT}(\alpha)\ \mathrm{Pf}'M\ e^{F_1}$
Born-Infeld:	$\det' A\ \mathrm{Pf}'M\ e^{F_1}$
IIASupergravity:	$\det' M\ e^{F_1+\tilde{F}_1}$
IIB Supergravity:	$\det' M\ e^{F_2}$
Heterotic Supergravity:	$\det' M\ e^{F_1}$.

We can also define the Einstein-Yang-Mills superamplitudes of heterotic supergravity by using the corresponding Einstein-Yang-Mills integrands of [11]. All formulae are manifestly supersymmetric, and reduce to the correct bosonic amplitudes.

Factorization. For CHY-like amplitudes, the scattering equations relate factorization – a crucial check on any amplitude representation – to behaviour at the boundary of the moduli space $\widehat{\mathfrak{M}}_{0,n}$ of n-points on the Riemann sphere up to Mobius transformations [22]:

$$\partial\widehat{\mathfrak{M}}_{0,n} \simeq \widehat{\mathfrak{M}}_{0,n_L+1} \times \widehat{\mathfrak{M}}_{0,n_R+1} \,. \tag{3.20}$$

The factorization of our formulae here follows very much analogously to the factorization of the analogous 6d formulae as proved in [23] and the reader is referred there for full details

of factorization in a closely analogous context. Parametrizing the moduli space around this boundary divisor by

$$(x - x_L)(\sigma - \sigma_R) = \varepsilon\,, \quad \text{with } x \in \Sigma_L,\, \sigma \in \Sigma_R\,, \tag{3.21}$$

and $\varepsilon \ll 1$, the polarized scattering equations allow us to introduce spinor data at the junction points $u_{RaA}\epsilon^A_{R\alpha} = \sum_{i\in L} u_{iaA}\epsilon^A_{i\alpha}$, such that $\lambda(\sigma)\sqrt{d\sigma}$ descends to the component spheres $\Sigma_{L,R}$. Moreover, since $\lambda(\sigma)\sqrt{d\sigma}$ is invariant under the inversion (3.21), u_i behave as worldsheet spinors of the local bundles at the marked points, $u_{iaA} = i\varepsilon^{1/2}x_{iL}^{-1}w_{iaA}$. Putting this together, the supersymmetry factors e^F factorize as

$$e^{F_N} = \int d^{4N}q_L d^{4N}q_R\, e^{F_N^{(L)}+F_N^{(R)}}\, G(q_L, q_R)\,, \tag{3.22}$$

where the exponential 'gluing factor' G is given by

$$G(q_L, q_R) = \det{}^N\big(\epsilon_L\epsilon_R\big)\, e^{-i(\epsilon_L\epsilon_R)^{-1}_{AB}\, q_L^A q_R^B}\,, \tag{3.23}$$

with $(\epsilon_L\epsilon_R)^{AB} = \epsilon_L^{aA}\epsilon_{Ra}^B$. This is the correct factorization behaviour for the exponential supersymmetry representation: the exponential in G is dictated by supersymmetry invariance, and the norm ensures agreement with the bosonic sum over states. We have thus verified that all supersymmetric amplitudes factorize correctly.

Reduction to 4d. In the following, we check that our formulae reduce to the correct 4d amplitudes, making contact with the ambitwistor representations [14], which are closely related to the twistor string amplitudes [5, 6, 7, 16, 17]. To implement the reduction, denote the 2-component spinor indices by $_A$ and $_{\dot A}$, and replace the six-dimensional SU(4) spinor indices A, B by $I, J = 1, \ldots, 4$, which will now play the role of $SU(4)$ R-symmetry indices. In this notation, 10d spinors decompose in $(4 + 6)$d to

$$\lambda_\alpha = \Big(\lambda_{AI}, \tilde\lambda^I_{\dot A}\Big)\,. \tag{3.24}$$

The gamma matrices and vectors decompose as

$$\gamma_\mu^{\alpha\beta}\lambda_\alpha\lambda_\beta = (\lambda_{AI}\tilde\lambda^I_{\dot A}, \lambda_{A[I}\lambda^A_{J]} + \frac{1}{2}\varepsilon_{IJKL}\tilde\lambda^K_{\dot A}\tilde\lambda^{\dot A L})\,. \tag{3.25}$$

For null 4d momenta such as $k_\mu = (\kappa_A\tilde\kappa_{\dot A}, 0)$, we can perform a global little group normalization $\xi_a = (\xi_I, \xi^I)$ so that 4-momenta $\kappa_A\tilde\kappa_{\dot A}$ and $\lambda_A\tilde\lambda_{\dot A}$ give rise to

$$\kappa^a_\alpha = \begin{pmatrix} 0 & \tilde\kappa_{\dot A}\,\delta^I_J \\ \kappa_A\,\delta^J_I & 0 \end{pmatrix}, \quad \lambda^a_\alpha(\sigma) = \begin{pmatrix} 0 & \tilde\lambda_{\dot A}(\sigma)\,\delta^I_J \\ \lambda_A(\sigma)\,\delta^J_I & 0 \end{pmatrix}. \tag{3.26}$$

Using $+$ and $-$ to denote self-dual and anti-self-dual particles respectively, we find that for $+$ the tiny group index can be normalized to be an upper $SU(4)$ index and for $-$ a lower one,

$$\epsilon^J_{ia} = \big(\epsilon\delta^J_I, 0\big),\ i \in +\,, \qquad \tilde\epsilon_{iaJ} = \big(0, \tilde\epsilon\,\delta^I_J\big),\ i \in -\,, \tag{3.27a}$$

$$\xi^a_{iI} = \frac{1}{\epsilon}\big(0, \delta^J_I\big),\ i \in +\,, \qquad \tilde\xi^{aI}_i = \frac{1}{\tilde\epsilon}\big(\,\delta^I_J, 0\big),\ i \in -\,. \tag{3.27b}$$

where the prefactors of ξ follow from the normalization condition (3.12) and the scalar ϵ and $\tilde{\epsilon}$ are all that's left of the polarization data with our choices. These identifications lead to

$$u_{iI}^a = u_i(0, \delta_I^J)\,,\ i \in +\,, \quad u_i^{aI} = \tilde{u}_i(\delta_J^I, 0)\,,\ i \in - \tag{3.28}$$

with identical expressions for v in place of ξ due to the normalization conditions. With this, (3.6) reduces to (3.26) with λ_A and $\tilde{\lambda}_{\dot{A}}$ given by

$$\lambda_A(\sigma) = \sum_{i\in-} \frac{u_i \epsilon_{iA}}{\sigma - \sigma_i}\,, \quad \tilde{\lambda}_{\dot{A}}(\sigma) = \sum_{p\in+} \frac{\tilde{u}_p \tilde{\epsilon}_{p\dot{A}}}{\sigma - \sigma_p}\,. \tag{3.29}$$

and the polarized scattering equations (3.7) reduce to

$$\Big(\tilde{u}_i \lambda_A(\sigma_i),\ u_i \tilde{\lambda}_{\dot{A}}(\sigma_i)\Big) = \left(\frac{\kappa_{iA}}{\tilde{\varepsilon}_i},\ \frac{\tilde{\kappa}_{i\dot{A}}}{\varepsilon_i}\right), \tag{3.30}$$

subject to $(u_p, v_p) = 0$ for $p \in +$ and $(\tilde{u}_i, \tilde{v}_i) = 0$ for $i \in -$. These are the familiar 4d refined scattering equations of [14] for the N^{k-2}MHV sector, where k denotes the number of negative helicity particles. They have $\left\langle {n-3 \atop k-2} \right\rangle$ solutions, where $\left\langle {P \atop Q} \right\rangle$ denotes the (P, Q) Eulerian number. Summing over all sectors, (3.28) incorporates all $(n-3)!$ solutions of the polarised scattering equations.

On the N^kMHV sector given by (3.28), the 10d supersymmetry generators reduce to the familiar 4d generators $Q_\alpha = (Q_{AI}, \tilde{Q}_{\dot{A}}^I)$, with

$$(Q_{AI}, \tilde{Q}_{\dot{A}}^I) = \begin{cases} \big(\epsilon_{iA} \frac{\partial}{\partial q_i^I},\ \frac{\tilde{\kappa}_{i\dot{A}}}{\epsilon_i}\, q_i^I\big) & i \in - \\ \big(\frac{\kappa_{i\dot{A}}}{\tilde{\epsilon}_i}\, q_{iI},\ \tilde{\epsilon}_{i\dot{A}} \frac{\partial}{\partial q_{iI}}\big) & i \in + \end{cases}. \tag{3.31}$$

Thus the supermomenta are chiral on the self-dual particles and antichiral on the anti-self-dual particles. In this MHV sector we have

$$\eta_a(\sigma) = \frac{1}{2}\left(\sum_{p\in+} \frac{\tilde{u}_p q_{pJ}}{\sigma - \sigma_p}, \sum_i \frac{u_i q_i^J}{\sigma - \sigma_i}\right), \tag{3.32}$$

and the terms proportional $q_i^I q_i^J$ in the F vanish due to $\xi_{iaI} v_{iJ}^a = 0$. The supersymmetry factors then become $\exp F_4^{(4\mathrm{d})}$ for $\mathcal{N} = 4$ super Yang-Mills and $\exp F_8^{(4\mathrm{d})}$ for $\mathcal{N} = 8$ supergravity, with $J = 1, \ldots, \mathcal{N}$ and

$$F_{\mathcal{N}}^{(4\mathrm{d})} = \sum_{\substack{i\in- \\ p\in+}} \frac{u_i \tilde{u}_p\, q_i^J q_{pJ}}{\sigma_{ip}}\,. \tag{3.33}$$

This is a standard representation for supersymmetry in four dimensions, known as the link representation [24].

The integrands can be identified with the 4d integrands of [14] after dimensional reduction [25], with the CHY Pfaffian playing a double role: as the reduced determinant required for the gravity amplitude, as well as the Jacobian from integrating out the u_i's. Thus our formulae reduce correctly to the known 4d formulae.

4. Discussion

Much underpinning theory for these equations is likely to follow analogously to that for the polarized scattering equations in 6d [15, 23] including factorization, BCFW proofs, the existence and uniqueness of solutions, reductions to other theories in $d < 10$, for example to the recent 6d superamplitudes of [15, 26, 27, 28]. In particular, in higher dimensions we can define an analogue of H_{ij}, but each term is now a matrix, $H_{ij}^{ab} = \epsilon_{i\mathfrak{a}}^{a}\epsilon_{j}^{\mathfrak{a}b}$ for $i \neq j$, $H_{ii}^{ab} = -e_i \cdot P(\sigma_i)\delta^{ab}$ in 11d. Thought of as an $n \times n$ matrix with 8×8 matrix entries, it is possible to introduce a reduced quasi-determinants that are equivalent to the CHY reduced determinants and Pfaffians, we plan to follow up with some of these details in due course [21].

Other directions include making contact with the semi-pure spinors of [1], the pure spinor framework in 10d and to extend the formulae to include brane degrees of freedom [19]. The distinction between our constraints and those of [19] is that ours are intended to restrict to pure ambitwistor degrees of freedom, i.e., those of the space of complex null geodesics, whereas, as pointed out in [29], weaker versions of the constraints might allow one to say something more profound about other M-theory degrees of freedom.

In the 11d and 10d ambitwistor string models, it remains a key issue to conduct a full study of the BRST structure of the constraints and associated anomalies. This is of special interest in 10d, where the corresponding RNS model for supergravity is critical. In the spinorial model, we expect that criticality requires a spinorial version of the integrand, similar to the reduced determinant of a matrix H_{ij} forming the integrand in 4d and 6d [15]. We can indeed define an analogue of H_{ij} in $d = 10, 11$, where each entry is now a matrix, $H_{ij}^{ab} = \epsilon_{i\mathfrak{a}}^{a}\epsilon_{j}^{\mathfrak{a}b}$, with the integrand a reduced quasi-determinant, [21].

Acknowledgments

Acknowledgements: We would like to thank Nathan Berkovits, Eduardo Casali, and Max Guillen for useful discussions on 11d. YG gratefully acknowledges support from the CUniverse research promotion project "Toward World-class Fundamental Physics" of Chulalongkorn University (grant reference CUAASC), as well as support from the National Science Foundation Grant PHY-1606531 and the Association of Members of the Institute for Advanced Study (AMIAS). LJM is grateful to the EPSRC for support under grant EP/M018911/1.

References

[1] N. Berkovits, *Towards covariant quantization of the supermembrane*, *JHEP* **09** (2002) 051, [hep-th/0201151].

[2] L. Mason and D. Skinner, *Ambitwistor strings and the scattering equations*, *JHEP* **1407** (2014) 048, [1311.2564].

[3] T. Adamo, E. Casali and D. Skinner, *Ambitwistor strings and the scattering equations at one loop*, *JHEP* **1404** (2014) 104, [1312.3828].

[4] E. Casali, Y. Geyer, L. Mason, R. Monteiro and K. A. Roehrig, *New Ambitwistor String Theories*, *JHEP* **11** (2015) 038, [1506.08771].

[5] E. Witten, *Perturbative gauge theory as a string theory in twistor space*, *Commun.Math.Phys.* **252** (2004) 189–258, [hep-th/0312171].

[6] N. Berkovits, *An Alternative string theory in twistor space for N=4 superYang-Mills*, *Phys.Rev.Lett.* **93** (2004) 011601, [hep-th/0402045].

[7] R. Roiban, M. Spradlin and A. Volovich, *On the tree level S matrix of Yang-Mills theory*, *Phys.Rev.* **D70** (2004) 026009, [hep-th/0403190].

[8] F. Cachazo, L. Mason and D. Skinner, *Gravity in Twistor Space and its Grassmannian Formulation*, 1207.4712.

[9] D. Skinner, *Twistor Strings for N=8 Supergravity*, 1301.0868.

[10] F. Cachazo, S. He and E. Y. Yuan, *Scattering of Massless Particles in Arbitrary Dimensions*, *Phys.Rev.Lett.* **113** (2014) 171601, [1307.2199].

[11] F. Cachazo, S. He and E. Y. Yuan, *Scattering Equations and Matrices: From Einstein To Yang-Mills, DBI and NLSM*, *JHEP* **07** (2015) 149, [1412.3479].

[12] N. Berkovits, *Infinite Tension Limit of the Pure Spinor Superstring*, *JHEP* **1403** (2014) 017, [1311.4156].

[13] H. Gomez and E. Y. Yuan, *N-point tree-level scattering amplitude in the new Berkovits' string*, *JHEP* **04** (2014) 046, [1312.5485].

[14] Y. Geyer, A. E. Lipstein and L. J. Mason, *Ambitwistor Strings in Four Dimensions*, *Phys. Rev. Lett.* **113** (2014) 081602, [1404.6219].

[15] Y. Geyer and L. Mason, *The polarized scattering equations for 6d superamplitudes*, 1812.05548.

[16] E. Witten, *Parity invariance for strings in twistor space*, *Adv.Theor.Math.Phys.* **8** (2004) 779–796, [hep-th/0403199].

[17] Y. Geyer, *Ambitwistor Strings: Worldsheet Approaches to perturbative Quantum Field Theories*, Ph.D. thesis, Oxford U., Inst. Math., 2016. 1610.04525.

[18] R. H. Boels and D. O'Connell, *Simple superamplitudes in higher dimensions*, *JHEP* **06** (2012) 163, [1201.2653].

[19] I. Bandos, *Twistor/ambitwistor strings and null-superstrings in spacetime of D=4, 10 and 11 dimensions*, *JHEP* **09** (2014) 086, [1404.1299].

[20] R. Penrose and W. Rindler, *Spinors and Space-Time: Spinor And Twistor Methods In Space-time Geometry*, vol. 2. Cambridge University Press, 1986.

[21] Y. Geyer and L. Mason *to appear* .

[22] L. Dolan and P. Goddard, *Proof of the Formula of Cachazo, He and Yuan for Yang-Mills Tree Amplitudes in Arbitrary Dimension*, *JHEP* **1405** (2014) 010, [1311.5200].

[23] G. Albonico, Y. Geyer and L. Mason, *Recursion and worldsheet formulae for 6d superamplitudes*, 2001.05928.

[24] S. He, *A Link Representation for Gravity Amplitudes*, *JHEP* **10** (2013) 139, [1207.4064].

[25] Y. Zhang, *CHY formulae in 4d*, *JHEP* **07** (2017) 069, [1610.05205].

[26] M. Heydeman, J. H. Schwarz and C. Wen, *M5-Brane and D-Brane Scattering Amplitudes*, *JHEP* **12** (2017) 003, [1710.02170].

[27] F. Cachazo, A. Guevara, M. Heydeman, S. Mizera, J. H. Schwarz and C. Wen, *The S Matrix of 6D Super Yang-Mills and Maximal Supergravity from Rational Maps*, *JHEP* **09** (2018) 125, [1805.11111].

[28] M. Heydeman, J. H. Schwarz, C. Wen and S.-Q. Zhang, *All Tree Amplitudes of 6D* $(2,0)$ *Supergravity: Interacting Tensor Multiplets and the* $K3$ *Moduli Space*, 1812.06111.

[29] I. Bandos, *On polarized scattering equations for superamplitudes of 11D supergravity and ambitwistor superstring*, *JHEP* **11** (2019) 087, [1908.07482].