

Statistical Modeling of Atmospheric Mean Temperature in sub—Sahel West Africa

Falaiye O. A.^{*}, Sukam Y. M.^{*} and Abimbola O. J.^{**}

^{*}Department of Physics, Faculty of Physical Sciences, University of Ilorin, Nigeria

^{**}Department of Physics, Faculty of Science, Federal University Lafia, Nigeria

Abstract

Atmospheric mean temperature T_m , is a vital parameter in the evaluation of precipitable water vapor (PWV) through the analysis of GPS signal, it is therefore important to have a good way of evaluation of T_m for the eventual accurate evaluation of PWV using GPS. Simple statistical models exist for various regions of the world for the evaluation of T_m using surface temperature T_s , in the form $T_m = aT_s + b$ where a and b are constants. For West Africa, where atmospheric data is usually very scarce, there is a minimal attempt at finding a statistical model for T_m , as a function of T_s . In this work attempt has been made to find such a model using data from the Climate Monitoring Satellite Facilities (CM-SAF) of the European Meteorological Satellites (EUMETSAT). The model derived was found to compare well with that obtained using radiosonde data with root-mean-square error of 1.189 and mean-biased error of 0.0952 between the two models.

Keywords: West Africa, atmospheric mean temperature, precipitable water vapor.

I. INTRODUCTION

Precipitable water vapor (PWV) is an important atmospheric parameter essential in both weather and climatic prediction. Knowledge of the variability of PWV is also very important in astronomy as water vapor interact with the incoming electromagnetic waves in the atmosphere.

GPS meteorology offers a real-time continuous measurement of PWV. From the GPS data, it is possible to estimate the zenith total delay (ZTD). The zenith hydrostatic delay (ZHD), which is the delay error caused by the dry component of the atmosphere could be estimated using models such as the Saastamoinen [7] model. The delay error introduced by the water vapor in the atmosphere, called the zenith wet delay (ZWD) is calculated thus:

$$ZWD = ZTD - ZHD \quad (1)$$

The ZWD is usually transformed into the PWV using the transformation equation as given below

$$PWV = \Pi \times ZWD \quad (2)$$

The transformation constant Π , is given by

$$\Pi = \left[10^{-6} \left(\frac{k_3}{T_m} + k'_2 \right) \rho_v R_v \right]^{-1} \quad (3)$$

here, ρ_v is the liquid water density = 1000 kgm^{-3} , R_v is water vapor gas constant = $461.524 \text{ JK}^{-1} \text{ kg}^{-1}$, $k_3 = 377600 \text{ K}^2 \text{ hPa}$, $k'_2 = 22.1 \text{ KhPa}^{-1}$ and T_m is called weighted atmospheric mean temperature determination of which is very crucial to accurate transformation of ZWD into the PWV.

The weighted mean atmospheric temperature T_m , could be defined as

$$T_m = \frac{\int_{p_2}^{p_1} T d(\ln p)}{\int_{p_2}^{p_1} d(\ln p)} = \frac{\int_z^{\infty} \frac{e}{T} dz}{\int_z^{\infty} \frac{e}{T^2} dz} \quad (4)$$

where p_1 and p_2 are pressures at two different layers of the atmosphere. T is the ambient temperature. e is the vapor pressure while z is the height above the ground. From Eq. 4. T_m could be estimated from the radiosonde data taken at different layers of the atmosphere. However, such radiosonde data are usually not readily available across the globe, especially Africa, hence several attempts have been made to statistically relate T_m with the surface temperature T_s which is readily available as a basic meteorological parameter at any weather observing station. Bevis *et al.*[1] used 8700 radiosonde profiles from 13 stations in the United States of America (USA) to obtain a statistical model of the form $T_m = aT_s + b$. The Bevis *et al.*[1] model was adopted in many studies (e.g., Raju *et al.*[2]; Fernandez *et al.*[3]; Musa *et al.*[4]; Abimbola *et al.*[5]) across the globe despite the fact that it was obtained for the USA region. Mendes *et al.*[6] and Solbrig[8], using radiosonde data, obtained similar linear relation for Germany, Raju *et al.*[2] for Indian sub-continent, Shoji[9] for Japan and Isioye *et al.*[10] for West Africa. Meanwhile, Schuler *et al.*[11], using a numerical weather prediction data, obtained a model which is more global in outlook relating T_m to T_s as shown in Eq. 5

$$T_m = 0.65T_s + 86.9 \quad (5)$$

From satellite data: which is of higher resolution spatio-temporally than the radiosonde data, it is the aim of this work to find a suitable statistical model for the evaluation of atmospheric mean temperature T_m in West African region.

II. METHODOLOGY

i. Study Area

This study was conducted for locations in West African region as shown in Figure 1. The selected locations in West Africa for this research work are: Dakar (14.76 °N Latitude, 17.36 °W Longitude), Conakry (9.64 °N Latitude, 13.57 °W Longitude), Bamako (12.63 °N Latitude, 8.00 °W Longitude), Abidjan (5.36 °N Latitude, 4.00 °W Longitude), Niamey (13.51 °N Latitude, 12.12 °E Longitude), and Abuja (9.07 °N Latitude, 7.39 °E Longitude).




Figure 1. Map of West Africa showing Locations of study area (Encyclopaedia Britannica, 2018).

For ease of analysis of the data sets and climatic consideration, the study locations were further divided into (1) Hinterland, that is those locations in the interior of the study region and (2) Coastal, that is those regions close to the Atlantic Ocean. These divisions are shown in Table 1.

Table 1. *Division of the Study Area into Hinterland and Coastal Regions.*

Station		Country	Longitude	Latitude	Period of data
Dakar	Coastal Region	Senegal	17.366 °W	14.765 °N	2004 – 2016
Conakry		Guinea	13.578 °W	09.641 °N	2004 – 2016
Abidjan		Cote d’ivoire	04.008 °W	05.360 °N	2004 – 2016
Bamako	Hinter- land Region	Mali	08.003 °W	12.639 °N	2004 – 2016
Niamey		Niger	12.125 °E	13.512 °N	2004 – 2016
Abuja		Nigeria	07.399 °E	09.077 °N	2004 – 2016

ii. Data Analysis

The data used in this work was obtained from the archive of the Climate Monitoring Satellite Application Facilities (CM-SAF) of the European Meteorological Satellites (EUMETSAT) at <https://www.cmsaf.eu>. The data was obtained as a monthly average for a time period spanning from 2004 to 2016 in Network Common Data Format (netCDF). The data obtained at six different pressure levels (level1 = 200 *hPa*; level2 = 300 *hPa*; level3 = 500 *hPa*; level4 = 700 *hPa*; level5 = 850 *hPa* and level6 (surface) = 1000 *hPa*). are the specific humidity q_m (*g/kg*) and the ambient temperature T (*K*). The vapor pressure e (in *hPa*) was estimated from the specific humidity using:

$$e = \frac{0.5q_m}{10^{-3}q_m + 0.622} \quad (6)$$

Panoply (v4.3.1), open source software from NOAA was used for the processing of the data while Jupyter notebook, an environment for running Python code, was used for data plotting, curve fittings and other statistical analysis.

Goodness of fit and correlational analysis of the derived empirical equations were done using coefficient of determination (R^2), mean-biased error (*MBE*) and root-mean-square error (*RMSE*).

III. RESULTS AND DISCUSSION

The derived empirical equations relating the mean atmospheric temperature T_m to the surface temperature T_s for each of the locations considered in this work within the West African region are shown in Table 2. All the stations considered show good linear correlations between T_m and T_s Except for Conakry with a below average coefficient of determination. It will be observed from Table 2 that the coastal region generally does not show good linear correlation between T_m and T_s , whereas the hinterland region shows very good linear correlation: also it was observed that the

Table 2. Statistics for each of the stations in West Africa.

Location		Empirical Model $T_m = aT_s + b$		Coefficient of Determination R^2
		a	b	
Dakar	Coastal Region	0.57	123.15	0.575
Conakry		1.19	63.28	0.442
Abidjan		0.87	32.68	0.741
Bamako	Hinterland Region	0.68	89.96	0.646
Niamey		0.81	52.23	0.763
Abuja		0.85	40.88	0.902

ambient surface temperature at the coast is generally much lower than that at the hinterland of West Africa.

Figures 2 and 3 show the combined plot and linear fit to the dataset of coastal and hinterland respectively. For these figures empirical equations Eq. 7 and Eq. 8 were derived for the coastal and hinterland respectively. It will be observed again from Figures 2 and 3 as well as the derived Eqs. 7 and 8 that the model fit to the hinterland data has better correlation than the fit to the coastal data.




Figure 2. Coastal Region data plot and linear fit.

$$T_m = 0.60T_s + 112.73 \quad R^2 = 0.459 \quad (7)$$




Figure 3. Hinterland Region data plot and linear fit.

$$T_m = 0.72T_s + 77.89 \quad R^2 = 0.752 \quad (8)$$

Figure 4 shows a plot of all the combined data for West Africa. The statistical model derived from Figure 4 is given in Eq. 9 with the corresponding coefficient of determination:

$$T_m = 0.617T_s + 108.49 \quad R^2 = 0.709 \quad (9)$$

Isioye *et. al.*[10] using radiosonde data covering 2009 to 2013 for West Africa obtained a corresponding statistical model as given in Eq. 10:

$$T_m = 0.5743T_s + 116.60 \quad R^2 = 0.436 \quad (10)$$

It will be noted that Eq. 9 shows a better statistical performance than Eq. 10, though the two statistical models could be observed to be quite similar. A statistical comparison of Eq. 9 with Eq. 10 yields root-mean-square error (*RMSE*) of 1.189 and mean-biased error (*MBE*) of 0.0953,

further showing that the two statistical models are comparable. Satellite data has wider spatio-temporal coverage than radiosonde data hence, the better coefficient of determination R^2 , observed for Eq. 9 as compared to Eq. 10.




Figure 4. A combined West African data plot and linear fit.

IV. CONCLUSION

Using satellite data from the Satellite Application Facility (CM-SAF) of the EUMETSAT a suitable statistical model has been derived to estimate weighted atmospheric mean temperature T_m . The statistical model derived is a simple linear model of T_m as a function of surface temperature T_s . The derived model was compared with a similar statistical model which had earlier been derived from the radiosonde data in West Africa. The correspondence between the two models were found to be significant.

Acknowledgment

This work was done using data from EUMETSAT's Satellite Application Facility on Climate Monitoring (CM SAF). DOI: 10.5676/EUM_SAF_CM/WVT_ATOVS/V001.

REFERENCES

- [1] Bevis, M., Businger, S., Herring, T. A., Rocken, C., Anthes, R. A. and Ware R. H. (1992). GPS Meteorology: Remote Sensing of Atmospheric Water Vapor using the Global Positioning System. *Journ. Geophys. Res.*, 97. doi:10.1029/92JD01517.
- [2] Raju, C. S., Saha, K., Thampi, B. V. and Parameswaran, K. (2007). Empirical Model for Mean Temperature for Indian Zone and Estimation of Precipitable Water Vapor from ground based GPS Measurements. *Annal of Geophysics*, 25, 1935-1948.
- [3] Fernandez, W., KhaChatryan, V., Tumasyan, A. (2010). Observation of Long range, Near-side Angular Correlations in Proton-proton Collisions at the LHC *Journal of High Energy Physics*, 2010:91.
- [4] Musa, T. A., Amir, S., Othman, R., Ses, S., Omar, K., Abdullah, K., Lim, S. and Rizos, C. (2011). GPS Meteorology in a low Latitude Region: Remote Sensing of Atmospheric Vapor over the Malaysian Peninsula. *Journal of Atmospheric Physics*, 73:2410-2422.
- [5] Abimbola, O. J., O. A. Falaiye and J. Omojola (2017). Estimation of precipitable water vapor in Nigeria using NIGNET GNSS/GPS, NCEP-DOE Reanalysis II and surface meteorological data. *J. Phys. Sci.*, 28(2), 19 – 29. doi: 10.21315/jps2017.28.2.2.
- [6] Mendes, V. B., Prates, G., Santos, L. and Langley, R. B. (2000). An Evaluation of the Accuracy of Models of the Determination of the Weighted Mean Temperature of the Atmosphere. *Proceedings of ION, National Technical Meeting*, Jan. 26-28, 2000, 2.2. Pacific Hotel Disneyland, Anaheim, CA.
- [7] Saastamoinen, J. (1972). Atmospheric correction for the troposphere and stratosphere in radio ranging of satellites. In Henriksen, W. et al. The use of artificial satellites for geodesy, geophysics monograph series, vol. 15. Washington DC: *American Geophysical Union*, 247, <https://doi.org/10.1029/GM015p0247>.
- [8] Solbrig, P. (2000). Untersuchungenuber die Nutzungnumerischer Wettermodellezur wasser dampfbestimmungmitHilfe des Global Positioning System. *Diploma Thesis, Institute of Geodesy and Navigation, University FAF Munich,Germany*, Feb. 2000.
- [9] Shoji, Y. (2010). Accurate Estimation of Precipitable Water Vapor using Ground based GPS Observation Network and its data assimilation into a mesoscale numerical weather prediction model. *PhD Thesis, Kyoto University*.
- [10] Isiye, O. A., L. Combrinck and J. Botai (2016). Modelling weighted mean temperature in the West African region: implications for GNSS meteorology. *Meteorol. Appl.* 23, 614 – 632. doi: 10.1002/met.1584.
- [11] Schuler, T., Posfay, A., Hein, G. W. and Biberger, R. (2001). A global analysis of the mean atmospheric temperature for GPS water vapor estimation. *Inproc. 14th Int. Tech. Meeting Satell. – 558 Atmospheric Effects*, 10NGPS, P.559.