

Atom-only descriptions of the driven-dissipative Dicke model

François Damanet,¹ Andrew J. Daley,¹ and Jonathan Keeling²

¹*Department of Physics and SUPA, University of Strathclyde, Glasgow G4 0NG, United Kingdom.*

²*SUPA, School of Physics and Astronomy, University of St Andrews, St Andrews, KY16 9SS, United Kingdom.*

(Dated: March 29, 2019)

We investigate how to describe the dissipative spin dynamics of the driven-dissipative Dicke model, describing N two-level atoms coupled to a cavity mode, after adiabatic elimination of the cavity mode. To this end, we derive a Redfield master equation which goes beyond the standard secular approximation and large detuning limits. We show that the secular (or rotating wave) approximation and the large detuning approximation both lead to inadequate master equations, that fail to predict the Dicke transition or the damping rates of the atomic dynamics. In contrast, the full Redfield theory correctly predicts the phase transition and the effective atomic damping rates. Our work provides a reliable framework to study the full quantum dynamics of atoms in a multimode cavity, where a quantum description of the full model becomes intractable.

I. INTRODUCTION

Placing ultracold atoms in a high finesse optical cavity provides an ideal platform to study quantum many body physics out of equilibrium. As a many body open quantum system, it also provides a severe test for theoretical modelling, as the problem size scales as the square of the Hilbert space dimension, and the Hilbert space dimension grows exponentially with the number of atoms and cavity modes involved. For this reason, much theoretical work has been restricted to modelling single-mode cavities [1–3], and cases where all atoms behave identically, so that mean-field descriptions can be applied, or permutation symmetry can be exploited. However, to fully explore many body physics one must move beyond mean-field descriptions, and consider multimode optical cavities [4–14]. Modelling such systems beyond a semiclassical approximation is a major challenge. However, a separation of energy scales naturally exists, with fast cavity degrees of freedom coupled to slower atomic motion. This suggests adiabatic elimination could be used to significantly shrink the Hilbert space. In this paper we show this is indeed possible, but to capture the resulting dissipative dynamics of atoms requires Redfield theory.

Ultracold atoms in optical cavities provide a versatile platform to study a wide variety of questions about engineering and controlling many-body non-equilibrium systems. In particular, one can produce controllable coherent atom-cavity interactions by using a Raman driving scheme, where atoms in the cavity scatter light between an external pump laser and the cavity modes [1]. This controllable interaction can be combined with multimode optical cavities, which support degenerate or near-degenerate families of transverse modes. This allows tuning the spatial structure of the interactions. Indeed, experiments have realized this using a tunable-length cavity [8], which can be used both in the non-degenerate [9] and nearly degenerate confocal [11, 13, 14] regime. This has enabled the creation of tunable-range [11] sign-changing [13, 14] interactions between atoms. A wide variety of applications of such multimode cavity exper-

iments have been considered. These include realization of quantum liquid crystalline states [4, 5], simulating dynamical gauge fields and the Meissner effect [15], realization of spin-glass phases [6], and creating of associative memories [7]. Related to this last concept there have also been a number of proposals of information processing using such systems, including proposing alternate routes to realize Hopfield associative memory [10], and to solve specific NP hard problems such as the N -queens problem [12]. Other quantum generalizations of the Hopfield associative memory [16] have also been studied.

Much of the work listed above on multimode cavities has made use of semiclassical equations, describing the amplitude of the cavity modes and the classical spin state of the atoms. Modelling the full quantum dynamics of the multimode system is a significant challenge, as the multimode structures require keeping track of the quantum state of each atom and each mode of light. Moreover, since the system is driven and dissipative, a full quantum description generally requires a density matrix approach, or an equivalent stochastic approach. Since the dynamics of the cavity modes are generally faster than those of the atoms, it would be highly advantageous to eliminate the cavity modes and consider a master equation for the atomic dynamics only.

To explore the properties of different approximations in deriving an atom-only description, we consider a model for which the correct behavior is well known, namely the open Dicke model [1]. This model describes N two-level atoms coupled to a single cavity mode; it has been extensively studied because this model has a ground state transition to a superradiant¹ state [21–23]. While the existence of this ground-state phase transition has his-

¹ Note the use of the term “superradiant” in this paper refers to a ground-state or steady-state of the open system in which there is a macroscopic photon field. This is distinct from the transient superradiance first discussed by Dicke [17] and reviewed by Gross and Haroche [18]. It is also distinct from the steady state superradiant laser [19, 20]. See Kirton *et al.* [3] for further discussion.

torically been questioned [24], more recent works [25–30] suggest such a transition is indeed possible, but with subtleties regarding gauge choice. No such issues however occur when considering the driven-dissipative realization of the Dicke model [3], and indeed the phase transition has been seen experimentally [2]. As a single-mode problem, the behavior of this model is well understood both through mean-field approaches [1, 31, 32], as well as through exact approaches based on permutation symmetry [33–35].

To derive an atom-only master equation, we consider both the cavity mode and the extra-cavity light as forming a structured bath, with a frequency-dependent density of states. Despite this structure, it is nonetheless possible to produce a time-local (i.e., Markovian) equation of motion for the system density matrix, as long as the effective damping rate due to coupling to the bath is smaller than the energy scale over which the bath density of states varies. This holds in the limit of weak enough matter-light coupling, where the Born-Markov approximation holds [36]. A time-local description means memory effects are neglected, allowing for an efficient computation of the atomic dynamics, while capturing the leading order effects of the cavity loss.

Directly integrating out the bath, and using the Born-Markov equation leads to an equation known as the Redfield master equation [37, 38]. Such an equation is not necessarily of Lindblad form [39], and so does not always preserve the positivity of the reduced density matrix for all time [40–42], yielding in some situations negative and/or diverging populations. The equation is of Lindblad form if the system-bath coupling terms all sample the bath at the same frequency, or if the bath has no structure. However in most cases (including the problem we consider), this is not true. In order to overcome this potential positivity violation, it is a common practice to use in addition the secular approximation, introduced by Wangsness and Bloch [43]. This approximation amounts to neglecting the non-resonant transitions induced by the system-bath dynamics, i.e. it removes the coupling between populations and coherences related to states of different energies.

In many cases in quantum optics, this approximation holds very well — the energy (or frequency) differences are very large compared with the dynamical frequency scales for evolution of the system, and so the neglected terms have very high frequencies in the interaction picture. Indeed, in the quantum optics literature the rotating wave approximation is used, neglecting all counter-rotating terms in the system-bath coupling, and this has an effect identical to the secular approximation. The master equation can then be put into the standard form [44] which, following Lindblad [39], guarantees the positivity of the density matrix for all times. However, neglecting the couplings between populations and coherences can have a dramatic effect and completely remove important physical processes. Indeed it is known that, compared to exactly solvable problems, secular master

equations can lead to wrong results where nonsecular Redfield theory gives qualitatively correct behavior [45–48]. In this paper, we show that the secular approximation is also inadequate to describe the dissipative dynamics of the Dicke model in the thermodynamic limit.

In this paper we present a variety of atom-only descriptions for the open Dicke model, in the form of effective master equations. These different forms correspond to making or not making the secular approximation, or making an approximation based on the small ratio of atomic energy vs cavity linewidth (i.e., the large bandwidth limit). We will see that these various approximations significantly modify the attractors of the dynamics, and that only the full Redfield theory correctly captures the known behavior of the driven Dicke model. Moreover, we will show how semiclassical equations derived from the full model capture dissipative processes which are lost if one first writes semiclassical equations of the Dicke model, and then adiabatically eliminates photons. By comparison to known results we demonstrate that we can derive a master equation for the atom-only system which captures all the required dissipative dynamics. This provides a firm foundation for future work to model the atom-only dynamics in multimode cavities, making use of advanced numerical methods [49–53].

The remainder of the article is arranged as follows. In Sec. II we introduce the open Dicke model, and review the well known behavior of this model, both in terms of its steady states, and the dissipative approach to those states in the limit where atomic energies are much smaller than the cavity linewidth. Section III then presents the atom-only equations of motion, and discusses the form that these take with and without various approximations. The results of each of these different approximations are given in Sec. IV, giving the exact solution in some cases, and numerical and analytic approximations for the full (unsecularized) model. Finally, in Sec. V we summarize our results, and discuss some potential future applications enabled by this work.

II. MODEL AND BACKGROUND

A. Raman-driven realization of Dicke model

We consider the Dicke model [21–23], describing N identical two-level atoms collectively coupled to a single-mode lossy cavity [3]. As described in Ref. [1], such a model can be realized as an effective low energy description of atoms with Raman driving. That is, transitions between two low lying atomic states are driven by scattering a pump photon into a cavity mode, or vice versa (see Fig. 1).

In this context, working in the rotating frame of the pump, one realizes the Dicke Hamiltonian, combined with optical losses from the cavity mode [1, 32]. The




FIG. 1. Cartoon of the Dicke model. Left: Many two-level systems placed in a lossy cavity. Right: Raman driving scheme; transitions between two states $|\downarrow\rangle$ and $|\uparrow\rangle$ involve a virtual transition due to scattering between a pump and a cavity photon.

problem is thus described by the master equation:

$$\partial_t \rho = -i[H_{\text{Dicke}}, \rho] + \kappa \mathcal{L}[a], \quad (1)$$

where $\mathcal{L}[X] = 2X\rho X^\dagger - \{X^\dagger X, \rho\}$ describes photon losses with a rate κ and where the Dicke Hamiltonian is given by:

$$H_{\text{Dicke}} = \omega_0 S^z + \omega a^\dagger a + 2g(a + a^\dagger) S^x. \quad (2)$$

The first term describes the level splitting ω_0 of the two low-lying atomic states, where $S^\alpha = (1/2) \sum_{i=1}^N \sigma_i^\alpha$ ($\alpha = x, y, z$) are collective spin operators written in terms of the standard single-spin Pauli operators σ_i^α ($i = 1, \dots, N$). The second term describes the cost of scattering photons into the cavity, where a is the annihilation operator of a cavity photon, and ω the detuning of the cavity mode from the pump frequency. The final term results from the Raman process, leading to an effective interaction between the atoms and the cavity field, where $g = g_0 \Omega / \Delta_a$ in terms of the bare coupling g_0 , the Rabi frequency Ω of the transverse pump and the atomic detuning Δ_a . These definitions are chosen to match the Hamiltonian in Ref. [32], for ease of comparison to the semiclassical results presented there.

B. Review of dynamics of the dissipative Dicke model

For completeness, in this section we briefly summarize the well-known properties of the model described by Eq. (1) and Eq. (2). In the thermodynamic limit (i.e., large S), the behavior of the model is well-described by the semiclassical equations of motion [32]:

$$\begin{aligned} \partial_t \langle S^- \rangle &= -i\omega_0 \langle S^- \rangle + 2ig (\langle a \rangle + \langle a^\dagger \rangle) \langle S^z \rangle, \\ \partial_t \langle S^z \rangle &= 2g (\langle a \rangle + \langle a^\dagger \rangle) \langle S^y \rangle, \\ \partial_t \langle a \rangle &= -(\kappa + i\omega) \langle a \rangle - 2ig \langle S^x \rangle. \end{aligned} \quad (3)$$

Following these equations, one may see this model shows a phase transition between two classes of steady state attractor: normal states, where $\langle S^x \rangle = \langle a + a^\dagger \rangle = 0$, and an ordered state where these expectations are non-zero. This ordered state spontaneously breaks the Z_2

symmetry of the model under the transformation $S^x \rightarrow -S^x, a \rightarrow -a$. By analogy to the ground state phase transition in the Dicke model [21–23], this ordered state is known as a superradiant state. For the open system [1, 32] the transition occurs when $g > g_c$ where $4g_c^2 N = \omega_0(\omega^2 + \kappa^2)/\omega$.

The dynamics in both states is dissipative, i.e., there is damped relaxation towards the given steady state. As discussed in detail in Bhaseen *et al.* [32], this can be characterized by considering the semiclassical equations of motion, and then linearizing around a given steady state. In general this procedure leads to a quartic equation for the eigenvalues, but this equation can be solved in the limit where $\omega_0 \ll \omega, \kappa$. In the normal state, the eigenvalues of this linearized analysis take the form

$$\lambda = -\frac{4\kappa\omega_0\omega g^2 N}{(\omega^2 + \kappa^2)^2} \pm i\omega_0 \sqrt{1 - \left(\frac{g}{g_c}\right)^2}, \quad (4)$$

(see Eq. (18) of Ref. [32]). Thus, for $g < g_c$, the normal state is absolutely stable, but with a decay rate that can be much smaller than the bare cavity loss rate, particularly in the experimentally relevant regime $\omega_0 \ll \kappa$. When the system becomes superradiant, one must linearize around the new superradiant solution. This (using Eqs. (19) and (20) of Ref. [32]) gives instead the eigenvalues:

$$\lambda = -\frac{\kappa\omega_0^2}{\omega^2 + \kappa^2} \pm i\omega_0 \sqrt{\left(\frac{g}{g_c}\right)^4 - 1}, \quad (5)$$

which corresponds to damped oscillations around the steady state.

The results above come from an analysis of the linearized semiclassical equations of the full model i.e., both atomic and photon degrees of freedom, as given in Eq. (3). If one performs adiabatic elimination of the photon mode on these semiclassical equations, i.e. using $\langle a \rangle = -2ig \langle S^x \rangle / (\kappa + i\omega)$ one may note that the other equations depend only on the combination $\langle a \rangle + \langle a^\dagger \rangle = -4g \langle S^x \rangle \omega / (\omega^2 + \kappa^2)$. Inserting this into the other two equations yields purely conservative dynamics of the collective spin expectation $\langle \mathbf{S} \rangle$. Indeed, as noted in [31], this semiclassical spin dynamics corresponds to that from a Lipkin-Meshkov-Glick [54–56] Hamiltonian

$$H = \omega_0 S^z - \frac{4g^2\omega}{\omega^2 + \kappa^2} (S^x)^2. \quad (6)$$

This Hamiltonian does reproduce the existence of a phase transition at the correct g_c , but since the dynamics is purely conservative, this atom-only semiclassical theory cannot describe the correct damped decay toward the steady state attractors. In the following we will see that a correct atom-only semiclassical theory can however be derived by eliminating cavity photons first, and then taking a semiclassical limit.

III. ATOM-ONLY REDFIELD MASTER EQUATION

A. Derivation of the Redfield equation

In this section, we treat the matter-light coupling as a weak system-reservoir coupling and derive a Redfield master equation for the atom-only dynamics. That is, we derive a description purely in terms of atomic operators, which nonetheless captures the effects of the dissipative cavity mode.

In order to formally derive the atom-only equations, it is useful to note that the starting model described in Eq. (1) can also be written as a purely Hamiltonian problem. That is, one could alternatively describe the same system by enlarging the Hilbert space to describe coupling between the cavity mode and a flat bath of extra-cavity radiation modes, $H_{\text{total}} = H_{\text{Dicke}} + \sum_k g_k (a^\dagger A_k + \text{H.c.}) + \mu_k A_k^\dagger A_k$, where the bath spectral density satisfies $\sum_k g_k^2 \pi \delta(\nu_k - \nu) = \kappa(\nu) \equiv \kappa$. In such a description, adiabatic elimination of the cavity modes means regarding both the cavity and extra-cavity modes as forming the bath. By diagonalizing these coupled harmonic oscillators, one finds the density of states for this effective bath, and can use this to write the open-system description of the atom-only problem.

Using such an approach, we perform the standard derivation of the Redfield equation, first dividing $H_{\text{Dicke}} = H_0 + H_1$ and then working in the interaction picture with respect to H_0 . In this interaction picture, the interaction Hamiltonian H_1 takes the form:

$$H_1(t) = gX(t)[S^+(t) + S^-(t)], \quad (7)$$

where $X(t) = a(t) + a^\dagger(t)$, and $S^\pm(t) = S^\pm e^{\pm i\omega_0 t}$ with $S^\pm = S^x \pm iS^y$. The master equation then becomes:

$$\dot{\rho} = - \int_0^t dt' \text{Tr}_B ([H_1(t), [H_1(t'), \rho]]). \quad (8)$$

To evaluate this, we need to find the correlation function of the cavity photons, thus one needs the time dependence of $a(t)$. This can be found either from the Green's function resulting from diagonalizing the cavity and extra-cavity modes, or alternatively by using Heisenberg-Langevin equations [36] for the cavity modes. Using the assumed flat spectral density of the extra-cavity modes finds the cavity photon correlation function:

$$\text{Tr}_B (X(\tau)X(0)\rho_B) = e^{-i\omega\tau - \kappa|\tau|}. \quad (9)$$

Performing the integrals over time after extending the limit of integration to infinity, we can then write the Redfield equation for the density matrix. It is convenient to introduce the quantities:

$$Q_\pm = \frac{g^2}{\kappa + i(\omega \pm \omega_0)}, \quad (10)$$

in terms of which, the Redfield equation in the Schrödinger picture takes the form:

$$\begin{aligned} \dot{\rho} = & -i[\omega_0 S^z, \rho] - \\ & \{Q_+ (S^+ S^+ \rho - S^+ \rho S^+) + Q_- (S^+ S^- \rho - S^- \rho S^+) \\ & Q_+ (S^- S^+ \rho - S^+ \rho S^-) + Q_- (S^- S^- \rho - S^- \rho S^-) \\ & Q_-^* (\rho S^+ S^+ - S^+ \rho S^+) + Q_-^* (\rho S^+ S^- - S^- \rho S^+) \\ & Q_+^* (\rho S^- S^+ - S^+ \rho S^-) + Q_+^* (\rho S^- S^- - S^- \rho S^-)\}. \end{aligned} \quad (11)$$

In writing Eq. (11), we have not made the secular approximation. To make this additional approximation we would neglect those terms which are time-dependent in the interaction picture. In the above equation, it is the terms with two operators S^- or two operators S^+ which oscillate at a frequency $\pm 2\omega_0$ respectively. In the following, we will compare a number of different approximations with and without secularization. To enable this we will introduce a prefactor ξ in front of those terms which are time dependent in the interaction picture, so that $\xi = 0$ corresponds to the secular approximation and $\xi = 1$ to making no approximation.

B. Master equation in operator form

The master equation given in Eq. (11) can be written in a more compact and convenient form:

$$\dot{\rho} = -i[H_{\text{eff}}, \rho] + \sum_{ij} L_{ij} \left(2C_j \rho C_i^\dagger - \{C_i^\dagger C_j, \rho\} \right), \quad (12)$$

$$H_{\text{eff}} = \omega_0 S^z + \sum_{ij} H_{ij} C_i^\dagger C_j. \quad (13)$$

Here the two-component vector C has components $C_{1,2} = S^\pm$. To write the Hamiltonian and Lindblad-Kossakowski matrices H, L it is convenient to first define Q_0 and Q_1 through $Q_\pm = Q_0 \pm Q_1$. We then find:

$$H = Q_0'' \begin{pmatrix} 1 & \xi \\ \xi & 1 \end{pmatrix} + \begin{pmatrix} Q_1'' & i\xi Q_1' \\ -i\xi Q_1' & -Q_1'' \end{pmatrix}, \quad (14)$$

$$L = Q_0' \begin{pmatrix} 1 & \xi \\ \xi & 1 \end{pmatrix} + \begin{pmatrix} Q_1' & i\xi Q_1'' \\ -i\xi Q_1'' & -Q_1' \end{pmatrix}, \quad (15)$$

where Q_i', Q_i'' refers to real and imaginary parts of these quantities.

The quantities Q_0 and Q_1 can be thought of as corresponding to the mean and difference of the quantities arising from the co- and counter-rotating terms in the matter-light coupling. In the following, we will compare this full master equation with the results making a number of commonly used approximations. Specifically, we will consider the limit $Q_1 = 0$, which is relevant when $\omega_0 \ll \omega, \kappa$ (i.e. the large detuning limit), and the limit $\xi = 0$ corresponding to secularization. We next briefly summarize the simplifications that occur to the effective master equation in these various limiting cases:

1. Secularized master equation

In the case $\xi = 0$, the Hamiltonian and Lindblad–Kossakowski matrix both become diagonal. As expected from secularization, this latter has positive entries $Q'_\pm = Q'_0 \pm Q'_1 = g^2\kappa/(\kappa^2 + (\omega \pm \omega_0)^2)$ guaranteeing complete positivity. We find the effective Hamiltonian takes the form of a Lipkin–Meshkov–Glick [54–56] model with XY symmetry

$$H_{\text{eff}} = (\omega_0 - 2Q''_1)S^z + 2Q''_0 [(S^x)^2 + (S^y)^2], \quad (16)$$

accompanied by simple spin raising and lowering rates:

$$\partial_t \rho = -i[H_{\text{eff}}, \rho] + Q'_+ \mathcal{L}[S^+] + Q'_- \mathcal{L}[S^-]. \quad (17)$$

Because the effective Hamiltonian has XY symmetry, it conserves the number of excited spins. This conservation is an expected consequence of secularization, as the interaction picture with respect to $H_0 = \omega_0 S^z$ will give time dependence to any term that is not diagonal in the S^z basis. As we will show below, this means the steady state density matrix is diagonal in the S^z basis, and we find particularly simple steady states arising from the competition of the spin raising and lowering processes.

If we combine this secular limit with the large detuning limit where Q_1 may be neglected, the equation simplifies further, giving equal rates $Q'_\pm = Q'_0$ for spin raising and lowering processes.

2. Large detuning limit

If we consider the limit where Q_1 may be neglected, but avoiding secularization (so $\xi = 1$), we also find a simple form of the master equation. In this case:

$$H_{\text{eff}} = \omega_0 S^z + 4Q''_0 (S^x)^2, \quad (18)$$

$$\partial_t \rho = i[H_{\text{eff}}, \rho] + 4Q'_0 \mathcal{L}[S^x]. \quad (19)$$

In this case, despite the lack of secularization, we still find a completely positive master equation [41]. This is not surprising, as dropping Q_1 corresponds to neglecting the energy differences $\pm\omega_0$, so that all operators sample the bath at the same frequency. As also discussed further below, this equation also has a simple steady state — since the jump operator is Hermitian, the steady state is a fully mixed density matrix.

3. Full model

While the two limiting cases mentioned above lead to completely positive master equations, this is not true for the full model. We may see this directly by considering the eigenvalues of the Lindblad–Kossakowski matrix L . Specifically, the eigenvalues of this matrix are $Q'_0 \pm \sqrt{(Q'_0)^2 + |Q_1|^2}$ which indeed involves a negative eigenvalue for any non-zero Q_1 . However, as we will show

below, despite this non-positivity, this full master equation is capable of describing the known behavior of the open Dicke model. This is in contrast to both the limiting cases which cannot reproduce the known behavior at the superradiance transition.

C. Master equation in the Dicke basis

Since the master equation is written only in terms of collective spin operators, it is convenient to write the master equation in the Dicke basis spanned by the Dicke states $|SM\rangle$ with $S = N/2$ and $M = -S, \dots, S$ which satisfy:

$$S^z |SM\rangle = M |SM\rangle \quad S^\pm |SM\rangle = f_\pm^M |SM \pm 1\rangle, \quad (20)$$

where $f_\pm^M = \sqrt{(S \mp M)(S \pm M + 1)}$. In this Dicke basis, the density matrix can be decomposed as

$$\rho(t) = \sum_{M=-S}^S \rho_{M,M'}(t) |SM\rangle \langle SM'|, \quad (21)$$

with $(N+1)^2$ matrix elements given by $\rho_{M,M'}(t) = \langle SM | \rho(t) | SM' \rangle$. Noting that $f_-^M = f_+^{M-1}$ we use only f_+^M and suppress the subscript + from hereon. The master equation (11) for these matrix elements reads

$$\begin{aligned} \dot{\rho}_{M,M'} = & -i\omega_0(M - M')\rho_{M,M'} - \\ & \left\{ \xi Q_+ \left(f^{M-1} f^{M-2} \rho_{M-2,M'} - f^{M-1} f^{M'} \rho_{M-1,M'+1} \right) + \right. \\ & Q_- \left(f^{M-1} f^{M-1} \rho_{M,M'} - f^M f^{M'} \rho_{M+1,M'+1} \right) + \\ & Q_+ \left(f^M f^M \rho_{M,M'} - f^{M-1} f^{M'-1} \rho_{M-1,M'-1} \right) + \\ & \xi Q_- \left(f^M f^{M+1} \rho_{M+2,M'} - f^M f^{M'-1} \rho_{M+1,M'-1} \right) + \\ & \xi Q_-^* \left(f^{M'+1} f^{M'} \rho_{M,M'+2} - f^{M-1} f^{M'} \rho_{M-1,M'+1} \right) + \\ & Q_-^* \left(f^{M'-1} f^{M'-1} \rho_{M,M'} - f^M f^{M'} \rho_{M+1,M'+1} \right) + \\ & Q_+^* \left(f^{M'} f^{M'} \rho_{M,M'} - f^{M-1} f^{M'-1} \rho_{M-1,M'-1} \right) + \\ & \left. \xi Q_+^* \left(f^{M'-2} f^{M'-1} \rho_{M,M'-2} - f^M f^{M'-1} \rho_{M+1,M'-1} \right) \right\}. \end{aligned} \quad (22)$$

D. Atom-only semiclassical dynamics

In the following sections, to understand the behavior in the thermodynamic limit, it is useful to write semiclassical equations of motion, found by writing equations for $\langle S^\alpha \rangle$ and then replacing $\langle S^\alpha S^\beta \rangle \rightarrow \langle S^\alpha \rangle \langle S^\beta \rangle$. It is generally easier to extract the equation of motion directly from Eq. (11), but introducing the factors of ξ . Considering a

general operator A we find:

$$\begin{aligned} \partial_t \langle A \rangle = & -i\omega_0 \langle [A, S^z] \rangle - \left\langle Q_+ [A, \xi S^+ + S^-] S^+ \right. \\ & + Q_- [A, S^+ + \xi S^-] S^- + Q_-^* S^+ [\xi S^+ + S^-, A] \\ & \left. + Q_+^* S^- [S^+ + \xi S^-, A] \right\rangle. \end{aligned} \quad (23)$$

We need only consider equations for $\langle S^z \rangle$ and $\langle S^- \rangle$, since $\langle S^+ \rangle$ follows by complex conjugation. We thus find:

$$\begin{aligned} \partial_t \langle S^z \rangle = & - \left\langle Q_+ (\xi S^+ - S^-) S^+ + Q_- (S^+ - \xi S^-) S^- \right. \\ & \left. + Q_-^* S^+ (-\xi S^+ + S^-) + Q_+^* S^- (-S^+ + \xi S^-) \right\rangle. \end{aligned} \quad (24)$$

and

$$\begin{aligned} \partial_t \langle S^- \rangle = & -i\omega_0 \langle S^- \rangle + 2 \left\langle Q_+ \xi S^z S^+ + Q_- S^z S^- \right. \\ & \left. - Q_-^* S^+ \xi S^z - Q_+^* S^- S^z \right\rangle. \end{aligned} \quad (25)$$

Further simplification of these equations depends on whether $\xi = 0$ or $\xi = 1$.

IV. SPIN DYNAMICS OF ATOM-ONLY MODEL

Having introduced the general model in the previous section, in this section, we analyse the dissipative spin dynamics of this model and each of its limiting cases. In several of these cases, we can exactly solve the model in closed form.

A. Secularized master equation

For the secularized case, $\xi = 0$, the populations $\rho_{M,M}(t)$ are decoupled from the coherences $\rho_{M,M' \neq M}(t)$, as can be seen from Eq.(22). Physically, this corresponds to the fact that the Hamiltonian is diagonal in the S^z basis, and the dissipative terms only create or destroy excitations. The equations for the populations (i.e. diagonal elements, $P_M = \rho_{M,M}$) read:

$$\begin{aligned} \dot{P}_M = & 2Q_-' \left[(S+M+1)(S-M)P_{M+1} \right. \\ & \left. - (S+M)(S-M+1)P_M \right] \\ & + 2Q_+' \left[(S+M)(S-M+1)P_{M-1} \right. \\ & \left. - (S+M+1)(S-M)P_M \right]. \end{aligned} \quad (26)$$

One may solve this explicitly for the steady state using a detailed balance condition, where gain and loss terms must be equilibrated [57]. The only consistent way of doing this consists in equating the 1st and 4th terms. This implies the 2nd and 3rd must then also match, and moreover, one may see that the 2nd and 3rd terms relate

to the 1st and 4th by replacing $M \rightarrow M-1$. Thus, the only required condition is $P_{M+1}/P_M = Q_+'/Q_-'$. This condition is identical to that for a thermal equilibrium magnet of moment μ in a Zeeman magnetic field B , for which $P_M \propto \exp(M\beta\mu B)$, but with the replacement $\beta\mu B \rightarrow \ln(Q_+'/Q_-')$. We can thus use the standard results of such a model and obtain $\langle S^z \rangle = SB_S(x)$, where $x = S \ln(Q_+'/Q_-')$ and $B_S(x)$ is the Brillouin function [58]:

$$B_S(x) = \frac{2S+1}{2S} \coth\left(\frac{2S+1}{2S}x\right) - \frac{1}{2S} \coth\left(\frac{1}{2S}x\right). \quad (27)$$

In the limit $S \rightarrow \infty$ the existence of the factor of S in the argument of the Brillouin function means there is a sharp dependence of the result on the ratio Q_+'/Q_-' . Namely, $\langle S^z \rangle = \pm S$ depending on whether $Q_+' > Q_-'$ or vice versa. Intermediate values of S only occur when $|Q_+'/Q_-' - 1| \lesssim 1/S$, a vanishing region at large S .

We thus see that in this approach, we never describe a superradiant state, but instead have a state purely determined by the ratio of spin flip rates. This is consistent with our observation from the effective Hamiltonian: the Hamiltonian conserved number of excitations, so could not modify the effects of gain or loss. It is notable that if one considered the effective Hamiltonian on its own, the ground state of this Lipkin-Meshkov-Glick model does have a ground-state phase transition when Q_0' is negative. That is, for $2|Q_0''|N > \omega_0 - 2Q_1'$, there is a transition to a state with a finite component of spin in the xy plane. The effects of dissipation however destroy this transition, and leave only a transition between states aligned along $+z$ and $-z$ axes.

One may also verify that in this limit, the semiclassical equations (24) and (25) support this result. The equations for $\langle S^- \rangle$ becomes

$$\partial_t \langle S^- \rangle = i\omega_0 \langle S^- \rangle + 2 \left\langle Q_- S^z S^- - Q_+^* S^- S^z \right\rangle. \quad (28)$$

which we may rewrite by symmetrizing expressions in the second term as:

$$\partial_t \langle S^- \rangle = -(i\omega_0 + Q_- - Q_+^*) \langle S^- \rangle + 2(Q_- - Q_+^*) \langle S^z \rangle \langle S^- \rangle. \quad (29)$$

This can be seen to describe overdamped oscillations of $\langle S^- \rangle$. Regardless of $\langle S^z \rangle$, we always find the steady state obeys $\langle S^- \rangle = 0$. The equation for $\langle S^z \rangle$ similarly becomes

$$\partial_t \langle S^z \rangle = 2(Q_+' - Q_-') (S^2 - \langle (S^z)^2 \rangle) - 2(Q_+' + Q_-') \langle S^z \rangle. \quad (30)$$

In the large S limit, the second term can be neglected, and we find the only steady state is $\langle S^z \rangle = \pm S$, corresponding to the sharp switch noted above.

If we combine the secular limit with the large detuning limit, where $Q_{\pm} = Q_0$, we immediately find all probabilities must be equal and so normalization implies $P_M = 1/(N+1)$, i.e. a fully mixed state. Thus, in this case $\langle S^z \rangle = 0$ independent of all parameters.

B. Large detuning unsecularized master equation

In the large detuning limit $Q_{\pm} = Q_0, Q_1 = 0$ (but without secularization), the decoupling of populations and coherences no longer occurs, i.e. the terms in Eq. (22) with $M' = M$ couple to other terms with $M' \neq M$. However, the form of the Master equation in Eq. (19) suggests the solution nonetheless remains straightforward. Namely, since the jump operator in the master equation is Hermitian, a general result [36] states that for Hermitian jump operators, $\rho \propto \mathbb{1}$ is a steady state². In such a state we find all expectations of $\langle S^{\alpha} \rangle$ vanish, and there is no transition as a function of parameters, as obtained in [59].

C. Full model

If we consider the full model with $\xi = 1$ and $Q_1 \neq 0$, no simple solution exists. Nevertheless, the total number of connected density matrix elements containing the populations is $\lceil (n+1)^2/2 \rceil$ (since Eq. (22) connects matrix elements according to a checkerboard pattern), i.e. grows only quadratically with the number of spins, which makes it possible to solve the equations numerically for moderate N . In the large N limit, we may also use the semiclassical equations of motion to obtain an analytic expression of the expectation value of the collective spin. In this thermodynamic limit, we find, in contrast to the previous two cases, that there is a superradiant transition.

In the case $\xi = 1$ the semiclassical equations simplify considerably, since we may in general write:

$$\begin{aligned} \partial_t \langle A \rangle &= -i\omega_0 \langle [A, S^z] \rangle - 2 \langle [A, S^x] (Q_+ S^+ + Q_- S^-) \rangle \\ &\quad - 2 \langle (Q_+ S^+ + Q_- S^-)^\dagger [S^x, A] \rangle. \end{aligned} \quad (31)$$

In this case it is clearest to write equations for $\langle S^{\alpha=x,y,z} \rangle$ explicitly, rather than for $\langle S^{\pm} \rangle$. If we symmetrize all products of operators before taking expectations (i.e. writing $\langle AB \rangle = \frac{1}{2} \langle \{A, B\} + [A, B] \rangle = \langle A \rangle \langle B \rangle + \frac{1}{2} \langle [A, B] \rangle$), we then find the following equations:

$$\partial_t \langle S^x \rangle = -\omega_0 \langle S^y \rangle, \quad (32)$$

$$\begin{aligned} \partial_t \langle S^y \rangle &= \omega_0 \langle S^x \rangle - 4Q'_0 \langle S^y \rangle - 8Q''_0 \langle S^z \rangle \langle S^x \rangle \\ &\quad - 4Q'_1 \langle S^x \rangle - 8Q'_1 \langle S^z \rangle \langle S^y \rangle, \end{aligned} \quad (33)$$

$$\partial_t \langle S^z \rangle = -4Q'_0 \langle S^z \rangle + 8Q''_0 \langle S^x \rangle \langle S^y \rangle + 8Q'_1 \langle S^y \rangle^2. \quad (34)$$

We may note that not all these terms are extensive in the thermodynamic limit, where we assume $g^2 N$ is finite so $g^2 \propto 1/N$. Specifically, those terms involving

Q_i multiplying a single spin operator (the terms arising from commutators) scale as g^2 and so vanish in the limit $N \rightarrow \infty$. In contrast, those terms multiplying two spin operators scale as $g^2 N$ and so remain finite in this limit. Neglecting such terms is therefore consistent with considering the semiclassical (i.e. mean-field) limit, where fluctuations are suppressed by $1/N$. Since real experiments involve finite numbers of atoms and finite cavity volumes, the practical distinction is that some of the terms in this equation are N -fold smaller, and for a typical $N \simeq 10^5$, that difference is significant. Neglecting these smaller terms then gives a simplified equation of motion:

$$\begin{aligned} \partial_t \langle S^x \rangle &= -\omega_0 \langle S^y \rangle \\ \partial_t \langle S^y \rangle &= \omega_0 \langle S^x \rangle - 8Q''_0 \langle S^z \rangle \langle S^x \rangle - 8Q'_1 \langle S^z \rangle \langle S^y \rangle \\ \partial_t \langle S^z \rangle &= 8Q''_0 \langle S^x \rangle \langle S^y \rangle + 8Q'_1 \langle S^y \rangle^2. \end{aligned} \quad (35)$$

In the limit where $\omega_0 \ll \omega, \kappa$, but where it remains finite, we may approximate that the two combinations of Q_i appearing here are:

$$Q''_0 \simeq -\frac{g^2 \omega}{\omega^2 + \kappa^2}, \quad Q'_1 \simeq -\frac{2g^2 \kappa \omega \omega_0}{(\omega^2 + \kappa^2)^2}. \quad (36)$$

It is notable that this procedure (eliminating photon modes from the quantum theory, and then deriving the semiclassical equations) does not match the result derived in [31] and reviewed at the end of Sec. II B. Namely, if one first makes a semiclassical approximation for the full model, and then eliminates the photon mode, the resulting equations do not match Eq. (35); such equations are missing the term proportional to Q'_1 which describes damping. Thus, the approach described here of eliminating photons first and making a semiclassical approximation second appears to restore the missing damping. We discuss the consequences of this in the remainder of this section.

1. Steady state.

To first check the semiclassical theory, we consider the steady state and its comparison to exact solution of the full atom-only model. The steady state of Eq. (35) is satisfied by $\langle S^y \rangle = 0$ along with $\langle S^x \rangle (\omega_0 - 8Q''_0 \langle S^z \rangle) = 0$. This indeed describes two distinct states, a normal state with $\langle S^x \rangle = 0$, or a superradiant state that becomes possible once $4N|Q''_0| > \omega_0$, allowing a solution with $\langle S^z \rangle > -N/2$. Using the above result for Q''_0 we indeed see this expression matches the location of the superradiance transition in the full Dicke model. Using this definition of threshold $4g_c^2 N = \omega_0(\omega^2 + \kappa^2)/\omega$ we see that above threshold the solution is

$$\langle S^x \rangle = \pm \frac{N}{2} \sqrt{1 - \frac{g_c^4}{g^4}}, \quad \langle S^z \rangle = -\frac{N}{2} \frac{g_c^2}{g^2}. \quad (37)$$

Figure 2 shows the semiclassical solution for $\langle S^z \rangle$ in comparison to results of the exact numerics for finite size

² The proof follows from the fact that the identity always commutes with the Hamiltonian, $[H, \mathbb{1}] = 0$, and the identity can make the Lindblad form vanish, i.e. $2X\mathbb{1}X^\dagger - \{X^\dagger X, \mathbb{1}\} = 0$ if $X^\dagger = X$.




FIG. 2. Evolution of the steady state value of $\langle S^z \rangle$ with matter-light coupling $g\sqrt{N}$ for $\omega_0 = 0.1$, and $\kappa = 1$ (in units chosen so that $\omega = 1$). Curves are rescaled by $N/2$ so as to produce a finite limit as $N \rightarrow \infty$. The blue curves show the exact numerical simulations [based on the resolution of the master equation Eq. (22)] for various values of N as indicated in the legend. The dashed black curve shows the semiclassical solution (valid as $N \rightarrow \infty$). The position of $g_c\sqrt{N}$, which marks the transition between the normal state and the superradiant state, is indicated by a red vertical dashed line.

calculations. We see that these match well in the large N limit, and that in that limit, a sharp cusp develops in the exact solution. We focus on $\langle S^z \rangle$, as the finite size calculation never shows symmetry breaking, however, as discussed elsewhere [60], signatures of the transition nonetheless survive.

2. Linear stability analysis.

To understand the role of the damping term Q'_1 appearing in the equations of motion, we consider linear stability of the normal and superradiant states.

a. Normal state. For the normal state, $\langle S^x \rangle = \langle S^y \rangle = 0$, $\langle S^z \rangle = -N/2$, we can easily see that fluctuations of $\langle S^z \rangle$ decouple from those of $\langle S^{x,y} \rangle$. Denoting fluctuations of $\langle S^{x,y} \rangle$ by x, y , we find these obey the coupled equations:

$$\partial_t \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 & -\omega_0 \\ \omega_0 + 4Q''_0 N & 4Q'_1 N \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \quad (38)$$

The eigenvalues of this matrix are:

$$\lambda = 2Q'_1 N \pm i\sqrt{\omega_0(\omega_0 + 4Q''_0 N) - (2Q'_1 N)^2}. \quad (39)$$

We can use this to recover the behavior below and at threshold. When g is small, the square root is real, corresponding to oscillating modes, with damping caused by Q'_1 . Instability of the normal state occurs when an eigenvalue crosses zero. This requires $\omega_0 + 4Q''_0 N = 0$.

Inserting the expression for Q''_0 we see this again matches the expected threshold, $g > g_c$ for superradiance of the Dicke model. Using this definition of g_c we find that away from this transition, we can write the eigenvalues as

$$\lambda \simeq -\frac{4g^2 N \kappa \omega \omega_0}{(\omega^2 + \kappa^2)^2} \pm i\omega_0 \sqrt{1 - \left(\frac{g}{g_c}\right)^2}, \quad (40)$$

which matches precisely the linear stability analysis of the full model as presented in Sec. II B. For $g < g_c$, the eigenvalues have a small negative real part (given by the first term) and an imaginary part (given by the square root). Hence, the solution is stable. For $g > g_c$, one of the eigenvalues becomes positive. The solution $\langle S^z \rangle = -N/2$ is thus unstable.

b. Superradiant state. One may similarly perform a linear stability analysis around the ordered states. In this case we consider e.g. $\langle S^x \rangle = \langle S^x \rangle_{ss} + x$, where $\langle S^x \rangle_{ss}$ is the steady state value. A similar replacement holds for $\langle S^z \rangle$, while $\langle S^y \rangle = y$ as we may note that $\langle S^y \rangle_{ss} = 0$. This then gives:

$$\begin{aligned} \partial_t x &= -\omega_0 y, \\ \partial_t y &= \omega_0 x - 8Q''_0 \left(\langle S^z \rangle_{ss} x + \langle S^x \rangle_{ss} z \right) - 8Q'_1 \langle S^z \rangle_{ss} y, \\ \partial_t z &= 8Q''_0 \langle S^x \rangle_{ss} y. \end{aligned} \quad (41)$$

The second equation can be simplified by noting that the steady state condition implies that $\omega_0 = 8Q''_0 \langle S^z \rangle_{ss}$. After this, one finds the equations for y, z no longer depend on x , allowing one to directly read out the eigenvalues:

$$\lambda = -4Q'_1 \langle S^z \rangle_{ss} \pm \sqrt{(4Q'_1 \langle S^z \rangle_{ss})^2 - (8Q''_0 \langle S^x \rangle_{ss})^2}. \quad (42)$$

We may note that since $Q'_1 \ll Q''_0$, the first term in square root can be neglected. Inserting the expressions for the steady state values (37) we then find:

$$\lambda \simeq -\frac{\kappa \omega_0^2}{\omega^2 + \kappa^2} \pm i\omega_0 \sqrt{\frac{g^4}{g_c^4} - 1}. \quad (43)$$

This once again matches the linear stability analysis of the full model reviewed in Sec. II B.

V. CONCLUSION

In summary, we have shown that it is possible to produce an atom-only description of the dissipative Dicke model which correctly describes the dissipation rates and the steady states of the system. To recover such a theory it is necessary to use an unsecularized Redfield master equation, accounting for the variation of effective bath density of states at system frequencies $\pm\omega_0$. Such a theory is not of Lindblad form. Attempts to put it in Lindblad form, either by neglecting the small detuning ω_0 or by secularization lead to a master equation that is qualitatively incorrect, i.e. it fails to produce the phase transition expected. As a result, constructing a time-local

equation of motion that captures the dissipative physics of the open Dicke model in an atom-only description appears to require such a non-Lindblad form. Although non-Lindblad master equations may lead to positivity violation, this violation of positivity is restricted to specific initial conditions, and does not typically occur for the steady state of the system. Indeed, in other contexts [47] it can be shown that violation of positivity of the density matrix only occurs as a transient behavior at early time starting from certain initial states; this transient non-positivity does not cause problems for the later time evolution.

The ability to describe the dissipative dynamics with an atom-only theory is a crucial step to understand the quantum dynamics in more complicated situations. Specifically, for multimode cavities [4–14], it enables one to adiabatically eliminate the bosonic cavity modes (which massively increases the size of the Hilbert space), and provides a description of the key slow dynamics with only atomic variables. Such a description can then form a basis for numerical methods, such as matrix-product-density-operator [49–51], corner-space-renormalization [52], or cluster expansions [53], to be applied, allowing a full quantum description of the problem. Moreover, as described above, the above quantum

theory has a semiclassical limit that correctly captures the effects of dissipation. In addition, the methods described here may be particularly useful when considering fermionic atoms in optical cavities [61–63] and the possibility of cavity mediated superconductivity [64], as fermions do not admit the same semiclassical approaches as used for bosonic atoms. Future work will make use of the methods above applied to multimode problems to explore the evolution of entanglement and quantum correlations in this complex open quantum system.

All data underpinning this publication are openly available from the University of Strathclyde Knowledge-Base [65].

ACKNOWLEDGMENTS

J. K. acknowledges useful discussions with Monika Schleier-Smith and Benjamin Lev, and in particular acknowledges Monika Schleier-Smith for asking the question that prompted this work. F. D. and A. D. acknowledge support from the EPSRC Programme Grant DesOEQ (EP/P009565/1), and by the EOARD via AFOSR grant number FA9550-18-1-0064. J. K. acknowledges support from SU2P.

-
- [1] F Dimer, B Estienne, A. S Parkins, and H. J Carmichael, “Proposed realization of the Dicke-model quantum phase transition in an optical cavity QED system,” *Phys. Rev. A* **75**, 013804 (2007).
 - [2] Kristian Baumann, Christine Guerlin, Ferdinand Brennecke, and Tilman Esslinger, “Dicke quantum phase transition with a superfluid gas in an optical cavity,” *Nature* **464**, 1301–1306 (2010).
 - [3] Peter Kirton, Mor M. Roses, Jonathan Keeling, and Emanuele G. Dalla Torre, “Introduction to the Dicke Model: From Equilibrium to Nonequilibrium, and Vice Versa,” *Adv. Quantum Technol.* **2**, 1800043 (2018).
 - [4] Sarang Gopalakrishnan, Benjamin L. Lev, and Paul M. Goldbart, “Emergent Crystallinity and Frustration with Bose–Einstein Condensates in Multimode Cavities,” *Nat. Phys.* **5**, 845 (2009).
 - [5] Sarang Gopalakrishnan, B.L. Lev, and P.M. Goldbart, “Atom-light crystallization of Bose-Einstein condensates in multimode cavities: Nonequilibrium classical and quantum phase transitions, emergent lattices, supersolidity, and frustration,” *Phys. Rev. A* **82**, 043612 (2010).
 - [6] Sarang Gopalakrishnan, Benjamin L. Lev, and Paul M. Goldbart, “Frustration and Glassiness in Spin Models with Cavity-Mediated Interactions,” *Phys. Rev. Lett.* **107**, 277201 (2011).
 - [7] Sarang Gopalakrishnan, Benjamin L Lev, and Paul M Goldbart, “Exploring models of associative memory via cavity quantum electrodynamics,” *Philos. Mag.* **92**, 353–361 (2012).
 - [8] Alicia J Kollár, Alexander T Papageorge, Kristian Baumann, Michael A Armen, and Benjamin L Lev, “An adjustable-length cavity and Bose–Einstein condensate apparatus for multimode cavity QED,” *New J. Phys.* **17**, 043012 (2015).
 - [9] Alicia J Kollár, Alexander T Papageorge, Varun D Vaidya, Yudan Guo, Jonathan Keeling, and Benjamin L Lev, “Supermode-Density-Wave-Polariton Condensation,” *Nat. Commun.* **8**, 14386 (2017).
 - [10] Valentin Torggler, Sebastian Krämer, and Helmut Ritsch, “Quantum annealing with ultracold atoms in a multimode optical resonator,” *Phys. Rev. A* **95**, 032310 (2017).
 - [11] V.D. Vaidya, Y. Guo, R.M. Kroeze, K.E. Ballantine, A.J. Kollár, J. Keeling, and B.L. Lev, “Tunable-Range, Photon-Mediated Atomic Interactions in Multimode Cavity QED,” *Phys. Rev. X* **8**, 011002 (2018).
 - [12] Valentin Torggler, Philipp Aumann, Helmut Ritsch, and Wolfgang Lechner, “A Quantum N-Queens Solver,” (2018), preprint, 1803.00735.
 - [13] Yudan Guo, Ronen M Kroeze, Varun V Vaidya, Jonathan Keeling, and Benjamin L Lev, “Sign-changing photon-mediated atom interactions in multimode cavity qed,” (2018), preprint, 1810.11086.
 - [14] Yudan Guo, Varun D Vaidya, Ronen M Kroeze, Rhianon A Lunney, Benjamin L Lev, and Jonathan Keeling, “Emergent and broken symmetries of atomic self-organization arising from gouy phase shifts in multimode cavity qed,” (2018), preprint, 1810.11085.
 - [15] Kyle E. Ballantine, Benjamin L. Lev, and Jonathan Keeling, “Meissner-like effect for a synthetic gauge field in multimode cavity qed,” *Phys. Rev. Lett.* **118**, 045302 (2017).
 - [16] P Rotondo, M Marcuzzi, J P Garrahan, I Lesanovsky,

- and M Müller, “Open quantum generalisation of hopfield neural networks,” *J. Phys. A* **51**, 115301 (2018).
- [17] R. H Dicke, “Coherence in Spontaneous Radiation Processes,” *Phys. Rev.* **93**, 99–110 (1954).
- [18] M. Gross and S. Haroche, “Superradiance: An essay on the theory of collective spontaneous emission,” *Phys. Rep.* **93**, 301–396 (1982).
- [19] D Meiser, Jun Ye, D R Carlson, and M J Holland, “Prospects for a Millihertz-Linewidth Laser,” *Phys. Rev. Lett.* **102**, 163601 (2009).
- [20] Justin G. Bohnet, Zilong Chen, Joshua M. Weiner, Dominic Meiser, Murray J. Holland, and James K. Thompson, “A steady-state superradiant laser with less than one intracavity photon,” *Nature* **484**, 78–81 (2012).
- [21] K Hepp and E H Lieb, “On the Superradiant Phase Transition for Molecules in a Quantized Radiation Field: the Dicke Maser Model,” *Ann. Phys.* **76**, 360 (1973).
- [22] Y K Wang and F T Hioe, “Phase Transition in the Dicke Model of Superradiance,” *Phys. Rev. A* **7**, 831 (1973).
- [23] Klauss Hepp and Elliott Lieb, “Equilibrium statistical mechanics of matter interacting with the quantized radiation field,” *Phys. Rev. A* **8**, 2517–2525 (1973).
- [24] K. Rzażewski, K. Wódkiewicz, and W. Żakowicz, “Phase Transitions, Two-Level Atoms, and the A^2 Term,” *Phys. Rev. Lett.* **35**, 432–434 (1975).
- [25] András Vukics and Peter Domokos, “Adequacy of the Dicke model in cavity QED: A counter-no-go statement,” *Phys. Rev. A* **86**, 053807 (2012).
- [26] András Vukics, Tobias Grieser, and Peter Domokos, “Elimination of the A -Square Problem from Cavity QED,” *Phys. Rev. Lett.* **112**, 73601 (2014).
- [27] András Vukics, Tobias Grieser, and Peter Domokos, “Fundamental limitation of ultrastrong coupling between light and atoms,” *Phys. Rev. A* **92**, 43835 (2015).
- [28] Tobias Grieser, András Vukics, and Peter Domokos, “Depolarization shift of the superradiant phase transition,” *Phys. Rev. A* **94**, 33815 (2016).
- [29] Daniele De Bernardis, Tuomas Jaako, and Peter Rabl, “Cavity quantum electrodynamics in the nonperturbative regime,” *Phys. Rev. A* **97**, 43820 (2018).
- [30] Daniele De Bernardis, Philipp Pilar, Tuomas Jaako, Simone De Liberato, and Peter Rabl, “Breakdown of gauge invariance in ultrastrong-coupling cavity QED,” *Phys. Rev. A* **98**, 053819 (2018).
- [31] J. Keeling, M.J. Bhaseen, and B.D. Simons, “Collective dynamics of Bose-Einstein condensates in optical cavities,” *Phys. Rev. Lett.* **105**, 043001 (2010).
- [32] M. J. Bhaseen, J Mayoh, B. D Simons, and J Keeling, “Dynamics of nonequilibrium Dicke models,” *Phys. Rev. A* **85**, 013817 (2012).
- [33] Minghui Xu, D A Tieri, and M J Holland, “Simulating open quantum systems by applying $SU(4)$ to quantum master equations,” *Phys. Rev. A* **87**, 62101 (2013).
- [34] Peter Kirton and Jonathan Keeling, “Superradiant and lasing states in driven-dissipative Dicke models,” *New J. Phys.* **20**, 015009 (2018).
- [35] Nathan Shammah, Shah Nawaz Ahmed, Neill Lambert, Simone De Liberato, and Franco Nori, “Open quantum systems with local and collective incoherent processes: Efficient numerical simulations using permutational invariance,” *Phys. Rev. A* **98**, 063815 (2018).
- [36] H.-P. Breuer and F Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2002).
- [37] A. G. Redfield, “On the Theory of Relaxation Processes,” *IBM J. Res. Dev.* **1**, 19–31 (1957).
- [38] F. Bloch, “Generalized theory of relaxation,” *Phys. Rev.* **105**, 1206–1222 (1957).
- [39] G. Lindblad, “On the generators of quantum dynamical semigroups,” *Commun. Math. Phys.* **48**, 119–130 (1976).
- [40] E. B. Davies, “Markovian master equations,” *Comm. Math. Phys.* **39**, 91 (1974).
- [41] R. Duemcke, H Spohn, and R Dümcke, “The proper form of the generator in the weak coupling limit,” *Z. Physik B* **34**, 419–422 (1979).
- [42] Robert S Whitney, “Staying positive: going beyond Lindblad with perturbative master equations,” *J. Phys. A Math. Theor.* **41**, 175304 (2008).
- [43] R. Wangsness and F. Bloch, “The Dynamical Theory of Nuclear Induction,” *Phys. Rev.* **89**, 728–739 (1953).
- [44] V Gorini, A Kossakowski, and E. C. G. Sudarshan, “Completely positive dynamical semigroups of N -level systems,” *J. Math. Phys.* **17**, 821 (1976).
- [45] Jan Jeske, David Ing, Martin B. Plenio, Susana F. Huelga, and Jared H. Cole, “Bloch-Redfield equations for modeling light-harvesting complexes,” *J. Chem. Phys.* **142**, 064104 (2015).
- [46] H. M. Cammack, P. Kirton, T. M. Stace, P. R. Eastham, J. Keeling, and B. W. Lovett, “Coherence protection in coupled quantum systems,” *Phys. Rev. A* **97**, 022103 (2018).
- [47] P. R. Eastham, P. Kirton, H. M. Cammack, B. W. Lovett, and J. Keeling, “Bath-induced coherence and the secular approximation,” *Phys. Rev. A* **94**, 012110 (2016).
- [48] A. Dodin, T. Tscherbul, R. Alicki, A. Vutha, and P. Brumer, “Secular versus nonsecular Redfield dynamics and Fano coherences in incoherent excitation: An experimental proposal,” *Phys. Rev. A* **97**, 013421 (2018).
- [49] A J Daley, C Kollath, U Schollwöck, and G Vidal, “Time-dependent density-matrix renormalization-group using adaptive effective hilbert spaces,” *J. Stat. Mech. Theory Exp.* **2004**, P04005 (2004).
- [50] F. Verstraete, J. J. García-Ripoll, and J. I. Cirac, “Matrix product density operators: Simulation of finite-temperature and dissipative systems,” *Phys. Rev. Lett.* **93**, 207204 (2004).
- [51] Michael Zwolak and Guifré Vidal, “Mixed-state dynamics in one-dimensional quantum lattice systems: A time-dependent superoperator renormalization algorithm,” *Phys. Rev. Lett.* **93**, 207205 (2004).
- [52] S. Finazzi, A. Le Boité, F. Storme, A. Baksic, and C. Ciuti, “Corner-space renormalization method for driven-dissipative two-dimensional correlated systems,” *Phys. Rev. Lett.* **115**, 080604 (2015).
- [53] Jiasen Jin, Alberto Biella, Oscar Viyuela, Leonardo Mazza, Jonathan Keeling, Rosario Fazio, and Davide Rossini, “Cluster mean-field approach to the steady-state phase diagram of dissipative spin systems,” *Phys. Rev. X* **6**, 031011 (2016).
- [54] H J Lipkin, N Meshkov, and A J Glick, “Validity of many-body approximation methods for a solvable model: (i). exact solutions and perturbation theory,” *Nucl. Phys.* **62**, 188 (1965).
- [55] N Meshkov, A J Glick, and H J Lipkin, “Validity of many-body approximation methods for a solvable model: (ii). linearization procedures,” *Nucl. Phys.* **62**, 199 (1965).

- [56] A J Glick, H J Lipkin, and N Meshkov, “Validity of many-body approximation methods for a solvable model: (iii). diagram summations,” *Nucl. Phys.* **62**, 211 (1965).
- [57] M O Scully and M S Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, 1997).
- [58] Patrik Fazekas, *Lecture notes on electron correlation and magnetism* (World Scientific, Singapore, 1999).
- [59] Ryosuke Imai and Yoshiya Yamanaka, “Dynamical properties of the finite-size dicke model coupled to a thermal reservoir,” *Journal of the Physical Society of Japan* **88**, 024401 (2019), <https://doi.org/10.7566/JPSJ.88.024401>.
- [60] Peter Kirton and Jonathan Keeling, “Suppressing and restoring the dicke superradiance transition by dephasing and decay,” *Phys. Rev. Lett.* **118**, 123602 (2017).
- [61] J. Keeling, M. J. Bhaseen, and B. D. Simons, “Fermionic superradiance in a transversely pumped optical cavity,” *Phys. Rev. Lett.* **112**, 143002 (2014).
- [62] Francesco Piazza and Philipp Strack, “Umklapp superradiance with a collisionless quantum degenerate fermi gas,” *Phys. Rev. Lett.* **112**, 143003 (2014).
- [63] Yu Chen, Zhenhua Yu, and Hui Zhai, “Superradiance of degenerate fermi gases in a cavity,” *Phys. Rev. Lett.* **112**, 143004 (2014).
- [64] E. Colella, R. Citro, M. Barsanti, D. Rossini, and M.-L. Chiofalo, “Quantum phases of spinful Fermi gases in optical cavities,” *Phys. Rev. B* **97**, 134502 (2018).
- [65] <http://dx.doi.org/10.15129/56063567-ad36-466d-9d45-2a64dac4b1f9>.