

On orthogonal transformations of Christoffel equations

Len Bos*, Michael A. Slawinski†, Theodore Stanoev‡, Maurizio Vianello§

Abstract: The purpose of this paper is to prove the equivalence—under rotations of distinct terms—of different forms of a determinantal equation that appears in the studies of wave propagation in Hookean solids, in the context of the Christoffel equations. To do so, we prove a general proposition that is not limited to \mathbb{R}^3 , nor is it limited to the elasticity tensor with its index symmetries. Furthermore, the proposition is valid for orthogonal transformations, not only for rotations. The sought equivalence is a corollary of that proposition.

1 Introduction

The existence and properties of three waves that propagate in a Hookean solid are a consequence of the Christoffel equations (e.g., Slawinski, 2015, Chapter 9), whose solubility condition is

$$\det \left[\sum_{j=1}^3 \sum_{\ell=1}^3 c_{ijkl} p_j p_\ell - \delta_{ik} \right] = 0, \quad i, k = 1, 2, 3,$$

which is a cubic polynomial, whose roots are the eikonal equations (e.g., Slawinski, 2015, Section 7.3). Let us examine the matrix therein,

$$\left[\sum_{j=1}^3 \sum_{\ell=1}^3 c_{ijkl} p_j p_\ell \right] \in \mathbb{R}^{3 \times 3},$$

where c_{ijkl} is a density-normalized elasticity tensor, whose units are $\text{km}^2 \text{s}^{-2}$, and p is the wavefront-slowness vector, whose units are s km^{-1} .

Studies of Hookean solids by Ivanov and Stovas (2016, equations (7)–(12)) and Ivanov and Stovas (2017, equations (10)–(11)) invoke a property that we state as Corollary 1, which is a consequence of Proposition 1. Ivanov and Stovas (2016, 2017) verify the equivalence of equations given in Corollary 1, without a general proof, hence, this paper.

The purpose of this paper is to prove Proposition 1 and, hence, Corollary 1. In doing so, we gain an insight into a tensor-algebra property that results in this corollary. The equivalence of the aforementioned equations is not a result of the invariance of a determinant, as suggested by Ivanov (pers. comm., 2018); it is a consequence of two orthogonal transformations of c_{ijkl} and p_i that result in two matrices that are similar to one another.

*Dipartimento di Informatica, Università di Verona, Italy; leonardpeter.bos@univr.it

†Department of Earth Sciences, Memorial University of Newfoundland, Canada; msslawins@mac.com

‡Department of Earth Sciences, Memorial University of Newfoundland, Canada; theodore.stanoev@gmail.com

§Dipartimento di Matematica, Politecnico di Milano, Italy; maurizio.vianello@polimi.it

2 Proposition and its corollary

Proposition 1. Consider a tensor, c_{ijkl} , in \mathbb{R}^d . Also, consider a vector, p_i , in \mathbb{R}^d , and an orthogonal transformation, $A \in \mathbb{R}^{d \times d}$. It follows that matrices

$$\left[\sum_{j=1}^d \sum_{\ell=1}^d c_{ijkl} \hat{p}_j \hat{p}_\ell \right]_{1 \leq i, k \leq d} \in \mathbb{R}^{d \times d} \quad (1)$$

and

$$\left[\sum_{j=1}^d \sum_{\ell=1}^d \check{c}_{ijkl} p_j p_\ell \right]_{1 \leq i, k \leq d} \in \mathbb{R}^{d \times d}, \quad (2)$$

where

$$\hat{p}_i := \sum_{j=1}^d A_{i,j} p_j \quad \text{and} \quad \check{c}_{ijkl} := \sum_{m=1}^d \sum_{n=1}^d \sum_{o=1}^d \sum_{q=1}^d A_{m,i} A_{n,j} A_{o,k} A_{q,\ell} c_{mnoq},$$

are similar to one another and, consequently, have the same spectra.

Proof. The fourth-rank tensor, c_{ijkl} , in \mathbb{R}^d can be viewed as a $d \times d$ matrix, whose entries are $d \times d$ matrices,

$$C = [C_{ik}]_{1 \leq i, k \leq d} \in (\mathbb{R}^{d \times d})^{d \times d},$$

with $C_{ik} \in \mathbb{R}^{d \times d}$ and $(C_{ik})_{j\ell} := c_{ijkl}$. Thus, matrix (1) can be written as

$$[\hat{p}^t C_{ik} \hat{p}]_{1 \leq i, k \leq d} = [(Ap)^t C_{ik} (Ap)]_{1 \leq i, k \leq d} = [p^t (A^t C_{ik} A) p]_{1 \leq i, k \leq d} \in \mathbb{R}^{d \times d}, \quad (3)$$

where t denotes the transpose.

We claim that matrix (2) can be written as

$$A^t [p^t (A^t C_{ik} A) p]_{1 \leq i, k \leq d} A \in \mathbb{R}^{d \times d}. \quad (4)$$

To see this, we let matrix (4) be

$$M = A^t X A,$$

where $X := [p^t (A^t C_{ik} A) p]_{1 \leq i, k \leq d}$, to write

$$M_{i,k} = \sum_{m=1}^d \sum_{o=1}^d (A^t)_{i,m} X_{m,o} A_{o,k} = \sum_{m=1}^d \sum_{o=1}^d A_{m,i} X_{m,o} A_{o,k}.$$

Defining $Y := A^t C_{mo} A$, we have

$$X_{m,o} = \sum_{j=1}^d \sum_{\ell=1}^d Y_{j,\ell} p_j p_\ell,$$

where

$$Y_{j,\ell} = \sum_{n=1}^d \sum_{q=1}^d (A^t)_{j,n} (C_{mo})_{nq} A_{q,\ell} = \sum_{n=1}^d \sum_{q=1}^d A_{n,j} A_{q,\ell} c_{mnoq}.$$

Hence,

$$X_{m,o} = \sum_{j=1}^d \sum_{\ell=1}^d \left(\sum_{n=1}^d \sum_{q=1}^d A_{n,j} A_{q,\ell} c_{mnoq} \right)$$

and, in turn,

$$M_{i,k} = \sum_{j=1}^d \sum_{\ell=1}^d \left(\sum_{m=1}^d \sum_{n=1}^d \sum_{o=1}^d \sum_{q=1}^d A_{m,i} A_{n,j} A_{o,k} A_{q,\ell} c_{mnoq} \right) p_j p_\ell = \sum_{j=1}^d \sum_{\ell=1}^d \check{c}_{ijkl} p_j p_\ell, \quad i, k = 1, \dots, d,$$

which is matrix (2), as required. \square

Corollary 1. *From Proposition 1—and the aforementioned fact that the similar matrices share the same spectrum, as well as the fact that the similarity of matrices is not affected by subtracting from them the identity matrices—it follows that*

$$\det \left[\sum_{j=1}^3 \sum_{\ell=1}^3 c_{ijkl} \hat{p}_j \hat{p}_\ell - \delta_{ik} \right] = \det \left[\sum_{j=1}^3 \sum_{\ell=1}^3 \check{c}_{ijkl} p_j p_\ell - \delta_{ik} \right], \quad i, k = 1, 2, 3,$$

and, hence, equations

$$\det \left[\sum_{j=1}^3 \sum_{\ell=1}^3 c_{ijkl} \hat{p}_j \hat{p}_\ell - \delta_{ik} \right] = 0, \quad i, k = 1, 2, 3, \quad (5)$$

and

$$\det \left[\sum_{j=1}^3 \sum_{\ell=1}^3 \check{c}_{ijkl} p_j p_\ell - \delta_{ik} \right] = 0, \quad i, k = 1, 2, 3, \quad (6)$$

are equivalent to one another.

Corollary 1 is valid even without requiring the index symmetries of Hookean solids. Also, $A \in O(3)$, not only $A \in SO(3)$, which is more general than the property invoked by Ivanov and Stovas (2016, 2017).

3 Numerical example

Consider an orthotropic tensor (Ivanov and Stovas, 2016, Table 2), whose components are

$$\begin{aligned} c_{1111} &= 6.3, & c_{2222} &= 6.9, & c_{3333} &= 5.4, \\ c_{1122} &= c_{2211} = 2.7, & c_{1133} &= c_{3311} = 2.2, & c_{2233} &= c_{3322} = 2.4, \\ c_{1212} &= c_{2112} = c_{2121} = c_{1221} = 1.5, & c_{1313} &= c_{3113} = c_{3131} = c_{1331} = 0.8, & c_{2323} &= c_{3223} = c_{3232} = c_{2332} = 1.0. \end{aligned}$$

Also, consider vector $p = \left[0, 0, \sqrt{\frac{1}{c_{3333}}} \right]$. Rotating this vector by

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix}, \quad (7)$$

with an arbitrary angle of $\theta = \frac{\pi}{5}$, and the tensor by A^t , we obtain

$$\hat{\Gamma} := \sum_{j=1}^3 \sum_{\ell=1}^3 c_{ijkl} \hat{p}_j \hat{p}_\ell = \begin{bmatrix} 0.192934 & 0 & 0 \\ 0 & 0.331749 & -0.0184241 \\ 0 & -0.0184241 & 0.949406 \end{bmatrix}$$

and

$$\check{\Gamma} := \sum_{j=1}^3 \sum_{\ell=1}^3 \check{c}_{ijkl} p_j p_\ell = \begin{bmatrix} 0.192934 & 0 & 0 \\ 0 & 0.562667 & -0.299407 \\ 0 & -0.299407 & 0.718488 \end{bmatrix},$$

respectively. The eigenvalues of these matrices are the same, $\lambda_1 = 0.949955$, $\lambda_2 = 0.33120$ and $\lambda_3 = 0.192934$, as required for similar matrices. Their corresponding eigenvectors are related by transformation (7).

Herein, $\det[\hat{\Gamma} - I] = -0.027013 = \det[\check{\Gamma} - I]$. In general, the two determinants are equal to one another. Hence, if $\det[\hat{\Gamma} - I] = 0$, so does $\det[\check{\Gamma} - I]$, and *vice versa*.

The equivalence of equations (5) and (6) does not imply their equivalence to

$$\det \left[\sum_{j=1}^3 \sum_{\ell=1}^3 c_{ijkl} p_j p_\ell - \delta_{ik} \right] = 0, \quad i, k = 1, 2, 3.$$

The eigenvalues of

$$\Gamma := \sum_{j=1}^3 \sum_{\ell=1}^3 c_{ijkl} p_j p_\ell = \begin{bmatrix} 0.148148 & 0 & 0 \\ 0 & 0.185185 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

are $\lambda_1 = 0.148148$, $\lambda_2 = 0.185185$ and $\lambda_3 = 1$, which are distinct from the eigenvalues of $\hat{\Gamma}$ and $\check{\Gamma}$. Herein—in view of p and c_{ijkl} representing, respectively, the slowness vector along the x_3 -axis and its corresponding elasticity tensor— $\det[\Gamma - I] = 0$, which results in the eikonal equations. We emphasize, however, that Proposition 1 and Corollary 1 are valid for arbitrary vectors and fourth-rank tensors, even though, in this example, they are related by the Christoffel equations.

Acknowledgments

The authors wish to acknowledge Igor Ravve for bringing this issue to their attention, Yuriy Ivanov for presenting and clarifying aspects of this problem, Sandra Forte for fruitful discussions, and David Dalton for insightful proofreading.

This research was performed in the context of The Geomechanics Project supported by Husky Energy. Also, this research was partially supported by the Natural Sciences and Engineering Research Council of Canada, grant 202259.

References

- Ivanov, Y. and Stovas, A. (2016). Normal moveout velocity ellipse in tilted orthorhombic media. *Geophysics*, 81(6):319–336.
- Ivanov, Y. and Stovas, A. (2017). Traveltime parameters in tilted orthorhombic medium. *Geophysics*, 82(6):187–200.
- Slawinski, M. A. (2015). *Waves and rays in elastic continua*. World Scientific, 3 edition.