

Viewpoint – Utilizing Data from Prevalent Technologies for Atmospheric Research

Noam David¹

Institute of Industrial Science, The University of Tokyo, Tokyo, Japan

Over the past decade, the Internet of Things (IoT) and smart devices have become increasingly common as part of the technological infrastructure that surrounds us. The flow of data generated by these systems is characterized by enormous granularity, availability and coverage. As a result, new opportunities are being opened to utilize the newly available information for various needs and, in particular, for atmospheric research. If we consider the data generated by these means, we may notice that many produce measurements with high environmental value. To name some examples - surveillance cameras that operate in the visible light spectrum are positioned in a vast number of locations. Previous works have shown that they can be used for monitoring the temporal patterns of fine atmospheric particulate matter (Wong et al., 2007). Lab experiments indicate a direct link between the speed of movement of car wipers, and the rainfall intensity. Hence, advanced vehicles that store these data can, in essence, be used as moving rain gauges (Rabiei et al., 2013). Kawamura et al (2017) have revealed a novel technique for monitoring atmospheric humidity using terrestrial broadcasting waves, based on propagation delays due to water vapor. Data shared as open source from social networks have been found to be potentially effective in improving automatic weather observations. Indeed, for the most part, the initial weather observation is conducted

¹Correspondence to: Dr. Noam David

E mail: nd363@cornell.edu

automatically by dedicated sensors, however, some weather conditions are still better detected by the human eye. On the other hand, millions of 'human observers' around the world use applications such as Twitter, which allows them to report publicly on subjects that are relevant to them, and in particular on weather phenomena (Cox and Plale, 2011). As was recently reviewed by Price et al (2018), in 2020 there will be more than 20 billion smartphones carried by the public worldwide. These mobile devices are equipped with sensors that can be used for environmental monitoring on a multisource basis. Recent works indicate the ability to obtain atmospheric temperature information for the urban canopy layer (Overeem et al., 2013a), to measure atmospheric pressure (Mass and Madaus, 2014), or to study atmospheric tides (Price et al., 2018). Additional studies point to the potential of using any camera enabled smart mobile device to monitor air quality (Pan et al., 2017).

Given the comprehensive coverage of the new 'virtual sensors' from all land locations across the whole globe, this low cost solution introduces a wide range of possibilities which previously could not be offered through existing technologies.

A key example of utilizing exiting data sources for atmospheric monitoring is the use of measurements acquired by commercial microwave links (CMLs) that comprise the infrastructure for wireless data transport between cellular base stations. This technology has essentially become a valuable weather monitoring tool over the past decade (e.g. Overeem et al., 2013b; Alpert et al., 2016; Gosset et al., 2016; David and Gao, 2017).

Weather phenomena and atmospheric conditions interfere with the electromagnetic waves transmitted by CMLs. Thus, these networks provide, in essence, an already existing environmental monitoring facility. The various works done in the field indicate

the ability to detect and map rainfall (e.g. Messer et al., 2006; Berne and Uijlenhoet, 2007; Chwala et al., 2012; Overeem et al., 2013b; Fencel et al., 2015), monitor fog, atmospheric moisture and dew (e.g. Chwala et al., 2014; David, 2014; David et al., 2016). Recent research revealed the ability of these radio links to identify atmospheric refractive index variations and to indirectly detect air pollution conditions (David and Gao, 2016; 2018).

Indeed, the new data available from the various means (smartphones, social networks, etc) and particularly from CMLs can provide observations with considerable spatial coverage and with minimal cost. However, the accuracy of each 'sensor' is lower than that of a dedicated instrument. This being the case, is it possible to produce significant information compared to that derived from specialized tools?

It can be assumed that the 'virtual sensors' are not a substitute for conventional monitoring means, whenever those exist in the field. The correct approach, then, is to consider the newly available sources of data as complementary measures to dedicated measurements and as a substitute during the many cases in which conventional monitoring tools are not available. However, the data acquired by prevalent technologies, even when taken by itself, often holds enormous potential.

In order to demonstrate the added value which lies in IoT data and prevalent technologies, let us focus on CMLs as an example of such a system.

Atmospheric moisture is more poorly characterized than wind or even precipitation due to the difficulty in observing the humidity field. Therefore, questions such as the magnitude of small-scale variability of moisture in the boundary layer, and its effect on convection initiation are still unanswered (Weckwerth et al., 2004). As a result,

prediction of convective precipitation, on the storm scale, is limited. However, for significantly improving convection initiation measurements, one will need moisture measurements at meso- γ resolution with accuracies of up to 1 gr/m^3 (Fabry, 2006). Notably, such type of observations can be acquired using CMLs (David et al., 2019). High resolution precipitation distribution maps can be generated using CMLs, therefore the relationship between pollutant wash-off and rainfall provides an opportunity to potentially acquire important spatial information about air quality, as discussed in recent research (David and Gao, 2016). Moreover, Liquid Water Content (LWC) constitutes a major parameter in fog research. Fog LWC changes in space, altitude, and over time, and is dependent on surface and atmospheric conditions (Gultepe et al., 2007). However, conventional sensors for acquiring LWC estimates are limited in the spatial range they can cover and in their availability. It has been shown that CMLs are able to provide fog LWC estimates across large spatial regions where dedicated sensors are nonexistent. Indeed, the availability of various spectral channels from satellites provides the possibility to observe clouds, aerosols, Earth surface and fog - in particular (Lensky and Rosenfeld, 2008; Michael et al., 2018). However, CMLs have also been found to have potential advantages for detecting fog under challenging conditions where satellite retrievals are limited, e.g when high altitude clouds cover the fog as observed from the satellite vantage point (David, 2018). Alternatively, the ability to monitor rainfall in areas where radars suffer from clutter effects (Goldshtein et al., 2009) or are blocked by complex topography (David, 2014), has also been demonstrated.

The possibilities for monitoring atmospheric phenomena utilizing the new observational powers are many, the available information vast, and the cost minimal, since the

'opportunistic sensors' are already deployed in the field. As a result, this mean of monitoring the environment is becoming advantageous for atmospheric research. Notably, the newly available 'virtual sensors' open the possibility of assimilation of their measurements into high-resolution numerical prediction models which could lead to improvements in the forecasting capabilities that exist today. In the applicative aspect, this novel approach could lay the groundwork for developing new early warning systems against natural hazards, generating a range of products required for a wide range of fields and, thus, the overall potential contribution to public health and safety may be invaluable.

Acknowledgments

I wish to gratefully acknowledge Associate Professor Dr. Yoshihide Sekimoto and his research team for the fruitful discussions and for hosting me in their laboratory as a Visiting Scientist at the Institute of Industrial Science of the University of Tokyo, Japan during 2018-2019.

References

Alpert P, Messer H, David N. Meteorology: Mobile networks aid weather monitoring. *Nature*. 2016 Sep 28;537(7622):617.

Berne A, Uijlenhoet R. Path-averaged rainfall estimation using microwave links: Uncertainty due to spatial rainfall variability. *Geophysical research letters*. 2007 Apr 1;34(7).

Chwala C, Gmeiner A, Qiu W, Hipp S, Nienaber D, Siart U, Eibert T, Pohl M, Seltmann J, Fritz J, Kunstmann H. Precipitation observation using microwave backhaul links in the alpine and pre-alpine region of Southern Germany. *Hydrology and Earth System Sciences*. 2012 Aug 13;16(8):2647-61.

Chwala C, Kunstmann H, Hipp S, Siart U. A monostatic microwave transmission experiment for line integrated precipitation and humidity remote sensing. *Atmospheric research*. 2014 Jul 1;144:57-72.

Cox J, Plale B. Improving automatic weather observations with the public Twitter stream. IU School of Informatics and Computing. 2011 Feb 22.

David N. Water vapor, fog and rainfall monitoring using commercial microwave network measurements: Noam, David. Universitat Tel-Aviv; 2014.

David N, Harel O, Alpert P, Messer H. Study of attenuation due to wet antenna in microwave radio communication. In Acoustics, Speech and Signal Processing (ICASSP), 2016 IEEE International Conference on 2016 Mar 20 (pp. 4418-4422). IEEE.

David N, Gao HO. Using cellular communication networks to detect air pollution. Environmental science & technology. 2016 Aug 18;50(17):9442-51.

David N, Gao HO. Atmospheric Monitoring Using Commercial Microwave Networks. 15th International Conference on Environmental Science And Technology (CEST). Rhodes, Greece, August 31–September 2, 2017. https://cest.gnest.org/sites/default/files/presentation_file_list/cest2017_01386_oral_paper.pdf

David N, Gao HO. Using Cell-Phone Tower Signals for Detecting the Precursors of Fog. Journal of Geophysical Research: Atmospheres. 2018 Jan 27;123(2):1325-38.

David N. Utilizing microwave communication data for detecting fog where satellite retrievals are challenged. Natural Hazards. 2018 Nov 1;94(2):867-82.

David N, Sendik O, Rubin Y, Messer H, Gao HO, Rostkier-Edelstein D, Alpert P. Analyzing the ability to reconstruct the moisture field using commercial microwave network data. Atmospheric Research. 2019 Jan 4. doi: 10.1016/j.atmosres.2018.12.025

Fabry F. The spatial variability of moisture in the boundary layer and its effect on convection initiation: Project-long characterization. Monthly Weather Review. 2006 Jan;134(1):79-91.

Fencl M, Rieckermann J, Sýkora P, Stránský D, Bareš V. Commercial microwave links instead of rain gauges: fiction or reality?. Water Science and Technology. 2015 Jan 1;71(1):31-7.

Goldshtein O, Messer H, Zinevich A. Rain rate estimation using measurements from commercial telecommunications links. IEEE Transactions on signal processing. 2009 Apr;57(4):1616-25.

Gosset M, Kunstmann H, Zougmore F, Cazenave F, Leijnse H, Uijlenhoet R, Chwala C, Keis F, Doumounia A, Boubacar B, Kacou M. Improving rainfall measurement in gauge poor regions thanks to mobile telecommunication networks. Bulletin of the American Meteorological Society. 2016 Mar;97(3):ES49-51.

Gultepe I, Tardif R, Michaelides SC, Cermak J, Bott A, Bendix J, Müller MD, Pagowski M, Hansen B, Ellrod G, Jacobs W. Fog research: A review of past achievements and future perspectives. *Pure and Applied Geophysics*. 2007 Jun 1;164(6-7):1121-59.

Kawamura S, Ohta H, Hanado H, Yamamoto MK, Shiga N, Kido K, Yasuda S, Goto T, Ichikawa R, Amagai J, Imamura K. Water vapor estimation using digital terrestrial broadcasting waves. *Radio Science*. 2017 Mar 1;52(3):367-77.

Lensky IM, Rosenfeld D. Clouds-aerosols-precipitation satellite analysis tool (CAPSAT). *Atmospheric Chemistry and Physics*. 2008 Nov 24;8(22):6739-53.

Mass CF, Madaus LE. Surface pressure observations from smartphones: A potential revolution for high-resolution weather prediction?. *Bulletin of the American Meteorological Society*. 2014 Sep;95(9):1343-9.

Messer H, Zinevich A, Alpert P. Environmental monitoring by wireless communication networks. *Science*. 2006 May 5;312(5774):713-.

Michael Y, Lensky I, Brenner S, Tchetchik A, Tessler N, Helman D. Economic Assessment of Fire Damage to Urban Forest in the Wildland–Urban Interface Using Planet Satellites Constellation Images. *Remote Sensing*. 2018 Sep;10(9):1479.

Overeem A, Robinson JC, Leijnse H, Steeneveld GJ, Horn BK, Uijlenhoet R. Crowdsourcing urban air temperatures from smartphone battery temperatures. *Geophysical Research Letters*. 2013(a) Aug 16;40(15):4081-5.

Overeem A, Leijnse H, Uijlenhoet R. Country-wide rainfall maps from cellular communication networks. *Proceedings of the National Academy of Sciences*. 2013(b) Feb 19;110(8):2741-5.

Pan Z, Yu H, Miao C, Leung C. Crowdsensing Air Quality with Camera-Enabled Mobile Devices. In *AAAI 2017* Feb 8 (pp. 4728-4733).

Price C, Maor R, Shachaf H. Using smartphones for monitoring atmospheric tides. *Journal of Atmospheric and Solar-Terrestrial Physics*. 2018 Sep 1;174:1-4.

Rabiei E, Haberlandt U, Sester M, Fitzner D. Rainfall estimation using moving cars as rain gauges-laboratory experiments. *Hydrology and Earth System Sciences* 17 (2013), Nr. 11. 2013;17(11):4701-12.

Weckwerth TM, Parsons DB, Koch SE, Moore JA, LeMone MA, Demoz BB, Flamant C, Geerts B, Wang J, Feltz WF. An overview of the International H₂O Project (IHOP_2002) and some preliminary highlights. *Bulletin of the American Meteorological Society*. 2004 Feb;85(2):253-78.

Wong CJ, MatJafri MZ, Abdullah K, Lim HS, Low KL. Temporal air quality monitoring using surveillance camera. In Geoscience and Remote Sensing Symposium, 2007. IGARSS 2007. IEEE International 2007 Jul 23 (pp. 2864-2868). IEEE.