

Maximal randomness from partially entangled states

Erik Woodhead^{*1,2}, Jędrzej Kaniewski^{†3,4}, Boris Bourdoncle², Alexia Salavrakos², Joseph Bowles², Antonio Acín^{2,5}, and Remigiusz Augusiak⁴

¹*Laboratoire d'Information Quantique, CP 225, Université libre de Bruxelles,
Avenue F. D. Roosevelt 50, 1050 Bruxelles, Belgium*

²*ICFO – Institut de Ciències Fotoniques, The Barcelona Institute of Science and Technology,
08860 Castelldefels (Barcelona), Spain*

³*Faculty of Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland*

⁴*Center for Theoretical Physics, Polish Academy of Sciences, Aleja Lotników 32/46, 02-668 Warsaw, Poland*

⁵*ICREA – Institutio Catalana de Recerca i Estudis Avançats, Lluís Companys 23, 08010 Barcelona, Spain*

(Dated: 13 November 2020)

We investigate how much randomness can be extracted from a generic partially entangled pure state of two qubits in a device-independent setting, where a Bell test is used to certify the correct functioning of the apparatus. For any such state, we first show that two bits of randomness are always attainable both if projective measurements are used to generate the randomness globally or if a nonprojective measurement is used to generate the randomness locally. We then prove that the maximum amount of randomness that can be generated using nonprojective measurements globally is restricted to between approximately 3.58 and 3.96 bits. The upper limit rules out that a bound of four bits potentially obtainable with extremal qubit measurements can be attained. We point out this is a consequence of the fact that nonprojective qubit measurements with four outcomes can only be self-tested to a limited degree in a Bell experiment.

Although it was not the original motivation [1], Bell's theorem [2] allows for a very strong test of quantum randomness. By preparing an entangled quantum system and exhibiting a Bell inequality violation with it, we can immediately know that the measurement outcomes were not the result of an underlying deterministic process. This observation is the basis of a class of quantum cryptography protocols, called *device independent*, that incorporate a Bell test as a self-test of the correct functioning of the apparatus. The class includes device-independent versions of quantum key distribution and random number generation [3–6].

This perspective prompts an obvious question: How much randomness can we extract from a given quantum system, and how might this depend on the degree of entanglement? Previous work (see table 1) has indicated that the two do not seem strongly related; we cannot necessarily get more randomness from a maximally entangled state than a weakly entangled one of the same dimension. This point was first made in [7] where it was shown that, with the help of a suitable Bell test, a uniformly random bit could be generated from the result of a projective measurement performed on one part of any partially entangled pure state of two qubits. Ref. [7] also considered the possibility of generating more randomness from the joint outcomes of projective measurements performed on both subsystems. In this case, [7] found that the maximum of two uniformly random bits could be generated, but only confirmed that this was attainable using a maximally entangled state $|\phi_+\rangle = (|00\rangle + |11\rangle)/\sqrt{2}$ or one could get arbitrarily close to it using a very weakly entangled state of the form $|\psi_\theta\rangle = \cos(\theta/2)|00\rangle + \sin(\theta/2)|11\rangle$ in the limit $\theta \rightarrow 0$ where it becomes separable. Between these two

		$ \phi_+\rangle$	$ \psi_\theta\rangle$
Local	PROJ	1 bit [5]	1 bit [7]
	POVM	2 bits [8]	2 bits
Global	PROJ	2 bits [7]	2 bits
	POVM	[3.58, 3.96] bits	[3.58, 3.96] bits

Table 1: Maximum amount of randomness (quantified by the min-entropy) extractable from one (local) or jointly from two (global) projective (PROJ) or nonprojective (POVM) measurements from the maximally ($|\phi_+\rangle$) and any partially ($|\psi_\theta\rangle$) entangled two-qubit pure state. Square brackets indicate a range (rounded outward) within which the optimal amount of randomness is known to lie. The results proved in this work are highlighted.

extremes, determining the amount of randomness that can be generated remains an open problem.

As well as projective measurements, it is also possible to perform nonprojective measurements on quantum systems. Nonprojective measurements can potentially generate more randomness as they can have more outcomes than the dimension of the quantum system they act on. Extremal qubit measurements in particular may have up to four outcomes [9]. In a bipartite Bell-type experiment this means that potentially up to two bits of randomness could be generated locally or up to four bits globally using nonprojective measurements. The first limit is known to be attainable: Ref. [8] describes a way in which two bits of randomness can be generated locally using a single

^{*}erik.woodhead@ulb.ac.be

[†]jkaniewski@fuw.edu.pl

measurement on one side. But it is currently an open question whether the second limit of four bits is attainable globally. The same work, [8], only confirmed numerically that at least around 2.8997 bits of randomness can be generated this way. Both results were established only for the maximally entangled state.

In this work, we solve the question of how much randomness can be generated using projective measurements from a generic pure entangled state of two qubits and show that the upper limit of two bits is always attainable regardless of how strong or weak the entanglement is. We also show that, alternatively, two bits of randomness can be extracted locally from any such state using a nonprojective measurement. It turns out however, as we will detail below, that nonprojective measurements can only be reconstructed to a limited degree from the correlations observed in a Bell experiment and this limits the amount of randomness that can be generated globally. We rule out that any scheme can generate more than about 3.9527 bits of randomness in this way, proving that the potential upper limit of four bits is not attainable. We nevertheless show that at least around 3.5850 bits of randomness can be generated globally with suitable nonprojective measurements from any partially entangled state.

The Bell test.— To introduce the problem, we begin by considering the form of an arbitrary partially entangled state of two qubits. Such a state can always be expressed in its Schmidt decomposition as

$$|\psi_\theta\rangle = \cos\left(\frac{\theta}{2}\right)|00\rangle + \sin\left(\frac{\theta}{2}\right)|11\rangle \quad (1)$$

for an angle θ that, without loss of generality, we can and hereafter will take to be in the range $0 < \theta \leq \frac{\pi}{2}$. The same state is equivalently represented by its density operator $\psi_\theta = |\psi_\theta\rangle\langle\psi_\theta|$, which we can express as

$$\begin{aligned} \psi_\theta = \frac{1}{4} \Big[& \mathbb{1} \otimes \mathbb{1} + \cos(\theta)(\mathbb{1} \otimes Z + Z \otimes \mathbb{1}) \\ & + \sin(\theta)(X \otimes X - Y \otimes Y) + Z \otimes Z \Big] \end{aligned} \quad (2)$$

in terms of the identity and Pauli operators $\mathbb{1}$, X , Y , and Z acting on each subsystem. We can see that Alice and Bob will have to perform measurements in the X - Y plane, for example $A = X$ and $B = Y$, in order to extract two uniformly random bits from this state, since this is the only way to have $\langle A \rangle = \langle A \otimes B \rangle = \langle B \rangle = 0$. We would, however, intuitively expect the maximum violation of a Bell inequality on ψ_θ to be attained with measurements having a component in the Z direction, since the correlation terms involving Z in (2) are larger in magnitude than the analogous terms involving X and Y . As such, we anticipate that we will need a Bell experiment engineered to exploit the entire Bloch sphere.

To this end, we propose the following Bell test in which Alice and Bob perform ± 1 -valued measurements A_x , $x = 1, 2, 3$ and B_y , $y = 1, \dots, 6$, in each round. They use the statistics to estimate the values of three Bell expressions.

The first two,

$$I_\beta = \langle \beta A_1 + A_1(B_1 + B_2) + A_2(B_1 - B_2) \rangle, \quad (3)$$

$$J_\beta = \langle \beta A_1 + A_1(B_3 + B_4) + A_3(B_3 - B_4) \rangle, \quad (4)$$

are modified CHSH expressions of the kind introduced in [7], while the third,

$$S = \langle A_2(B_5 + B_6) + A_3(B_5 - B_6) \rangle, \quad (5)$$

is an ordinary CHSH [10, 11] expression. We choose

$$\beta = \frac{2 \cos(\theta)}{\sqrt{1 + \sin(\theta)^2}} \quad (6)$$

for the value of the parameter β in the definitions of I_β and J_β , depending on the angle θ that identifies the intended state $|\psi_\theta\rangle$. Alice and Bob should in particular check that these Bell expressions attain the values

$$I_\beta = 2\sqrt{2}\sqrt{1 + \beta^2/4}, \quad (7)$$

$$J_\beta = 2\sqrt{2}\sqrt{1 + \beta^2/4}, \quad (8)$$

$$S = 2\sqrt{2}\sin(\theta). \quad (9)$$

The Bell expectation values (7), (8), and (9) can be attained by measuring

$$A_1 = Z, \quad A_2 = X, \quad A_3 = \pm Y \quad (10)$$

on Alice's side and performing suitable measurements on Bob's side on $|\psi_\theta\rangle$ [7]. Crucially for the intended application to randomness generation this is, up to trivial modifications such as local changes of bases and extensions to higher dimension, essentially the only way to attain these expectation values. More precisely, in appendix A we establish the following, which holds regardless of the Hilbert-space dimension.

Lemma 1. *The conditions $I_\beta = J_\beta = 2\sqrt{2}\sqrt{1 + \beta^2/4}$ and $S = 2\sqrt{2}\sin(\theta)$ identify an extremal point in the quantum set and if they are attained there is a choice of local bases in which:*

i) *the underlying state has the form*

$$\rho = \psi_\theta \otimes \sigma_{A'B'}, \quad (11)$$

where ψ_θ is the partially entangled state (2) and $\sigma_{A'B'}$ is an ancillary state which may be of any dimension;

ii) *Alice's measurements act on the state according to*

$$A_1 = Z \otimes \mathbb{1}_{A'}, \quad (12)$$

$$A_2 = X \otimes \mathbb{1}_{A'}, \quad (13)$$

$$A_3 = Y \otimes A' \quad (14)$$

where A' is a ± 1 -valued Hermitian operator;

iii) *Bob's measurements act on the state according to*

$$\frac{B_1 + B_2}{\sqrt{2\lambda_+}} = \frac{B_3 + B_4}{\sqrt{2\lambda_+}} = Z \otimes \mathbb{1}_{B'}, \quad (15)$$

$$\frac{B_1 - B_2}{\sqrt{2\lambda_-}} = \frac{B_5 + B_6}{\sqrt{2}} = X \otimes \mathbb{1}_{B'}, \quad (16)$$

$$-\frac{B_3 - B_4}{\sqrt{2\lambda_-}} = -\frac{B_5 - B_6}{\sqrt{2}} = Y \otimes B', \quad (17)$$

where $\lambda_{\pm} = 1 \pm \beta^2/4$ and B' is a ± 1 -valued Hermitian operator;

iv) the ancillary state $\sigma_{A'B'}$ in (11) and operators A' and B' are related in such a way that

$$\langle A' \otimes B' \rangle_{\sigma_{A'B'}} = 1. \quad (18)$$

The operators A' and B' appearing in lemma 1 are inevitable and reflect the fact that we cannot distinguish a set of qubit measurements from their complex conjugates on both sides [12]. We should also remark that, strictly speaking, *ii*) and *iii*) give the form of Alice's and Bob's measurements only on the supports of the respective marginals $\rho_A = \text{Tr}_B[\rho]$ and $\rho_B = \text{Tr}_A[\rho]$ of the underlying state. This is not a problem for us since any action the measurements may have on part of the Hilbert space not containing the state cannot have any impact on the correlations. In the following we will assume, for simplicity, that the marginals are of full rank.

Randomness with projective measurements.— Lemma 1 makes it straightforward to show that we can device-independently extract up to two bits of randomness using projective measurements. To do this, we simply add a seventh measurement, B_7 , to the Bell test on Bob's side and check that its correlation with A_2 is

$$\langle A_2 B_7 \rangle = \sin(\theta). \quad (19)$$

Using $A_2 = X \otimes \mathbb{1}_{A'}$ and $\rho = \psi_{\theta} \otimes \sigma_{A'B'}$ from lemma 1 and tracing out everything on Alice's side, we can rewrite the correlation on the left as

$$\langle A_2 B_7 \rangle = \sin(\theta) \text{Tr}[B_7 \frac{1}{2} X \otimes \sigma_{B'}]. \quad (20)$$

The operator $\frac{1}{2} X \otimes \sigma_{B'}$ has a trace norm of 1 and, since we are assuming $\sigma_{B'}$ is of full rank, the only way for the right sides of (19) and (20) to be equal is with

$$B_7 = X \otimes \mathbb{1}_{B'}. \quad (21)$$

With this information we can prove that the results of measuring A_3 and B_7 are maximally random. The probabilities of the four possible outcomes are

$$P(ab|37) = \frac{1}{4} \langle (\mathbb{1} + aA_3) \otimes (\mathbb{1} + bB_7) \rangle, \quad (22)$$

$a, b \in \{\pm 1\}$. Direct calculation with $A_3 = Y \otimes A'$ and $B_7 = X \otimes \mathbb{1}$ gives

$$P(ab|37) = \frac{1}{4}. \quad (23)$$

Importantly, the fact that we can derive $P(ab|37) = 1/4$ from $I_{\beta} = J_{\beta} = 2\sqrt{2}\sqrt{1+\beta^2/4}$, $S = 2\sqrt{2}\sin(\theta)$, and $\langle A_2 B_7 \rangle = \sin(\theta)$ shows that these conditions together are extremal, i.e., they cannot be attained by averaging quantum strategies that give different values of these quantities. This rules out the possibility of a more detailed underlying explanation of the correlations that might allow better predictions to be made about A_3 and B_7 .

Tomographic reconstruction of POVMs.— POVMs performed on qubit systems can have more than two outcomes and can potentially be used to generate more randomness than projective measurements. The nature of the device-independent scenario means we will only be interested in POVMs that are extremal, i.e., that cannot be expressed as convex combinations of other POVMs. The extremal qubit POVMs were classified in [9] and the only nontrivial ones consist of at most four rank-one elements $\alpha_a = |\alpha_a\rangle\langle\alpha_a|$ that are linearly independent.

We can certify the randomness of *some* POVMs device-independently by using a form of tomography to partially reconstruct them. To see how this works note first that, in the device-dependent setting, we can reconstruct any extremal qubit POVM $\{\alpha_a\}$ on (for example) Alice's side from the correlations it produces with the Pauli operators on Bob's side. That is, it turns out that the expectation values $\langle \alpha_a \otimes \sigma_{\nu} \rangle_{\psi_{\theta}}$, for $\sigma_{\nu} = (1, X, Y, Z)$ on Bob's side, contain sufficient information to uniquely identify $\{\alpha_a\}$ on Alice's side provided that the underlying state $|\psi_{\theta}\rangle$ is known.

In the device-independent scenario, we do not know that the quantum system we are manipulating is limited to a pair of qubits. However, according to lemma 1 we can verify that Alice is performing Pauli-type measurements up to complex conjugation, and the linear combinations of Bob's measurements in (15)–(17) effectively give us such operators on Bob's side. With this, we can check that a POVM $\{R_a\}$ on (for example, again) Alice's side produces correlations consistent with an extremal qubit one, i.e., that

$$\langle R_a \otimes B_{\nu} \rangle_{\psi_{\theta} \otimes \sigma} = \langle \alpha_a \otimes \sigma_{\nu} \rangle_{\psi_{\theta}}, \quad (24)$$

with $B_{\nu} = (\mathbb{1} \otimes \mathbb{1}, X \otimes \mathbb{1}, Y \otimes B', Z \otimes \mathbb{1})$, where $\{\alpha_a\}$ is some ideal reference qubit POVM. In appendix B, we prove that this allows us to infer the following on the form of $\{R_a\}$.

Lemma 2. *If the correlations obtained from a POVM $\{R_a\}$ match those obtainable from an extremal reference qubit POVM $\{\alpha_a\}$ according to (24), then the elements R_a must be of the form*

$$R_a = \alpha_a \otimes A'_+ + \alpha_a^* \otimes A'_- + |\alpha_a\rangle\langle\alpha_a^*| \otimes K'_a + |\alpha_a^*\rangle\langle\alpha_a| \otimes K_a'^{\dagger}, \quad (25)$$

where $|\alpha_a^*\rangle$ is the complex conjugate of $|\alpha_a\rangle$, $A'_{\pm} = (\mathbb{1} \pm A')/2$ are projectors on the positive and negative parts of A' from lemma 1, and the K'_a satisfy the operator inequalities

$$K'_a K_a'^{\dagger} \leq A'_+, \quad K_a'^{\dagger} K'_a \leq A'_- \quad (26)$$

and the condition

$$\sum_a |\alpha_a\rangle\langle\alpha_a^*| \otimes K'_a = 0. \quad (27)$$

Furthermore, if $\{R_a\}$ has three outcomes or less then $K'_a = 0$ and (25) simplifies to

$$R_a = \alpha_a \otimes A'_+ + \alpha_a^* \otimes A'_-. \quad (28)$$

In other words, our Bell test allows us to reconstruct, up to complex conjugation, extremal POVMs with two or three outcomes but we can only partially constrain the form of POVMs with four outcomes. As we elaborate on in the appendix, the place where the number of outcomes makes a difference is in the condition (27): when there are three outcomes or less, the off-diagonal $|\alpha_a\rangle\langle\alpha_a^*|$ terms in (25) are always linearly independent and thus (27) can only be satisfied with $K'_a = 0$. On the other hand, a simple calculation shows that $\langle\alpha^*|Y|\alpha\rangle = 0$ for any qubit state vector; this means that the $|\alpha_a\rangle\langle\alpha_a^*|$ s are restricted to the three-dimensional space of operators spanned by $\{\mathbb{1}, X, Z\}$ and they can never be linearly independent if there are four of them. In that case it is always possible to satisfy (27) with nonzero K'_a s.

Randomness with POVMs.— As we stated earlier, the maximum amount of randomness that could potentially be generated if both parties use extremal POVMs is limited to four bits. It is indeed possible to find extremal qubit POVMs that can generate arbitrarily close to this amount of randomness from any partially entangled state $|\psi_\theta\rangle$. Unfortunately, the fact that we cannot fully self-test POVMs means that this bound is not attainable in the device-independent setting. To see this, let us suppose that Alice and Bob unknowingly try to generate their random results using four-outcome POVMs $\{R_a\}$ and $\{S_b\}$ which are related to some ideal extremal qubit POVMs $\{\alpha_a\}$ and $\{\beta_b\}$ by

$$\begin{aligned} R_a &= \alpha_a \otimes |+\rangle\langle +|_{A'} + \alpha_a^* \otimes |-\rangle\langle -|_{A'} \\ &\quad + \lambda_a |\alpha_a\rangle\langle\alpha_a^*| \otimes |+\rangle\langle -|_{A'} \\ &\quad + \lambda_a^* |\alpha_a^*\rangle\langle\alpha_a| \otimes |-\rangle\langle +|_{A'}, \end{aligned} \quad (29)$$

$$\begin{aligned} S_b &= \beta_b \otimes |+\rangle\langle +|_{B'} + \beta_b^* \otimes |-\rangle\langle -|_{B'} \\ &\quad + \mu_b |\beta_b\rangle\langle\beta_b^*| \otimes |+\rangle\langle -|_{B'} \\ &\quad + \mu_b^* |\beta_b^*\rangle\langle\beta_b| \otimes |-\rangle\langle +|_{B'}, \end{aligned} \quad (30)$$

where λ_a and μ_b are some complex coefficients with magnitudes less than 1, while an eavesdropper at each round randomly and equiprobably chooses and prepares one of two states $\psi_\theta \otimes \chi'_+$ or $\psi_\theta \otimes \chi'_-$ with different ancillary parts $|\chi'_\pm\rangle = (|++\rangle \pm |--\rangle)/\sqrt{2}$. Using that $\langle\alpha^*\beta^*|\psi_\theta\rangle = \langle\alpha\beta|\psi_\theta\rangle^*$, we can work out that the joint probability of Alice's and Bob's outcomes, conditioned on either ancillary state being chosen by Eve, is

$$\begin{aligned} \langle R_a \otimes S_b \rangle_{\psi_\theta \otimes \chi'_\pm} &= |\langle\alpha_a\beta_b|\psi_\theta\rangle|^2 \pm \text{Re}[\lambda_a^* \mu_b^* \langle\alpha_a\beta_b|\psi_\theta\rangle^2]. \end{aligned} \quad (31)$$

These probabilities average out to the ideal joint probabilities $|\langle\alpha_a\beta_b|\psi_\theta\rangle|^2$ that would be obtained from the reference qubit POVMs on $|\psi_\theta\rangle$; hence, Alice and Bob have no way to detect, device independently, that they are measuring $\{R_a\}$ and $\{S_b\}$ rather than $\{\alpha_a\}$ and $\{\beta_b\}$. Eve, however, knowing which ancilla state she chose, also knows which of the two joint distributions in (31) was actually prepared in each round and can use this to make a more informed guess about what the outcome will be.

Let us see how this could help Eve in the worst case. As we pointed out above, the off-diagonal terms $|\alpha_a\rangle\langle\alpha_a^*|$

and $|\beta_b\rangle\langle\beta_b^*|$ are never linearly independent and, thus, the coefficients λ_a and μ_b can be chosen nonzero. We are free to scale them such that the largest coefficient on each side is of magnitude one. By also exploiting the freedom to choose their phases we can arrange that, for at least one pair (a, b) of outputs, $\text{Re}[\lambda_a^* \mu_b^* \langle\alpha_a\beta_b|\psi_\theta\rangle^2] = |\langle\alpha_a\beta_b|\psi_\theta\rangle|^2$. In other words, we are certain we can arrange for at least one of the probabilities $\langle R_a \otimes S_b \rangle_{\psi_\theta \otimes \chi'_\pm}$ to be zero. This means that the probability of the most likely joint outcome, conditioned on Eve choosing $|\chi'_-\rangle$, cannot be lower than $1/15$. It follows that the randomness that can be certified device-independently for the entire protocol can never be higher than

$$-\log_2 \left[\frac{1}{2} \left(\frac{1}{15} + \frac{1}{16} \right) \right] \approx 3.9527 \text{ bits} \quad (32)$$

regardless of the state and POVMs that Alice and Bob try to use.

On a more positive note, the above-described complication does not manifest if only one of the parties uses a measurement with four outcomes and, in that case, the amount of randomness that can be generated device-independently is the same as the amount of randomness that can be generated using extremal qubit POVMs performed on $|\psi_\theta\rangle$. This means it is potentially possible to generate up to two bits of randomness locally, or alternatively potentially up to

$$-\log_2(1/12) \approx 3.5850 \text{ bits} \quad (33)$$

of randomness globally using a four-outcome POVM on one side and a three-outcome POVM on the other. We give explicit constructions of POVMs that yield these amounts of randomness (or arbitrarily close) in appendix C.

Conclusion.— Our work reinforces the observation that the amount of randomness obtainable from a quantum system does not in general increase with the degree of entanglement. In two versions of the problem, we have confirmed that an upper limit of two bits of randomness is always obtainable from any partially entangled pure state of two qubits. In the global case using POVMs, although we do not know the optimal amount of extractable randomness we have significantly narrowed the range to between about 3.58 and 3.96 bits for any state. The nontrivial latter limit establishes that the upper bound of four bits is not attainable in this case.

Our results rely on the fact that we can reconstruct the underlying quantum state and measurements in our Bell test sufficiently well to conclude that the outcomes are genuinely random. This adds to a growing literature showing that we can often infer substantial information about the quantum resources available from a Bell test [13–17]. Previous work has notably shown that the partially entangled state [18, 19] or measurements spanning the entire Bloch sphere (up to complex conjugation) [8, 20, 21] can be self-tested, although before now not together in the same test.

Our work also led us to investigate whether it is possible to self-test nonprojective measurements in quantum

physics and we found that qubit POVMs with four outcomes can only be self-tested to a limited extent. The ambiguity with respect to complex conjugation can thus, as we found here, make a significant difference in the device-independent setting. It will be interesting to further explore this problem, both for qubit systems and in higher dimension. In particular, closing the gap on optimal randomness generation with POVMs is likely to require developing a better understanding of the general form that we found POVMs may take in lemma 2.

Note added.— While completing this work, we learned that the authors of [22] had independently found using a similar approach that two bits of randomness can be generated globally using projective measurements from the partially entangled state.

Acknowledgements.— This work was supported by the Spanish MINECO (Severo Ochoa grant SEV-2015-0522), the Generalitat de Catalunya (CERCA Program and SGR 1381), the Fundació Privada Cellex and Mir-Puig, the AXA Chair in Quantum Information Science, the ERC AdG CERQUITE and CoG QITBOX, and the EU Quantum Flagship project QRANGE. J. K. acknowledges support from the National Science Centre, Poland (grant no. 2016/23/P/ST2/02122). This project is carried out under POLONEZ programme which has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 665778. B. B. acknowledges support from the Secretaria d’Universitats i Recerca del Departament d’Economia i Coneixement de la Generalitat de Catalunya and the European Social Fund (FEDER). R. A. acknowledges support from the Foundation for Polish Science through the First Team project (First TEAM/2017-4/31) cofinanced by the European Union under the European Regional Development Fund.

REFERENCES

- [1] T. Norsen, *Am. J. Phys.* **79**, 1261 (2011).
- [2] J. S. Bell, *Physics* **1**, 195 (1964).
- [3] A. Acín, N. Brunner, N. Gisin, S. Massar, S. Pironio, and V. Scarani, *Phys. Rev. Lett.* **98**, 230501 (2007), [arXiv:quant-ph/0702152](#).
- [4] R. Colbeck, *Quantum And Relativistic Protocols For Secure Multi-Party Computation*, Ph.D. thesis, University of Cambridge (2006), [arXiv:0911.3814 \[quant-ph\]](#).
- [5] S. Pironio, A. Acín, S. Massar, A. Boyer de La Giroday, D. N. Matsukevich, P. Maunz, S. Olmschenk, D. Hayes, L. Luo, T. A. Manning, and C. Monroe, *Nature* **464**, 1021 (2010), [arXiv:0911.3427 \[quant-ph\]](#).
- [6] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, *Rev. Mod. Phys.* **86**, 419 (2014), [arXiv:1303.2849 \[quant-ph\]](#).
- [7] A. Acín, S. Massar, and S. Pironio, *Phys. Rev. Lett.* **108**, 100402 (2012), [arXiv:1107.2754 \[quant-ph\]](#).
- [8] A. Acín, S. Pironio, T. Vértesi, and P. Wittek, *Phys. Rev. A* **93**, 040102(R) (2016), [arXiv:1505.03837 \[quant-ph\]](#).
- [9] G. M. D’Ariano, P. Lo Presti, and P. Perinotti, *J. Phys. A: Math. Gen.* **38**, 5979 (2005), [arXiv:quant-ph/0408115](#).
- [10] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, *Phys. Rev. Lett.* **23**, 880 (1969).
- [11] B. S. Cirel’son, *Lett. Math. Phys.* **4**, 93 (1980).
- [12] M. McKague and M. Mosca, in *Theory of Quantum Computation, Communication, and Cryptography*, Lecture Notes in Computer Science, Vol. 6519 (Springer Berlin Heidelberg, Berlin, Heidelberg, 2011) pp. 113–130, [arXiv:1006.0150 \[quant-ph\]](#).
- [13] L. A. Khalfin and B. S. Tsirelson, in *Symposium on the Foundations of Modern Physics* (World Scientific, Singapore, 1985) pp. 441–460.
- [14] S. L. Braunstein, A. Mann, and M. Revzen, *Phys. Rev. Lett.* **68**, 3259 (1992).
- [15] V. Scarani and N. Gisin, *J. Phys. A: Math. Gen.* **34**, 6043 (2001), [arXiv:quant-ph/0103068](#).
- [16] D. Mayers and A. Yao, *Quantum Inf. Comput.* **4**, 273 (2004), [arXiv:quant-ph/0307205](#).
- [17] M. McKague, T. H. Yang, and V. Scarani, *J. Phys. A: Math. Theor.* **45**, 455304 (2012), [arXiv:1203.2976 \[quant-ph\]](#).
- [18] T. H. Yang and M. Navascués, *Phys. Rev. A* **87**, 050102(R) (2013), [arXiv:1210.4409](#).
- [19] C. Bamps and S. Pironio, *Phys. Rev. A* **91**, 052111 (2015), [arXiv:1504.06960 \[quant-ph\]](#).
- [20] J. Kaniewski, *Phys. Rev. A* **95**, 062323 (2017), [arXiv:1702.06845 \[quant-ph\]](#).
- [21] O. Andersson, P. Badziąg, I. Bengtsson, I. Dumitru, and A. Cabello, *Phys. Rev. A* **96**, 032119 (2017), [arXiv:1706.02130 \[quant-ph\]](#).
- [22] R. Ramanathan and S. Pironio, (unpublished).

A. SELF-TEST OF STATE AND MEASUREMENTS

The core behind our randomness results in the main text is a self-test, lemma 1, that we prove here. To recapitulate, we let

$$I_\beta = \langle \beta A_1 + A_1(B_1 + B_2) + A_2(B_1 - B_2) \rangle, \quad (34)$$

$$J_\beta = \langle \beta A_1 + A_1(B_3 + B_4) + A_3(B_3 - B_4) \rangle, \quad (35)$$

$$S = \langle A_2(B_5 + B_6) + A_3(B_5 - B_6) \rangle \quad (36)$$

denote the expectation values of three Bell operators involving three ± 1 -valued observables A_x on Alice's side and six ± 1 -valued observables B_y on Bob's side acting on a shared underlying quantum system. We suppose, furthermore, that the expectation values

$$I_\beta = 2\sqrt{2}\sqrt{1 + \beta^2/4}, \quad (37)$$

$$J_\beta = 2\sqrt{2}\sqrt{1 + \beta^2/4}, \quad (38)$$

$$S = 2\sqrt{2}\sin(\theta) \quad (39)$$

are attained, where θ is related to the parameter β by

$$\beta = \frac{2\cos(\theta)}{\sqrt{1 + \sin^2(\theta)}}. \quad (40)$$

According to lemma 1, this is only possible if there is a choice of bases in which the quantum state is of the form

$$\rho = \psi_\theta \otimes \sigma_{A'B'}, \quad (41)$$

where

$$\begin{aligned} \psi_\theta = \frac{1}{4} \Big[& \mathbb{1} \otimes \mathbb{1} + \cos(\theta)(Z \otimes \mathbb{1} + \mathbb{1} \otimes Z) \\ & + \sin(\theta)(X \otimes X - Y \otimes Y) + Z \otimes Z \Big], \end{aligned} \quad (42)$$

and Alice's and Bob's observables act on the supports of their marginals of ρ according to

$$A_1 = Z \otimes \mathbb{1}, \quad A_2 = X \otimes \mathbb{1}, \quad A_3 = Y \otimes A' \quad (43)$$

and

$$\frac{B_1 + B_2}{\sqrt{2\lambda_+}} = \frac{B_3 + B_4}{\sqrt{2\lambda_+}} = Z \otimes \mathbb{1}, \quad (44)$$

$$\frac{B_1 - B_2}{\sqrt{2\lambda_-}} = \frac{B_5 + B_6}{\sqrt{2}} = X \otimes \mathbb{1}, \quad (45)$$

$$-\frac{B_3 - B_4}{\sqrt{2\lambda_-}} = -\frac{B_5 - B_6}{\sqrt{2}} = Y \otimes B', \quad (46)$$

where A' and B' are ± 1 -valued Hermitian operators and $\lambda_\pm = 1 \pm \beta^2/4$. Furthermore, the ancillary state $\sigma_{A'B'}$ in (41) and operators A' and B' are such that

$$\text{Tr}[(A' \otimes B')\sigma_{A'B'}] = 1. \quad (47)$$

We will proceed by deriving constraints on the state and measurements implied by each of the three Bell expectation values (37), (38), and (39) in turn.

A. I_β constraint

The first condition $I_\beta = 2\sqrt{2}\sqrt{1 + \beta^2/4}$ alone already allows us to self-test the state as well as the measurements A_1 , A_2 , B_1 , and B_2 . This was essentially proved in the course of deriving the Tsirelson bound for the more general family of I_α^β expressions done in [7], particularly the steps around Eqs. (14)–(16). (The result is also closely related to the self-test based on I_β in [19], although the formulation is slightly different.) Since [7] is not very explicit about this we will give an alternative and explicit proof here.

I_β is the expectation value of its corresponding Bell operator,

$$\mathcal{I}_\beta = \beta A_1 + A_1(B_1 + B_2) + A_2(B_1 - B_2). \quad (48)$$

Without loss of generality, we take the measurements to be projective, such that $A_x^2 = B_y^2 = \mathbb{1}$. This allows us to use the Jordan lemma to decompose the measurements as

$$A_x = \sum_j A_{x|j} \otimes |j'\rangle\langle j'| \oplus A_{x\perp}, \quad (49)$$

$$B_y = \sum_k B_{y|k} \otimes |k'\rangle\langle k'| \oplus B_{y\perp}, \quad (50)$$

where we absorb any 1×1 Jordan blocks as well as 2×2 blocks where one of the measurements is ± 1 into $A_{x\perp}$ and $B_{y\perp}$. In other words, $A_{x\perp}$ and $B_{y\perp}$ contain the Jordan blocks that we already know cannot contribute to a violation of the I_β inequality because at least one of the measurements in the blocks is deterministic. We correspondingly decompose the Bell operator as

$$\mathcal{I}_\beta = \sum_{jk} \mathcal{I}_{\beta|jk} \otimes |j'k'\rangle\langle j'k'| \oplus \mathcal{I}_{\beta\perp}, \quad (51)$$

where

$$\begin{aligned} \mathcal{I}_{\beta|jk} = & \beta A_{1|j} + A_{1|j}(B_{1|k} + B_{2|k}) \\ & + A_{2|j}(B_{1|k} - B_{2|k}) \end{aligned} \quad (52)$$

and $\mathcal{I}_{\beta\perp}$ is the part of $\mathcal{I}_\beta$ containing ' $\perp$ ' blocks on one or the other side or both.

In order to attain the quantum bound, the measurements $A_{x|j}$ and $B_{y|k}$ in each block must be optimised such that every $\mathcal{I}_{\beta|jk}$ has an eigenvalue equal to $2\sqrt{2}\sqrt{1 + \beta^2/4}$ or the underlying state must have zero presence in the corresponding part of the Hilbert space. From now, let us redefine $A_{x\perp}$ and $B_{y\perp}$ to also absorb Jordan blocks where the state has no presence and remove them from further consideration. The remaining blocks must have measurements optimised to attain the quantum bound.

Let us concentrate on a particular pair of remaining blocks (j, k) and drop the indices jk while we do this to lighten the notation. Squaring $\mathcal{I}_\beta$ and using that $A_x^2 = B_y^2 = \mathbb{1}$ gives

$$\begin{aligned} \mathcal{I}_\beta^2 = & (4 + \beta^2)\mathbb{1} + 2\beta\mathbb{1} \otimes (B_1 + B_2) \\ & + \beta\{A_1, A_2\} \otimes (B_1 - B_2) \\ & - [A_1, A_2] \otimes [B_1, B_2]. \end{aligned} \quad (53)$$

We are free to choose the bases on both sides in such a way that

$$A_1 = Z, \quad (54)$$

$$A_2 = \cos(\lambda)Z - \sin(\lambda)X, \quad (55)$$

$$B_1 + B_2 = 2\cos(\frac{\mu}{2})Z, \quad (56)$$

$$B_1 - B_2 = 2\sin(\frac{\mu}{2})X \quad (57)$$

with all of the coss and sins nonnegative. With this choice,

$$\begin{aligned} \mathcal{I}_\beta^2/4 &= (1 + \beta^2/4)\mathbb{1} + \beta\cos(\frac{\mu}{2})\mathbb{1} \otimes Z \\ &+ \beta\cos(\lambda)\sin(\frac{\mu}{2})\mathbb{1} \otimes X \\ &+ \sin(\lambda)\sin(\mu)Y \otimes Y. \end{aligned} \quad (58)$$

Notice here that only $\mathbb{1}$ and Y , which are simultaneously diagonalisable, appear on Alice's side. We can use this to identify the eigenvalues of $\mathcal{I}_\beta^2/4$. The largest one (with multiplicity two) is

$$\begin{aligned} &1 + \beta^2/4 \\ &+ \sqrt{\beta^2\cos(\frac{\mu}{2})^2 + \beta^2\cos(\lambda)^2\sin(\frac{\mu}{2})^2 + \sin(\lambda)^2\sin(\mu)^2}. \end{aligned} \quad (59)$$

This is maximised with either $\cos(\lambda)^2 = 1$ or $\sin(\lambda)^2 = 1$, depending on whether $\beta^2\sin(\mu/2)^2$ or $\sin(\mu)^2$ is larger. We can ignore the former since it would mean that A_1 and A_2 commute and we would have already absorbed them into $A_{x\perp}$, and thus use $\sin(\lambda) = 1$ which fixes $A_2 = X$. The term under the square root then is

$$\beta^2\cos(\frac{\mu}{2})^2 + 4\cos(\frac{\mu}{2})^2\sin(\frac{\mu}{2})^2 \quad (60)$$

which we can rewrite as

$$(1 + \beta^2/4)^2 - \left(2\cos(\frac{\mu}{2})^2 - (1 + \beta^2/4)\right)^2. \quad (61)$$

This is maximised with

$$\cos(\frac{\mu}{2}) = \sqrt{\frac{1 + \beta^2/4}{2}}, \quad (62)$$

which is less than one in the range $\beta < 2$ where I_β has a quantum violation, for which the maximum eigenvalues (59) of $\mathcal{I}_\beta^2/4$ become $2(1 + \beta^2/4)$.

Since the other two eigenvalues for the optimal measurements are zero and $\mathcal{I}_\beta$ is traceless we can safely infer that its only two nonzero eigenvalues are $\pm 2\sqrt{2}\sqrt{1 + \beta^2/4}$ and we can identify the corresponding eigenstates. Substituting in the now-known-to-be-optimal qubit measurements, the Bell operator in our Jordan block is

$$\begin{aligned} \mathcal{I}_\beta &= \beta Z \otimes \mathbb{1} + \sqrt{2}\sqrt{1 + \beta^2/4} Z \otimes Z \\ &+ \sqrt{2}\sqrt{1 - \beta^2/4} X \otimes X. \end{aligned} \quad (63)$$

To identify the eigenstates we express $\mathcal{I}_\beta$ in spectral form as

$$\mathcal{I}_\beta = 2\sqrt{2}\sqrt{1 + \beta^2/4}(\psi_\theta - \phi_\theta) \quad (64)$$

where

$$\begin{aligned} \psi_\theta &= \frac{1}{4} \left[\mathbb{1} \otimes \mathbb{1} + \cos(\theta)(Z \otimes \mathbb{1} + \mathbb{1} \otimes Z) \right. \\ &\quad \left. + \sin(\theta)(X \otimes X - Y \otimes Y) + Z \otimes Z \right], \end{aligned} \quad (65)$$

$$\begin{aligned} \phi_\theta &= \frac{1}{4} \left[\mathbb{1} \otimes \mathbb{1} - \cos(\theta)(Z \otimes \mathbb{1} - \mathbb{1} \otimes Z) \right. \\ &\quad \left. - \sin(\theta)(X \otimes X + Y \otimes Y) - Z \otimes Z \right] \end{aligned} \quad (66)$$

are rank-one projectors corresponding to pure states

$$|\psi_\theta\rangle = \cos(\frac{\theta}{2})|00\rangle + \sin(\frac{\theta}{2})|11\rangle, \quad (67)$$

$$|\phi_\theta\rangle = \sin(\frac{\theta}{2})|01\rangle - \cos(\frac{\theta}{2})|10\rangle. \quad (68)$$

Precisely, the expressions (63) and (64) for $\mathcal{I}_\beta$ coincide if

$$\cos(\theta) = \sqrt{\frac{2\beta^2/4}{1 + \beta^2/4}}, \quad \sin(\theta) = \sqrt{\frac{1 - \beta^2/4}{1 + \beta^2/4}}, \quad (69)$$

which fixes how β and μ must be related to θ .

Returning back to the full Bell operator, in order to be able to attain its quantum bound we find that it must be of the form

$$\mathcal{I}_\beta = 2\sqrt{2}\sqrt{1 + \beta^2/4}(\psi_\theta - \phi_\theta) \otimes \mathbb{1}_{A'B'} \oplus \mathcal{I}_{\beta\perp}, \quad (70)$$

and the underlying state must have no support on the $\mathcal{I}_{\beta\perp}$ part. The problem

$$\text{Tr}[\mathcal{I}_\beta \rho] = 2\sqrt{2}\sqrt{1 + \beta^2/4} \quad (71)$$

can thus be solved with, and only with, a state of the form

$$\rho = \psi_\theta \otimes \sigma_{A'B'}. \quad (72)$$

To verify this, we express an arbitrary state ρ in spectral form,

$$\rho = \sum_k q_k \Psi_k, \quad (73)$$

with $q_k > 0$. Then

$$\text{Tr}[\mathcal{I}_\beta \rho] = \sum_k p_k \text{Tr}[\mathcal{I}_\beta \Psi_k] = 2\sqrt{2}\sqrt{1 + \beta^2/4} \quad (74)$$

is only possible if

$$\text{Tr}[\mathcal{I}_\beta \Psi_k] = 2\sqrt{2}\sqrt{1 + \beta^2/4} \quad (75)$$

for all k . The left side is more precisely

$$\begin{aligned} \text{Tr}[\mathcal{I}_\beta \Psi_k] &= 2\sqrt{2}\sqrt{1 + \beta^2/4} \text{Tr}[(\psi_\theta - \phi_\theta) \otimes \mathbb{1}_{A'B'} \Psi_k] \\ &+ \text{Tr}[\mathcal{I}_{\beta\perp} \Psi_k]. \end{aligned} \quad (76)$$

From this we can see that, to attain the Tsirelson bound, the second term $\text{Tr}[\mathcal{I}_{\beta\perp} \Psi_k]$ must be zero and we must have

$$\begin{aligned} &\text{Tr}[(\psi_\theta - \phi_\theta) \otimes \mathbb{1}_{A'B'} \Psi_k] \\ &= \langle \psi_\theta | \Psi_{2|k} | \psi_\theta \rangle - \langle \phi_\theta | \Psi_{2|k} | \phi_\theta \rangle \\ &= 1, \end{aligned} \quad (77)$$

where $\Psi_{2|k} = \text{Tr}_{A'B'}[\Psi_k]$ in the second expression is the qubit marginal of Ψ_k . The above problem is only solvable if $\Psi_{2|k} = \psi_\theta$, i.e., if the $|\Psi_k\rangle$ are of the form

$$|\Psi_k\rangle = |\psi_\theta\rangle \otimes |\chi'_k\rangle \quad (78)$$

for all k . This finally gives

$$\rho = \psi_\theta \otimes \left(\sum_k q_k \chi'_k \right) = \psi_\theta \otimes \sigma_{A'B'}. \quad (79)$$

Furthermore, the marginal states $\sigma_{A'}$ and $\sigma_{B'}$ are of full rank in the Hilbert spaces we implicitly introduced to contain the Jordan block indices, since we absorbed blocks where the state has no presence into $\mathcal{I}_{\beta\perp}$.

In the remainder of this proof we will restrict our attention to the Hilbert space containing the state and take ρ to be such that its marginals ρ_A and ρ_B are of full rank.

B. J_β constraint

The second Bell expression,

$$J_\beta = \langle \beta A_1 + A_1(B_3 + B_4) + A_3(B_3 - B_4) \rangle, \quad (80)$$

is the same as I_β except with different measurements and the second condition $J_\beta = 2\sqrt{2}\sqrt{1 + \beta^2/4}$ implies analogous conditions for the measurements, although not necessarily in the same bases. In particular, the second condition implies that, like A_1 and A_2 , A_1 and A_3 must anticommute.

Having already identified ρ and fixed A_1 , we can derive the most general A_3 that anticommutes with A_1 . In general we may expand A_3 as

$$A_3 = \mathbb{1} \otimes A'_1 + X \otimes A'_X + Y \otimes A'_Y + Z \otimes A'_Z \quad (81)$$

for some Hermitian operators A'_1 , A'_X , A'_Y , and A'_Z . By imposing the conditions $A_3^2 = \mathbb{1} \otimes \mathbb{1}$ and $\{A_1, A_3\} = 0$ for $A_1 = Z \otimes \mathbb{1}$ on A_3 we derive constraints on the A' operators. To satisfy the constraints we find that we must take $A'_1 = A'_Z = 0$ and A_3 of the form

$$A_3 = X \otimes A'_X + Y \otimes A'_Y, \quad (82)$$

with A'_X and A'_Y Hermitian and satisfying

$$A'^2_X + A'^2_Y = \mathbb{1}, \quad (83)$$

$$[A'_X, A'_Y] = 0. \quad (84)$$

To state the same result differently, A_3 is of the form

$$A_3 = (\cos(\varphi_j)X + \sin(\varphi_j)Y) \otimes |j'\rangle\langle j'|, \quad (85)$$

which we obtain by explicitly decomposing the co-diagonal operators A'_X and A'_Y .

C. S constraint

As the final step in the proof of our self-test, we will now prove that satisfying the third condition,

$$S = \langle A_2(B_5 + B_6) + A_3(B_5 - B_6) \rangle = 2\sqrt{2}\sin(\theta), \quad (86)$$

forces us to set $A_X = 0$ in (82). We start by writing the expectation value explicitly as

$$S = \text{Tr}[(A_2 + A_3)B_5 + (A_2 - A_3)B_6] \psi_\theta \otimes \sigma_{A'B'}. \quad (87)$$

Using the form (42) of ψ_θ in terms of Pauli operators and (85) of A_3 and doing the trace on Alice's side first we can rewrite the CHSH expectation value as

$$S = \sin(\theta)(\text{Tr}[\rho_{B+}B_5] + \text{Tr}[\rho_{B-}B_6]), \quad (88)$$

where we introduce

$$\begin{aligned} \rho_{B\pm} &= \frac{1}{\sin(\theta)} \text{Tr}_A[(A_2 \pm A_3) \psi_\theta \otimes \sigma_{A'B'}] \\ &= \frac{1}{2} \sum_j p_j ((1 \mp c_j)X \mp s_j Y) \otimes \sigma_{B'|j} \end{aligned} \quad (89)$$

and, in turn, c_j and s_j are shorthand for $\cos(\varphi_j)$ and $\sin(\varphi_j)$ and p_j and $\sigma_{B'|j}$ are defined such that

$$p_j \sigma_{B'|j} = \text{Tr}_{A'}[|j'\rangle\langle j'| \otimes \mathbb{1}_{B'}] \sigma_{A'B'} \quad (90)$$

and $\text{Tr}[\sigma_{B'|j}] = 1$. Note that this implies $\sum_j p_j = 1$. We can then upper bound the CHSH expectation value by

$$\begin{aligned} \frac{S}{\sin(\theta)} &= \text{Tr}[\rho_{B+}B_5] + \text{Tr}[\rho_{B-}B_6] \\ &\leq \|\rho_{B+}\|_1 + \|\rho_{B-}\|_1 \\ &\leq \sum_j p_j \left(\frac{1}{2} \|(1 - c_j)X - s_j Y\|_1 \|\sigma_{B'|j}\|_1 \right. \\ &\quad \left. + \frac{1}{2} \|(1 + c_j)X + s_j Y\|_1 \|\sigma_{B'|j}\|_1 \right) \\ &= \sum_j p_j \left(\sqrt{(1 - c_j)^2 + s_j^2} + \sqrt{(1 + c_j)^2 + s_j^2} \right) \\ &= \sum_j p_j \left(\sqrt{2 - 2c_j} + \sqrt{2 + 2c_j} \right) \\ &\leq \sum_j p_j \sqrt{2} \sqrt{(2 - 2c_j) + (2 + 2c_j)} \\ &= 2\sqrt{2}. \end{aligned} \quad (91)$$

To determine the optimal A_3 , observe that to attain (91) all of the inequalities applied to derive it must hold with equality. In particular, equality between the third and second last lines requires

$$\sqrt{2 + 2c_j} = \sqrt{2 - 2c_j}, \quad (92)$$

or $c_j = 0$ and $s_j = \pm 1$. We deduce that

$$A_3 = Y \otimes A' \quad (93)$$

where $A' = A'_+ - A'_-$ is the difference of two orthogonal projectors that sum to the identity. We can then extract B_5 and B_6 by reevaluating CHSH with

$$A_2 = X \otimes (A'_+ + A'_-), \quad (94)$$

$$A_3 = Y \otimes (A'_+ - A'_-). \quad (95)$$

The result can be written

$$\frac{S}{\sin(\theta)} = \frac{1}{2} \text{Tr}[\rho_{B+} B_5] + \frac{1}{2} \text{Tr}[\rho_{B-} B_6] \quad (96)$$

where

$$\rho_{B'\pm} = p_+(X \mp Y) \otimes \sigma_{B'|+} + p_-(X \pm Y) \otimes \sigma_{B'|-} \quad (97)$$

and in turn

$$p_{\pm} \sigma_{B'|\pm} = \text{Tr}_{A'}[A'_{\pm} \sigma_{A'B'}] \quad (98)$$

are Bob's partial traces conditioned on projection on the $A'_{\pm}$ subspaces on Alice's side. Satisfying the CHSH constraint requires that B_5 , B_6 , and $\sigma_{B'|\pm}$ be such that

$$\frac{1}{2} \text{Tr}[\rho_{B+} B_5] = \frac{1}{2} \|\rho_{B+}\|_1 = \sqrt{2}, \quad (99)$$

$$\frac{1}{2} \text{Tr}[\rho_{B-} B_6] = \frac{1}{2} \|\rho_{B-}\|_1 = \sqrt{2}. \quad (100)$$

This is only possible if $\sigma_{B'|\pm}$ have orthogonal support and

$$B_5 = \frac{X-Y}{\sqrt{2}} \otimes B'_+ + \frac{X+Y}{\sqrt{2}} \otimes B'_-, \quad (101)$$

$$B_6 = \frac{X+Y}{\sqrt{2}} \otimes B'_+ + \frac{X-Y}{\sqrt{2}} \otimes B'_-, \quad (102)$$

where $B'_{\pm}$ are orthogonal projectors on the supports of $\sigma_{B'|\pm}$. The above forms for B_5 and B_6 can alternatively be written

$$B_5 = \frac{1}{\sqrt{2}} (X \otimes \mathbb{1} - Y \otimes B'), \quad (103)$$

$$B_6 = \frac{1}{\sqrt{2}} (X \otimes \mathbb{1} + Y \otimes B') \quad (104)$$

with $B' = B'_+ - B'_-$, and orthogonality of $\sigma_{B'|\pm}$ compactly as the condition

$$\text{Tr}[(A' \otimes B') \sigma_{A'B'}] = 1 \quad (105)$$

on $\sigma_{A'B'}$.

Similarly, computing the previous Bell expression, J_{β} , with $A_1 = Z \otimes \mathbb{1}_{A'}$, $A_3 = Y \otimes A'$, and $\rho = \psi_{\theta} \otimes \sigma_{A'B'}$ allows the optimal measurements B_3 and B_4 to be identified as those given at the beginning of this section.

B. SELF-TESTING EXTREMAL POVMS

The self-test we have described, which among other things allows the Pauli measurements X , Y , and Z to be identified up to complex conjugation, allows us to perform a form of tomography with nonprojective measurements. It is stated as lemma 2 in the main text. To recall the problem: we suppose that Alice performs a POVM $\{R_a\}$ and Bob measures

$$B_{\nu} = (\mathbb{1}, X \otimes \mathbb{1}, Y \otimes B', Z \otimes \mathbb{1}) \quad (106)$$

on the state $\psi_{\theta} \otimes \sigma_{A'B'}$ from our Bell test, and we check that the correlations obtained match those that could be obtained from an ideal reference qubit POVM, i.e., that

$$\langle R_a \otimes B_{\mu} \rangle_{\psi_{\theta} \otimes \sigma} = \langle \alpha_a \otimes \sigma_{\nu} \rangle_{\psi_{\theta}}, \quad (107)$$

where $\{\alpha_a\}$ is a given extremal qubit POVM and $\sigma_{\nu} = (\mathbb{1}, X, Y, Z)$. We will prove here that this implies that the POVM elements R_a must be of the form asserted in lemma 2 in the main text, i.e.,

$$R_a = \alpha_a \otimes A'_+ + \alpha_a^* \otimes A'_- + |\alpha_a\rangle\langle\alpha_a^*| \otimes K'_a + |\alpha_a^*\rangle\langle\alpha_a| \otimes K_a'^{\dagger}, \quad (108)$$

where $|\alpha_a^*\rangle$ is the complex conjugate of $|\alpha_a\rangle$, $A'_{\pm} = (\mathbb{1} \pm A')/2$, and the operators K'_a are such that

$$K'_a K_a'^{\dagger} \leq A'_+, \quad K_a'^{\dagger} K'_a \leq A'_- \quad (109)$$

and

$$\sum_a |\alpha_a\rangle\langle\alpha_a^*| \otimes K'_a = 0. \quad (110)$$

To begin with, we remark that we can always decompose the reference qubit POVM $\{\alpha_a\}$ in the basis of the identity and Pauli operators as

$$\alpha_a = r_a^{\mu} \sigma_{\mu} \quad (111)$$

for some coefficients r_a^{μ} , where we use implicit summation over the repeated Greek index μ . The correlation on the left side of (107) can be written as

$$\langle \alpha_a \otimes \sigma_{\nu} \rangle = \eta_{\mu\nu} r_a^{\mu} \quad (112)$$

where $\eta_{\mu\nu} = \langle \sigma_{\mu} \otimes \sigma_{\nu} \rangle$ make up the components of a matrix that can be read off the expression (42) of ψ_{θ} in terms of the Pauli operators. Importantly, for $\theta \neq 0$, the matrix $(\eta_{\mu\nu})$ is invertible (for example, its determinant is $-\sin(\theta)^4$), so that the correlations $\langle \alpha_a \otimes \sigma_{\nu} \rangle$ uniquely identify the coefficients r_a^{μ} and the POVM elements α_a . We can then combine (111) and (112) to express the POVM elements directly in terms of the correlations as

$$\alpha_a = \sigma^{\nu} \langle \alpha_a \otimes \sigma_{\nu} \rangle_{\psi_{\theta}}, \quad (113)$$

where σ^{ν} are operators defined to be related to σ_{μ} by $\eta_{\mu\nu} \sigma^{\nu} = \sigma_{\mu}$.

By hypothesis, the correlations in (113) are the same as those obtained with R_a and B_{ν} according to (107); we can thus reexpress α_a as

$$\alpha_a = \sigma^{\nu} \langle R_a \otimes B_{\nu} \rangle_{\psi_{\theta} \otimes \sigma_{A'B'}}. \quad (114)$$

Now we note that we can always decompose R_a as

$$R_a = \sigma_{\mu} \otimes R_a'^{\mu} \quad (115)$$

and we can write the operators B_{ν} together as

$$B_{\nu} = \sigma_{\nu} \otimes B'_+ + \sigma_{\nu}^* \otimes B'_-. \quad (116)$$

Using these and developing,

$$\begin{aligned} \alpha_a &= \sigma^{\nu} \langle \sigma_{\mu} \otimes \sigma_{\nu} \rangle_{\psi_{\theta}} \langle R_a'^{\mu} \otimes B'_+ \rangle_{\sigma_{A'B'}} \\ &\quad + \sigma^{\nu} \langle \sigma_{\mu} \otimes \sigma_{\nu}^* \rangle_{\psi_{\theta}} \langle R_a'^{\mu} \otimes B'_- \rangle_{\sigma_{A'B'}} \\ &= \sigma^{\nu} \eta_{\mu\nu} \langle A'_+ R_a'^{\mu} A'_+ \otimes \mathbb{1}_{B'} \rangle_{\sigma_{A'B'}} \\ &\quad + \sigma^{\nu*} \eta_{\mu\nu} \langle A'_- R_a'^{\mu} A'_- \otimes \mathbb{1}_{B'} \rangle_{\sigma_{A'B'}} \\ &= \sigma_{\mu} \text{Tr}[R_a'^{\mu} A'_+ \sigma_{A'} A'_+] \\ &\quad + \sigma_{\mu}^* \text{Tr}[R_a'^{\mu} A'_- \sigma_{A'} A'_-], \end{aligned} \quad (117)$$

where we used that $\sigma^{\nu*} = \pm\sigma^\nu$ in the same way as $\sigma_\nu^* = \pm\sigma_\nu$ and the property

$$A'_\pm \otimes B'_\mp \sigma_{A'B'} = 0 \quad (118)$$

to get to the second expression.

Let us now introduce diagonal decompositions

$$A'_+ \sigma_{A'} A'_+ = \sum_k p_{k+} |k'_+\rangle \langle k'_+|, \quad (119)$$

$$A'_- \sigma_{A'} A'_- = \sum_l p_{l-} |l'_-\rangle \langle l'_-|, \quad (120)$$

with $\sum_k p_{k+} + \sum_l p_{l-} = 1$, of the two orthogonal operators $A'_\pm \sigma_{A'} A'_\pm$ appearing in the traces. We then obtain

$$\alpha_a = \sum_k p_{k+} R_{a|k+} + \sum_l p_{l-} R_{a|l-}^* \quad (121)$$

where one can verify that

$$R_{a|k+} = \sigma_\mu \langle k'_+ | R_a^{\mu\mu} | k'_+ \rangle, \quad (122)$$

$$R_{a|l-} = \sigma_\mu \langle l'_- | R_a^{\mu\mu} | l'_- \rangle \quad (123)$$

are POVM elements. Since $\{\alpha_a\}$ is by hypothesis extremal, (121) is only possible if

$$R_{a|k+} = R_{a|l-}^* = \alpha_a. \quad (124)$$

Put differently, this means that our uncharacterised POVM $\{R_a\}$ must be such that

$$\text{Tr}_{A'} [R_a (\mathbb{1} \otimes |k'_+\rangle \langle k'_+|)] = \alpha_a, \quad (125)$$

$$\text{Tr}_{A'} [R_a (\mathbb{1} \otimes |l'_-\rangle \langle l'_-|)] = \alpha_a^*. \quad (126)$$

To further constrain $\{R_a\}$, we introduce states

$$|\varphi'_\pm\rangle = \frac{1}{\sqrt{2}} (|j'_+\rangle \pm |k'_+\rangle), \quad (127)$$

with $j < k$, and compute

$$\begin{aligned} & (\mathbb{1} \otimes \langle \varphi'_\pm |) R_a (\mathbb{1} \otimes |\varphi'_\pm\rangle) \\ &= \alpha_a \pm \text{Tr}_{A'} [R_a (\mathbb{1} \otimes X_{jk}^\pm)], \end{aligned} \quad (128)$$

where

$$X_{jk}^\pm = \frac{1}{2} (|j'_+\rangle \langle k'_+| + |k'_+\rangle \langle j'_+|). \quad (129)$$

The left side of (128) is by construction positive semidefinite, which implies

$$\pm \text{Tr}_{A'} [R_a (\mathbb{1} \otimes X_{jk}^\pm)] \leq \alpha_a. \quad (130)$$

Since α_a is of rank one, this is only possible if the partial trace term is itself a multiple of α_a , i.e., if

$$\text{Tr}_{A'} [R_a (\mathbb{1} \otimes X_{jk}^\pm)] = \lambda_a \alpha_a. \quad (131)$$

Since $\{R_a\}$ is a POVM and $\sum_a R_a = \mathbb{1}$,

$$\sum_a \text{Tr}_{A'} [R_a (\mathbb{1} \otimes X_{jk}^\pm)] = \sum_a \lambda_a \alpha_a = 0. \quad (132)$$

Linear independence of the α_a s means that the second part of this equality can only be solved with $\lambda_a = 0$, from which we obtain the constraint

$$\text{Tr}_{A'} [R_a (\mathbb{1} \otimes X_{jk}^\pm)] = 0. \quad (133)$$

Repeating this reasoning starting with states

$$|\phi'_\pm\rangle = \frac{1}{\sqrt{2}} (|l'_-\rangle \pm |m'_-\rangle), \quad (134)$$

and

$$|\phi'_\pm\rangle = \frac{1}{\sqrt{2}} (|j'_+\rangle \pm i|k'_+\rangle), \quad (135)$$

$$|\phi'_\pm\rangle = \frac{1}{\sqrt{2}} (|l'_-\rangle \pm i|m'_-\rangle), \quad (136)$$

we obtain analogous constraints

$$\text{Tr}_{A'} [R_a (\mathbb{1} \otimes X_{lm}^-)] = 0, \quad (137)$$

$$\text{Tr}_{A'} [R_a (\mathbb{1} \otimes Y_{jk}^+)] = 0, \quad (138)$$

$$\text{Tr}_{A'} [R_a (\mathbb{1} \otimes Y_{lm}^-)] = 0 \quad (139)$$

for operators

$$X_{lm}^- = \frac{1}{2} (|l'_-\rangle \langle m'_-| + |m'_-\rangle \langle l'_-|), \quad (140)$$

$$Y_{jk}^+ = \frac{1}{2} (-i|j'_+\rangle \langle k'_+| + i|k'_+\rangle \langle j'_+|), \quad (141)$$

$$Y_{lm}^- = \frac{1}{2} (-i|l'_-\rangle \langle m'_-| + i|m'_-\rangle \langle l'_-|). \quad (142)$$

The sets of operators $\{|k'_+\rangle \langle k'_+|, X_{jk}^+, Y_{jk}^+\}$ and $\{|l'_-\rangle \langle l'_-|, X_{lm}^-, Y_{lm}^-\}$ are bases of the spaces of Hermitian operators acting on the supports of A'_+ and A'_- , so the constraints (125), (126), and (133)–(142) can be written together as the single operator constraint

$$\mathbb{1} \otimes \$'(R_a) = \alpha_a \otimes A'_+ + \alpha_a^* \otimes A'_-, \quad (143)$$

where the superoperator $\$'$ acts on operators on $\mathcal{H}_{A'}$ as

$$\$(X') = A'_+ X' A'_+ + A'_- X' A'_-. \quad (144)$$

The most general operator R_a that satisfies (143) is one of the form

$$R_a = \alpha_a \otimes A'_+ + \alpha_a^* \otimes A'_- + K_a + K_a^\dagger \quad (145)$$

where the off-diagonal operator K_a is identified by whatever

$$K_a = (\mathbb{1} \otimes A'_+) R_a (\mathbb{1} \otimes A'_-) \quad (146)$$

is.

Now we reintroduce that R_a is supposed to be a POVM element. The property $R_a \geq 0$ means, by definition, that we must have $\langle \phi | R_a | \phi \rangle \geq 0$ for any state. Imposing this for certain choice states allows us to further constrain the form of R_a . Dropping in the following the subscript ‘ a ’ for readability, we start with a family of states

$$|\phi\rangle = \frac{1}{\sqrt{2}} (|\alpha_\perp\rangle |+\rangle + e^{i\varphi} |\alpha_\perp^*\rangle |-\rangle), \quad (147)$$

where φ is some phase, $|\alpha_\perp\rangle$ and $|\alpha_\perp^*\rangle$ are the unique (up to a phase) qubit states orthogonal to $|\alpha\rangle$ and $|\alpha^*\rangle$, and $|\pm'\rangle$ are any states in the support of $A'_\pm$, such that $A'_\pm|\pm'\rangle = |\pm'\rangle$ and $A'_\mp|\pm'\rangle = 0$. For the minimising phase $\varphi = \varphi_0$ we get

$$\begin{aligned}\langle\phi|R|\phi\rangle &= \text{Re}[e^{i\varphi_0}\langle\alpha_\perp|\langle+'|K|\alpha_\perp^*\rangle|-\rangle] \\ &= -|\langle\alpha_\perp|\langle+'|K|\alpha_\perp^*\rangle|-\rangle|, \end{aligned} \quad (148)$$

from which we conclude that $\langle\alpha_\perp|\langle+'|K|\alpha_\perp^*\rangle|-\rangle = 0$. Since we also have $\langle\alpha_\perp|\langle-'|K = 0$ and $K|\alpha_\perp^*\rangle|+\rangle$, we can generalise this to the operator equality

$$(\langle\alpha_\perp| \otimes \mathbb{1}_{A'})K(|\alpha_\perp^*\rangle \otimes \mathbb{1}_{A'}) = 0. \quad (149)$$

Next, for

$$|\phi\rangle = \cos(\frac{\theta}{2})|\alpha\rangle|+\rangle + e^{i\varphi}\sin(\frac{\theta}{2})|\alpha_\perp^*\rangle|-\rangle \quad (150)$$

and the minimising $(\theta, \varphi) = (\theta_0, \varphi_0)$,

$$\begin{aligned}\langle\phi|R|\phi\rangle &= \frac{1 + \cos(\theta_0)}{2}\|\alpha\|^2 \\ &\quad + \sin(\theta_0)\text{Re}[e^{i\varphi_0}\langle\alpha|\langle+'|K|\alpha_\perp^*\rangle|-\rangle] \\ &= \frac{1}{2}\|\alpha\|^2 - \frac{1}{2}\sqrt{\|\alpha\|^4 + 4|\langle\alpha|\langle+'|K|\alpha_\perp^*\rangle|-\rangle|^2}, \end{aligned} \quad (151)$$

where $\|\alpha\| = \text{Tr}[\alpha]$. The second expression is negative unless $\langle\alpha|\langle+'|K|\alpha_\perp^*\rangle|-\rangle = 0$; as above, this lets us conclude

$$(\langle\alpha| \otimes \mathbb{1}_{A'})K(|\alpha_\perp^*\rangle \otimes \mathbb{1}_{A'}) = 0. \quad (152)$$

Similarly, starting with a state

$$|\phi\rangle = \cos(\frac{\theta}{2})|\alpha_\perp\rangle|+\rangle + e^{i\varphi}\sin(\frac{\theta}{2})|\alpha^*\rangle|-\rangle \quad (153)$$

we obtain

$$(\langle\alpha_\perp| \otimes \mathbb{1}_{A'})K(|\alpha^*\rangle \otimes \mathbb{1}_{A'}) = 0. \quad (154)$$

These three constraints on K mean that the only remaining possibility is that it is of the form

$$K = |\alpha\rangle\langle\alpha^*| \otimes K', \quad (155)$$

and that R is of the form

$$\begin{aligned}R &= \alpha \otimes A'_+ + \alpha^* \otimes A'_- \\ &\quad + |\alpha\rangle\langle\alpha^*| \otimes K' + |\alpha^*\rangle\langle\alpha| \otimes K'^\dagger \end{aligned} \quad (156)$$

for some operator K' taking states from the support of A'_- to the support of A'_+ , i.e., satisfying

$$A'_+K' = K'A'_- = K' \quad (157)$$

and

$$A'_-K' = K'A'_+ = 0. \quad (158)$$

We can finally ensure that R is positive semidefinite by requiring that all of its eigenvalues are nonnegative. We write

$$K' = \sum_j \kappa_j |j'_+\rangle\langle j'_-|, \quad (159)$$

where the $\kappa_j > 0$ are singular values of K' and $|j'_+\rangle$ and $|j'_-\rangle$ are states in the supports of A'_+ and A'_- . It is then a straightforward exercise to check that

$$|\phi\rangle = |\alpha\rangle|j'_+\rangle \pm |\alpha^*\rangle|j'_-\rangle \quad (160)$$

are eigenstates of R with the form (156) above with eigenvalues

$$\|\alpha\|(1 \pm \kappa_j), \quad (161)$$

from which we extract $\kappa_j \leq 1$. This and the conditions (157) and (158) above together are equivalent to the operator inequalities

$$K'^\dagger K' \leq A'_+ \quad \text{and} \quad K'^\dagger K' \leq A'_-. \quad (162)$$

Conversely, any R that satisfies these conditions is necessarily positive semidefinite since the only other possible eigenvalues of R are $\|\alpha\|$, associated to possible additional eigenstates of A'_+ or A'_- .

Reintroducing the subscript 'a', the only remaining requirement for $\{R_a\}$ with the form above to be a valid POVM is $\sum_a R_a = \mathbb{1}$; this translates directly to the second condition,

$$\sum_a |\alpha_a\rangle\langle\alpha_a^*| \otimes K'_a = 0, \quad (163)$$

stated at the beginning of this section. This is as far as we can go toward identifying $\{R_a\}$ if there are four outcomes. As we pointed out in the main text, $\text{Tr}[|\alpha_a\rangle\langle\alpha_a^*|Y] = 0$ and the operators $|\alpha_a\rangle\langle\alpha_a^*|$ are thus linear combinations of $\mathbb{1}$, X , and Z and cannot be linearly independent if there are four of them. In that case it is always possible to satisfy (163) with nonzero K'_a s.

On the other hand, the operators $|\alpha_a\rangle\langle\alpha_a^*|$ are always linearly independent if there are no more than three of them provided that the α_a are linearly independent. We can see this from the fact that one can always change the basis such that all three α_a are real so that $|\alpha_a\rangle\langle\alpha_a^*| = \alpha_a$. To be precise, let U be a unitary such that, for example, $U\alpha_1U^\dagger$ and $U\alpha_2U^\dagger$ are real. In terms of the Bloch expressions

$$\alpha_a = \frac{1}{2}\|\alpha_a\|(\mathbb{1} + \mathbf{n}_a \cdot \boldsymbol{\sigma}) \quad (164)$$

of α_a , this is any unitary that rotates the normal vector $\mathbf{n}_1 \times \mathbf{n}_2$ onto the y axis, which necessarily rotates $\mathbf{n}_1$ and $\mathbf{n}_2$ onto the x-z plane. Then, necessarily, $U\alpha_3U^\dagger = \mathbb{1} - U\alpha_1U^\dagger - U\alpha_2U^\dagger$ is also real. Applied to the kets, $U|\alpha_a\rangle$ are real up to global phases which are a matter of convention, although we allow for them explicitly anyway. This means that

$$e^{-i\varphi_a}U|\alpha_a\rangle = e^{i\varphi_a}U^*|\alpha_a^*\rangle \quad (165)$$

are real for some phases φ_a . We can use this to relate $|\alpha_a\rangle\langle\alpha_a^*|$ to α_a by

$$e^{-2i\varphi_a}U|\alpha_a\rangle\langle\alpha_a^*|U^T = \alpha_a. \quad (166)$$

This tells us that the $|\alpha_a\rangle\langle\alpha_a^*|$ must necessarily be linearly independent if the α_a are, since if there were nonzero parameters λ'_a such that $\sum_a \lambda'_a |\alpha_a\rangle\langle\alpha_a^*| = 0$ then (166) would imply that $\sum_a \lambda_a \alpha_a = 0$ with the nonzero parameters $\lambda_a = e^{2i\varphi_a} \lambda'_a$.

C. RANDOMNESS GENERATION WITH POVMS

Having identified the extent to which we can self-test extremal qubit POVMs we can return to our original problem of randomness generation. Here we prove that the parties can generate two bits of randomness locally and arbitrarily close to $\log_2(12) \approx 3.5850$ bits of randomness globally by adding POVMs to the Bell test. Before confirming that there are indeed POVMs that yield these amounts of randomness we verify, provided that only one of the parties uses a four-outcome measurement, that the randomness of the outcomes is the same as the randomness yielded by reference qubit POVMs.

A. Reduction to qubits

In an adversarial scenario, Alice and Bob may share an extension ρ_{ABE} of their quantum state prepared by an eavesdropper, Eve. From the Bell test, we can infer that such a state must have the form

$$\rho_{ABE} = \psi_\theta \otimes \sigma_{A'B'E}, \quad (167)$$

where $\sigma_{A'B'E}$ is an extension of the ancillary state $\sigma_{A'B'}$ from the self-test. By performing a measurement $\{\Pi_e\}$ on her part, Eve can effectively prepare one of a number of different ancillary states for Alice and Bob, given by

$$p_e \sigma_{A'B'|e} = \text{Tr}_E[(\mathbb{I}_{A'B'} \otimes \Pi_e) \sigma_{A'B'E}] \quad (168)$$

with $p_e = \text{Tr}[\Pi_e \sigma_E]$, depending on the outcome e she obtains. Alice and Bob's part then is the average

$$\sigma_{A'B'} = \sum_e p_e \sigma_{A'B'|e} \quad (169)$$

of these and the joint probabilities of outcomes of POVMs $\{R_a\}$ and $\{S_b\}$ they perform are averages

$$\langle R_a \otimes S_b \rangle = \sum_e p_e \langle R_a \otimes S_b \rangle_{|e} \quad (170)$$

of the corresponding joint probabilities

$$\langle R_a \otimes S_b \rangle_{|e} = \text{Tr}[(R_a \otimes S_b) \psi_\theta \otimes \sigma_{A'B'|e}] \quad (171)$$

conditioned on e . As we pointed out in the main text, with such a strategy an eavesdropper can improve her chance of correctly guessing Alice and Bob's joint outcome if they both use four-outcome POVMs.

Now let us suppose that Alice uses a four-outcome measurement $\{R_a\}$ and Bob a measurement $\{S_b\}$ with three outcomes or less. By checking their correlations with the other measurements in the Bell test, Alice and Bob infer that the elements of their POVMs are of the forms

$$R_a = \alpha_a \otimes A'_+ + \alpha_a^* \otimes A'_- + |\alpha_a\rangle\langle\alpha_a^*| \otimes K'_a + |\alpha_a^*\rangle\langle\alpha_a| \otimes K_a'^\dagger, \quad (172)$$

and

$$S_b = \beta_b \otimes B'_+ + \beta_b^* \otimes B'_- \quad (173)$$

for some reference extremal qubit POVMs $\{\alpha_a\}$ and $\{\beta_b\}$. The joint probability for an underlying state $\psi_\theta \otimes \sigma_{A'B'|e}$ expands to an expression,

$$\begin{aligned} \langle R_a \otimes S_b \rangle_{|e} = & \langle \alpha_a \otimes \beta_b \rangle_{\psi_\theta} \langle A'_+ \otimes B'_+ \rangle_{\sigma_{A'B'|e}} \\ & + \langle \alpha_a^* \otimes \beta_b \rangle_{\psi_\theta} \langle A'_- \otimes B'_+ \rangle_{\sigma_{A'B'|e}} \\ & + \langle |\alpha_a\rangle\langle\alpha_a^*| \otimes \beta_b \rangle_{\psi_\theta} \langle K'_a \otimes B'_+ \rangle_{\sigma_{A'B'|e}} \\ & + \langle |\alpha_a^*\rangle\langle\alpha_a| \otimes \beta_b \rangle_{\psi_\theta} \langle K_a'^\dagger \otimes B'_+ \rangle_{\sigma_{A'B'|e}} \\ & + \langle \alpha_a \otimes \beta_b^* \rangle_{\psi_\theta} \langle A'_+ \otimes B'_- \rangle_{\sigma_{A'B'|e}} \\ & + \langle \alpha_a^* \otimes \beta_b^* \rangle_{\psi_\theta} \langle A'_- \otimes B'_- \rangle_{\sigma_{A'B'|e}} \\ & + \langle |\alpha_a\rangle\langle\alpha_a^*| \otimes \beta_b^* \rangle_{\psi_\theta} \langle K'_a \otimes B'_- \rangle_{\sigma_{A'B'|e}} \\ & + \langle |\alpha_a^*\rangle\langle\alpha_a| \otimes \beta_b^* \rangle_{\psi_\theta} \langle K_a'^\dagger \otimes B'_- \rangle_{\sigma_{A'B'|e}}, \end{aligned} \quad (174)$$

which is quite lengthy but we can use properties of the state and POVMs derived in the previous two sections to show that most of the terms vanish. We recall first that $\sigma_{A'B'}$ cannot be arbitrary and must be such that $\langle A' \otimes B' \rangle_{\sigma_{A'B'}} = 1$, which requires that

$$\langle A' \otimes B' \rangle_{\sigma_{A'B'|e}} = 1 \quad (175)$$

for all of Eve's possible outcomes e . In terms of the projectors $A'_\pm$ and $B'_\pm$ this means that

$$\langle A'_+ \otimes B'_+ \rangle_{\sigma_{A'B'|e}} + \langle A'_- \otimes B'_- \rangle_{\sigma_{A'B'|e}} = 1 \quad (176)$$

and

$$\langle A'_+ \otimes B'_- \rangle_{\sigma_{A'B'|e}} = \langle A'_- \otimes B'_+ \rangle_{\sigma_{A'B'|e}} = 0. \quad (177)$$

The second pair (177) of constraints already means that the terms involving $\langle \alpha_a^* \otimes \beta_b \rangle_{\psi_\theta}$ and $\langle \alpha_a \otimes \beta_b^* \rangle_{\psi_\theta}$ vanish. (177) furthermore implies the operator constraints

$$(A'_+ \otimes B'_-) \sqrt{\sigma_{A'B'|e}} = 0, \quad (178)$$

$$(A'_- \otimes B'_+) \sqrt{\sigma_{A'B'|e}} = 0, \quad (179)$$

while we recall that the K'_a s satisfy

$$A'_+ K'_a = K'_a A'_- = K'_a. \quad (180)$$

These constraints can be used to show, for example,

$$\begin{aligned} \langle K'_a \otimes B'_+ \rangle_{\sigma_{A'B'|e}} &= \text{Tr}[(K'_a A'_- \otimes B'_+) \sigma_{A'B'|e}] \\ &= 0, \end{aligned} \quad (181)$$

with similar manipulations implying

$$\langle K_a'^\dagger \otimes B'_+ \rangle_{\sigma_{A'B'|e}} = 0, \quad (182)$$

$$\langle K'_a \otimes B'_- \rangle_{\sigma_{A'B'|e}} = 0, \quad (183)$$

$$\langle K_a'^\dagger \otimes B'_- \rangle_{\sigma_{A'B'|e}} = 0. \quad (184)$$

In the joint probabilities we are thus left with only

$$\begin{aligned} \langle R_a \otimes S_b \rangle_{|e} = & \langle \alpha_a \otimes \beta_b \rangle_{\psi_\theta} \langle A'_+ \otimes B'_+ \rangle_{\sigma_{A'B'|e}} \\ & + \langle \alpha_a^* \otimes \beta_b^* \rangle_{\psi_\theta} \langle A'_- \otimes B'_- \rangle_{\sigma_{A'B'|e}}, \end{aligned} \quad (185)$$

which simplify to

$$\langle R_a \otimes S_b \rangle_e = \langle \alpha_a \otimes \beta_b \rangle_{\psi_\theta} \quad (186)$$

because

$$\langle \alpha_a^* \otimes \beta_b^* \rangle_{\psi_\theta} = \langle \alpha_a \otimes \beta_b \rangle_{\psi_\theta^*} = \langle \alpha_a \otimes \beta_b \rangle_{\psi_\theta} \quad (187)$$

due to the state $|\psi_\theta\rangle$ being real.

We conclude that the joint probability of Alice and Bob obtaining outcomes a and b using the POVMs $\{R_a\}$ and $\{S_b\}$, even conditioned on the outcome of a measurement by Eve, is identical to the probability of obtaining the same outcomes using the reference qubit POVMs $\{\alpha_a\}$ and $\{\beta_b\}$ on $|\psi_\theta\rangle$. This considerably simplifies the problem of determining how much randomness we can generate using POVMs provided that a four-outcome POVM is used only on one side.

B. Two bits of local randomness

We suppose here that Alice wishes to extract two bits of randomness in the context of our Bell test. Following the analysis of the previous subsection, we need only confirm that there exist extremal qubit POVMs that produce a uniformly random result from the ideal partially entangled state $|\psi_\theta\rangle$.

In the ideal qubit setting, Alice's marginal state is

$$\psi_{\theta A} = \frac{1}{2}(\mathbb{1} + \cos(\theta)Z) \quad (188)$$

and we are looking to construct a four-outcome POVM $\{\alpha_a\}$ where the POVM elements α_a are of rank one, are linearly independent, and yield the correct probabilities

$$\text{Tr}[\alpha_a \psi_{\theta A}] = \frac{1}{4}. \quad (189)$$

Such POVMs can be constructed by adjusting a tetrahedral POVM, which gives the correct amount of randomness in the special case of the maximally-mixed state. A specific example that works is to take

$$\alpha_1 = \frac{\lambda_1}{2}(\mathbb{1} + Z) \quad (190)$$

and

$$\alpha_a = \frac{\lambda_a}{2} \left(\mathbb{1} + \cos(\gamma)Z + \sin(\gamma)(\cos(\delta_a)X + \sin(\delta_a)Y) \right) \quad (191)$$

with weights

$$\lambda_1 = \frac{1}{2 + 2\cos(\theta)}, \quad (192)$$

$$\lambda_a = \frac{3 + 4\cos(\theta)}{6 + 6\cos(\theta)} \quad \text{for } a = 2, 3, 4 \quad (193)$$

and, in (191), an angle γ such that

$$\cos(\gamma) = -\frac{1}{3 + 4\cos(\theta)} \quad (194)$$

and for example $\delta_2, \delta_3, \delta_4 = 0, 2\pi/3, 4\pi/3$.

C. $\log_2(12)$ bits of global randomness

In this second variant, we suppose Alice and Bob both use POVMs to generate randomness from their joint outcomes. If one of the parties, say, Bob, uses a POVM with three outcomes then it is in principle possible to generate up to $\log_2(12)$ bits of randomness. In the ideal qubit setting, at least one way to get arbitrarily close to attaining this is for Bob to perform a modified version of a Mercedes-Benz POVM in the X-Z plane,

$$\beta_1 = \frac{\lambda_1}{2}(\mathbb{1} + Z), \quad (195)$$

$$\beta_2 = \frac{\lambda_2}{2}(\mathbb{1} - \mu Z + \sqrt{1 - \mu^2}X), \quad (196)$$

$$\beta_3 = \frac{\lambda_3}{2}(\mathbb{1} - \mu Z - \sqrt{1 - \mu^2}X) \quad (197)$$

for parameters related to θ by

$$\lambda_1 = \frac{2}{3 + 3\cos(\theta)}, \quad (198)$$

$$\lambda_2 = \lambda_3 = \frac{2 + 3\cos(\theta)}{3 + 3\cos(\theta)}, \quad (199)$$

$$\mu = \frac{1}{2 + 3\cos(\theta)}, \quad (200)$$

while Alice performs a measurement

$$\alpha_1 = \frac{1}{4}(\mathbb{1} + \sqrt{1 - \varepsilon^2}Y + \varepsilon Z), \quad (201)$$

$$\alpha_2 = \frac{1}{4}(\mathbb{1} + \sqrt{1 - \varepsilon^2}Y - \varepsilon Z), \quad (202)$$

$$\alpha_3 = \frac{1}{4}(\mathbb{1} - \sqrt{1 - \varepsilon^2}Y + \varepsilon X), \quad (203)$$

$$\alpha_4 = \frac{1}{4}(\mathbb{1} - \sqrt{1 - \varepsilon^2}Y - \varepsilon X) \quad (204)$$

whose elements deviate only a small amount from the Y axis. Importantly, Alice's measurement is extremal for any nonzero ε . For small ε , these measurements give joint probabilities

$$\langle \alpha_a \otimes \beta_b \rangle_{\psi_\theta} \approx \frac{1}{12}, \quad (205)$$

with equality in the limit $\varepsilon \rightarrow 0$.