

Reply to comment on “Interaction of a BEC with a gravitational wave”

Ralf Schützhold

*Helmholtz-Zentrum Dresden-Rossendorf, Bautzner Landstraße 400, 01328 Dresden, Germany, and
Institut für Theoretische Physik, Technische Universität Dresden, 01062 Dresden, Germany.*

(Dated: February 1, 2022)

This reply contains a brief response to the comment by R. Howl, D. Rätzel, and I. Fuentes.

In case of conflicting points of view, a proper scientific debate can be very useful since it can help to address an issue from several angles and to identify the core of a problem. Thus, let me respond to the comment [1] on my publication [2]. First, I would like to stress that my work in [2] is not meant to be a suggestion of a special (e.g., alternative to [3]) detection scheme, but rather a general investigation of the interaction between gravitational waves and Bose-Einstein condensates (BEC) together with a general order-of-magnitude estimate of the interaction effects (as already stated in [2]). In the comment [1], I did not find any argument which invalidates the main results of [2]. Instead, I found several points of agreement – so it might be best to start with those.

As is well-known, for non-classical states (such as squeezed states or NOON states), the sensitivity can scale with the number of particles (e.g., photons, atoms, or phonons) instead of its square root. There also seems to be agreement that inhomogeneities in the condensate (e.g., weak inhomogeneities such as sound waves or strong inhomogeneities such as vortices) can enhance its interaction with gravitational waves. Furthermore, we seem to agree that the number of phonons (as linear excitations of the dilute BEC) should not exceed the number of atoms (even though the squeezing parameter $r = 10$ proposed in [3] appears to suggest otherwise because it would correspond to a mean phonon number of $\sinh^2 r > 10^8$).

Now, if the phonon number is well below the atom number, we may estimate the scaling of any perturbation Hamiltonian \hat{H}_{int} describing the interaction with a gravitational wave (including that stemming from the trap potential) which is bi-linear in the field operators $\hat{\Psi}$ by inserting the mean-field approximation $\hat{\Psi} \approx \psi_{\text{cond}} + \hat{\chi}$, where ψ_{cond} is the condensate wave-function and $\hat{\chi}$ describes the Bogoliubov-de Gennes (phonon) modes. We find the following hierarchy [2] of contributions: To leading order, the action of the perturbation Hamiltonian \hat{H}_{int} scales with the atom number. To sub-leading order, it scales with the geometric mean of the atom and phonon number (or, if no phonons are present, with the square root of the atom number). In this sequence, the effect considered in [3] would be next-to-sub-leading order as it scales with the number of phonons. Again, I found no argument in [1] which invalidates this picture.

It is correct that the response of the trap potential to the gravitational wave is (apart from general estimates as sketched above) not considered explicitly in [2].

This would require a microscopic analysis of the trapping mechanism and how it changes due to the gravitational wave. However, this is also not done in [3], which is based on the simplified assumption of a rigid and uniform box trap potential. Also, the direct creation of phonons by the trap (instead of the amplification of already existing phonons, see also [4]) is not considered in [3].

Since the number of atoms or phonons in present-day BEC made of ultra-cold atomic vapor are not sufficient to reach the sensitivity required for gravitational wave detection, further large numbers are necessary. This could be large ratios of length or time scales (or a large number of detectors). The associated challenges can be exemplarily illustrated by considering Eq. (1) in [1]

$$\Delta\epsilon \propto \frac{1}{N\sqrt{\omega_m\omega_n\tau t N_d}}, \quad (1)$$

where $\Delta\epsilon$ measures the sensitivity, N is the phonon number (denoted by n in [2]), ω_m and ω_n are phonon frequencies, t is the run-time, τ the integration time, and N_d denotes the number of detectors. Unfortunately, the authors of [1] did not provide explicit numbers, so let us insert example values suggested in the proposal [3]. Using the atom number of 10^6 as assumed in [3] as an upper bound for the phonon number N (see above), the term within the square root in (1) should exceed 10^{30} in order to reach a sensitivity of 10^{-21} , cf. [5]. As suggested in [3], let us insert comparably large phonon frequencies in the $2\pi \times 5$ kHz regime and a rather long run-time t of 1000 seconds (with all the problems, see also [2, 4]). Still, we obtain an accumulated integration time $N_d\tau$ exceeding the age of our Universe, which seems to require an astronomical number N_d of detectors/condensates.

In summary, even though it would be extremely nice to detect gravitational waves with small present-day Bose-Einstein condensates made of ultra-cold atomic vapor, it appears to be a tremendously challenging task.

-
- [1] R. Howl, D. Rätzel, and I. Fuentes, [arXiv:1811.10306v2](#).
 - [2] R. Schützhold, [arXiv:1807.07046v1](#), see also [arXiv:1807.07046v2](#); Phys. Rev. D **98**, 105019 (2018).
 - [3] C. Sabín, D.E. Bruschi, M. Ahmadi and I. Fuentes, New J. Phys. **16**, 085003 (2014).
 - [4] D. Rätzel, R. Howl, J. Lindkvist and I. Fuentes, New J. Phys. **20**, 073044 (2018).
 - [5] B. P. Abbott *et al*, Phys. Rev. Lett. **116**, 061102 (2016).