# **Two-body problems with magnetic interactions**

## Hesham Mansour<sup>+</sup> and Ahmed Gamal<sup>\*</sup>

Physics Department, faculty of science, Cairo University, Giza, Egypt
Correspondence: Ahmed Gamal, Department of Physics, Cairo University, Giza, Egypt.
E-mail: <u>mansourhesham@gmail.com</u>
\* corresponder author

+ FInstP

## Abstract

In the present work, we present different two-body potentials which have oscillatory shapes with magnetic interactions. The eigenvalues and eigenfunctions are obtained for one of those problems using Nikiforov-Uvarov method.


#### **Keywords**

The non-relativistic quantum theory; The relativistic two-body equation; The Nikiforov-Uvarov method.

### (1)-Introduction

In the present work, we shall be particularly interested in a potential well as the one shown in figure (1).

Magnetic forces between spin 1/2 particles lead to the effective radial potentials of this type [1-3], with one or more deep narrow wells. Magnetic interactions are studied for various problems ranging from macroscopic to microscopic scales [4-10]. The



**Figure (1),** show the relation between the potential V(r) and the distance r at different distance

purpose of our work is to use analytic techniques to study the eigenvalues and eigenfunctions for such cases. Effective potentials between two-fermions taking into

account form factors and full relativistic spinor kinematics have similar shapes. The motivation for doing that is as we shall show later on the existence of such potentials in several physical applications. In such cases, one would need to understand better the existence and to characterize the quantum numbers of such possible solutions using these potentials.

Let us consider, for example, a relativistic charged spinless particle m in a field of a fixed (quantum ) magnetic moment  $\vec{\mu}$  [11], or alterentivly , a charged spin 1/2 particle of mass m and magntic moment  $\vec{\mu}$ , in the field of a fixed charge [12,13]. In both cases the radial equation has the form

$$\left[-\frac{d^2}{dr^2} + V - \lambda^2\right]U(r) = 0\tag{1}$$

Where the effective potential is given, apart from the Coulomb potential  $\frac{\alpha}{r}$ , by

$$V(r) = \frac{A}{r^2} + \frac{B}{r^3} + \frac{C}{r^4}$$
(2)

Clearly, if one solves the same problem with a Dirac equation and give also an anomalous magnetic moment  $\vec{a}$  to the particle, then additional terms are added to equation (2). Further models may also treat the magnetic moment of both particles. The potential in equation (2) is treated in atomic phenomena as well as in the quark models as a perturbation. This is justified only if the energies are of the order of the Coulomb energies. At very short distances, the form of the potentials is quite uncertain, and these potentials must be modified by form factors. Form factors must then be calculated in a non-perturbative and self-cosistent method from the wave functions which are localized around each well. In the next section, we will sketch some of the physical problems in which case such a form of this potential appears.

In section (3) we will give a short description of the relativistic two -body equation which is used to describe two-body systems, and finally, in section (4), we will show a method for solving equation (1) over the whole range of the independent variable.

#### (2)-Dynamical models

#### (2.1)-Magnetic interactions in nonrelativistic quantum theory

Consider two charged particles  $e_1$  and  $e_2$  with magnic moments  $\overrightarrow{\mu_1}$  and  $\overrightarrow{\mu_2}$  respectively the Hamiltonian in this case is

$$H = \frac{1}{2m_1} \left[ \overrightarrow{P_1} - e_1 \frac{\overrightarrow{\mu_2} \times (\overrightarrow{r_1} - \overrightarrow{r_2})}{|\overrightarrow{r_1} - \overrightarrow{r_2}|^3} \right]^2 + \frac{1}{2m_2} \left[ \overrightarrow{P_2} - e_2 \frac{\overrightarrow{\mu_1} \times (\overrightarrow{r_2} - \overrightarrow{r_1})}{|\overrightarrow{r_2} - \overrightarrow{r_1}|^3} \right]^2 + \frac{e_1 e_2}{|\overrightarrow{r_2} - \overrightarrow{r_1}|} + \overrightarrow{\mu_1}$$
$$\cdot \overrightarrow{\mu_2} \frac{(3\overrightarrow{\sigma_1} \cdot \overrightarrow{r_1}) (3\overrightarrow{\sigma_2} \cdot \overrightarrow{r_2}) - \overrightarrow{\sigma_1} \cdot \overrightarrow{\sigma_2}}{r^3} + \frac{8\pi}{3} (\overrightarrow{\mu_1} \cdot \overrightarrow{\mu_2}) (\overrightarrow{\sigma_1} \cdot \overrightarrow{\sigma_2}) \delta(\overrightarrow{r_1} - \overrightarrow{r_2})$$
(3)

Or In the center of mass frame

$$H = \frac{1}{2\mu}P^{2} - \vec{P} \cdot \frac{\vec{a} \times \vec{r}}{r^{3}} + \frac{A}{r^{4}} + \frac{e_{1} \cdot e_{2}}{r} - \mu_{1} \cdot \mu_{2} S_{12}(\vec{r})$$
(4)  
$$\vec{a} = \frac{e_{1}}{m_{1}}\overline{\mu_{2}} + \frac{e_{2}}{m_{2}}\overline{\mu_{1}} , \qquad \vec{P} = \frac{m_{2}\overline{P_{1}} - m_{1}\overline{P_{2}}}{m_{1} + m_{2}} ,$$

Where

$$\vec{a} = \frac{e_1}{m_1} \overrightarrow{\mu_2} + \frac{e_2}{m_2} \overrightarrow{\mu_1} , \qquad \vec{P} = \frac{m_2 \overrightarrow{P_1} - m_1 \overrightarrow{P_2}}{m_1 + m_2}$$
$$A = \frac{e_1^2}{m_1} \mu_2^2 + \frac{e_2^2}{m_2} \mu_1^2 , \qquad \frac{1}{\mu} = \frac{1}{m_1} + \frac{1}{m_2}$$

And  $S_{12}(\vec{r})$  is the so-called tensor and dipole-dipole interaction potential

In some special cases, the problem becomes easier to solve e.g.

a)  $m_2 \gg m_1$  ,  $\mu_1 = 0$  or b)  $m_2 = m_1$  ( $e \cdot g \ pp$  ,  $p\bar{p}$  ,  $e^-e^+$  ,  $e^-e^-$  ,... )

#### (2.2)-The Dirac particle with an anomalous magnetic moment in a dipole field.

Consider now a relativistic Dirac particle with an anomalous magnetic moment a, charge e in the field of a fixed quantum dipole moment  $\vec{\mu}_2$  and charge  $e_2$  the equation (in natural units)[1,2] is

$$[\gamma^{\mu}(p_{\mu} - e_{1}A_{\mu}) - m]\Psi = -a\frac{e_{1}}{4\pi}\gamma^{\mu}\gamma^{\nu}F_{\mu\nu}\Psi$$

$$A_{\mu} = \left(\frac{e_{2}}{r}, \mu_{2}\frac{\vec{\sigma}\times\vec{r}}{r^{3}}\right)$$
(5)

With

$$\left[\vec{\alpha}.\left(\vec{p}-e_{1}\vec{A}\right)-\left[E-\epsilon\frac{\alpha}{r}\right]+\beta m\right]\Psi=-a\epsilon\frac{\alpha}{2m}\frac{1}{r^{2}}i\beta\alpha_{r}\Psi\tag{6}$$


Where  $\epsilon = sign(e_1 \cdot e_2)$  and  $\alpha_r = \frac{\vec{\alpha} \cdot \vec{r}}{r}$ 

When  $\vec{A} = 0$ , the above equation can be transformed to two- uncoupled second order Sturm-Liouville eigen value equations[1,15]; viz;

$$\left[\frac{d^2}{dr^2} + k^2 - V_i(r)\right] \Psi_i = 0$$
(7)

Where  $k^2 = E^2 - m^2$ 

And the effective energy and angular momentum potential  $V_i(r)$  will have the same shape as shown in figure (2)



**Figure (2),** the trend of the potential V(r) with distance r at maxima and minima for different regions

#### (2.3)-The 4-component neutrino with a magnetic moment

An interesting limit of the equation in the previous section is when

$$e 
ightarrow 0$$
 ,  $m 
ightarrow 0$  such that  $rac{e}{m} = \mu = ext{magntic moment}$ 

One gets an equation which is exactly solvable[14,15]; viz,

$$\left[\frac{d^2}{dy^2} + \lambda^2 - \left[\frac{k(k+1)}{y^2} + \epsilon \frac{2(k+1)}{y^3} + \frac{1}{y^4}\right]\right]\Psi = 0$$
(8)

Where  $k = \pm (j + \frac{1}{2})$ 

Also for a vv system, we have the Hamiltionian which includes the recoil and spin-spin terms

$$H = \alpha^{(1)} \cdot \overrightarrow{p_1} + \alpha^{(2)} \cdot \overrightarrow{p_2} + \frac{3(\overrightarrow{\sigma_1} \cdot \overrightarrow{r})(\overrightarrow{\sigma_2} \cdot \overrightarrow{r}) - \overrightarrow{\sigma_1} \cdot \overrightarrow{\sigma_2}}{r^3} + \frac{8\pi}{3} \overrightarrow{\sigma_1} \cdot \overrightarrow{\sigma_2} \,\delta(\overrightarrow{r}) \tag{9}$$

This Hamiltonian is interesting to study .To this one may add also the Hamiltonian for a dyonium i.e two objects having each a magnetic and an electric charge

$$H = \boldsymbol{\alpha}^{(1)} \cdot \left( \overrightarrow{p_1} - e_1 \overrightarrow{A_1} \right) + \boldsymbol{\alpha}^{(2)} \cdot \left( \overrightarrow{p_2} - e_2 \overrightarrow{A_2} \right)$$
(10)  
$$\overrightarrow{A_1} = \mu_2 \frac{\boldsymbol{\sigma}^{(2)} \times \overrightarrow{r}}{r^3} \quad . \quad \overrightarrow{A_2} = \mu_1 \frac{\boldsymbol{\sigma}^{(1)} \times \overrightarrow{r}}{r^3}$$

Magnetic monopoles have not yet been seen [17,18]. They have influenced fields like high energy physics e.g. quark confinement, superstring theory and supersymmetric quantum field theories. This has guided condensed matter physicists to discover anomalous states and excitations in systems such as Bose-Einstein condensates [19] and spinices [20]. The above-mentioned models may be treated within the framework of relativistic two-body problem in the so-called one-time formulation[21]. Also one must add the various potentials obtained from the Bethe-Salpeter type of equations which should be treated in a nonperturbative manner. In the next section, we will discuss the two-body equations.

#### (3)-The relativistic two-body equations

In a previous works, we derived a new set of the relativistic radial equations for two spin 1/2 particles [22,23]. The 16 amplitudes are re-expressed in terms of four scalars and four vectors which satisfy coupled differential equations. These equations are reduced further to sets of coupled differential equations according to parity and total angular momentum. The equations are solved in the positronium case and the results obtained are similar to that of Schrödinger equation for the hydrogen-like atom.

In quantum field theory, the work on the two-body problem is based on two approaches. The first approach is a covariant formulation ( the so-called Bethe-Salpeter equation ) for describing the relativistic two-body systems [24], but owing to the complex structure of this equation no general solution has been found as yet [25-31]. Other approaches have been used by several authors (the one-time formulation) in order to give a physical interpretation for the two-body amplitudes. The reason for using our equations is that they are exactly solvable. A simplification and reduction of the system of coupled equations for the 16 amplitudes were achieved by the explicit imposition of Lorentz and charge conjugation invariance and parity conservation.

The number of the coupled equations is not more than eight and can be fewer depending on the interaction used. For instantaneous potential, the equations reduce to three- sets of tractable coupled differential equations. These equations are easier and analytically solvable [23].

Physically the use of the concept of potential for the relativistic problems at short distances may be questionable, however, one must remember that we are dealing with interactions of highly localized states. In such case for the bound state, the potential concept is proven to be superior to the use of the perturbation theory [32]. The interset in such class of interactions stems from the existence of high mass resonances e.g. the superpositrium in QED.

The effective static potential between two fermions is known and leads to the Brietequation which is valid for weak binding. However, because of the strong binding nature of the potentials one may need some extrapolation of such potentials.

To do that one may introduce form factors at the vertices to take into account the selfenergy and radiative corrections in an already renormalized form. Such a program has been discussed previously[19], where they incorporated form factors for both particles using a relativistic two-body one-time formalism. The effective potential is given by equation (2.12) in that reference. The effect of form factors has been derived in previous works [22,23].


#### (4)-Methods of solution

In general one may recourse to the numerical solution to any of the above-mentioned problems but, here we will sketch the method for the simple case i.e equations (1) and (2) as shown in figure (3)

For such a potential, we have three zeros at

$$V(r) = 0 \tag{11}$$

and the positions of minima and maxima can be obtained from the equation



**Figure (3),** the potential V(r) with the distance r showing the positions at V(r)=0 and the maxima and minima according equation (2)

$$\dot{V}(r) = 0 \tag{12}$$

Equations (11) and (12) are a cubic equations in the general form, viz;

$$\dot{a}r^3 + \dot{b}r^2 + \dot{c}r + \dot{d} = 0 \tag{13}$$

By replacing r by r + y, this can be transformed into

$$r^3 + Fr + H \tag{14}$$

For which the three roots are

$$-a - b \quad . \quad -aw^2 - bw \quad . \quad -aw - bw^2$$
 (15)

Where

$$w^3 = 1 \quad . \ 1 + w + w^2 = 0 \tag{16}$$

$$2a^3 = H + \sqrt{H^2 + \frac{4}{27}F^3} \tag{17}$$

$$2b^3 = H - \sqrt{H^2 + \frac{4}{27}F^3} \tag{18}$$

And from here one can proceed to find the solutions around the extremum points and the WKB like solutions as in a previous work[33,34]. For the solutions around the zeros one can use a generalized procedure which has been adopted by Shorupski[35].

In the present work, we will use the Nikiforov -Uvarov method as described in the next section.

#### (4.1)-The Nikiforov -Uvarov method[36-38].

The Nikiforov-Uvarov (NU) method is based on solving the hypergeometric-type secondorder differential equation.

$$\ddot{\Psi}(s) + \frac{\tilde{\tau}(s)}{\sigma(s)}\dot{\Psi}(s) + \frac{\tilde{\sigma}(s)}{\sigma^2(s)}\Psi(s) = 0.$$
(19)

Where  $\sigma(s)$  and  $\tilde{\sigma}(s)$  are second-degree polynomials,  $\tilde{\tau}(s)$  is a first-degree polynomial and  $\psi(s)$  is a function of the hypergeometric-type.

By taking

$$\Psi(s) = \varphi(s)\Upsilon(s) \tag{20}$$

(23)

And substituting in equation (19), we get

$$\ddot{Y}(s) + \left[2\frac{\dot{\varphi}(s)}{\varphi(s)} + \frac{\tilde{\tau}(s)}{\sigma(s)}\right]Y(s) + \left[\frac{\ddot{\varphi}(s)}{\varphi(s)} + \frac{\dot{\varphi}(s)}{\varphi(s)}\frac{\tilde{\tau}(s)}{\sigma(s)} + \frac{\tilde{\sigma}(s)}{\sigma^2(s)}\right]Y(s) = 0.$$
(21)

By taking 
$$2\frac{\dot{\varphi}(s)}{\varphi(s)} + \frac{\tilde{\tau}(s)}{\sigma(s)} = \frac{\tau(s)}{\sigma(s)} \quad and \quad \frac{\dot{\varphi}(s)}{\varphi(s)} = \frac{\pi(s)}{\sigma(s)} \quad .$$
 (22)

We get 
$$\tau(s) = \tilde{\tau}(s) + 2\pi(s)$$
.

Where both,  $\pi(s)$  and  $\tau(s)$  are polynomials of degree at most one.

Also one can take 
$$\frac{\ddot{\varphi}(s)}{\varphi(s)} + \frac{\dot{\varphi}(s)}{\varphi(s)}\frac{\tilde{\tau}(s)}{\sigma(s)} + \frac{\tilde{\sigma}(s)}{\sigma^2(s)} = \frac{\bar{\sigma}(s)}{\sigma^2(s)}.$$
 (24)

$$\frac{\ddot{\varphi}(s)}{\varphi(s)} = \left[\frac{\dot{\varphi}(s)}{\varphi(s)}\right]^2 + \left[\frac{\dot{\varphi}(s)}{\varphi(s)}\right]^2 = \left[\frac{\pi(s)}{\sigma(s)}\right]^2 + \left[\frac{\pi(s)}{\sigma(s)}\right]^2.$$
(25)

And

Where

$$\bar{\sigma}(s) = \tilde{\sigma}(s) + \pi^2(s) + \pi(s)[\tilde{\tau}(s) - \dot{\sigma}(s)] + \dot{\pi}(s)\sigma(s).$$
<sup>(26)</sup>

So equation (21) becomes

$$\ddot{Y}(s) + \frac{\tau(s)}{\sigma(s)}Y(s) + \frac{\bar{\sigma}(s)}{\sigma^2(s)}Y(s) = 0.$$
(27)

An algebraic transformation from equation (19) to equation (27) is systematic. Hence one can divide  $\bar{\sigma}(s)$  by  $\sigma(s)$  to obtain a constant  $\lambda$ . i.e.

$$\bar{\sigma}(s) = \lambda \sigma(s) . \tag{28}$$

Equation (27) can be reduced to a hypergeometric equation in the form

$$\sigma(s)\ddot{Y}(s) + \tau(s)Y(s) + \lambda Y(s) = 0.$$
<sup>(29)</sup>

If one substitutes from equation (28) in equation (26) and solve the quadratic equation for  $\pi(s)$ , we get

$$\pi^2(s) + \pi(s)[\tilde{\tau}(s) - \dot{\sigma}(s)] + \tilde{\sigma}(s) - k\sigma(s) = 0.$$
(30)

Where

$$k = \lambda - \dot{\pi}(s) \,. \tag{31}$$

$$\pi(s) = \frac{\dot{\sigma}(s) - \tilde{\tau}(s)}{2} \pm \sqrt{\left(\frac{\dot{\sigma}(s) - \tilde{\tau}(s)}{2}\right)^2 - \tilde{\sigma}(s) + k\sigma(s)}$$
(32)

The possible solutions for  $\pi(s)$  depend on the parameter k according to the plus and minus signs of  $\pi(s)$ . Since  $\pi(s)$  is a polynomial of degree at most one, so the expression under the square root must be the square of a polynomial. In this case, an equation of the quadratic form is available for the constant k. To determine the parameter k one must set the discriminant of this quadratic expression to be equal to zero. After determining the values of k one can find the values of  $\pi(s)$ ,  $\lambda$  and  $\tau(s)$ .

Applying the same systematic way for equation (28), we get

$$\lambda_n = -n\dot{\tau}(s) - \frac{n(n-1)}{2}\ddot{\sigma}(s) \tag{33}$$

Where n is the principle quantum number.

By comparing equations (31) and (33), we get the energy eigenvalues equation.

To get the eigenfunction  $\Psi(s)$ , one must know  $\varphi(s)$  and  $\Upsilon(s)$ .

We can obtian  $\varphi(s)$  from equation (22).

Now, we will obtian  $\Upsilon(s)$  by

$$\Upsilon(s) = \frac{B_n}{\rho(s)} \frac{d^n}{ds^n} [\sigma^n(s) \ \rho(s)] \tag{34}$$

We can use the Rodrigues' formula of the associated Laguerre polynomials, where

$$L_n^z(r) \equiv \Upsilon(s)$$
 and  $B_n = \frac{1}{n!}$  (35)

 $B_n$  is a normalization constant and the weight function  $\rho(s)$  must satisfy the condition below

$$\frac{\dot{\rho}(s)}{\rho(s)} = \frac{\tau(s) - \dot{\sigma}(s)}{\sigma(s)} \tag{36}$$

Now, we will apply this method to solve equation (1) in our case.

First, we will give the solution for a general inverse polynomial potential form using Nikiforov-Uvarov (NU) method.

let U(r) = r R(r) and subtituting in equation (1)[39 - 47], we get

$$\frac{d^2R}{dr^2} + \frac{2}{r} * \frac{dR}{dr} + [\lambda^2 - V(r)]R = 0$$
(37)

The generalized potential

$$V(r) = \sum_{h=0}^{h} A_{-h} r^{-h} = A_0 + A_{-1} r^{-1} + A_{-2} r^{-2} + A_{-3} r^{-3} + A_{-4} r^{-4} + \cdots$$
(38)

Substituting into equation (37), one gets

$$\frac{d^2 R}{dr^2} + \frac{2}{r} * \frac{dR}{dr} + \left[\lambda^2 - \sum_{h=0}^h A_{-h} r^{-h}\right] R = 0$$
(39)

$$\frac{d^2R}{dr^2} + \frac{2}{r} * \frac{dR}{dr} + \frac{V_1(r)}{r^2}R = 0$$
(40)

Where  $V_1(r) = -[B_0 r^2 + A_{-1}r + A_{-2} + A_{-3}r^{-1} + A_{-4}r^{-2} + A_{-5}r^{-3} + \cdots]$  (41)

And  $B_0 = -\lambda^2 + A_0$ 

By substituting equation (41) in equation (40), we get

$$\frac{d^2R}{dr^2} + \frac{2}{r} * \frac{dR}{dr} + \frac{1}{r^2} \left[ -B_0 r^2 - A_{-1} r - A_{-2} - \sum_{h=1}^h A_{-h-2} \left(\frac{1}{r}\right)^h \right] R = 0$$
(42)

Let  $r = y + r_0$  (where  $r_0$  is the smallest distance from the origin to the particle) and take Maclurin expansion for the summation terms to the second order, one gets

$$\sum_{h=1}^{h} A_{-h-2} \left(\frac{1}{r}\right)^{h} = \sum_{h=1}^{h} \frac{A_{-h-2}}{(y+r_{0})^{h}} = \sum_{h=1}^{h} \frac{A_{-h-2}}{r_{0}^{h}} \left[1 + \frac{y}{r_{0}}\right]^{-h}$$
(43)

$$\sum_{h=1}^{h} \frac{A_{-h-2}}{r_0^h} \left[ 1 + \frac{y}{r_0} \right]^{-h} \approx \sum_{h=1}^{h} \frac{A_{-h-2}}{r_0^h} \left[ 1 - \frac{h}{r_0} y + \frac{h(h+1)}{2r_0^2} y^2 \right]$$
(44)

$$\sum_{h=1}^{h} A_{-h-2} \left(\frac{1}{r}\right)^{h} \approx \sum_{h=1}^{h} \frac{A_{-h-2}}{r_{0}^{h}} \left[ \left[\frac{(h+2)(h+1)}{2}\right] - \left[\frac{h(h+2)}{r_{0}}\right]r + \frac{h(h+1)}{2r_{0}^{2}}r^{2} \right]$$
(45)

By substituting equation (45) into equation (42), we get

$$\frac{d^{2}R}{dr^{2}} + \frac{2}{r} * \frac{dR}{dr} + \frac{1}{r^{2}} \left[ -B_{0}r^{2} - A_{-1}r - A_{-2} - \sum_{h=1}^{h} \frac{A_{-h-2}}{r_{0}^{h}} \left[ \left[ \frac{(h+2)(h+1)}{2} \right] - \left[ \frac{h(h+2)}{r_{0}} \right]r + \frac{h(h+1)}{2r_{0}^{2}}r^{2} \right] \right] R = 0$$
(46)

Rearranging equation (46), we get

$$\frac{d^{2}R}{dr^{2}} + \frac{2}{r} * \frac{dR}{dr} + \frac{1}{r^{2}} \left[ -\left(A_{-2} + \frac{1}{2} \sum_{h=1}^{h} \frac{(h+2)(h+1)}{r_{0}^{h}} A_{-h-2}\right) + \left(\sum_{h=1}^{h} \frac{h(h+2)A_{-h-2}}{r_{0}^{h+1}} - A_{-1}\right) r - \left(B_{0} + \frac{1}{2} \sum_{h=1}^{h} \left[\frac{h(h+1)A_{-h-2}}{r_{0}^{h+2}}\right]\right) r^{2} \right] R = 0$$

$$(47)$$

let 
$$A_{-2} + \frac{1}{2} \sum_{h=1}^{h} \frac{(h+2)(h+1) A_{-h-2}}{r_0^h} = q$$
,  
 $\sum_{h=1}^{h} \frac{h(h+2) A_{-h-2}}{r_0^{h+1}} - A_{-1} = w$ ,  $B_0 + \frac{1}{2} \sum_{h=1}^{h} \left[ \frac{h(h+1) A_{-h-2}}{r_0^{h+2}} \right] = z$ 

we obtain

$$\frac{d^2R}{dr^2} + \frac{2}{r} * \frac{dR}{dr} + \frac{1}{r^2} [-q + wr - zr^2]R = 0$$
(48)

Now, we follow Nikiforov -Uvarov method

$$ilde{ au}=2$$
 ,  $\sigma=r$  ,  $ilde{\sigma}=-q+rw-r^2z$ 

Equation (32) is used to obtain  $\pi(r)$ 

$$\pi(r) = -\frac{1}{2} \pm \sqrt{r^2 z + (k - w)r + q + \frac{1}{4}}$$
(49)

Now, we calculate the value of the parameter k

$$\Delta = b^{2} - 4ac = 0 \qquad \to \qquad (k - w)^{2} - 4z\left(q + \frac{1}{4}\right) = 0$$
$$(k - w)^{2} = 4z\left(q + \frac{1}{4}\right) \qquad \to \qquad k = \pm \sqrt{4z\left(q + \frac{1}{4}\right)} + w \qquad (50)$$

In equation (50), we will deal with two cases of k.

By substituting the values of k in equation (49) and take the negative value of  $\pi(r)$ , one gets

$$\pi(r) = -\frac{1}{2} - \sqrt{\left(\sqrt{z}r \pm \sqrt{q + \frac{1}{4}}\right)^2} = -\frac{1}{2} - \sqrt{z}r \mp \sqrt{q + \frac{1}{4}}$$
(51)

Using equation (23), one obtains the value of  $\tau(r)$ 

$$\tau(r) = 2 - 1 - 2\sqrt{z}r \mp 2\sqrt{q + \frac{1}{4}} = 1 - 2\sqrt{z}r \mp 2\sqrt{q + \frac{1}{4}}$$
(52)

Also using equation (31) to get  $\hat{\lambda}$ 

$$\lambda = k + \dot{\pi}(r) = \pm \sqrt{4z\left(q + \frac{1}{4}\right)} + w - \sqrt{z} = w \pm \sqrt{4z\left(q + \frac{1}{4}\right)} - \sqrt{z}$$
(53)

 $\lambda_n$  can be obtained from equation (33)

$$\lambda_n = 2n\sqrt{z} \tag{54}$$

Comparing equations (53) and (54), one gets the energy eigenvalue equation, hence

$$z = \frac{w^2}{\left[1 + 2n \pm 2\sqrt{\left(q + \frac{1}{4}\right)}\right]^2}$$
(55)

By substituting the values of z, w and q, we get the energy eigenvalue equation for a general inverse polynomial potential

$$\lambda^{2} = A_{0} + \frac{1}{2} \sum_{h=1}^{h} \left[ \frac{h(h+1)A_{-h-2}}{r_{0}^{h+2}} \right] - \frac{\left(\sum_{h=1}^{h} \frac{h(h+2)A_{-h-2}}{r_{0}^{h+1}} - A_{-1}\right)^{2}}{\left[ (2n+1) \pm 2\sqrt{\left(A_{-2} + \frac{1}{2}\sum_{h=1}^{h} \frac{(h+2)(h+1)A_{-h-2}}{r_{0}^{h}} + \frac{1}{4}\right)} \right]^{2}}$$

One can also calculate the eigenfunctions for the general inverse polynomial potential from equation (20) using the negative value of k. So first, we calculate  $\varphi(s)$  from equation (22)

$$\int \frac{d\varphi(\mathbf{r})}{\varphi(\mathbf{r})} = \int \left[ -\sqrt{z} + \frac{-\frac{1}{2} + \sqrt{q + \frac{1}{4}}}{r} \right] dr$$
(56)

$$\varphi(\mathbf{r}) = r^{-\frac{1}{2} + \sqrt{q + \frac{1}{4}}} e^{-\sqrt{z}r}$$
(57)

Second, we calculate  $\Upsilon(r)$  from equation (36) and equation (34)

$$\int \frac{d\rho(\mathbf{r})}{\rho(\mathbf{r})} = \int \left[ -2\sqrt{z} + \frac{2\sqrt{q+\frac{1}{4}}}{r} \right] d\mathbf{r}$$
(58)

$$\rho(\mathbf{r}) = r^{2\sqrt{q+\frac{1}{4}}} e^{-2\sqrt{z}r}$$
(59)

$$Y(\mathbf{r}) = Y_n(r) = j_n r^{-2\sqrt{q+\frac{1}{4}}} e^{2\sqrt{zr}} \frac{d^n}{ds^n} \left[ r^{n+2\sqrt{q+\frac{1}{4}}} e^{-2\sqrt{zr}} \right]$$
(60)

where  $j_n$  is a normalization constant of Y(r) which can be expressed as an associated Laguerre polynomial, where

$$L_n^{2\sqrt{q+\frac{1}{4}}}(r) = \frac{1}{n!}r^{-2\sqrt{q+\frac{1}{4}}} e^{2\sqrt{z}r}\frac{d^n}{ds^n} \left[r^{n+2\sqrt{q+\frac{1}{4}}}e^{-2\sqrt{z}r}\right]$$
(61)

And  $j_n = \frac{1}{n!}$ 

From equation (20), the eigenfunction becomes

$$\Psi(\mathbf{r}) = \varphi(\mathbf{r})\Upsilon(\mathbf{r}) = N_n r^{-\frac{1}{2} + \sqrt{q + \frac{1}{4}}} e^{-\sqrt{zr}} L_n^{2\sqrt{q + \frac{1}{4}}}(r)$$
(62)

Where  $N_n$  is a normalization constant of the eigenfunction  $\Psi(\mathbf{r})$ By substituting the values of z, w and q, we get the eigenfunction for a general inverse polynomial potential

$$\Psi(\mathbf{r}) = R(r)$$

$$= N_{n} r^{\frac{-1}{2} + \sqrt{A_{-2} + \frac{1}{2} \sum_{h=1}^{h} \frac{(h+2)(h+1)}{r_{0}^{h}} A_{-h-2} + \frac{1}{4}} e^{-\sqrt{A_{0} - \lambda^{2} + \frac{1}{2} \sum_{h=1}^{h} \left[\frac{h(h+1)A_{-h-2}}{r_{0}^{h+2}}\right]}} r \sum_{L_{n}}^{2} \sqrt{A_{-2} + \frac{1}{2} \sum_{h=1}^{h} \frac{(h+2)(h+1)}{r_{0}^{h}} A_{-h-2} + \frac{1}{4}} (r)$$

Hence,

$$U(r) = N_n r^{\frac{1}{2} + \sqrt{A_{-2} + \frac{1}{2} \sum_{h=1}^{h} \frac{(h+2)(h+1)}{r_0^h} \frac{A_{-h-2}}{4} + \frac{1}{4}} e^{-\sqrt{A_0 - \lambda^2 + \frac{1}{2} \sum_{h=1}^{h} \left[\frac{h(h+1)A_{-h-2}}{r_0^{h+2}}\right]}} r \sum_{L_n}^{2} \sqrt{A_{-2} + \frac{1}{2} \sum_{h=1}^{h} \frac{(h+2)(h+1)}{r_0^h} \frac{A_{-h-2}}{4} + \frac{1}{4}} (r)$$

For the potential in equation (2)

$$V(r) = \frac{\alpha}{r} + \frac{A}{r^2} + \frac{B}{r^3} + \frac{C}{r^4}$$

Where  $A_{-1} = \alpha$  ,

$$A_{-2} = A$$
 ,  $A_{-3} = B$  ,  $A_{-4}$   
= C and the rest of the summation parameters in V<sub>1</sub>(r) is equal to zero

The energy eigenvalues become

$$\lambda^{2} = \left[Br_{0}^{-3} + 3Cr_{0}^{-4}\right] - \frac{\left[3Br_{0}^{-2} + 8Cr_{0}^{-3} - \alpha\right]^{2}}{\left[(2n+1) \pm 2\sqrt{\left(A + \left[3Br_{0}^{-1} + 6Cr_{0}^{-2}\right] + \frac{1}{4}\right)}\right]^{2}}$$

And, the state eigenfunctions are given as

$$U(r) = N_n r^{\frac{1}{2} + \sqrt{A + [3Br_0^{-1} + 6Cr_0^{-2}] + \frac{1}{4}}} e^{-\sqrt{[Br_0^{-3} + 3Cr_0^{-4}] - \lambda^2}} r L_n^{\sqrt{A + [3Br_0^{-1} + 6Cr_0^{-2}] + \frac{1}{4}}}(r)$$

#### Conclusion

In the present paper, we discussed some special cases of the two-body potentials which have an oscillatory shape. The energy eigenvalues and eigenfunctions for bound states of such potentials have been obtained by Nikiforov-Uvarov method. The obtained results are useful in nuclear physics and quantum mechanics for magnetic interactions.

#### References

**[1]-**A. O. Barut and J. Kraus, Resonances in e<sup>+</sup>,e<sup>-</sup> system due to anomalous magnetic moment interactions, Physics Letters B**59**,175, 1975.

**[2]-** A. O. Barut and J. Kraus, Solution of the Dirac equation with Coulomb and magnetic moment interactions, Journal of Mathematical Physics **17**, 506, 1976.

**[3]-** A. O. Barut and J. Kraus, Form-factor corrections to superpositronium and short-distance behavior of the magnetic moment of the electron. , physical review D **16**, 161, 1977.

**[4]-** Alberto Sadum and Masoud Asadi-Zeydabadi, A Study of the Two-body Problem with Magnetic Interactions, Journal of physics, astrophysics and physical cosmology, **3**, Issue 1, 2009.

**[5]**- Bruno Gruber, two body problems with magnetic interaction "Symmetries in Science VIII", Springer Science, 1994.

[6]- M. Naghdi , Nucleon-Nucleon Interaction: A Typical/Concise Review, arXiv:nucl-th/0702078

**[7]**- Bernard Schaeffer, Electromagnetic Theory of the Nuclear Interaction, World Journal of Nuclear Science and Technology, **6**, no.4, 199, 2016.

**[8]-** F.BLOCH, Le moment magnétique du neutron, Annales de l'institut Henri Poincaré, **8**, no. 1, 63, 1938.

**[9]**-Richard P Feynman; Robert B Leighton, Matthew Linzee Sands, the Feynman lectures on physics, San Francisco, Pearson addition Wesley, 2006.

**[10]**-Rajmund Krivec, Mitja Rosina, Bojan Golli and Simon Širca, Few-Body Problems in Physics, Springer Wein New York, 2003; W.T. Grandy, book: Relativistic Quantum Mechanics of Leptons and Fields, Springer Science & Business Media,1990

[11]-A. O. Barut and G.strobel, Large Magnetic Moment Effects at High Energies and Composite Particle Models, Field Theory in Elementary Particles. Studies in the Natural Sciences (A Series from the Center for Theoretical Studies), Springer, Boston, 19, 323, 1983.
[12]-C. Prion, F. Reuse, Relativistic dynamics for the spin 1/2 particle, Helvetica Physica Act, 51, 146, 1978.

**[13]-**F. Reuse, A new relativistic model for the hydrogen atom, Helvetica Physica Acta, **51**, 157, 1978.

**[14]**-A. O. Barut, lectures on magnetic interaction of stable particles and magnetic resonances, mathematical reviews, American mathematical society, P 4933, 1982.

**[15]**-A. O. Barut, Magnetic resonances between massive and massless spin- 1/2 particles with magnetic moments, Journal of Mathematical Physics **21**, 568, 1980.

**[16]-**A. O. Barut, radial equations for Dirac particle with anomalous magnetic moment in a dipole field, quantum theory groups, fields and particles, mathematical physics studies, 1983.

**[17]**-G. Aad *et al.* (ATLAS Collaboration), Search for magnetic monopoles and stable particles with high electric charges in 8 TeV pp collisions with the ATLAS detector, Phys. Rev. D **93**, 052009,2016.

**[18]-**The MoEDAL collaboration, Search for magnetic monopoles with the MoEDAL prototype trapping detector in 8 TeV proton-proton collisions at the LHC, Journal of High Energy Physics, **8**, 67, 2016.

**[19]-**M. W. Ray, E. Ruokokoski, S. Kandel, M. Möttönen & D. S. Hall, Observation of Dirac monopoles in a synthetic magnetic field, Nature, **505**, 657, 2014.

[20]-C. Castelnovo, R. Moessner & S. L. Sondhi, Magnetic monopoles in spin ice, Nature 451, 42, 2008.

**[21]-**A.O. Barut, R. Rączka, Two-Fermion Equation with Form-Factors and Electromagnetic High Mass Resonances, acta physica polonica B**10**, 687,1979.

**[22]-**M.M. Bakri, H.M.M. Mansour, The Relativistic Two-Fermion Equations (I), acta physica polonica B**11**, 413, 1980.

**[23]-**M.M. Bakri, H.M.M. Mansour, The Relativistic Two-Fermion Equations (II), acta physica polonica B**11**, 543, 1980.

**[24]-**E. E. Salpeter and H. A. Bethe, A Relativistic Equation for Bound-State Problems, Physical Review, **84**, 1232, 1951.

**[25]-**W. Królikowski, J. Rzewuski, Covariant one-time formulation of the many-body problem in quantum theory, Il Nuovo Cimento, **2**, Issue 2, 203, 1955

**[26]-**W. Królikowski, J. Rzewuski, On " potentials " in the theory of quantized fields, Il Nuovo Cimento, **3**, Issue 2, 260, 1956.

**[27]-**W. Królikowski, J. Rzewuski, One-time formulation of the relativistic two-body problem. Separation of angular variables, Il Nuovo Cimento, **4**, Issue 5, 975, 1956.

**[28]-**W. Królikowski, J. Rzewuski, Remarks on composite and unstable objects in quantum field theory, Il Nuovo Cimento B, **25**, Issue 2, 739, 1975.

**[29]-**M. Hossein Partovi, New treatment of the bound-state problem in quantum field theory, physical review D **12**, 3887,1975.

**[30]**-H. Suura, Relativistic Two-Body Wave Equation, and Meson Spectrum, Physical Review Letters, **38**, 636,1977.

**[31]-**G .Preparata ,N.S .Craigie, A space-time description of quarks and hadrons, Nuclear Physics B **102**, Issue 3, 478, 1976.

**[32]**-A. O. Barut, M. Berrondo, and G. García-Calderón, Narrow resonances as an eigenvalue problem and applications to high energy magnetic resonances: An exactly soluble model, Journal of Mathematical Physics **21**, 1851, 1980.

**[33]**-H. M. M. Mansour, and H. J. W. Müller-Kirsten, Perturbative technique as an alternative to the WKB method applied to the double-well potential, Journal of Mathematical Physics **23**, 1835 ,1982.

**[34]-**H. M. M. Mansour and H. J. W. Müller-Kirsten, Matching of WKB solutions to other solutions, physical review, D **24**, 1033, 1981.

**[35]**-Andrzej A.Skorupski, Reports on Mathematical Physics, Improved higher order phaseintegral approximations of the JWKB type in the vicinity of zeros and singularities of the "wave number", **17**, Issue 2, 161, 1980.

[36] Nouredine Zettili, Quantum Mechanics Concepts and Applications, John Wiley, 2009.

[37] Cüneyt Berkdemir, Theoretical concept of quantum mechanics, In Tech, 2012.

[38] F. Nikiforov and V. B. Uvarov, Special Functions of Mathematical Physics, Birkhäauser, Basel, Switzerland, 1988

[39]-E. Eichten, K. Gottfried, T. Kinoshita, J. Kogut, K. D. Lane, and T. M. Yan, "Spectrum of charmed quark-antiquark bound states," Physical Review Letters, 34, no. 6, 369, 1975
[40]-S. M. Kuchin and N. V. Maksimenko, "Theoretical Estimations of the Spin – Averaged Mass Spectra of Heavy Quarkonia and Bc Mesons," Universal Journal of Physics and Application, 7, 295, 2013.

**[41]**-S. Laachir and A. Laaribi, "Exact Solutions of the Helmholtz equation via the Nikiforov-Uvarov Method," International Journal of Mathematical, Computational, Physical, Electrical and Computer Engineering, **7**, no. 1, 2013.

**[42]**-H. Ciftci, R. L. Hall, and N. Saad, "Iterative solutions to the Dirac equation," Physical Review A: Atomic, Molecular and Optical Physics, **72**, no. 2, part A, 022101, 2005.

**[43]**-M. Alberg and L. Wilets, "Exact solutions to the Schrödinger equation for potentials with Coulomb and harmonic oscillator terms," Physics Letters A, **286**, no. 1, 7, 2001.

**[44]**-B. Ita, P. Tchoua, E. Siryabe, and G. E. Ntamack, "Solutions of the Klein-Gordon Equation with the Hulthen Potential Using the Frobenius Method," International Journal of Theoretical and Mathematical Physics, **4**, no. 5, 173, 2014.

[45]-H. Goudarzi and V. Vahidi, "Supersymmetric approach for Eckart potential using the

{NU} method," Advanced Studies in Theoretical Physics, 5, no. 9, 469, 2011.
[46]-N. V. Masksimenko and S. M. Kuchin, "Determination of the mass spectrum of quarkonia by the Nikiforov–Uvarov method," Russian Physics Journal, 54, no. 57, 54, 2011.
[47]-Hesham Mansour and Ahmed Gamal, Bound State of Heavy Quarks Using a General Polynomial Potential, Advances in High Energy Physics, 1, Article ID 7269657, 2018.