

STUDY OF SOME HOLOMORPHIC CURVES IN \mathbb{C}^3 AND THEIR PROJECTION INTO THE COMPLEX PROJECTIVE SPACE $\mathbb{C}P^2$

FATHI HAGGUI AND ABDESSAMI JALLED

ABSTRACT. We study holomorphic curves $f : \mathbb{C} \rightarrow \mathbb{C}^3$ avoiding four complex hyperplanes and a real subspace of real dimension four or five in \mathbb{C}^3 . We show that the projection of f into the complex projective space $\mathbb{C}P^2$ is not necessarily constant.

keywords Complex projective space, Holomorphic curves, Kobayashi hyperbolicity

1. INTRODUCTION

The classical Picard Theorem [1] (see also [2]) states that every holomorphic map from the complex Euclidean space \mathbb{C} to $\mathbb{C}P^1$ that avoids three points, is constant. This Theorem has been extended to higher dimension by M.Green [3] who provided with examples of complex Kobayashi hyperbolic manifolds. We note that if H_1, \dots, H_m are complex hyperplanes in $\mathbb{C}P^n$, then they are said to be in general position if $m \geq n+1$ and any $(n+1)$ of these hyperplanes are linearly independent. Let us recall the Green Theorem:

Theorem (Green, [3]). Let C be a union of $2n+1$ complex hyperplanes in general position in $\mathbb{C}P^n$. Then, any holomorphic curve $f : \mathbb{C} \rightarrow \mathbb{C}P^n \setminus C$ is constant.

In particular, for $n=2$, any holomorphic curve $f : \mathbb{C} \rightarrow \mathbb{C}P^2 \setminus C$ is constant, where C is a union of five complex lines in general position in $\mathbb{C}P^2$.

As a direct consequence of the Green Theorem, the canonical projection into the complex projective space $\mathbb{C}P^2$ of any holomorphic map $f : \mathbb{C} \rightarrow \mathbb{C}^3$ which avoids five complex hyperplanes in \mathbb{C}^3 is constant, since its image avoids the projections of the five complex hyperplanes, which are complex projective lines in general position in $\mathbb{C}P^2$ (see Lemma 2.1). Our main goal is to study the projection into $\mathbb{C}P^2$ of a holomorphic curve $f : \mathbb{C} \rightarrow \mathbb{C}^3$ which avoids four complex hyperplanes in general position in \mathbb{C}^3 and a real subspace H of real dimension four or five and check if the projection remains constant.

Throughout the paper we identify \mathbb{R}^6 , endowed with its standard complex structure J_{st} , to \mathbb{C}^3 .

Definition 1.1. Let $n \geq 3$ and let $\mathcal{H} = (H_1, \dots, H_n)$ be a family of real subspaces of \mathbb{R}^6 such that $\text{codim}_{\mathbb{R}} H_j = 2$ for $j = 1, \dots, n$. Then \mathcal{H} is said to be in general position if for every 3-tuple (i, j, k) of distinct integers $i, j, k \in \{1, \dots, n\}$,

$$\text{Span}_{\mathbb{R}}(H_i^\perp, H_j^\perp, H_k^\perp) = \mathbb{R}^6.$$

Here, if H is a real subspace in \mathbb{R}^6 , then H^\perp denotes the orthogonal complement of H with respect to the Euclidean metric.

We first study the case of four real dimensional subspaces in \mathbb{C}^3 . We have the following

Theorem 1.

(i) Let H_1, \dots, H_n be n complex hyperplanes in \mathbb{C}^3 in general position ($n \geq 5$), then there

exists a non constant holomorphic curve $f : \mathbb{C} \rightarrow \mathbb{C}^3$ which avoid these n hyperplanes and $\pi(f)$ is constant.

(ii) Let H_1, H_2, H_3, H_4 be four complex hyperplanes in \mathbb{C}^3 . Then there exists a real subspace H of \mathbb{R}^6 , of real dimension four, such that (H, H_j, H_k) are in general position for all $j \neq k$, $j, k \in \{1, \dots, 4\}$, and there exists $f : \mathbb{C} \rightarrow \mathbb{C}^3$ holomorphic, such that

$$f(\mathbb{C}) \cap \left(\bigcup_{i=1}^4 H_i \cup H \right) = \emptyset$$

and $\pi(f)$ is non constant.

Remark 1.1. Here π denotes the canonical projection from $\mathbb{C}^3 \setminus \{0\}$ into $\mathbb{C}P^2$ and $\pi(f) := \pi \circ f$. Notice that $\pi(f)$ is well-defined in Theorem 1 (ii) since $f(\mathbb{C}) \subset \mathbb{C}^3 \setminus \{0\}$. In case (i), according to the Green Theorem and to Lemma 2.1 (see below), $\pi(f)$ is constant.

We study then the case of a subspace in \mathbb{C}^3 of real dimension five. We have the following:

Theorem 2. Let H_1, H_2, H_3, H_4 be four complex hyperplanes in \mathbb{C}^3 and let H be a real subspace of \mathbb{R}^6 of real dimension five. Let \tilde{H} be a complex hyperplane of \mathbb{C}^3 such that $\tilde{H} \subset H$. Then:

(1) If (\tilde{H}, H_j, H_k) are in general position for all $j \neq k$, $j, k \in \{1, \dots, 4\}$, then every holomorphic map $f : \mathbb{C} \rightarrow \mathbb{C}^3$ such that $f(\mathbb{C}) \cap \left(\bigcup_{i=1}^4 H_i \cup H \right) = \emptyset$ is constant.

(2) If there exist H_j, H_k , $j \neq k$, $j, k \in \{1, \dots, 4\}$, such that (\tilde{H}, H_j, H_k) are not in general position, then there exists $f : \mathbb{C} \rightarrow \mathbb{C}^3$, holomorphic, such that $f(\mathbb{C}) \cap \left(\bigcup_{i=1}^4 H_i \cup H \right) = \emptyset$ and $\pi(f)$ is non constant.

Remark 1.2.

- (a) The existence and uniqueness of $\tilde{H} \subset H$ is explained in the proof of Theorem 2.
- (b) The condition " (\tilde{H}, H_j, H_k) are not in general position" is equivalent to the condition " $\dim_{\mathbb{R}} \text{Span}_{\mathbb{R}}(\tilde{H}^{\perp}, H_j^{\perp}, H_k^{\perp}) = 4$ ".
- (c) The fact of considering four complex hyperplanes is an optimal condition (see the end of section two for more details).

The paper is organized as follows. In the first section, we give some results and properties. In section two, we prove Theorem 1. Finally, in section three, we prove Theorem 2.

2. PRELIMINARIES AND PROPERTIES

In 1972, Fujimoto [4] (see also M.Green[3] and [5]) showed a statement that characterizes the image of a holomorphic map $f : \mathbb{C} \rightarrow \mathbb{C}P^n$ omitting $(n+p)$ hyperplanes in general position. He proved the following

Theorem (Fujimoto [4], Serge Lang [5] pp 196). Let $f : \mathbb{C} \rightarrow \mathbb{C}P^n$ be holomorphic. Assume that the image of f lies in the complement of $n+p$ hyperplanes in general position, then this image is contained in a complex projective subspace of complex dimension $\leq [n/p]$.

The version of the Green Theorem stated in the introduction is a particular case of the previous Theorem, with $p = n + 1$.

In the remaining of the paper we will need the following properties satisfied by the canonical projection in \mathbb{CP}^2 of a holomorphic curve $f : \mathbb{C} \rightarrow \mathbb{C}^3$. For H a real subspace of \mathbb{R}^6 , we denote by H^* the set $H \setminus \{0\}$. Then, we have the following Lemma

Lemma 2.1. *Let $\pi : \mathbb{C}^3 \setminus \{0\} \rightarrow \mathbb{CP}^2$ be the canonical projection. Then:*

- (1) *If H is a complex hyperplane in \mathbb{C}^3 , then $\pi(H^*)$ is a complex projective line in \mathbb{CP}^2 .*
- (2) *If $f : \mathbb{C} \rightarrow \mathbb{C}^3$ is holomorphic and H is a complex hyperplane in \mathbb{C}^3 , then*

$$f(\mathbb{C}) \cap H = \emptyset \Rightarrow \pi(f)(\mathbb{C}) \cap \pi(H^*) = \emptyset.$$

- (3) *If H_1, H_2, H_3 are complex hyperplanes in general position in \mathbb{C}^3 , then $\pi(H_1^*), \pi(H_2^*), \pi(H_3^*)$ are in general position in \mathbb{CP}^2 .*

Notation: if $Z \in \mathbb{CP}^2$, we denote $[z_1 : z_2 : z_3]$ its homogeneous coordinates, where $(z_1, z_2, z_3) \in \mathbb{C}^3$.

Proof.

Point (1). We may assume that $H = \{(z_1, z_2, z_3) \in \mathbb{C}^3 / a_1 z_1 + a_2 z_2 + a_3 z_3 = 0\}$, with $a_1, a_2, a_3 \in \mathbb{C}, a_3 \neq 0$. Then

$$\begin{aligned} \pi(H^*) &= \{[1 : z_2 : z_3] \in \mathbb{CP}^2 / a_1 + a_2 z_2 + a_3 z_3 = 0\} \cup \{[0 : 1 : -\frac{a_2}{a_3}]\} \\ &= \{[1 : z : -\frac{a_1 + a_2 z}{a_3}], z \in \mathbb{C}\} \cup \{[0 : 1 : -\frac{a_2}{a_3}]\}. \end{aligned}$$

We notice that $[0 : 1 : -\frac{a_2}{a_3}]$ corresponds to $[\frac{1}{\infty} : 1 : -\frac{a_1 + a_2 \infty}{a_3 \infty}]$. Hence $\pi(H^*)$ is a projective complex line in \mathbb{CP}^2 .

Point (2). We first notice that $\pi(f)$ is well defined since, by assumption $f(\mathbb{C}) \cap H = \emptyset$, which implies that $f(\mathbb{C}) \subset \mathbb{C}^3 \setminus \{0\}$. Assume now, to get a contradiction, that $\pi(f)(\mathbb{C}) \cap \pi(H^*) \neq \emptyset$. Then there are two possibilities.

Case (α). There exists $z \in \mathbb{C}$ and there exists $\lambda \in \mathbb{C}$ such that

$$\pi(f)(z) = \left[1 : \lambda : -\frac{a_1 + a_2 \lambda}{a_3}\right].$$

Then, there exists $c_z \in \mathbb{C}^*$ such that $f(z) = (c_z, \lambda c_z, -\frac{a_1 + a_2 \lambda}{a_3} c_z)$. In particular $a_1 f_1(z) + a_2 f_2(z) + a_3 f_3(z) = 0$, where $f = (f_1, f_2, f_3)$. Hence, $f(z) \in H$. This is a contradiction.

Case (β). There exists $z \in \mathbb{C}$ such that

$$\pi(f)(z) = \left[0 : 1 : -\frac{a_2}{a_3}\right].$$

Then, there exists $c_z \in \mathbb{C}^*$ such that $f(z) = (0, c_z, -\frac{a_2}{a_3} c_z)$ and $a_1 f_1(z) + a_2 f_2(z) + a_3 f_3(z) = 0$. We obtain again that $f(z) \in H$: this is a contradiction.

Point (3). Since H_1, H_2, H_3 are complex hyperplanes in \mathbb{C}^3 , then there is a linear change

of coordinates such that the hyperplanes are defined by equations

$$\begin{aligned} H_1 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_1 = 0\}, \\ H_2 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_2 = 0\}, \\ H_3 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_3 = 0\}. \end{aligned}$$

Now by projection into $\mathbb{C}P^2$, we get

$$\begin{aligned} \pi(H_1^*) &= \{[0 : 1 : z]; z \in \mathbb{C} \cup \{\infty\}\} \cup [0 : 0 : 1], \\ \pi(H_2^*) &= \{[1 : 0 : z]; z \in \mathbb{C} \cup \{\infty\}\} \cup [0 : 0 : 1], \\ \pi(H_3^*) &= \{[1 : z : 0]; z \in \mathbb{C} \cup \{\infty\}\} \cup [0 : 1 : 0]. \end{aligned}$$

Hence $\pi(H_1^*) \cap \pi(H_2^*) \cap \pi(H_3^*) = \emptyset$, meaning that $\pi(H_1^*), \pi(H_2^*), \pi(H_3^*)$ are in general position since there is no triple point. \square

3. PROOF OF THEOREM 1

To prove theorem 1, we need the following Lemma which characterize the image of a holomorphic map $f : \mathbb{C} \rightarrow \mathbb{C}P^n$ avoiding $2n$ complex hyperplanes in general position. This precises the result of H.Fujimoto [4], [5] pp 196.

Definition 3.1. Let H_1, \dots, H_m , $m \geq 2n$, be hyperplanes of $\mathbb{C}P^n$. We call diagonal, a line passing through the two points $\bigcap_{i \in I} H_i$ and $\bigcap_{j \in J} H_j$, where $\text{card}(I) = \text{card}(J) = n$ and $I \cap J = \emptyset$.

Lemma 3.1. Let H_1, \dots, H_{2n} be $(2n)$ projective hyperplanes in general position in $\mathbb{C}P^n$. Then there are $\frac{1}{2}C_{2n}^n$ diagonals $\Delta_1, \dots, \Delta_{\frac{1}{2}C_{2n}^n}$ such that for every holomorphic curve $f : \mathbb{C} \rightarrow \mathbb{C}P^n \setminus \bigcup_{i=1}^{2n} H_i$, there exists $k_f \in \{1, \dots, \frac{1}{2}C_{2n}^n\}$ such that $f(\mathbb{C}) \subset \Delta_{k_f}$.

Proof. The proof is inspired by the Fujimoto Theorem, [5] pp 196.

Let $f : \mathbb{C} \rightarrow \mathbb{C}P^n$ be holomorphic, such that $f(\mathbb{C}) \cap \left(\bigcup_{i=1}^{2n} H_i\right) = \emptyset$.

Let L_1, \dots, L_{2n} be linear forms defining the hyperplanes H_1, \dots, H_{2n} , namely $H_k = L_k^{-1}(\{0\})$ for $k = 1, \dots, 2n$. If $f = [f_1 : \dots : f_{n+1}]$, we denote

$$h_k := H_k(f), \quad k = 1, \dots, 2n.$$

Let $I = \{1, \dots, 2n\}$ be the set of indices and \sim be the equivalence relation defined by $i \sim j$ if h_i/h_j is constant. We take a partition of the set of indices according to \sim . First, we know that the complement of a given class S has at most n elements (see [5] pp 197). Hence S has at least n elements and there are at most two classes.

The case of one class is not possible. In fact, There exists $\alpha_2, \dots, \alpha_{2n} \in \mathbb{C}$ such that

$$(S) \quad \begin{cases} h_2 &= \alpha_2 h_1 \\ h_3 &= \alpha_3 h_1 \\ &\vdots \\ h_{2n} &= \alpha_{2n} h_1 \end{cases}$$

Hence $f(\mathbb{C}) \subset (\bigcap_{k=2}^{n+1} H_k) \bigcap H_1 = \bigcap_{k=1}^{n+1} H_k = \emptyset$, which is impossible. Hence there are exactly two classes S_1 and S_2 .

We know that each of the two classes S_1, S_2 contains n elements. Then there exists a permutation $\sigma : \{1, \dots, 2n\} \rightarrow \{1, \dots, 2n\}$ such that

$$S_1 = \{\sigma(1), \dots, \sigma(n)\}, \quad S_2 = \{\sigma(n+1), \dots, \sigma(2n)\}.$$

Hence There exists $\alpha_2, \dots, \alpha_n, \beta_{n+1}, \dots, \beta_{2n-1} \in \mathbb{C}$ such that h_1, \dots, h_{2n} satisfy the systems:

$$(S_1) \begin{cases} h_{\sigma(2)} &= \alpha_2 h_{\sigma(1)} \\ h_{\sigma(3)} &= \alpha_3 h_{\sigma(1)} \\ \vdots & \\ h_{\sigma(n)} &= \alpha_n h_{\sigma(1)} \end{cases} \quad (S_2) \begin{cases} h_{\sigma(n+1)} &= \beta_{n+1} h_{\sigma(2n)} \\ h_{\sigma(n+2)} &= \beta_{n+2} h_{\sigma(2n)} \\ \vdots & \\ h_{\sigma(2n-1)} &= \beta_{2n-1} h_{\sigma(2n)} \end{cases}$$

Hence

$$\begin{cases} f(\mathbb{C}) \subset (\bigcap_{k=2}^n H_k) \bigcap H_1 &= \bigcap_{k=1}^n H_k \\ f(\mathbb{C}) \subset (\bigcap_{k=n+1}^{2n-1} H_k) \cap H_{2n} &= \bigcap_{k=n+1}^{2n} H_k \end{cases}$$

Then $f(\mathbb{C}) \subset \Delta_\sigma$, where Δ_σ is the unique diagonal (line) passing through the two points

$$\bigcap_{k=1}^n H_{\sigma(k)} \quad \text{and} \quad \bigcap_{k=n+1}^{2n} H_{\sigma(k)}.$$

Now the two points, and consequently Δ_σ , are completely determined by $S_1 = \{\sigma(1), \dots, \sigma(n)\}$ since S_2 is automatically fixed once S_1 is chosen. Hence Δ_σ is completely determined by a choice of a partition of $\{1, \dots, 2n\}$ into two subsets, each of them containing n elements.

There are exactly $\frac{1}{2} C_{2n}^n$ such partitions. This proves the Lemma. \square

We may prove now Theorem 1.

We denote by $z = (z_1, z_2, z_3)$ the coordinates in \mathbb{C}^3 , where $z_j = x_j + iy_j$, $j = 1, 2, 3$. Hence $(x_1, y_1, x_2, y_2, x_3, y_3)$ denote the coordinates in \mathbb{R}^6 .

Point (i). Consider first the case $n = 5$. By a linear change of coordinates, we take the hyperplanes H_1, H_2, H_3, H_4 and H_5 in standard form defined by the following equations

$$\begin{aligned} H_1 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_1 = 0\}, \\ H_2 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_2 = 0\}, \\ H_3 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_3 = 0\}, \\ H_4 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_1 + z_2 + z_3 = 0\}, \\ H_5 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / a_1 z_1 + a_2 z_2 + a_3 z_3 = 0\}, \quad a_j \in \mathbb{R} \setminus \{0\} \quad \forall j = 1, 2, 3. \end{aligned}$$

By hypothesis $f(\mathbb{C}) \cap (\bigcup_{i=1}^5 H_i) = \emptyset$. Then there exists $h_1, h_2, h_3 : \mathbb{C} \rightarrow \mathbb{C}$, holomorphic, such that

$$f = (e^{h_1}, e^{h_2}, e^{h_3}).$$

Moreover, since $\pi(f)(\mathbb{C})$ omits $\pi(H_i)$ for $i = 1, \dots, 5$ (see Lemma 2.1) and $\pi \circ f$ is constant by Green (see [3]), there exists $(\omega_1, \omega_2, \omega_3) \neq (0, 0, 0)$ such that for all $z \in \mathbb{C}$,

$$\left[e^{h_1(z)} : e^{h_2(z)} : e^{h_3(z)} \right] = [\omega_1 : \omega_2 : \omega_3].$$

Therefore

$$\left[1 : \frac{e^{h_2(z)}}{e^{h_1(z)}} : \frac{e^{h_3(z)}}{e^{h_1(z)}} \right] = \left[1 : \frac{\omega_2}{\omega_1} : \frac{\omega_3}{\omega_1} \right]$$

which implies that

$$\begin{cases} e^{h_2(z)-h_1(z)} = \frac{\omega_2}{\omega_1} \\ e^{h_3(z)-h_1(z)} = \frac{\omega_3}{\omega_1} \end{cases} \Rightarrow \begin{cases} e^{h_2(z)} = \frac{\omega_2}{\omega_1} e^{h_1(z)} \\ e^{h_3(z)} = \frac{\omega_3}{\omega_1} e^{h_1(z)} \end{cases}$$

Hence $f = (e^{h_1}, c_2 e^{h_1}, c_3 e^{h_1})$, with $1 + c_2 + c_3 \neq 0$, and f is not constant.

Essentially the same type of argument works in general. Let H_1, \dots, H_n , $n \geq 5$, be n hyperplanes defined by:

$$H_k := \left\{ Z \in \mathbb{C}^3 / \sum_{i=1}^3 \alpha_i^k z_i = 0, \alpha_i^k \in \mathbb{C}, 1 \leq k \leq n \right\}.$$

By hypothesis $f(\mathbb{C}) \cap \left(\bigcup_{i=1}^n H_i \right) = \emptyset$, then in particular $f(\mathbb{C}) \cap \left(\bigcup_{i=1}^5 H_i \right) = \emptyset$ and $f = (e^h, c_2 e^h, c_3 e^h)$ is not constant, where h is holomorphic from \mathbb{C} to \mathbb{C} .

Hence, in order that f avoids H_1, \dots, H_n , it is sufficient to choose $c_2, c_3 \in \mathbb{C}$ such that for every $k = 1, \dots, n$

$$\alpha_1^k + \alpha_2^k c_2 + \alpha_3^k c_3 \neq 0.$$

We point out that what precedes proves more generally that given a countable set of complex hyperplanes in \mathbb{C}^3 passing through the origin, there exists $f : \mathbb{C} \rightarrow \mathbb{C}^3$ not constant and avoiding each hyperplane. This proves Point (i).

Point (ii). Let H_1, H_2, H_3 and H_4 be four complex hyperplanes in general position in \mathbb{C}^3 . We know that there is a linear change of coordinate such that H_1, H_2, H_3 and H_4 are defined in standard form by :

$$\begin{aligned} H_1 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_1 = 0\}, \\ H_2 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_2 = 0\}, \\ H_3 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_3 = 0\}, \\ H_4 &= \{(z_1, z_2, z_3) \in \mathbb{C}^3 / z_1 + z_2 + z_3 = 0\}, \end{aligned}$$

Then

$$\begin{aligned} H_1^\perp &= \text{Span}_{\mathbb{R}}[(1, 0, 0, 0, 0); (0, 1, 0, 0, 0)], \\ H_2^\perp &= \text{Span}_{\mathbb{R}}[(0, 0, 1, 0, 0); (0, 0, 0, 1, 0)], \\ H_3^\perp &= \text{Span}_{\mathbb{R}}[(0, 0, 0, 0, 1); (0, 0, 0, 0, 1)], \\ H_4^\perp &= \text{Span}_{\mathbb{R}}[(1, 0, 1, 0, 1); (0, 1, 0, 1, 1)]. \end{aligned}$$

We pose now

$$H = \begin{cases} X_1 - X_2 &= 0 \\ X_1 - X_3 &= 0 \end{cases}$$

Then $H^\perp = \text{Span}_{\mathbb{R}}[(1, 0, 1, 0, 0, 0); (1, 0, 0, 0, 1, 0)]$, which of course satisfies the condition $\text{Span}_{\mathbb{R}}(H^\perp, H_j^\perp, H_k^\perp) = \mathbb{R}^6$ for all $j \neq k$, $j, k \in \{1, \dots, 4\}$.

Since $f(\mathbb{C}) \cap \left(\bigcap_{i=1}^4 H_i \right) = \emptyset$, then there exists holomorphic functions $f_i : \mathbb{C} \rightarrow \mathbb{C}$, $i = 1, 2, 3$ such that

$$f = (e^{f_1}, e^{f_2}, e^{f_3}).$$

Then, by Lemma 2.1 (2), $g := \pi(f)$ satisfies $g(\mathbb{C}) \subset \mathbb{C}P^2 \setminus \bigcup_{j=1}^4 \pi(H_j^\star)$. Hence g has the following form

$$(1) \quad g = [1 : e^{g_2} : e^{g_3}],$$

where $g_2 = f_2 - f_1$ and $g_3 = f_3 - f_1$. According to Lemma 3.1 there exists $\frac{1}{2}C_4^2 = 3$ diagonals $\Delta_{12,34}, \Delta_{13,24}, \Delta_{14,23}$ such that $g = \pi(f(\mathbb{C}))$ is contained in one of these diagonals, where $\Delta_{ij,kl}$ is the diagonal line passing through $(\pi(H_i^\star) \cap \pi(H_j^\star))$ and $(\pi(H_k^\star) \cap \pi(H_l^\star))$. We recall that

$$\begin{aligned} \pi(H_i^\star) &= \{[z_1 : z_2 : z_3] \in \mathbb{C}P^2 : z_i = 0\} \text{ For } j = 1, 2, 3, \\ \pi(H_4^\star) &= \{[z_1 : z_2 : z_3] \in \mathbb{C}P^2 : z_1 + z_2 + z_3 = 0\}. \end{aligned}$$

Hence $\Delta_{12,34}, \Delta_{13,24}, \Delta_{14,23}$ are given by

$$(2) \quad \begin{aligned} \Delta_{12,34} &= \{[z_1 : z_2 : z_3] \in \mathbb{C}P^2 : z_1 + z_2 = 0\}, \\ \Delta_{13,24} &= \{[z_1 : z_2 : z_3] \in \mathbb{C}P^2 : z_2 + z_3 = 0\}, \\ \Delta_{14,23} &= \{[z_1 : z_2 : z_3] \in \mathbb{C}P^2 : z_1 + z_3 = 0\}. \end{aligned}$$

Suppose that $g(\mathbb{C})$ is contained in $\Delta_{12,34}$, the cases $g(\mathbb{C}) \subset \Delta_{13,24}$ or $g(\mathbb{C}) \subset \Delta_{14,23}$ being similar. Then

$e^{g_2} + 1 = 0 \Rightarrow e^{g_2} = -1 \Rightarrow g = [1 : -1 : e^{g_3}]$, where $g_3 = f_3 - f_1$. Hence

$$(3) \quad f = (e^{f_1}, -e^{f_1}, e^{f_3}).$$

On another hand $f(\mathbb{C}) \cap H = \emptyset \Leftrightarrow \forall z \in \mathbb{C}, \begin{cases} \text{Re}(e^{f_1(z)}) & \neq 0 \\ \text{or} \\ \text{Re}(e^{f_1(z)} - e^{f_3(z)}) & \neq 0. \end{cases}$

We pose $f_3 = 2f_1$, then $f = (e^{f_1}, -e^{f_1}, e^{2f_1})$ avoids H . In fact

$$\text{Re}(e^{f_3}) = \text{Re}(e^{2f_1}) = \text{Re}(e^{f_1}e^{f_1}) = \text{Re}(e^{f_1})^2 - \text{Im}(e^{f_1})^2.$$

Now if $\text{Re}(e^{f_1(z)}) = 0$ for some $z \in \mathbb{C}$, then $\text{Im}(e^{f_1(z)}) \neq 0$ and consequently $\text{Re}(e^{f_3(z)}) \neq 0$. Hence $f(\mathbb{C}) \cap H = \emptyset$.

Finally, $\pi(f) = [1 : -1 : e^{f_1}]$ is not constant and $f(\mathbb{C}) \cap \left(\bigcap_{j=1}^4 H_j \cup H \right) = \emptyset$. This concludes the proof of Theorem 1.

4. PROOF OF THEOREM 2

Let H be a real subspace of \mathbb{C}^3 such that $\dim_{\mathbb{R}} H = 5$, then H contains a unique complex hyperplane \tilde{H} of \mathbb{C}^3 . Indeed, there exists $(a_1, b_1, a_2, b_2, a_3, b_3) \in \mathbb{R}^6 \setminus \{0\}$ such that

$$\begin{aligned}
H &= \left\{ (x_1, y_1, x_2, y_2, x_3, y_3) \in \mathbb{R}^6 / \sum_{j=1}^3 (a_j x_j + b_j y_j) = 0 \right\} \\
&= \left\{ z \in \mathbb{C}^3 / \operatorname{Re} \left(\sum_{j=1}^3 (a_j - ib_j) z_j \right) = 0 \right\}.
\end{aligned}$$

Hence $\tilde{H} := \left\{ z \in \mathbb{C}^3 / \sum_{j=1}^3 (a_j - ib_j) z_j = 0 \right\}$ is a complex hyperplane in \mathbb{C}^3 , contained in H .

Point (1). Assume that (\tilde{H}, H_j, H_k) are in general position for some $j \neq k$, $j, k \in \{1, \dots, 4\}$. Since $\tilde{H} \subset H$, where \tilde{H} is a complex hyperplane of \mathbb{C}^3 , and

$$f(\mathbb{C}) \cap \left(\bigcup_{i=1}^4 H_i \cup H \right) = \emptyset \Rightarrow f(\mathbb{C}) \cap \left(\bigcup_{i=1}^4 H_i \cup \tilde{H} \right) = \emptyset,$$

then it follows from Theorem 1 (i) that there is $(c_1, c_2) \in (\mathbb{C}^*)^2$ which satisfies $1 + c_2 + c_3 \neq 0$ and there exists $h : \mathbb{C} \rightarrow \mathbb{C}$ holomorphic such that

$$f(z) = (e^h, c_2 e^h, c_3 e^h).$$

On another hand $H := \{(x_1, y_1, \dots, x_3, y_3) \in \mathbb{R}^6 / \sum_{j=1}^3 (a_j x_j + b_j y_j) = 0\}$. By hypothesis $f(\mathbb{C}) \cap H = \emptyset$ then for every $z \in \mathbb{C}$ we have,

$$\begin{aligned}
&a_1 \operatorname{Re}(e^{h(z)}) + a_2 \operatorname{Re}(c_2 e^{h(z)}) + a_3 \operatorname{Re}(c_3 e^{h(z)}) \\
&+ b_1 \operatorname{Im}(e^{h(z)}) + b_2 \operatorname{Im}(c_2 e^{h(z)}) + b_3 \operatorname{Im}(c_3 e^{h(z)}) \neq 0.
\end{aligned}$$

Thus, for every $z \in \mathbb{C}$

$$\begin{aligned}
&\operatorname{Re}(e^{h(z)}) [a_1 + a_2 \operatorname{Re}(c_2) + a_3 \operatorname{Re}(c_3) + b_2 \operatorname{Im}(c_2) + b_3 \operatorname{Im}(c_3)] \\
&+ \operatorname{Im}(e^{h(z)}) [b_1 + b_2 \operatorname{Re}(c_2) + b_3 \operatorname{Re}(c_3) - a_1 \operatorname{Im}(c_2) - a_3 \operatorname{Im}(c_3)] \neq 0.
\end{aligned}$$

We denote

$$a := [a_1 + a_2 \operatorname{Re}(c_2) + a_3 \operatorname{Re}(c_3) + b_2 \operatorname{Im}(c_2) + b_3 \operatorname{Im}(c_3)]$$

$$b := [b_1 + b_2 \operatorname{Re}(c_2) + b_3 \operatorname{Re}(c_3) - a_1 \operatorname{Im}(c_2) - a_3 \operatorname{Im}(c_3)]$$

then

$$f(\mathbb{C}) \cap H = \emptyset \Leftrightarrow e^{h(\mathbb{C})} \cap \{(x, y) \in \mathbb{R}^2 / ax + by = 0\} = \emptyset.$$

However $\{(x, y) \in \mathbb{R}^2 / ax + by = 0\}$ is either a real line or \mathbb{R}^2 , depending on the values of a and b . Then by the little Picard Theorem e^h is constant because it avoids an infinite number of points. Hence h is constant and f is then constant. We point out that the projection of f into $\mathbb{C}P^2$ is also constant.

Point (2). Suppose there exists $j \neq k$, $j, k \in \{1, \dots, 4\}$, such that

$\dim_{\mathbb{R}} \operatorname{Span}_{\mathbb{R}}(\tilde{H}^\perp, H_j^\perp, H_k^\perp) = 4$. Then:

$$\tilde{H}^\perp \subset \operatorname{Span}_{\mathbb{R}}(H_j^\perp, H_k^\perp).$$

In fact for all $i = 1, \dots, 4$, $\dim_{\mathbb{R}} H_i^{\perp} = 2$, then $\dim_{\mathbb{R}} \text{Span}_{\mathbb{R}}(H_i^{\perp}, H_j^{\perp}) = 4$. Suppose $\tilde{H}^{\perp} \subset \text{Span}_{\mathbb{R}}(H_1^{\perp}, H_2^{\perp})$ then there exists $\alpha_1, \alpha_2 \in \mathbb{C}$ such that

$$\tilde{H} = \{\alpha_1 z_1 + \alpha_2 z_2 = 0\}.$$

Since $f(\mathbb{C}) \cap (\bigcup_{i=1}^4 H_i) = \emptyset$, then by 3

$$f = (e^{f_1}, -e^{f_1}, e^{f_3}).$$

We take $f_1 = c$, $c \in \mathbb{C} \setminus \{0\}$, such that $\text{Re}(\alpha_1 e^c - \alpha_2 e^c) \neq 0$ and f_3 not constant. Then

$$f = (c, -c, e^{f_3})$$

avoids $\bigcup_{i=1}^4 H_i \bigcup H$, and $\pi(f)$ is not constant. This concludes the proof of Theorem 2. \square

By the end of the paper, we show the optimality of considering four complex hyperplanes. Let H_1, H_2, H_3 be three complex hyperplanes in \mathbb{C}^3 , then there exists H a real hyperplane in \mathbb{R}^6 and a complex hyperplane \tilde{H} contained in H , $(H_1, H_2, H_3, \tilde{H})$ are in general position, and there exists $f : \mathbb{C} \rightarrow \mathbb{C}^3$, holomorphic, such that $f(\mathbb{C}) \cap (\bigcup_{j=1}^3 H_j \bigcup H) = \emptyset$

and $\pi \circ f$ is not constant. In fact:

We pose $H = \{x_1 + x_2 + x_3 = 0\}$ and $\tilde{H} = \{z_1 + z_2 + z_3 = 0\}$, which is clearly contained in H .

Since $f(\mathbb{C}) \cap (\bigcup_{j=1}^3 H_j \bigcup H) = \emptyset$, then $f(\mathbb{C}) \cap (\bigcup_{j=1}^3 H_j \bigcup \tilde{H}) = \emptyset$ and $f = (e^{f_1}, e^{f_2}, e^{f_3})$.

Hence

$$g := \pi(f) = [1 : e^{g_2} : e^{g_3}],$$

where $g_2 = f_2 - f_1$ and $g_3 = f_3 - f_1$. By lemma 3.1, $g := \pi(f)$ is contained in one of diagonals $\Delta_{12,34}, \Delta_{13,24}, \Delta_{14,23}$ (see 2). Suppose $\pi(f)(\mathbb{C}) \subset \Delta_{13,24}$, then

$$\pi(f) = [1, e^{g_2}, -e^{g_2}],$$

Hence $f = (1, e^{g_2}, -e^{g_2})$ avoids $(\bigcup_{j=1}^3 H_j \bigcup H)$ and $\pi(f)$ is not constant.

REFERENCES

- [1] GREEN, M. Some Picard theorems for holomorphic maps to algebraic varieties. Amer. J. Math. 97 (1975), pp. 43-75.
- [2] Davis, B. Picard's theorem and Brownian motion. Trans. Amer. Math. Soc. 213 (1975), 353-362.
- [3] GREEN, M. Holomorphic maps into complex projective space omitting hyperplanes, Trans. Amer. Math. Soc., 169 (1972), pp. 89-103.
- [4] FUJIMOTO, H. On holomorphic maps into a taut complex space, Nagoya Math. J., 46 (1972), pp. 49-61.
- [5] Lang, S. Introduction to Complex Hyperbolic Spaces, New York, Inc. in 1987, pp. 196

E-mail address: fathi.haggui@gmail.com

jalled.abdessami90@gmail.com

Institut préparatoire aux études d'ingénieur de Monastir Rue Ibn Eljazzar - 5019 Monastir TUNISIE