

A simple approach to the quantum theory of nonlinear fiber optics

J. Bonetti, A. D. Sánchez, S. M. Hernandez, and D. F. Grosz
*Instituto Balseiro, Av. Bustillo km 9.500, Bariloche (R8402AGP), Argentina and
 Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina*

We put forth an approach to obtain a quantum master equation for the propagation of light in nonlinear fiber optics by relying on simple quantum pictures of the processes (linear and nonlinear) occurring along propagation in an optical fiber. This equation is shown to be in excellent agreement with the classical Generalized Nonlinear Schrödinger Equation and predicts the effects of self-steepening and spontaneous Raman scattering. Last, we apply these results to the analysis of two cases of relevance in quantum technologies: single-photon frequency translation and spontaneous four-wave mixing.

The nonlinear fiber optics realm is ruled by the Generalized Nonlinear Schrödinger Equation (GNLSE) [1], which describes the propagation of classical light pulses. This equation is not useful in quantum-technology applications since it cannot be applied to non-classical light [2–5]. For these cases, a master equation describing the propagation of the quantum state of light (ρ), instead of the classical field, is required

$$\frac{\partial \rho}{\partial z} = \mathcal{F}(\rho), \quad (1)$$

an equation that, to the best of our knowledge, has not yet been developed. A starting point could be the quantum theories for nonlinear fibers [6–11], namely

$$\frac{\partial \rho}{\partial t} = \text{Tr}_{\text{fiber}} \left\{ \frac{1}{i\hbar} [\hat{H}, \rho] \right\}, \quad (2)$$

where ρ is the density matrix of the system, including both light and optical fiber, $\rho = \text{Tr}_{\text{fiber}} \{ \rho \}$ is the reduced density matrix for the quantum state of light, in which we are interested, and $\hat{H} = \hat{H}_{\text{light}} + \hat{H}_{\text{fiber}} + \hat{H}_{\text{int}}$ [10, 11]. However, this equation is not easily solvable due to the complexity of the operators involved and the large dimension of ρ .

In this paper we propose a simple approach to Eq. (1), avoiding the difficult derivation from Eq. (2). Our proposal relies on simple quantum pictures of the different processes that occur while propagation in a nonlinear fiber in terms of creation and annihilation of photons [6]. Such an approach is shown not only to be in excellent agreement with the GNLSE in the classical limit but, also, to predict the effect of self-steepening [1], an aspect that has not been studied from a quantum point of view in the literature, and the effect of spontaneous Raman scattering (SpRS). Moreover, this master equation is readily applicable to relevant schemes found in quantum technologies. As an example, we apply it for a novel analysis of frequency translation of single-photons by Bragg scattering (BS) [3] and spontaneous four-wave mixing (SpFWM), a crucial process in heralded single-photon sources [5].

In the remainder of the text we use \int to represent $\int_{-\infty}^{\infty}$. We start by the GNLSE describing the propagation of the complex envelope of the electrical field $A(z, t)$ in a

polarization-preserving fiber

$$i \frac{\partial A}{\partial z} = \left(-i \frac{\hat{\alpha}}{2} + \hat{\beta} \right) A - \gamma A \int_0^{\infty} R(t') |A(t-t')|^2 dt', \quad (3)$$

where $|A|^2$ is the optical power, $\hat{\alpha}$ and $\hat{\beta}$ are linear operators defined as $\hat{\alpha} e^{-i\omega t} = \alpha_{\omega} e^{-i\omega t}$, $\hat{\beta} e^{-i\omega t} = -\beta_{\omega} e^{-i\omega t}$. α_{ω} and β_{ω} are the frequency profiles of loss and dispersion, respectively. γ is the fiber nonlinear coefficient and $R(t)$ is the nonlinear-response function including both the instantaneous (electronic) and delayed Raman response [1]. We replace $A(z, t) = (2\pi)^{-1/2} \int A_{\omega}(z) e^{-i\omega t} d\omega$ into Eq. (3) to obtain the frequency domain version of this equation,

$$\frac{\partial A_{\omega}}{\partial z} = \left(-\frac{\alpha_{\omega}}{2} + i\beta_{\omega} \right) A_{\omega} + i\tilde{\gamma} \iint R_{\mu} A_{\omega'}^* A_{\omega-\mu} A_{\omega'+\mu} d\mu, \quad (4)$$

where $\tilde{\gamma} = \gamma/2\pi$ and $R_{\omega} = \int R(t') e^{i\omega t'} dt'$.

In a similar way to the work done in Ref. [12] we perform the quantization of A (see Appendix A) by proposing the correspondence with the quantum operators \hat{A}_{ω} , related to the annihilation and creation operators by

$$\hat{A}_{\omega} = \sqrt{\frac{\hbar\omega}{\Delta\omega}} \hat{a}_{\omega}, \quad (5)$$

where $\omega = \omega_0 + w$ is the photon frequency of the envelope mode w , ω_0 is the envelope central frequency, $\Delta\omega = 2\pi/T$ and T is the quantization period. The number operators $\hat{n}_{\omega} = \hat{a}_{\omega}^{\dagger} \hat{a}_{\omega}$ are interpreted as the quantity of photons of frequency ω within T .

The master equation proposed in this work is a standard Lindblad equation [13] that reads

$$\begin{aligned} \frac{\partial \rho}{\partial z} = & i[\hat{H}_{\text{disp}} + \hat{H}_{\text{fwm}}, \rho] + \\ & \int \hat{L}_{\alpha\omega'} \rho \hat{L}_{\alpha\omega'}^{\dagger} - \frac{1}{2} \{ \rho, \hat{L}_{\alpha\omega'}^{\dagger} \hat{L}_{\alpha\omega'} \} d\omega' + \\ & \int_0^{\infty} \hat{L}_{R\mu} \rho \hat{L}_{R\mu}^{\dagger} - \frac{1}{2} \{ \rho, \hat{L}_{R\mu}^{\dagger} \hat{L}_{R\mu} \} d\mu, \quad (6) \end{aligned}$$

where

$$\hat{H}_{\text{disp}} = \int \frac{\beta_{w'}}{\hbar\omega'} \hat{A}_{w'}^\dagger \hat{A}_{w'} dw', \quad (7)$$

$$\hat{H}_{\text{fwm}} = \iiint \frac{\tilde{\gamma} R_\mu^R}{2\hbar\omega_0} \hat{A}_{w_1}^\dagger \hat{A}_{w_2}^\dagger \hat{A}_{w_1-\mu} \hat{A}_{w_2+\mu} dw_1 dw_2 d\mu, \quad (8)$$

$$\hat{L}_{\alpha w'} = \sqrt{\frac{\alpha_{w'}}{\hbar\omega'}} \hat{A}_{w'}, \quad (9)$$

$$\hat{L}_{R\mu} = \int \sqrt{\frac{2\tilde{\gamma} R_\mu^I}{\hbar\omega_0}} \hat{A}_{w'-\mu}^\dagger \hat{A}_{w'} dw'. \quad (10)$$

R_μ^R and R_μ^I are respectively the real and imaginary parts of R_μ . In this equation we can easily recognize the four basic processes of the GNLSE: loss (\hat{L}_α), dispersion (\hat{H}_{disp}), four-wave mixing (\hat{H}_{fwm}) and stimulated Raman scattering (SRS) (\hat{L}_R).

We test this master equation by calculating the evolution of the mean values $\langle \hat{A}_w \rangle$. For this it is important the commutation relations of the operators \hat{A}_w , that in the limit $T \rightarrow \infty$ read $[\hat{A}_w, \hat{A}_{w'}] = 0$ and $[\hat{A}_w, \hat{A}_{w'}^\dagger] = \hbar\omega\delta(w - w')$. The calculation is quite involved (see Appendix B) but straightforward and gives

$$\begin{aligned} \frac{\partial}{\partial z} \langle \hat{A}_w \rangle &= \left(-\frac{\alpha_w}{2} + i\beta_w \right) \langle \hat{A}_w \rangle + \\ &i\tilde{\gamma} \left(1 + \frac{w}{\omega_0} \right) \iint R_\mu \langle \hat{A}_{w'}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'+\mu} \rangle dw' d\mu - \\ &\tilde{\gamma} \left(1 + \frac{w}{\omega_0} \right) \int_0^\infty R_\mu^I \hbar(\omega - \mu) d\mu \langle \hat{A}_w \rangle. \end{aligned} \quad (11)$$

Observe that the evolution of the mean values is in excellent agreement with the GNLSE. Further, the factor $(1 + w/\omega_0)$, that introduces a frequency dependence of the nonlinear coefficient $\tilde{\gamma}$, does not represent a departure from the classical limit. Instead, it appears naturally in the derivation and accounts for the well-known phenomenon in nonlinear fiber optics of *self-steepening* [1]. Although often associated with the distortion of ultrashort pulses, recent work has shown this effect to be of importance in the context of modulation instability and its complex interplay with stimulated Raman scattering [14, 15]. It must be emphasized that the effect of self-steepening has been overlooked in earlier quantum theories underlying the GNLSE, but it is *built in* in Eq. (6).

The operator \hat{L}_R in the master equation gives a clear picture of the Raman scattering process, interpreted as the annihilation of a photon at the pump frequency w' and the creation of a photon at the signal frequency $w' - \mu$. This process can occur even when there are no




FIG. 1. Frequency configuration of a degenerate four-wave mixing process.




FIG. 2. Degenerate SpFWM without (a) and with (b) loss and SRS. Probabilities of off-diagonal elements in (b) seriously affect the heralded-photon scheme.

photons present at the signal, producing the effect known as spontaneous Raman scattering (SpRS). It is the source of the last term in the Eq. (11) and cannot be obtained from the classical GNLSE. We use this term to calculate the loss of a pump, at frequency ω_0 , induced by SpRS and obtain

$$\frac{\partial I_p}{\partial z} = -2\tilde{\gamma} \left[\int_0^\infty R_\mu^I \hbar(\omega_0 - \mu) d\mu \right] I_p. \quad (12)$$

This result is in agreement with earlier works on Raman scattering [16, 17].

Despite the large size of the density matrix, we can use the master equation to study quantum-technology devices. In these applications we are interested in few-photons cases and hence the density matrix is reduced to numerically manageable sizes. We analyze two relevant cases depicting the interaction of few-photons signals with strong pumps. The quantum propagation of these small signals is derived from Eq. (6) by considering the pumps as classical and undepleted, and regarding only frequencies of interest.

Degenerate SpFWM, showed in Fig. 1 is a parametric non-classical phenomenon by which two photons of a

noiseless CW pump (p) are annihilated and simultaneously two photons are created in the signal (s) and the idler (i). This effect can be exploited to design heralded-photon sources by detecting one of the created photons to infer the presence of the other [5]. Loss and SRS affect this scheme, producing false heralding or non-heralded photons.

We simplify the master equation by proposing $\hat{A}_w = \sqrt{2\pi P}e^{ik_p z}\delta(w - w_p) + \hat{A}_{w_s}\delta_{ww_s} + \hat{A}_{w_i}\delta_{ww_i}$, $\alpha_{w_p} = 0$ and the phase-matching condition to obtain (see Appendix C)

$$\frac{\partial \rho}{\partial z} = i[\hat{H}, \rho] + \sum_{\nu=1}^3 \hat{L}_\nu \rho \hat{L}_\nu^\dagger - \frac{1}{2}\{\rho, \hat{L}_\nu^\dagger \hat{L}_\nu\}, \quad (13)$$

where

$$\hat{H} = k_s \hat{b}_s^\dagger \hat{b}_s + k_i \hat{b}_i^\dagger \hat{b}_i + \gamma P R_\Omega^R (\hat{b}_s^\dagger \hat{b}_i^\dagger + \hat{b}_s \hat{b}_i), \quad (14)$$

$$\hat{L}_1 = \sqrt{\alpha_s} \hat{b}_s, \quad (15)$$

$$\hat{L}_2 = \sqrt{\alpha_i} \hat{b}_i, \quad (16)$$

$$\hat{L}_3 = \sqrt{2\gamma P R_\Omega^I} (\hat{b}_s + \hat{b}_i). \quad (17)$$

In this case $k_p = \beta_{w_p} + \gamma P$, $k_x = \beta_{w_x} + \gamma P(1 + R_\Omega^R)$, and phase-matching means $2k_p - k_s - k_i = 0$.

In Fig. 2 we show numerical results for a particular case. The parameters are those of the frequency translation example. The probability of measuring n_s photons in the frequency s and n_i photons in the frequency i is calculated by $\langle n_s n_i | \rho | n_s n_i \rangle$. Loss and SRS seriously affect the heralding scheme by the possibility of mismatch between the photon numbers n_s and n_i . This simple calculation proves useful to accurately evaluate the performance of a heralded-photon scheme in the presence of loss and SRS.

Bragg Scattering (BS), showed in Fig. 3 is a well-known four-wave mixing process by which two strong pumps (p_1 and p_2) interact with two small signals (signal s and idler i). The quantum mechanical description of BS implies the annihilation of one photon in s and the simultaneous creation of one photon in i . Particularly, single-photon inputs in BS processes are of significant importance in quantum technologies [3]. To the best of our knowledge, the analysis of single-photon translation including loss and SRS has not been reported in the literature before due to the difficulty in including these effects in quantum models [16].

As in the previous case, we simplify Eq. (6) by proposing $\hat{A}_w = \sqrt{2\pi P}e^{ik_{p1}z}\delta(w - w_{p1}) + \sqrt{2\pi P}e^{ik_{p2}z}\delta(w - w_{p2}) + \hat{A}_{w_s}\delta_{ww_s} + \hat{A}_{w_i}\delta_{ww_i}$, where P is the optical power of the pumps, and $k_{px} = \beta_{w_x} + \gamma P(2 + R_\phi^R)$. We assume that the pumps do not interact by SRS ($R_\phi^I = 0$) and




FIG. 3. Frequency configuration of the Bragg scattering process.

$\alpha_{w_{p1}} = \alpha_{w_{p2}} = 0$. The simplified master equation for this particular case reads

$$\frac{\partial \rho}{\partial z} = i[\hat{H}, \rho] + \sum_{\nu=1}^5 \hat{L}_\nu \rho \hat{L}_\nu^\dagger - \frac{1}{2}\{\rho, \hat{L}_\nu^\dagger \hat{L}_\nu\}, \quad (18)$$

where

$$\hat{H} = k_s \hat{b}_s^\dagger \hat{b}_s + k_i \hat{b}_i^\dagger \hat{b}_i + \gamma P (R_\phi^R + R_\Phi^R) (\hat{b}_s^\dagger \hat{b}_i + \hat{b}_i^\dagger \hat{b}_s), \quad (19)$$

$$\hat{L}_1 = \sqrt{\alpha_s} \hat{b}_s, \quad (20)$$

$$\hat{L}_2 = \sqrt{\alpha_i} \hat{b}_i, \quad (21)$$

$$\hat{L}_3 = \sqrt{2\gamma P R_{\Phi-\phi}^I} \hat{b}_s, \quad (22)$$

$$\hat{L}_4 = \sqrt{2\gamma P R_{\Phi+\phi}^I} \hat{b}_i, \quad (23)$$

$$\hat{L}_5 = \sqrt{2\gamma P R_\Phi^I} (\hat{b}_s + \hat{b}_i). \quad (24)$$

We introduced the annihilation operators $\hat{b}_x = e^{-ik_x z} \hat{a}_{w_x}$, with $k_s = \beta_{w_s} + 2\gamma P(R_\Phi^R + R_{\Phi-\phi}^R)$, $k_i = \beta_{w_i} + 2\gamma P(R_\Phi^R + R_{\Phi+\phi}^R)$ and assumed the perfect phase-matching condition $k_{p1} - k_{p2} + k_i - k_s = 0$.

We numerically solve this system for the single-photon input with $\gamma = 1 \text{ W}^{-1}\text{km}^{-1}$, $P = 1 \text{ W}$, $\alpha_{w_s} = \alpha_{w_i} = 0.01 \text{ km}^{-1}$, $R_\Phi^R = R_\phi^R = 1$, $R_\Phi^I = R_{\Phi-\phi}^I = R_{\Phi+\phi}^I = 0.1$, $\omega_s \simeq \omega_i \simeq \omega_0$ and a fiber length of 5 km. In Fig. 4 we show the probabilities of finding the photon at w_s or w_i , calculated by $\langle 10 | \rho | 10 \rangle$ and $\langle 01 | \rho | 01 \rangle$. This calculation can be used to evaluate the efficiency of the single-photon frequency translation process even when loss and SRS effects are taken into account.

In conclusion, we proposed a simple approach to a quantum theory of light propagation in nonlinear fiber optics. By computing the evolution of the field-operator mean values we showed excellent agreement with the




FIG. 4. Propagation of a single-photon in a frequency-translation scheme, without (a) and with (b) loss and SRS.

APPENDIX A

A quantization of the field A is carried out in a similar fashion as standard quantum optics texts, i. e., by establishing an analogue between optical modes and harmonic oscillators. Thus, we propose $A(t)$ to be a periodic solution of period T that can be represented by the (Fourier) series

$$A(t) = \sum_{m=-\infty}^{\infty} O_m(t) e^{i\omega_0 t},$$

where

$$O_m(t) = \frac{\Delta w}{\sqrt{2\pi}} A_m e^{-i\omega_m t},$$

with $\Delta w = 2\pi/T$, $\omega_m = \omega_0 + m\Delta w$, and ω_0 is the central frequency of the envelope A . We consider the $O_m(t)$ as solutions to oscillators

$$\frac{\partial}{\partial t} O_m = -i\omega_m O_m.$$

The energy carried by each oscillator within the period T is $E_m = T|O_m|^2$. By decomposing these solutions as $O_m = O_m^R + iO_m^I$, we obtain equations

$$\frac{\partial}{\partial t} O_m^R = \omega_m O_m^I, \quad \frac{\partial}{\partial t} O_m^I = -\omega_m O_m^R, \quad E_m = T \left((O_m^R)^2 + (O_m^I)^2 \right). \quad (25)$$

Introducing variables $x_m = \sqrt{\frac{2T}{M\omega_m^2}} O_m^R$ and $p_m = \sqrt{2MT} O_m^I$, where M is an arbitrary mass, and replacing them into Eqs. (25) we obtain a classical harmonic oscillator

$$\frac{\partial x_m}{\partial t} = \frac{p_m}{M}, \quad \frac{\partial p_m}{\partial t} = -M\omega_m^2 x_m, \quad E_m = \frac{M\omega_m^2}{2} x_m^2 + \frac{p_m^2}{2M},$$

whose quantization is given by the operators \hat{H}_m , \hat{x}_m and \hat{p}_m such that

$$\hat{H}_m = \frac{M\omega_m^2}{2} \hat{x}_m^2 + \frac{\hat{p}_m^2}{2M}, \quad [\hat{x}_m, \hat{p}_m] = i\hbar.$$

An annihilation operator is defined correspondingly as

$$\hat{a}'_m = \sqrt{\frac{M\omega_m}{2\hbar}} \left(\hat{x}_m + \frac{i}{M\omega_m} \hat{p}_m \right), \quad (26)$$

and the number operator $\hat{n}_m = \hat{a}'_m^\dagger \hat{a}'_m$ is interpreted as the quantity of photons of frequency ω_m within the period T .

From Eq. (26) we can obtain operators \hat{A}_m by noting that

$$\hat{a}'_m = \sqrt{\frac{T}{\hbar\omega_m}} \hat{O}_m = \sqrt{\frac{\Delta w}{\hbar\omega_m}} \hat{A}_m e^{-i\omega_m t},$$

and also deduce the commutation relations

$$[\hat{A}_m, \hat{A}_n] = [\hat{A}_m^\dagger, \hat{A}_n^\dagger] = 0, \\ [\hat{A}_m, \hat{A}_n^\dagger] = \frac{\hbar\omega_m}{\Delta w} \delta_{nm},$$

classical GNLS. Further, the proposed quantum master equation leads naturally to the effects of spontaneous Raman scattering and self-steepening, the latter being included for the first time in a quantum framework. Finally, we applied the proposed approach to two cases of singular relevance in quantum technologies: single-photon frequency translation and spontaneous four-wave mixing.

where δ_{nm} is the Kroenecker's delta. In the limit where $T \rightarrow \infty$, $\Delta w \rightarrow 0$ we can deduce that $[\hat{A}_w, \hat{A}_{w'}^\dagger] = \hbar\omega\delta(w - w')$.

When dealing with number states, it is useful to define annihilation operators

$$\hat{a}_m = \hat{a}'_m e^{i\omega_m t} = \sqrt{\frac{\Delta w}{\hbar\omega_m}} \hat{A}_m,$$

for as it can be readily noticed, $\hat{n}_m = \hat{a}'_m^\dagger \hat{a}'_m = \hat{a}_m^\dagger \hat{a}_m$.

APPENDIX B

In the following we derive the evolution of the mean value of \hat{A}_w . For the sake of clarity, we deal separately with the four quantum processes that add up to the right hand side of Eq. 6.

Dispersion

$$i[\hat{A}_w, \hat{H}_{\text{disp}}] = i \left[\hat{A}_w, \int \frac{\beta_{w'}}{\hbar\omega'} \hat{A}_{w'}^\dagger \hat{A}_{w'} \partial w' \right] = i \int \frac{\beta_{w'}}{\hbar\omega'} [\hat{A}_w, \hat{A}_{w'}^\dagger] \hat{A}_{w'} \partial w' = i \int \beta_{w'} \delta(w - w') \hat{A}_{w'} \partial w' = i\beta_w \hat{A}_w.$$

$$\frac{\partial}{\partial z} \langle A_w \rangle = \langle i[\hat{A}_w, \hat{H}_{\text{disp}}] \rangle = i\beta_w \langle \hat{A}_w \rangle.$$

Four-wave mixing

$$\begin{aligned} i[\hat{A}_w, \hat{H}_{\text{FWM}}] &= i \left[\hat{A}_w, \tilde{\gamma} \iiint \frac{R_\mu^R}{2\hbar\omega_0} \hat{A}_{w_1}^\dagger \hat{A}_{w_2}^\dagger \hat{A}_{w_1-\mu} \hat{A}_{w_2+\mu} \partial w_1 \partial w_2 \partial \mu \right] = i\tilde{\gamma} \iiint \frac{R_\mu^R}{2\hbar\omega_0} [\hat{A}_w, \hat{A}_{w_1}^\dagger \hat{A}_{w_2}^\dagger \hat{A}_{w_1-\mu} \hat{A}_{w_2+\mu}] \partial w_1 \partial w_2 \partial \mu = \\ & i\tilde{\gamma} \iiint \frac{R_\mu^R}{2\hbar\omega_0} \left([\hat{A}_w, \hat{A}_{w_1}^\dagger] \hat{A}_{w_2}^\dagger \hat{A}_{w_1-\mu} \hat{A}_{w_2+\mu} + \hat{A}_{w_1}^\dagger [\hat{A}_w, \hat{A}_{w_2}^\dagger] \hat{A}_{w_1-\mu} \hat{A}_{w_2+\mu} \right) \partial w_1 \partial w_2 \partial \mu = \\ & i\tilde{\gamma} \iiint \frac{R_\mu^R \omega}{2\omega_0} \left(\delta(w - w_1) \hat{A}_{w_2}^\dagger \hat{A}_{w_1-\mu} \hat{A}_{w_2+\mu} + \hat{A}_{w_1}^\dagger \delta(w - w_2) \hat{A}_{w_1-\mu} \hat{A}_{w_2+\mu} \right) \partial w_1 \partial w_2 \partial \mu = \\ & i\tilde{\gamma} \int \int \frac{R_\mu^R \omega}{2\omega_0} \hat{A}_{w_2}^\dagger \hat{A}_{w-\mu} \hat{A}_{w_2+\mu} \partial w_2 \partial \mu + i\tilde{\gamma} \int \int \frac{R_\mu^R \omega}{2\omega_0} \hat{A}_{w_1}^\dagger \hat{A}_{w_1-\mu} \hat{A}_{w+\mu} \partial w_1 \partial \mu = \\ & i\tilde{\gamma} \int \int \frac{R_\mu^R \omega}{2\omega_0} \hat{A}_{w'}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'+\mu} \partial w' \partial \mu + i\tilde{\gamma} \int \int \frac{R_\mu^R \omega}{2\omega_0} \hat{A}_{w'}^\dagger \hat{A}_{w+\mu} \hat{A}_{w'-\mu} \partial w' \partial \mu = \\ & i\tilde{\gamma} \int \int \frac{R_\mu^R \omega}{2\omega_0} \hat{A}_{w'}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'+\mu} \partial w' \partial \mu + i\tilde{\gamma} \int \int \frac{R_\nu^R \omega}{2\omega_0} \hat{A}_{w'}^\dagger \hat{A}_{w-\nu} \hat{A}_{w'+\nu} \partial w' \partial \nu = i\tilde{\gamma} \frac{\omega}{\omega_0} \int \int R_\mu^R \hat{A}_{w'}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'+\mu} \partial w' \partial \mu. \end{aligned}$$

where, in the second to the last line, we used the change of variable $\nu = -\mu$ in the second integral and, also, we used the Fourier transform property $R_{-\mu}^R = R_\mu^R$.

$$\frac{\partial}{\partial z} \langle A_w \rangle = \langle i[\hat{A}_w, \hat{H}_{\text{FWM}}] \rangle = i\tilde{\gamma} \frac{\omega}{\omega_0} \int \int R_\mu^R \langle \hat{A}_{w'}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'+\mu} \rangle \partial w' \partial \mu.$$

Losses

$$[\hat{L}_{\alpha w'}^\dagger, \hat{A}_w] \hat{L}_{\alpha w'} = \left[\sqrt{\frac{\alpha_{w'}}{\hbar\omega'}} \hat{A}_{w'}^\dagger, \hat{A}_w \right] \sqrt{\frac{\alpha_{w'}}{\hbar\omega'}} \hat{A}_{w'} = \frac{2\alpha_{w'}}{\hbar\omega'} [\hat{A}_{w'}^\dagger, \hat{A}_w] \hat{A}_{w'} = -2\alpha_{w'} \delta(w - w') \hat{A}_{w'}.$$

$$\hat{L}_{\alpha w'}^\dagger [\hat{L}_{\alpha w'}, \hat{A}_w] = \sqrt{\frac{\alpha_{w'}}{\hbar\omega'}} \hat{A}_{w'}^\dagger \left[\sqrt{\frac{\alpha_{w'}}{\hbar\omega'}} \hat{A}_{w'}, \hat{A}_w \right] = 0.$$

$$\frac{\partial}{\partial z} \langle \hat{A}_w \rangle = \frac{1}{2} \int \langle [\hat{L}_{\alpha w'}^\dagger, \hat{A}_w] \hat{L}_{\alpha w'} \rangle - \langle \hat{L}_{\alpha w'}^\dagger [\hat{L}_{\alpha w'}, \hat{A}_w] \rangle \partial w' = \frac{1}{2} \langle \int -\alpha_{w'} \delta(w - w') \hat{A}_{w'} \partial w' \rangle = -\frac{\alpha_w}{2} \langle \hat{A}_w \rangle.$$

Raman scattering

$$[\hat{L}_{R\mu}^\dagger, \hat{A}_w] = \left[\int \sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \hat{A}_{w'}^\dagger \hat{A}_{w'-\mu} \partial w', \hat{A}_w \right] = \int \sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} [\hat{A}_{w'}^\dagger \hat{A}_{w'-\mu}, \hat{A}_w] \partial w' =$$

$$-\sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \int \hbar\omega \delta(w-w') \hat{A}_{w'-\mu} \partial w' = -\sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \hbar\omega \hat{A}_{w-\mu}.$$

$$[\hat{L}_{R\mu}^\dagger, \hat{A}_w] \hat{L}_{R\mu} = -\sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \hbar\omega \hat{A}_{w-\mu} \int \sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \hat{A}_{w'}^\dagger \hat{A}_{w'-\mu} \hat{A}_{w'} \partial w' = -2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \int \hat{A}_{w-\mu} \hat{A}_{w'-\mu}^\dagger \hat{A}_{w'} \partial w' =$$

$$-2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \int [\hat{A}_{w-\mu}, \hat{A}_{w'-\mu}^\dagger] \hat{A}_{w'} \partial w' - 2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \int \hat{A}_{w'-\mu}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'} \partial w' =$$

$$-2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \left(\int \hbar(\omega-\mu) \delta(w-w') \hat{A}_{w'} \partial w' + \int \hat{A}_{w'-\mu}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'} \partial w' \right) =$$

$$-2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \left(\hbar(\omega-\mu) \hat{A}_w + \int \hat{A}_{w'}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'+\mu} \partial w' \right).$$

$$\int_0^\infty [\hat{L}_{R\mu}^\dagger, \hat{A}_w] \hat{L}_{R\mu} \partial \mu = - \int_0^\infty 2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \left(\hbar(\omega-\mu) \hat{A}_w + \int \hat{A}_{w'}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'+\mu} \partial w' \right) \partial \mu =$$

$$- \int_0^\infty \frac{2\tilde{\gamma}R_\mu^I \hbar\omega(\omega-\mu)}{\omega_0} \hat{A}_w \partial \mu - \int_0^\infty \int 2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \hat{A}_{w'}^\dagger \hat{A}_{w-\mu} \hat{A}_{w'+\mu} \partial w' \partial \mu.$$

$$[\hat{L}_{R\mu}, \hat{A}_w] = \left[\int \sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \hat{A}_{w'-\mu}^\dagger \hat{A}_{w'} \partial w', \hat{A}_w \right] = \int \sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} [\hat{A}_{w'-\mu}^\dagger \hat{A}_{w'}, \hat{A}_w] \partial w' =$$

$$-\sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \int \hbar\omega \delta(w-w'+\mu) \hat{A}_{w'} \partial w' = -\sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \hbar\omega \hat{A}_{w+\mu}.$$

$$\hat{L}_{R\mu}^\dagger [\hat{L}_{R\mu}, \hat{A}_w] = \int \sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \hat{A}_{w'}^\dagger \hat{A}_{w'-\mu} \partial w' \left(-\sqrt{\frac{2\tilde{\gamma}R_\mu^I}{\hbar\omega_0}} \hbar\omega \hat{A}_{w+\mu} \right) = -2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \int \hat{A}_{w'}^\dagger \hat{A}_{w'-\mu} \hat{A}_{w+\mu} \partial w' = -2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \int \hat{A}_{w'}^\dagger \hat{A}_{w+\mu} \hat{A}_{w'-\mu} \partial w'.$$

$$\int_0^\infty \hat{L}_{R\mu}^\dagger [\hat{L}_{R\mu}, \hat{A}_w] \partial \mu = - \int_0^\infty 2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \int \hat{A}_{w'}^\dagger \hat{A}_{w+\mu} \hat{A}_{w'-\mu} \partial w' \partial \mu =$$

$$\int_0^{-\infty} \int 2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \hat{A}_{w'}^\dagger \hat{A}_{w-\nu} \hat{A}_{w'+\nu} \partial w' \partial \nu =$$

$$\int_{-\infty}^0 \int 2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \hat{A}_{w'}^\dagger \hat{A}_{w+\mu} \hat{A}_{w'-\mu} \partial w' \partial \mu.$$

$$\int_0^\infty [\hat{L}_{R\mu}^\dagger, \hat{A}_w] \hat{L}_{R\mu} - \hat{L}_{R\mu}^\dagger [\hat{L}_{R\mu}, \hat{A}_w] \partial \mu = - \iint 2\tilde{\gamma} \frac{\omega}{\omega_0} R_\mu^I \hat{A}_{w'}^\dagger \hat{A}_{w+\mu} \hat{A}_{w'-\mu} \partial w' \partial \mu - \int_0^\infty \frac{2\tilde{\gamma}R_\mu^I \hbar\omega(\omega-\mu)}{\omega_0} \hat{A}_w \partial \mu.$$

$$\frac{\partial}{\partial z} \langle \hat{A}_w \rangle = \frac{1}{2} \int \langle [\hat{L}_{R\mu}^\dagger, \hat{A}_w] \hat{L}_{R\mu} \rangle - \langle \hat{L}_{R\mu}^\dagger [\hat{L}_{R\mu}, \hat{A}_w] \rangle \partial w' = -\tilde{\gamma} \frac{\omega}{\omega_0} \iint R_\mu^I \langle \hat{A}_{w'}^\dagger \hat{A}_{w+\mu} \hat{A}_{w'-\mu} \rangle \partial w' \partial \mu - \tilde{\gamma} \frac{\omega}{\omega_0} \int_0^\infty R_\mu^I \hbar(\omega-\mu) \partial \mu \langle \hat{A}_w \rangle.$$

APPENDIX C

We consider a modified version of Eq. 6 for periodic signals, i.e.,

$$\frac{\partial \rho}{\partial z} = i \left[\hat{H}_{\text{disp}} + \hat{H}_{\text{fwm}}, \rho \right] + \sum_m \hat{L}_{\alpha m} \rho \hat{L}_{\alpha m}^\dagger - \frac{1}{2} \{ \rho, \hat{L}_{\alpha m}^\dagger \hat{L}_{\alpha m} \} \Delta w + \sum_{n=0}^{\infty} \hat{L}_{Rn} \rho \hat{L}_{Rn}^\dagger - \frac{1}{2} \{ \rho, \hat{L}_{Rn}^\dagger \hat{L}_{Rn} \} \Delta w,$$

where

$$\begin{aligned} \hat{H}_{\text{disp}} &= \sum_m \frac{\beta_{w_m}}{\hbar \omega_m} \hat{A}_m^\dagger \hat{A}_m \Delta w, \\ \hat{H}_{\text{fwm}} &= \sum_{m,n,j} \frac{\tilde{\gamma} R_{w_j}^R}{2\hbar \omega_0} \hat{A}_m^\dagger \hat{A}_n^\dagger \hat{A}_{m-j} \hat{A}_{n+j} \Delta w^3, \\ \hat{L}_{\alpha m} &= \sqrt{\frac{\alpha_{w_m}}{\hbar \omega_m}} \hat{A}_m, \\ \hat{L}_{Rn} &= \sum_m \sqrt{\frac{2\tilde{\gamma} R_{w_n}^I}{\hbar \omega_0}} \hat{A}_{m-n}^\dagger \hat{A}_m \Delta w. \end{aligned}$$

We propose the operator

$$\hat{A}_m = \frac{\sqrt{2\pi P}}{\Delta w} e^{ik_p z} \delta_{m0} + \hat{A}_s \delta_{ms} + \hat{A}_i \delta_{mi}.$$

The dispersion hamiltonian reads

$$\begin{aligned} \hat{H}_{\text{disp}} &= \sum_m \frac{\beta_{w_m}}{\hbar \omega_m} \left(\frac{\sqrt{2\pi P}}{\Delta w} e^{-ik_p z} \delta_{m0} + \hat{A}_s^\dagger \delta_{ms} + \hat{A}_i^\dagger \delta_{mi} \right) \left(\frac{\sqrt{2\pi P}}{\Delta w} e^{ik_p z} \delta_{m0} + \hat{A}_s \delta_{ms} + \hat{A}_i \delta_{mi} \right) \Delta w, \\ \hat{H}_{\text{disp}} &= \frac{2\pi P \beta_{w_0}}{\hbar \omega_0} + \frac{\beta_{w_s} \hat{A}_s^\dagger \hat{A}_s}{\hbar \omega_s} \Delta w + \frac{\beta_{w_i} \hat{A}_i^\dagger \hat{A}_i}{\hbar \omega_i} \Delta w. \end{aligned} \quad (27)$$

The first term can be neglected since it is a constant and does not affect the motion equations for \hat{A}_s and \hat{A}_i . Equation 27 can be expressed more succinctly as

$$\hat{H}_{\text{disp}} = \beta_{w_s} \hat{a}_s^\dagger \hat{a}_s + \beta_{w_i} \hat{a}_i^\dagger \hat{a}_i.$$

The FWM hamiltonian reads

$$\begin{aligned} \hat{H}_{\text{fwm}} &= \sum_{m,n,j} \frac{\tilde{\gamma} R_{w_j}^R}{2\hbar \omega_0} \left(A_0^* \delta_{m0} + \hat{A}_s^\dagger \delta_{ms} + \hat{A}_i^\dagger \delta_{mi} \right) \\ &\quad \left(A_0^* \delta_{n0} + \hat{A}_s^\dagger \delta_{ns} + \hat{A}_i^\dagger \delta_{ni} \right) \left(A_0 \delta_{(m-j)0} + \hat{A}_s \delta_{(m-j)s} + \hat{A}_i \delta_{(m-j)i} \right) \\ &\quad \left(A_0 \delta_{(n+j)0} + \hat{A}_s \delta_{(n+j)s} + \hat{A}_i \delta_{(n+j)i} \right) \Delta w^3, \end{aligned}$$

where

$$A_0 = \frac{\sqrt{2\pi P}}{\Delta w} e^{ik_p z}.$$

Just as done with the dispersion hamiltonian, we neglect the constant term. Also, we neglect terms with products of three to four operators \hat{A}_s or \hat{A}_i , since signal or idler optical powers are negligible against pump's powers. Hence, using the relation $R_{-w}^R = R_w^R$ we obtain

$$\begin{aligned} \hat{H}_{\text{fwm}} &= \frac{\tilde{\gamma}}{2\hbar \omega_0} \left(2(1 + R_{w_s}^R) |A_0|^2 (\hat{A}_s^\dagger \hat{A}_s + \hat{A}_i^\dagger \hat{A}_i) + 2R_{w_s}^R (A_0^2 \hat{A}_s^\dagger \hat{A}_i^\dagger + (A_0^*)^2 \hat{A}_s \hat{A}_i) \right) \Delta w^3, \\ \hat{H}_{\text{fwm}} &= \frac{\gamma P}{\hbar \omega_0} \left((1 + R_{w_s}^R) (\hat{A}_s^\dagger \hat{A}_s + \hat{A}_i^\dagger \hat{A}_i) + R_{w_s}^R (e^{2ik_p z} \hat{A}_s^\dagger \hat{A}_i^\dagger + e^{-2ik_p z} \hat{A}_s \hat{A}_i) \right) \Delta w. \end{aligned}$$

Replacing with creation and annihilation operators, and assuming that the frequencies ω_s and ω_i are very close to ω_0 , we obtain

$$\hat{H}_{\text{fwm}} = \gamma P \left((1 + R_{w_s}^R) (\hat{a}_s^\dagger \hat{a}_s + \hat{a}_i^\dagger \hat{a}_i) + R_{w_s}^R (e^{2ik_p z} \hat{a}_s^\dagger \hat{a}_i^\dagger + e^{-2ik_p z} \hat{a}_s \hat{a}_i) \right).$$

We introduce new creation-annihilation operators, defined as

$$\begin{aligned} \hat{b}_s &= e^{-ik_s z} \hat{a}_s, \\ \hat{b}_i &= e^{-ik_i z} \hat{a}_i, \end{aligned}$$

where

$$\begin{aligned} k_s &= \beta_{w_s} + \gamma P (1 + R_{w_s}^R), \\ k_i &= \beta_{w_i} + \gamma P (1 + R_{w_i}^R). \end{aligned}$$

Having in mind the phase-matching condition, $2k_p - k_s - k_i = 0$, the hamiltonian reads

$$\hat{H}_{\text{fwm}} = \gamma P \left((1 + R_{w_s}^R) (\hat{b}_s^\dagger \hat{b}_s + \hat{b}_i^\dagger \hat{b}_i) + R_{w_s}^R (\hat{b}_s^\dagger \hat{b}_i^\dagger + \hat{b}_s \hat{b}_i) \right).$$

This two hamiltonians can be compacted into a unique operator,

$$\hat{H} = \hat{H}_{\text{disp}} + \hat{H}_{\text{fwm}} = k_s \hat{b}_s^\dagger \hat{b}_s + k_i \hat{b}_i^\dagger \hat{b}_i + \gamma P R_{w_s}^R (\hat{b}_s^\dagger \hat{b}_i^\dagger + \hat{b}_s \hat{b}_i).$$

Attenuation operators, only relevant for s -and- i -indexed modes, read

$$\hat{L}_{\alpha s} = \sqrt{\frac{\alpha w_s}{\hbar \omega_s}} \hat{A}_s.$$

$$\hat{L}_{\alpha i} = \sqrt{\frac{\alpha w_i}{\hbar \omega_i}} \hat{A}_i,$$

We use the property

$$\left(\hat{L}_{\alpha m} \rho \hat{L}_{\alpha m}^\dagger - \frac{1}{2} \{ \rho, \hat{L}_{\alpha m}^\dagger \hat{L}_{\alpha m} \} \right) \Delta w = \hat{L}'_{\alpha m} \rho \hat{L}'_{\alpha m}^\dagger - \frac{1}{2} \{ \rho, \hat{L}'_{\alpha m}^\dagger \hat{L}'_{\alpha m} \},$$

where

$$\hat{L}'_{\alpha m} = \sqrt{\alpha w_m} \hat{b}_m.$$

Raman-scattering operators read

$$\hat{L}_{Rn} = \sum_m \sqrt{\frac{2\tilde{\gamma} R_{wn}^I}{\hbar \omega_0}} \left(A_0^* \delta_{(m-n)0} + \hat{A}_s^\dagger \delta_{(m-n)s} + \hat{A}_i^\dagger \delta_{(m-n)i} \right) \left(A_0 \delta_{m0} + \hat{A}_s \delta_{ms} + \hat{A}_i \delta_{mi} \right) \Delta w.$$

We neglect operators with \hat{A}_s^2 , \hat{A}_i^2 or $\hat{A}_s \hat{A}_i$, then we only keep

$$\hat{L}_{Rs} = \sum_m \sqrt{\frac{2\tilde{\gamma} R_{ws}^I}{\hbar \omega_0}} \left(A_0^* \delta_{(m-s)0} + \hat{A}_s^\dagger \delta_{(m-s)s} + \hat{A}_i^\dagger \delta_{(m-s)i} \right) \left(A_0 \delta_{m0} + \hat{A}_s \delta_{ms} + \hat{A}_i \delta_{mi} \right) \Delta w,$$

$$\hat{L}_{R,s} = \sqrt{\frac{2\tilde{\gamma} R_{ws}^I}{\hbar \omega_0}} \left(A_0^* \hat{A}_s + A_0 \hat{A}_i^\dagger \right) \Delta w,$$

$$\hat{L}_{R,s} = \sqrt{\frac{2\gamma P R_{ws}^I}{\hbar \omega_0}} \left(e^{-ik_p z} \hat{A}_s + e^{ik_p z} \hat{A}_i^\dagger \right),$$

It can be easily shown that

$$\left(\hat{L}_{R,s} \rho \hat{L}_{R,s}^\dagger - \frac{1}{2} \{ \rho, \hat{L}_{R,s}^\dagger \hat{L}_{R,s} \} \right) \Delta w = \hat{L}'_{R,s} \rho \hat{L}'_{R,s}^\dagger - \frac{1}{2} \{ \rho, \hat{L}'_{R,s}^\dagger \hat{L}'_{R,s} \},$$

where

$$\hat{L}'_{R,s} = \sqrt{2\gamma P R_{ws}^I} \left(\hat{b}_s + \hat{b}_i^\dagger \right).$$

We assumed that $\omega_s \simeq \omega_i \simeq \omega_0$.

Finally, the master equation for this special case is

$$\frac{\partial \rho}{\partial z} = i[\hat{H}, \rho] + \sum_{\nu=1}^3 \hat{L}_\nu \rho \hat{L}_\nu^\dagger - \frac{1}{2} \{ \rho, \hat{L}_\nu^\dagger \hat{L}_\nu \},$$

with

$$\hat{H} = k_s \hat{b}_s^\dagger \hat{b}_s + k_i \hat{b}_i^\dagger \hat{b}_i + \gamma P R_{ws}^R \left(\hat{b}_s^\dagger \hat{b}_i^\dagger + \hat{b}_s \hat{b}_i \right),$$

$$\hat{L}_1 = \sqrt{\alpha_s} \hat{b}_s,$$

$$\hat{L}_2 = \sqrt{\alpha_i} \hat{b}_i,$$

$$\hat{L}_3 = \sqrt{2\gamma P R_{ws}^I} \left(\hat{b}_s + \hat{b}_i^\dagger \right).$$

-
- [1] Govind P Agrawal. *Nonlinear fiber optics*. Academic press, 2007.
 - [2] Xiaoying Li, Paul L. Voss, Jay E. Sharping, and Prem Kumar. Optical-fiber source of polarization-entangled photons in the 1550 nm telecom band. *Phys. Rev. Lett.*, 94:053601, Feb 2005.
 - [3] HJ McGuinness, MG Raymer, CJ McKinstrie, and S Radic. Quantum frequency translation of single-photon states in a photonic crystal fiber. *Physical review letters*, 105(9):093604, 2010.
 - [4] F. Kaiser, B. Fedrici, A. Zavatta, V. D'Auria, and S. Tanzilli. A fully guided-wave squeezing experiment for fiber quantum networks. *Optica*, 3(4):362–365, Apr 2016.
 - [5] Wang Peng-Xiang, Zhou Qiang, Zhang Wei, Huang Yi-Dong, and Peng Jiang-De. High-quality fiber-based heralded single-photon source at 1.5 μ m. *Chinese Physics Letters*, 29(5):054215, 2012.
 - [6] Robert W. Boyd. *Nonlinear Optics, Third Edition*. Academic Press, Inc., Orlando, FL, USA, 3rd edition, 2008.
 - [7] K. J. Blow, R. Loudon, and S. J. D. Phoenix. Exact solution for quantum self-phase modulation. *J. Opt. Soc. Am. B*, 8(8):1750–1756, Aug 1991.
 - [8] L. Boivin, F. X. Kärtner, and H. A. Haus. Analytical solution to the quantum field theory of self-phase modulation with a finite response time. *Phys. Rev. Lett.*, 2016.

- 73:240–243, Jul 1994.
- [9] Yinchieh Lai and Shinn-Sheng Yu. General quantum theory of nonlinear optical-pulse propagation. *Phys. Rev. A*, 51:817–829, Jan 1995.
- [10] S. J. Carter. Quantum theory of nonlinear fiber optics: Phase-space representations. *Phys. Rev. A*, 51:3274–3301, Apr 1995.
- [11] P. D. Drummond and J. F. Corney. Quantum noise in optical fibers. i. stochastic equations. *J. Opt. Soc. Am. B*, 18(2):139–152, Feb 2001.
- [12] Y. Lai and H. A. Haus. Quantum theory of solitons in optical fibers. i. time-dependent hartree approximation. *Phys. Rev. A*, 40:844–853, Jul 1989.
- [13] Philip Pearle. Simple derivation of the lindblad equation. *European Journal of Physics*, 33(4):805, 2012.
- [14] A. D. Sánchez, S. M. Hernandez, J. Bonetti, P. I. Fierens, and D. F. Grosz. Tunable raman gain in mid-ir waveguides. *J. Opt. Soc. Am. B*, 35(1):95–99, Jan 2018.
- [15] A. D. Sánchez, P. I. Fierens, S. M. Hernandez, J. Bonetti, G. Brambilla, and D. F. Grosz. Anti-stokes raman gain enabled by modulation instability in mid-ir waveguides. *J. Opt. Soc. Am. B*, 35(11):2828–2832, Nov 2018.
- [16] Søren M. M. Friis, Lasse Mejling, and Karsten Rottwitt. Effects of raman scattering and attenuation in silica fiber-based parametric frequency conversion. *Opt. Express*, 25(7):7324–7337, Apr 2017.
- [17] Karsten Rottwitt, Jake Bromage, Andrew J. Stentz, Lufeng Leng, Malcolm E. Lines, and Henrik Smith. Scaling of the raman gain coefficient: Applications to germanosilicate fibers. *J. Lightwave Technol.*, 21(7):1652, Jul 2003.