

Splitting groups with cubic Cayley graphs of connectivity two

Babak Miraftab Konstantinos Stavropoulos

Department of Mathematics
University of Hamburg

February 21, 2022

Abstract

A group G splits over a subgroup C if G is either a free product with amalgamation $A *_C B$ or an HNN-extension $G = A *_C \langle t \rangle$. We invoke Bass-Serre theory to classify all infinite groups which admit cubic Cayley graphs of connectivity two in terms of splittings over a subgroup.

1 Introduction

The study of the structure of groups in terms of the connectivity of their Cayley graphs was started by Jung and Watkins. They characterized infinite transitive graphs of connectivity one whose automorphism groups act on their vertex sets as primitive and regular permutation groups [12]. Later, Watkins [17] characterized all Cayley graphs of connectivity one.

A topic that has been already paired with the connectivity of Cayley graphs in order to study them is the planarity of infinite Cayley graphs. A finitely generated group G is called *planar* if it admits a generating set S such that the Cayley graph $\text{Cay}(G, S)$ is planar. In that case, S is called a *planar generating set*. For the first time, in 1896, Maschke [13] characterized all finite groups admitting planar Cayley graphs. Infinite planar groups attracted more attention, as some of them are related to surface and Fuchsian groups [18, Section 4.10] which play a substantial role in complex analysis, see survey [18]. Hamann [10] uses a combinatorial method in order to show that planar groups are finitely presented. His method is based on tree-decompositions, a crucial tool of graph minor theory which we also utilize extensively in this paper.

In [7], Droms, B. Servatius, and H. Servatius characterized planar groups with low connectivity in terms of the fundamental group of the graph of groups. Indeed, they showed:

Theorem. [7, Theorem 4.4] *If a group G has planar connectivity¹ 2, then either G is a finite cyclic or dihedral group, or it is the fundamental group of a graph of groups whose edge groups all have order two or less and whose vertex groups*

¹The planar connectivity $\kappa(G)$ of a planar group G is the minimum connectivity of all its planar Cayley graphs.

all have planar connectivity at least three. In the latter case, the vertex groups have planar generating sets which include the nontrivial elements of the incident edge groups.

Later, Georgakopoulos in [9] determines all presentations of groups which admit planar cubic Cayley graphs with connectivity two. Furthermore, he provides partial information about the presentations of non-planar ones. Our result provides full information not only for the planar, but also the non-planar groups with cubic Cayley graphs of connectivity two.

More specifically, Georgakopoulos' method does not assert anything regarding (and is, in a sense, independent of) splitting the group over subgroups to obtain its structure. By combining tree-decompositions and Bass-Serre theory, we give a simple proof for the full characterization of all groups with cubic Cayley graphs of connectivity two via the following theorem:

Theorem 1.1. *Let $G = \langle S \rangle$ be a group such that $\Gamma = \text{Cay}(G, S)$ is a cubic graph of connectivity two. Then G is isomorphic to one of the following groups:*

- (i) $\mathbb{Z}_n * \mathbb{Z}_2 = \langle a, b \mid b^2, (ba)^n \rangle$ or $\langle a, b, c \mid a^2, b^2, c^2, (bcb)^n \rangle$,
- (ii) $D_{2n} *_{\mathbb{Z}_2} (t) = \langle a, b \mid b^2, (ba^{-1}ba)^n \rangle$,
- (iii) $D_{2n} *_{\mathbb{Z}_2} D_{2m} = \langle a, b, c \mid a^2, b^2, c^2, (ba)^n, (bc)^m \rangle$ or $\langle a, b, c \mid a^2, b^2, c^2, (bc)^{2n}, (a(bc)^n)^m \rangle$
or $\langle a, b, c \mid a^2, b^2, c^2, (bc)^n, (a(bc)^k b)^m \rangle$,
- (iv) $\mathbb{Z}_{2n} *_{\mathbb{Z}_2} D_{2m} = \langle a, b \mid b^2, a^{2n}, (ba^n)^m \rangle$,
- (v) $D_\infty *_{\mathbb{Z}_2} D_{2m} = \langle a, b, c \mid a^2, b^2, c^2, (a(bc)^n b)^m \rangle$.

Theorem 1.1 is a direct consequence of Theorems 4.3, 4.5, 5.4 and 5.8, where we also discuss in detail the planarity of the corresponding Cayley graphs in each case, as well as their presentations. This allows us to obtain as a corollary the results of [9], as well as full presentations for the non-planar groups with cubic Cayley graphs of connectivity two.

Compared to the methods in [9], we believe that our graph theoretical arguments are simplified, while we inevitably spend more time to recover the full algebraic structure of the group in terms of splitting this time around. Moreover, even though the planar part of our result can be relatively quickly recovered from [9] by applying Tietze transformations accordingly, such an approach usually works only provided that one knows or guesses beforehand the new desired presentation (in our case, the one that expresses the splitting of the group) in order to apply the correct Tietze transformations. By applying Bass-Serre theory, we naturally determine the structure of the group in terms of splitting avoiding the nuisance above, which was also the way we originally obtained it.

2 Preliminaries

Our terminology of groups and graphs is standard. We refer the reader to [15] for Bass-Serre theory and [6] for graph theory for any notation missing.

2.1 Graphs

Throughout this paper, Γ always denotes a connected locally finite graph with vertex set $V(\Gamma)$ and edge set $E(\Gamma)$. A *ray* is a one-way infinite path and a *tail* of a ray is an infinite subpath of the ray. Two rays R_1 and R_2 are equivalent if there is no finite set S of vertices such that R_1 and R_2 have tails in different components of $G \setminus S$. The equivalence classes of rays are called *ends*. We refer the reader to surveys [4, 5] for a detailed study of the end structure of graphs.

For a subset $U \subseteq V(\Gamma)$, we denote by $\Gamma[U]$ the subgraph induced by the vertices of U . A *separation* of Γ is an ordered pair (A, B) , where $A, B \subseteq V(\Gamma)$, such that $\Gamma[A] \cup \Gamma[B] = \Gamma$ and there is no edge between $A \setminus B$ and $B \setminus A$. The *order* of (A, B) is the size of $A \cap B$ and we denote it by $|(A, B)|$. If $|(A, B)| = k$, we say that (A, B) is a *k-separation*. The set of separations of Γ can be equipped with the following partial order: $(A, B) \leq (C, D)$ if $A \subseteq C$ and $B \supseteq D$. We say that (A, B) is *nested* with (C, D) if (A, B) is comparable to either (C, D) or (D, C) . Otherwise, we say that the two separations *cross*. We say that a vertex set X separates vertex sets U and W if there exists a separation (A, B) such that $U \subseteq A$, $W \subseteq B$ and $X = A \cap B$.

Let S be a set of vertices of Γ . The set of neighbors of S is denoted by $N(S) := \bigcup_{s \in S} N(s) \setminus S$, whereas $N[S]$ denotes $S \cup N(S)$. A component C of $G \setminus S$ is called *tight* if $N(C) = S$. A separation (A, B) is called *tight* if both $A \setminus B$ and $B \setminus A$ have tight components. A separation (A, B) *distinguishes* two ends ω_1 and ω_2 if a ray $R_1 \in \omega_1$ has a tail in $A \setminus B$ and a ray $R_2 \in \omega_2$ has a tail in $B \setminus A$ or vice versa. Moreover, it distinguishes ω_1 and ω_2 *efficiently* if there is no separation (C, D) distinguishing ω_1 and ω_2 such that $|(C, D)| < |(A, B)|$. Two ends ω_1 and ω_2 are *k-distinguishable* if there is a separation of order at most k distinguishing ω_1 and ω_2 efficiently.

A separation is *splitting* if it distinguishes at least two ends efficiently. We note that if (A, B) is splitting, then (A, B) is a tight separation. Let (A, B) be a splitting k -separation. The *crossing number* $\text{cn}(A, B)$ of (A, B) is the cardinality of the set containing all crossing tight ℓ -separations distinguishing at least two ends, where $\ell \leq k$ (which can be seen to be finite [3]).

Let Γ be an arbitrary connected graph. A *tree-decomposition* of Γ is a pair (T, \mathcal{V}) of a tree T and a family $\mathcal{V} = (V_t)_{t \in V(T)}$ of vertex sets $V_t \subseteq V(\Gamma)$, which are called *parts*, such that:

- (T1) $V(\Gamma) = \bigcup_{t \in T} V_t$,
- (T2) for every edge $e \in E(\Gamma)$, there exists a $t \in V(T)$ such that both ends of e lie in V_t ,
- (T3) $V_{t_1} \cap V_{t_3} \subseteq V_{t_2}$ whenever t_2 lies on the (t_1, t_3) -path in T .

In order to distinguish them from the vertices of the graph Γ , we will usually refer to the vertices of the underlying tree T of a tree decomposition of Γ as nodes.

An *adhesion set* of (T, \mathcal{V}) is a set of the form $V_t \cap V_{t'}$, where $tt' \in E(T)$. The *adhesion* of (T, \mathcal{V}) is the maximum over the sizes of its adhesion sets. It is not hard to see that each adhesion set leads to a separation of Γ . More precisely, assume that T_t and $T_{t'}$ are the components of $T - tt'$ containing t and t' respectively. Then the adhesion set $V_t \cap V_{t'}$ induces the separation $(W_{t \setminus t'}, W_{t' \setminus t})$ of Γ , where $W_{t \setminus t'} = \bigcup_{s \in T_t} V_s$ and $W_{t' \setminus t} = \bigcup_{s \in T_{t'}} V_s$. When every such separation is

tight, we call the tree-decomposition *tight* as well. Finally, a tree decomposition is *reduced* if no part is contained in another one.

The following folklore fact about tree decompositions and nested set of separations is well-known, see [2].

Remark 2.1. *Every nested set \mathcal{N} of separations gives rise to a reduced tree-decomposition whose adhesion sets are exactly the elements of \mathcal{N} . On the other hand, each adhesion set of a tree-decomposition induces a separation and the set of all induced separations of adhesion sets of a tree-decomposition is a nested set of separations.*

Let Γ be a locally finite graph with a tree-decomposition (T, \mathcal{V}) . We call the *torso* of a part V_t the supergraph of $\Gamma[V_t]$ obtained by adding to it all possible edges in the adhesion sets incident to V_t . The following general lemma for tree-decompositions is folklore.

Lemma 2.2. *Let (T, \mathcal{V}) be a tree-decomposition of a connected graph Γ and $t \in V(T)$ such that every adhesion set of t induces a connected subgraph. Then $\Gamma[V_t]$ is connected. In particular, the torso of every part of (T, \mathcal{V}) is connected.*

In this paper, we are studying groups admitting cubic Cayley graphs of connectivity two.

2.2 Groups

Let G be a group acting on a set X . Then the *setwise stabilizer* of a subset Y of X is the set of all elements $g \in G$ stabilizing Y setwise, i.e

$$\text{St}_G(Y) := \{g \in G \mid gY = Y\}.$$

Let G be a group acting on a graph Γ . Then this action induces an action on $E(\Gamma)$. We say that G acts without *inversion* on Γ if $g(uv) \neq vu$ for all $uv \in E(\Gamma)$ and $g \in G$. In the case that $g(uv) = vu$, we say that g *inverts* u, v . Notice that when G acts transitively with inversion on the set $E(T)$ of edges of a tree T without leaves, it must also act transitively on the set $V(T)$ of its vertices.

Let $G_1 = \langle S_1 \mid R_1 \rangle$ and $G_2 = \langle S_2 \mid R_2 \rangle$ be two groups. Suppose that a subgroup H_1 of G_1 is isomorphic to a subgroup H_2 of G_2 , say via an isomorphic map $\phi: H_1 \rightarrow H_2$. The *free-product with amalgamation* of G_1 and G_2 over H_1 is

$$G_1 *_{{H_1}} G_2 = \langle S_1 \cup S_2 \mid R_1 \cup R_2 \cup h\phi(h)^{-1}, \forall h \in H_1 \rangle.$$

If H_1 and $\phi(H_1)$ are isomorphic subgroups of G_1 , then the HNN-extension of G_1 over H_1 with respect to ϕ is

$$G_1 *_{{H_1}}(t) = \langle S_1, t \mid R_1 \cup tht^{-1}\phi(h)^{-1}, \forall h \in H_1 \rangle.$$

The crux of Bass-Serre theory is captured in the next lemma which determines the structure of groups acting on trees.

Lemma 2.3. [15] *Let G act without inversion on a tree which has no vertices of degree one and let G act transitively on the set of (undirected) edges. If G acts transitively on the vertices of the tree, then G is an HNN-extension of the stabilizer of a vertex over the stabilizer of an edge. If there are two orbits on the vertices of the tree, then G is the free product of the stabilizers of two adjacent vertices with amalgamation over the stabilizer of an edge.*

There is a standard way to deal with the case where we cannot apply Lemma 2.3 directly when G acts with inversion on the tree.

Lemma 2.4. *Let G act transitively with inversion on the edges of a tree T without leaves. Then G is the free product of the stabilizer of a vertex and the stabilizer of an edge with amalgamation over their intersection.*

Proof. Subdivide every edge tt' of T to obtain a tree T' and let $v_{tt'}$ be the corresponding new node. Notice that G now acts transitively on $E(T')$ without inversion and with two orbits on $V(T')$. Each old node t of T has the same pointwise stabilizer in T' . Observe that for each new node $v_{tt'}$ we have $\text{St}_G(v_{tt'}) = \text{St}_G(e)$, where $tt' = e \in E(T)$. The result follows from Lemma 2.3. \square

Finally, \mathbb{Z}_n denotes the *cyclic group of order n* . A finite *dihedral* group is defined by the presentation $\langle a, b \mid b^2, a^n, (ba)^2 \rangle$ and denoted by D_{2n} . Moreover, the infinite dihedral group D_∞ is defined by $\langle a, b \mid b^2, (ba)^2 \rangle$.

3 General structure of the tree-decomposition

Our key tool is the canonical tree decomposition (T, \mathcal{V}) of Lemma 3.3, which will allow us to translate the action of G on Γ to an action of G on T and apply Bass-Serre theory. The lemma follows easily by the following results of [3], which we slightly reformulate for our needs.

Theorem 3.1. [3, Corollary 1.2.] *Let Γ be a locally finite graph with more than one end. For each $\ell \in \mathbb{N}$, there is a canonical tree-decomposition distinguishing all ℓ -distinguishable ends efficiently. Moreover, its adhesion sets induce splitting separations of minimum crossing number.*

As a last preparatory step before we state and prove Lemma 3.3, we need the following result which ensures that our Cayley graph is multi-ended.

Lemma 3.2. [1, Lemma 2.4] *Let Γ be a connected vertex-transitive d -regular graph. Assume Γ has one end. Then the connectivity of Γ is $\geq 3(d+1)/4$.*

For the rest of the paper, we assume that $G = \langle S \rangle$ is an infinite finitely generated group such that $\Gamma = \text{Cay}(G, S)$ is cubic with connectivity two.

The proof idea of the following lemma is similar to [11, Corollary 4.3].

Lemma 3.3. *Let $G = \langle S \rangle$ be an infinite finitely generated group such that $\Gamma = \text{Cay}(G, S)$ is cubic with connectivity two. Then there exists a reduced tree-decomposition (T, \mathcal{V}) with the following properties:*

- (i) *The adhesion sets of (T, \mathcal{V}) have size exactly 2 and induce splitting separations of minimum crossing number.*
- (ii) *The action of G on Γ induces an action on $V(T)$ and a transitive action on the set of separations corresponding to the adhesion sets.*

Proof. Since Γ is an infinite graph of connectivity exactly 2, Lemma 3.2 implies that Γ has at least two ends. Let (T', \mathcal{V}') be a canonical tree-decomposition obtained by Theorem 3.1 which distinguishes all 2-distinguishable ends and since Γ is 2-connected, its adhesion sets have size 2. We consider the set \mathcal{S} of all

induced separations by (T, \mathcal{V}) and choose a separation (A, B) of \mathcal{S} . It follows from [3, Lemma 2.1] that the orbit of (A, B) under the action of G is a nested set \mathcal{N} of separations of order 2. Now by Remark 2.1, the set \mathcal{N} gives rise to a tree-decomposition (T, \mathcal{V}) which has the desired properties. \square

Notice that the transitive action on the set of separations in Lemma 3.3 (ii) implies at most two orbits for $\Gamma[\mathcal{V}] := \{\Gamma[V_t] \mid t \in V(T)\}$ under the action of G . Moreover, we can translate the action of item (ii) to an action of G on T in the natural way (and G will clearly act transitively on $E(T)$):

$$gt = t' \Leftrightarrow gV_t = V_{t'}.$$

Let \mathcal{N} be a nested set of separations of order two in such a way that \mathcal{N} gives a tree-decomposition as in Lemma 3.3. It is easy to see that every 2-separation of Γ such that $A \cap B$ is a proper subset of A and B distinguishes at least two ends, see [8, Lemma 3.4]. For an arbitrary element $(A, B) \in \mathcal{N}$, there are three cases in terms of the degrees of the vertices of the separator in each side of the separation:




Figure 1: The three types of splitting 2-separations in cubic Cayley graphs of connectivity 2.

First, we dismiss the case of TYPE III separations. This follows as an easy corollary of the following lemma.

Lemma 3.4. *There is always a splitting 2-separation of crossing number 0 in Γ .*

Proof. Any TYPE I separation has crossing number 0, as the two vertices of the separator are connected with an edge and thus they are inseparable. Hence we can assume that every splitting 2-separation is not of TYPE I. We show that there is always a splitting 2-separation (A, B) on $A \cap B = \{x, y\}$ such that there are at least two internally disjoint (x, y) -paths in $\Gamma[A]$. This will directly imply that there is no 2-separation (C, D) crossing such an (A, B) as in that case the single vertex in $C \cap D \cap A$ would separate x and y in $\Gamma[A]$, so $\text{cn}(A, B) = 0$.

First, we note that for every 2-separation (A, B) both $A \setminus B, B \setminus A$ are tight components of $G \setminus (A \cap B)$: otherwise, by the 2-connectivity of Γ there are two tight connected components in either $A \setminus B$ or $B \setminus A$, hence there are two internally disjoint (x, y) -paths in $\Gamma[A]$ or $\Gamma[B]$. Then (A, B) or (B, A) is the desired separation.

Now, consider a splitting separation (A, B) on $A \cap B = \{x, y\}$ such that there exist single vertices separating x, y in $\Gamma[A]$ and let S_A be the (non-empty)

set of these cut vertices in $\Gamma[A]$. Let P be a shortest (x, y) -path in $\Gamma[A]$. Then $S_A \subseteq V(P)$. It is easily verified that any two consecutive vertices of $S_A \cup \{x, y\}$ in the ordering inherited by P constitute the separator of a 2-separation (C, D) in Γ nested with (A, B) , and suppose w.l.o.g. that $B \subseteq D$. Since (A, B) is a splitting separation, say distinguishing ω_1, ω_2 , there is a separation (C, D) as above that is also splitting, distinguishing ω_1, ω_2 as well. Then (C, D) is the desired separation as any vertex separating the vertices of $C \cap D$ in $\Gamma[C]$ must also separate x, y in $\Gamma[A]$, contradicting the fact that $S_A \cap (C \setminus D) = \emptyset$. \square

Lemma 3.5. *Any tree decomposition of Γ as in Lemma 3.3 is either of TYPE I or TYPE II.*

Proof. Let (A, B) be a TYPE III separation with $A \cap B = \{x, y\}$. We can assume that $|N(x) \cap A| = 1$ and $|N(x) \cap B| = 2$. Let x' be the unique neighbor of x in A and y' be the unique neighbor of y in B . Then $(A', B') := (A \cup \{y'\} \setminus \{x\}, B \cup \{x'\} \setminus \{y\})$ is a tight TYPE III separation on $\{x', y'\}$, clearly crossing (A, B) and distinguishing efficiently the same ends. The lemma is now a direct consequence of Lemma 3.3 and Lemma 3.4. \square

In what follows, (T, \mathcal{V}) will always be as in Lemma 3.3, either of TYPE I or TYPE II if not explicitly stated otherwise. For a node $t \in V(T)$, we define

$$n(t) := \Gamma \left[\bigcup_{t \in N_T[t]} V_t \right].$$

Recall that every adhesion set $V_t \cap V_{t'}$ of (T, \mathcal{V}) induces the separation $(W_{t \setminus t'}, W_{t' \setminus t})$ of Γ . Assume that (T, \mathcal{V}) and the separations $(W_{t \setminus t'}, W_{t' \setminus t})$ it induces are of TYPE II. We call such a separation $(W_{t \setminus t'}, W_{t' \setminus t})$ *small* if the vertices of the separator $V_t \cap V_{t'}$ have degree 1 in $W_{t' \setminus t}$ and *big* if they have degree 2 in $W_{t' \setminus t}$.

One of our main goals towards the general structure of the tree-decomposition of Γ is to eventually prove in Lemma 3.8 that all adhesion sets of (T, \mathcal{V}) are disjoint. As a preparatory step for that, we need the following Lemma.

Lemma 3.6. *Every vertex u belongs in at least one and at most two different adhesion sets of (T, \mathcal{V}) (as subsets of $V(\Gamma)$ and not as intersections of different pairs of parts). Moreover, for every node t of T and every $t_1, t_2 \in N_T(t)$, we have $|V_{t_1} \cap V_{t_2}| \leq 1$.*

Proof. The lower bound for the first assertion of the lemma follows directly from the transitivity of the actions of G on Γ and $E(T)$. For the upper bound, let $\{x, u\}$ and $\{y, u\}$ be two adhesion sets of the tree-decomposition meeting on u . Since G acts transitively on $E(T)$, there is a $1 \neq g \in G$ such that $g\{x, u\} = \{y, u\}$. Observe that since $g \neq 1$, we must have $gx = u$ and $gu = y$, from which we obtain $ux^{-1}u = y$. Since $\{x, u\}$ and $\{y, u\}$ were arbitrary adhesion sets containing u , the upper bound follows.

For the second assertion, we clearly have that $|V_{t_1} \cap V_{t_2}| \leq 2$. Suppose that $|V_{t_1} \cap V_{t_2}| = 2$. It follows from the definition of a tree-decomposition that $V_{t_1} \cap V_{t_2} \subseteq V_t$ and so $V_{t_1} \cap V_{t_2}$ is a subset of both $V_t \cap V_{t_1}$ and $V_t \cap V_{t_2}$. Therefore, we have $V_{t_1} \cap V_t = V_{t_2} \cap V_t = V_{t_1} \cap V_{t_2} := S$. We observe that the 2-connectivity of Γ implies that all components of $\Gamma \setminus S$ are tight (and in particular, the one containing $V_t \setminus S$).

Let T_S be the subtree of T whose corresponding parts contain S . Then $|V(T_S)| \geq 3$ and assume that $|V(T_S)| \geq 4$. Observe that $\Gamma \setminus S$ then has at least four tight components, which contradicts the fact that Γ is cubic. Hence, $|V(T_S)| = 3$ and so $V(T_S) = \{t, t_1, t_2\}$. Consequently, we see that $C_1 = W_{t_1 \setminus t} \setminus S$, $C_2 = W_{t_2 \setminus t} \setminus S$ and $C_3 = (W_{t \setminus t_1}) \setminus (W_{t_2 \setminus t}) = (W_{t \setminus t_2}) \setminus (W_{t_1 \setminus t})$ must be the components of $G \setminus S$, all of them tight.

This means that both vertices of S must have degree one in each of C_1, C_2, C_3 , respectively, and that S induces an independent set. Since G acts transitively on Γ and $E(T)$ and t was an arbitrary node of t , it follows that every vertex has degree at most one in every part it belongs in. We conclude that every part of \mathcal{V} induces at most a matching where every pair of vertices in the same adhesion set is unmatched. It easily follows that Γ is the disjoint union of two infinite cubic trees, contradicting the fact that Γ is connected. \square

Let H be an arbitrary graph with a set $U \subseteq V(H)$ and a subgraph H' of H . The set U is called *connected in H'* if for every pair of vertices $u, u' \in U$ there is a (u, u') -path in H' .

Lemma 3.7. *Let t be an arbitrary vertex of T . Then for every $t' \in N_T(t)$, the following holds:*

- (i) *The adhesion set $V_t \cap V_{t'}$ is connected in at least one of $V_t, V_{t'}$.*
- (ii) *V_t is connected in $n(t)$.*

Proof. (i) Let $V_t \cap V_{t'} = \{u, u'\}$ and P be a path between u and u' . Suppose that P is contained in $W_{t \setminus t'}$ and consider the tree decomposition (T', \mathcal{V}') of P by restricting (T, \mathcal{V}) on the parts that contain at least two vertices of P . Notice that every adhesion set of (T', \mathcal{V}') has size exactly 2 and root T' on t . Since $T' \subseteq T$, the second assertion of Lemma 3.6 holds for (T', \mathcal{V}') as well, therefore we have that every part of (T', \mathcal{V}') contains at least one new vertex of P compared to its predecessor in the tree-order of T' when rooted on t . Since P is finite, it follows that T' is finite as well. We eventually find a part V_s of (T, \mathcal{V}) — in particular a leaf of T' — such that $P' = V(P) \cap V_s$ is a subpath of P whose end vertices constitute exactly one of the adhesion sets S of V_s . Recall that G acts transitively on the set of adhesion sets of (T, \mathcal{V}) . Hence, we can map S to $V_t \cap V_{t'}$, say $gS = V_t \cap V_{t'}$. Then $gs \in \{t, t'\}$. Thus, gP' is a (u, u') -path that either lies in V_t or $V_{t'}$.

- (ii) Since Γ is connected, the torso of V_t is a connected graph. The result follows by replacing the virtual edges of a path within the torso of V_t by paths obtained by (i). \square

The next crucial lemma implies that all adhesion sets in \mathcal{N} are disjoint. Recall that (T, \mathcal{V}) is a tree-decomposition as in Lemma 3.3.

Lemma 3.8. *Let t be a node of T . Then for every $t_1, t_2 \in N_T(t)$, we have $V_{t_1} \cap V_{t_2} = \emptyset$.*

Proof. By Lemma 3.6, we have that $|V_{t_1} \cap V_{t_2}| \leq 1$. Suppose that $|V_{t_1} \cap V_{t_2}| = 1$. Let $V_{t_1} \cap V_{t_2} = \{x, y\}$, $V_{t_2} \cap V_t = \{x, z\}$. By Lemma 3.6, these are the only adhesion sets of V_t containing x . We can assume that (T, \mathcal{V}) is of TYPE II:

indeed, assume that (T, \mathcal{V}) is of TYPE I. By the tightness of all separations in \mathcal{N} , we have that x has at least one neighbor in each of $V_{t_1} \setminus V_t$ and $V_{t_2} \setminus V_t$ in addition to y and z , a contradiction to Γ being cubic. Hence, (T, \mathcal{V}) is of TYPE II.

Now, by the transitive action on $E(T)$ we have that $(W_{t \setminus t_1}, W_{t_1 \setminus t})$ is either isomorphic to $(W_{t \setminus t_2}, W_{t_2 \setminus t})$ or its inverse (as an ordered separation). In the latter case, assume there are $g \in G$, $t \in V(T)$, such that

$$(W_{t \setminus t_1}, W_{t_1 \setminus t}) = (gW_{t_2 \setminus t}, gW_{t \setminus t_2}). \quad (1)$$

Recall the definition of small and big separations from Page 7. We can assume w.l.o.g. that the above separations are small (a similar situation arises in case they are big separations). We observe that this implies that $\deg_{W_{t_1 \setminus t}}(x) = 1$, $\deg_{W_{t_2 \setminus t}}(x) = 2$ and the degree of x in the component of $\Gamma \setminus \{x, y, z\}$ containing $V_t \setminus \{x, y, z\}$ is 0. By Lemma 3.7, there is an (x, y) -path P lying completely within V_{t_1} , but by (1) we have that the (x, y) -path gP lies within V_t , which yields a contradiction to the degree of x .

Otherwise, for every $t \in V(T)$ there is a $g \in G$ such that

$$(W_{t \setminus t_1}, W_{t_1 \setminus t}) = (gW_{t \setminus t_2}, gW_{t_2 \setminus t}).$$

Let C'_1, C'_2, C'_3 be the components of $\Gamma \setminus \{x, y, z\}$ corresponding to V_{t_1}, V_{t_2}, V_t . Then we directly observe that both separations $(W_{t \setminus t_1}, W_{t_1 \setminus t})$, $(W_{t \setminus t_2}, W_{t_2 \setminus t})$ must be small. The only way for this to happen is when x has degree one in each of C_1, C_2, C_3 and therefore degree at most one in V_{t_1}, V_{t_2}, V_t (in other words, every part that contains it). By the transitivity of Γ and the fact that t was arbitrary, we conclude that every vertex of Γ has degree at most one in every part that contains it, to obtain a contradiction to Lemma 3.7 exactly as before. \square

Lemma 3.8 has some important consequences. Combined with Lemma 3.6, we immediately obtain the following.

Corollary 3.9. *Every vertex u of Γ is contained in exactly two parts $t, t' \in V(T)$. In addition, $N_\Gamma(u) \subseteq V_t \cup V_{t'}$ and every part is the disjoint union of its adhesion sets.* \square

Moreover, let $\{x, y\}$ be an adhesion set. Observe that $xy^{-1}\{x, y\}$ is again an adhesion set containing x , so $xy^{-1}\{x, y\} = \{x, y\}$ with $xy^{-1}x = y$. We obtain:

Lemma 3.10. *For every adhesion set $\{x, y\}$, we have $(xy^{-1})^2 = 1$.* \square

Lemma 3.10 implies the following Corollary for the edge stabilizers of T .

Corollary 3.11. *Let $tt' \in E(T)$. Then $\text{St}_G(V_t \cap V_{t'}) \cong \mathbb{Z}_2$.* \square

Lastly, we will invoke the following folklore Lemma from the well-known theory of tree decompositions into 3-connected components (see [14, 16] as an example) when we argue about the planarity of Γ and G in each case that arises.

Lemma 3.12. *Let (T, \mathcal{V}) be a tight tree-decomposition of a (locally finite) connected graph H with finite parts and adhesion at most 2. Then Γ is planar if and only if the torso of every part of (T, \mathcal{V}) is planar.*

Proof. The forward implication follows from the fact that the torso of a part in (T, \mathcal{V}) is a topological minor of H : for every virtual edge of the part realized by an adhesion set of size exactly two, there is always a path outside of the part that connects the two vertices of the adhesion set.

For the backward implication, we will inductively embed the planar torsos of the parts of (T, \mathcal{V}) on the plane by gluing (the torso of) each new bag at the appropriate adhesion set. For the inductive step, we consider the adhesion set $V_t \cap V_{t'} = \{x, y\}$ of the new bag V_t along which it is going to be amalgamated with the already embedded torso of $V_{t'}$. We simply replace the edge xy with a planar embedding of the torso of V_t (restricted inside a thin disk around the embedded xy), keeping the edge xy in the so far embedded part of Γ depending on whether it is an actual edge of Γ or a virtual edge of the torsos of $V_t, V_{t'}$, accordingly. It is straightforward to check that the above inductive strategy to combine the planar embeddings of the torsos along the adhesion sets produces an embedding of Γ on the plane, since each virtual edge of the torsos is replaced with a new bag at some inductive step. \square

Our goal in the following sections is to determine the structure of the parts of the tree-decomposition of Γ obtained by Corollary 3.3 in order to compute their stabilizers and apply Lemma 2.3 or 2.4.

4 Tree-decomposition of TYPE I

In this section, we assume that (T, \mathcal{V}) is of TYPE I. Suppose that b is the label of the edge induced by the adhesion sets of (T, \mathcal{V}) , which by Lemma 3.10 is an involution. It will be enough to study two neighboring parts $V_t, V_{t'}$ to obtain the general structure of (T, \mathcal{V}) . In order to simplify this, we can assume w.l.o.g that $V_t \cap V_{t'} = \{1, b\}$, so $\text{St}_G(V_t \cap V_{t'}) = \langle b \rangle$.

Notice that if G acts on (T, \mathcal{V}) with inversion, there is an element in $g \in \text{St}_G(V_t \cap V_{t'}) = \langle b \rangle$ that inverts $V_t, V_{t'}$. Let us express this easy fact with the following lemma.

Lemma 4.1. *G acts with inversion on (T, \mathcal{V}) if and only if b inverts V_t and $V_{t'}$.* \square

Lemma 4.2. *Every part of \mathcal{V} induces a finite cycle.*

Proof. Let $t \in V(T)$. Since every adhesion set induces a connected subgraph, we conclude by Lemma 2.2 that $\Gamma[V_t]$ is connected. Moreover, Corollary 3.9 implies that $\Gamma[V_t]$ is 2-regular. It follows that $\Gamma[V_t]$ is either a finite cycle or a double ray. Recall that by Lemma 3.8 all adhesion sets are disjoint. The conclusion follows by observing that every vertex of V_t is a cut vertex when V_t induces a double ray and hence, the graph Γ is not 2-connected. \square

It will be clear by Lemma 3.12 that we will obtain planar Cayley graphs in all subcases.

4.1 Two Generators

Assume that $G = \langle a, b \rangle$, where b is an involution. We distinguish the following cases depending on the colors of the edges incident to the adhesion sets, depicted as Figure 2.




Figure 2: Cases of TYPE I with two generators

4.1.1 Case I

Suppose that the edges incident to each adhesion set inducing a separation in \mathcal{N} are as in Case I of Figure 2. Observe that $\{a^{-1}, ba\} \subseteq V_t$ and $\{a, ba^{-1}\} \subseteq V_{t'}$ are the neighbors of 1 and b in V_t and $V_{t'}$, respectively. Since $b\{a^{-1}, ba\} = \{a, ba^{-1}\}$, it must be that $bV_t = V_{t'}$ and $bV_{t'} = V_t$. Lemma 4.1 implies that G acts on $E(T)$ with inversion (and hence transitively on $V(T)$).

By Lemma 4.2, there is an $n \in \mathbb{N}$ such that $(ba)^n = 1$ and

$$V_t = \{1, b, ba, \dots, (ba)^{n-1}b = a^{-1}\}.$$

This gives a partition $\langle ba \rangle \sqcup \langle ba \rangle b$ of V_t . We next conclude that $\text{St}_G(V_t) \subseteq V_t$ by noting that $1 \in V_t$. Clearly, we have $\langle ba \rangle \subseteq \text{St}_G(V_t)$. Moreover, for the element $ba \in V_t$, we observe that

$$(ba)^i b(ba) = (ba)^i a \notin V_t.$$

Since $V_t = \langle ba \rangle \sqcup \langle ba \rangle b$, we conclude that $\text{St}_G(V_t) = \langle ba \rangle \cong \mathbb{Z}_n$. Moreover, $\text{St}_G(V_t) \cap \text{St}_G(V_t \cap V_{t'}) = \langle ba \rangle \cap \langle b \rangle = 1$.

We apply Lemma 2.4 and obtain that

$$G \cong \mathbb{Z}_n * \mathbb{Z}_2.$$

4.1.2 Case II

By the structure of the neighbourhood of $\{1, b\}$ and Lemma 4.1 we see that b cannot invert V_t and $V_{t'}$, hence G acts on (T, \mathcal{V}) without inversion.

Now, consider the adhesion set $a^{-1}\{1, b\} = (a^{-1}V_t) \cap (a^{-1}V_{t'})$. From the fact that $a^{-1}\{1, b\} \subseteq V_t$ we deduce that either $a^{-1}V_t = V_t$ or $a^{-1}V_{t'} = V_t$. Since the adhesion set $\{1, b\}$ has ingoing a -edges but $a\{1, b\}$ has outgoing a -edges in V_t , we cannot have that $a^{-1}V_t = V_t$. Consequently, it must be that $a^{-1}V_{t'} = V_t$. The fact that two adjacent parts lie in the same orbit under the action of G implies that G acts transitively on \mathcal{V} (and $V(T)$).

By Lemma 4.2, there is in this case an $n \in \mathbb{N}$ such that $(ba^{-1}ba)^n = 1$ and

$$V_t = \{1, b, ba^{-1}, ba^{-1}b, \dots, (ba^{-1}ba)^{n-1}ba^{-1}b = a^{-1}\}.$$

In other words, $\langle ba^{-1}ba \rangle \sqcup \langle ba^{-1}ba \rangle b \sqcup \langle ba^{-1}ba \rangle ba^{-1} \sqcup \langle ba^{-1}ba \rangle ba^{-1}b$ forms a partition of V_t . Notice that $\langle ba^{-1}ba \rangle$ is the trivial group when $ba^{-1}ba = 1$. As before, since $1 \in V_t$ we infer that $\text{St}_G(V_t) \subseteq V_t$. Clearly, we have

$\langle ba^{-1}ba \rangle \subseteq \text{St}_G(V_t)$. Moreover, we see that $\langle ba^{-1}ba \rangle ba^{-1} \not\subseteq \text{St}_G(V_t)$ because we have $(ba^{-1}ba)^i ba^{-1}(ba^{-1}ba) \notin V_t$ and that $\langle ba^{-1}ba \rangle ba^{-1}a \not\subseteq \text{St}_G(V_t)$ because $(ba^{-1}ba)^i ba^{-1}b(a^{-1}ba) \notin V_t$.

Lastly, observe that since b is an involution and all adhesion sets induce a b -edge, we have that the action of b on Γ fixes every adhesion set. Hence, we have that $b \in \text{St}_G(V_t)$. It follows that $\langle ba^{-1}ba, b \rangle \subseteq \text{St}_G(V_t)$. Therefore, we conclude that

$$\text{St}_G(V_t) = \langle ba^{-1}ba, b \mid b^2, (ba^{-1}ba)^n, (a^{-1}ba)^2 \rangle \cong D_{2n}.$$

By Lemma 2.3, we have that

$$G \cong D_{2n} *_{\mathbb{Z}_2} (t).$$

We collect both cases in the following theorem.

Theorem 4.3. *If (T, \mathcal{V}) is of TYPE I with two generators, then G satisfies one of the following cases:*

(i) $G \cong \mathbb{Z}_n * \mathbb{Z}_2$.

(ii) $G \cong D_{2n} *_{\mathbb{Z}_2} (t)$. □

The definitions of a free product with amalgamation, an HNN-extension and the proof of Theorem 4.3 immediately imply:

Corollary 4.4. [9, Theorem 1.1] *If (T, \mathcal{V}) is of TYPE I with two generators, then G has one of the following presentations:*

(i) $G = \langle a, b \mid b^2, (ba)^n \rangle$.

(ii) $G = \langle a, b \mid b^2, (ba^{-1}ba)^n \rangle$. □

4.2 Three Generators

Let $G = \langle a, b, c \rangle$, where a, b and c are involutions. Suppose that the edges induced by the adhesion sets with corresponding separations in \mathcal{N} are colored with b . Up to rearranging a, b, c , there are two cases for the local structure of the adhesion sets with separations in \mathcal{N} , as in the following figure:




Figure 3: Cases of TYPE I with three generators

4.2.1 Case I

First, we observe by Lemma 4.1 that G acts on T without inversion, since by the structure of the neighbourhood of $\{1, b\}$ we see that b must stabilize both V_t and $V_{t'}$. Consequently, G must act with two orbits O_1, O_2 on $\Gamma[\mathcal{V}]$, where the parts in O_1 contain the a -edges and the parts in O_2 contain the c -edges. By Lemma 4.2 we deduce that there exist n, m with $(ba)^n = 1$ and $(bc)^m = 1$ and so $V_t = \langle ba \rangle \sqcup \langle ba \rangle b$ and $V_{t'} = \langle bc \rangle \sqcup \langle bc \rangle b$.

To compute the stabilizers of the parts, observe that we can escape a part in O_1 only with c -edges. Hence, we have $\text{St}_G(V_t) = V_t = \langle ba, b \mid b^2, (ba)^n, a^2 \rangle \cong D_{2n}$ and similarly $\text{St}_G(V_{t'}) = V_{t'} = \langle bc, b \mid b^2, (bc)^m, c^2 \rangle \cong D_{2m}$. Therefore, by Lemma 2.3 we obtain

$$G \cong D_{2n} \underset{\mathbb{Z}_2}{*} D_{2m}.$$

4.2.2 Case II

In this case, we see that b inverts V_t and $V_{t'}$, so G acts on T with inversion by Lemma 4.1. Hence, G also acts transitively on $V(T)$.

Let $x := bcba$. By Lemma 4.2 we see that $(bcba)^n = 1$ and that $\langle x \rangle \sqcup \langle x \rangle b \sqcup \langle x \rangle bc \sqcup \langle x \rangle bcb$ is a partition of V_t . Clearly, we have that $\langle bcba \rangle \subseteq \text{St}_G(V_t)$. We show that we actually have equality:

- $x^i b \cdot bc = x^i c \notin V_t$, hence $\langle x \rangle b \notin \text{St}_G(V_t)$,
- $x^i bc \cdot a \notin V_t$, hence $\langle x \rangle bc \notin \text{St}_G(V_t)$,
- $x^i bcb \cdot c \notin V_t$, hence $\langle x \rangle bcb \notin \text{St}_G(V_t)$.

We conclude that $\text{St}_G(t) = \langle bcba \rangle \cong \mathbb{Z}_n$ and consequently we also have that $\text{St}_G(V_t) \cap \text{St}_G(V_t \cap V_{t'}) = \langle bcba \rangle \cap \langle b \rangle = 1$. It follows from Lemma 2.4 that

$$G \cong \mathbb{Z}_n * \mathbb{Z}_2.$$

In conclusion, we have proved:

Theorem 4.5. *If (T, \mathcal{V}) is of TYPE I with three generators, then G satisfies one of the following cases:*

(i) $G \cong D_{2n} \underset{\mathbb{Z}_2}{*} D_{2m}.$

(ii) $G \cong \mathbb{Z}_n * \mathbb{Z}_2.$

□

Corollary 4.6. [9, Theorem 1.1] *If (T, \mathcal{V}) is of TYPE I with three generators, then G has one of the following presentations:*

(i) $G = \langle a, b, c \mid a^2, b^2, c^2, (ba)^n, (bc)^m \rangle.$

(ii) $G = \langle a, b, c \mid a^2, b^2, c^2, (bcba)^n \rangle.$

□

5 Tree-decomposition of TYPE II

Even though at first glance there can be several cases for TYPE II separations, we will in fact be able to quickly exclude most of them using appropriately the following lemma.

Lemma 5.1. *Let $G = \langle a, b, c \rangle$ (with possibly $c = a^{-1}$), where b is an involution and let $\{x, y\}$ be the adhesion set of a TYPE II separation in (T, \mathcal{V}) of Γ as in Lemma 3.3. Let v_1, v_2, v_3 be any consecutive vertices in a shortest (x, y) -path P with at least two edges and suppose there is $g \in G$ such that $gv_2 \in \{x, y\}$. Then gv_1 and gv_3 lie in the same component of $\Gamma \setminus \{x, y\}$.*

Proof. Suppose not. We observe that gx, gy must then lie in different components of $\Gamma \setminus \{x, y\}$ as well: if not, then gx, gy lie in the same component, say, C . The fact that gv_1, gv_3 lie in different components implies that gP leaves C , therefore we have that both $x, y \in V(gP)$. Since $gv_2 \in \{x, y\}$ is an inner vertex of gP , the subpath of gP from x to y contradicts the choice of P .

Hence, $g\{x, y\}$ is a separator where gx, gy lie in different components of $\Gamma \setminus \{x, y\}$. It easily follows that $\{x, y\}$ and $\{gx, gy\}$ are not nested, a contradiction to Lemma 3.1. \square

Now, let $W_{2n+1, 2k}$, where $n \geq 1, k \geq 3, n \leq k$, denote the cubic graph obtained by the $2k$ -cycle with vertices $\{0, 1, \dots, 2k-1\}$ along with “chord” edges of the form $\{2i, 2i + 2n + 1\} \pmod{2k}$ forming a matching. Moreover, we define $V_{2n}, n \geq 2$ as the cubic graph obtained by the $2n$ -cycle along with the “diagonal” edges $\{i, i + n\} \pmod{2n}$. We note that for $k = 2n + 1$ we have $W_{2n+1, 4n+2} = V_{4n+2}$ (Fig. 4).




Figure 4: The graphs $W_{5,10} = V_{10}$ and $W_{5,8}$.

Lastly, let R_{2m+1} be the cubic graph obtained by a double ray with vertex set \mathbb{Z} (defined in the natural way) and by adding the edges of the form $\{2i, 2i + 2m + 1\}$ (Fig. 5).




Figure 5: The graph R_5 .

We note that we will see in the next subsections that the tree-decomposition of Γ obtained by Lemma 3.3 will have two orbits of parts and that the torsos of the parts of one of the two orbits will always be isomorphic to either $W_{2n+1,2k}$, V_{2n} or R_{2n+1} . The fact that $W_{2n+1,2k}$ is never planar, whereas V_{2n} and R_{2m+1} are planar if and only if $n = 2$ and $m = 1$, respectively, will allow us by Lemma 3.12 to determine exactly when Γ will be planar.

5.1 Two generators

Let $G = \langle a, b \rangle$, where b is an involution. Let (T, \mathcal{V}) the corresponding tree-decomposition obtained by Lemma 3.3 and \mathcal{N} the set of separations obtained from its adhesion sets. Then we have the following cases for the neighborhood of such an adhesion set $\{x, y\}$:




Figure 6: Cases of TYPE II with two generators.

Lemma 5.2. *The adhesion sets of (T, \mathcal{V}) satisfy Case III.*

Proof. Let $\{x, y\}$ be an adhesion set. First, observe that no path in Γ contains two consecutive b -edges, hence every path of length two contains at least one a -edge. Let P be a shortest (x, y) -path², necessarily of length at least two.

Assume that either Case I or Case II happen. Notice that –in both cases– for every possible edge-coloring of a path of length two there exists a path Q of length two whose middle vertex belongs in $\{x, y\}$ and its two endpoints lie in different components of $\Gamma \setminus \{x, y\}$ that realizes the same edge-coloring. Consider an arbitrary subpath $P' = v_1 v_2 v_3$ of P of length two and an appropriate Q as above that realizes the edge-coloring of P' . Let w be the middle vertex of Q and $g = wv_2^{-1}$. Then $gP = Q$ and gv_1, gv_3 lie in different components of $\Gamma \setminus \{x, y\}$, contradicting Lemma 5.1. \square

Consequently, we can assume for the rest of this subsection that only Case III happens. It follows that no part of (T, \mathcal{V}) contains edges of all colors: otherwise, by Corollary 3.9 and the fact that no adhesion set contains both a vertex incident

²By Lemma 3.7(i) we can see that P lies completely within V_t or $V_{t'}$, but this is irrelevant to the proof of the Lemma.

with a -edges as well as a vertex incident with b -edges in a part V_t , we see that the a -edges and the b -edges induce different connected components in the torso of V_t , a contradiction to the connectivity of Γ .

Hence, (T, \mathcal{V}) has two orbits of parts O_1, O_2 , where parts in O_1 contain only edges colored with a and parts in O_2 contain edges colored with b . Moreover, G acts on (T, \mathcal{V}) without inversion. The structure of the parts in O_2 is clear: their edges induce a perfect b -matching in the part. We are ready to obtain the full structure of the parts in O_1 as well.

Lemma 5.3. *There is an $n \geq 2$, such that for every adhesion set $\{x, y\}$ we have $x = ya^n$ or $x = ya^{-n}$. Moreover, every part in O_1 induces an a -cycle of length $2n$.*

Proof. Let $V_t \in O_1$ and $\{x, y\} = V_t \cap V_{t'}$ be an adhesion set of t . For every $s \in N_T(t)$, we have that $V_s \in O_2$ and consequently that V_s induces a b -matching. By Lemma 3.7(ii), it follows that $\Gamma[V_t]$ is connected.

Consider an (x, y) -path P within V_t and let $n \geq 2$ be its length. Hence, $x = ya^n$ or $x = ya^{-n}$. By Lemma 3.10, we have $(xy^{-1})^2 = 1$, from which we obtain $a^{2n} = 1$ after substituting x .

We have inferred that the 2-regular graph $\Gamma[V_t]$ is connected. Notice that $P, xy^{-1}P$ are internally disjoint paths with the same ends living inside V_t , therefore their concatenation induces a cycle in V_t . Recall that a has order $2n$. This directly implies the Lemma. \square

Observe that the torso of a part $V_s \in O_2$ induces a connected, 2-regular graph. It cannot be a double ray: in that case every vertex is a cut vertex (as is easily seen), which violates the 2-connectivity of Γ . Hence, the torso of V_s induces a finite cycle, whose edges we can label by Lemma 5.3 with a^n (corresponding to the virtual edges of the torso) and b in an alternating fashion. Therefore, there is an $m \geq 2$ such that $(ba^n)^m = 1$.

It remains to compute the vertex stabilizers of T .

Let $V_{t_1} \in O_1$ such that $1 \in V_{t_1}$. By Lemma 5.3, we clearly have $\langle a \rangle = V_{t_1}$ and therefore $\text{St}_G(V_{t_1}) = \langle a \rangle \cong \mathbb{Z}_{2n}$. Next, let $V_{t_2} \in O_2$ such that $1 \in V_{t_2}$. Recall that $(ba^n)^m = 1$ and notice that $(b(ba^n))^2 = a^{2n} = 1$. By the structure of the torso of V_{t_2} , we observe that the elements of V_{t_2} form a group generated by b and ba^n with presentation $\langle ba^n, b \mid ((ba^n)^m, b^2, (b(ba^n))^2) \rangle$. Since V_{t_2} forms a subgroup of G , we deduce that

$$\text{St}_G(V_{t_2}) = V_{t_2} = \langle ba^n, b \mid ((ba^n)^m, b^2, (b(ba^n))^2) \rangle \cong D_{2m}.$$

Finally, by Lemma 2.3 we obtain $G \cong \mathbb{Z}_{2n} \underset{\mathbb{Z}_2}{*} D_{2m}$.

We observe that the torso of V_{t_1} is isomorphic to V_{2n} . Since V_{2n} is planar if and only if $n = 2$, we conclude by Lemma 3.12 that Γ is planar if and only if $n = 2$. We have obtained the following theorem, along with its corollary by the definition of a free product with amalgamation:

Theorem 5.4. *If (T, \mathcal{V}) is of TYPE II with two generators, then*

$$G \cong \mathbb{Z}_{2n} \underset{\mathbb{Z}_2}{*} D_{2m}.$$

In particular, G is planar if and only if $n = 2$.

Corollary 5.5. [9, Theorem 1.1] *If (T, \mathcal{V}) is of TYPE II with two generators, then*

$$G = \langle a, b \mid b^2, a^{2n}, (ba^n)^m \rangle.$$

In particular, G is planar if and only if $n = 2$.

5.2 Three generators

Let $G = \langle a, b, c \rangle$, where a, b and c are involutions. Then –up to rearranging a, b, c – we have the following cases for the separations in \mathcal{N} :




Figure 7: TYPE II cases with three generators

As in Subsection 5.1, by properly applying Lemma 5.1 we obtain the analogue of Lemma 5.2 for three generators with exactly the same proof.

Lemma 5.6. *The adhesion sets of (T, \mathcal{V}) satisfy Case II.* □

Since the torso of every part of (T, \mathcal{V}) is a connected graph, we deduce that the tree-decomposition has two orbits of parts: parts in O_1 contain only b - and c -edges and parts in O_2 induce perfect a -matchings. Clearly, G then acts on (T, \mathcal{V}) without inversion. Let us quickly obtain the analogue of Lemma 5.3.

Lemma 5.7. *Every part in O_1 induces an alternating (b, c) -cycle of even length or an alternating double (b, c) -ray.*

Proof. Let $V_t \in O_1$ and $\{x, y\} = V_t \cap V_{t'}$ be an adhesion set of t . Since all neighbours of t induce an a -matching, it follows by Lemma 3.7(ii) that $\Gamma[V_t]$ is connected.

Hence, there exists an (x, y) -path P of length i within V_t , necessarily alternating with b - and c -edges. Then, either $x = y(bc)^n$ or $x = y(bc)^nb$, up to swapping b and c . To obtain the structure of the 2-regular, connected graph V_t we distinguish the following cases.

- (i) If $x = y(bc)^n$, then the (x, y) -path $xy^{-1}P$ intersects P only in x, y and by Lemma 3.10, we obtain $(bc)^{2n} = 1$. In this case, V_t induces an alternating (b, c) -cycle of length $4n$.

If $x = y(bc)^nb$, then $xy^{-1}P = P$ and:

- (ii) either V_t induces an alternating (b, c) -cycle,
- (iii) or V_t induces an alternating double (b, c) -ray. □

By the 2-connectivity of Γ , the connected, 2-regular torso of a part $V_s \in O_2$ must be a finite cycle. Depending on which of the cases of Lemma 5.7 we have, we can label its edges with $(bc)^n$ or $(bc)^nb$ (corresponding to the virtual edges of the torso) and a in an alternating fashion. Therefore, there is an $m \geq 2$ such that $(a(bc)^n)^m = 1$ or $(a(bc)^nb)^m = 1$. It remains to infer the structure of G in each case.

- (i) Assume that every part in O_1 is an alternating (b, c) -cycle of length $4n$ and $(a(bc)^n)^m = 1$.

In order to compute the vertex stabilizers of T , let $V_{t_1} \in O_1$ with $1 \in V_{t_1}$. Since $(b(bc))^2 = c^2 = 1$, we have that

$$V_{t_1} = \langle bc \rangle \cup \langle bc \rangle b = \langle bc, b \mid (bc)^{2n}, b^2, (b(bc))^2 \rangle \cong D_{4n}.$$

Then $\text{St}_G(V_{t_1}) = V_{t_1} \cong D_{4n}$, as V_{t_1} forms a group. Next, let $V_{t_2} \in O_2$ with $1 \in V_{t_2}$. Notice that $(a(bc)^n)^m = a^2 = 1$ and $(a(a(bc)^n))^2 = (bc)^{2n} = 1$. We can deduce that V_{t_2} is a group (and hence $\text{St}_G(V_{t_2}) = V_{t_2}$), along with its presentation:

$$\text{St}_G(V_{t_2}) = V_{t_2} = \langle a(bc)^n, a \mid (a(bc)^n)^m, a^2, (a(a(bc)^n))^2 \rangle \cong D_{2m}.$$

By Lemma 2.3, we have

$$G \cong D_{4n} *_{\mathbb{Z}_2} D_{2m}.$$

In this case, the torso of V_{t_1} is isomorphic to V_{4n} , which is planar if and only if $n = 1$.

- (ii) Suppose that parts in O_1 induce an alternating (b, c) -cycle of even length and that $(a(bc)^nb)^m = 1$.

Let $V_{t_1} \in O_1$ and $V_{t_2} \in O_2$ both containing 1 in the respective parts. We see that

$$\text{St}_G(V_{t_1}) = V_{t_1} = \langle bc, b \mid (bc)^k, b^2, (b(bc))^2 \rangle \cong D_{2k},$$

$$\text{St}_G(V_{t_2}) = V_{t_2} = \langle a(bc)^nb, a \mid (a(bc)^nb)^m, a^2, (a(a(bc)^nb))^2 \rangle \cong D_{2m}.$$

By Lemma 2.3,

$$G \cong D_{2k} *_{\mathbb{Z}_2} D_{2m},$$

Notice that the torso of V_{t_1} is isomorphic to $W_{2n+1, 2k}$, which is not planar.

- (iii) In this case, every part in O_1 is an alternating double (b, c) -ray and $(a(bc)^nb)^m = 1$.

Let $V_{t_1} \in O_1$ and $V_{t_2} \in O_2$ with 1 contained in their common adhesion set. Similarly, we have that

$$\text{St}_G(V_{t_1}) = V_{t_1} = \langle bc, b \mid b^2, (b(bc))^2 \rangle \cong D_\infty,$$

$$\text{St}_G(V_{t_2}) = V_{t_2} = \langle a(bc)^nb, a \mid (a(bc)^nb)^m, a^2, (a(a(bc)^nb))^2 \rangle \cong D_{2m}.$$

By Lemma 2.3, we have

$$G \cong D_\infty \underset{\mathbb{Z}_2}{*} D_{2m},$$

to conclude that the torso of V_{t_1} is isomorphic to R_{2n+1} , which is planar if and only if $n = 1$.

By Lemma 3.12 and the above discussion, we have deduced:

Theorem 5.8. *If (T, \mathcal{V}) is of TYPE II with three generators, then G satisfies one of the following cases:*

- (i) $G \cong D_{4n} \underset{\mathbb{Z}_2}{*} D_{2m}$.
- (ii) $G \cong D_{2k} \underset{\mathbb{Z}_2}{*} D_{2m}$
- (iii) $G \cong D_\infty \underset{\mathbb{Z}_2}{*} D_{2m}$. □

Corollary 5.9. [9, Theorem 1.1] *If (T, \mathcal{V}) is of TYPE II with three generators, then G has one of the following presentations:*

- (i) $G = \langle a, b, c \mid a^2, b^2, c^2, (bc)^{2n}, (a(bc)^n)^m \rangle$ and Γ is planar if and only if $n = 1$.
- (ii) $G = \langle a, b, c \mid a^2, b^2, c^2, (bc)^k, (a(bc)^n b)^m \rangle$. Γ is not planar.
- (iii) $G = \langle a, b, c \mid a^2, b^2, c^2, (a(bc)^n b)^m \rangle$ and Γ is planar if and only if $n = 1$. □

6 Open Questions

Having obtained the full characterization of groups admitting cubic Cayley graphs of connectivity two, some further open questions can naturally be raised. In light of Lemma 3.2, we can ask the following.

Problem 1. *Characterize all groups admitting 4-regular Cayley graphs of connectivity at most three in terms of splitting over subgroups.*

Let \mathcal{G} be the family of graphs containing all cycles and all graphs of the form $W_{2n+1,k}$, V_{2n} or R_{2m+1} . A graph is called quasi-transitive if it has a finite number of orbits under the action of its automorphism group. Looking back at Theorem 1.1, we see that cubic Cayley graphs of connectivity two can be expressed as a tree decomposition whose torsos induce graphs from \mathcal{G} . The main tools from our proof seem to go through to support that this is in general the case for every cubic transitive graph of connectivity two. We can go a step further and ask the following question:

Problem 2. *Characterize all cubic quasi-transitive graphs of connectivity two in terms of “canonical” tree decompositions with the property that the automorphism group of the graph acts transitively on the set of the adhesion sets.*

References

- [1] L. Babai. The growth rate of vertex-transitive planar graphs. *Association for Computing Machinery, New York, NY (United States)*, pages 564–573, 1997.
- [2] J. Carmesin, R. Diestel, F. Hundertmark, and M. Stein. Connectivity and tree structure in finite graphs. *Combinatorica*, 34(1):1–35, 2014.
- [3] J. Carmesin, M. Hamann, and B. Miraftab. Canonical trees of tree-decompositions. *arXiv preprint arXiv:2002.12030*.
- [4] R. Diestel. Locally finite graphs with ends: A topological approach, II. applications. *Discrete Mathematics*, 311(15):2750–2765, 2010.
- [5] R. Diestel. Locally finite graphs with ends: A topological approach, I. basic theory. *Discrete Mathematics*, 311(15):1423–1447, 2011.
- [6] R. Diestel. *Graph Theory* (Fifth edition). Springer-Verlag, 2017.
- [7] C. Droms, B. Servatius, and H. Servatius. Connectivity and planarity of Cayley graphs. *Beiträge Algebra Geom*, 39(no. 2):269–282, 1998.
- [8] A. Georgakopoulos. Characterising planar Cayley graphs and Cayley complexes in terms of group presentations. *European Journal of Combinatorics*, 36:282–293, 2014.
- [9] A. Georgakopoulos. The planar cubic Cayley graphs of connectivity 2. *European Journal of Combinatorics*, 64:152–169, 2017.
- [10] M. Hamann. Planar transitive graphs. *Electronic Journal of Combinatorics*, 25(04):25pp, 2018.
- [11] M. Hamann, F. Lehner, B. Miraftab, and T. Rühmann. A Stallings’ type theorem for quasi-transitive graphs. *arXiv preprint arXiv:1812.06312*, 2018.
- [12] H. A. Jung and M. E. Watkins. On the structure of infinite vertex-transitive graphs. *Discrete Mathematics*, 18(1):45–53, 1977.
- [13] H. Maschke. The representation of finite groups, especially of the rotation groups of the regular bodies of three-and four-dimensional space, by Cayley’s color diagrams. *American Journal of Mathematics*, 18(2):156–194, 1896.
- [14] R. B. Richter. Decomposing infinite 2-connected graphs into 3-connected components. *Electronic Journal of Combinatorics*, 11(1):25, 2004.
- [15] J. P. Serre. *Trees*. Springer-Verlag, Berlin–New York, 1980.
- [16] W. T. Tutte. Connectivity in graphs, mathematical expositions, no. 15, 1966.
- [17] M. E. Watkins. Les graphes de cayley de connectivité un. In *Problemes Combinatoires et Théorie des Graphes, Proc. International Colloquium on Graph Theory, Orsay, France*, pages 419–422, 1976.

- [18] H. Zieschang, E. Vogt, and H. D. Coldewey. *Surfaces and planar discontinuous groups*, volume 835. Springer, 2006.