

First detection of oscillations in the Halo giant HD 122563: validation of seismic scaling relations and new parameters

O. Creevey^{1,*}, F. Grundahl², F. Thévenin¹, E. Corsaro³, P. L. Pallé⁴, D. Salabert^{5,6,**},
B. Pichon¹, R. Collet², L. Bigot¹, V. Antoci², and M. F. Andersen²

¹ Université Côte d’Azur, Observatoire de la Côte d’Azur, CNRS, Laboratoire Lagrange, France
e-mail: ocreevey@oca.eu

² Stellar Astrophysics Centre, Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, 8000 Aarhus C, Denmark

³ INAF - Osservatorio Astrofisico di Catania, via S. Sofia 78, 95123-I, Catania, Italy

⁴ Instituto de Astrofísica de Canarias, 38205 La Laguna, Tenerife, Spain

⁵ IRFU, CEA, Université Paris-Saclay, F-91191, Gif-sur-Yvette, France

⁶ Université Paris Diderot, AIM, Sorbonne Paris Cité, CEA, CNRS, F-91191 Gif-sur-Yvette, France

Received November 26, 2018; accepted November 26, 2018

ABSTRACT

Aims. The nearby metal-poor giant HD 122563 is an important astrophysical laboratory for which to test stellar atmospheric and interior physics. It is also a benchmark star for which to calibrate methods to apply to large scale surveys. Recently it has been remeasured using various methodologies given the new high precision instruments at our disposal. However, inconsistencies in the observations and models have been found. In order to better characterise this star using complementary techniques we have been measuring its radial velocities since 2016 using the Hertzprung telescope (SONG network node) in order to detect oscillations.

Methods.

Results. In this work we report the first detections of sun-like oscillations in this star, and to our knowledge, a detection in the most metal-poor giant to date. We apply the classical seismic scaling relation to derive a new surface gravity for HD 122563 of $\log g_v = 1.39 \pm 0.01$ dex. Reasonable constraints on the mass imposed by its PopII giant classification then yields a radius of $30.8 \pm 1.0 R_\odot$. By coupling this new radius with recent interferometric measurements we infer a distance to the star of 306 ± 9 pc, which places it further away than previously thought and inconsistent with the Hipparcos parallax. Independent data from the Gaia mission corroborates the distance hypothesis ($d_{\text{GDR2}} = 290 \pm 5$ pc), and thus the updated fundamental parameters.

Conclusions. We confirm the validity of the seismic scaling relation without corrections for surface gravity in metal-poor and evolved star regimes. The remaining discrepancy of 0.04 dex between $\log g_{\text{GDR2}}$ ($= 1.43 \pm 0.03$) reduces to 0.02 dex by applying corrections to the scaling relations based on the mean molecular weight and adiabatic exponent. The new constraints on the HR diagram ($L_{\star,v} = 381 \pm 26$) significantly reduce the disagreement between the stellar parameters and evolution models, however, a discrepancy on the order of 150 K still exists. Fine-tuned stellar evolution calculations show that this can be reconciled by changing the mixing-length parameter by an amount (-0.35) that is in agreement with predictions from recent 3D simulations and empirical results. Asteroseismic measurements are continuing, and analysis of the full frequency data complemented by a distance estimate promises to bring important constraints on our understanding of this star and of the accurate calibration of the seismic scaling relations in this regime.

Key words. asteroseismology – stars: individual HD 122563 – stars: fundamental parameters – stars: Population II – stars: distances
methods: observational

1. Introduction

HD 122563 ($V=6.2$ mag, $14^{\text{h}}02^{\text{m}}31.8^{\text{s}}$, $+09^{\circ}41'09.95''$) is one of the brightest and closest metal-poor $[M/H] = -2.4$ (Collet et al. 2018; Prákavicius et al. 2017) giant stars that offers the possibility to be observed and analysed using many different methodologies. There are few stars for which such a complete set of observations can be obtained. The advantage of this is that we can obtain robust (almost) model-independent determinations of many of its fundamental parameters, but many of these can also be compared by using different methodologies, and so it allows us to investigate sources of systematic errors. For example, it serves as a benchmark star for testing stellar astrophysics, such as, non-LTE effects in stellar atmospheres (Heiter et al. 2015;

Thévenin & Idiart 1999; Collet et al. 2005), or 3D stellar atmosphere structure (Prákavicius et al. 2017; Collet et al. 2018). In stellar evolution models strong assumptions on its age, mass and initial helium abundance can be made which allows one to investigate tunable parameters or details of the interior physics. HD 122563 also contributes to calibrating large Galactic surveys, e.g. Gilmore et al. (2012), which aim to understand the evolution of the Milky Way (Jofré et al. 2017).

In Creevey et al. (2012b), from hereon C12, strong constraints were placed on the position of this star in the Hertzprung-Russell diagram, in particular with the determination of its interferometric diameter. It was found that classical stellar evolution models were unable to reproduce its position within the error box. One of the proposed solutions by these authors was to change the mixing-length parameter α in the stellar evolution codes, and they managed to produce a model which

* ocreevey@oca.eu

** Previous affiliation

correctly fit the observational data. Such changes are not unrealistic e.g. (Bonaca et al. 2012; Creevey et al. 2017; Tayar et al. 2017; Joyce & Chaboyer 2018; Viani et al. 2018b), however, the size of the modification suggested that something should be addressed either in the models for very metal-poor stars or in the observations. Even with such constraints on the model parameters, its derived age was not well constrained, due to degeneracies between the unobservable mass and initial helium mass fraction, see Lebreton et al. (1999) for a discussion. C12 proposed that asteroseismic observations could help to constrain these other parameters, which would improve the age determination and perhaps shed some light on the difficulty of matching the HR diagram constraints with classical models.

HD 122563 has also been the subject of several recent atmospheric studies (Amarsi et al. 2018), and some discrepancies in analyses were noted, for example, the 3D non-LTE analysis of hydrogen lines shows a spread of T_{eff} values depending on the line used. Collet et al. (2018) suggested that a lowering of the surface gravity from their adopted $\log g = 1.61 \pm 0.07$ would reduce the oxygen abundance discrepancy between molecular and atomic species. C12 also predicted a value $\log g = 1.60 \pm 0.04$ based on the measured diameter, available parallax, and a model-constrained mass.

Given the current questions and indications that the models or the observations are not entirely consistent, we proposed to observe the star in multi-epoch spectrography in order to detect stellar oscillations. Asteroseismic analysis would provide a fresh new perspective, and hopefully help to unravel the current inconsistencies, while also allowing us to investigate the scaling relation in the non-solar regime. In this paper we report the first detection of stellar oscillations in HD 122563 using the Hertzprung SONG telescope located on the Observatorio del Teide. This is the most metal-poor star (to our knowledge) to have sun-like oscillations detected¹. We measure the global seismic quantity ν_{max} for HD122563 (Sect. 2) and along with complementary information we derive a new surface gravity, radius and distance to the star (Sect. 3). New data from the Gaia mission (Gaia Collaboration et al. 2018) corroborate our results. We then discuss the implications of our results on the seismic scaling relation for $\log g$, the position of the star in the HR diagram and 1D stellar models (Sect. 4).

2. Observations of HD 122563

2.1. New asteroseismic observations from radial velocities.

We obtained time series radial velocity observations with the 1-m Hertzprung SONG telescope equipped with an echelle spectrograph and located at the Observatorio del Teide. The Hertzprung telescope is a node of the Stellar Observations Network Group (SONG). From April 2016 to December 2017 we obtained an average of one spectrum per night when the object is visible. The spectra were reduced and calibrated using the SONG pipeline. Details of the Hertzprung telescope characteristics and reduction pipeline are given in Andersen et al. (2014) and Grundahl et al. (2017). All observations were obtained using an iodine cell for precise wavelength calibration. A spectral resolution of 80 000 and an exposure time of 900s was used throughout.

The radial velocity (RV) time series is presented in Figure 1. It comprises 387 data points over a total of 449 nights. The typical uncertainty on the RV was found to be in the 11-14 m/s range,

¹ Epstein et al. (2014) report a list of 9 metal-poor stars with detected oscillations with abundances larger than -2.2 dex.

depending on the signal-to-noise ratio in the observed spectrum. However, as can be seen from the figure, there is a long-term trend. We believe that this could be evidence of convection, rotation or activity. As the trend is on the order of 300 days, far from the expected intrinsic pulsation periods, we perform a frequency analysis directly on the time series produced by the pipeline.

2.1.1. Time-series analysis




Fig. 1. Radial velocity time series of HD 122563. The oscillations are on the timescale of a few days, and a long-term trend on the order of 300 days is visible. See Sect. 2.1.1.

The power spectrum of the velocity time-series was initially calculated as an unweighted least-squares fit of sinusoids (Frandsen et al. 1995; Arentoft et al. 1998; Bedding et al. 2004; Kjeldsen et al. 2005; Corsaro et al. 2012), and converted into power spectral density (PSD) by normalizing for the spectral resolution, namely the integral of the spectral window, of $\sim 0.06 \mu\text{Hz}$. We also tested the case of a weighted least-squares fit to check for possible improvements in the signal-to-noise. For this purpose we used a weight assigned to each point of the radial velocity time-series according to the corresponding uncertainty estimate obtained from the SONG pipeline². The radial velocity uncertainties were previously rescaled in order to correct for the presence of possible outliers, following the approach presented by Butler et al. (2004) (see also Corsaro et al. 2012). Finally we measured the amplitude of the noise level in the amplitude spectrum, in the region $4\text{--}6 \mu\text{Hz}$, outside the power excess due to oscillations, for both the weighted and unweighted case. We found that the amplitude of the noise is lower in the un-weighted case, reaching down to 11.8 m s^{-1} . We therefore decided to adopt the un-weighted spectrum for further analysis.

2.1.2. Extraction of global seismic parameters ν_{max}

We used the DIAMONDS Bayesian Inference tool (Corsaro & De Ridder 2014, App. A) to model the power spectral density (PSD) of the star. The PSD and the best-fit model are shown in Fig. 2, and incorporates a flat noise component, two Harvey-like profiles to account for granulation-driven signal, and a Gaussian envelope to model the oscillation power excess (Corsaro et al. 2015). A clear excess of power due to the oscillations is detected at $3 \mu\text{Hz}$ (see Table 1), this is referred to as ν_{max} . We note that the width of the power excess is narrow, and this seems to follow the trend presented by Yu et al. (2018), although their sample only goes as low as $40 \mu\text{Hz}$. The results reported in Table 1 are obtained from calculating the 16, 50, and 84 percentiles of the marginalised distribution of ν_{max} . We note that lower and upper confidence intervals are strictly formal uncertainties, with-

² It is noted in the SONG documentation that the uncertainties reported on the RV data should be considered with caution.




Fig. 2. Power spectral density and background model fit with DIAMONDS to determine ν_{\max} . The red line shows the total fit, including the oscillation power excess. See Sect. 2.1.2.

Table 1. Observed Properties of HD 122563 used in this work

Property	Value	Source
ν_{\max} [μHz]	$3.07^{+0.05}_{-0.04}$	this work
$\theta_{\text{LD,A}}$ [mas]	0.940 ± 0.011	C12
$F_{\text{bol,A}}$ [erg s ⁻¹ cm ⁻²]	$13.16 \pm 0.36 \text{ e-}8$	C12
$T_{\text{eff,A}}$ [K]	4598 ± 41	C12
$\theta_{\text{LD,B}}$ [mas]	0.926 ± 0.011	K18
$F_{\text{bol,B}}$ [erg s ⁻¹ cm ⁻²]	$13.20 \pm 0.29 \text{ e-}8$	K18
$T_{\text{eff,B}}$ [K]	4636 ± 36	K18
π_{HIP} [mas]	4.22 ± 0.36	van Leeuwen (2007)
π_{GDR2} [mas]	3.444 ± 0.063	Lindegren et al. (2018)

out consideration of possible systematic errors. As some possible sources of errors, we also determined ν_{\max} 1) after filtering for the low-frequency signal, and 2) using a flat background in the power spectrum. In both cases our results are consistent with those reported in Table 1.

2.2. Literature values of the effective temperature of HD 122563

The T_{eff} of HD 122563 has been derived using many independent methods, with all methods showing agreement within a range of $\pm 100\text{K}$ around 4600K. A recent compilation of spectroscopically derived T_{eff} is given in Heiter et al. (2015) who recommend 4587 ± 60 K. Casagrande et al. (2014) used the (quasi-) model independent Infra Red Flux Method (IRFM) and derived a $T_{\text{eff}} = 4600 \pm 47$ K. Two recent analyses by C12 and Karovicova et al. (2018) (K18 hereafter) using interferometry obtain results also in agreement, 4598 ± 41 K and 4636 ± 36 K. These values have been obtained using independent determinations of the angular diameter θ and the bolometric flux F_{bol} (Table 1). For the latter both adapt a value of extinction³ of $A_V = 0.01$ mag.

³ In this work we rederived F_{bol} using the iterative method described in C12, but adopting the bolometric corrections from Houdashelt et al. (2000). This was done with the aim of exploring the effect of a non-negligible interstellar extinction (Sect. 4). The new F_{bol} change slightly which result in a $T_{\text{eff}} = 4610/4629$ K for C12/K18 for $A_V = 0.01$ mag.




Fig. 3. Distributions of $\log g$ derived from asteroseismic data using the seismic scaling relation. The blue and red represent the results using the observed properties from C12 and K18, respectively, and $f_{\nu_{\max}} = 1.0$. The grey and green lines show $\log g$ using the different solar reference values and a revised scaling relation (Viani et al. 2017), respectively. See Sect. 3.1 for details.

The agreement between all of the determinations provides good confidence in the T_{eff} and the assumed low value of extinction.

In this work we do not rederive T_{eff} but we adopt the two independent interferometric determinations from C12 and K18. However, we use their reported bolometric flux F_{bol} and measured angular diameter θ in order to correctly propagate the uncertainties on all of our inferences. Furthermore, θ is used along with a parallax to perform a similar exercise in order to compare our results (see Sect. 3.2). By adopting both C12 and K18 one can investigate the effect of a possible source of systematic error from θ , and considering an extreme value of extinction we investigate a source of error in F_{bol} , and consequently its T_{eff} (Sect. 4).

3. Surface gravity and distance of HD 122563

3.1. Surface gravity and distance from asteroseismic observations

It has been demonstrated in many papers that the surface gravity of a star can be derived with very high precision using asteroseismic observations e.g. Brown & Gilliland (1994); Chaplin et al. (2011); Creevey et al. (2012a). Even using simple scaling relations, such as that proposed by Kjeldsen & Bedding (1995) (KB95 hereafter), $\log g$ can be easily derived. Direct comparisons between $\log g$ derived from seismology and from other methods have also demonstrated its accuracy (Morel & Miglio 2012; Hekker et al. 2013). For non-evolved stars typical errors, including systematics on the input parameters and accuracy, are on the order of 0.04 dex (Creevey et al. 2013). However, these scaling relations are based on the Sun, and as the star begins to differ from the Sun, the relations may begin to deviate from solar-scaled values. Several authors have addressed this issue recently (Hekker et al. 2013; Coelho et al. 2015; Sharma et al. 2016; Viani et al. 2017; Kallinger et al. 2018; Brogaard et al. 2018) and propose modifications to the scaling relations.

The classical relation for the asteroseismic quantity ν_{\max} is

$$\frac{\nu_{\max}}{\nu_{\max,\odot}} = f_{\nu_{\max}} \frac{g}{g_{\odot}} \sqrt{\frac{T_{\text{eff},\odot}}{T_{\text{eff}}}} \quad (1)$$

where $f_{\nu_{\max}} = 1.0$, $\nu_{\max,\odot} = 3\,050\mu\text{Hz}$ and $T_{\text{eff},\odot} = 5\,777$ K (Kjeldsen & Bedding 1995). In this work we adopt $\log g_{\odot}$




Fig. 4. Distances to HD 122563 derived using parallaxes (black and grey) and asteroseismic inferences (blue, red and green)

= 4.438 dex from the IAU convention⁴ (Prša et al. 2016), and thus we consequently adopt $T_{\text{eff},\odot} = 5772$ K. Revised relations have been presented in some of the references cited above where $f_{\nu_{\text{max}}} \neq 1.00$, or $\nu_{\text{max},\odot} \neq 3\,050\mu\text{Hz}$, and these are both discussed below.

Recent analysis by Viani et al. (2017) (V17 hereafter) replaces $f_{\nu_{\text{max}}}$ explicitly with terms associated with the mean molecular weight and the adiabatic exponent, the theoretical basis for which has been studied in e.g. Belkacem et al. (2011). However, as these terms can not be derived without a stellar model, we are interested in testing the classical relation, but we do consider both cases. We also note that while these corrections can be important, they do not fully account for the effects of changes in composition on the atmospheric opacities and on the structure of the outer layers (as acknowledged by the authors). There are important deviations between 1D and 3D stellar models in terms of the surface stratifications at very low metallicity regime and differences between 1D and 3D stratifications contribute to this surface effect (Trampedach et al. 2017).

Using equation 1 in its classical form we calculate the surface gravity of HD 122563 from ν_{max} and T_{eff} . To correctly propagate the uncertainties reported in C12 and K18, and the determination of ν_{max} , we performed Monte-Carlo-like simulations to derive $\log g$. More specifically for each simulation we perturb the observed quantity (F_{bol} , θ) by adding a random number drawn from a Normal distribution with width 1 scaled by its symmetric uncertainty. For ν_{max} we used the marginal distribution directly from Sect. 2. The distributions of the resulting values of $\log g$ from the simulations are shown in Fig. 3 adopting the two referenced interferometric measurements (blue and red, respectively). The value of $\log g_{\nu}$ for K18 (red) is 1.393 ± 0.007 dex, where the subscript ν denotes an asteroseismically-derived value. The median values using C12 and K18 differ by 0.0018 dex. A possible source of systematic error arises from the definition of the solar parameters, providing a change of -0.0002 dex for $T_{\text{eff},\odot}$ and -0.0056 for $\nu_{\text{max},\odot}$, and the combined change is shown for C12 by the grey histogram.

For the more recent investigations of the scaling relation, $f_{\nu_{\text{max}}}$ is replaced by terms associated with the mean molecular weight (μ/μ_{\odot})^{1/2} or a combination of μ and the adiabatic exponent ($\Gamma_1/\Gamma_{1,\odot}$)^{1/2}. This is justified by the fact that ν_{max} is expected to scale with the cut-off frequency in the atmosphere (see refer-

ences above). In practice, these values are not readily obtained and require the use of stellar models. For μ this may be estimated by making assumptions about the helium abundance of the star and using the observed metallicity. For Γ_1 a stellar model is needed. Using the stellar models from Sect. 4 we calculated $\mu_{\star} = 0.5904$ and $\Gamma_{1,\star} = 1.545$, and adopted the solar values, $\mu_{\odot} = 0.6159$ and $\Gamma_{1,\odot} = 5/3$, in order to derive the corrections to the scaling relation.

Applying the correction associated only with μ results in an increased $\log g$ of 0.007 dex. However, applying the correction associated with Γ_1 has a more significant impact. The resulting distribution for K18 is represented by the green histogram in Fig. 3. It results in a systematic change of $+0.0256$ dex to yield a $\log g_{\nu} = 1.418 \pm 0.007$ (see Table 2).

Given cosmological constraints (upper age) and our knowledge of stellar evolution, we can assume that the mass of this evolved star is likely between 0.80 and $0.90 M_{\odot}$ (C12). By adopting a conservative prior on the mass of $0.85 \pm 0.05 M_{\odot}$, the radius R can be inferred from $\log g$ and the mass prior. This then gives us access to the distance of the star, because the angular diameter has been measured. From the classical relation with $f_{\nu_{\text{max}}} = 1.0$ we derive a radius of $30.8 \pm 1.0 R_{\odot}$ implying a distance of 305 ± 10 pc for C12 (see Table 2). The smallest derived distance is 296 ± 9 pc using C12+V17. These distances are larger than that proposed by van Leeuwen (2007) who measured its parallax of 4.22 ± 0.35 mas using data from the Hipparcos mission. The derived distances are illustrated in Fig. 4 using the same color-code as Fig. 3, with the latter denoted by the grey dashed lines.

3.2. The surface gravity and distance from Gaia DR2 measurements

The Gaia DR2 catalogue (Gaia Collaboration et al. 2018; Lindgren et al. 2018) provides a new and more precise parallax for HD 122563 of 3.444 ± 0.063 mas. The distance to the star inferred from this newer parallax is then 290 ± 5 pc if we assume no prior, and thus 30% further away than previously thought (see Fig. 4, black distribution). This value is consistent at the 1σ level with the values obtained using asteroseismology. Following the methodology from Sect. 3.1, using the angular diameter and the parallax measurement and adopting a mass prior, we infer (a new radius) and $\log g$ for HD 122563. This results in $\log g_{\text{GDR2}} = 1.432^{+0.030}_{-0.033}$ and $1.445^{+0.031}_{-0.033}$ using C12 and K18, respectively. We furthermore use F_{bol} to derive luminosity L_{\star} (Sect. 4). The distance d , radius, surface gravity and L_{\star} are summarized in Table 2 under the heading ‘GDR2 Parallax’.

A potential systematic error of $+0.029$ mas on the parallax has been documented in the Gaia Second Data Release (Luri et al. 2018). This zeropoint corresponds to the difference between the median value of the quasars observed by Gaia, which are assumed to have no parallactic motion, and zero. For completeness, we apply this error also in our analysis, and calculate the corresponding parameters (Table 2). A larger parallax implies a smaller distance and radius, and a higher value of $\log g$. This is further away from the seismic value. However, as this zeropoint is a value derived for faint quasars, there is no reason to expect it to apply to the brighter end of the Gaia spectrum.

4. Observational constraints in the HR diagram

We plot the position of the new observational constraints in the HR diagram in Fig. 5 using the results from the classical scaling

⁴ https://www.iau.org/static/resolutions/IAU2015_English.pdf Resolution B3

Table 2. Derived properties of HD 122563 using asteroseismology (left) and Gaia parallaxes (right). For the seismology parameters, we show the results using the classic scaling relation (KB95) and that given in V17, using both C12 and K18 measurements. For the results using the GDR2 parallax we show the results without and with a systematic error $s(\pi)$ of $+0.029$ mas on the parallax, again for C12 and K18.

		Seismology				GDR2 Parallax			
		KB95		V17		$s(\pi) = 0.0$		$s(\pi) \neq 0.0$	
		C12	K18	C12	K18	C12	K18	C12	K18
$\log g$	[dex]	1.391^{+8}_{-7}	1.393^{+7}_{-7}	1.416^{+8}_{-7}	1.418^{+7}_{-7}	1.433^{+30}_{-32}	1.446^{+30}_{-32}	1.441^{+30}_{-33}	1.453^{+31}_{-33}
d	[pc]	305^{+10}_{-10}	308^{+10}_{-10}	296^{+9}_{-9}	299^{+10}_{-10}	290^{+5}_{-5}	—	288^{+5}_{-5}	—
R_\star	[R_\odot]	30.8^{+10}_{-10}	30.7^{+9}_{-9}	29.9^{+9}_{-9}	29.8^{+9}_{-9}	29.4^{+6}_{-7}	28.9^{+7}_{-6}	29.1^{+7}_{-6}	29.7^{+6}_{-6}
L_\star	[L_\odot]	381^{+26}_{-26}	392^{+27}_{-26}	359^{+25}_{-24}	370^{+25}_{-24}	346^{+16}_{-15}	347^{+15}_{-15}	340^{+15}_{-16}	341^{+14}_{-14}




Fig. 5. HR diagram presenting the revised positions of HD 122563 based on asteroseismic data (red and blue) and Gaia data (black) for K18 and C12 (continuous and dashed lines, respectively). The value presented in C12 using the parallax from van Leeuwen (2007) is shown as the grey dashed box. The red vector represents a potential shift in the median values of L_\star and T_{eff} if we consider an extinction $A_V = 0.08$ mag (the reference value is 0.01 mag). Classical evolution tracks for a 0.75 and 0.90 M_\odot model from BASTI (Pietrinferni et al. 2004) are shown by the green lines, while a fine-tuned 0.86 M_\odot model using the CESAM2K evolution code (Morel 1997) with a reduced value of the mixing-length parameter compared to the solar one is shown by the grey continuous curve. See Sect. 4 for details.

relation and considering no systematic error on the parallax. We also show BASTI evolutionary tracks (Pietrinferni et al. 2004) for a 0.75 and 0.90 M_\odot star using standard solar scaled physics (non-canonical, alpha-enhanced shifts the tracks to hotter temperatures). These tracks are frequently used in the literature. The new HR diagram constraints are shown using the same color-code as Fig. 3 with the grey dashed lines representing C12 constraints.

The discrepancy between the observational error box and the evolution tracks already reduces from $> 300\text{K}$ (C12) to the order of 100K (considering the uncertainties) using the new constraints. However, assuming these models to be correct, we would still require $L_\star \sim 550 L_\odot$, assuming the T_{eff} is correct. Given that we have two independent measures of the distance suggesting similar luminosities it is likely that it is the stellar models that need to be adjusted, based on the assumed $A_V = 0.01$ mag (and therefore T_{eff}).

We investigate the effect of assuming non-negligible interstellar extinction⁵ of $A_V = 0.08$ mag by considering the maximum values from (Lallement et al. 2014), although most studies indicate that this should not be the case. This would increase F_{bol} and consequently increase T_{eff} , for a fixed θ . The increase in the median values of T_{eff} and L_\star as a result of imposing $A_V = 0.08$ mag is indicated by the red vector in Fig. 5 for K18. This could explain some of the discrepancy with evolution models, although such a strong absorption seems unlikely and would also bring the various T_{eff} determinations into disagreement.

In this work we have assumed a conservative mass of $0.85 \pm 0.05 M_\odot$. However, using stellar models we can further constrain the mass by assuming a limited age range to be consistent with a Pop II star. We performed computations of evolutionary tracks with the CESAM2k code (Morel 1997) assuming $[Z/X] = -2.4$ which includes an assumed $[a/\text{Fe}] \sim +0.25$, similar to those described in C12. In order to match the constraints, we are required to lower the mixing-length parameter by ~ 0.35 compared to the solar one, to 1.55 ± 0.03 . Only models with masses in the range of $0.85 - 0.87 M_\odot$ reach the L_\star at an age between 10 and 12 Gyr. A new representative stellar model (0.86 M_\odot , $\alpha = 1.55$) calculated from CESAM2k is shown in grey in Fig. 5.

5. Discussion

We determined an asteroseismic $\log g$ value of 1.39 ± 0.01 and 1.42 ± 0.01 by using the ν_{max} seismic scaling relation without and with corrections for the mean molecular weight and adiabatic exponent. These values are in statistical agreement with those derived using a parallax from GDR2 (1.43 ± 0.03). A recent determination from the APOGEE survey (DR15, Majewski et al. 2017) yields a *calibrated*⁶ $\log g$ of 1.43 dex. These agreements imply that the seismic scaling relation for surface gravity even without corrections is valid in the evolved, metal-poor regime. Our work also validates the use of the corrective terms, although some discrepancy still remains. This has an important consequence, because today we have access to thousands of giant stars which show oscillation signatures such as ν_{max} , and this implies that seismic constraints in spectroscopic analyses can be

⁵ <http://stilism.obspm.fr/reddening?frame=icrs&vlong=210.63268943&ulong=deg&vlat=09.68609665&ulat=deg&valid=>

⁶ The APOGEE stellar parameters provide calibrated and uncalibrated stellar parameters, where the calibration is done independently of the other stellar parameter. For surface gravity, the calibrated value is obtained by adding a constant that is derived from seismic calibrations of spectra. For T_{eff} however this is obtained by adding a constant derived from photometrical calibrations with models. For the T_{eff} , the calibrated value is not valid in our case.

safely used. Fixing this parameter allows more precise determinations of other spectroscopic parameters, and can help to uncover systematic errors associated with these analyses.

A recent analysis of the CNO abundances of HD 122563 using 3D hydrodynamical atmospheres by Collet et al. (2018) found that their imposed surface gravity of 1.61 ± 0.07 resulted in discrepant oxygen abundances between molecular and atomic species. They suggested that a downward revision on the order of 0.3 dex would relieve this tension. In this work, the surface gravity has been revised downward by 0.2 dex, close to that proposed by the authors.

Using stellar models and the constraints from the revised luminosity, we refined the mass to within 0.85 and $0.87 M_{\odot}$. Nevertheless, we still find differences compared to standard stellar models. We looked into the possibility of having a possible increase in interstellar extinction. This would bring the error box in the HR diagram closer to the BASTI stellar evolution tracks (red vector in Fig. 5), because the T_{eff} would increase. However, then all different methodologies (spectroscopic, interferometric, IRFM) would no longer be in agreement. A more recent measurement of the T_{eff} with APOGEE spectra yields an uncalibrated (i.e. derived from the spectra) T_{eff} of 4594 K, also in agreement with the literature T_{eff} presented here. An alternative is to increase the L_{\star} only, i.e. a star that is much further away. From this work we have two independent determinations of the distance yielding similar results. This would also be unlikely. This leaves the only possibility to investigate the stellar models.

In order to match the observational constraints with stellar models, we could increase the metallicity by about 1.0 dex, however, this has been measured by many different authors and this is a very unlikely solution. We could also decrease the initial helium abundance to an extremely low value, but this would be inconsistent with predictions of the primordial abundances, e.g. Tytler et al. (2000). One of the solutions of this discrepancy is to lower the mixing-length parameter α . This decreases the T_{eff} , without influencing the L_{\star} . We estimated a shift of α of -0.35 compared to the solar-calibrated one by calculating refined stellar models. Such a value is in agreement with estimations from 3D simulations. Magic et al. 2015 suggest a reduction of ~ 0.2 compared to the solar value for a star of this g , $[M/H]$, and T_{eff} (their Fig. 3), where the reduction is primarily due to the lower value of $\log g$.

More recently, Tayar et al. (2017), Viani et al. (2018a) and Creevey et al. (2017) investigated the mixing-length parameter empirically by studying samples of stars. For the second two, their analysis concentrated on main sequence stars and subgiants, with the lowest metallicity values close to -0.60 dex, so their results are not applicable to HD 122563. However, they do find a relation that depends on $\log g$, T_{eff} and metallicity. Tayar et al. (2017) also restricted their studies to stars with $[Fe/H] > -1.0$ dex, but looked at the specific case of a sample of 3,000 giants, with the aim of studying the metallicity dependence. They found that a correction to α on the order of 0.2 per dex is needed. If we apply this correction, we would require a reduction of ~ 0.5 dex in α compared to the solar one, not far from what we find. Both empirical and model results indeed support that α needs to be modified in the 1D stellar models. This then would support our new parameter determinations, which consequently validates the seismic scaling relation for $\log g$ in the metal-poor giant regime.

6. Conclusions

In this work we described the first detection of oscillations in the metal-poor giant HD 122563. We determined its surface gravity using the detected ν_{max} along with scaling relations. By comparing with the value derived using a Gaia parallax we validated the classical ($f_{\nu_{\text{max}}} = 1.0$) seismic scaling relations in such a metal-poor and non-solar regime. We found a non-significant difference of 0.04 dex. While these relations are valid, applying the corrections for mean molecular weight and the adiabatic exponent results in a smaller discrepancy with the surface gravity derived from the Gaia parallax (0.02 dex).

We derived updated surface gravity, radius, and luminosity for HD 122563. These new parameters are quite different from previous determinations. These permit us to make a new estimate of the mass by using stellar models. Only models with masses between 0.85 and $0.87 M_{\odot}$ satisfy the new constraints. The updated luminosity along with the literature T_{eff} provide a new error box in the HR diagram. The large difference on the T_{eff} -axis found in earlier works has significantly reduced, to the order of 100 K. This final difference can be rectified by modifying the mixing-length parameter used in the models. We needed a change of -0.35 compared to the solar-calibrated value, a value in agreement with 3D simulations and empirically derived values.

SONG radial velocity observations are continuing in order to determine $\langle \Delta \nu \rangle$ and resolve the individual frequencies. These observations will help to test scaling relations for the radius and the mass using $\langle \Delta \nu \rangle$ outside of the solar regime, and bring important constraints on the knowledge of the fundamental parameters of this star including its age. Complementing these data with high precision parallax measurements will allow us to derive accurate masses and radii independent of models.

Acknowledgements. We are grateful for the Programme National de Physique Stellaire for financial support for this research project. This work is based on observations made with the Hertzprung SONG telescope operated on the Spanish Observatorio del Teide on the island of Tenerife by the Aarhus and Copenhagen Universities and by the Instituto de Astrofísica de Canarias. Support for the construction of the Hertzprung SONG Telescope from the Instituto de Astrofísica de Canarias, the Villum Foundation, the Carlsberg Foundation and the Independent Research Fund Denmark is gratefully acknowledged. E.C. is funded by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 664931. D.S. acknowledges the financial support from the CNES GOLF grant and the Observatoire de la Côte d'Azur for support during his stays. We also deeply acknowledge the Gaia team for their huge efforts in bringing us high quality data.

Appendix A: Software and Observations

This work made use of the following free public data and software

- The radial velocity data were obtained using the Hertzprung SONG telescope, which is operated on the Spanish Observatorio del Teide on the island of Tenerife by the Aarhus and Copenhagen Universities and by the Instituto de Astrofísica de Canarias.
- Parallaxes from the ESA Gaia Space Mission Data Release 2.
- The VizieR catalogue access tool, CDS, Strasbourg, France. The original description of the VizieR service was published in A&AS 143, 23.
- The radial velocities were analysed using the SONG reduction pipeline.
- Extinction map tools from the Stilism project <https://stilism.obspm.fr/>

- The frequency analysis was done using the DIAMONDS code. This code is available at <https://github.com/EnricoCorsaro/DIAMONDS>.
- This article was prepared using overleaf.
- The figures were prepared using Jupyter-Notebook.

References

- Amarsi, A. M., Nordlander, T., Barklem, P. S., et al. 2018, *A&A*, 615, A139
- Andersen, M. F., Grundahl, F., Christensen-Dalsgaard, J., et al. 2014, in *Revista Mexicana de Astronomía y Astrofísica Conference Series*, Vol. 45, *Revista Mexicana de Astronomía y Astrofísica Conference Series*, 83–86
- Arentoft, T., Kjeldsen, H., Nuspl, J., et al. 1998, *A&A*, 338, 909
- Bedding, T. R., Kjeldsen, H., Butler, R. P., et al. 2004, *ApJ*, 614, 380
- Belkacem, K., Goupil, M. J., Dupret, M. A., et al. 2011, *A&A*, 530, A142
- Bonaca, A., Tanner, J. D., Basu, S., et al. 2012, *ApJ*, 755, L12
- Brogaard, K., Hansen, C. J., Miglio, A., et al. 2018, *MNRAS*, 476, 3729
- Brown, T. M. & Gilliland, R. L. 1994, *ARA&A*, 32, 37
- Butler, R. P., Bedding, T. R., Kjeldsen, H., et al. 2004, *ApJ*, 600, L75
- Casagrande, L., Portinari, L., Glass, I. S., et al. 2014, *MNRAS*, 439, 2060
- Chaplin, W. J., Kjeldsen, H., Christensen-Dalsgaard, J., et al. 2011, *Science*, 332, 213
- Coelho, H. R., Chaplin, W. J., Basu, S., et al. 2015, *MNRAS*, 451, 3011
- Collet, R., Asplund, M., & Thévenin, F. 2005, *A&A*, 442, 643
- Collet, R., Nordlund, Å., Asplund, M., Hayek, W., & Trampedach, R. 2018, *MNRAS*, 475, 3369
- Corsaro, E. & De Ridder, J. 2014, *A&A*, 571, A71
- Corsaro, E., De Ridder, J., & García, R. A. 2015, *A&A*, 579, A83
- Corsaro, E., Grundahl, F., Leccia, S., et al. 2012, *A&A*, 537, A9
- Creevey, O. L., Doğan, G., Frasca, A., et al. 2012a, *A&A*, 537, A111
- Creevey, O. L., Metcalfe, T. S., Schultheis, M., et al. 2017, *A&A*, 601, A67
- Creevey, O. L., Thévenin, F., Basu, S., et al. 2013, *MNRAS*, 431, 2419
- Creevey, O. L., Thévenin, F., Boyajian, T. S., et al. 2012b, *A&A*, 545, A17
- Epstein, C. R., Elsworth, Y. P., Johnson, J. A., et al. 2014, *ApJ*, 785, L28
- Frandsen, S., Jones, A., Kjeldsen, H., et al. 1995, *A&A*, 301, 123
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1
- Gilmore, G., Randich, S., Asplund, M., et al. 2012, *The Messenger*, 147, 25
- Grundahl, F., Fredslund Andersen, M., Christensen-Dalsgaard, J., et al. 2017, *ApJ*, 836, 142
- Heiter, U., Jofré, P., Gustafsson, B., et al. 2015, *A&A*, 582, A49
- Hekker, S., Elsworth, Y., Mosser, B., et al. 2013, *A&A*, 556, A59
- Houdashelt, M. L., Bell, R. A., & Sweigart, A. V. 2000, *AJ*, 119, 1448
- Jofré, P., Heiter, U., Worley, C. C., et al. 2017, *A&A*, 601, A38
- Joyce, M. & Chaboyer, B. 2018, *ApJ*, 856, 10
- Kallinger, T., Beck, P. G., Stello, D., & García, R. A. 2018, *A&A*, 616, A104
- Karovicova, I., White, T. R., Nordlander, T., et al. 2018, *MNRAS*, 475, L81
- Kjeldsen, H. & Bedding, T. R. 1995, *A&A*, 293, 87
- Kjeldsen, H., Bedding, T. R., Butler, R. P., et al. 2005, *ApJ*, 635, 1281
- Lallement, R., Vergely, J.-L., Valette, B., et al. 2014, *A&A*, 561, A91
- Lebreton, Y., Perrin, M.-N., Cayrel, R., Baglin, A., & Fernandes, J. 1999, *A&A*, 350, 587
- Lindgren, L., Hernández, J., Bombrun, A., et al. 2018, *A&A*, 616, A2
- Luri, X., Brown, A. G. A., Sarro, L. M., et al. 2018, *A&A*, 616, A9
- Magic, Z., Weiss, A., & Asplund, M. 2015, *A&A*, 573, A89
- Majewski, S. R., Schiavon, R. P., Frinchaboy, P. M., et al. 2017, *AJ*, 154, 94
- Morel, P. 1997, *A&AS*, 124, 597
- Morel, T. & Miglio, A. 2012, *MNRAS*, 419, L34
- Pietrinferni, A., Cassisi, S., Salaris, M., & Castelli, F. 2004, *ApJ*, 612, 168
- Prakapavičius, D., Kučinskas, A., Dobrovolskas, V., et al. 2017, *A&A*, 599, A128
- Prša, A., Harmanec, P., Torres, G., et al. 2016, *AJ*, 152, 41
- Sharma, S., Stello, D., Bland-Hawthorn, J., Huber, D., & Bedding, T. R. 2016, *ApJ*, 822, 15
- Tayar, J., Somers, G., Pinsonneault, M. H., et al. 2017, *ApJ*, 840, 17
- Thévenin, F. & Idart, T. P. 1999, *ApJ*, 521, 753
- Trampedach, R., Aarslev, M. J., Houdek, G., et al. 2017, *MNRAS*, 466, L43
- Tytler, D., O’Meara, J. M., Suzuki, N., & Lubin, D. 2000, *Physica Scripta Volume T*, 85, 12
- van Leeuwen, F. 2007, *A&A*, 474, 653
- Viani, L. S., Basu, S., Chaplin, W. J., Davies, G. R., & Elsworth, Y. 2017, *ApJ*, 843, 11
- Viani, L. S., Basu, S., Joel Ong J., M., Bonaca, A., & Chaplin, W. J. 2018a, *ApJ*, 858, 28
- Viani, L. S., Basu, S., Joel Ong J., M., Bonaca, A., & Chaplin, W. J. 2018b, *ApJ*, 858, 28
- Yu, J., Huber, D., Bedding, T. R., et al. 2018, *ApJS*, 236, 42