

Comment on “Quantization of the damped harmonic oscillator”
[Serhan et al, J. Math. Phys. **59**, 082105 (2018)]

Zafar Ahmed^{1,*}, Sachin Kumar², and Abhijit Baishya³

¹*Nuclear Physics Division, Bhabha Atomic Research Centre, Mumbai 400 085, India*

^{*}*Homi Bhabha National Institute, Mumbai 400 094 , India*

²*Theoretical Physics Section, Bhabha Atomic Research Centre, Mumbai 400 085, India*

³*Human Resource Development Division,*

*Bhabha Atomic Research Centre, Mumbai 400 085, India**

(Dated: November 29, 2021)

Abstract

A recent paper [J. Math. Phys. **59**, 082105 (2018)] constructs a Hamiltonian for the (dissipative) damped harmonic oscillator. We point out that non-Hermiticity of this Hamiltonian has been ignored to find real discrete eigenvalues which are actually non-real. We emphasize that non-Hermiticity in Hamiltonian is crucial and it is a quantal signature of dissipation.

*Electronic address: 1:zahmed@barc.gov.in, 2: sachinv@barc.gov.in, 3: abhijitb498@gmail.com,

Heuristically, by making a Hamiltonian non-Hermitian one can account for dissipation of energy and matter. Presently interesting frameworks are being developed [1] wherein classical dissipative systems can be associated with non-Hermitian quantum Hamiltonians. The simplest model of dissipative systems is classical damped harmonic oscillator (DHO) whose equation of motion is

$$\ddot{q} + \lambda \dot{q} + \omega^2 q = 0. \quad (1)$$

Recently, this has been transformed to a Hamiltonian H [2].

$$H = \frac{p_y^2}{2m} + \frac{1}{2}m\omega^2 y^2 + \frac{\lambda y p_y}{2}. \quad (2)$$

We point out that the non-Hermiticity of H due to $(y p_y)^\dagger = y p_y - i\hbar$ has been ignored and real discrete eigenvalues

$$E_n = (n + 1/2)\hbar\sqrt{\omega^2 - \lambda^2/4} \quad (3)$$

have been obtained [2] by some N-U method [2]. Using $y p_y - p_y y = i\hbar$, H can be re-written as

$$H = \frac{p_y^2}{2m} + \frac{1}{2}m\omega^2 y^2 + \frac{\lambda(y p_y + p_y y)}{4} + \frac{i\hbar\lambda}{4}. \quad (4)$$

This can be re-written as

$$H = \frac{(p_y + m\lambda y/2)^2}{2m} + \frac{1}{2}m(\omega^2 - \lambda^2/4)y^2 + \frac{i\hbar\lambda}{4}. \quad (5)$$

Consider the eigenvalue equation: $H\psi = E\psi \Rightarrow \eta H \eta^{-1} \eta\psi = E \eta\psi$ (see [3]). Let us choose $\eta = \exp(im\lambda y^2/4\hbar)$. We find that

$$\begin{aligned} e^{(im\lambda y^2/4\hbar)} [p_y + m\lambda y/2] e^{-(im\lambda y^2/4\hbar)} &= p_y, \\ e^{(im\lambda y^2/4\hbar)} [p_y + m\lambda y/2]^n e^{-(im\lambda y^2/4\hbar)} &= p_y^n. \end{aligned} \quad (6)$$

Consequently,

$$\eta H \eta^{-1} = \frac{p_y^2}{2m} + \frac{1}{2}m(\omega^2 - \lambda^2/4)y^2 + \frac{i\hbar\lambda}{4}. \quad (7)$$

Finally the quantization of (7) leads to

$$E_n = (n + 1/2)\hbar\sqrt{\omega^2 - \lambda^2/4} + \frac{i\hbar\lambda}{4}, \quad (8)$$

which are complex and it is rightly so, as the Hamiltonian (2) is non-Hermitian. One needs to highlight that the Hamiltonian (2) for the damped harmonic oscillator (1) is crucially non-Hermitian, this will help in the development of quantization of dissipative systems.

References

1. Eva-Maria Graefe, Michael Honing, and Hans Jurgen Korsch, J. Phys. A: Math. Theor. **43** 075306 (2010).
2. M. Serhan, M. Abusini, Ahmed Al-Jamel, H. El-Nasser, and Eqab M. Rabei, J. Math. Phys. J. Math. Phys. **59**, 082105 (2018).
3. Z. Ahmed, Phys. Lett. A **294** 287 (2002).