

Assessing the quantumness of the annealing dynamics via Leggett Garg inequalities: a weak measurement approach

V. Vitale,¹ G. De Filippis,^{1,2} A. De Candia,¹ A. Tagliacozzo,^{1,2} V. Cataudella,^{1,2} and P. Lucignano^{2,1,a}

¹*Dipartimento di Fisica "Ettore Pancini", Università di Napoli "Federico II", Monte S. Angelo, I-80126 Napoli, Italy*

²*CNR-SPIN, Monte S. Angelo via Cinthia, I-80126 Napoli, Italy*

(Dated: March 11, 2022)

Adiabatic quantum computation (AQC) [1] is a promising counterpart of universal quantum computation [2], based on the key concept of quantum annealing (QA) [3]. QA is claimed to be at the basis of commercial quantum computers [4] and benefits from the fact that the detrimental role of decoherence and dephasing seems to have poor impact on the annealing towards the ground state. While many papers [5–7] show interesting optimization results with a sizable number of qubits, a clear evidence of a full quantum coherent behavior during the whole annealing procedure is still lacking. In this paper we show that quantum non-demolition (weak) measurements [8] of Leggett Garg inequalities can be used to efficiently assess the quantumness of the QA procedure. Numerical simulations based on a weak coupling Lindblad approach [9] are compared with classical Langevin simulations to support our statements.

Although any quantum algorithm can be run on adiabatic quantum computers [10], the interest of the scientific community is focused on decision and optimization problems that are very difficult to handle on classical computers, because their computational time, most of the times, grows exponentially with the number of bits. Optimization problems can be mapped onto complex many body hamiltonians [11], hence AQC is also of outmost interest from the fundamental point of view, as it may provide insights into longstanding problems in modern condensed matter physics such as physics of the strongly correlated cuprate materials [12] and of the spin glasses [13].

AQC is founded on QA, a slow quantum dynamics that proceeds from an initial Hamiltonian with a trivial ground state (easy to prepare), to a final Hamiltonian whose ground state encodes the solution of the computational problem. The adiabatic theorem guarantees that the system will track the instantaneous ground state if the Hamiltonian varies sufficiently slowly [14]. QA can perform better than thermal annealing [15, 16], however, to the date, there are only a few problems where this quantum speed-up has been clearly demonstrated. Furthermore, in the last few years, there has been a renewed interest in QA [17–21], because physical superconducting implementations of "quantum" annealers [5–7] up to thousand of spins have been used to obtain the GS

of interacting many body Hamiltonians. Moreover also optical techniques have been successfully used [22, 23], though with a smaller number of qubits. In real devices, the presence of the environment introduces decoherence and thermalization time scales that have to be longer than the annealing time [9]. This condition does not strictly guarantee that the dynamics can be assumed to be quantum for the whole evolution. Moreover, as the presence of a classicizing environment is unavoidable, the questions we want to address in the present letter are the following:

To what extent, the QA can be considered a quantum coherent dynamics, in the presence of a dissipative environment?

Once we choose an annealing time t_f at which a certain target state is (almost) reached, is it possible to find a "quantumness" estimator of the evolution towards such state?

Giving a clear answer to the previous questions, is not only a matter of semantics. It is well known that quantum speed up [2] can only be accessed if quantum mechanical coherence is preserved during the whole dynamics.

In order to answer to these questions, we propose to evaluate the Leggett-Garg's inequalities (LGI) during the QA. LGI were developed in 1985, to study quantum coherence properties of macroscopic quantum systems [24], that have been recently studied to assess the quantumness of a damped two level system [25]. They are Bell's-like inequalities in time, and predict anomalous values for some correlation functions that are only possible if the system behaves according to quantum mechanics. They also provide sharp bounds for classical correlation functions [26].

In this paper we focus on a simple model i.e. we study the QA of a single qubit.

Of course, the dynamics of a single qubit does not contain all the relevant ingredients of an interacting multi-qubit system. Phenomena like many-body tunneling, entanglement, and many body localization are, of course, not included. Although the generalization of the proposed approach to multi-qubits is straightforward (see Supplementary Information) we believe that the simple

single-qubit case is the fundamental first step for understanding how quantum mechanical coherent behaviors are spoiled by the system-bath interaction, even if it neglects the aforementioned many-qubit phenomena. However, even if a full analysis is beyond the aim of the present work, in the Supplementary Information we show that the proposed procedure is still meaningful for two interacting qubits providing that the coupling is not very strong. Hence, in the following, for sake of clarity, we give a detailed description of the physics of the single qubit. The QA is described by the following time-dependent Hamiltonian:

$$H(s) = (1-s)\frac{\Gamma_x}{2}\sigma_x + s\frac{\Gamma_z}{2}\sigma_z. \quad (1)$$

where $s = t/t_f$, $s \in [0, 1]$, t_f is so-called annealing time and σ_i ($i = x, z$) are the Pauli matrices describing the qubit as a quantum two level system. At the initial time $s = 0$ the system is prepared into the GS that is the σ_x eigenstate $|GS(0)\rangle = 1/\sqrt{2}(|\uparrow\rangle - |\downarrow\rangle)$, and eventually at the final time $s = 1$ it is annealed towards $|GS(1)\rangle = |\downarrow\rangle$. Of course this single qubit QA does not entail the complexity of a many qubit problem, however it is a simple prototypical model to address decoherence and relaxation phenomena out of equilibrium. Following Ref. [27], we set our time/energy scale choosing $\Gamma_x = \Gamma_z = 1\text{GHz}$ (that is the typical working frequency of the experimentally relevant annealers based on superconducting flux qubits [4]) and express all the energies in units of Γ_x (times in units of $1/\Gamma_x$, $\hbar = 1$). The qubit environment is described by a bath of bosonic harmonic oscillators in thermal equilibrium at the inverse temperature $\beta = 1/k_B T$ (k_B is the Boltzmann constant). The qubit bath coupling is described by an ohmic spectral density whose effective interaction strength is the dimensionless parameter α . At $\alpha = 1$ (for $s = 0$) the system undergoes the Leggett transition [28], however in this paper we will focus on the weak coupling limit ($\alpha \ll 1$) far away from this critical point. For a detailed description of the system bath interaction please refer to the Supplementary Information.

The reduced density matrix ρ_Q describing the qubit only is obtained by tracing over the environment degrees of freedom. We adopt a Lindblad approach (for details see Ref.s [9, 29]) that gives the following master equation for the density matrix:

$$\frac{d\rho_Q(t)}{dt} = -i[H(t) + H_{LS}(t), \rho_Q(t)] + \mathcal{D}_t[\rho_Q(t)], \quad (2)$$

where $H_{LS}(t)$ is the Lamb shift Hamiltonian and \mathcal{D}_t is the dissipator, responsible for the non unitary dynamics. They are both described in terms of local (in time) Lindblad operators $L(t)$. A detailed description of $H_{LS}(t)$, \mathcal{D}_t and the Lindblad operators is given in the Supplementary Information.

In this paper we focus on two estimators.

The *residual energy* ϵ_{res} tells us whether our adiabatic dynamics is successful (or not) in reaching the target




FIG. 1. Log-log plot of the residual energy of the system as a function of the variance D and of the times at which the measurements are performed: $t_1 = \tau$ and $t_2 = 2\tau$. We set $t_f = 14$. We observe that the residual energy decreases eventually going to $5.49 \cdot 10^{-4}$ which is the residual energy of the system in the absence of measurements. The inset in the right top corner shows a diagram of the system-ancilla ensemble.

state and is defined as the difference between the energy of the system at the final time t_f and the exact ground state $E_0(t_f)$ of the target Hamiltonian $H(t_f)$.

$$\epsilon_{\text{res}} = \text{Tr}[\rho_Q H(t_f)] - E_0(t_f), \quad (3)$$

Of course, due to the adiabatic theorem [30] ϵ_{res} tends to zero if $t_f \rightarrow \infty$, if the evolution is unitary.

The *Leggett-Garg's correlation functions* tell us if the system behaves quantum coherently during its dynamics [26]. We focus on the third-order Leggett-Garg's function:

$$K_3^a = C_{12} + C_{23} - C_{13} \quad (4)$$

$$C_{i,j} = \langle \sigma_z(t_i) \sigma_z(t_j) \rangle$$

and other nonequivalent third order functions $K_3^b = -C_{12} - C_{23} - C_{13}$, and $K_3^c = -C_{12} + C_{13} + C_{23}$ obtained by nontrivial cycling of the 1, 2, 3 indexes (all the other permutations are trivially reduced to one of these three). If the system behaves classically then:

$$-3 \leq K_3^i \leq 1, \quad i \in \{a, b, c\}; \quad (5)$$

Hence, in the following, we seek for violation of Leggett Garg's inequalities during the annealing to make sure that our system behaves quantum mechanically up to the annealing time t_f .

Studying the LGI requires the evaluation of two times correlation functions, which implies measuring the system twice during the annealing dynamics. Conventional




FIG. 2. Plot of the Leggett-Garg's function, in the absence of coupling to the environment. The lines are obtained performing projective measurements. The dots are calculated with weak measurements considering $N = 10^6$, $D = 20$. K_3^a in red, K_3^b in black and K_3^c in green. The orange line marks the bound of the LGI. Here $\tau = t_2 - t_1 = t_3 - t_2$, $\tau \in [0, t_f/2]$.

projective measurements [2] are detrimental for the adiabatic quantum computation, indeed after the measurement the system may populate instantaneously many excited states (see Supplementary Information) reducing the fidelity very close to zero. By contrast, we adopt the paradigm of weak measurements [8], to gain relevant information from the LGI with negligible effect on the annealing results.

In particular we propose adopting the measurement approach designed in Ref. [31] to weakly measure the LGI. The qubit described by a pseudospin degree of freedom σ_z is coupled to an ancilla. Measuring the ancilla we can extract information on the qubit state. The stronger the coupling between the two systems, the larger the influence on the quantum annealing dynamics (for a detailed discussion on the measurement scheme see Supplementary Information). In order to specify an experimentally accessible weak measurement setup, following Ref. [31], we describe a superconducting flux qubit (based, for instance on a RF SQUID) inductively coupled to a hysteretic DC SQUID acting as an ancilla device as shown in the inset of Fig. 1. The ancilla is biased close to its critical current via a short current pulse. The interaction between the two loops is given by $H_I = MI_p \sigma_z J$ where I_p is the current circulating in the qubit, J the current in the ancilla and M the mutual inductance coefficient. At this point the ancilla has a certain probability of switching to the dissipative state, depending on the value of σ_z , hence on the qubit state. We propose to measure the output voltage of the ancilla SQUID, which hence is intimately connected to the qubit state. If the ancilla relaxation time is shorter than the, so called, discrimination time (for which the signal to noise ratio is close to

unity [32]), one cannot determine with certainty which is the output voltage, then σ_z . Therefore one may consider the value of the voltage as distributed according to the following probability distribution $P(V)$ as in [33]:

$$P(V) = \rho_{Q\downarrow\downarrow} P_-(V) + \rho_{Q\uparrow\uparrow} P_+(V). \quad (6)$$

Here V is binormally distributed, peaked around the two values it could assume in the case of projective measurements, say V_0 and \bar{V} (P_{\pm} are the gaussian distributions centred around the two voltage states, V_0 and \bar{V}). Then the qubit state σ_z can be derived from the output V and will be distributed according to $P(\sigma_z)$ as well, peaked around $(-1,1)$, with a variance D proportional to the ratio between the the ancilla relaxation time and the discrimination time. The quantities $\rho_{Q\downarrow\downarrow}$ and $\rho_{Q\uparrow\uparrow}$ are the diagonal elements of the qubit density matrix ρ_Q in the computational basis ($|\uparrow\rangle, |\downarrow\rangle$).

Within this approach, it is also possible to calculate how measuring the ancilla influences the qubit. Indeed, the update rule of the density matrix ρ_Q becomes very simple (see Supplementary Information) and it is evident that the shorter is the pulse, the less the system is perturbed. The density matrix ρ'_Q after the measurement is related to the one before the measurement ρ_Q by the update rule [34, 35]:

$$\rho'_Q = \frac{1}{\rho_{Q\downarrow\downarrow} e^{\gamma} + \rho_{Q\uparrow\uparrow} e^{-\gamma}} \begin{pmatrix} \rho_{Q\downarrow\downarrow} e^{\gamma} & \rho_{Q\downarrow\uparrow} \\ \rho_{Q\uparrow\downarrow}^* & \rho_{Q\uparrow\uparrow} e^{-\gamma} \end{pmatrix}, \quad (7)$$

where $\gamma = \sigma_z(t)/D$.

In order to extract meaningful information on the qubit, one must repeat the same evolution many times and evaluate $\sigma_z(t)$ as the average of the different results.

This approach allows us to simultaneously measure spin-spin correlation functions, with negligible effect on the annealing dynamics, hence on the residual energy as it will be clearer in the following (detail on our weak measurement scheme in Supplementary Information).

Results and Discussion

In our model hamiltonian, we choose $t_f = 14$ as annealing time. In the unitary limit, this guarantees that the adiabatic condition is fully satisfied. Adiabaticity is evident by analysing the behaviour of the residual energy as a function of t/t_f at fixed $t_f = 14$ and the corresponding ground state population (see Supplementary Information Fig. 1 bottom left panel). At the final time the residual energy is very close to zero $\epsilon_{res} = 5.49 \times 10^{-4}$ and the fidelity (ground state population at the annealing time) is $\rho_{Q\downarrow\downarrow}(t_f) = 0.999$, when no spin-spin correlation function is "measured" during the annealing dynamics, in agreement with Ref. [27].

As the Hamiltonian is time dependent, we expect that $C_{i,j}$ will have a generic dependance on the two times (t_i, t_j) , $C_{i,j} = C_{i,j}(t_i, t_j)$. However in the following we study the LGI as a function of $\tau = t_2 - t_1 = t_3 - t_2$, because maximum violation is expected (in the absence

of dissipative baths) when the three times are equally spaced.

When evaluating $C_{i,j}$ we perform two weak measurements per annealing and then calculate their expectation values averaging over N repeated dynamics. The larger the variance D , the less the system is affected by the measurement, and the closer is the residual energy to its unperturbed value.

To analyse how the measurement procedure affects the dynamics, in Fig. 1 we show the residual energy for the annealing Hamiltonian of Eq.(1), with no interaction with the environment, obtained for different choices of the variance D , at different values of τ . The results are presented repeating the evolution $N = 10^6$ times.

The residual energy ϵ_{res} at $D = 20$ ranges from 7.0×10^{-3} to 2.5×10^{-2} depending on τ . Correspondingly, the fidelity ranges from 0.973 to 0.993. That makes us comfortable about the convergence of the annealing procedure towards the target state, hence we choose $D = 20$ for all the following simulations.

In Fig.2 we show the Leggett-Garg's functions during the annealing dynamics in the unitary case, to be compared with the outcomes in the presence of a decoherence bath. The bold lines are obtained in the case of projective measurements, while the dots represent weak measurement results. While in a single "weak" measurement procedure, occasional violations may be driven by the system-detector interaction, averaging over N repeated measures, assures the convergence to the projective results. The agreement between the correlation functions calculated by means of projective and weak measurements demonstrate that our method reproduce the same correlations without affecting in a significant way the residual energy. Hence the quantum computation can be successful even during the LGI testing.

We clearly see that all the Leggett-Garg's functions go beyond the unitary "classical" limit. K_3^b and K_3^c are violated at intermediate times while K_3^a for short and long times. Remarkably, at short times, K_3^a and K_3^b show complementary violations, as well as in the case of a single qubit oscillating between its two states, at its own frequency (without annealing) [26], that is possibly due to the fact that, at short times, the qubit hamiltonian is "mostly" proportional to σ_x [25]. Moreover, at $\tau = t_f/2 = 7$ there is a significant violation of K_3^a . This value corresponds to a final measurement time $t_3 = t_f = 14$. Hence a violation of K_3^a at $\tau = t_f/2$, is particularly relevant, because it is a direct proof of quantum coherence until the end of the annealing dynamics. Hence in the following we will focus on the long time ($\tau \sim t_f/2$) behavior of K_3^a .

Now we move to discuss the same results in the presence of the dissipative environment. Hence, we turn on the system bath interaction and again calculate the LGIs fixing $D = 20$, $t_f = 14$. The value of K_3^a drops below

the classical bound for the LGIs at the final time both increasing the temperature (see Fig. 3 a) and increasing the coupling α (see Fig. 3 b). That corresponds to a lost quantum coherence at the annealing time t_f .

It is evident that the temperature plays a key role in the detrimental effect of the thermal bath. For low temperatures the quantum behavior persists during the whole evolution even in the presence a finite of coupling with the environment. By contrast, increasing the temperature, the time during which the system shows quantum features decreases, eventually going to zero for very high temperatures.

Since we have used a master equation in the Lindblad form, we must consider that this approach guarantees reliable results only in the weak coupling limit. Therefore the results shown for very high temperatures, and strong coupling, might be beyond our approximation and have to be considered with care. Moreover, at high temperatures, also the description of a SQUID flux-qubit as a two level system ceases to be correct. Hence we focus on results at low temperatures ($\beta = 10$) and weak-intermediate couplings. In this case (Fig. 3b)) the K_3^a function violate the unitary limit at long times from $\alpha = 0$ up to $\alpha = 10^{-3}$. However increasing the coupling to $\alpha = 10^{-2}$ the value of K_3^a is very close to its "classical" upper bound. Hence, in this case, we decided to simulate the system also using a classical Langevin dynamics to assess the LG functions in the absence of quantum correlations, as a comparison case (see Supplementary Information). Results are presented in the bottom panels of Figs 3c), 3d). First of all we notice that, as expected, the LG functions never violates their classical upper bounds (dashed line). Interestingly enough, the high temperature results (see Fig.s 3a) and c)) show agreement between classical and quantum dynamics. Of course we cannot make strong claims comparing Langevin and Lindblad dynamics, however this result points to poor quantum correlations in the Lindblad dynamics at these couplings. In the case of low temperatures (see Fig.s 3b) and d)), at weak coupling we show a completely different behavior of the K_3^a that can be hence used as quantum estimator. At intermediate couplings, quantum dynamics does not allow for full violation of LGI, nonetheless, the behavior of K_3^a show remarkable and qualitative differences with respect to the classical case. In particular we notice that the ordering of the curves as a function of the coupling at the annealing time (corresponding to $\tau = 7$) is reversed between the quantum and the classical case, except for the cases $\alpha = 0, 10^{-3}$. Hence, even in these "border line" cases, the evaluation of LGI's could be relevant to asses whether the dynamics has occurred via a quantum or a classical path. A comparative analysis of long time dynamics is shown in the Supplementary Information.

In conclusion, the presence of a dissipative environ-




FIG. 3. Plot of the Leggett-Garg's function K_3^a during the annealing dynamics. The black dashed line highlights the upper bound for the LGIs. The LG's functions are plotted as a function of the difference of the times at which the measurements are performed: $t_2 - t_1 = t_3 - t_2 = \tau$ (in units of \hbar/Γ_x with $\hbar = 1$). The time τ goes from 0 to $t_f/2$ so that it scans the whole evolution ($t_f = 14$). Top panels present results of quantum simulations, bottom panels of classical Langevin dynamics described in the Supplementary Information. In a) and c) we set $\alpha = 10^{-3}$, in b) and d) $\beta = 10$.

ment can modify the dynamics of a quantum two level system during the QA. In general, it is detrimental even if, under particular circumstances, it may also improve annealing performances [36]. This does not forbid, in principle, to successfully reach the target state at the annealing time. The question whether the dynamics has driven the system through a quantum or a classical path, up to now, remained unanswered and only partially addressed in the literature.

In this paper we propose a new, powerful and experimentally accessible method to assess the quantumness of a system during its adiabatic evolution, based on the LGI that we evaluate in the framework of weak measurements.

It allows us to show that time correlations can be measured without perturbing the annealing dynamics and that LGIs hold information about the interaction with the environment and can be used as witness of quantum coherence.

Do our results allow us to answer the point raised at the beginning of this paper? Namely: are real annealers, claimed to work performing adiabatic quantum computation, really quantum annealers? Are their outcomes macroscopic manifestations of quantum mechanics?

Our results show, for a very simple model, that if one

measures the LGIs along the adiabatic dynamics, a possible, yet non trivial, outcome could be that $K_3^a(t_f/2)$ contains all the information we need to assess if the final result of the computation is quantum or not. In the case of long time violation $K_3^a(t_f/2) > 1$ the result has to be considered as quantum, even in the presence of a dissipative bath. In borderline cases $K_3^a(t_f/2) \sim 1$, a careful analysis of the K_3^a behaviour as a function of time, could unveil the characteristics of the dynamics, namely if it was quantum, classical, or due to a non trivial occurrence of quantum and classical mechanisms. However this approach is still at its infancy. Extending it to more complicated ensembles like N spin Ising chains would be a fascinating way along which to proceed.

Data availability

The data that support the plots within this paper and other findings of this study are available from the corresponding author upon request.

Competing interests

The authors have no potential financial or non-financial conflicts of interest.

- ^a procolo.lucignano@spin.cnr.it
- [1] T. Albash and D. A. Lidar, “Adiabatic quantum computing,” *arXiv preprint arXiv:1611.04471*, 2016.
 - [2] M. A. Nielsen and I. Chuang, *Quantum Computation and Quantum Information*. Cambridge University Press, 2000.
 - [3] T. Kadowaki and H. Nishimori, “Quantum annealing in the transverse ising model,” *Phys. Rev. E*, vol. 58, pp. 5355–5363, 1998. [Online]. Available: <http://link.aps.org/doi/10.1103/PhysRevE.58.5355>
 - [4] M. W. Johnson, P. Bunyk, F. Maibaum, E. Tolkacheva, A. J. Berkley, E. M. Chapple, R. Harris, J. Johansson, T. Lanting, I. Perminov, E. Ladizinsky, T. Oh, and G. Rose, “A scalable control system for a superconducting adiabatic quantum optimization processor,” *Superconductor Science and Technology*, vol. 23, no. 6, p. 065004, 2010. [Online]. Available: <http://stacks.iop.org/0953-2048/23/i=6/a=065004>
 - [5] —, “A scalable control system for a superconducting adiabatic quantum optimization processor,” *Superconductor Science and Technology*, vol. 23, no. 6, p. 065004, 2010. [Online]. Available: <http://stacks.iop.org/0953-2048/23/i=6/a=065004>
 - [6] R. Harris, M. W. Johnson, T. Lanting, A. J. Berkley, J. Johansson, P. Bunyk, E. Tolkacheva, E. Ladizinsky, N. Ladizinsky, T. Oh, F. Cioata, I. Perminov, P. Spear, C. Enderud, C. Rich, S. Uchaikin, M. C. Thom, E. M. Chapple, J. Wang, B. Wilson, M. H. S. Amin, N. Dickson, K. Karimi, B. Macready, C. J. S. Truncik, and G. Rose, “Experimental investigation of an eight-qubit unit cell in a superconducting optimization processor,” *Phys. Rev. B*, vol. 82, p. 024511, Jul 2010. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevB.82.024511>
 - [7] J. King, S. Yarkoni, M. M. Nevisi, J. P. Hilton, and C. C. McGeoch, “Benchmarking a quantum annealing processor with the time-to-target metric,” *ArXiv e-prints*, Aug. 2015.
 - [8] Y. Aharonov, D. Z. Albert, and L. Vaidman, “How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100,” *Phys. Rev. Lett.*, vol. 60, pp. 1351–1354, Apr 1988. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevLett.60.1351>
 - [9] T. Albash, S. Boixo, D. A. Lidar, and P. Zanardi, “Quantum adiabatic markovian master equations,” *New Journal of Physics*, vol. 14, no. 12, p. 123016, 2012. [Online]. Available: <http://stacks.iop.org/1367-2630/14/i=12/a=123016>
 - [10] A. Das and B. K. Chakrabarti, “Colloquium,” *Rev. Mod. Phys.*, vol. 80, pp. 1061–1081, Sep 2008. [Online]. Available: <https://link.aps.org/doi/10.1103/RevModPhys.80.1061>
 - [11] A. Lucas, “Ising formulations of many np problems,” *Front. Phys.*, vol. 2, p. 5, 2014. [Online]. Available: <http://journal.frontiersin.org/article/10.3389/fphy.2014.00005>
 - [12] P. A. Lee, N. Nagaosa, and X.-G. Wen, “Doping a mott insulator: Physics of high-temperature superconductivity,” *Reviews of modern physics*, vol. 78, no. 1, p. 17, 2006.
 - [13] F. Barahona, “On the computational complexity of ising spin glass models,” *Journal of Physics A: Mathematical and General*, vol. 15, no. 10, p. 3241, 1982.
 - [14] E. Farhi, J. Goldstone, S. Gutmann, and M. Sipser, “Quantum computation by adiabatic evolution,” *arXiv preprint arXiv:quant-ph/0001106*, 2000.
 - [15] G. E. Santoro, R. Martoňák, E. Tosatti, and R. Car, “Theory of quantum annealing of an ising spin glass,” *Science*, vol. 295, no. 5564, pp. 2427–2430, 2002. [Online]. Available: <http://science.sciencemag.org/content/295/5564/2427>
 - [16] R. Martoňák, G. E. Santoro, and E. Tosatti, “Quantum annealing of the traveling-salesman problem,” *Physical Review E*, vol. 70, no. 5, p. 057701, 2004. [Online]. Available: <https://journals.aps.org/pre/abstract/10.1103/PhysRevE.70.057701>
 - [17] R. Harris, A. J. Berkley, J. Johansson, P. Bunyk, E. M. Chapple, C. Enderud, J. P. Hilton, K. Karimi, E. Ladizinsky, N. Ladizinsky, T. Oh, I. Perminov, C. Rich, M. C. Thom, E. Tolkacheva, C. J. S. Truncik, S. Uchaikin, J. Wang, B. Wilson, and G. Rose, “Quantum annealing with manufactured spins,” *Nature*, vol. 473, pp. 194–198, 2011. [Online]. Available: <http://dx.doi.org/10.1038/nature10012>
 - [18] T. F. Rønnow, Z. Wang, J. Job, S. Boixo, S. V. Isakov, D. Wecker, J. M. Martinis, D. A. Lidar, and M. Troyer, “Defining and detecting quantum speedup,” *Science*, vol. 345, no. 6195, pp. 420–424, 2014.
 - [19] S. Boixo, T. Albash, F. M. Spedalieri, N. Chancellor, and D. A. Lidar, “Experimental signature of programmable quantum annealing,” *Nature communications*, no. 4, p. 2067, 2013.
 - [20] S. Boixo, T. F. Rønnow, S. V. Isakov, Z. Wang, D. Wecker, D. A. Lidar, J. M. Martinis, and M. Troyer, “Evidence for quantum annealing with more than one hundred qubits,” *Nature physics*, vol. 10, pp. 218–224, 2014.
 - [21] S. W. Shin, G. Smith, J. A. Smolin, and U. Vazirani, “How ‘quantum’ is the d-wave machine?” *arXiv preprint arXiv:1401.7087*, 2014.
 - [22] P. Hauke, L. Bonnes, M. Heyl, and W. Lechner, “Probing entanglement in adiabatic quantum optimization with trapped ions,” *Frontiers in Physics*, vol. 3, p. 21, 2015. [Online]. Available: <https://www.frontiersin.org/article/10.3389/fphy.2015.00021>
 - [23] A. W. Glaetzle, R. M. W. van Bijnen, P. Zoller, and W. Lechner, “A coherent quantum annealer with rydberg atoms,” *Nature Communications*, vol. 8, pp. 15813 EP –, Jun 2017, article. [Online]. Available: <https://doi.org/10.1038/ncomms15813>
 - [24] A. J. Leggett and A. Garg, “Quantum mechanics versus macroscopic realism: Is the flux there when nobody looks?” *Phys. Rev. Lett.*, vol. 54, pp. 857–860, Mar 1985. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevLett.54.857>
 - [25] A. Friedenberger and E. Lutz, “Assessing the quantumness of a damped two-level system,” *Phys. Rev. A*, vol. 95, p. 022101, Feb 2017. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevA.95.022101>
 - [26] C. Emary, N. Lambert, and F. Nori, “Leggett-garg inequalities,” *Reports on Progress in Physics*, vol. 77, no. 1, p. 016001, 2014. [Online]. Available: <http://stacks.iop.org/0034-4885/77/i=1/a=016001>
 - [27] T. Albash and D. A. Lidar, “Decoherence in adiabatic quantum computation,” *Phys. Rev. A*, vol. 91, p.

- 062320, 2015. [Online]. Available: <http://link.aps.org/doi/10.1103/PhysRevA.91.062320>
- [28] A. J. Leggett, S. Chakravarty, A. Dorsey, M. P. Fisher, A. Garg, and W. Zwerger, “Dynamics of the dissipative two-state system,” *Reviews of Modern Physics*, vol. 59, no. 1, p. 1, 1987.
- [29] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems*. OUP Oxford, 2007.
- [30] S. Morita and H. Nishimori, “Mathematical foundation of quantum annealing,” *Journal of Mathematical Physics*, vol. 49, no. 12, p. 125210, 2008. [Online]. Available: <http://aip.scitation.org/doi/abs/10.1063/1.2995837>
- [31] T. Picot, R. Schouten, C. Harmans, and J. Mooij, “Quantum nondemolition measurement of a superconducting qubit in the weakly projective regime,” *Physical review letters*, vol. 105, no. 4, p. 040506, 2010.
- [32] A. Lupascu, C. Verwijs, R. Schouten, C. Harmans, and J. Mooij, “Nondestructive readout for a superconducting flux qubit,” *Physical review letters*, vol. 93, no. 17, p. 177006, 2004.
- [33] N. S. Williams and A. N. Jordan, “Weak values and the leggett-garg inequality in solid-state qubits,” *Phys. Rev. Lett.*, vol. 100, p. 026804, Jan 2008. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevLett.100.026804>
- [34] A. N. Jordan and A. N. Korotkov, “Qubit feedback and control with kicked quantum nondemolition measurements: A quantum bayesian analysis,” *Physical Review B*, vol. 74, no. 8, p. 085307, 2006.
- [35] A. N. Jordan, A. N. Korotkov, and M. Büttiker, “Leggett-garg inequality with a kicked quantum pump,” *Phys. Rev. Lett.*, vol. 97, p. 026805, Jul 2006. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevLett.97.026805>
- [36] G. Passarelli, G. De Filippis, V. Cataudella, and P. Lucignano, “Dissipative environment may improve the quantum annealing performances of the ferromagnetic p -spin model,” *Phys. Rev. A*, vol. 97, p. 022319, Feb 2018. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevA.97.022319>

Acknowledgments

The authors acknowledge stimulating discussions with Pino Falci, Rosario Fazio and Giuseppe E. Santoro as well as Gianluca Passarelli for his invaluable support in optimizing the performances of the Lindblad code.

Author contributions

GD, VC, PL, AT set the theoretical framework. PL and VV wrote the Lindblad code to weakly measure LGI during the QA. VV performed the quantum simulations. AD performed the classical-quantum mapping and the classical simulations. PL wrote the paper, with help from all co-authors.

Additional Information

Supplementary information accompanies this paper. Reprints and permissions information is available online. Correspondence and requests for materials should be addressed to P.L.