

Optimal stopping of oscillating Brownian motion

Ernesto Mordecki* and Paavo Salminen†

April 1, 2024

Abstract

We solve optimal stopping problems for an oscillating Brownian motion, i.e. a diffusion with positive piecewise constant volatility changing at the point $x = 0$. Let σ_1 and σ_2 denote the volatilities on the negative and positive half-lines, respectively. Our main result is that continuation region of the optimal stopping problem with reward $((1+x)^+)^2$ is disconnected, if and only if $\sigma_1^2 < \sigma_2^2 < 2\sigma_1^2$. Based on the fact that the skew Brownian motion in natural scale is an oscillating Brownian motion, the obtained results are translated into corresponding results for the skew Brownian motion.

AMS Subject Classification: 60J60, 60J65, 62L15

Keywords: excessive function, integral representation of excessive functions.

1 Introduction

The optimal stopping problems of diffusions with exceptional points has attracted interest in recent years. These include cases where the underlying diffusion has sticky points, skew points, or discontinuities in the diffusion coefficients. One of the first findings is that in the presence of sticky points the classical smooth fit principle does not necessarily hold, even for differentiable payoff functions (as found in Croce and Mordecki [2], and Salminen and Ta [15]). A second finding is that if the diffusion has a skew point, it can be the case that this point is in the continuation region for all discount values, as found by Alvarez and Salminen [1] and Presman [12]. A third one is that the continuation region in these cases can be disconnected, as observed in [1] for the skew Brownian motion, and also found recently by Mordecki and Salminen [9] for a diffusion with discontinuous drift and payoff function is $(1+x)^+$. General verification results for diffusions with discontinuous coefficients were obtained by

*Universidad de la República, Facultad de Ciencias, Centro de Matemática, Iguá 4225, 11400, Montevideo, Uruguay, e-mail: mordecki@cmat.edu.uy

†Åbo Akademi University, Faculty of Science and Engineering, FIN-20500 Åbo, Finland, e-mail: phsalmin@abo.fi.

Rüschendorf and Urusov [13]. An exposition of the general theory of optimal stopping (including historical comments) can be found in Shiryaev [16] and Peskir and Shiryaev [11].

In this paper the focus is on the case when the underlying diffusion has discontinuous infinitesimal variance. We then consider the optimal stopping problem for the oscillating Brownian motion (OBM), a diffusion with positive piecewise constant volatility changing at the origin. For details and further results on OBM, see Keilson and Wellner [5], Lejay and Pigato [7], and the references therein. Our main results are the following: Firstly, for the payoff $(1+x)^+$ the solution of the optimal stopping problem for the OBM is one sided for all values of the parameters, but for the payoff $((1+x)^+)^2$ the continuation region is disconnected for some values of the parameters. Hence, this latter situation is similar as the one in [9]. Secondly, based on the fact that the skew Brownian motion (SBM) in natural scale is an OBM, we obtain a result that connects the solutions of the respective optimal stopping problems for SBM and OBM, finding that the non-differentiability of the scale function of SBM at the origin plays a key rôle in understanding some of the phenomena that appear in the solutions of these problems.

2 Diffusions and optimal stopping.

Consider a conservative and regular one-dimensional (or linear) diffusion $X = (X_t)_{t \geq 0}$ taking values in $\mathbb{R}$, in the sense of Itô and McKean [4] (see also Borodin and Salminen [3]). Let $\mathbb{P}_x$ and $\mathbb{E}_x$ denote the probability and the expectation associated with X when starting from x , respectively; m denotes the speed measure and S the scale function. For $r \geq 0$ let φ_r (ψ_r) be the decreasing (increasing) positive fundamental solution of the generalized ODE

$$\frac{d}{dm} \frac{d}{dS} u = ru, \quad (1)$$

satisfying the appropriate boundary conditions (see [3] II.10 p.18). Denote by $\mathcal{M}$ the set of all stopping times in the filtration $(\mathcal{F}_t)_{t \geq 0}$, the usual augmentation of the natural filtration generated by X . Given a continuous reward function $g: \mathbb{R} \rightarrow [0, \infty)$ and a discount factor $r \geq 0$, consider the optimal stopping problem consisting of finding a function V_r and a stopping time $\tau^* \in \mathcal{M}$, such that

$$V_r(x) = \mathbb{E}_x[e^{-r\tau^*} g(X_{\tau^*})] = \sup_{\tau \in \mathcal{M}} \mathbb{E}_x[e^{-r\tau} g(X_\tau)], \quad (2)$$

where on the set $\{\tau = \infty\}$

$$e^{-r\tau} g(X_\tau) := \limsup_{t \rightarrow \infty} e^{-rt} g(X_t).$$

The *value function* V_r and the *optimal stopping time* τ^* constitute the solution of the problem. The optimal stopping time τ^* in (2), can be characterized (see Theorem 3, Section 3.3 in [16]) as the first entrance time into the stopping region

$$\Gamma_r := \{x: V_r(x) = g(x)\}. \quad (3)$$

The set $C_r := \mathbb{R} \setminus \Gamma_r$ is called the continuation region.

Our main tools to solve the optimal stopping problem for OBM are the representation theory for excessive functions, and the following two results from the theory of optimal stopping. The first one (Theorem 1) –formulated here for a left boundary point of the stopping region– is the smooth fit theorem, proof of which can be found in [14] or [10]; the second one (Proposition 1) is a verification result, for the proof see Corollary on p. 124 in [16].

Theorem 1. *Let z be a left boundary point of Γ_r , i.e., $[z, z + \varepsilon_1) \subset \Gamma_r$ and $(z - \varepsilon_2, z) \subset C_r$ for some positive ε_1 and ε_2 . Assume that the reward function g and the fundamental solutions φ_r and ψ_r are differentiable at z . Then the value function V_r in (2) is differentiable at z and it holds $V_r'(z) = g'(z)$.*

Proposition 1. *Let $A \subset \mathbb{R}$ be a nonempty Borel subset of $\mathbb{R}$ and*

$$H_A := \inf\{t : X_t \in A\}.$$

Assume that the function

$$\widehat{V}(x) := \mathbb{E}_x \left[e^{-r H_A} g(X_{H_A}) \right]$$

is r -excessive and dominates g . Then $\widehat{V}$ coincides with the value function of OSP (2) and H_A is an optimal stopping time.

3 Oscillating Brownian motion

Consider the diffusion satisfying the stochastic differential equation

$$X_t = x + \int_0^t \sigma(X_s) W_s,$$

where

$$\sigma(x) = \begin{cases} \sigma_1, & x < 0, \\ \sigma_2, & x \geq 0, \end{cases}$$

and $(W_t)_{t \geq 0}$ is a standard Brownian motion. The diffusion X is called an oscillating Brownian motion (OBM). Notice that this process is in natural scale, i.e. the scale function is $S(x) = x$, and the speed measure is

$$m(dx) = \begin{cases} (2/\sigma_1^2)dx, & x < 0, \\ (2/\sigma_2^2)dx, & x > 0. \end{cases}$$

(by definition there is no mass at $x = 0$). Let

$$\lambda_1^\pm = \pm \frac{\sqrt{2r}}{\sigma_1}, \quad \lambda_2^\pm = \pm \frac{\sqrt{2r}}{\sigma_2}.$$

The decreasing fundamental solution is

$$\varphi_r(x) = \begin{cases} A_1 \exp(\lambda_1^- x) + A_2 \exp(\lambda_1^+ x), & x < 0, \\ \exp(\lambda_2^- x), & x \geq 0, \end{cases} \quad (4)$$

where the constants A_1 and A_2 are determined so that φ_r is continuous and differentiable at 0. Hence,

$$A_1 = \frac{\lambda_1^+ - \lambda_2^-}{\lambda_1^+ - \lambda_1^-} = \frac{1 + \sigma_1/\sigma_2}{2}, \quad A_2 = \frac{\lambda_2^- - \lambda_1^-}{\lambda_1^+ - \lambda_1^-} = \frac{1 - \sigma_1/\sigma_2}{2}.$$

Analogously, the increasing solution is

$$\psi_r(x) = \begin{cases} \exp(\lambda_1^+ x), & x < 0, \\ B_1 \exp(\lambda_2^+ x) + B_2 \exp(\lambda_2^- x), & x \geq 0, \end{cases} \quad (5)$$

with

$$B_1 = \frac{\lambda_1^+ - \lambda_2^-}{\lambda_2^+ - \lambda_2^-} = \frac{1 + \sigma_2/\sigma_1}{2}, \quad B_2 = \frac{\lambda_2^- - \lambda_1^-}{\lambda_2^+ - \lambda_2^-} = \frac{1 - \sigma_2/\sigma_1}{2}.$$

4 Optimal stopping of OBM

We first analyze the optimal stopping problem (2) for the diffusion introduced above and the reward function

$$g_1(x) = \begin{cases} 0, & x \leq -1, \\ 1 + x, & x > -1. \end{cases}$$

The following result shows that the solution of this problem is one sided.

Proposition 2. *Consider the OSP problem (2) with payoff g_1 . For all values of $r > 0$, σ_1 and σ_2 the continuation region is given by*

$$C_r = (-\infty, c),$$

where $c = c(r) > -1$ is the unique solution of the equation

$$\psi_r'(x)(1+x) - \psi_r(x) = 0. \quad (6)$$

Furthermore

$$2r \leq \sigma_1^2 \Rightarrow c(r) \leq 0. \quad (7)$$

Proof. To prove the first statement, consider the functions (cf. [14])

$$H_-(x) := \psi_r'(x)g(x) - \psi_r(x)g'(x) = \psi_r'(x)(1+x) - \psi_r(x), \quad (8)$$

$$H_+(x) := \varphi_r(x)g'(x) - \varphi_r'(x)g(x) = \varphi_r(x) - \varphi_r'(x)(1+x), \quad (9)$$

and their derivatives for $x > -1$ and $x \neq 0$ can be expressed as

$$\begin{aligned} H'_-(x) &= m(x) \frac{d}{dm} H_-(x) = m(x) \psi_r(x) \left(r(1+x) - \frac{d}{dm} \frac{d}{dx} (1+x) \right) \\ &= m(x) \psi_r(x) r(1+x), \end{aligned}$$

and, similarly,

$$H'_+(x) = -m(x) \varphi_r(x) r(1+x),$$

where it is used that φ_r and ψ_r solve (1). Observe now that the function H_- in (8) has a unique positive root, since for $x > -1$ the derivative is strictly positive, $H_-(-1) = -\psi_r(-1) < 0$, and $H_-(x) \rightarrow \infty$ as $x \rightarrow \infty$. Therefore, equation (6) has a unique solution as claimed. The rest of the proof is standard, see for instance [14] or the detailed proof of Proposition 3 below. Statement (7) follows since $H_-(0) = \sqrt{2r}/\sigma_1 - 1$. $\square$

Next we study the OSP (2) for OBM with $0 < \sigma_1 \leq \sigma_2$ and the reward function

$$g(x) = \begin{cases} (1+x)^2, & x > -1, \\ 0, & x \leq -1. \end{cases} \quad (10)$$

In this situation it is seen that, for some specific values of the parameters, the continuation region is disconnected. The approach to analyze this problem is similar as the one in [9].

Define now

$$\begin{aligned} G_-(x) &:= \psi'_r(x)g(x) - \psi_r(x)g'(x) \\ &= (1+x) (\psi'_r(x)(1+x) - 2\psi_r(x)), \end{aligned} \quad (11)$$

$$\begin{aligned} G_+(x) &:= \varphi_r(x)g'(x) - \varphi'_r(x)g(x) \\ &= 2\varphi_r(x)(1+x) - \varphi'_r(x)(1+x)^2. \end{aligned} \quad (12)$$

These functions are used below to verify the excessivity of the proposed value function. The derivatives for $x > -1$ and $x \neq 0$ are

$$G'_-(x) = m(x)\psi_r(x) \begin{cases} r(1+x)^2 - \sigma_1^2, & x < 0, \\ r(1+x)^2 - \sigma_2^2, & x > 0, \end{cases} \quad (13)$$

$$G'_+(x) = m(x)\varphi_r(x) \begin{cases} \sigma_1^2 - r(1+x)^2, & x < 0, \\ \sigma_2^2 - r(1+x)^2, & x > 0. \end{cases} \quad (14)$$




Figure 1: The sign of the derivative G'_- is ruled by the depicted above function $r(1+x)^2 - \sigma(x)^2$. Here the parameters are $r = 1.5$, $\sigma_1 = 1$, $\sigma_2 = 2$.

Proposition 3. *In case $0 < r \leq \sigma_1^2 \leq \sigma_2^2$ the continuation region for OSP (2) is given by*

$$C_r = (-\infty, c),$$

where $c = c(r)$ is the unique positive solution of the equation

$$\psi'_r(x)(1+x) - 2\psi_r(x) = 0, \quad x \geq -1. \quad (15)$$

Proof. We show first that equation (15) has a unique positive solution. For this consider for $x > -1$ the function G_- defined in (11), the claim is equivalent with the statement that G_- has a unique positive zero. In fact, we claim a bit more; namely that the function G_- attains the global minimum at $x_0 := \sigma_2/\sqrt{r} - 1 > 0$, is negative and decreasing for $x \leq x_0$, is increasing for $x > x_0$, and has, therefore, a unique zero. Analyzing G'_- as given in (13), it is straightforward to deduce, since $0 < r \leq \sigma_1^2 \leq \sigma_2^2$, the claimed properties of G_- .

Let

$$H_c := \inf\{t : X_t \geq c\},$$

where c is the unique solution of (15), and define

$$\widehat{V}(x) := \mathbb{E}_x[e^{-rH_c}g(X_{H_c})] = \begin{cases} \frac{\psi_r(x)}{\psi_r(c)}g(c), & x \leq c, \\ g(x), & x \geq c. \end{cases} \quad (16)$$

If $\widehat{V}$ is an r -excessive majorant of g it follows from Proposition 1 that $\widehat{V}$ is the value function of OSP (2). The excessivity can be checked with the method based on the representation theory of excessive functions (cf. [14] Section 3). This boils down to study for $x \neq -1$ the functions

$$\begin{aligned} I_V(x) &:= \psi'_r(x)\widehat{V}(x) - \psi_r(x)\widehat{V}'(x), \\ D_V(x) &:= \varphi_r(x)\widehat{V}'(x) - \widehat{V}(x)\varphi'_r(x). \end{aligned}$$

Clearly, $I_V(x) = 0$ for $x \leq c$ and increasing for $x > c$. Notice that $I_V = G_-$ on $[c, +\infty)$. Studying the derivative (w.r.t. the speed measure) of D_V it is easily seen that D_V is positive and decreasing to 0 on $[c, +\infty)$. Consequently, I_V and D_V induce a (probability) measure which represents $\widehat{V}$ proving that $\widehat{V}$ is r -excessive. To prove that $\widehat{V}$ is a majorant of g consider for $-1 < x < c$

$$\widehat{V}(x) \geq g(x) \quad \Leftrightarrow \quad \frac{\psi_r(x)}{g(x)} \geq \frac{\psi_r(c)}{g(c)}.$$

The right hand side inequality holds since the derivative of $\psi_r(x)/g(x)$ is $G_-(x)$ which is negative for $-1 < x < c$, as is shown above. $\square$

If the volatilities are close enough, the problem is one sided for all discount values. This is made precise in the next result.

Proposition 4. *In case $0 \leq \sigma_1^2 \leq \sigma_2^2 \leq 2\sigma_1^2$ the continuation region for the OSP (2) is given by*

$$C_r = (-\infty, c),$$

where $c = c(r)$ is the unique solution of equation (15). As r increases from 0 to $+\infty$, $c(r)$ decreases monotonically from $+\infty$ to -1 . In particular, $c(r) = 0$ for $r = 2\sigma_1^2$.

Proof. If $r \leq \sigma_1^2$ the statement is the same as in Proposition 3. We assume next that $r \geq \sigma_2^2$. The proof in this case is very similar to the proof of Proposition 3. It can be seen that G_- attains the global minimum at $x_1 := \sigma_1/\sqrt{r} - 1 < 0$, is negative and decreasing for $x \leq x_1$, is increasing

for $x > x_1$, and has, therefore, a unique zero. Consequently, this root can be taken to be an optimal stopping point $c = c(r)$ and the analogous function $\hat{V}$ as in (16) can be proved to be the value of OSP (2). Finally, assume $\sigma_1^2 < r < \sigma_2^2 \leq 2\sigma_1^2$. In this case, G_- has a local maximum at 0, which is negative since

$$G_-(0) = \psi'_r(0) - 2\psi_r(0) = \lambda_1^+ - 2 = \frac{\sqrt{2r}}{\sigma_1} - 2 \leq 0. \quad (17)$$

Clearly, $G_-(0) = 0$ (and then $c(r) = 0$) when $r = 2\sigma_1^2$. Hence, equation (15) has a unique positive root and the proof can be completed as in the previous cases. $\square$

Proposition 5. *Assume $0 \leq \sigma_1^2 \leq \sigma_2^2 \leq 2\sigma_1^2$. For $r \geq 2\sigma_1^2$ there exist A and B such that the function*

$$F(x) := \begin{cases} A \exp(\lambda_1^+ x) + B \exp(\lambda_1^- x), & x \leq 0, \\ (1+x)^2, & x \geq 0, \end{cases} \quad (18)$$

satisfies the principle of smooth fit at 0, i.e., $F'(0-) = F'(0+) = 2$. The function F is r -harmonic (and positive) on $(-\infty, 0)$ but not r -excessive if $r < \sigma_2^2$. For $r < 2\sigma_1^2$ the coefficient B is negative and the function $F(x) \rightarrow -\infty$ as $x \rightarrow -\infty$ (and the function is not r -excessive).

Proof. We study only the case $r = r_0 := 2\sigma_1^2$ and leave the details of the other cases to the reader. In this case $\lambda_1^+ = \sqrt{2r}/\sigma_1 = 1$, and, obviously,

$$F(x) := \begin{cases} 2e^x, & x \leq 0, \\ (1+x)^2, & x \geq 0, \end{cases}$$

satisfies smooth fit at 0. Consequently, F is r_0 -harmonic (and positive) on $(-\infty, 0)$ and it remains to prove that F is not r_0 -excessive. For this, consider the representing function (this corresponds G_- in (11))

$$x \mapsto \psi'_{r_0}(x)F(x) - \psi_{r_0}(x)F'(x).$$

The claim is that this function is not non-decreasing. Indeed, differentiate w.r.t. the speed measure to obtain

$$\begin{aligned} & \frac{d}{dm} \left(\psi'_{r_0}(x)F(x) - \psi_{r_0}(x)F'(x) \right) \\ &= F(x) \frac{d}{dm} \frac{d}{dx} \psi_{r_0}(x) - \psi_{r_0}(x) \frac{d}{dm} \frac{d}{dx} F(x) \\ &= \psi_{r_0}(x) \begin{cases} 0, & x < 0, \\ r_0(1+x)^2 - \sigma_2^2, & x > 0. \end{cases} \end{aligned}$$

Since $r_0 = 2\sigma_1^2 < \sigma_2^2$ this derivative is negative, e.g., for small positive x -values; therefore, F is not r_0 -excessive. $\square$

For the theorem to follow, which can be seen as our main result concerning OSP (2), we need the following technical result.

Lemma 1. Consider a family $\{h_r: \mathbb{R} \rightarrow [0, \infty); r \in I\}$ such that for each $r \in I \subset \mathbb{R}$ the function h_r is r -excessive. Assume that this family is dominated by a function $\hat{h}$ (i.e. $h_r \leq \hat{h}$) such that $\mathbb{E}_x(\hat{h}(X_t)) < \infty$ for all $t \geq 0$ and $x \in \mathbb{R}$. Then, if for some r_0 the limit

$$\lim_{r \rightarrow r_0} h_r(x) \rightarrow h_0(x),$$

exists for all $x \in \mathbb{R}$, the function h_0 is r_0 -excessive.

Proof. Consider

$$\begin{aligned} \mathbb{E}_x [e^{-r_0 t} h_0(X_t)] &= \mathbb{E}_x \left[\lim_{r \rightarrow r_0} e^{-rt} h_r(X_t) \right] = \lim_{r \rightarrow r_0} \mathbb{E}_x [e^{-rt} h_r(X_t)] \\ &\leq \lim_{r \rightarrow r_0} h_r(x) = h_0(x), \end{aligned}$$

where in the second step we use the dominated convergence theorem which is applicable since $e^{-rt} h_r(X_t) \leq \hat{h}(X_t)$. $\square$

The theorem below states that if $r \in (2\sigma_1^2, \sigma_2^2)$ but is close enough to σ_2^2 then C_r has a bubble (i.e. an isolated bounded interval in the continuation region). However, the bubble disappears when r becomes larger than σ_2^2 or tends to $2\sigma_1^2$.

Theorem 2. In case $0 < \sigma_1^2 < 2\sigma_1^2 < \sigma_2^2$ there exists $r_0 \in (2\sigma_1^2, \sigma_2^2)$ with the following properties:

- (a) If $r \in [r_0, \sigma_2^2)$ the continuation region is given by

$$C_r = (-\infty, c_1) \cup (c_2, c_3),$$

where $c_i = c_i(r)$, $i = 1, 2, 3$, are such that $-1 < c_1 \leq c_2 \leq 0 < c_3$. In particular, for $r = r_0$ it holds $c_1 = c_2 < 0$.

- (b) If $r \geq \sigma_2^2$ the continuation region is explicitly given by

$$C_r = (-\infty, c_-),$$

where

$$c_- = c_-(r) = \frac{2\sigma_1}{\sqrt{2r}} - 1 < 0,$$

i.e. c_- is the unique solution of (15).

- (c) If $r < r_0$ the continuation region is given by

$$C_r = (-\infty, c_+),$$

where $c_+ = c_+(r) > 0$ is the unique solution of (15).

Proof. The proof of (b) is as the proof of Proposition 4 when $r \geq \sigma_2^2$. Notice, however, that in the present case $c(r) < 0$ for all $r \geq \sigma_2^2$.

We consider next (c) in case $r \leq 2\sigma_1^2$. Studying G'_- and $G_-(0)$ in (13) and (17) respectively, it is seen, as in the proof of Proposition 3, that equation $G_-(x) = 0$ has for $r < 2\sigma_1^2$ one (and only one) root $\rho = \rho(r) > \sigma_2/\sqrt{r} - 1 > 0$. In case $r = 2\sigma_1^2$ there are two roots $\rho_1 = 0$ and $\rho_2 > \sigma_2/(\sqrt{2}\sigma_1) - 1 > 0$. Proceeding as in the proof of Proposition 3 it

is seen that the stopping region is as claimed with $c_+ = \rho$ if $r < 2\sigma_1^2$ and $c_+ = \rho_2$ if $r = 2\sigma_1^2$.

Finally we consider (a). Assume now that there does not exist a bubble for any $r \in [2\sigma_1^2, \sigma_2^2]$. Then for all $r \in [2\sigma_1^2, \sigma_2^2]$ we can find $c = c(r)$ such that $\Gamma_r = [c, +\infty)$. Knowing that $c(r) > 0$ for $r = 2\sigma_1^2$ and $c(r) < 0$ for $r = \sigma_2^2$ we remark first there does not exist r such that $c(r) = 0$. Indeed, by Theorem 1, the value should satisfy the smooth fit principle at 0 but from Proposition 5 we know that such functions are not r -excessive. Next, using $\Gamma_{r_1} \subseteq \Gamma_{r_2}$ for $r_1 < r_2$ (cf. Proposition 1 in [9]) it is seen that $r \mapsto c(r)$ is non-increasing, and has, hence, left and right limits. Consequently, there exists a unique point r_0 such that

$$\hat{c}_+ := \lim_{r \uparrow r_0} c(r) > 0 \quad \text{and} \quad \hat{c}_- := \lim_{r \downarrow r_0} c(r) < 0.$$

Under the assumption that there is no bubble the value function is of the form given in (16), i.e.,

$$\begin{aligned} V_r(x) &= \begin{cases} \psi_r(x) \frac{(1+c(r))^2}{\psi_r(c(r))}, & x \leq c(r), \\ (1+x)^2, & x \geq c(r). \end{cases} \\ &= \mathbb{E}_x(e^{-rH_c} (1 + X_{H_c})^2), \end{aligned} \quad (19)$$

where $H_c := \inf\{y : X_t \geq c(r)\}$. For $\hat{r}$ small enough, there exists an excessive majorant $h_{\hat{r}}$, so that we can apply Proposition 1, since $\mathbb{E}_x e^{-\hat{r}t} h_{\hat{r}}(X_t) \leq h(x) < \infty$. Then, letting in (19) $r \uparrow r_0$ yields an r_0 -excessive function which by Proposition 1 is the value of the corresponding OSP (2). Similarly, letting $r \downarrow r_0$ yields an r_0 -excessive function which should also be the value of the same OSP. However, the functions are clearly different and since the value is unique we have reached a contradiction showing that there exists at least one bubble, i.e. a bounded open interval $(x_1, x_2) \subseteq C_r$ with endpoints x_1 and x_2 in the stopping set (see [9]). Proceeding similarly as in Proposition 6 in [9], it can be seen that if there is a bubble, there is at most one bubble, and it contains the origin, completing the proof. $\square$

We end this section by considering the case $\sigma_1^2 \geq \sigma_2^2$ with quadratic reward (10), and show that the stopping region in this case is always one sided.

Proposition 6. *Consider the OSP problem (2) for OBM with $\sigma_1^2 \geq \sigma_2^2$, $r > 0$ and $g(x)$ in (10). For all values of $r > 0$, the continuation region for OSP (2) is given by*

$$C_r = (-\infty, c),$$

where $c = c(r) > -1$ is the unique solution of the equation (15). Furthermore

$$r \leq 2\sigma_1^2 \quad \Rightarrow \quad c(r) \leq 0.$$

Proof. The proof follows the general lines of [14] developed in the previous proofs of Propositions 2 and 3. Consider then the functions defined in (11) and (12), with their respective derivatives in (13) and (14). As $\sigma_1^2 \geq \sigma_2^2$, the derivative $G'_-(x)$ changes sign only once, from negative to positive.

Hence, equation (15) has only one root $c(r) > -1$, and the proof of (a) follows as claimed. For (b), notice that the root $c(r) = 0$ when $r = 2\sigma_1^2$, the result follows from the monotonicity of the function $c(r)$. This concludes the proof. $\square$

5 OSP for skew Brownian motion

Consider a SBM $(\hat{X}_t)_{t \geq 0}$ with index $\beta \in (0, 1)$ starting at $\hat{x} \in \mathbb{R}$ (see [4], [18], [6]). This diffusion can be characterized by scale function

$$\hat{S}(x) = \begin{cases} x/(2(1-\beta)), & x < 0, \\ x/(2\beta), & x \geq 0, \end{cases}$$

and speed measure

$$m(dx) = \begin{cases} 4(1-\beta)dx, & x < 0, \\ 4\beta dx, & x \geq 0. \end{cases}$$

It is known (see e.g. [7]) that $(\hat{S}(\hat{X}_t))_{t \geq 0} \stackrel{d}{=} (X_t)_{t \geq 0}$, i.e. the composition of SBM with its scale function has the same law as OBM with $\sigma_1 = 1/(2(1-\beta))$ and $\sigma_2 = 1/(2\beta)$ and starting point $x = \hat{S}(\hat{x})$. In other words, we can say that SBM in natural scale is OBM. We use this relationship to obtain conclusions about the OSP problem (2). Given a payoff function $g: \mathbb{R} \rightarrow [0, \infty)$ we introduce the payoff function

$$\hat{g}(x) = (g \circ \hat{S})(x). \quad (20)$$

Due to the fact that $\hat{S}$ is not differentiable at the origin, both functions g and $\hat{g}$ can not be differentiable at the origin. The next result connects the optimal stopping problems for OBM and SBM.

Proposition 7. *For $\beta \in (0, 1)$ the optimal stopping problem (2) for OBM X with parameters $\sigma_1 = 1/(2(1-\beta))$, $\sigma_2 = 1/(2\beta)$ and continuous reward g has value function $V(x)$ and stopping region Γ if and only if the optimal stopping problem for SBM $\hat{X}$ with index $\beta \in (0, 1)$ and reward $\hat{g}$ in (20) has stopping region $\hat{\Gamma} = \hat{S}^{-1}(\Gamma)$ and value function $\hat{V}(y) = V(\hat{S}(y))$.*

Proof. It holds

$$\begin{aligned} V(x) &= \sup_{\tau} \mathbb{E}_x (e^{-r\tau} g(X_\tau)) = \sup_{\tau} \mathbb{E}_x (e^{-r\tau} \hat{g}(\hat{S}^{-1}(X_\tau))) \\ &= \sup_{\tau} \mathbb{E}_{\hat{S}^{-1}(x)} (e^{-r\tau} \hat{g}(\hat{X}_\tau)) = \hat{V}(\hat{S}^{-1}(x)), \end{aligned}$$

and this yields $\hat{V}(y) = V(\hat{S}(y))$. Let $\hat{\Gamma} = \{y: \hat{g}(y) = \hat{V}(y)\}$, and consider

$$\begin{aligned} \hat{S}(\hat{\Gamma}) &= \{x: \exists y \in \hat{\Gamma} \text{ such that } x = \hat{S}(y)\} \\ &= \{x: \hat{g}(\hat{S}^{-1}(x)) = \hat{V}(\hat{S}^{-1}(x))\} \\ &= \{x: g(x) = V(x)\} = \Gamma. \end{aligned}$$

This concludes the proof. $\square$

From this proposition it follows that $0 \notin \Gamma$ if and only if $0 \notin \widehat{\Gamma}$, because $\widehat{S}(0) = 0$. Furthermore, Γ is disconnected if and only if $\widehat{\Gamma}$ is disconnected, as the function $\widehat{S}$ is strictly increasing and continuous.

Example. Consider the problem (2) for SBM with $\beta > 1/2$ and reward $\widehat{g}(x) = (1+x)^+$ (cf. [1]). The corresponding OSP for OBM has $\sigma_1 = 1/(2(1-\beta))$, $\sigma_2 = 1/(2\beta)$, and reward

$$g(x) = \begin{cases} (1 + 2(1 - \beta)x)^+, & x < 0, \\ 1 + 2\beta x, & x \geq 0. \end{cases} \quad (21)$$

Notice that $g'(0-) = 2(1 - \beta) < 2\beta = g'(0+)$ (see Figure 2). As the scale function of OBM is $S(x) = x$, any r -excessive function h satisfies (see p. 93 in [14])

$$h'(x-) \geq h'(x+), \text{ for all } x \in \mathbb{R}. \quad (22)$$

If $0 \in \Gamma$ for some $r \geq 0$, then $V(0) = g(0)$, hence $V'(0-) < V'(0+)$ violating condition (22). We conclude that $0 \notin \Gamma$ for any value of r , hence $0 \notin \widehat{\Gamma}$ for any value of r for the SBM problem. This is a particular case of the result obtained in Proposition 1 in [1].




Figure 2: The reward g in (21) for OBM, with $\sigma_1 = 2$ and $\sigma_2 = 2/3$ ($\beta = 3/4$).

Acknowledgement. This research has been partially supported by grants from Magnus Ehrnrooths stiftelse, Finland. The authors acknowledge also the hospitality of Åbo Akademi University, Turku-Åbo, Finland and Universidad de la República, Montevideo, Uruguay.

References

- [1] Alvarez E., L. and Salminen, P.: Timing in the Presence of Directional Predictability: Optimal Stopping of Skew Brownian Motion. *Mathematical Methods of Operations Research*, Vol. 86, pp. 377-400, 2017.

- [2] Croce, F., Mordecki, E. Explicit solutions in one-sided optimal stopping problems for one-dimensional diffusions. *Stochastics* Vol. 86(3), pp. 491–509, 2014.
- [3] Borodin, A.N. and Salminen, P.: *Handbook of Brownian motion—Facts and formulae*, 2nd. ed., 2nd corr. print., Birkhäuser Verlag, 2015.
- [4] Itô, K. and McKean, H. P.: *Diffusion Processes and their Sample Paths*, Springer-Verlag, Berlin, 1974.
- [5] Keilson, J., and Wellner, J. A.: Oscillating Brownian motion. *Journal of Applied Probability*, Vol. 15(2), pp. 300–310, 1978.
- [6] Lejay, A. On the constructions of the skew Brownian motion, *Probability Surveys*, Vol. 3, pp. 413–466, 2006.
- [7] Lejay, A. and Pigato, P. Statistical estimation of the Oscillating Brownian Motion. *Bernoulli*, Vol. 24(4B), pp. 3568–3602, 2018.
- [8] Mikhalevich, V. S. A Bayes test of two hypotheses concerning the mean of a normal process (in Ukrainian). *Visnik Kiiv. Univ.* 1, 101–104 1958.
- [9] Mordecki, E., Salminen, P. Optimal stopping of Brownian motion with broken drift. *ArXiv:1811.05738* (2019)
- [10] Peskir, G.: Principle of smooth fit and diffusions with angles. *Stochastics*, Vol. 79, pp. 293–302, 2007.
- [11] Peskir, G. and Shiryaev, A.N.: *Optimal Stopping and Free-Boundary Problems*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2006.
- [12] Presman E.: Solution of the optimal stopping problem of one-dimensional diffusion based on a modification of payoff function. *Honor of Yuri V. Prokhorov, Prokhorov and contemporary probability theory*. Springer Proceedings in Mathematics and Statistics, vol 33, pp 371–403. Springer Verlag, 2012,
- [13] Rüschendorf, L. and Urusov, M.A.: On a class of optimal stopping stopping problems for diffusions with discontinuous coefficients. *Annals of Applied Probability*, Vol. 18, pp. 847878, 2008.
- [14] Salminen, P.: Optimal stopping of one-dimensional diffusions. *Math. Nachr.* Vol. 124 , pp. 85–101, 1985
- [15] Salminen, P. and Ta, B. Q. : Differentiability of excessive functions of one-dimensional diffusions and the principle of smooth fit, *Banach Center Publications*, Vol. 104, pp. 181–199, 2015.
- [16] Shiryaev, A.N.: *Optimal Stopping Rules*. Springer-Verlag, New York, 1978.
- [17] Taylor, H. : Optimal stopping in a Markov process. *Ann. Math. Stat.* Vol. 39, pp. 1333-1344, 1968.
- [18] Walsh, J. B.: A diffusion with a discontinuous local time *Astérisque*, Vol. 52-53, pp. 37–45, 1978.