HILBERT SCHEMES, COMMUTING MATRICES, AND HYPERKÄHLER GEOMETRY

ROGER BIELAWSKI AND CAROLIN PETERNELL

ABSTRACT. We represent algebraic curves via commuting matrix polynomials. This allows us to show that the Hilbert scheme of cohomologically stable nonplanar curves of genus 0 and degree d in $\mathbb{P}^3 \setminus \mathbb{P}^1$ is isomorphic to a complexified hyperkähler quotient of an open subset of a vector space by a nonreductive Lie group.

It has been observed in [4] that the Hilbert scheme of real cohomologically stable (i.e. satisfying $h^0(\mathcal{N}(-2)) = 0$) nonplanar curves of fixed genus and degree in \mathbb{P}^3 , not intersecting a fixed real line, carries a natural pseudo-hyperkähler structure. Here "real" means invariant under a fixed point free antilinear involution of \mathbb{P}^3 . In the case of g = 0 and d = 3 this pseudo-hyperkähler structure was shown in [4] to be flat, and in fact, the manifold of cohomologically stable pure-dimensional Cohen-Macaulay nonplanar curves in $\mathbb{P}^3 \setminus \mathbb{P}^1$ with Hilbert polynomial 3n + 1 with its natural complexified hyperkähler structure was shown there to be isomorphic to $\mathbb{C}^{12} \simeq \mathbb{C}^3 \otimes \operatorname{Mat}_{2\times 2}(\mathbb{C})$. The proof of this relies on such curves being ACM and so clearly different methods are needed to study the pseudo-hyperkähler geometry of the corresponding open subset of Hilb_{d,g} for other values of d and g.

In the present article we present such a method via a correspondence between algebraic curves equipped with a flat projection onto \mathbb{P}^1 and commuting matrix polynomials. This correspondence allows us to describe the locus of cohomologically stable nonplanar curves of arithmetic genus 0 and degree d in $\mathbb{P}^3 \setminus \mathbb{P}^1$ as a complexified hyperkähler quotient of (an open subset of) a vector space by a *nonreductive* Lie group (Theorem 5.6). Formally, our moment map equations are very similar to the complex ADHM equations used by Frenkel and Jardim [10] in their construction of admissible torsion-free sheaves on \mathbb{P}^3 . The main difference is that the Lie group acting on solutions is no longer reductive.

Restricting this description to real curves, we obtain the above pseudo-hyperkähler structure for g = 0 and any odd d (for even d there are no real rational curves in the above sense) as a hyperkähler quotient of an open subset of a flat quaternionic vector space by a nonreductive Lie group. There is also an analogous description of the natural hypersymplectic structure on cohomologically stable nonplanar curves of genus 0 and any degree which are invariant under an antilinear involution of \mathbb{P}^3 , the fixed point set of which is \mathbb{RP}^3 .

We briefly describe the structure and the content of the paper. In the next section we provide alternative (to the one given by Nakajima [22]) descriptions of the Hilbert scheme $(\mathbb{C}^2)^{[n]}$ and of its open subset of non-collinear points. In §2 we discuss the above-mentioned correspondence between algebraic curves and commuting matrix polynomials. In §3 we restrict our attention to space curves, and show

²⁰¹⁰ Mathematics Subject Classification. 14C05, 14H50, 53C26, 53C28.

how to construct a twistor space $Z_d \to \mathbb{P}^1$, the sections of which can be identified with either a) algebraic curves of fixed degree, or b) equivalence classes of certain commuting pairs of polynomials. Section 4 discusses complexified hyperkähler and hypercomplex structures and their quotients. Finally, in §5 we apply these ideas and results to genus 0 space curves and show that in this case the description b) of sections of Z_d is equivalent to a complexified hyperkähler quotient.

Remark 0.1. Throughout the paper a *curve* means a projective Cohen-Macaulay scheme of pure dimension 1.

Acknowledgement. This work has been carried out while both authors were members of, and the second author was fully funded by the DFG Priority Programme 2026 "Geometry at infinity", the support of which is gratefully acknowledged. The authors also thank Michael Bulois and Israel Vainsencher for pointing out mistakes in an earlier version of the paper. Last, but not least, we are grateful to the anonymous referees for many helpful comments, which resulted in a greatly improved presentation.

1. Hilbert schemes of points in \mathbb{C}^2 and commuting matrices

We begin with the following easy observation (cf. [22] if k = 2 and [16, Thm. 2.5] for general k):

Proposition 1.1. There exists a natural set-theoretic bijection between the Hilbert scheme $(\mathbb{C}^k)^{[n]}$ of n points in \mathbb{C}^k and $GL(n, \mathbb{C})$ -orbits of k-tuples (A_1, \ldots, A_k) of $n \times n$ matrices such that $\mathbb{C}[A_1, \ldots, A_k]$ is an n-dimensional commutative algebra which is conjugate to its image in $\operatorname{End}(\mathbb{C}^n)$ under the regular representation.

Proof. A point in the Hilbert scheme $(\mathbb{C}^k)^{[n]}$ is defined by an ideal $I \subset \mathbb{C}[z_1, \ldots, z_k]$ of length n, i.e. such that $\dim \mathbb{C}[z_1, \ldots, z_k]/I = n$. Multiplication by z_i , $i = 1, \ldots, k$, defines an endomorphism A_i of $\mathbb{C}[z_1, \ldots, z_k]/I$. Clearly the A_i commute and $\dim \mathbb{C}[A_1, \ldots, A_k] = \dim \mathbb{C}[z_1, \ldots, z_k]/I = n$. Moreover, directly from the construction, $\mathbb{C}[A_1, \ldots, A_k]$ is conjugate to its image under the regular representation. Conversely, given k commuting matrices A_1, \ldots, A_k with $\dim \mathbb{C}[A_1, \ldots, A_k] = n$ we define a homomorphism $\mathbb{C}[z_1, \ldots, z_k] \to \mathbb{C}^n \simeq \mathbb{C}[A_1, \ldots, A_k]$ via $p(z_1, \ldots, z_k) \mapsto$ $p(A_1, \ldots, A_k)$. Clearly it is surjective and its kernel is an ideal of length n. Moreover this ideal does not change under simultaneous conjugation of A_1, \ldots, A_k , and so we obtain a well-defined point of $(\mathbb{C}^k)^{[n]}$. Since we assume that $\mathbb{C}[A_1, \ldots, A_k]$ is conjugate to its image under the regular representation, the two maps are inverse to each other.

Remark 1.2. The condition that the algebra $\mathcal{A} = \mathbb{C}[A_1, \ldots, A_k]$ is conjugate to its image under the regular representation is equivalent to the existence of a cyclic vector for \mathcal{A} . This is the same argument as in [22, p.8]. Also, the set of k-tuples (A_1, \ldots, A_k) of commuting $n \times n$ matrices with dim $\mathbb{C}[A_1, \ldots, A_k] = n$ and having a cyclic vector is open in the set of all commuting k-tuples. Indeed, due to the Cayley-Hamilton theorem, any element of $\mathbb{C}[A_1, \ldots, A_k]$ belongs to the linear span of $A_1^{i_1} \ldots A_k^{i_k}$ for $i_1, \ldots, i_k \leq n-1$. Thus the condition of not having cyclic vector is equivalent to dim $\langle A_1^{i_1} \ldots A_k^{i_k} v; i_1, \ldots, i_k \leq n-1 \rangle \leq n-1$ for any v. This is a closed condition. Henni and Jardim show that the quotient of the variety of k commuting matrices together with a choice of a cyclic vector by $GL(n, \mathbb{C})$ is a geometric quotient [16, §2], and that the resulting scheme is isomorphic to $(\mathbb{C}^k)^{[n]}$ [16, Cor. 4.9]. For k = 2, these results are due to Nakajima [22, §1.2].

We shall now modify this description of $(\mathbb{C}^2)^{[n]}$ in several ways. First of all, we shall want to eliminate the dependence on a cyclic vector. We recall the following theorem of Neubauer and Saltman [23]:

Theorem 1.3 (Neubauer-Saltman). Let A and B be two commuting $n \times n$ matrices. Then the following conditions are equivalent:

- (1) dim $\mathbb{C}[A, B] = n$.
- (2) dim Z(A, B) = n, where Z(A, B) is the centraliser of the pair (A, B).
- (3) (A, B) is a nonsingular point of the variety of commuting matrices.

The equivalence between (1) and (2) will be used repeatedly throughout the paper.

Let us write M_n for the variety of pairs of commuting $n \times n$ matrices and define

$$M_n^0 = \{ (A, B); A, B \in \operatorname{Mat}_{n \times n}(\mathbb{C}), [A, B] = 0, \dim Z(A, B) = n \},\$$

and its open subset M_n^{reg} consisting of those $(A, B) \in M_n^0$ for which $\mathbb{C}[A, B]$ is conjugate to its image in $\operatorname{Mat}_{n \times n}(\mathbb{C})$ under the regular representation.

Proposition 1.4. $(\mathbb{C}^2)^{[n]}$ is the geometric quotient of M_n^{reg} by $GL(n, \mathbb{C})$.

Proof. It is the matter of checking the conditions that a geometric quotient has to satisfy [21, Def. 0.6]. The only one which perhaps requires an argument is that the structure sheaf of $(\mathbb{C}^2)^{[n]}$ is equal to the $GL(n, \mathbb{C})$ -invariant part of the structure sheaf of M_n^{reg} . Nakajima [22] shows that the Hilbert scheme of n points in \mathbb{C}^2 is isomorphic to the geometric quotient of the (smooth) variety

$$H_n = \{(A, B, v); [A, B] = 0, v \text{ is a cyclic vector for } (A, B)\}$$

by the (free) action of $GL(n,\mathbb{C})$. Hence $\mathcal{O}_{(\mathbb{C}^2)^{[n]}} = (\mathcal{O}_{\tilde{H}_n})^{GL(n,\mathbb{C})}$.

We have the natural forgetful map $p: \tilde{H}_n \to M_n^{\text{reg}}, (A, B, v) \mapsto (A, B)$, and the proof of Theorem 1.9 in [22] shows that p is a submersion. Observe now that the fibre $p^{-1}(A, B)$ is isomorphic to the stabiliser of (A, B) in $GL(n, \mathbb{C})$. Indeed, the stabiliser is *n*-dimensional and it acts freely on the fibre. Since, owing to Proposition 1.1, the projection $\tilde{H}_n \to (\mathbb{C}^2)^{[n]}$ factors (set-theoretically) through M_n^{reg} , the action of the stabiliser on the fibre must be transitive. Therefore, for any open subset $U \subset M_n^{\text{reg}}, \mathcal{O}(U)^{GL(n,\mathbb{C})} = \mathcal{O}(p^{-1}(U))^{GL(n,\mathbb{C})}$, and so $\mathcal{O}_{M_n^{\text{reg}}} = (\mathcal{O}_{\tilde{H}_n})^{GL(n,\mathbb{C})}$.

We shall now present another description of $(\mathbb{C}^2)^{[n]}$: as a symplectic quotient of an open dense subset of pairs of matrices by a nonreductive group. The relevant subset K_n consists of pairs (A, B) of $n \times n$ matrices, such that the vector $e_1 =$ $(1, 0, \ldots, 0)^T$ is cyclic for the pair A, B. The group G is the subgroup of $GL(n, \mathbb{C})$ preserving the cyclic vector e_1 , i.e. the first column of elements of G is equal to e_1 . The symplectic form on K_n is tr $dA \wedge dB$, and the corresponding moment map $\mu: K_n \to \mathfrak{g}^*$ is the projection of [A, B] onto the last n - 1 rows.

Theorem 1.5. The action of G on $\mu^{-1}(0) \subset K_n$ is free and proper, and the symplectic quotient $\mu^{-1}(0)/G$ is biholomorphic to $(\mathbb{C}^2)^{[n]}$.

Proof. First of all, we claim that $\mu(A, B) = 0$ implies [A, B] = 0. Indeed, $\mu(A, B) = 0$ means that [A, B] can have nonzero entries only in the first row. Suppose that $[A, B] \neq 0$. Then $\text{Im}[A, B] = \langle e_1 \rangle$ is a 1-dimensional subspace which is cyclic for (A, B). On the other hand, according to [13] (see also [9, Lemma 12.7] for a simple proof) A and B can be simultaneously conjugated into upper-triangular matrices. But then any vector in Im[A, B] has the last coordinate equal to zero and since no such vector can be cyclic for a pair of upper-triangular matrices, we obtain a contradiction. Hence [A, B] = 0.

The action of G on $\mu^{-1}(0)$ is free, since the action of $GL(n, \mathbb{C})$ on \tilde{H}_n is free. This implies, in particular, that $\mu^{-1}(0)$ is smooth. To show that the action of G on $\mu^{-1}(0)$ is proper it is enough to show that the action of $GL(n, \mathbb{C})$ on \tilde{H}_n is proper, since K_n is closed in \tilde{H}_n and G is closed in $GL(n, \mathbb{C})$. The properness of the $GL(n, \mathbb{C})$ on \tilde{H}_n is equivalent to \tilde{H}_n being a principal $GL(n, \mathbb{C})$ -bundle over $(\mathbb{C}^2)^{[n]}$. Nakajima [22, Theorem 3.24 & Cor. 3.42] shows that $(\mathbb{C}^2)^{[n]}$ is a (real) symplectic quotient $\nu^{-1}(c)/U(n)$ of \tilde{H}_n , where ν denotes the moment map. Since U(n) is compact, $\nu^{-1}(c)$ is a principal U(n)-bundle over $(\mathbb{C}^2)^{[n]}$. This means that $\nu^{-1}(c) \to (\mathbb{C}^2)^{[n]}$ admits local sections. Such a local section gives a local section of $\tilde{H}_n \to (\mathbb{C}^2)^{[n]}$, and, consequently, \tilde{H}_n is a principal $GL(n, \mathbb{C})$ -bundle over $(\mathbb{C}^2)^{[n]}$. As explained above, this implies that the action of G on $\mu^{-1}(0)$ is proper. It follows that $\mu^{-1}(0)/G$ is a complex manifold [7, Ch.III, Prop.10], biholomorphic to $\tilde{H}_n/GL(n, \mathbb{C})$.

1.1. $(\mathbb{C}^2)^{[n]}$ and torsion-free sheaves on \mathbb{P}^2 . We shall now give a description of an open subset of $(\mathbb{C}^2)^{[n]}$ consisting of 0-dimensional subschemes not contained in any line (thus, necessarily, $n \geq 3$). It is closely related to the description of the moduli space $\mathcal{M}(2, n-1)$ of framed torsion-free sheaves on \mathbb{P}^2 of rank 2 and $c_2 = n-1$ in terms of the ADHM equations [22, Theorem 2.1]. In order to simplify the notation, set k = n - 1. The moduli space $\mathcal{M}(2, k)$ is biholomorphic to the $GL(k, \mathbb{C})$ -quotient of the set U of stable solutions to the equation

(1.1)
$$[X,Y] + ij = 0, \quad X,Y \in \operatorname{Mat}_{k,k}(\mathbb{C}), \ i \in \operatorname{Mat}_{k,2}(\mathbb{C}), \ j \in \operatorname{Mat}_{2,k}(\mathbb{C}).$$

Here a quadruple (X, Y, i, j) is called *stable* if there is no proper subspace S of \mathbb{C}^k such that $\operatorname{Im} i \subset S$, $XS \subset S$, $YS \subset S$, and the framing of a sheaf \mathcal{F} is a trivialisation on the line $l_{\infty} \subset \mathbb{P}^2$ (in particular $c_1 = 0$).

Lemma 1.6. Let $\mathcal{F} \in \mathcal{M}(2, k)$. The following conditions are equivalent:

- (i) the rank of *i* is equal to 1;
- (ii) $H^0(\mathbb{P}^2, \mathcal{F}) \neq 0;$
- (iii) \mathcal{F} is an extension of the form $0 \to \mathcal{O}_{\mathbb{P}^2} \to \mathcal{F} \to \mathcal{I}_Z \to 0$, where \mathcal{I}_Z is the ideal sheaf of $Z \in (\mathbb{P}^2 \setminus l_\infty)^{[k]}$.

Proof. Given $c_2(\mathcal{F}) = k$, the conditions (ii) and (iii) are clearly equivalent. We prove the equivalence of (i) and (iii). If *i* has rank 1, then $ij = i_0(\alpha_1 j_1 + \alpha_2 j_2)$ for some $i_0 \in \mathbb{C}^k$, $\alpha_1, \alpha_2 \in \mathbb{C}$, where j_1, j_2 are the two rows of *j*. Equation (1.1) implies that [X, Y] has rank 1, and therefore $\alpha_1 j_1 + \alpha_2 j_2 = 0$ [22, Prop. 2.8]. Thus *j* has also rank 1 and we can write $\mathbb{C}^2 \simeq W_1 \oplus W_2$, where $W_1 = \text{Im } j = \text{Ker } i$. The sheaf \mathcal{F} is then isomorphic to Ker b/Im a, where (a, b) is the monad given on p. 23 in [22]. The embedding $W_1 \otimes \mathcal{O}_{\mathbb{P}^2} \hookrightarrow \text{Ker } b$ induces an injective morphism $\mathcal{O}_{\mathbb{P}^2} \to \mathcal{F}$. Its cokernel is the sheaf obtained from the monad with W replaced by W_2 and j = 0. This is a rank one framed torsion free sheaf with $c_2 = k$, i.e. the ideal sheaf of a $Z \in (\mathbb{P}^2 \setminus l_\infty)^{[k]}$. Conversely, given \mathcal{F} as in the statement, we obtain from \mathcal{I}_Z a stable solution $(X, Y, i_0, 0)$ to the ADHM equation with $i_0 \in \mathbb{C}^k$. Let j_0^T be the extension class of \mathcal{F} in $\text{Ext}^1(\mathcal{I}_Z, \mathcal{O}_{\mathbb{P}^2}) \simeq \mathbb{C}^k$. Then $(X, Y, i \oplus 0, 0 \oplus j)$ is a solution to the ADHM equation which yields (the isomorphism class of) \mathcal{F} .

We denote by $\mathcal{M}(2,k)^{\circ}$ the complement of the (isomorphism classes of) sheaves described in this lemma. If (X, Y, i, j) is the ADHM-data corresponding to a point in $\mathcal{M}(2,k)^{\circ}$, then *i* has rank 2 and we can use the action of $GL(k, \mathbb{C})$ to fix *i* to be

(1.2)
$$i = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \end{pmatrix}^T$$

The stabiliser G_0 of *i* consists of matrices of the form

$$\begin{pmatrix} 1_{2\times 2} & * \\ 0 & * \end{pmatrix},$$

and $\mathcal{M}(2,k)^{\circ}$ is also biholomorphic to the quotient of the space U_0 of such (X,Y,i,j)by G_0 . Here "quotient" means that U_0 is a principal G_0 -bundle over $\mathcal{M}(2,k)^{\circ}$. Indeed, a well-known description of $\mathcal{M}(2,k)$ as a hyperkähler quotient means that U is a principal $GL(k, \mathbb{C})$ -bundle over $\mathcal{M}(2,k)$ (cf. the proof of Theorem 1.5) and, consequently, the action of $GL(k, \mathbb{C})$ on U is free and proper. This implies that the action of G_0 on U_0 is also free and proper, so that U_0 is a principal G_0 -bundle over $\mathcal{M}(2,k)^{\circ}$. In fact, j is now superfluous: it is simply given by the first two rows of [X,Y]. Equation (1.1) is now equivalent to the last k-2 rows of [X,Y]being identically zero. Since the projection onto the last k-2 rows is precisely the moment map μ for the action of G_0 on pairs of matrices X, Y with respect to the symplectic form tr $dX \wedge dY$, we conclude:

Proposition 1.7. Let V^s be the set of pairs $(X, Y) \in \operatorname{Mat}_{k,k}(\mathbb{C})^2$ such that (X, Y, i) is stable¹, where *i* is given by (1.2). The group G_0 acts freely and properly on $\mu^{-1}(0) \subset V^s$ and the symplectic quotient $\mu^{-1}(0)/G_0$ is biholomorphic to $\mathcal{M}(2, k)^o$. \Box

Remark 1.8. Clearly, there is an analogous description of an open dense subset of $\mathcal{M}(r,k)$ for sheaves of higher rank r.

Given a quadruple (X, Y, i, j) (not necessarily satisfying (1.1) or (1.2)), we define a pair of $(k + 1) \times (k + 1)$ -matrices as follows (cf. [5]):

(1.3)
$$\hat{X} = \begin{pmatrix} 0 & -j_2 \\ i_1 & X \end{pmatrix}, \quad \hat{Y} = \begin{pmatrix} 0 & j_1 \\ i_2 & Y \end{pmatrix},$$

where i_s (resp. j_s) denotes the s-th column (resp. s-th row) of i (resp. of j).

Lemma 1.9. (X, Y, i, j) satisfies (1.1) if and only if $[\hat{X}, \hat{Y}]$ has nonzero entries only in the first row or the first column. Moreover (X, Y, i, j) is stable if and only if e_1 is cyclic for (\hat{X}, \hat{Y}) .

Proof. The first statement is obvious. For the second one, observe that if S is a destabilising subspace for (X, Y, i, j), then $S \oplus \langle e_1 \rangle$ is invariant for (\hat{X}, \hat{Y}) . Conversely, suppose that \hat{S} contains e_1 and is invariant for (\hat{X}, \hat{Y}) . Let $p : \mathbb{C}^{k+1} \to \mathbb{C}^k =$

¹Stability does not depend on j.

 $\mathbb{C}^{k+1}/\langle e_1 \rangle$ be the projection and set $S = p(\hat{S})$. Then $i_1 = p(\hat{X}e_1)$, $i_2 = p(\hat{Y}e_1)$ belong to S. Moreover, if $v \in S$, then there is an $\alpha \in \mathbb{C}$ such that $\alpha e_1 + v \in \hat{S}$. Since

$$\hat{X}(\alpha e_1 + v) = \alpha i_1 + Xv \mod \langle e_1 \rangle$$

and similarly for Y, it follows that $XS \subset S$ and $YS \subset S$. Therefore S is a destabilising subspace for (X, Y, i, j).

We now fix i to be (1.2). Let \hat{G}_0 denote the subgroup of $GL(k+1, \mathbb{C})$, consisting of matrices of the form

$$\begin{pmatrix} 1 & u \\ 0 & h \end{pmatrix}, \quad h \in G_0, \ u_1 = u_2 = 0,$$

i.e. matrices, the first 3 columns of which are (e_1, e_2, e_3) . \hat{G}_0 is a semidirect product of G_0 and \mathbb{C}^{k-2} . Its action by conjugation on \hat{X}, \hat{Y} defined in (1.3) is given by

$$\hat{X} \mapsto \begin{pmatrix} 0 & -j_2h^{-1} + uXh^{-1} \\ i_1 & -i_1uh^{-1} + hXh^{-1} \end{pmatrix}, \quad \hat{Y} \mapsto \begin{pmatrix} 0 & j_1h^{-1} + uYh^{-1} \\ i_2 & -i_2uh^{-1} + hYh^{-1} \end{pmatrix}.$$

This means that \hat{G}_0 acts on pairs of matrices X, Y via

(1.4)
$$(X,Y) \mapsto (-i_1 u h^{-1} + h X h^{-1}, -i_2 u h^{-1} + h Y h^{-1}).$$

Owing to Lemma 1.9, \hat{G}_0 preserves $V^{\rm s}$ of Proposition 1.7.

We consider the following codimension 2 subset of V^{s} :

$$V_0^{\rm s} = \{(X, Y) \in V^{\rm s}; X_{12} = Y_{11}, X_{22} = Y_{21}\}.$$

It is \hat{G}^0 -invariant and the restriction of the symplectic form $\operatorname{tr} dX \wedge dY$ to V_0^s is nondegenerate.

Theorem 1.10. Let $\hat{\mu}$ denote the moment map for the action of \hat{G}^0 on V_0^s . The action of \hat{G}^0 on $\hat{\mu}^{-1}(0)$ is free and proper and the symplectic quotient $\hat{\mu}^{-1}(0)/\hat{G}_0$ is biholomorphic to the open subset of $(\mathbb{C}^2)^{[k+1]}$ consisting of 0-dimensional subschemes not contained in any line.

Proof. The moment map for the action of $\hat{G}_0 \simeq G_0 \ltimes \mathbb{C}^{k-2}$ is equal to $\hat{\mu}(X,Y) = (\mu(X,Y), \alpha(X,Y))$, where

$$\alpha(X,Y) = (X_{32} - Y_{31}, \dots, X_{k2} - Y_{k1})^T.$$

Thus, given the definition of V_0^s , the condition $\alpha(X,Y) = 0$ means that the first column of Y is equal to the second column of X. This in turn implies that the first column of $[\hat{X}, \hat{Y}]$ is equal to 0. On the other hand $\mu(X, Y) = 0$ means that the last k - 2 rows of [X, Y] are zero. If we now take j_1 to be the first row of [Y, X] and j_2 the second row of [X, Y], then $[\hat{X}, \hat{Y}]$ has nonzero entries only in the first row. The argument in the proof of Theorem 1.5 together with Lemma 1.9 imply that $[\hat{X}, \hat{Y}] = 0$. Let K_{k+1}^0 be the set of pairs $(A, B) \in \text{Mat}_{k+1,k+1}(\mathbb{C})^2$ such that e_1 is cyclic for (A, B), the first column of A is equal to e_2 , and the first column of B is equal to e_3 . The above discussion shows that the map $V_0^s \to K_{k+1}^0$, sending (X, Y) to the matrices \hat{X}, \hat{Y} defined above, restricts to a \hat{G}_0 -equivariant isomorphism between $\hat{\mu}^{-1}(0)$ and $T = \{[A, B] \in K_{k+1}^0; [A, B] = 0\}$. The action of \hat{G}_0 on T is free and proper, owing to the same argument as in the proof of Theorem 1.5. It remains to show that T/\hat{G}_0 is biholomorphic to the open subset of $(\mathbb{C}^2)^{[k+1]}$ consisting of 0-dimensional subschemes not contained in any line. Let $H^0 \subset (\mathbb{C}^2)^{[k+1]}$ be the set of 0-dimensional subschemes not contained in any line. If $D \in H^0$ has ideal I, then $1, z_1, z_2$ are linearly independent elements of $\mathbb{C}[z_1, z_2]/I$. Therefore we can choose a basis $\mathbb{C}[z_1, z_2]/I$ such that $1, z_1, z_2$ are its first 3 elements. This means that the resulting commuting matrices A, B belong to K_n^0 . The group preserving $K_n^0 \subset K_n$ is precisely \hat{G}_0 , so that $H^0 \subset T/\hat{G}_0$. Conversely, let D be a 0-dimensional subscheme of \mathbb{C}^2 represented by an element of T/\hat{G}_0 . The ideal of D is given by polynomials $p(z_1, z_2)$ vanishing on (A, B). Given the form of the first column of A and of B, no linear polynomial vanishes on (A, B).

2. Curves and commuting matrices

We now wish to describe a correspondence between finite coverings of \mathbb{P}^1 and certain commuting matrix polynomials.

2.1. Flat projections. Let C be a connected curve (cf. Remark 0.1) of arithmetic genus g and $\pi : C \to \mathbb{P}^1$ a flat projection of degree d. We consider the sheaf of algebras $\pi_*\mathcal{O}_C$. As a sheaf of $\mathcal{O}_{\mathbb{P}^1}$ -modules it is locally free, i.e. a vector bundle of rank d, which we denote by E_{π} . It has a trivial summand and all remaining summands have a negative degree. Moreover the degree of E_{π} is equal to -(d+g-1)(since $\chi(E_{\pi}) = \chi(\mathcal{O}_C) = 1 - g$). Consider now the regular representation of $\pi_*\mathcal{O}_C$ on itself. It gives a global injective morphism $\pi_*\mathcal{O}_C \to \mathcal{E}nd(E_{\pi})$, i.e. a global section of $\mathcal{E}nd(E_{\pi}) \otimes E_{\pi}^*$. Suppose that

(2.1)
$$E_{\pi} \simeq \bigoplus_{i=1}^{d} \mathcal{O}_{\mathbb{P}^{1}}(-k_{i})$$

where $0 = k_1 < k_2 \leq \cdots \leq k_{d-1}$. Then a global section of $\mathcal{E}nd(E_{\pi})\otimes E_{\pi}^*$ corresponds to $d \neq d$ matrices A_1, \ldots, A_d of polynomials in one variable with $A_1 = 1$ and the (i, j) entry of A_l having degree $k_j - k_i + k_l$. Moreover these matrices commute, since \mathcal{O}_C is commutative. Finally, since each stalk of $\pi_*\mathcal{O}_C$ is a *d*-dimensional algebra, we have dim $\mathbb{C}[A_2(t), \ldots, A_d(t)] = d$ for each *t*. Clearly the freedom in choosing the matrices A_l is equivalent to the choice of the isomorphism (2.1).

Conversely, let R be such a d-dimensional commutative subalgebra of $\operatorname{Mat}_{d \times d}(\mathbb{C}[t])$ with identity. Then R is integral over $\mathbb{C}[t]$. Consider the matrices A'_l over $\mathbb{C}[1/t]$ obtained by conjugating A_l by diag $(t^{k_1}, \ldots, t^{k_d})$ and dividing by t^{k_l} . We obtain a commutative ring R', integral over $\mathbb{C}[1/t]$. The affine curves $\operatorname{Spec} R$ and $\operatorname{Spec} R'$ have an obvious gluing and we obtain a curve C together with a projection $C \to \mathbb{P}^1 = \operatorname{Spec} \mathbb{C}[t] \cup \operatorname{Spec} \mathbb{C}[1/t]$. Each fibre C_t is a 0-dimensional scheme of length d and so the projection $C \to \mathbb{P}^1$ is flat. It is clear that the constructions are inverse to each other.

Proposition 2.1. There exists a natural 1 - 1 correspondence between:

(A) connected curves C with a flat projection π of degree d onto \mathbb{P}^1 and a fixed isomorphism

$$\pi_*\mathcal{O}_C\simeq\mathcal{O}_{\mathbb{P}^1}\oplus\bigoplus_{i=2}^d\mathcal{O}_{\mathbb{P}^1}(-k_i),$$

(B) d-1-tuples of commuting $d \times d$ matrix polynomials $A_2(t), \ldots, A_d(t)$ such that

- (i) the first column of A_l is the constant vector e_l ;
- (ii) the (i, j) entry of A_l has degree $k_j k_i + k_l$;
- (iii) dim $\mathbb{C}[A_2(t), \ldots, A_d(t)] = d$ for any $t \in \mathbb{P}^1$.

2.2. Curves in projective spaces. Suppose now that C is a connected curve of degree d in $\mathbb{P}^r \setminus \mathbb{P}^{r-2}$, not contained in any hyperplane, and let the projection π be the restriction of the projection

$$[z_1,\ldots,z_{r+1}]\mapsto [z_r,z_{r+1}].$$

Each z_i , i = 1, ..., r - 1, defines a direct summand of $\pi_* \mathcal{O}_C$, isomorphic to $\mathcal{O}_{\mathbb{P}^1}(-1)$. Moreover, since C is not contained in a hyperplane, these summands are linearly independent, and, consequently, $k_2 = \cdots = k_r = 1$ in (2.1). Since \mathcal{O}_{C_t} is generated by z_1, \ldots, z_{r-1} , the matrices $A_2(t), \ldots, A_r(t)$ generate $\mathbb{C}[A_2(t), \ldots, A_d(t)]$ for each t. We conclude:

Proposition 2.2. There exists a natural 1 - 1 correspondence between:

(A) pairs (C, ϕ) , where C is a connected curve C of degree d in $\mathbb{P}^r \setminus \mathbb{P}^{r-2}$ not contained in any hyperplane and such that the projection π onto the complementary \mathbb{P}^1 is flat, while ϕ is a fixed isomorphism

$$\pi_*\mathcal{O}_C\simeq\mathcal{O}\oplus\mathcal{O}(-1)^{\oplus r-1}\oplus\bigoplus_{i=r+1}^d\mathcal{O}(-k_i),$$

(B) (r-1)-tuples of commuting $d \times d$ matrix polynomials $A_2(t), \ldots, A_r(t)$ such that

- (i) the first column of A_l is the constant vector $e_l, l = 2, ..., r$;
- (ii) the (ij)-entry of A_l has degree $k_j k_i + k_l$ (here $k_1 = 0$ and $k_2 = \cdots = k_r = 1$);
- (iii) for any $t \in \mathbb{P}^1$, $\mathbb{C}[A_2(t), \ldots, A_r(t)]$ has dimension d and is conjugate to its image in $\operatorname{End}(\mathbb{C}^d)$ under the regular representation.

Example 2.3. Let C be a smooth curve of degree r in $\mathbb{P}^r \setminus \mathbb{P}^{r-2}$. Such a curve is cut out by the 2×2 minors of a $2 \times r$ matrix of linear forms in homogeneous coordinates z_1, \ldots, z_{r+1} . Thus we obtain $\binom{r}{2}$ quadratic equations. Since we assume that the projection onto $[z_r, z_{r+1}]$ is flat, the matrix of coefficients of $z_i z_j$, $i, j \leq r - 1$, is invertible and we can write the equations in affine coordinates $t = z_r/z_{r+1}$, $x_i = z_i/z_{r+1}$, $i \leq r - 1$, as

$$x_i x_j = \sum_{k=1}^{r-1} a_{ij}^k(t) x_k + b_{ij}(t), \quad i, j = 1, \dots, r-1,$$

where a_{ij} are linear in t and b_{ij} is quadratic in t. It follows that the commuting matrices $A_2(t), \ldots, A_r(t)$ are given by

$$A_{l}(t) = \begin{pmatrix} 0 & b_{l1}(t) & \dots & b_{l,r-1}(t) \\ 0 & a_{l1}^{1}(t) & \dots & a_{l,r-1}^{1}(t) \\ \vdots & \vdots & & \vdots \\ 1 & a_{l1}^{l}(t) & \dots & a_{l,r-1}^{l}(t) \\ \vdots & \vdots & & \vdots \\ 0 & a_{l1}^{r-1}(t) & \dots & a_{l,r-1}^{r-1}(t) \end{pmatrix}.$$

8

and

Example 2.4. The correspondence in Proposition 2.2 commutes with projections. Thus, if C is a curve of degree d in $\mathbb{P}^r \setminus \mathbb{P}^{r-2}$ obtained by projection onto the first r coordinates from a curve \hat{C} in $\mathbb{P}^n \setminus \mathbb{P}^{n-2}$, then the matrix polynomials $A_2(t), \ldots, A_r(t)$ corresponding to the curve C are simply the first r-1 matrix polynomials corresponding to the curve \hat{C} .

Example 2.5. Let C be a rational curve of degree 4 in \mathbb{P}^3 parametrised by $[u^3v, v^3u, u^4, v^4]$. We can either use the previous example or proceed directly as follows: if we set $x = z_1/z_4, y = z_2/z_4, t = z_3/z_4$, then the ideal of C is generated by

$$xy - t$$
, $t^2y - x^3$, $ty^2 - x^2$, $y^3 - x$.

It follows that $\mathcal{O}(C_t)$ is spanned by $1, x, y, y^2$ for $t \neq \infty$ and by $1, x, y, x^2$ for $t \neq 0$. Computing the endomorphisms $x \cdot ()$ and $y \cdot ()$ gives:

$$A = \begin{pmatrix} 0 & 0 & t & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & t \\ 0 & t & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & t & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

Example 2.6. Let C a complete intersection of two quadrics in \mathbb{P}^3 , given by equations $Q_1(x, y, t) = 0$, $Q_2(x, y, t) = 0$. Let $\tilde{Q}_1(x, y), \tilde{Q}_2(x, y)$ be the parts involving only x, y and let $\tilde{Q}_3(x, y)$ be the quadratic polynomial in x, y independent of \tilde{Q}_1, \tilde{Q}_2 . Then $\mathcal{O}(C_t)$ is spanned by $1, x, y, \tilde{Q}_3$, and hence $E_{\pi} \simeq \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-2)$. In particular, the arithmetic genus of C is 1. Generically, such a C is a smooth elliptic curve, but if Q_1 and Q_2 are two of the three quadrics defining a twisted cubic, then C is the union of this cubic and a line, intersecting in two points.

3. Space curves

We consider the fibrewise Hilbert scheme Z_d of d points for the projection $\pi : \mathbb{P}^3 \setminus \mathbb{P}^1 \to \mathbb{P}^1$ (see [1, Ch. 1, §7] for a definition and properties of relative Hilbert schemes). Locally, over an open subset U of \mathbb{P}^1 , it is just $U \times (\mathbb{C}^2)^{[d]}$. It is therefore a (2d+1)-dimensional complex manifold with a holomorphic projection $p: Z_d \to \mathbb{P}^1$.

 Z_d satisfies the necessary conditions to be the twistor space of a pseudo-hyperkähler manifold (i.e. a pseudo-Riemannian manifold M with a fibrewise action of quaternions on TM, parallel with respect to the Levi-Civita connection):

- (i) it has an antiholomorphic involution σ covering the antipodal map on P¹, induced from the standard antiholomorphic involution on the total space of O_{P¹}(1) ⊕ O_{P¹}(1) ≃ P³\P¹;
- (ii) it has an $\mathcal{O}_{\mathbb{P}^1}(2)$ -valued symplectic form ω along the fibres of π , again induced² from the $\mathcal{O}_{\mathbb{P}^1}(2)$ -valued fibrewise symplectic form on the total space of $\mathcal{O}_{\mathbb{P}^1}(1) \oplus \mathcal{O}_{\mathbb{P}^1}(1)$.

The pseudo-hyperkähler manifold is then the Kodaira moduli space of σ -invariant sections of p, the normal bundle of which splits as $\mathcal{O}_{\mathbb{P}^1}(1)^{\oplus 2d}$ [18, §3(F)].

 $^{^2 \}mathrm{Recall}$ [2] that the Hilbert scheme of d points on a symplectic surface has a canonical symplectic structure.

3.1. Normal sheaf of space curves.

Proposition 3.1. There exists a natural isomorphism between an open subset of the Hilbert scheme of $\mathbb{P}^3 \setminus \mathbb{P}^1$ consisting of degree d curves which are flat over \mathbb{P}^1 , and the Hilbert scheme of sections of $p: Z_d \to \mathbb{P}^1$.

Remark 3.2. The Hilbert scheme of curves of degree d is the union of all components of Hilb($\mathbb{P}^3 \setminus \mathbb{P}^1$) which have the Hilbert polynomial of the form $h(n) = dn + c, c \in \mathbb{Z}$. The Hilbert scheme of sections of $p: \mathbb{Z}_d \to \mathbb{P}^1$ is an open subscheme of Hilb(\mathbb{Z}_d).

Proof. If C is a degree d curve in $\mathbb{P}^3 \setminus \mathbb{P}^1$, flat over \mathbb{P}^1 , then its scheme-theoretic intersection with each fibre of $\mathbb{P}^3 \setminus \mathbb{P}^1 \to \mathbb{P}^1$ yields a section of Z_d . The inverse map is defined as follows: given a section of Z_d , pullback the universal family over the relative Hilbert scheme Z_d to \mathbb{P}^1 . A flat family of curves W in $\mathbb{P}^3 \setminus \mathbb{P}^1$ parameterised by a scheme T is sent to a flat family of sections of p (viewed as a flat family of \mathbb{P}^1 's in Z_d). Similarly, the inverse map sends a flat family of sections to a flat family of degree d curves. The functorial interpretation of Hilbert schemes implies that both maps are morphisms of schemes.

We can relate the normal bundle N_{s/Z_d} of a section to the normal sheaf $\mathcal{N}_{C/\mathbb{P}^3}$ of the corresponding curve as follows:

Lemma 3.3. $N_{s/Z_d} \simeq \pi_* \mathcal{N}_{C/\mathbb{P}^3}$.

Proof. The normal bundle of a section is isomorphic to the restriction of the vertical tangent bundle Ker dp of Z_d to the section. Since Z_d is the fibrewise Hilbert scheme of points, the fibre of the vertical tangent bundle at $D \in Z_d$ is $H^0(D, \mathcal{N}_{D/F})$, where $F \simeq \mathbb{C}^2$ is the fibre containing D. On the other hand the ideal of D in F is just $J_C \otimes \mathcal{O}_F$, where J_C is the ideal of C in \mathbb{P}^3 . Thus $\mathcal{N}_{D/F} = \mathcal{N}_{C/\mathbb{P}^3} \otimes \mathcal{O}_F$, and hence the function $\mathbb{P}^1 \ni z \to h^0(\pi^{-1}(z), \mathcal{N}_{C/\mathbb{P}^3}|_{\pi^{-1}(z)})$ is constant. It follows from a result of Grauert [14, Cor. III.12.9] that $\pi_* \mathcal{N}_{C/\mathbb{P}^3}$ is locally free. Therefore both $\pi_* \mathcal{N}_{C/\mathbb{P}^3}$ and N_{s/Z_d} are vector bundles and the natural map $\pi_* \mathcal{N}_{C/\mathbb{P}^3} \to N_{s/Z_d}$ is an isomorphism on fibres.

Corollary 3.4. The normal sheaf of a Cohen-Macaulay curve in \mathbb{P}^3 is torsion-free.

Proof. We can find a projective line $l \subset \mathbb{P}^3$ such that the projection $\pi : C \to l$ is flat. Were $\mathcal{N}_{C/\mathbb{P}^3}$ not torsion free, neither would be $\pi_*\mathcal{N}_{C/\mathbb{P}^3}$, contradicting the above lemma.

Lemma 3.3 implies that the normal bundle of a section splits as $\mathcal{O}_{\mathbb{P}^1}(1)^{\oplus 2d}$ if and only if $H^*(C, \mathcal{N}_{C/\mathbb{P}^3}(-2)) = 0$. As mentioned at the beginning of the section, the parameter space of curves satisfying this condition has a natural complexified pseudo-hyperkähler structure and on its σ -invariant part a genuine pseudohyperkähler structure [4]. We now want to investigate this parameter space via commuting matrix polynomials.

If E is a rank d vector bundle on \mathbb{P}^1 , we denote by $\mathcal{K}(E)$ the subsheaf of $(\mathcal{E}nd(E)\otimes \mathcal{O}_{\mathbb{P}^1}(1))^{\oplus 2}$ defined by

$$\mathcal{K}(E)(U) = \left\{ (A(t), B(t)) \in \left(\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(1) \right)^{\oplus 2}(U); \, \forall_{t \in U} \ (A(t), B(t)) \in M_d^{\mathrm{reg}} \right\}.$$

We then denote by K(E) the total space of $\mathcal{K}(E)$, i.e. the subset of the total space of $(\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(1))^{\oplus 2}$ consisting of points through which passes a local section of $\mathcal{K}(E)$. Since \mathbb{Z}_d is the relative Hilbert scheme for the projection $\mathcal{O}_{\mathbb{P}^1}(1) \oplus \mathcal{O}_{\mathbb{P}^1}(1) \to$ \mathbb{P}^1 , Proposition 1.4 implies that Z_d is biholomorphic to the quotient of $K(\mathcal{O}_{\mathbb{P}^1}^{\oplus d})$ by the fibrewise action of $GL(d, \mathbb{C})$. In fact we have:

Lemma 3.5. Let E be a vector bundle of rank d on \mathbb{P}^1 . Then the quotient of K(E) by the fibrewise action of $GL(d, \mathbb{C})$ is biholomorphic to Z_d .

Proof. Let $E \simeq \bigoplus \mathcal{O}_{\mathbb{P}^1}(\lambda_i)$. Then $\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(1)$ is isomorphic to two copies $\mathbb{C} \times \operatorname{Mat}_{d \times d}(\mathbb{C})$ glued over $t \neq 0, \infty$ by $(\tilde{t}, \tilde{A}) = (1/t, t^{\lambda}At^{-\lambda})$, where $\lambda = \operatorname{diag}(\lambda_1, \ldots, \lambda_d)$. Thus, after taking the fibrewise quotient by $GL(d, \mathbb{C})$, we obtain the same complex manifold, independently of the choice of E. \Box

From now on $d \geq 3$. Let C be a connected nonplanar curve of degree d in $\mathbb{P}^3 \setminus \mathbb{P}^1$, flat over \mathbb{P}^1 . If C satisfies $\pi_* \mathcal{O}_C \simeq E$, then the corresponding section of $p : Z_d \to \mathbb{P}^1$ arises as the projection of a global section of $\mathcal{K}(E) \to \mathbb{P}^1$, i.e. it can be represented by a pair of commuting matrix polynomials, the degrees of which satisfy the constraints of part B in Proposition 2.2. We should like to remark that this gives constraints for possible degrees of commuting pairs of polynomials A(t), B(t), such that dim Z(A(t), B(t)) = d for each t.

Let \mathcal{Z} denote the bundle of centralisers of (A, B), i.e. a subbundle of $\mathcal{E}nd(E)$, the fibre of which over each t is spanned by Z(A(t), B(t)). Let \mathcal{T} denote the kernel of the homomorphism

$$(3.1) \quad D: \left(\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(1)\right)^{\oplus 2} \to \mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(2), \quad (a,b) \to [A,b] + [a,B].$$

In other words \mathcal{T} is the fibrewise tangent bundle to K(E). We obtain a short exact sequence of locally free sheaves on \mathbb{P}^1 :

$$(3.2) 0 \to \mathcal{E}nd(E)/\mathcal{Z} \to \mathcal{T} \to N \to 0$$

where N is the normal bundle of the section in Z_d and the first map is given by $\rho \mapsto ([\rho, A], [\rho, B])$, i.e. its image consists of fundamental vector fields for the action of $GL(d, \mathbb{C})$.

Lemma 3.6. $\mathcal{Z} \simeq E$ and if we write $E \simeq \mathcal{O}_{\mathbb{P}^1} \oplus E'$, then the embedding $\mathcal{Z} \hookrightarrow \mathcal{E}nd(E) \simeq \mathcal{O}_{\mathbb{P}^1} \oplus E' \oplus (E')^* \oplus \mathcal{E}nd(E')$ is the isomorphism with the direct sum of the first two summands.

Proof. Let C be the curve in $\mathbb{P}^3 \setminus \mathbb{P}^1$ determined by A(t), B(t). We know that $\pi_* \mathcal{O}_C \simeq E$. Let $E' \simeq \bigoplus_{i=2}^d \mathcal{O}_{\mathbb{P}^1}(\lambda_i)$. Each summand defines a section $A_i(t)$ of $\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(-\lambda_i)$ and these matrix polynomials commute. Moreover the first column of each $A_i(t)$ is just e_i . The bundle \mathcal{Z} is spanned over each t by 1 and the $A_i(t)$, i.e. it is isomorphic to the first column of $\mathcal{E}nd(E)$. The claim follows. \Box

The image sheaf of D is also locally free and can be identified with the annihilator of \mathcal{Z} in $\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(2)$ (as vector bundles):

$$\rho \in \operatorname{Im} D \iff \operatorname{tr} \rho z = 0 \ \forall z \in \mathcal{Z}.$$

Let us write $\mathcal{L} = (E')^* \oplus \mathcal{E}nd(E') \simeq \mathcal{E}nd(E)/\mathcal{Z}$. As a matrix of endomorphisms of E, it has the first column equal to 0. From the above characterisation of Im Dit follows that Im $D \simeq \mathcal{L}^*(2)$, and, consequently, we can write the two short exact sequences (i.e. (3.2) and the sequence induced by (3.1)) as:

$$(3.3) 0 \to \mathcal{L} \to \mathcal{T} \to N \to 0,$$

(3.4)
$$0 \to \mathcal{T} \to \left(\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(1)\right)^{\oplus 2} \to \mathcal{L}^*(2) \to 0.$$

These sequence describe Z_d , at least infinitesimally, as a fibrewise symplectic quotient of the total space of $(\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(1))^{\oplus 2}$ by a Lie group, the Lie algebra of which is \mathcal{L} . Our idea, on how to study the hyperkähler structure of the Hilbert scheme of curves in $\mathbb{P}^3 \setminus \mathbb{P}^1$, is essentially to describe the hyperkähler manifold of sections of Z_d as a "hyperkähler quotient" of the (open dense subset of) vector space $H^0(\mathbb{P}^1, (\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(1))^{\oplus 2})$ by a Lie group, the Lie algebra of which is $H^0(\mathbb{P}^1, \mathcal{L})$. This cannot quite work in this form for various reasons, the most obvious of which is that the sections of $(\mathcal{E}nd(E) \otimes \mathcal{O}_{\mathbb{P}^1}(1))^{\oplus 2}$ have the wrong normal bundle in order to form a hyperkähler manifold. In §5 we shall show how to modify and implement this idea for curves of genus 0. For higher genera, one probably has to work with the \mathcal{P} -structures of Gindikin [12], and with their quotients.

Remark 3.7. The above sequences provide numerical restrictions on curves which are cohomologically stable. Indeed, tensor (3.3) and (3.4) with $\mathcal{O}(-2)$ and put:

$$a = h^{0}((\mathcal{L}(-2)) = h^{1}(\mathcal{L}^{*}), \ b = h^{1}((\mathcal{L}(-2)) = h^{0}(\mathcal{L}^{*}),$$
$$c = h^{0}(\mathcal{E}nd(E)(-1)) = h^{1}(\mathcal{E}nd(E)(-1)).$$

Then $h^0(\mathcal{T}(-2)) \geq 2c - b$ and, consequently, if C is cohomologically stable, i.e. $h^0(N(-2)) = h^1(N(-2)) = 0$, then $a \geq 2c - b$. Let

$$\pi_*\mathcal{O}_C\simeq\mathcal{O}_{\mathbb{P}^1}\oplus \bigoplus_{i=1}^s\mathcal{O}_{\mathbb{P}^1}(-i)^{m_i}.$$

Then $a = \sum_{j \ge i+2} (j-i-1)m_i m_j + \sum_{i \ge 2} (i-1)m_i$, $b = \sum_{j \ge i} (j-i+1)m_i m_j$, $c = \sum_{j \ge i+1} (j-i)m_i m_j + \sum_i im_i$. Thus

$$0 \ge 2c - a - b = \sum_{i} (i+1)m_i - \sum_{i} m_i^2.$$

For example a curve such that $m_i \leq i + 1$ for all i and $m_i \leq i$ for at least one i cannot be cohomologically stable (cf. Example 2.6).

3.2. Twistor space revisited. Given the constraints of Proposition 2.2 on the matrices A, B representing a space curve, we can replace Z_d for $d \ge 3$ by its open dense subset Z'_d , the fibres of which consist of 0-dimensional subschemes of \mathbb{C}^2 not contained in any line. We have an analogue of Proposition 3.1, the proof of which is the same:

Proposition 3.8. Let $d \geq 3$. There exists a natural isomorphism between an open subset of the Hilbert scheme of $\mathbb{P}^3 \setminus \mathbb{P}^1$ consisting of degree d curves which are flat over \mathbb{P}^1 , and the Hilbert scheme of sections of $p: Z'_d \to \mathbb{P}^1$. \Box

Theorem 1.10 tells us how to construct Z'_d from a vector bundle on \mathbb{P}^1 . Let E' be a vector bundle of rank d-1 over \mathbb{P}^1 of the form $E' \simeq \bigoplus_{i=1}^{d-1} \mathcal{O}_{\mathbb{P}^1}(-k_i)$ with $k_1 = k_2 = 1$ and all k_i positive. Define $\mathcal{K}^0(E')$ to be the subsheaf of $(\mathcal{E}nd(E') \otimes \mathcal{O}_{\mathbb{P}^1}(1))^{\oplus 2}$, the local section of which are pairs (X(t), Y(t)) such that for all t:

- (i) the first column of Y(t) is equal to the second column of X(t);
- (ii) there is no subspace S of \mathbb{C}^{d-1} such that $e_1, e_2 \in S, X(t)S \subset S, Y(t)S \subset S$;
- (iii) [X(t), Y(t)] has nonzero entries only in the first two rows.

We denote by $K^0(E)$ the total space of $\mathcal{K}^0(E)$ (defined as for K(E)). We define Z(E') to be the fibrewise quotient of $K^0(E')$ by the action (1.4) of the group \hat{G}^0 defined in §1.1. Theorem 1.10 implies that $Z(E') \simeq Z'_d$ and that is a smooth manifold fibreing over \mathbb{P}^1 and we can conclude:

Proposition 3.9. $Z(E') \simeq Z'_d$.

Proof. This is the same argument as in the proof of Lemma 3.5.

Consequently, sections of Z'_d corresponding to curves with $\pi_* \mathcal{O}_C \simeq \mathcal{O}_{\mathbb{P}^1} \oplus E'$ arise as projections of global sections of K(E'), and we can represent them by pairs (X(t), Y(t)) of matrix polynomials satisfying the above constraints.

4. INTERLUDE: COMPLEXIFIED HYPERKÄHLER STRUCTURES

The complexified algebra of quaternions is isomorphic to $\operatorname{Mat}_{2,2}(\mathbb{C})$. Consequently, a complexification of a real manifold with a geometry based on quaternions (such as hyperkähler, hypercomplex, quaternionic) or on split quaternions (hypersymplectic geometry) will posses a geometry based on this algebra (called *algebra* of biquaternions by Hamilton). Such geometries have been considered in the past, in particular by Jardim and Verbitsky in [19]. They call a complexified hyperkähler structure a trisymplectic structure generating an $SL(2, \mathbb{C})$ -web. Although there are good reasons for this terminology, we prefer the shorter name \mathbb{C} -hyperkähler.

Definition 4.1. A complex manifold M is called almost \mathbb{C} -hypercomplex if its holomorphic tangent bundle TM decomposes as $TM \simeq E \otimes \mathbb{C}^2$, where E is a holomorphic vector bundle. It is \mathbb{C} -hypercomplex if, in addition, for any $v \in \mathbb{C}^2$ the subbundle $E \otimes v$ defines an integrable distribution on M.

Definition 4.2. A \mathbb{C} -hypercomplex manifold is called \mathbb{C} -hyperkähler if it is equipped with a holomorphic section g of S^2T^*M , such that:

- (i) at any $m \in M$, g_m is nondegenerate,
- (ii) for any $A \in Mat_{2,2}(\mathbb{C}), g(A \cdot, \cdot) = g(\cdot, A_{adj} \cdot)$, and
- (iii) for any $A \in \mathfrak{sl}(2,\mathbb{C})$, the holomorphic 2-form $\omega_A = g(A, \cdot)$ is closed.

Given a \mathbb{C} -hypercomplex manifold we can define an integrable distribution \mathcal{D} on $M \times \mathbb{P}^1$ by $\mathcal{D}_{|M \times \{z\}} = E \otimes h$, where h is the highest weight vector for the maximal torus in $SL(2, \mathbb{C})$ corresponding to $z \in \mathbb{P}^1$. If \mathcal{D} is simple³, then the leaf space Z is the twistor space of M, and points of M correspond to sections of $Z \to \mathbb{P}^1$ with normal bundles splitting as $\bigoplus \mathcal{O}_{\mathbb{P}^1}(1)$. If the distribution \mathcal{D} is not simple, we need to view Z in terms of foliated geometry.

If M is \mathbb{C} -hyperkähler, then its twistor space is equipped with a fibrewise $\mathcal{O}_{\mathbb{P}^1}(2)$ -valued complex symplectic form.

We shall now discuss \mathbb{C} -hypercomplex and \mathbb{C} -hyperkähler quotients. In the case of a reductive Lie group, \mathbb{C} -hyperkähler quotients have been introduced and studied by Jardim and Verbitsky [19] under the name "trisymplectic reduction". For us the case of nonreductive groups will be of paramount importance.

³We call a distribution \mathcal{D} on a manifold Y simple, if the leaf space X of the corresponding foliation is a manifold and the map $Y \to X$ is a submersion.

Definition 4.3. Let M be a \mathbb{C} -hypercomplex manifold equipped with a holomorphic action of a complex Lie group G preserving the \mathbb{C} -hypercomplex structure. A \mathbb{C} hypercomplex moment map is a G-equivariant holomorphic map $\mu : M \to \mathfrak{g}^* \otimes \mathfrak{sl}(2, \mathbb{C})^*$, such that there exists a holomorphic \mathfrak{g}^* -valued 1-form ϕ with the property that $\Phi = \phi \cdot 1_{2\times 2} + d\mu : TM \to \mathfrak{g}^* \otimes \operatorname{Mat}_{2,2}(\mathbb{C})^*$ satisfies $\langle \Phi(Av), B \rangle = \langle \Phi(v), A_{\operatorname{adj}} B \rangle$ for any $A \in \operatorname{Mat}_{2,2}(\mathbb{C})$.

Definition 4.4. Let M be a \mathbb{C} -hyperkähler manifold equipped with a holomorphic action of a complex Lie group G preserving the \mathbb{C} -hyperkähler structure. A \mathbb{C} hyperkähler moment map is a G-equivariant holomorphic map $\mu : M \to \mathfrak{g}^* \otimes \mathfrak{sl}(2,\mathbb{C})^*$, such that, for any $A \in \mathfrak{sl}(2,\mathbb{C})$ and any fundamental vector field X_{ρ} , $\rho \in \mathfrak{g}, \langle d\mu(\cdot), \rho \otimes A \rangle = \omega_A(X_{\rho}, \cdot).$

Remark 4.5. A C-hyperkähler moment map is also a C-hypercomplex moment map, with $\langle \phi(v), \rho \rangle = g(X_{\rho}, v), \ \rho \in \mathfrak{g}, \ v \in TM$.

The \mathbb{C} -hypercomplex or \mathbb{C} -hyperkähler reduction proceeds now along the usual lines: given G and a moment map $\mu : M \to \mathfrak{g}^* \otimes \mathfrak{sl}(2, \mathbb{C})^*$, choose a G-invariant element $c \in \mathfrak{g}^* \otimes \mathfrak{sl}(2, \mathbb{C})^*$. Unlike in the hyperkähler case, the freeness of the action of G on $\mu^{-1}(c)$ does not imply that $\mu^{-1}(c)$ is smooth. Moreover, even if $\mu^{-1}(c)$ is smooth and the action of G on $\mu^{-1}(c)$ is free and proper, then although $\mu^{-1}(c)/G$ is a complex manifold [7, Ch.III, Prop.10], it is not necessarily a \mathbb{C} -hypercomplex or \mathbb{C} -hyperkähler manifold. As observed by several authors in related settings (especially [20, §3], [17], [8, §4]), both smoothness of $\mu^{-1}(c)$ and the existence of induced geometry on $\mu^{-1}(c)/G$ are guaranteed by a single nondegeneracy condition. Namely, we have:

Theorem 4.6. Let $\mu : M \to \mathfrak{g}^* \otimes \mathfrak{sl}(2, \mathbb{C})^*$ be a \mathbb{C} -hypercomplex (resp. \mathbb{C} -hyperkähler) moment map and let $c \in \mathfrak{g}^* \otimes \mathfrak{sl}(2, \mathbb{C})^*$ be *G*-invariant. Suppose that the action of *G* on $\mu^{-1}(c)$ is free and proper and that, at any $m \in \mu^{-1}(c)$, $\phi_m(X_\rho) \neq 0$ for any $\rho \in \mathfrak{g}$ (resp. the restriction of *g* to the subspace of $T_m M$ generated by fundamental vector fields is nondegenerate). Then $\mu^{-1}(c)/G$ is a \mathbb{C} -hypercomplex (resp. \mathbb{C} -hyperkähler) manifold. \Box

However, the assumptions in Theorem 4.6 are not the most general ones, under which one obtains a nondegenerate \mathbb{C} -hyperkähler quotient:

Proposition 4.7. Let $\mu : M \to \mathfrak{g}^* \otimes \mathfrak{sl}(2, \mathbb{C})^*$ be a \mathbb{C} -hyperkähler moment map and let $c \in \mathfrak{g}^* \otimes \mathfrak{sl}(2, \mathbb{C})^*$ be *G*-invariant. Suppose that $\mu^{-1}(c)$ is smooth and the action of *G* on $\mu^{-1}(c)$ is free and proper. Then $\mu^{-1}(c)/G$ is a \mathbb{C} -hyperkähler manifold if and only if $\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp}$ is $\operatorname{Mat}_{2,2}(\mathbb{C})$ -invariant along $\mu^{-1}(c)$, where $\check{\mathfrak{g}}$ denotes the subspace of $T_m M$ generated by fundamental vector fields, and $\check{\mathfrak{g}}^{\perp} = \{v; g(X_{\rho}, v) = 0 \ \forall \rho \in \mathfrak{g}\}.$

Proof. Let $I, J, K \in \mathfrak{sl}(2, \mathbb{C})$ denote the standard basis of quaternions. Hitchin [17] (see also [8, p.102]) shows that along $\mu^{-1}(c)$,

$$\operatorname{Ker} \omega_I = \check{\mathfrak{g}} + J\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp} + K\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp},$$

and cyclically in I, J, K. Thus ω_I descends to a nondegenerate symplectic form on $\mu^{-1}(c)/G$ if and only if $J\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp} + K\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp} \subset \check{\mathfrak{g}}$. Therefore $\mu^{-1}(c)/G$ is a \mathbb{C} -hyperkähler manifold if and only if $\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp}$ is $Mat_{2,2}(\mathbb{C})$ -invariant. \Box

Proposition 4.8. Suppose that the conditions of the above proposition are satisfied. Then $\dim \check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp} = k$ is constant along $\mu^{-1}(c)$ and

$$\dim \mu^{-1}(c)/G = \dim M - 4 \dim G + 2k.$$

Proof. The smoothness of $\mu^{-1}(c)$ implies that $(\operatorname{Ker} d\mu)^{\perp} = I\check{\mathfrak{g}} + J\check{\mathfrak{g}} + K\check{\mathfrak{g}}$ has constant dimension along $\mu^{-1}(c)$, where \perp denotes the orthogonal "complement" with respect to the form g. Suppose that $IX_{\rho_1} + JX_{\rho_2} + KX_{\rho_3} = 0$. Taking the scalar products (for the form g) with $IX_{\rho}, JX_{\rho}, KX_{\rho}$ shows that $X_{\rho_1}, X_{\rho_2}, X_{\rho_3} \in$ $\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp}$. Therefore dim $(\operatorname{Ker} d\mu)^{\perp} = 3 \dim G - 2 \dim \check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp}$ and it follows that dim $\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp}$ must be constant along $\mu^{-1}(c)$.

We shall use the above results in the following situation.

Corollary 4.9. Let $\mu = \mu_H \oplus \mu_L : M \to \mathfrak{g}^* \otimes \mathfrak{sl}(2, \mathbb{C})^*$ be a \mathbb{C} -hyperkähler moment map for a semidirect product $G \simeq H \ltimes L$ and let $c = (c_H, c_L) \in \mathfrak{g}^* \otimes \mathfrak{sl}(2, \mathbb{C})^*$ be *G*-invariant. Suppose that the following conditions are satisfied:

- (i) the action of G on $\mu^{-1}(c)$ is free and proper;
- (ii) $\mu_L^{-1}(c_L)$ is smooth and the action of L on $\mu_L^{-1}(c_L)$ is free and proper; (iii) \check{I} is $\operatorname{Mat}_{2,2}(\mathbb{C})$ -invariant along $\mu^{-1}(c_L)$;

(iv) $\check{\mathfrak{l}} \subset \check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp}$ along $\mu^{-1}(c)$, with equality holding generically.

Then:

a) $\mu^{-1}(c)/G$ is smooth if and only if dim $I\check{\mathfrak{h}} + J\check{\mathfrak{h}} + K\check{\mathfrak{h}} = 3 \dim H$ along $\mu^{-1}(c)$.

b) $\mu^{-1}(c)/G$ is a \mathbb{C} -hyperkähler manifold if and only if $\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp} = \check{\mathfrak{l}}$ along $\mu^{-1}(c)$.

In both a) and b), the dimension of $\mu^{-1}(c)/G$ is dim $M - 4 \dim H - 2 \dim L$.

Proof. Conditions (ii) and (iii) together with Proposition 4.7 imply that $M_L =$ $\mu_L^{-1}(c_L)/L$ is a \mathbb{C} -hyperkähler manifold (of dimension dim $M-2 \dim L$). Since L is normal in G, we obtain a \mathbb{C} -hyperkähler action of H on M_L and a \mathbb{C} -hyperkähler moment map $\bar{\mu}_H : M_L \to \mathfrak{h}^* \otimes \mathfrak{sl}(2,\mathbb{C})^*$. The \mathbb{C} -hyperkähler quotient of M_L by H is isomorphic to the \mathbb{C} -hyperkähler quotient of M by G. Statement a) follows, since dim $I\dot{\mathfrak{h}} + J\dot{\mathfrak{h}} + K\dot{\mathfrak{h}} = 3 \dim H$ is equivalent to $\bar{\mu}_H$ being a submersion at points of $\bar{\mu}_{H}^{-1}(c_{H})$. On the other hand, the condition in b) means that the triple $(M_L, H, \bar{\mu}_H)$ satisfies conditions of Theorem 4.6, and is therefore sufficient. Conversely, if $\mu^{-1}(c)/G$ is a \mathbb{C} -hyperkähler manifold, then, owing to a), $\dim I\check{\mathfrak{h}} + J\check{\mathfrak{h}} + K\check{\mathfrak{h}} = 3\dim H$, which means that $\bar{\mu}_{H}^{-1}(c_{H})$ is smooth. The neccessity of $\check{\mathfrak{g}} \cap \check{\mathfrak{g}}^{\perp} = \check{\mathfrak{l}}$ follows now from Proposition 4.7.

5. Genus zero space curves

We return to the situation discussed in §3 and consider in detail the case of genus 0 space curves. We denote by R_d an open subset of the Hilbert scheme Hilb_{d.0} of subschemes in \mathbb{P}^3 with Hilbert polynomial h(n) = dn + 1, consisting of those $C \in \operatorname{Hilb}_{d,0}$ which satisfy

- (i) C is contained in $\mathbb{P}^3 \setminus \{[z_0, z_1, 0, 0]\};$
- (ii) the projection π of C onto $\{[0, 0, z_2, z_3]\}$ is flat;
- (iii) C is nonplanar.

Such a C is automatically Cohen-Macaulay and connected. We denote by R_d^i , i = 0, 1, 2, an open subset of R_d consisting of those $C \in \text{Hilb}_{d,0}$ which satisfy, in addition, $h^1(\mathcal{N}_{C/\mathbb{P}^3}(-i)) = 0$. We have $R^2_d \subset R^1_d \subset R^0_d$ and R^0_d is precisely the smooth locus of R_d . Moreover, the results of Ghione and Sacchiero [11, Cor. 1.4] imply that any immersed rational curve in R_d belongs to R_d^1 . For a $C \in R_d$ we have $\pi_* \mathcal{O}_C \simeq \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus (d-1)}$ (since deg $\pi_* \mathcal{O}_C =$

-(d+g-1), g=0, the degrees of the summands are nonpositive, and $h^0(\mathcal{O}_C)=1$

and thus curves in R_d correspond to sections of $p: Z_d \to \mathbb{P}^1$ which arise as projections of sections of $K(E) \to \mathbb{P}^1$ with $E = \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus (d-1)}$ (and the map associating a section to a curve is a biholomorphism between R_d and the corresponding open subset of the Kodaira moduli space of sections of p, cf. Proposition 3.1). Equivalently, owing to Proposition 3.9, curves in R_d can be obtained as sections of Z(E') with $E' \simeq \mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus (d-1)}$.

If $C \in R_d^2$, then the normal bundle of the corresponding section of Z_d is isomorphic to $\mathcal{O}_{\mathbb{P}^1}(1)^{\oplus 2d}$ and, hence, R_d^2 comes equipped with a \mathbb{C} -hyperkähler structure. Moreover, given a real structure σ on $|\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)|$, covering a real structure on \mathbb{P}^1 , we obtain a pseudo-hyperkähler or hypersymplectic structure on $(R_d^2)^{\sigma}$ (depending on whether σ is fixed-point free or not).

Example 5.1. Let C be a twisted rational curve of degree 3 not meeting the \mathbb{P}^1 . Its ideal is generated by the minors of the 3×2 matrix

$$\begin{pmatrix} x & 0\\ y & x\\ 0 & y \end{pmatrix} - C(t)$$

where the entries of C are constant or linear in t. Thus

 $x^2 = (c_{11} + c_{22})x - c_{12}y - C_3$, $xy = c_{32}x + c_{11}y - C_2$, $y^2 = (c_{21} + c_{32})y - c_{31}x - C_1$, where C_i denotes the determinant of a 2 × 2 matrix obtained by deleting the *i*-th row from *C*. Thus the matrices *A* and *B* are

$$A(t) = \begin{pmatrix} 0 & -C_3 & -C_2 \\ 1 & c_{11} + c_{22} & c_{32} \\ 0 & -c_{12} & c_{11} \end{pmatrix}, \quad B(t) = \begin{pmatrix} 0 & -C_2 & -C_1 \\ 0 & c_{32} & -c_{31} \\ 1 & c_{11} & c_{21} + c_{32} \end{pmatrix}.$$

The metric is given as the coefficient of t in power series expansion of

$$\operatorname{tr} dA \wedge dB = d(c_{11} + c_{22}) \wedge dc_{32} + dc_{32} \wedge dc_{11} + dc_{12} \wedge dc_{31} + da_{11} \wedge d(c_{21} + c_{32}) = d(c_{11} + c_{22}) \wedge d(c_{32} + c_{21}) - dc_{22} \wedge dc_{21} + dc_{12} \wedge dc_{31}.$$

The antiholomorphic involution σ , covering the antipodal map on \mathbb{P}^1 , acts on linear polynomials $c_{ij}(t)$ as

$$\begin{pmatrix} c_{11}(t) & c_{12}(t) \\ c_{21}(t) & c_{22}(t) \\ c_{31}(t) & c_{32}(t) \end{pmatrix} \mapsto t \overline{\begin{pmatrix} c_{32}(-1/\bar{t}) & c_{31}(-1/\bar{t}) \\ -c_{22}(-1/\bar{t}) & c_{21}(-1/\bar{t}) \\ -c_{12}(-1/\bar{t}) & -c_{11}(-1/\bar{t}) \end{pmatrix}}.$$

Restricting the above formula for tr $dA \wedge dB$ to σ -invariant sections shows that the hyperkähler metric on the space of real twisted cubics is flat with signature (8, 4).

We shall now describe R_d^2 , $d \geq 4$, as a \mathbb{C} -hyperkähler quotient of a flat \mathbb{C} -hyperkähler manifold. Let $M = \operatorname{Mat}_{d-1,d-1}(\mathbb{C}) \otimes \mathbb{C}^4$ and write its elements as (X_0, X_1, Y_0, Y_1) . M is a \mathbb{C} -hyperkähler manifold with the action of $\operatorname{Mat}_{2,2}(\mathbb{C})$ on TM given by the left multiplication on $\operatorname{Mat}_{2,2}(\mathbb{C}) \simeq \mathbb{C}^4$, i.e.:

(5.1)
$$\begin{pmatrix} X_0 & Y_0 \\ X_1 & Y_1 \end{pmatrix} \mapsto \begin{pmatrix} aI & bI \\ cI & dI \end{pmatrix} \begin{pmatrix} X_0 & Y_0 \\ X_1 & Y_1 \end{pmatrix}$$

The symmetric holomorphic (2, 0)-tensor g is

(5.2)
$$g = tr(dX_1 dY_0 - dX_0 dY_1).$$

The twistor space Z of M is the total space of $\operatorname{Mat}_{d-1,d-1}(\mathbb{C}) \otimes \mathcal{O}_{\mathbb{P}^1}(1)^{\oplus 2}$, and an element (X_0, X_1, Y_0, Y_1) of M is identified with the section

$$(X(t), Y(t)) = (X_0 + tX_1, Y_0 + tY_1)$$

of $Z \to \mathbb{P}^1$. The twisted fibrewise symplectic form on Z is tr $d(X_0+tX_1) \wedge d(Y_0+tY_1)$.

We define a \mathbb{C} -hyperkähler submanifold M_d of M to consist of (X_0, X_1, Y_0, Y_1) such that:

(i) $\forall_{t \in \mathbb{P}^1} (X(t), Y(t), \langle e_1, e_2 \rangle)$ is stable.

(ii) $\forall_{t \in \mathbb{P}^1} X_{12}(t) = Y_{11}(t), \ X_{22}(t) = Y_{21}(t).$

Next we define the relevant group \hat{G}_d acting on M_d . It consists of invertible $d \times d$ matrices of the form

(5.3)
$$g(t) = \begin{pmatrix} 1 & 0 & 0 & u_1(t) & \dots & u_{d-3}(t) \\ 0 & 1 & 0 & * & \dots & * \\ 0 & 0 & 1 & * & \dots & * \\ 0 & 0 & 0 & * & \dots & * \\ \vdots & \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & * & \dots & * \end{pmatrix},$$

where asterisks denote complex numbers, and $u_i(t)$, $i = 1, \ldots, d-3$, are linear polynomials. The group structure is given by matrix multiplication. Observe that if we fix t, we obtain a group $\hat{G}_d(t)$ isomorphic to the group \hat{G}_0 of §1.1. We have $\hat{G}_d \simeq G_0 \ltimes L$, where G_0 is obtained by deleting the first row and the first column of (5.3) and $L \simeq \mathbb{C}^{2(d-3)}$ is the additive group of d-3 linear polynomials. If we write an element of L as $u(t) = (0, 0, u_1(t), \ldots, u_{d-3}(t))$, then the action of \hat{G}_d on pairs of linear matrix polynomials (X(t), Y(t)) can be written as (cf. (1.4)):

(5.4)
$$(g_0, u(t)).(X(t), Y(t)) = (g_0 X(t)g_0^{-1} - e_1 u(t)g_0^{-1}, g_0 Y(t)g_0^{-1} - e_2 u(t)g_0^{-1}).$$

This action preserves the \mathbb{C} -hyperkähler structure of M_d and it lifts to an action on the twistor space of M_d : an element g(t) acts as $g(t_0)$ on the fibre over t_0 . We easily compute the \mathbb{C} -hyperkähler moment map $\mu : M_d \to \hat{\mathfrak{g}}_d^* \otimes \mathfrak{sl}(2, \mathbb{C}) \simeq$ $(\mathfrak{g}_0^* \oplus \mathfrak{l}^*) \otimes \mathfrak{sl}(2, \mathbb{C})$:

(5.5)
$$\mu_{ij} = \begin{cases} \frac{1}{2} \left(\pi([X_0, Y_1] + [X_1, Y_0]), (X_1 + tX_0)e_2 - (Y_1 + tY_0)e_1 \right) & \text{if } ij = 11, \\ \left(\pi([X_0, Y_0]), X_0e_2 - Y_0e_1 \right) & \text{if } ij = 21, \\ \left(\pi([X_1, Y_1]), t(X_1e_2 - Y_1e_1) \right) & \text{if } ij = 12, \end{cases}$$

where π : Mat_{d-1,d-1} $\rightarrow \mathfrak{g}_0^*$ is the projection onto the last d-3 rows. We conclude:

Lemma 5.2. An element (X_0, X_1, Y_0, Y_1) of M_d belongs to $\mu^{-1}(0)$ if and only if, for every $t \in \mathbb{P}^1$, $X(t)e_2 = Y(t)e_1$ and [X(t), Y(t)] has nonzero entries only in the first two rows.

Remark 5.3. We should like to remark that the moment map equations for \hat{G}_d are formally similar to the complex ADHM equations considered in [10]. The difference is that the $((d-1) \times 2, 2 \times (d-1))$ -part (i, j) (cf. §1.1) of the ADHM-datum is no longer linear in t; i is now constant, while j is quadratic.

Lemma 5.4. The action of \hat{G}_d on $\mu^{-1}(0)$ is free and proper.

Proof. If $g \in \hat{G}_d$ fixes a point (X_0, X_1, Y_0, Y_1) of M_d , then $g(t_0)$ fixes $(X(t_0), Y(t_0))$, for every t_0 . Choosing a t_0 such that $g(t_0) \neq 1$ contradicts Theorem 1.10. Therefore \hat{G}_d acts freely. The moment map for the twisted symplectic form tr $dX(t) \wedge dY(t)$ is $\mu(t) = \mu_{21} + 2\mu_{11}t + \mu_{12}t^2$. Setting $X_0 = \mu(0)^{-1}(0)$, $X_1 = \mu(1)^{-1}(0)$, $X_2 =$ $\mu(\infty)^{-1}(0)$, we have $\mu^{-1}(0) = X_0 \cap X_1 \cap X_2$. For every t_0 , the group $\hat{G}_d(t_0)$ (obtained by evaluating (5.3) at t_0) acts properly on $\mu(t_0)^{-1}(0)$, owing to Theorem 1.10. It follows that, if $m_i \in \mu^{-1}(0)$, $g_i = (g_i^0, u_i(t)) \in \hat{G}_d$, $i \in \mathbb{N}$, and $m_i \to m$, $g_i m_i \to m'$, then both $(g_i^0, u_i(0))$ and $(g_i^0, u_i(\infty))$ have convergent subsequences. Therefore (g_i) has a convergent subsequence and \hat{G}_d acts properly on $\mu^{-1}(0)$. \Box

This lemma implies that $\mu^{-1}(0)/\hat{G}_d$ is a Hausdorff topological space, and that the smooth part of $\mu^{-1}(0)$ is a principal \hat{G}_d -bundle over a complex manifold. Observe now that the fundamental vector field corresponding to $u_0 + tu_1 \in \mathfrak{l}$ is

$$X_u = (-e_1u_0, -e_1u_1, -e_2u_0, -e_2u_1)$$

and, consequently, the subspace of these vector fields is $\operatorname{Mat}_{2,2}(\mathbb{C})$ -invariant, owing to (5.1). Observe also that $g(X_u, X_\rho) = 0$ at points of $\mu^{-1}(0)$, for every $\rho \in \hat{\mathfrak{g}}_d$. Corollary 4.9 implies:

Proposition 5.5. Let $m = (X_0, X_1, Y_0, Y_1) \in \mu^{-1}(0)$ and let \bar{m} denote its image in $\mu^{-1}(0)/\hat{G}_d$.

- (i) m̄ is a smooth point of μ⁻¹(0)/Ĝ_d if and only if the dimension of the subspace of T_mM_d spanned by AX_ρ, A ∈ sl(2, C), ρ ∈ g₀, equals 3 dim g₀;
- (ii) \bar{m} is a smooth point of $\mu^{-1}(0)/\hat{G}_d$ and the \mathbb{C} -hyperkähler structure is nondegenerate at \bar{m} if and only if the quadratic form (5.2) is nondegenerate on the subspace generated by $X_{\rho}, \rho \in \mathfrak{g}_0$.

Let us denote by M_d^0 (resp. M_d^2) the subset of M_d where the condition (i) (resp. (ii)) is satisfied. Thus $\mu^{-1}(0) \cap M_d^0/\hat{G}_d$ is a complex manifold and $\mu^{-1}(0) \cap M_d^2/\hat{G}_d$ is a \mathbb{C} -hyperkähler manifold. We can now state and prove our main result:

Theorem 5.6. There is a natural bijection between R_d and $\mu^{-1}(0)/\hat{G}_d$. This bijection is a biholomorphism between R_d^0 and $(\mu^{-1}(0) \cap M_d^0)/\hat{G}_d$ and a \mathbb{C} -hyperkähler isomorphism between R_d^2 and $(\mu^{-1}(0) \cap M_d^2)/\hat{G}_d$.

Proof. Propositions 3.8 and 3.9 show that there is a natural isomorphism between R_d and sections of Z(E') with $E' \simeq \mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus (d-1)}$. From the construction, Z(E') is the twistor space of $\mu^{-1}(0)/\hat{G}_d$. Sections of Z(E') are projections of sections of K(E'), which in turn correspond to points of $\mu^{-1}(0)$ (as shown in Lemma 5.2). The smooth locus of the (component of) Hilbert scheme of sections of Z(E') is then the smooth locus of $\mu^{-1}(0)/\hat{G}_d$, i.e. $\mu^{-1}(0) \cap M_d^0/\hat{G}_d$, and the locus of sections, the normal bundle of which splits as $\mathcal{O}_{\mathbb{P}^1}(1)^{\oplus 2d}$, is $(\mu^{-1}(0) \cap M_d^2)/\hat{G}_d$. The result follows.

Remark 5.7. The bijection in the first statement of this theorem is an isomorphism if we give $\mu^{-1}(0)/\hat{G}_d$ the scheme structure of the Hilbert scheme of sections of Z(E'). A natural question is whether this scheme structure can be obtained via nonreductive invariant theory [3]. Remark 5.8. One can compute the restriction of (5.2) to the subspace generated by fundamental vector fields, and thus characterise M_d^2 as consisting of points msuch that certain linear operator $L_m : \mathfrak{g}_0 \to \mathfrak{g}_0^*$ is invertible.

5.1. **Real structures.** Up to the action of $PGL(4, \mathbb{C})$, \mathbb{P}^3 has two antilinear involutions:

$$\begin{aligned} \sigma[z_0, z_1, z_2, z_3] &= [-\bar{z}_1, \bar{z}_0, -\bar{z}_3, \bar{z}_2], \\ \sigma'[z_0, z_1, z_2, z_3] &= [\bar{z}_1, \bar{z}_0, \bar{z}_3, \bar{z}_2]. \end{aligned}$$

Both of them preserve $\mathbb{P}^1 = \{[z_0, z_1, 0, 0]\}$, and the space of invariant lines in $\mathbb{P}^3 \setminus \mathbb{P}^1$ is diffeomorphic to \mathbb{R}^4 . In the case of σ , the \mathbb{C} -hyperkähler structure on the space of lines restricts to a flat hyperkähler structure on \mathbb{R}^4 , while in the case of σ' , it restricts to a flat hypersymplectic structure [8]. The involutions σ, σ' act on curves in $\mathbb{P}^3 \setminus \mathbb{P}^1$ and we obtain a pseudo-hyperkähler (resp. hypersymplectic) structure on manifolds of σ -invariant (resp. σ' -invariant) cohomologically stable curves of fixed genus and degree. We want to describe these manifolds in the case of genus 0 curves, i.e. $(R_d^2)^{\sigma}$ and $(R_d^2)^{\sigma'}$. First of all we have:

Proposition 5.9. R_d^{σ} is empty if d is even.

Proof. For C in R_d and the corresponding $\pi : C \to \mathbb{P}^1$ we have, as observed before, $\pi_*\mathcal{O}_C \simeq \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus (d-1)}$. The involution σ induces an antilinear involution on $W = \mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus (d-1)}$, which covers the antipodal map on \mathbb{P}^1 . This, in turn, induces an antilinear involution on $\Lambda^{d-1}W^* \simeq \mathcal{O}_{\mathbb{P}^1}(d-1)$, which covers the antipodal map. Since this involution has no fixed points, the number of zeros of any section of $\Lambda^{d-1}W^*$ must be even.

If d is odd, then there is an induced involution σ on $E' \simeq \mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus (d-1)}$. Modulo conjugation we can write σ in the standard trivialisation over $t \neq \infty$ as

$$\sigma(t; f_1, \dots, f_{d-1}) = (-1/\bar{t}; -\bar{t}\bar{f}_2, \bar{t}\bar{f}_1, -\bar{t}\bar{f}_4, \bar{t}\bar{f}_3, \dots, -\bar{t}\bar{f}_{d-1}, \bar{t}\bar{f}_{d-2}).$$

This, in turn, yields an antiholomorphic involution on $\operatorname{End}(E')$: $M \mapsto \sigma M \sigma$ and, finally, on M_d :

$$(X(t), Y(t)) \mapsto (-\sigma Y(t)\sigma/\bar{t}, \sigma X(t)\sigma/\bar{t}).$$

It follows that:

$$(X_0, X_1, Y_0, Y_1) \mapsto \left(-\tau \bar{Y}_1 \tau, \tau \bar{Y}_0 \tau, \tau \bar{X}_1 \tau, -\tau \bar{X}_0 \tau\right),$$

(5.6) where

$$\tau_{ij} = \begin{cases} -1 & \text{if } j = i+1 \text{ and } i \text{ is odd,} \\ 1 & \text{if } j = i-1 \text{ and } i \text{ is even,} \\ 0 & \text{otherwise.} \end{cases}$$

Consequently:

$$M_d^{\sigma} = \{ (X_0, X_1, Y_0, Y_1) \in M_d; Y_0 = \tau \bar{X}_1 \tau, Y_1 = -\tau \bar{X}_0 \tau \}$$

An easy computation shows that the quadratic form (5.2) restricted to M_d^{σ} is a pseudo-Riemannian metric of signature $(2(d-1)^2 + 2(d-3), 2(d-1)^2 - 2(d-1))$. The subgroup \hat{G}_d^{σ} commuting with σ consists of elements $(h, u) \in G_0 \ltimes L$, where $h = \tau \bar{h} \tau^{-1}$ and $u_{2i-1}(-1/\bar{t}) = -\overline{u_{2i}(t)}/\bar{t}$, $i = 1, \ldots, (d-3)/2$. We conclude from Theorem 5.6: **Theorem 5.10.** Let d be odd. The pseudo-hyperkähler manifold $(R_d^2)^{\sigma}$ of cohomologically stable connected σ -invariant Cohen-Macaulay curves of arithmetic genus 0 in $\mathbb{P}^3 \setminus \mathbb{P}^1$ is isomorphic to the hyperkähler quotient of $(M_d^2)^{\sigma}$ by the group \hat{G}_d^{σ} .

Remark 5.11. It follows from the main result of [6] that the signature of the pseudo-hyperkähler metric on $(R_d^2)^{\sigma}$ is (2d+2, 2d-2).

There is an analogous description of the hypersymplectic manifold $(R_d^2)^{\sigma}$. Essentially, we just have to remove all the minus signs in the above formulae (in particular, d can be arbitrary). We leave the details to the interested reader.

References

- E. Arbarello, M. Cornalba, and Ph.A. Griffiths, *Geometry of algebraic curves, vol. II*, Springer, Berlin, 2011.
- [2] A. Beauville, 'Variétés Kähleriennes dont la première classe de Chern est nulle', J. Differential Geom. 18(4) (1983), 755–782.
- [3] G. Bérczi, B. Doran, T. Hawes, and F. Kirwan, 'Constructing quotients of algebraic varieties by linear algebraic group actions', in: *Handbook of Group Actions, Vol. IV*, International Press, Boston, 2018, 341–446.
- [4] R. Bielawski, 'Hyperkähler manifolds of curves in twistor spaces', SIGMA 10 (2014).
- [5] R. Bielawski and V. Pidstrygach, 'On the symplectic structure of instanton moduli spaces' Adv. Math. 226 (2011), no. 3, 2796–2824.
- [6] R. Bielawski and N. Zhang, 'Hyperkähler geometry of rational curves in twistor spaces', preprint.
- [7] N. Bourbaki, Lie Groups and Lie Algebras, Chapters I-III, Hermann, Paris, 1975.
- [8] A. Dancer and A. Swann, 'Hypersymplectic manifolds', in: Recent Developments in Pseudo-Riemannian Geometry, ESI Lect. Math. Phys., EMS, Zürich, 2008, 97–111.
- [9] P. Etingof and V. Ginzburg, 'Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra', *Invent. Math.* 147 (2002), 243–348.
- [10] B. Frenkel and M. Jardim, 'Complex ADHM equations and sheaves on P³', J. Algebra 319 (2008), 2913–2937.
- [11] F. Ghione and G. Sacchiero, 'Normal bundles of rational curves in P³', Manuscripta Math. 33 (1980), 111–128.
- [12] S.G. Gindikin, 'Generalized conformal structures', in: Twistors in mathematics and physics, Cambridge University Press, Cambridge, 1990, 36–52.
- [13] R. Guralnick, 'A note on pairs of matrices with rank 1 commutator', Lin. & Multilin. Alg. 8 (1979), 97–99.
- [14] R. Hartshorne, Algebraic Geometry, Springer, New York, 1977.
- [15] R. Hartshorne, Deformation Theory, Springer, New York, 2010.
- [16] A.A. Henni and M. Jardim, 'Commuting matrices and the Hilbert scheme of points on affine spaces', Adv. Geom. 18 (2018), 467–482.
- [17] N.J. Hitchin, 'Hypersymplectic quotients', Acta Acad. Sci. Tauriensis, 124 Supp. (1990), 169–180.
- [18] N.J. Hitchin, A. Karlhede, U. Lindström, and M. Roček, 'Hyperkähler metrics and supersymmetry', Comm. Math. Phys. 108(4) (1987), 535–589.
- [19] M. Jardim and M. Verbitsky, 'Trihyperkähler reduction and instanton bundles on CP³', Compositio Math. 150 (2014), 1836–1868.
- [20] D. Joyce, 'The hypercomplex quotient and the quaternionic quotient', Math. Ann. 290 (1991), 323–340.
- [21] D. Mumford, J. Fogarty, and F. Kirwan, Geometric invariant theory, Springer, Berlin, 1994.
- [22] H. Nakajima, Lectures on Hilbert Schemes of Points on Surfaces, AMS, Providence, 1999.
- [23] M.G. Neubauer and D.J. Saltman, 'Two-generated commutative subalgebras of $M_n(F)$ ', J. Algebra 164(2) (1994), 545–562.

INSTITUT FÜR DIFFERENTIALGEOMETRIE, LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1, 30167 HANNOVER, GERMANY