

NONDEGENERACY OF THE BUBBLE FOR THE CRITICAL p -LAPLACE EQUATION

ANGELA PISTOIA AND GIUSI VAIRA

ABSTRACT. We prove the non-degeneracy of the extremals of the Sobolev inequality

$$\int_{\mathbb{R}^N} |\nabla u|^p dx \geq \mathcal{S}_p \int_{\mathbb{R}^N} |u|^{\frac{Np}{N-p}} dx, \quad u \in \mathcal{D}^{1,p}(\mathbb{R}^N)$$

when $1 < p < N$, as solutions of a critical quasilinear equation involving the p -Laplacian.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

In this paper we establish the *linear non-degeneracy* of the extremals of the optimal classical Sobolev inequality

$$\mathcal{S}_p \|u\|_{L^{p^*}(\mathbb{R}^N)} \leq \|\nabla u\|_{L^p(\mathbb{R}^N)} \quad \text{for any } u \in \mathcal{D}^{1,p}(\mathbb{R}^N), \quad (1.1)$$

where $p^* := \frac{Np}{N-p}$ and $1 < p < N$.

Aubin [1] and Talenti [24] found the optimal constant and the extremals for inequality (1.1). Indeed, equality is achieved precisely by the functions

$$U_{\delta,\xi}(x) := \delta^{-\frac{N-p}{p}} U\left(\frac{x-\xi}{\delta}\right) \quad \text{where } \delta > 0, \xi \in \mathbb{R}^N \quad (1.2)$$

where

$$U(x) = \left(\frac{\alpha_{N,p}}{1 + |x - \xi|^{\frac{p}{p-1}}} \right)^{\frac{N-p}{p}} \quad \text{with } \alpha_{N,p} := N^{\frac{1}{p}} \left(\frac{N-p}{p-1} \right)^{\frac{p-1}{p}}, \quad (1.3)$$

which solve the critical equation

$$-\Delta_p u = u^{p^*-1} \text{ in } \mathbb{R}^N, \quad u > 0 \text{ in } \mathbb{R}^N, \quad u \in \mathcal{D}^{1,p}(\mathbb{R}^N). \quad (1.4)$$

All the solutions to the equation (1.4) are indeed the only ones of (1.2). Caffarelli, Gidas and Spruck proved the claim when $p = 2$. The case $p \neq 2$ has been firstly solved by Guedda and Veron [15] in the radial case, where the authors classified all the positive radial solutions and successively by Damascelli, Merchán, Montoro and Sciunzi [8] when $\frac{2N}{N+2} \leq p < 2$, by Vetóis [25] and Damascelli and Ramaswamy [7] when $1 < p < 2$ and finally by Sciunzi [23] in the remaining cases, namely when $2 < p < N$.

Here we are interested in the linear non-degeneracy of the solutions (1.2) to equation (1.4). Let us point out that equation (1.4) is invariant by scaling and by translations. Therefore, if we differentiate the equation

$$-\Delta_p U_{\delta,\xi} = U_{\delta,\xi}^{p^*-1} \text{ in } \mathbb{R}^N$$

with respect to the parameters δ and ξ_1, \dots, ξ_N at $\delta = 1$ and $\xi = 0$ we see that the functions

$$Z_0(x) := -\partial_\delta U_{\delta,\xi}|_{\delta=1,\xi=0} = \frac{N-p}{p} U + x \cdot \nabla U \quad (1.5)$$

2010 *Mathematics Subject Classification.* 35J60 (primary), and 35B33, 35J20 (secondary).

and

$$Z_i(x) := \partial_{\xi_i} U_{\delta, \xi}|_{\delta=1, \xi=0} = -\partial_{x_i} U, \quad i = 1, \dots, N \quad (1.6)$$

annihilate the linearized operator around the function U defined in (1.3), namely they solve the linear equation

$$-\operatorname{div}(|\nabla U|^{p-2} \nabla \phi) - (p-2) \operatorname{div}(|\nabla U|^{p-4} (\nabla U, \nabla \phi) \nabla U) = (p^* - 1) U^{p^*-2} \phi \text{ in } \mathbb{R}^N. \quad (1.7)$$

We say that U is non-degenerate if the kernel of the associated linearized operator (1.7) is spanned only by the functions Z_i 's defined in (1.5) and (1.6). This property is true when $p = 2$ as it was established by Rey in [22]. Our main result extends the non-degeneracy of the solution U to any $p \in (1, N)$ in the weighted Sobolev space $\mathcal{D}_*^{1,2}(\mathbb{R}^N)$, which is defined as the completion of $C_c^1(\mathbb{R}^N)$ with respect to the norm

$$\|\phi\| := \left(\int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx \right)^{1/2}.$$

Theorem 1.1. *The solution*

$$U(x) = \left(\frac{\alpha_{N,p}}{1 + |x|^{\frac{p}{p-1}}} \right)^{\frac{N-p}{p}} \quad \text{with } \alpha_{N,p} := N^{\frac{1}{p}} \left(\frac{N-p}{p-1} \right)^{\frac{p-1}{p}}$$

of equation (1.4) is non-degenerate in the sense that all the solutions of the equation (1.7) in the space $\mathcal{D}_*^{1,2}(\mathbb{R}^N)$ are linear combination of the functions

$$Z_0(x) = \frac{N-p}{p} U + x \cdot \nabla U, \quad Z_1(x) = \partial_{x_1} U(x), \dots, \quad Z_N(x) = \partial_{x_N} U(x).$$

The structure of the linearized equation (1.7) strongly suggests to introduce the space $\mathcal{D}_*^{1,2}(\mathbb{R}^N)$. A similar first order Sobolev space with weight was introduced by Damascelli and Sciunzi in [9] to study a linearized operator on a bounded domain. Here the situation is much more delicate due to the unboundness of the domain. Section 2 is devoted to prove some properties of $\mathcal{D}_*^{1,2}(\mathbb{R}^N)$ which are essential to get Theorem 1.1 whose proof is carried out in Section 3.

Quasilinear equations with critical growth involving the p -Laplace operator have been widely studied in recent years using a variational framework, starting from the quasilinear version of the classical Brezis-Nirenberg problem (see [2]) studied by Guedda and Veron in [16]. In particular, we would like to focus on the problem of the existence of sign-changing solutions to the critical equation

$$-\Delta_p u = |u|^{p^*-2} u \text{ in } \Omega, \quad (1.8)$$

where Ω is either the whole space \mathbb{R}^N or a bounded smooth domain in \mathbb{R}^N in which case we assume homogeneous Dirichlet boundary conditions. As far as we know the only result concerning existence of sign-changing solutions to (1.8) in the whole space is due to Clapp and Lopez Rios in [5], where they prove that (1.8) has a certain finite number (depending on the dimension N) of non-radial sign-changing solutions. On the other hand if $p = 2$ del Pino, Musso, Pacard, and Pistoia in [10, 11] used the Lyapunov-Schmidt procedure to build infinitely many sign-changing solutions which look like a positive bubble crowned by an arbitrary large number of negative bubbles arranged on a regular polygon. It would be interesting to check if it is possible to build this kind of solutions in the quasilinear case. When Ω is a bounded domain, the existence of solutions is a more delicate issue. Indeed if Ω is starshaped the problem does not have any solutions because of a Pohozaev identity obtained by Guedda and Veron in [16]. The existence of a positive solution has been proved by Mercuri, Sciunzi and Squassina in [18] when the domain has a small hole, in the same spirit of Coron's result [6] when $p = 2$. The existence of a sign-changing solution has been

obtained by Mercuri and Pacella in [17] when the domain Ω has either a small hole and little symmetry or a hole of any size and more symmetry. On the other hand, if $p = 2$ and Ω has a small hole, Musso and Pistoia in [20] (see also [13, 12]) used the Lyapunov-Schmidt procedure to built sign-changing solutions which look like the superposition of bubbles with alternating sign whose number becomes arbitrary large as the size of the hole approaches zero. It is natural to ask if this kind of solutions do exist also in the quasilinear case. In both cases the understanding of the linear non-degeneracy of the bubble is the first step in the application of the Ljapunov-Schmidt procedure.

Acknowledgements. We wish to thank professor Berardino Sciunzi for many helpful comments and discussions. Moreover, we warmly thank the anonymous referee for his/her valuable comments which allow us to improve the presentation of the paper.

2. A SUITABLE WEIGHTED SOBOLEV SPACE

First of all, let us point out the following fact.

Lemma 2.1. (i) *If $p \in (1, 2)$ there exists $C > 0$ such that*

$$\left(\int_{\mathbb{R}^N} |\phi|^{\frac{Np}{N-p}} dx \right)^{\frac{N-p}{p}} \leq C \left(\int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx \right)^{\frac{1}{2}} \text{ for any } \phi \in C_c^1(\mathbb{R}^N).$$

(ii) *If $p \in (2, N)$ for any $R > 0$ there exists $C(R) > 0$ such that*

$$\left(\int_{\mathbb{R}^N \setminus B_R(0)} |x|^{-\frac{Np+p-2N}{p-1}} |\phi|^2 dx \right)^{\frac{1}{2}} \leq C(R) \left(\int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx \right)^{\frac{1}{2}} \text{ for any } \phi \in C_c^1(\mathbb{R}^N).$$

Proof. To get (i) it is useful to recall the Caffarelli-Kohn-Nirenberg inequality (see [4]): if $r, q \geq 1$, $\frac{1}{r} + \frac{\gamma}{N} = \frac{1}{q} + \frac{\alpha-1}{N} > 0$ then for any $\phi \in C_c^1(\mathbb{R}^N)$

$$\| |x|^\gamma \phi \|_{L^r(\mathbb{R}^N)} \leq c \| |x|^\alpha |\nabla \phi| \|_{L^q(\mathbb{R}^N)}. \quad (2.1)$$

Then we apply (2.1) with $\gamma = 0$, $r = \frac{Np}{N-p}$, $q = 2$ and $\alpha = \frac{N(2-p)}{2p}$

$$\left(\int_{\mathbb{R}^N} |\phi|^{\frac{Np}{N-p}} dx \right)^{\frac{(N-p)}{Np}} \leq c \left(\int_{\mathbb{R}^N} |x|^{\frac{N(2-p)}{p}} |\nabla \phi|^2 dx \right)^{1/2} \leq c \left(\int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx \right)^{\frac{1}{2}}.$$

and the claim follows since if $p < 2$ there exists a constant c such that

$$|x|^{\frac{N(2-p)}{p}} \leq c \frac{|x|^{\frac{p-2}{p-1}}}{\left(1 + |x|^{\frac{p}{p-1}}\right)^{\frac{N(p-2)}{p}}} \text{ for any } x \in \mathbb{R}. \quad (2.2)$$

To get (ii) it is useful to recall the weighted Hardy-Sobolev inequality (see for example Lemma 2.3 in [7]):

if $q \geq 1$, $s > q - N$ and $R \geq 0$ then

$$\int_{\mathbb{R}^N \setminus B_R(0)} |x|^{s-q} |\varphi|^q \leq c(N, q, s) \int_{\mathbb{R}^N \setminus B_R(0)} |x|^s |\nabla \varphi|^q dx \text{ for any } \varphi \in C_c^1(\mathbb{R}^N). \quad (2.3)$$

Then we apply (2.3) with $s = -\frac{(N-1)(p-2)}{p-1}$ and $q = 2$ (note that $s - q > -N$ because $p < N$)

$$\int_{\mathbb{R}^N \setminus B_R(0)} |x|^{-\frac{(N-1)(p-2)}{p-1}-2} |\varphi|^2 \leq c \int_{\mathbb{R}^N \setminus B_R(0)} |x|^{-\frac{(N-1)(p-2)}{p-1}} |\nabla \varphi|^2 \leq c(R) \int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2.$$

and the claim follows since if $p > 2$ for any $R > 0$ there exists $c(R)$ such that

$$|\nabla U(x)| \geq c|x|^{\frac{N-1}{p-1}} \text{ if } |x| \geq R.$$

□

Lemma 2.1 allows us to define the Hilbert space $\mathcal{D}_*^{1,2}(\mathbb{R}^N)$, which is defined as the completion of $C_c^1(\mathbb{R}^N)$ with respect to the norm $\|\phi\| := \left(\int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx \right)^{1/2}$ induced by the scalar product

$$\langle \phi, \psi \rangle := \int_{\mathbb{R}^N} |\nabla U|^{p-2} (\nabla \phi, \nabla \psi) dx.$$

Now, we can look for a weak solution $\phi \in \mathcal{D}_*^{1,2}(\mathbb{R}^N)$ to the linear equation (1.7), namely

$$\begin{aligned} & \int_{\mathbb{R}^N} |\nabla U|^{p-2} (\nabla \phi, \nabla \psi) dx + (p-2) \int_{\mathbb{R}^N} |\nabla U|^{p-4} (\nabla U, \nabla \phi) (\nabla U, \nabla \psi) dx \\ &= \frac{Np - N + p}{N - p} \int_{\mathbb{R}^N} U^{\frac{Np-2N+2p}{N-p}} \phi \psi dx \text{ for any } \psi \in \mathcal{D}_*^{1,2}(\mathbb{R}^N). \end{aligned} \quad (2.4)$$

All the integrals involved in (2.4) are finite. Indeed, the integrals in the L.H.S. can be easily estimated using Hölder inequality and Cauchy-Schwarz inequality. The finiteness of the integral in the R.H.S. is more delicate and follows by the continuous embedding of the weighted space $\mathcal{D}_*^{1,2}(\mathbb{R}^N)$ into the weighted space

$$I_*^2(\mathbb{R}^N) = \left\{ \phi : \int_{\mathbb{R}^N} U^{\frac{Np-2N+2p}{N-p}} \phi^2 dx < +\infty \right\}, \quad (2.5)$$

which is stated in the following result.

Proposition 2.2. *There exists $C > 0$ such that*

$$\int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx \geq C \int_{\mathbb{R}^N} U^{\frac{Np}{N-p}-2} \phi^2 dx \text{ for any } \phi \in \mathcal{D}_*^{1,2}(\mathbb{R}^N). \quad (2.6)$$

Proof. We will prove (2.6) for any $\phi \in C_c^1(\mathbb{R}^N)$. The statement will follow by a density argument. Throughout the proof c will denote a constant (possibly depending on the parameters) which may change from line to line. Although we will not estimate the constants explicitly, it will be clear from the arguments that our claims hold.

It is useful to remind that

$$U(x) = c \frac{1}{\left(1 + |x|^{\frac{p}{p-1}}\right)^{\frac{N(p-2)+2p}{p}}} \quad \text{and} \quad |\nabla U(x)|^{p-2} = c \frac{|x|^{\frac{p-2}{p-1}}}{\left(1 + |x|^{\frac{p}{p-1}}\right)^{\frac{N(p-2)}{p}}}.$$

We distinguish 3 cases.

- The case $\frac{2N}{N+2} < p < 2$.

We remark that since $\frac{2N}{N+2} < p$ Hölder's inequality implies

$$\int_{\mathbb{R}^N} \frac{1}{\left(1 + |x|^{\frac{p}{p-1}}\right)^{\frac{N(p-2)+2p}{p}}} |\phi|^2 \leq \left(\int_{\mathbb{R}^N} \frac{1}{\left(1 + |x|^{\frac{p}{p-1}}\right)^N} \right)^{\frac{N(p-2)+2p}{Np}} \left(\int_{\mathbb{R}^N} |\phi|^{\frac{Np}{N-p}} \right)^{\frac{2(N-p)}{Np}}.$$

Now, we apply Caffarelli-Kohn-Nirenberg's inequality (2.1) (with $\gamma = 0$, $r = \frac{Np}{N-p}$, $q = 2$ and $\alpha = \frac{N(2-p)}{2p}$) and we get

$$\left(\int_{\mathbb{R}^N} |\phi|^{\frac{Np}{N-p}} \right)^{\frac{(N-p)}{Np}} \leq c \left(\int_{\mathbb{R}^N} |x|^{\frac{N(2-p)}{p}} |\nabla \phi|^2 \right)^{1/2}.$$

The claim follows because of (2.2).

- *The case $1 < p \leq \frac{2N}{N+2}$.*

In this case $\frac{N(p-2)+2p}{p} \leq 0$ and so

$$\left(1 + |x|^{\frac{p}{p-1}} \right)^{-\frac{N(p-2)+2p}{p}} \leq c \left(1 + |x|^{-\frac{N(p-2)+2p}{(p-1)}} \right).$$

Then

$$\int_{\mathbb{R}^N} \frac{1}{\left(1 + |x|^{\frac{p}{p-1}} \right)^{\frac{N(p-2)+2p}{p}}} |\phi|^2 \leq c \int_{\mathbb{R}^N} |\phi|^2 + c \int_{\mathbb{R}^N} |x|^{-\frac{N(p-2)+2p}{(p-1)}} |\phi|^2.$$

Now, we apply Caffarelli-Kohn-Nirenberg's inequality and we get (with $\gamma = 0$, $r = q = 2$ and $\alpha = 2$)

$$\left(\int_{\mathbb{R}^N} |\phi|^2 \right)^{1/2} \leq c \left(\int_{\mathbb{R}^N} |x|^2 |\nabla \phi|^2 \right)^{1/2}.$$

and also (with $\gamma = -\frac{N(p-2)+2p}{2(p-1)}$, $r = q = 2$ and $\alpha = 1 - \frac{N(p-2)+2p}{2(p-1)} = \frac{-2-Np+2N}{2(p-1)}$)

$$\left(\int_{\mathbb{R}^N} |x|^{-\frac{N(p-2)+2p}{(p-1)}} |\phi|^2 \right)^{1/2} \leq c \left(\int_{\mathbb{R}^N} |x|^{\frac{-2-Np+2N}{(p-1)}} |\nabla \phi|^2 \right)^{1/2}.$$

Then

$$\int_{\mathbb{R}^N} \frac{1}{\left(1 + |x|^{\frac{p}{p-1}} \right)^{\frac{N(p-2)+2p}{p}}} |\phi|^2 \leq c \int_{\mathbb{R}^N} \left(|x|^2 + |x|^{\frac{-2-Np+2N}{(p-1)}} \right) |\nabla \phi|^2.$$

The claim follows because if $p \leq \frac{2N}{N+2}$ it is easy to check that there exists a constant c such that

$$|x|^2 + |x|^{\frac{-2-Np+2N}{(p-1)}} \leq c \frac{|x|^{\frac{p-2}{p-1}}}{\left(1 + |x|^{\frac{p}{p-1}} \right)^{\frac{N(p-2)}{p}}} \text{ for any } x \in \mathbb{R}.$$

- *The case $p > 2$.*

The proof in this case is much more delicate because the weight $|\nabla U|^{p-2}$ has different decay as $|x| \rightarrow 0$ or $|x| \rightarrow \infty$. Let $m \geq 1$ be a fixed integer. We can write

$$\int_{\mathbb{R}^N} U^{p^*-2} \phi^2 dx = \underbrace{\int_{\mathbb{R}^N \setminus B_{2^m}(0)} U^{p^*-2} \phi^2 dx}_{(I)} + \underbrace{\int_{B_{2^m}(0)} U^{p^*-2} \phi^2 dx}_{(II)},$$

where $B_{2^m}(0)$ is the ball centered at the origin with radius 2^m .

First, we estimate (I). We remark that there exists constants c_1, \dots, c_4 such that

$$\frac{c_1}{|x|^{\frac{N-p}{p-1}}} \leq U(x) \leq \frac{c_2}{|x|^{\frac{N-p}{p-1}}} \text{ and } \frac{c_3}{|x|^{\frac{N-1}{p-1}}} \leq |\nabla U| \leq \frac{c_4}{|x|^{\frac{N-1}{p-1}}} \text{ if } |x| \geq 2^m. \quad (2.7)$$

Therefore

$$\begin{aligned} & \int_{\mathbb{R}^N \setminus B_{2^m}(0)} U^{p^*-2} \phi^2 dx \\ & \quad (\text{we use (2.7)}) \\ & \leq c \int_{\mathbb{R}^N \setminus B_{2^m}(0)} \frac{1}{|x|^{\frac{N-p}{p-1}(p^*-2)}} \phi^2 dx \\ & \quad (\text{we set } s = -\frac{N-1}{p-1}(p-2) > 2-N \text{ and } \beta = (p^*-2)\frac{N-p}{p-1} + s - 2 > 0) \\ & = c \int_{\mathbb{R}^N \setminus B_{2^m}(0)} \frac{1}{|x|^\beta} |x|^{s-2} \phi^2 dx \\ & \quad (\text{we use (2.3) with } q = 2 \text{ and } R = 2^m) \\ & \leq c \int_{\mathbb{R}^N \setminus B_{2^m}(0)} |x|^s |\nabla \phi|^2 dx \\ & \quad (\text{we use (2.7)}) \\ & \leq c \int_{\mathbb{R}^N \setminus B_{2^m}(0)} |\nabla U|^{p-2} |\nabla \phi|^2 dx \end{aligned}$$

and so

$$\int_{\mathbb{R}^N \setminus B_{2^m}(0)} U^{p^*-2} \phi^2 dx \leq c \int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx \quad (2.8)$$

Now, we estimate (II). Firstly, given $A := \{x \in \mathbb{R}^N : 1 < |x| \leq 2\}$, it is useful to recall the standard interpolation inequality (see for example [19])

$$\int_A |u - u_A|^m dx \leq c \left(\int_A |\nabla u|^r dx \right)^{\frac{m}{r}} \quad (2.9)$$

where

$$\frac{1}{m} = \frac{1}{r} - \frac{1}{N} > 0 \text{ and } u_A := \frac{1}{|A|} \int_A u dx.$$

Therefore, if $r = 2$, $m = \frac{2N}{N-2} > 0$ by Hölder inequality we immediately deduce

$$\int_A |u - u_A|^2 dx \leq \left(\int_A 1^{\frac{m}{m-2}} dx \right)^{\frac{m-2}{m}} \left(\int_A |u - u_A|^m dx \right)^{\frac{2}{m}} \leq c |A|^{\frac{2}{N}} \int_A |\nabla u|^2 dx.$$

Moreover, if $\lambda > 0$ and $\lambda A := \{\lambda x : x \in A\}$, a simply scaling gives

$$\int_{\lambda A} |u - u_{\lambda A}|^2 dx \leq c |A|^{\frac{2}{N}} \lambda^2 \int_{\lambda A} |\nabla u|^2 dx \quad (2.10)$$

Now, let us introduce a sequence of disjoint annuli $A_k := \{x \in \mathbb{R}^N : 2^k < |x| \leq 2^{k+1}\}$ which covers the ball $B_{2^m}(0)$, namely $A_k \cap A_h = \emptyset$ for $h \neq k$ and

$$B_{2^m}(0) = \bigcup_{k=-\infty}^{m-1} A_k,$$

so that

$$\int_{B_{2^m}(0)} U^{p^*-2} \phi^2 dx = \sum_{k=-\infty}^m \int_{A_k} U^{p^*-2} \phi^2 dx. \quad (2.11)$$

We are going to estimate each term in the sum of R.H.S. of (2.11), taking into account that

$$\inf_{A_k} |\nabla U|^{p-2} \geq c \frac{2^{k \frac{p-2}{p-1}}}{(1 + 2^{k \frac{p}{p-1}})^{\frac{N}{p}(p-2)}}. \quad (2.12)$$

We have

$$\begin{aligned} & \int_{A_k} U^{p^*-2} \phi^2 dx \\ & \text{(since } p \geq 2, \sup_{x \in \mathbb{R}^N} |x|^{\frac{p}{p-1}} U^{p^*-2} = L) \\ & \leq L \int_{A_k} |x|^{-\frac{p}{p-1}} \phi^2 dx \\ & \leq c \int_{A_k} |x|^{-\frac{p}{p-1}} |\phi - \phi_{A_k}|^2 dx + c \int_{A_k} |x|^{-\frac{p}{p-1}} |\phi_{A_k}|^2 dx \\ & \leq c \int_{A_k} |x|^{-\frac{p}{p-1}} |\phi - \phi_{A_k}|^2 dx + c 2^{k(-\frac{p}{p-1}+N)} |\phi_{A_k}|^2 \end{aligned} \quad (2.13)$$

and

$$\begin{aligned} & \int_{A_k} |x|^{-\frac{p}{p-1}} |\phi - \phi_{A_k}|^2 dx \leq c 2^{-k \frac{p}{p-1}} \int_{A_k} |\phi - \phi_{A_k}|^2 dx \\ & \text{(we use (2.10) with } \lambda = 2^k) \\ & \leq c 2^{k(-\frac{p}{p-1}+2)} \int_{A_k} |\nabla \phi|^2 dx \\ & \text{(we use (2.12))} \\ & \leq c 2^{k(-\frac{p}{p-1}+2)} \frac{(1 + 2^{k \frac{p}{p-1}})^{\frac{N}{p}(p-2)}}{2^{k \frac{p-2}{p-1}}} \int_{A_k} |\nabla U|^{p-2} |\nabla \phi|^2 dx \\ & = c(1 + 2^{k \frac{p}{p-1}})^{\frac{N}{p}(p-2)} \int_{A_k} |\nabla U|^{p-2} |\nabla \phi|^2 dx. \end{aligned} \quad (2.14)$$

Combining (2.13) and (2.14) we get

$$\int_{A_k} U^{p^*-2} \phi^2 dx \leq c(1 + 2^{k \frac{p}{p-1}})^{\frac{N}{p}(p-2)} \int_{A_k} |\nabla U|^{p-2} |\nabla \phi|^2 dx + c 2^{k(-\frac{p}{p-1}+N)} |\phi_{A_k}|^2$$

and summing upon k

$$\begin{aligned}
 & \int_{B_{2^m}} U^{p^*-2} \phi^2 dx \\
 &= \sum_{k=-\infty}^{m-1} \int_{A_k} U^{p^*-2} \phi^2 dx \\
 &\leq c \sum_{k=-\infty}^{m-1} \left[\left(1 + 2^{k \frac{p}{p-1}}\right)^{\frac{N}{p}(p-2)} \int_{A_k} |\nabla U|^{p-2} |\nabla \phi|^2 dx \right] \\
 &+ c \sum_{k=-\infty}^{m-1} 2^{k(-\frac{p}{p-1}+N)} |\phi_{A_k}|^2 \\
 &\leq c \sum_{k=-\infty}^{m-1} \int_{A_k} |\nabla U|^{p-2} |\nabla \phi|^2 dx + c \sum_{k=-\infty}^{m-1} \left[2^{kN \frac{p-2}{p-1}} \int_{A_k} |\nabla U|^{p-2} |\nabla \phi|^2 dx \right] \\
 &+ c \sum_{k=-\infty}^{m-1} 2^{k(-\frac{p}{p-1}+N)} |\phi_{A_k}|^2 \\
 &\leq c \int_{B_{2^m}} |\nabla U|^{p-2} |\nabla \phi|^2 dx + c \underbrace{\sum_{k=-\infty}^{m-1} 2^{kN \frac{p-2}{p-1}}}_{<+\infty} \int_{B_{2^m}} |\nabla U|^{p-2} |\nabla \phi|^2 dx + \\
 &+ c \sum_{k=-\infty}^{m-1} 2^{k(-\frac{p}{p-1}+N)} |\phi_{A_k}|^2 \\
 &\leq c \int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx + c \sum_{k=-\infty}^{m-1} 2^{k(-\frac{p}{p-1}+N)} |\phi_{A_k}|^2
 \end{aligned} \tag{2.15}$$

It remains to estimate the last term of (2.15), namely $\sum_{k=-\infty}^{m-1} 2^{k(-\frac{p}{p-1}+N)} |\phi_{A_k}|^2$. Now

$$\begin{aligned}
 & \int_{A_k \cup A_{k+1}} |\phi - \phi_{A_k \cup A_{k+1}}|^2 dx \\
 &= \frac{1}{|A_k| + |A_{k+1}|} \left(\int_{A_k} |\phi - \phi_{A_k \cup A_{k+1}}|^2 dx + \int_{A_{k+1}} |\phi - \phi_{A_k \cup A_{k+1}}|^2 dx \right) \\
 &\geq \int_{A_k} |\phi - \phi_{A_k \cup A_{k+1}}|^2 dx \\
 &\geq \left| \int_{A_k} (\phi - \phi_{A_k \cup A_{k+1}}) dx \right|^2 \\
 &= \left| \int_{A_k} \phi dx - \frac{|A_k|}{|A_{k+1}| + |A_k|} \int_{A_k} \phi dx - \frac{|A_k|}{|A_{k+1}| + |A_k|} \int_{A_{k+1}} \phi dx \right|^2 \\
 &= \frac{1}{|A_k| + |A_{k+1}|} \left| |A_{k+1}| \int_{A_k} \phi dx - |A_k| \int_{A_{k+1}} \phi dx \right|^2 \\
 &= \frac{|A_k| |A_{k+1}|}{|A_k| + |A_{k+1}|} |\phi_{A_k} - \phi_{A_{k+1}}|^2
 \end{aligned}$$

and so

$$\begin{aligned}
|\phi_{A_k} - \phi_{A_{k+1}}|^2 &\leq c \frac{|A_k| + |A_{k+1}|}{|A_k||A_{k+1}|} \int_{A_k \cup A_{k+1}} |\phi - \phi_{A_k \cup A_{k+1}}|^2 dx \\
&\quad (\text{taking into account that } |A_k| = c2^{kN} (2^N - 1)) \\
&\leq c2^{-kN} \int_{A_k \cup A_{k+1}} |\phi - \phi_{A_k \cup A_{k+1}}|^2 dx \\
&\quad (\text{we use (2.10) with } \lambda = 2^k) \\
&\leq c2^{k(2-N)} \int_{A_k \cup A_{k+1}} |\nabla \phi|^2 dx \\
&\quad (\text{we use (2.12) }) \\
&\leq c2^{k(2-N)} (1 + 2^{k \frac{p}{p-1}})^{\frac{N(p-2)}{p}} 2^{-k \frac{p-2}{p-1}} \int_{A_k \cup A_{k+1}} |\nabla U|^{p-2} |\nabla \phi|^2 dx \\
&\leq c \left(2^{k \frac{p-Np+N}{p-1}} + 2^{k \frac{p-N}{p-1}} \right) \int_{A_k \cup A_{k+1}} |\nabla U|^{p-2} |\nabla \phi|^2 dx.
\end{aligned} \tag{2.16}$$

Now, we use the simple fact that for any $\eta > 0$ the following inequality holds

$$(a + b)^2 \leq (1 + \eta)a^2 + \left(\frac{\eta + 1}{\eta} \right) b^2 \text{ for any } a, b \in \mathbb{R}.$$

Then, if we choose $\eta > 0$ so that $1 + \eta = \eta_0 2^{-\frac{p}{p-1} + N}$ where $\eta_0 = \frac{2}{1 + 2^{-\frac{p}{p-1} + N}} < 1$, (this is possible because $N - \frac{p}{p-1} > 0$, since $p \geq 2$), we get

$$|\phi_{A_k}|^2 = |\phi_{A_k} - \phi_{A_{k+1}} + \phi_{A_{k+1}}|^2 \leq \eta_0 2^{-\frac{p}{p-1} + N} |\phi_{A_{k+1}}|^2 + \frac{\eta + 1}{\eta} |\phi_{A_k} - \phi_{A_{k+1}}|^2.$$

and using (2.16) we deduce

$$\begin{aligned}
&2^{k(-\frac{p}{p-1} + N)} |\phi_{A_k}|^2 \\
&\leq 2^{k(-\frac{p}{p-1} + N)} 2^{(k+1)(-\frac{p}{p-1} + N)} \eta_0 |\phi_{A_{k+1}}|^2 \\
&\quad + c2^{k(-\frac{p}{p-1} + N)} \left(2^{k \frac{p-Np+N}{p-1}} + 2^{k \frac{p-N}{p-1}} \right) \int_{A_k \cup A_{k+1}} |\nabla U|^{p-2} |\nabla \phi|^2 dx \\
&= 2^{(k+1)(-\frac{p}{p-1} + N)} \eta_0 |\phi_{A_{k+1}}|^2 + c \left(1 + 2^{kN \frac{p-2}{p-1}} \right) \int_{A_k \cup A_{k+1}} |\nabla U|^{p-2} |\nabla \phi|^2 dx
\end{aligned}$$

We sum upon k and we get

$$\begin{aligned}
&\sum_{k=-\infty}^{m-1} 2^{k(-\frac{p}{p-1} + N)} |\phi_{A_k}|^2 \\
&\leq \eta_0 \sum_{k=-\infty}^{m-1} 2^{(k+1)(-\frac{p}{p-1} + N)} |\phi_{A_{k+1}}|^2 + \sum_{k=-\infty}^{m-1} \left(1 + 2^{kN \frac{p-2}{p-1}} \right) \int_{A_k \cup A_{k+1}} |\nabla U|^{p-2} |\nabla \phi|^2 dx \\
&\leq \eta_0 \sum_{k=-\infty}^m 2^{k(-\frac{p}{p-1} + N)} |\phi_{A_k}|^2 + \underbrace{\left(1 + \sum_{k=-\infty}^{m-1} 2^{kN \frac{p-2}{p-1}} \right)}_{< +\infty} \int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx,
\end{aligned}$$

which implies

$$(1 - \eta_0) \sum_{k=-\infty}^{m-1} 2^{k(-\frac{p}{p-1} + N)} |\phi_{A_k}|^2 \leq c |\phi_{A_m}|^2 + c \int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx. \tag{2.17}$$

On the other hand, we have

$$\begin{aligned}
& |\phi_{A_m}|^2 \quad (\text{we use Hölder's inequality}) \\
& \leq c \int_{A_m} |x|^{-\frac{(N-1)(p-2)}{p-1}} \phi^2 dx \quad (\text{the annulus } A_m \subset \mathbb{R}^N \setminus B_{2^m(0)}) \\
& \leq c \int_{\mathbb{R}^N \setminus B_{2^m(0)}} |x|^{-\frac{(N-1)(p-2)}{p-1}} \phi^2 dx \quad (\text{we use (2.8)}) \\
& \leq c \int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx.
\end{aligned} \tag{2.18}$$

Finally, combining (2.15) with (2.17) (remember that $\eta_0 < 1$) and (2.18) we get

$$\int_{B_{2^m(0)}} U^{p^*-2} \phi^2 dx \leq c \int_{\mathbb{R}^N} |\nabla U|^{p-2} |\nabla \phi|^2 dx. \tag{2.19}$$

□

3. PROOF OF THEOREM 1.1

3.1. A wave decomposition. First of all, let us rewrite the linear equation (1.7) as

$$|x|^2 \Delta \phi + (p-2) \sum_{i,j=1}^N \partial_{ij}^2 \phi x_i x_j + \frac{(p-2)N}{1+|x|^{\frac{p}{p-1}}} (\nabla \phi, x) + \gamma_{N,p} \frac{|x|^{\frac{p}{p-1}}}{(1+|x|^{\frac{p}{p-1}})^2} \phi = 0 \tag{3.1}$$

where $\gamma_{N,p} := N \frac{Np-N+p}{p-1}$. Indeed a straightforward computation shows that

$$\begin{aligned}
& \operatorname{div} (|\nabla U|^{p-2} \nabla \phi) + (p-2) \operatorname{div} (|\nabla U|^{p-4} (\nabla U, \nabla \phi) \nabla U) \\
& = |\nabla U|^{p-2} \Delta \phi + (\nabla |\nabla U|^{p-2}, \nabla \phi) \\
& + (p-2) |\nabla U|^{p-4} (\nabla U, \nabla \phi) \Delta U \\
& + (p-2) (\nabla U, \nabla \phi) (\nabla |\nabla U|^{p-4}, \nabla U) \\
& + (p-2) |\nabla U|^{p-4} (\nabla (\nabla U, \nabla \phi), \nabla U).
\end{aligned}$$

and

$$\begin{aligned}
 \nabla U &= -c_{N,p} \frac{|x|^{\frac{2-p}{p-1}} x}{(1 + |x|^{\frac{p}{p-1}})^{\frac{N}{p}}} \\
 |\nabla U|^{p-4} &= c_{N,p}^{p-4} \frac{|x|^{\frac{p-4}{p-1}}}{(1 + |x|^{\frac{p}{p-1}})^{\frac{N(p-4)}{p}}} \\
 (\nabla |\nabla U|^{p-4}, \nabla U) &= -c_{N,p}^{p-3} \frac{p-4}{p-1} \frac{|x|^{-\frac{2}{p-1}}}{(1 + |x|^{\frac{p}{p-1}})^{\frac{N(p-4)}{p} + \frac{N}{p}}} + c_{N,p}^{p-3} N \frac{p-4}{p-1} \frac{|x|^{\frac{p-2}{p-1}}}{(1 + |x|^{\frac{p}{p-1}})^{\frac{N(p-4)}{p} + \frac{N}{p} + 1}} \\
 (\nabla U, \nabla \phi) &= -c_{N,p}^{p-3} \frac{|x|^{\frac{2-p}{p-1}}}{(1 + |x|^{\frac{p}{p-1}})^{\frac{N}{p}}} (\nabla \phi, x) \\
 (\nabla(\nabla U, \nabla \phi), \nabla U) &= \frac{c_{N,p}^2}{(1 + |x|^{\frac{p}{p-1}})^{\frac{2N}{p}}} (\nabla \phi, x) \left[\frac{1}{p-1} |x|^{2\frac{2-p}{p-1}} - \frac{N}{p-1} \frac{|x|^{\frac{4-p}{p-1}}}{1 + |x|^{\frac{p}{p-1}}} \right] \\
 &\quad + c_{N,p}^2 \frac{|x|^{2\frac{2-p}{p-1}}}{(1 + |x|^{\frac{p}{p-1}})^{\frac{2N}{p}}} \sum_{i,j} \partial_{ij}^2 \phi x_i x_j \\
 |\nabla U|^{p-4} (\nabla U, \nabla \phi) \Delta U &= \frac{c_{N,p}^{p-2}}{(1 + |x|^{\frac{p}{p-1}})^{\frac{N(p-4)}{p} + \frac{2N}{p}}} (\nabla \phi, x) \left[\left(\frac{2-p}{p-1} + N \right) |x|^{-\frac{p}{p-1}} - \frac{N}{p-1} \frac{1}{1 + |x|^{\frac{p}{p-1}}} \right] \\
 (\nabla |\nabla U|^{p-2}, \nabla \phi) &= \frac{c_{N,p}^{p-2}}{(1 + |x|^{\frac{p}{p-1}})^{\frac{N(p-2)}{p}}} (\nabla \phi, x) \left[\frac{p-2}{p-1} |x|^{-\frac{p}{p-1}} - \frac{N(p-2)}{p-1} \frac{1}{1 + |x|^{\frac{p}{p-1}}} \right]
 \end{aligned}$$

where $c_{N,p} := \alpha_{N,p}^{\frac{N-p}{p}} \frac{N-p}{p-1}$.

Now, since U is radial we can make a partial wave decomposition of (3.1), namely we can write

$$\phi(x) = \sum_{k=0}^{\infty} \phi_k(r) Y_k(\theta), \quad \text{where } \psi_k(r) = \int_{S^{N-1}} \phi(r, \theta) Y_k(\theta) d\theta, \quad (3.2)$$

where $r = |x|$, $\theta = \frac{x}{|x|} \in S^{N-1}$ and $Y_k(\theta)$ denotes the k -th spherical harmonic satisfying ($\Delta_{S^{N-1}}$ stands for the Laplace-Beltrami operator)

$$-\Delta_{S^{N-1}} Y_k = \lambda_k Y_k. \quad (3.3)$$

It is known that this equation has a sequence of eigenvalues

$$\lambda_k = k(N + k - 2), \quad k = 0, 1, 2, \dots, \quad (3.4)$$

whose multiplicity is finite. In particular $\lambda_0 = 0$ has multiplicity 1 and $\lambda_1 = N - 1$ has multiplicity N .

Let us write the equations satisfied by the radial functions ψ_k .

It is known that (hereafter $'$ stands for $\frac{d}{dr}$)

$$\Delta(\psi_k(r) Y_k(\theta)) = Y_k(\theta) \left(\psi_k'' + \frac{N-1}{r} \psi_k' \right) + \frac{1}{r^2} \psi_k(r) \Delta_{S^{N-1}} Y_k(\theta). \quad (3.5)$$

Now, we have to compute the other terms in (3.1). It is easy to see that

$$\partial_{x_i} \phi = \psi_k'(r) \frac{x_i}{r} Y_k(\theta) + \psi_k(r) \frac{\partial Y_k}{\partial \theta_h} \frac{\partial \theta_h}{\partial x_i}$$

and

$$\begin{aligned} \partial_{x_i x_j}^2 \phi &= \psi_k''(r) \frac{x_i x_j}{r^2} Y_k(\theta) + \psi_k'(r) \left(\frac{\delta_{ij}}{r} - \frac{x_i x_j}{r^3} \right) Y_k(\theta) + \psi_k'(r) \frac{x_i}{r} \frac{\partial Y_k}{\partial \theta_h} \frac{\partial \theta_h}{\partial x_j} \\ &\quad + \psi_k'(r) \frac{x_j}{r} \frac{\partial Y_k}{\partial \theta_h} \frac{\partial \theta_h}{\partial x_i} + \psi_k(r) \frac{\partial^2 Y_k}{\partial \theta_h \partial \theta_\ell} \frac{\partial \theta_\ell}{\partial x_j} \frac{\partial \theta_h}{\partial x_i} + \psi_k(r) \frac{\partial Y_k}{\partial \theta_h} \frac{\partial^2 \theta_h}{\partial x_i \partial x_j}. \end{aligned}$$

Hence

$$(\nabla \phi, x) = \sum_{i=1}^N x_i \partial_{x_i} \phi = \psi_k'(r) r Y_k(\theta) + \psi_k(r) \frac{\partial Y_k}{\partial \theta_h} \sum_{i=1}^N \frac{\partial \theta_h}{\partial x_i} x_i = \psi_k'(r) r Y_k(\theta) \quad (3.6)$$

and

$$\begin{aligned} \sum_{i,j=1}^N \partial_{x_i x_j}^2 \phi x_i x_j &= \psi_k''(r) r^2 Y_k(\theta) + 2\psi_k'(r) r \sum_{i=1}^N \frac{\partial Y_k}{\partial \theta_h} \frac{\partial \theta_h}{\partial x_i} x_i + \psi_k(r) \sum_{i,j=1}^N \frac{\partial^2 Y_k}{\partial \theta_h \partial \theta_\ell} \frac{\partial \theta_\ell}{\partial x_j} x_j \frac{\partial \theta_h}{\partial x_i} x_i \\ &\quad + \psi_k(r) \sum_{i,j=1}^N \frac{\partial Y_k}{\partial \theta_h} \frac{\partial^2 \theta_h}{\partial x_i \partial x_j} x_i x_j = \psi_k''(r) r^2 Y_k(\theta). \end{aligned} \quad (3.7)$$

because it holds true that

$$\sum_{i=1}^N \frac{\partial \theta_h}{\partial x_i} x_i = 0 \text{ and } \sum_{i,j=1}^N \frac{\partial^2 \theta_h}{\partial x_i \partial x_j} x_i x_j = 0, \quad h = 1, \dots, N-1.$$

Putting together (3.3), (3.5), (3.6) and (3.7) into (3.1) we get the following equations for any ψ_k , $k = 0, 1, 2, \dots$,

$$\psi_k'' + \frac{\psi_k'}{r} \left(\frac{N-1}{p-1} + \frac{(p-2)N}{p-1} \frac{1}{1+r^{\frac{p}{p-1}}} \right) - \frac{\lambda_k}{r^2} \psi_k + \gamma_{N,p} \frac{r^{\frac{p}{p-1}-2}}{(1+r^{\frac{p}{p-1}})^2} \psi_k = 0, \quad (3.8)$$

which can be rewritten in a weak form as

$$\mathcal{L}_k(\psi_k) = 0, \quad k = 0, 1, 2, \dots, \quad (3.9)$$

where the operator \mathcal{L}_k is defined by

$$\mathcal{L}_k(\psi) := (r^{N-1} |U'(r)|^{p-2} \psi')' + (p^* - 1) r^{N-1} (U(r))^{p^*-2} \psi - \lambda_k r^{N-3} |U'(r)|^{p-2} \psi.$$

Since we are concerned with solutions $\psi \in \mathcal{D}_*^{1,2}(\mathbb{R}^N)$ to the linear equation (1.7), we will look for solutions ψ_k to (3.8) or (3.9) in the space \mathcal{D}_k which is the completion of $C_c^1([0, +\infty))$ with respect to the norm

$$\|\psi\|_k := \left(\int_0^{+\infty} r^{N-1} |U'(r)|^{p-2} |\psi'(r)|^2 dr + \lambda_k \int_0^{+\infty} r^{N-3} |U'(r)|^{p-2} |\psi(r)|^2 dr \right)^{\frac{1}{2}}.$$

3.2. Solving the equations $\mathcal{L}_k(\psi) = 0$.

- *The case $k = 0$.*

We know that the function Z_0 defined in (1.5) as

$$Z_0(x) = \frac{N-p}{p(p-1)} \alpha_{N,p}^{\frac{N-p}{p}} \underbrace{\frac{p-1-|x|^{\frac{p}{p-1}}}{\left(1+|x|^{\frac{p}{p-1}}\right)^{\frac{N}{p}}}}_{:=\psi_0(|x|)}$$

solves the equation (3.1). We claim that all the solutions in \mathcal{D}_0 to $\mathcal{L}_0(\psi) = 0$ are given by $\psi = c\psi_0$, $c \in \mathbb{R}$. Indeed, for $k = 0$ we have that $\lambda_0 = 0$ and a straightforward

computation shows that $\psi_0 \in \mathcal{D}_0$ and $\mathcal{L}_0(\psi_0) = 0$.

We look for a second linearly independent solution of the form

$$w(r) = c(r)\psi_0(r).$$

Then we get

$$c''(r)\psi_0(r) + c'(r) \left[2\psi_0'(r) + \frac{\psi_0(r)}{r} \left(\frac{N-1}{p-1} + \frac{N(p-2)}{p-1} \frac{1}{1+r^{\frac{p}{p-1}}} \right) \right] = 0$$

and hence

$$\frac{c''(r)}{c'(r)} = -2 \frac{\psi_0'(r)}{\psi_0(r)} - \frac{1}{r} \left(\frac{N-1}{p-1} + \frac{N(p-2)}{p-1} \frac{1}{1+r^{\frac{p}{p-1}}} \right).$$

A direct computation shows that

$$c'(r) = A \frac{(1+r^{\frac{p}{p-1}})^{\frac{N(p-2)}{p}}}{(\psi_0(r))^2 r^{\frac{N-1+N(p-2)}{p-1}}} \text{ for some } A \in \mathbb{R} \setminus \{0\}.$$

Therefore

$$c(r) \sim Br^{\frac{N-p}{p-1}} \text{ and } w(r) = c(r)\psi_0(r) \sim B \text{ as } r \rightarrow +\infty \text{ with } B \neq 0.$$

However $w \notin \mathcal{D}_0$ because of Lemma 2.1.

- *The case $k = 1$.*

We know that the function Z_i defined in (1.6) as

$$Z_i(x) = \frac{N-p}{p-1} \alpha_{N,p}^{\frac{N-p}{p}} \frac{x_i}{|x|} \underbrace{\frac{|x|^{\frac{1}{p-1}}}{\left(1+|x|^{\frac{p}{p-1}}\right)^{\frac{N}{p}}}}_{:=\psi_1(|x|)}, \quad i = 1, \dots, N \quad (3.10)$$

solve the equation (3.1). We claim that all the solutions in \mathcal{D}_1 to $\mathcal{L}_1(\psi) = 0$ are given by $\psi = c\psi_1$, $c \in \mathbb{R}$. Indeed, for $k = 1$ we have that $\lambda_1 = N-1$ and a straightforward computation shows that $\psi_1 \in \mathcal{D}_1$ and $\mathcal{L}_1(\psi_1) = 0$.

As above, we look for a second linearly independent solution of the form

$$w(r) = c(r)\psi_1(r).$$

Then we get

$$c''(r)\psi_1(r) + c'(r) \left[2\psi_1'(r) + \frac{\psi_1(r)}{r} \left(\frac{N-1}{p-1} + \frac{N(p-2)}{p-1} \frac{1}{1+r^{\frac{p}{p-1}}} \right) \right] = 0$$

and a direct computation shows that

$$c'(r) = A \frac{(1+r^{\frac{p}{p-1}})^{\frac{N(p-2)}{p}}}{(\psi_1(r))^2 r^{\frac{N-1+N(p-2)}{p-1}}} \text{ for some } A \in \mathbb{R} \setminus \{0\}.$$

Therefore

$$c(r) \sim Br^{\frac{N-1}{p-1}+1} \text{ and } w(r) = c(r)\psi_1(r) \sim Br \text{ as } r \rightarrow +\infty \text{ with } B \neq 0.$$

However $w \notin \mathcal{D}_1$ because of Lemma 2.1.

- *The case $k \geq 2$.*

We claim that all the solutions in \mathcal{D}_k of $\mathcal{L}_k(\psi) = 0$ are identically zero if $k \geq 2$.

Assume there exists a function ψ_k such that $\mathcal{L}_k(\psi_k) = 0$, i.e. for any $r \geq 0$

$$(r^{N-1}|U'(r)|^{p-2}\psi_k')' + r^{N-1}(U(r))^{\frac{Np-2N+2p}{N-p}}\psi_k - \lambda_k r^{N-3}|U'(r)|^{p-2}\psi_k = 0. \quad (3.11)$$

We claim that $\psi_k \equiv 0$ if $k \geq 2$. We argue by contradiction. Without loss of generality, we can suppose that there exists $r_k > 0$ (possibly $+\infty$) such that $\psi_k(r) > 0$ for any $r \in (0, r_k)$ and $\psi_k(r_k) = 0$. In particular, $\psi'_k(r_k) \leq 0$.

Now, let $\psi_1(r) = U'(r)$ (see (3.10)) be the solution of $\mathcal{L}_1(\psi_1) = 0$, i.e. for any $r \geq 0$

$$(r^{N-1}|U'(r)|^{p-2}\psi'_1)' + r^{N-1}(U(r))^{\frac{Np-2N+2p}{N-p}}\psi_1 - \lambda_1 r^{N-3}|U'(r)|^{p-2}\psi_1 = 0. \quad (3.12)$$

We multiply (3.11) by ψ_1 , (3.12) by ψ_k , we integrate between 0 and r_k , we subtract the two expressions and we get

$$\begin{aligned} & (\lambda_k - \lambda_1) \int_0^{r_k} r^{N-3}|U'(r)|^{p-2}\psi_k\psi_1 dr \\ &= \int_0^{r_k} (r^{N-1}|U'(r)|^{p-2}\psi'_k)' \psi_1 dr - \int_0^{r_k} (r^{N-1}|U'(r)|^{p-2}\psi'_1)' \psi_k dr \end{aligned} \quad (3.13)$$

(we integrate by part and we use that $\psi_k(r_k) = 0$)

$$= r_k^{N-1}|U'(r_k)|^{p-2}\psi'_k(r_k)\psi_1(r_k)$$

and a contradiction arises when $\lambda_k > \lambda_1$, (that is $k \geq 2$), since $\psi'_k(r_k) \leq 0$, $\psi_1(r) < 0$

for any $r > 0$ and $\int_0^{r_k} r^{N-3}|U'(r)|^{p-2}\psi_k\psi_1 dr < 0$.

REFERENCES

- [1] T. Aubin, *Problemes isoperimtriques et espaces de Sobolev*, J. Differ. Geometry **11** (1976), 573–598.
- [2] H. Brzis and L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983), 437–77.
- [3] L. Caffarelli, B. Gidas and J. Spruck, *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical sobolev growth*. Comm. Pur. Appl. Math. **42** (1989) 271–297.
- [4] Caffarelli, L.; Kohn, R.; Nirenberg, L. First order interpolation inequalities with weights. Compositio Math. **53** (1984), no. 3, 259–275.
- [5] Clapp, M.; Lopez Rios, L. *Entire nodal solutions to the pure critical exponent problem for the p -Laplacian*. J. Differential Equations **265** (2018), no. 3, 891–905
- [6] Coron, J.-M. *Topologie et cas limite des injections de Sobolev*. C. R. Acad. Sci. Paris Sr. I Math. **299** (1984) 209–212
- [7] L. Damascelli and M. Ramaswamy, *Symmetry of C^1 - solutions of p -Laplace equations in \mathbb{R}^N* , Adv. Nonlinear Stud. **1** (2001), no. 1, 40–64.
- [8] L. Damascelli, S. Merchán, L. Montoro and B. Sciunzi, *Radial symmetry and applications for a problem involving the $-\Delta_p(\cdot)$ operator and critical nonlinearity in \mathbb{R}^N* , Adv. Math. **256** (2014), 313–335.
- [9] L. Damascelli, and B. Sciunzi, *Regularity, monotonicity and symmetry of positive solutions of m -Laplace equations*. J. Differential Equations **206** (2004), no. 2, 483–515.
- [10] del Pino, M. Musso, M.; Pacard, F.; Pistoia, A. *Torus action on S_n and sign-changing solutions for conformally invariant equations*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **12** (2013), no. 1, 209–237.
- [11] del Pino, M. Musso, M.; Pacard, F.; Pistoia, A. *Large energy entire solutions for the Yamabe equation*. J. Differential Equations **251** (2011), no. 9, 2568–2597.
- [12] Ge, Y.; Musso, M.; Pistoia, A.; Pollack, D. *A refined result on sign changing solutions for a critical elliptic problem*. Commun. Pure Appl. Anal. **12** (2013), no. 1, 125–155
- [13] Ge, Y.; Musso, M.; Pistoia, A. *Sign changing tower of bubbles for an elliptic problem at the critical exponent in pierced non-symmetric domains*. Comm. Partial Differential Equations **35** (2010), no. 8, 1419–1457.
- [14] B. Gidas, W.-M. Ni and L. Nirenberg, *Symmetry and related properties via the maximum principle*. Comm. Math. Phys. **68** (1979), 209–243.
- [15] M. Guedda, L. Veron, *Local and global properties of solutions of quasilinear elliptic equations*, J. Differential Equations **76** (1) (1988) 159–189
- [16] M. Guedda, L. Veron, *Quasilinear elliptic equations involving critical Sobolev exponents*, Nonlinear Anal. **13** (1989) 879–902.
- [17] Mercuri, C.; Pacella, F. *On the pure critical exponent problem for the p -Laplacian*. Calc. Var. Partial Differential Equations **49** (2014), no. 3-4, 1075–1090.

- [18] Mercuri, C.; Sciunzi, B.; Squassina, M. *On Coron's problem for the p -Laplacian*. J. Math. Anal. Appl. **421** (2015), no. 1, 362–369.
- [19] Nirenberg, L. *On elliptic partial differential equations*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) **13** (1959), 115–162.
- [20] Musso, M., Pistoia A. *Sign changing solutions to a nonlinear elliptic problem involving the critical Sobolev exponent in pierced domains*. J. Math. Pures Appl. **86** (2006) 510–528
- [21] Perera, K.; Squassina, M.; Yang, Y. *Bifurcation and multiplicity results for critical p -Laplacian problems*. Topol. Methods Nonlinear Anal. **47** (2016), no. 1, 187–194.
- [22] O. Rey, *The role of the Green's function in a nonlinear elliptic equation involving the critical Sobolev exponent*, J. Funct. Anal. **89** (1990), 1–52
- [23] B. Sciunzi, *Classification of positive $\mathcal{D}^{1,p}(\mathbb{R}^N)$ -solutions to the critical p -Laplace equation in \mathbb{R}^N* , Advances in Mathematics, **291** (2016), 12–23.
- [24] G. Talenti, *Best constant in Sobolev inequality*, Ann. Mat. Pura Appl. **110** (1976), 353–372.
- [25] J. Vétois, *A priori estimates and application to the symmetry of solutions for critical p -Laplace equations*, J. Differential Equations. **260** (2016), 149–161.

ANGELA PISTOIA, DIPARTIMENTO DI SCIENZE DI BASE E APPLICATE PER L'INGEGNERIA, SAPIENZA UNIVERSITÀ DI ROMA, VIA SCARPA 16, 00161 ROMA, ITALY
E-mail address: angela.pistoia@uniroma1.it

GIUSI VAIRA, DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DEGLI STUDI DELLA CAMPANIA "LUIGI VANVITELLI", VIALE LINCOLN 5, 81100 CASERTA, ITALY
E-mail address: giusi.vaira@unicampania.it