

Looking for the hidden-charm pentaquark resonances in $J/\psi p$ scattering

Hongxia Huang,^{*} Jun He,[†] and Jialun Ping[‡]

Department of Physics, Nanjing Normal University, Nanjing, Jiangsu 210023, China

In the framework of quark delocalization color screening model, the three new reported pentaquarks $P_c(4312)$, $P_c(4440)$, and $P_c(4457)$ can be identified as the hidden-charm molecular states $\Sigma_c D$ with $J^P = \frac{1}{2}^-$, $\Sigma_c D^*$ with $J^P = \frac{3}{2}^-$, and $\Sigma_c D^*$ with $J^P = \frac{1}{2}^-$, in the baryon-meson scattering process, respectively. Besides, the $\Sigma_c^* D^*$ of both $J^P = \frac{1}{2}^-$ and $J^P = \frac{3}{2}^-$ are also possible molecular pentaquarks. Moreover, the calculation is extended to the P_c -like molecular pentaquarks P_b . Several states with masses above 11 GeV and narrow width are obtained. All these heavy pentaquarks are worth searching in the future experiments.

PACS numbers: 13.75.Cs, 12.39.Pn, 12.39.Jh

I. INTRODUCTION

In 2015, the claim of two hidden-charm pentaquark states $P_c(4380)$ and $P_c(4450)$ by the LHCb Collaboration [13] attracted people's interesting in the pentaquarks with heavy quarks and inspired a lot of theoretical work on these two states, such as the baryon-meson molecules [14–24], the diquark-triquark pentaquarks [25, 26], the diquark-diquark-antiquark pentaquarks [27–30], the genuine multiquark states [31], the topological soliton [32], and the kinematical threshold effects in the triangle singularity mechanism [33–35], and so on. The lattice QCD simulation of NJ/ψ and $N\eta_c$ scattering is also performed to find these P_c states [36].

Four years later, at the Rencontres de Moriond QCD Conference, the LHCb Collaboration reported the observation of three new pentaquarks, named as $P_c(4312)$, $P_c(4440)$, and $P_c(4457)$ [37]. The $P_c(4312)$ was discovered with 7.3σ significance by analyzing the $J/\psi p$ invariant mass spectrum. The previously reported $P_c(4440)$ structure was resolved at 5.4σ significance into two narrow states: the $P_c(4440)$ and $P_c(4457)$. The masses and widths of these states are:

$$\begin{aligned} P_c(4312) : M &= 4311.9 \pm 0.7_{-0.6}^{+6.8} \text{ MeV}, \\ \Gamma &= 9.8 \pm 2.7_{-4.5}^{+3.7} \text{ MeV}, \\ P_c(4440) : M &= 4440.3 \pm 1.3_{-4.7}^{+4.1} \text{ MeV}, \\ \Gamma &= 20.6 \pm 4.9_{-10.1}^{+8.7} \text{ MeV}, \\ P_c(4457) : M &= 4457.3 \pm 0.6_{-1.7}^{+4.1} \text{ MeV}, \\ \Gamma &= 6.4 \pm 2.0_{-1.9}^{+5.7} \text{ MeV}. \end{aligned} \quad (1)$$

As mentioned in Ref. [37], since all three states are narrow and below the $\Sigma_c^+ \bar{D}^0$ and $\Sigma_c^+ \bar{D}^{*0}$ thresholds within plausible hadron-hadron binding energies, they provide the strongest experimental evidence to date for the existence of molecular states composed of a charmed baryon

and an anticharmed meson. Immediately after the report of the LHCb Collaboration, several theoretical work have been done to study the mass spectrum of these states [38–40]. Ref. [41] studied the isospin breaking decays of the molecular structure of the $P_c(4457)$.

Searching for the existence of multiquark states is an important issue of the hadron physics. To provide the necessary information for experiments, mass spectrum calculation alone is not enough. The study of hadron-hadron scattering, as well as the main production process of multiquark states, is indispensable. In the framework of the quark delocalization color screening model (QDCSM), the detail of which can be found in Refs. [42, 43], we apply the well developed resonating group method (RGM) [44] to calculate the baryon-meson scattering phase shifts and to find the hidden-charm and hidden-bottom pentaquark resonances. The wave function of the baryon-meson system is of the form

$$\Psi = \mathcal{A} \left[\hat{\phi}_A(\xi_1, \xi_2) \hat{\phi}_B(\xi_3) \chi_L(\mathbf{R}_{AB}) \right]. \quad (2)$$

where ξ_1 and ξ_2 are the internal coordinates for the baryon cluster A, and ξ_3 is the internal coordinate for the meson cluster B. $\mathbf{R}_{AB} = \mathbf{R}_A - \mathbf{R}_B$ is the relative coordinate between the two clusters. The $\hat{\phi}_A$ and $\hat{\phi}_B$ are the internal cluster wave functions of the baryon A (antisymmetrized) and meson B, and $\chi_L(\mathbf{R}_{AB})$ is the relative motion wave function between two clusters. The symbol \mathcal{A} is the anti-symmetrization operator defined as

$$\mathcal{A} = 1 - P_{14} - P_{24} - P_{34}, \quad (3)$$

where 1, 2, and 3 stand for the quarks in the baryon cluster and 4 stands for the quark in the meson cluster. For a bound-state problem, $\chi_L(\mathbf{R}_{AB})$ is expanded by gaussian bases

$$\begin{aligned} \chi_L(\mathbf{R}_{AB}) &= \frac{1}{\sqrt{4\pi}} \left(\frac{6}{5\pi b^2} \right)^{3/4} \sum_{i=1}^n C_i \\ &\times \int \exp \left[-\frac{3}{5b^2} (\mathbf{R}_{AB} - \mathbf{S}_i)^2 \right] Y_{LM}(\hat{\mathbf{S}}_i) d\hat{\mathbf{S}}_i \\ &= \sum_{i=1}^n C_i \frac{u_L(R_{AB}, S_i)}{R_{AB}} Y_{LM}(\hat{\mathbf{R}}_{AB}). \end{aligned} \quad (4)$$

^{*}Electronic address: hxhuang@njnu.edu.cn

[†]Electronic address: junhe@njnu.edu.cn

[‡]Electronic address: jlping@njnu.edu.cn

with

$$u_L(R_{AB}, S_i) = \sqrt{4\pi} \left(\frac{6}{5\pi b^2} \right)^{3/4} R_{AB} \times \exp \left[-\frac{3}{5b^2} (R_{AB}^2 - S_i^2) \right] i^L j_L \left(-i \frac{6}{5b^2} R_{AB} S_i \right). \quad (5)$$

where S_i is called the generating coordinate, C_i is expansion coefficients, n is the number of the gaussian bases, which is determined by the stability of the results, and j_L is the L -th spherical Bessel function.

For a scattering problem, the relative wave function is expanded as

$$\chi_L(\mathbf{R}_{AB}) = \sum_{i=1}^n C_i \frac{\tilde{u}_L(R_{AB}, S_i)}{R_{AB}} Y_{LM}(\hat{\mathbf{R}}_{AB}). \quad (6)$$

with

$$\tilde{u}_L(R_{AB}, S_i) = \begin{cases} \alpha_i u_L(R_{AB}, S_i), & R_{AB} \leq R_C \\ [h_L^-(k_{AB}, R_{AB}) - s_i h_L^+(k_{AB}, R_{AB})] & R_{AB}, R_{AB} \geq R_C \end{cases} \quad (7)$$

where h_L^\pm is the L -th spherical Hankel functions, k_{AB} is the momentum of relative motion with $k_{AB} = \sqrt{2\mu_{AB}E_{cm}}$, μ_{AB} is the reduced mass of two hadrons (A and B) of the open channel; E_{cm} is the incident energy, and R_C is a cutoff radius beyond which all the strong interaction can be disregarded. Besides, α_i and s_i are complex parameters which are determined by the smoothness condition at $R_{AB} = R_C$ and C_i satisfy $\sum_{i=1}^n C_i = 1$. After performing variational procedure, a L -th partial-wave equation for the scattering problem can be deduced as

$$\sum_{j=1}^n \mathcal{L}_{ij}^L C_j = \mathcal{M}_i^L \quad (i = 0, 1, \dots, n-1), \quad (8)$$

with

$$\mathcal{L}_{ij}^L = \mathcal{K}_{ij}^L - \mathcal{K}_{i0}^L - \mathcal{K}_{0j}^L + \mathcal{K}_{00}^L, \quad (9)$$

$$\mathcal{M}_i^L = \mathcal{K}_{00}^L - \mathcal{K}_{i0}^L, \quad (10)$$

and

$$\mathcal{K}_{ij}^L = \left\langle \hat{\phi}_A(\xi'_1, \xi'_2) \hat{\phi}_B(\xi'_3) \frac{\tilde{u}_L(R'_{AB}, S_i)}{R'_{AB}} Y_{LM}(\hat{\mathbf{R}}'_{AB}) \right| H - E \left| \hat{\phi}_A(\xi_1, \xi_2) \hat{\phi}_B(\xi_3) \frac{\tilde{u}_L(R_{AB}, S_j)}{R_{AB}} Y_{LM}(\hat{\mathbf{R}}_{AB}) \right\rangle. \quad (11)$$

By solving Eq.(8), we can obtain the expansion coefficients C_i . Then the S matrix element S_L and the phase shifts δ_L are given by

$$S_L \equiv e^{2i\delta_L} = \sum_{i=1}^n C_i s_i, \quad (12)$$

Resonances are unstable particles usually observed as bell-shaped structures in scattering cross sections of their corresponding open channels. For a simple narrow resonance, the peak position and the of the half-width of the bell shape are the mass M and the decay width Γ of the resonance. The cross-section σ_L and the scattering phase shifts δ_L have relations:

$$\sigma_L = \frac{4\pi}{k^2} (2L+1) \sin^2 \delta_L, \quad (13)$$

where $k = \sqrt{2\mu E_{cm}}/\hbar$; μ is the reduced mass of two hadrons of the open channel; E_{cm} is the incident energy. Therefore, by using the scattering phase shifts, the cross sections can be easily obtained. The scattering phase shifts are shown in our previous work [45]. Here, to compare with the experimental data, we calculate the cross sections of the $J/\psi p$ scattering channel, which are shown in Fig. 1. Based on our previous theoretical work, the new experimental information and the calculated cross sections, we discuss possible explanations of the three new pentaquarks: $P_c(4312)$, $P_c(4440)$, and $P_c(4457)$.

For $P_c(4312)$, our previous study showed that there was a narrow resonance state $\Sigma_c D$ with $J^P = \frac{1}{2}^-$ in the scattering channels of $\eta_c p$, $J/\psi p$, $\Lambda_c D$ and $\Lambda_c D^*$ [45]. The calculated mass of this resonance state is 4306.7 \sim 4311.3 MeV, and the decay width is 7.1 MeV. From the Fig. 1(a), we also see a sharp peak appearing at the mass of 4307.9 MeV with a very narrow partial width of about 1.2 MeV. It is obvious that both the mass and decay width are close to the experimental value of the $P_c(4312)$, which indicates that the newly reported $P_c(4312)$ state can be identified as the $\Sigma_c D$ molecular pentaquark with $J^P = \frac{1}{2}^-$ in our model calculation.

For $P_c(4440)$ and $P_c(4457)$, we assigned them as the resonances $\Sigma_c D^*$ with $J^P = \frac{1}{2}^-$ and $\Sigma_c D^*$ with $J^P = \frac{3}{2}^-$ according to our calculation [45]. These two resonances also appear as two sharp peaks in the cross section of the $J/\psi p$ channel (see Fig. 1 (a) and (b)). The masses and the partial decay widths can be read from Fig. 1, they are: $\Sigma_c D^*$ of $J^P = \frac{1}{2}^-$, 4459.7 MeV and 3.9 MeV; $\Sigma_c D^*$ of $J^P = \frac{3}{2}^-$, 4445.7 MeV and 1.5 MeV. Compared with experimental data, the $P_c(4440)$ is more possible to be the molecular pentaquark $\Sigma_c D^*$ of $J^P = \frac{3}{2}^-$, and the $P_c(4457)$ can be explained as the molecular pentaquark $\Sigma_c D^*$ of $J^P = \frac{1}{2}^-$.

Besides the three resonances discussed above, one may see the fourth peak showed in Fig. 1 (b). It is the molecular pentaquark $\Sigma_c^* D$ with $J^P = \frac{3}{2}^-$. The mass of this resonance is 4376.4 MeV, which is very close to the reported $P_c(4380)$. However, the decay width is only 1.5 MeV, much smaller than the experimental value. We propose the experiment to find whether there is any narrow resonance near the $P_c(4380)$.

In addition, in Fig. 1(a), we also find there is a cusp near the mass of 4527 MeV, which is the threshold of the $\Sigma_c^* D^*$. Our previous calculation showed that the single




FIG. 1: The cross section of the $J/\psi p$ channel with $J^P = \frac{1}{2}^-$ and $J^P = \frac{3}{2}^-$ respectively.

channel $\Sigma_c^* D^*$ of $J^P = \frac{1}{2}^-$ was bound, but it disappeared by coupling with the scattering channel $J/\psi p$, $\Sigma_c D$, and $\Sigma_c D^*$ channels [45]. So it only appears as a cusp in the cross section of the $J/\psi p$. This is consistent with the experimental result, there is no any distinct signal near the threshold of the $\Sigma_c^* D^*$ in the experimental data. However, this molecular pentaquark $\Sigma_c^* D^*$ of $J^P = \frac{1}{2}^-$ appeared as a resonance state in the $\eta_c p$ scattering channel [45]. The mass and width of it is 4525.8 MeV and 4.0 MeV, respectively. Besides, the $\Sigma_c^* D^*$ state of $J^P = \frac{3}{2}^-$ has the similar situation. It appears as a cusp in the cross section of the $J/\psi p$ (see Fig. 1(b)), but it turned to be a resonance state in the $\Lambda_c D^*$ scattering process, with a mass of 4523.0 MeV and a width of 1.0 MeV [45]. Although it is difficult for the experiment to perform the $\eta_c p$ or the $\Lambda_c D^*$ scattering, the $\Sigma_c^* D^*$ of both $J^P = \frac{1}{2}^-$ and $J^P = \frac{3}{2}^-$ are possible molecular pentaquarks, which are worth looking for.

Because of the heavy flavor symmetry, we also extend the study to the hidden-bottom pentaquarks. The results are similar to the hidden-charm molecular pentaquarks. Fig. 2 shows the cross sections of the Υp channel with $J^P = \frac{1}{2}^-$ and $J^P = \frac{3}{2}^-$ respectively. From Fig. 2(a), we can see three sharp peaks in the cross sections, which correspond to three resonance states: $\Sigma_b B$, $\Sigma_b B^*$ and $\Sigma_b^* B^*$ states with $IJ^P = \frac{1}{2} \frac{1}{2}^-$. The resonance mass of $\Sigma_b B$ state is 11077.5 MeV and the decay width is about 0.1 MeV; $\Sigma_b B^*$ has the mass of 11125.8 MeV and the decay width of 0.8 MeV; and $\Sigma_b^* B^*$ has the mass of 11153.5 MeV and the decay width of 3.0 MeV.




FIG. 2: The cross section of the Υp channel with $J^P = \frac{1}{2}^-$ and $J^P = \frac{3}{2}^-$ respectively.

For the hidden-bottom pentaquarks with $IJ^P = \frac{1}{2} \frac{3}{2}^-$, two pinnacles appear in the Fig. 2(b), corresponding to two resonance states: $\Sigma_b B^*$ and $\Sigma_b^* B$. The mass and the decay width of $\Sigma_b B^*$ is 11122.7 MeV and 0.2 MeV respectively. The mass and the decay width of $\Sigma_b^* B$ is 11103.6 MeV and 0.8 MeV, respectively. All these hidden-bottom pentaquarks have the similar properties with the hidden-charm pentaquarks, so we can call them P_c -like molecular pentaquarks P_b , which are also worth searching for by experiments.

To summarize, in the framework of QDCSM, we look for the hidden-charm and hidden-bottom pentaquark resonances by studying $J/\psi p$ and Υp scattering process. The three new narrow pentaquarks $P_c(4312)$, $P_c(4440)$, and $P_c(4457)$ observed in the process of $\Lambda_b \rightarrow J/\psi p K$ reported by LHCb can be interpreted as the hidden-charm molecular pentaquarks $\Sigma_c D$ with $J^P = \frac{1}{2}^-$, $\Sigma_c D^*$ with $J^P = \frac{3}{2}^-$, and $\Sigma_c D^*$ with $J^P = \frac{1}{2}^-$, respectively. Another molecular pentaquark $\Sigma_c^* D$ with $J^P = \frac{3}{2}^-$ is also existed in our calculation, the mass of which is close to the $P_c(4380)$, but the width is much smaller than it. Besides, the $\Sigma_c^* D^*$ of both $J^P = \frac{1}{2}^-$ and $J^P = \frac{3}{2}^-$ are possible molecular pentaquarks. All these narrow pentaquarks are worth searching for or being confirmed in future experiments. The Jefferson Lab has proposed to look for the hidden-charm pentaquarks by using photo-production of J/ψ at threshold in Hall C [46]. Moreover, the pentaquarks with charm quarks can also be observed by the PANDA/FAIR [47]. For the pentaquarks with

the hidden-bottom, we predict several P_c -like molecular pentaquarks P_b above 11 GeV with narrow width. We hope the proposed electron-ion collider (EIC) [48] and the upgraded facilities at Jefferson Lab [49] can play important role in discovering these interesting super-heavy pentaquarks.

Searching for multi-quark states is an important topic in hadron physics. To provide more information for experiments, the baryon-meson scattering process calculation is expected. Doing baryon-meson scattering is also a challenge for quark model. The model is needed to describe the baryon and meson spectra at the same time. With the accumulation of more experimental data on baryon-meson scattering and the pentaquark, the quark

model will be further updated. To use the simple picture to describe the natural phenomena is one of the goal of physics.

Acknowledgment

This work is supported partly by the National Science Foundation of China under Contract Nos. 11675080, 11775118 and 11535005, the Natural Science Foundation of the Jiangsu Higher Education Institutions of China (Grant No. 16KJB140006).

-
- [1] T. Nakano, *et al.* (LEPS Collaboration), Phys. Rev. Lett. **91**, 012002 (2003).
 - [2] T. Nakano, *et al.* (LEPS Collaboration), Phys. Rev. C **79**, 025210 (2009).
 - [3] Megumi Naruki for the J-PARC E19 Collaboration, Kyoto University, arXiv:1602.03951.
 - [4] J. J. Wu, R. Molina, E. Oset, and B. S. Zou, Phys. Rev. Lett. **105**, 232001 (2010); Phys. Rev. C **84**, 015202 (2011).
 - [5] Z. C. Yang, Z. F. Sun, J. He, X. Liu and S. L. Zhu, Chin. Phys. C **36**, 6 (2012).
 - [6] C. W. Xiao, J. Nieves and E. Oset, Phys. Rev. D **88**, 056012 (2013).
 - [7] T. Uchino, W. H. Liang and E. Oset, Eur. Phys. J. A **52**, 43 (2016).
 - [8] M. Karliner and J. L. Rosner, Phys. Rev. Lett. **115**, 122001 (2015).
 - [9] E. J. Garzon and J. J. Xie, Phys. Rev. C **92**, 035201 (2015).
 - [10] W. L. Wang, F. Huang, Z. Y. Zhang and B. S. Zou, Phys. Rev. C **84**, 015203 (2011).
 - [11] S. G. Yuan, K. W. We, J. He, H. S. Xu and B. S. Zou, Eur. Phys. J. A **48**, 61 (2012).
 - [12] Y. Huang, J. He, H. F. Zhang and X. R. Chen, J. Phys. G **41**, 115004 (2014).
 - [13] R. Aaij, *et al.* (LHCb Collaboration), Phys. Rev. Lett. **115**, 072001 (2015).
 - [14] R. Chen, X. Liu, X. Q. Li and S. L. Zhu, Phys. Rev. Lett. **115**, 132002 (2015).
 - [15] H. X. Chen, W. Chen, X. Liu, T. G. Steele and S. L. Zhu, Phys. Rev. Lett. **115**, 172001 (2015).
 - [16] L. Roca, J. Nieves and E. Oset, Phys. Rev. D **92**, 094003 (2015).
 - [17] J. He, Phys. Lett. B **753**, 547 (2016).
 - [18] H. X. Huang, C. R. Deng, J. L. Ping and F. Wang, Eur. Phys. J. C **76**, 624 (2016).
 - [19] G. Yang, J. L. Ping and F. Wang, Phys. Rev. D **95**, 014010 (2017).
 - [20] U. G. Meissner and J. A. Oller, Phys. Lett. B **751**, 59 (2015).
 - [21] C. W. Xiao and U. G. Meissner, Phys. Rev. D **92**, 114002 (2015).
 - [22] R. Chen, X. Liu and S. L. Zhu, Nucl. Phys. A **954**, 406 (2016).
 - [23] H. X. Chen, E. L. Cui, W. Chen, T. G. Steele, X. Liu and S. L. Zhu, Eur. Phys. J. C **76**, 572 (2016).
 - [24] K. Azizi, Y. Sarac and H. Sundu, Phys. Rev. D **95**, 094016 (2017).
 - [25] R. F. Lebed, Phys. Lett. B **749**, 454 (2015).
 - [26] R. Zhu and C. F. Qiao, Phys. Lett. B **756**, 259 (2016).
 - [27] L. Maiani, A. D. Polosa and V. Riquer, Phys. Lett. B **749**, 289 (2015).
 - [28] V. V. Anisovich, M. A. Matveev, J. Nyiri, A. V. Sarantsev and A. N. Semenova, arXiv:1507.07652.
 - [29] R. Ghosh, A. Bhattacharya and B. Chakrabarti, Phys. Part. Nucl. Lett. **14**, 550 (2017).
 - [30] Z. G. Wang, Eur. Phys. J. C **76**, 70 (2016).
 - [31] A. Mironov and A. Morozov, JETP Lett. **102**, 271 (2015).
 - [32] N. N. Scoccola, D. O. Riska and M. Rho, Phys. Rev. D **92**, 051501 (2015).
 - [33] F. K. Guo, Ulf-G. Meissner, W. Wang and Z. Yang, Phys. Rev. D **92**, 071502 (2015).
 - [34] X. H. Liu, Q. Wang and Q. Zhao, Phys. Lett. B **757**, 231 (2016).
 - [35] M. Mikhasenko, arXiv:1507.06552.
 - [36] U. Skerbis and S. Prelovsek, arXiv:1811.02285.
 - [37] T. Skwarnicki, Hadron spectroscopy and exotic states at LHCb, <http://moriond.in2p3.fr/QCD/2019/TuesdayMorning/Skwarnicki.pptx>, talk given on behalf of the LHCb Collaboration at the QCD and High Energy Interactions at the 54th Rencontres de Moriond, 23-30 March 2019.
 - [38] H. X. Chen, W. Chen and S. L. Zhu, arXiv:1903.11001.
 - [39] R. Chen, X. Liu, Z. F. Sun and S. L. Zhu, arXiv:1903.11013.
 - [40] M. Z. Liu, Y. W. Pan, F. Z. Peng, M. S. Sanchez, L. S. Geng, A. Hosaka and P. Valderrama, arXiv:1903.11560.
 - [41] F. K. Guo, H. J. Jing, Ulf-G. Meissner and S. Sakai, arXiv:1903.11503.
 - [42] F. Wang, G. H. Wu, L. J. Teng and T. Goldman, Phys. Rev. Lett. **69**, 2901 (1992); G. H. Wu, L. J. Teng, J. L. Ping, F. Wang and T. Goldman, Phys. Rev. C **53**, 1161 (1996).
 - [43] H. X. Huang, P. Xu, J. L. Ping and F. Wang, Phys. Rev. C **84**, 064001 (2011).
 - [44] M. Kamimura, Supp. Prog. Theo. Phys. **62**, 236 (1977).
 - [45] H. X. Huang and J. L. Ping, Phys. Rev. D **99**, 014010 (2019).

- (2019).
- [46] Z.-E. Meziani, *et al.* (JLab Collaboration), arXiv:1609.00676v2.
- [47] M. F. M. Lutz *et al.* (PANDA Collaboration), arXiv:0903.3905.
- [48] V. Ptitsyn, AIP Conf. Proc. **735**, 1149 (2009).
- [49] S. Joosten and Z.-E. Meziani, arXiv:1802.02616v2.