2d duality for orthogonal gauge theories with 8 supersymmetries

Eran Avraham a 1 and Oren Bergman a 2

^a Department of Physics, Technion, Israel Institute of Technology Haifa, 32000, Israel

ABSTRACT

We recently conjectured a set of dualities relating two-dimensional orthogonal gauge theories with $\mathcal{N}=(4,4)$ supersymmetry, analogous to Hori's dualities with $\mathcal{N}=(2,2)$ supersymmetry. Here we provide a quantitative test of this conjecture by computing the elliptic genera of the dual pairs and showing that they agree. The elliptic genus of orthogonal gauge theories has multiple topological sectors that depend on the global structure of the group and on the value of a discrete θ parameter. We derive the dependence on the θ parameter by determining whether a given sector has (S)Pin structure or not.

¹eranav@post.bgu.ac.il

²bergman@physics.technion.ac.il

Contents

1	Introduction	2
2	Flat $O(k)$ connections on T^2 2.1 Pin structure	4
3	$\mathcal{N} = (4,4) \text{ theories}$ $3.1 O(1) + n$ $3.2 O(2) + n$ $3.3 O(3) + n$ $3.4 \mathcal{N} = (4,4) \text{ duality}$ 3.5Twisting	8 9 9
4	Conclusions	12
5	Acknowledgements	13
\mathbf{A}	Rank one elliptic genus formula	13
В	Playing with Jacobi theta functions	13

1 Introduction

Two dimensional supersymmetric gauge theories have rich dynamics and exhibit IR dualities similar to their cousins in higher dimensions. In fact, there is a web of connections between the various dualities that goes all the way to six dimensions upon compactification. Deciphering this underlying structure is a long standing goal. A common theme is to use localization techniques to verify and solidify geometric intuition coming from string theory and M-theory realizations. Based on a such a string theory realization, we have recently conjectured a set of dualities relating two dimensional $\mathcal{N} = (4,4)$ supersymmetric gauge theories with orthogonal and symplectic gauge groups [1], generalizing a conjecture relating unitary gauge theories made by Brodie in [2]. In this work, we implement the elliptic genus machinery of [3, 4] to further study these dualities.

Gauge theories in two dimensions based on the so(k) algebra are particularly interesting and subtle for two reasons. First there are several possibilities for the global structure of the group, including SO(k), $O(k)_{\pm}$, Spin(k) and $Pin^{\pm}(k)$, which give rise to distinct gauge theories, with a distinct spectrum of local and line operators [5, 6].³ Second, except for Spin(k), the theories may possess multiple topological sectors that contribute in the path integral. For example an SO(k) theory (with k > 2) on S^2 has two topological sectors

The $O(k)_{\pm}$ theories correspond to \mathbb{Z}_2 orbifolds of the SO(k) theory by charge conjugation and by charge conjugation combined with $(-1)^F$, respectively [7].

corresponding to monopole number $n \in \{0,1\}$, weighted by a phase given by $\exp in\theta$, where θ is a discrete analog of a theta parameter taking values in $\{0,\pi\}$. Investigating these theories therefore requires one to classify the different gauge bundles on the given spacetime manifold.

Orthogonal gauge heories with $\mathcal{N} = (2, 2)$ supersymmetry and fundamental chiral multiplets have been argued to satisfy a set of Seiberg-like dualities given by [7]

$$SO(k) + Q_i \iff O(N - k + 1)_+ + q_i + S_{ij}$$
 (1)

$$O(k)_{+} + Q_{i} \iff SO(N - k + 1) + q_{i} + S_{ij}$$
 (2)

$$O(k)_{-} + Q_i \iff O(N - k + 1)_{-} + q_i + S_{ij}, \qquad (3)$$

where $i=1,\ldots,N$, and the theory on the RHS also has a gauge singlet S in the symmetric representation of the flavor U(N) symmetry, and a superpotential W=qSq. The value of the discrete theta parameter θ is fixed in each case such that the Coulomb branch is lifted. These dualities were further tested by [8, 9]. In particular the T^2 partition functions, or elliptic genus, of the dual theories for $k \leq 4$ were computed, and found to agree, in [9]. A brane realization of these dualities in M theory was subsequently provided in [1].

Interestingly, a very similar brane construction suggested an analogous set of dualities for $\mathcal{N} = (4,4)$ theories with fundamental hypermultiplets given roughly by

$$(S)O(k) + Q_i \iff (S)O(2n - k + 1) + q_i, \qquad (4)$$

where $i=1,\ldots,n$. There are no extra singlets in this case, and $\theta=\pi$ on both sides. The brane construction does not seem to differentiate O(k) from SO(k), or $O(k)_+$ from $O(k)_-$, so we were not able to make a more refined set of conjectures as in the $\mathcal{N}=(2,2)$ case. More concretely, the $\mathcal{N}=(4,4)$ duality is the statement that the two gauge theories flow to the same superconformal theory on the Higgs branch. This is supported in part by the fact that the central charges of the "magnetic" and "electric" Higgs branch SCFT's agree:

$$\widehat{c}_m = (2n - k + 1)n - \frac{1}{2}(2n - k + 1)(2n - k) = kn - \frac{1}{2}k(k - 1) = \widehat{c}_e.$$
 (5)

The aim of this work is to provide further evidence for, and to make more precise, the $\mathcal{N} = (4,4)$ dualities of orthogonal theories via the elliptic genus.

In section 2 we recall the structure of flat O(k) bundles on the torus, and classify them according to whether or not they admit Pin^+ structure, which is relevant for the dependence on the discrete theta parameter. Then in section 3 we test the $\mathcal{N}=(4,4)$ dualities for orthogonal gauge theories by comparing the elliptic genera for some low rank cases. We also test a twisted generalization of the dualities and show that the three pairs of dualities are connected by gauging of a global \mathbb{Z}_2 symmetry. In the appendix we briefly review the elliptic genus formulas and then analytically compare the elliptic genera of the lower rank dualities using the known identities of the Jacobi-theta functions.

2 Flat O(k) connections on T^2

Flat O(k) connections on T^2 were classified in [10]. There are eight sectors in general given by the following pairs of Wilson lines along the temporal and spatial circles of the torus, (U_1, U_2) . For O(2p) with $p \ge 2$:

$$(U_{1}, U_{2})_{ab}^{+} = \begin{cases} (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}), \operatorname{diag}(e^{iu_{2i}\sigma_{2}})) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1, 1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1, -1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1, -1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1, 1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1, -1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1, -1)) \end{cases}$$

$$(6)$$

$$(U_{1}, U_{2})_{ab}^{-} = \begin{cases} (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1, -1, -1, 1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1, 1, -1, -1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, -1, -1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1, -1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1, -1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, -1, -1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1, -1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, -1, 1)), \end{cases}$$

$$(7)$$

where (a, b) = (0, 0), (0, 1), (1, 0) and (1, 1), respectively, and for O(2p + 1) with $p \ge 1$:

$$(U_{1}, U_{2})_{ab}^{+} = \begin{cases} (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, -1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, -1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, -1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, -1)) \end{cases}$$
(8)

$$(U_{1}, U_{2})_{ab}^{-} = \begin{cases} (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, -1, -1, 1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1, -1, -1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, -1, -1, 1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1, -1, 1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1, -1, 1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, -1, -1, 1)) \\ (\operatorname{diag}(e^{iu_{1i}\sigma_{2}}, 1, 1, -1), \operatorname{diag}(e^{iu_{2i}\sigma_{2}}, 1, -1, 1)) . \end{cases}$$
(9)

For O(1) there are only the four sectors in (8),

$$(U_1, U_2)_{ab}^+ = (1, 1), (1, -1), (-1, 1), (-1, -1),$$
 (10)

and for O(2) the first sector in (7) is absent, leaving only seven sectors,

$$(U_1, U_2)_{ab}^+ = (e^{iu_1\sigma_2}, e^{iu_2\sigma_2}), (\mathbb{1}, \sigma_3), (\sigma_3, \mathbb{1}), (\sigma_3, \sigma_3)$$

$$(U_1, U_2)_{01,10,11}^- = (-\mathbb{1}, \sigma_3), (\sigma_3, -\mathbb{1}), (\sigma_3, -\sigma_3).$$

$$(11)$$

2.1 Pin structure

An O(k) bundle over X lifts to a $Pin^{\pm}(k)$ bundle over X provided that $w_2 = 0$ and $w_2 + w_1^2 = 0$, respectively, where w_1 and w_2 are the first and second Stiefel-Whitney classes of the O(k) bundle. We are specifically interested in bundles over T^2 with a flat connection. The corresponding sector in the path integral is accompanied by a phase $e^{i\theta w_2}$, where $\theta \in \{0, \pi\}$ is the discrete theta parameter. We therefore concentrate on the case

of $Pin^+(k)$. Namely we would like to classify flat O(k) connections on T^2 according to whether they lift to flat $Pin^+(k)$ connections on T^2 .

A closely related problem was considered in the context of the toroidal compactification of Type I string theory by Witten in [11]. Since the precise gauge symmetry of Type I string theory is $Spin(32)/\mathbb{Z}_2$, one can consider a compactification on T^2 with a flat connection that does not lift to a flat Spin(32) connection. This was called "toroidal compactification without vector structure". The condition for the existence of vector structure is $\tilde{w}_2 = 0$, where \tilde{w}_2 is defined in an analogous manner to w_2 . A flat $Spin(32)/\mathbb{Z}_2$ connection on T^2 is given by a pair U_1, U_2 satisfying $U_1U_2 = U_2U_1$. The lift to Spin(32) then satisfies $U_1U_2 = (-1)^{\tilde{w}_2}U_2U_1$. For our case, flat O(k) connections on T^2 lift to $Pin^+(k)$ connections satisfying $U_1U_2 = (-1)^{w_2}U_2U_1$.

We will show that the flat O(k) connections $(U_1, U_2)^+$ correspond to bundles with Pin^+ structure, namely $w_2 = 0$, whereas $(U_1, U_2)^-$ correspond to bundles without Pin^+ structure, namely $w_2 \neq 0$. The latter will therefore contribute to the T^2 partition function with a phase $e^{i\theta}$.

Let us recall the definition of Pin(k) [12]. The group Pin(k) is a double cover of the group O(k), in the same way that Spin(k) is a double cover of SO(k). There is a two-to-one map $\rho: Pin(k) \to O(k)$ that takes an element $\Lambda \in Pin(k)$ into the element $R \in O(k)$ given by

$$R_{ij} = \pm \frac{1}{k} \text{Tr}(\Lambda \gamma_i \Lambda^{\dagger} \gamma_j) , \qquad (12)$$

where the sign depends on whether Λ consists of an even or odd number of gamma matrices. More explicitly $Pin^{\pm}(k)$ is generated by $S_{ij} = \frac{1}{2}[\gamma_i, \gamma_j]$, together with a parity operator P that satisfies $\det(\rho(P)) = -1$, and $P^2 = \pm 1$. For $Pin^+(k)$ we can take $P = \gamma_i$ for some i for even k, and $P = \prod_{i=1}^k \gamma_i$ for odd k. In these cases $\rho(P) = \operatorname{diag}(+1, \ldots, -1, \ldots, +1)$ for even k and $\rho(P) = \operatorname{diag}(-1, \ldots, -1)$ for odd k.

Let us first consider the k=2,3,4 cases explicitly, and then generalize to all k. For k=2 we can choose $\gamma_1=\sigma_1, \ \gamma_2=\sigma_2, \ \text{and} \ P=\gamma_1=\sigma_1$. The map $Pin^+(2)\to O(2)$ takes the form

$$\Lambda(\phi) = e^{i\frac{\phi}{2}S} = \begin{pmatrix} e^{i\frac{\phi}{2}} & 0\\ 0 & e^{-i\frac{\phi}{2}} \end{pmatrix} \quad \mapsto \quad \rho(\Lambda) = R(\phi) = \begin{pmatrix} \cos(\phi) & \sin(\phi)\\ -\sin(\phi) & \cos(\phi) \end{pmatrix}$$
(13)

$$P = \sigma_1 \quad \mapsto \quad \rho(P) = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \,. \tag{14}$$

Using these relations one can easily find lifts to $Pin^+(2)$ of the commuting flat O(2) connections in (11). For $(U_1, U_2)^+$ we have

$$(\Lambda(u_1), \Lambda(u_2)) \mapsto (e^{iu_1\sigma_2}, e^{iu_2\sigma_2})$$

$$(\mathbb{1}, P\Lambda(\pi)) \mapsto (\mathbb{1}, \sigma_3)$$

$$(P\Lambda(\pi), \mathbb{1})) \mapsto (\sigma_3, \mathbb{1})$$

$$(P\Lambda(\pi), P\Lambda(\pi)) \mapsto (\sigma_3, \sigma_3).$$

$$(15)$$

The $Pin^+(2)$ holonomy pairs on the LHS all commute and so correspond to $w_2 = 0$. For $(U_1, U_2)^-$ on the other hand we find

$$(\Lambda(\pi), P\Lambda(\pi)) \mapsto (-1, \sigma_3)$$

$$(P\Lambda(\pi), \Lambda(\pi)) \mapsto (\sigma_3, -1)$$

$$(P\Lambda(\pi), P) \mapsto (\sigma_3, -\sigma_3),$$

$$(16)$$

which anti-commute in $Pin^+(2)$ and therefore correspond to $w_2 \neq 0$. For k = 3 we can take $\gamma_1 = \sigma_1 \otimes \mathbb{1}$, $\gamma_2 = \sigma_3 \otimes \mathbb{1}$, $\gamma_3 = \sigma_2 \otimes \sigma_2$, and $P = \gamma_1 \gamma_2 \gamma_3 = \mathbb{1} \otimes \sigma_2$. In this case $\rho(P) = \text{diag}(-1, -1, -1)$. The lifts to $Pin^+(3)$, up to signs, of the $(U_1, U_2)^+$ holonomies in (8) are given by the commuting pairs

$$(\Lambda_{12}(u_1), \Lambda_{12}(u_2)), (\Lambda_{12}(u_1), P\Lambda_{12}(u_2)), (P\Lambda_{12}(u_1), \Lambda_{12}(u_2)), (P\Lambda_{12}(u_1), P\Lambda_{12}(u_2)), (17)$$

and the lifts of $(U_1, U_2)^-$ are given by the anti-commuting pairs

$$(\Lambda_{12}(\pi), \Lambda_{23}(\pi)), (\Lambda_{12}(\pi), P\Lambda_{31}(\pi)), (P\Lambda_{31}(\pi), \Lambda_{12}(\pi)), (P\Lambda_{12}(\pi), P\Lambda_{31}(\pi)).$$
 (18)

As before, the former correspond to $w_2 = 0$ and the latter to $w_2 \neq 0$. For k = 4 we can take $\gamma_1 = i\sigma_2 \otimes i\sigma_2 \otimes \mathbb{1}$, $\gamma_2 = i\sigma_2 \otimes \sigma_1 \otimes i\sigma_2$, $\gamma_3 = i\sigma_2 \otimes \sigma_3 \otimes i\sigma_2$, $\gamma_4 = \sigma_3 \otimes \mathbb{1} \otimes \mathbb{1}$, and $P = \gamma_4$. In this case $\rho(P) = \text{diag}(1, 1, 1, -1)$. The O(4) pairs $(U_1, U_2)^+$ lift to the commuting pairs

$$(\Lambda_{12}(u_{11})\Lambda_{34}(u_{12}), \Lambda_{12}(u_{21})\Lambda_{34}(u_{22})), \quad (\Lambda_{12}(u_{11}), P\Lambda_{12}(u_{21})), \tag{19}$$

$$(P\Lambda_{12}(u_{11}), \Lambda_{12}(u_{21})), (P\Lambda_{12}(u_{11}), P\Lambda_{12}(u_{21})),$$
 (20)

and the pairs $(U_1, U_2)^-$ lift to the anti-commuting pairs

$$(\Lambda_{23}(\pi), \Lambda_{34}(\pi)), \quad (\Lambda_{12}(u_1)\Lambda_{34}(\pi), P\Lambda_{12}(u_2)),$$
 (21)

$$(P\Lambda_{12}(u_1), \Lambda_{12}(u_2)\Lambda_{34}(\pi)), (P\Lambda_{12}(u_1), P\Lambda_{12}(u_2)\Lambda_{34}(\pi)).$$
 (22)

The generalization to higher k is straightforward. In (6)-(9) we see that the O(2p) and O(2p+1) bundles decompose into a direct sum of commuting flat connections of $SO(2)^{p-1}$ and those of O(4) and O(3), respectively. The first part lifts trivially to a flat $Spin(2)^{p-1}$ bundle, since Spin(2) = SO(2). To conclude, the pairs $(U_1, U_2)^+$ define flat O(k) bundles on T^2 with trivial w_2 , while the pairs $(U_1, U_2)^-$ define flat O(k) bundles on T^2 with non-trivial w_2 .

3 $\mathcal{N} = (4,4)$ theories

Two dimensional gauge theories with $\mathcal{N}=(4,4)$ supersymmetry have a moduli space of vacua that generically has two branches, a Coulomb branch and a Higgs branch. The two branches decouple in the IR and the theory flows to two independent superconformal field theories with an $SU(2) \times SU(2)$ R-symmetry [13]. In some cases the Coulomb branch is lifted and one is left with just the Higgs branch SCFT. In particular this is the case for

O(k) or SO(k) with any number N of fundamental hypermultiplets and with $\theta = \pi$ [1]. This is somewhat surprising given the fact that in theories with eight supersymmetries in higher dimensions the Coulomb branch cannot be lifted. It is in fact very similar to what happens in $\mathcal{N} = (2, 2)$ supersymmetric gauge theories [7], and is due partly to an effective linear twisted superpotential on the Coulomb branch.⁴ We will also see that it is consistent with the results we will present below: the Witten index is a finite integer, as it should be for a regular theory.

As stated in the introduction, the brane construction of [1] suggests a duality between the O(k) (or SO(k)) theory with n fundamental hypermultiplets and the O(2n - k + 1) (or SO(2n - k + 1) theory with n fundamental hypermultiplets. We will provide more evidence for this conjecture by comparing the elliptic genera. Our analysis shows that, as in the $\mathcal{N} = (2, 2)$ theories, there are actually three dualities between $\mathcal{N} = (4, 4)$ theories:

$$SO(k) + Q_i \longleftrightarrow O(2n - k + 1)_+ + q_i$$
 (23)

$$O(k)_{+} + Q_{i} \iff SO(2n - k + 1) + q_{i}$$
 (24)

$$O(k)_{-} + Q_i \iff O(2n - k + 1)_{-} + q_i, \tag{25}$$

where $\theta = \pi$ in all cases.

To compute the elliptic genus of the $\mathcal{N}=(4,4)$ theories we use the results of [3, 4] for the $\mathcal{N}=(2,2)$ theories, add the contribution of the adjoint chiral superfield, and take into account the condition on the global symmetry and R-symmetry charges imposed by the superpotential $W=Q\Phi Q$. This was actually done for the U(k) theory in [4], so we will follow their notation convention. We denote the holonomies of the global symmetry $Sp(n) \times U(1)_{\Phi}$ by ξ_{α} , and λ , and that of the left-moving U(1) R-symmetry by z. As in [4] we assign R-charge 0 to all fields but include the holonomy λ . The superpotential will impose the constraint $\lambda=z$.

It will also be useful to label the flat connections $(U_1, U_2)^{\pm}$ by their exponents $u^{\pm} = (u_1^{\pm}, \dots, u_k^{\pm})$. In the 2×2 blocks $e^{iu_1i\sigma_2}$, $e^{iu_2i\sigma_2}$ with continuous elements, the two associated u parameters are given by the two eigenvalues $\pm (u_{1i} + \tau u_{2i})$. In the blocks with discrete numbers, we assign $u_i = 0$ for an eigenvalue pair (1,1), $u_i = \frac{1}{2}$ for an eigenvalue pair (-1,1), $u_i = \frac{\tau}{2}$ for (1,-1), and $u_i = \frac{1+\tau}{2}$ for (-1,-1). We therefore have

$$u_{ab}^{+} = \begin{cases} (\pm u_{1}, \dots, \pm u_{p}) \\ (\pm u_{1}, \dots, \pm u_{p-1}, 0, \frac{\tau}{2}) \\ (\pm u_{1}, \dots, \pm u_{p-1}, 0, \frac{1}{2}) \\ (\pm u_{1}, \dots, \pm u_{p-1}, 0, \frac{1+\tau}{2}) \end{cases} \quad u_{ab}^{-} = \begin{cases} (\pm u_{1}, \dots, \pm u_{p-2}, 0, \frac{1}{2}, \frac{1+\tau}{2}, \frac{\tau}{2}) \\ (\pm u_{1}, \dots, \pm u_{p-1}, \frac{1}{2}, \frac{1+\tau}{2}) \\ (\pm u_{1}, \dots, \pm u_{p-1}, \frac{\tau}{2}, \frac{1+\tau}{2}) \\ (\pm u_{1}, \dots, \pm u_{p-1}, \frac{1}{2}, \frac{\tau}{2}) \end{cases}$$
(26)

⁴The twisted superpotential lifts the part of the Coulomb branch corresponding to the scalar in the $\mathcal{N}=(2,2)$ vector multiplet. Presumably a linear superpotential is also generated for the adjoint chiral multiplet, though this has not been demonstrated yet.

for O(2p), and

$$u_{ab}^{+} = \begin{cases} (\pm u_1, \dots, \pm u_p, 0) \\ (\pm u_1, \dots, \pm u_p, \frac{\tau}{2}) \\ (\pm u_1, \dots, \pm u_p, \frac{1}{2}) \\ (\pm u_1, \dots, \pm u_p, \frac{1+\tau}{2}) \end{cases} \quad u_{ab}^{-} = \begin{cases} (\pm u_1, \dots, \pm u_{p-1}, \frac{1}{2}, \frac{1+\tau}{2}, \frac{\tau}{2}) \\ (\pm u_1, \dots, \pm u_{p-1}, \frac{1}{2}, \frac{1+\tau}{2}, 0) \\ (\pm u_1, \dots, \pm u_{p-1}, \frac{\tau}{2}, \frac{1+\tau}{2}, 0) \\ (\pm u_1, \dots, \pm u_{p-1}, \frac{1}{2}, \frac{\tau}{2}, 0) \end{cases}$$
(27)

for O(2p+1).

3.1 O(1) + n

In this case there is only the contribution of the matter fields:

$$Z_{ab}^{O(1),n} = \frac{1}{2} \prod_{\alpha=1}^{n} \frac{\theta_1(-z \pm \xi_\alpha + \frac{a+b\tau}{2})}{\theta_1(\pm \xi_\alpha + \frac{a+b\tau}{2})},$$
(28)

where the Jacobi's theta function definitions and the elliptic genus formulas are given in the appendix and we have also introduced the shorthand $\theta_1(A \pm B) \equiv \theta_1(A + B)\theta_1(A - B)$

3.2 O(2) + n

The continuous sector is given by

$$Z_{00}^{O(2),n} = -\frac{1}{2} \frac{i\eta(q)^3}{\theta_1(-z)} \frac{\theta_1(\lambda - z)}{\theta_1(\lambda)} \oint_{u_*} du \prod_{\alpha=1}^n \frac{\theta_1(u - z \pm \xi_\alpha)}{\theta_1(u \pm \xi_\alpha)} \frac{\theta_1(-u - z \pm \xi_\alpha)}{\theta_1(-u \pm \xi_\alpha)}$$
$$= \sum_{\beta=1}^n \frac{\theta_1(2\xi_\beta \pm z)}{\theta_1(2\xi_\beta)^2} \prod_{\alpha \neq \beta}^n \frac{\theta_1(\xi_\beta \pm \xi_\alpha - z)}{\theta_1(\xi_\beta \pm \xi_\alpha)} \frac{\theta_1(-\xi_\beta \pm \xi_\alpha - z)}{\theta_1(-\xi_\beta \pm \xi_\alpha)}$$

where we have picked up 2n residues at $u_* = \pm \xi_{\beta}$ and imposed the constraint $\lambda = z$. In particular the $\theta_1(0)$ factor in the denominator cancels with the $\theta_1(\lambda - z)$ factor in the numerator. The six discrete sectors are

$$Z_{ab\pm}^{O(2),n} = \frac{1}{4} \frac{\theta_1(u_{1ab}^{\pm} + u_{2ab}^{\pm})}{\theta_1(-z + u_{1ab}^{\pm} + u_{2ab}^{\pm})} \frac{\theta_1(\lambda - z + u_{1ab}^{\pm} + u_{2ab}^{\pm})}{\theta_1(\lambda + u_{1ab}^{\pm} + u_{2ab}^{\pm})} \prod_{\alpha=1}^n \prod_{i=1}^2 \frac{\theta_1(u_{iab}^{\pm} - z \pm \xi_{\alpha})}{\theta_1(u_{iab}^{\pm} \pm \xi_{\alpha})}$$

$$= \frac{1}{4} \frac{\theta_1(u_{1ab}^{\pm} + u_{2ab}^{\pm})^2}{\theta_1(\pm z + u_{1ab}^{\pm} + u_{2ab}^{\pm})} \prod_{\alpha=1}^n \prod_{i=1}^2 \frac{\theta_1(u_{iab}^{\pm} - z \pm \xi_{\alpha})}{\theta_1(u_{iab}^{\pm} \pm \xi_{\alpha})}$$
(29)

with (a,b) = (0,1), (1,0) and (1,1), where u_{iab}^{\pm} are given in (26).

3.3 O(3) + n

There are four continuous sectors given by

$$Z_{ab+}^{O(3),n} = -\frac{1}{4} \frac{i\eta(q)^3}{\theta_1(-z)} \oint_{u_*} du \left(\frac{\theta_1(u + \frac{a+b\tau}{2})}{\theta_1(u - z + \frac{a+b\tau}{2})} \frac{\theta_1(-u + \frac{a+b\tau}{2})}{\theta_1(-u - z + \frac{a+b\tau}{2})} \right) \times \frac{\theta_1(u + \lambda - z + \frac{a+b\tau}{2})}{\theta_1(u + \lambda + \frac{a+b\tau}{2})} \frac{\theta_1(-u + \lambda - z + \frac{a+b\tau}{2})}{\theta_1(-u + \lambda + \frac{a+b\tau}{2})} \frac{\theta_1(\lambda - z + \frac{a+b\tau}{2})}{\theta_1(\lambda + \frac{a+b\tau}{2})} \times \prod_{i=\alpha}^n \frac{\theta_1(u - z \pm \xi_\alpha)}{\theta_1(u \pm \xi_\alpha)} \frac{\theta_1(-u - z \pm \xi_\alpha)}{\theta_1(-u \pm \xi_\alpha)} \frac{\theta_1(-z \pm \xi_\alpha + \frac{a+b\tau}{2})}{\theta_1(\pm \xi_\alpha + \frac{a+b\tau}{2})} \right).$$
(30)

The JK poles are at $u_* = \{z - \frac{a+b\tau}{2}, -\lambda - \frac{a+b\tau}{2}, \pm \xi_{\beta}\}$. Evaluating the residues is straightforward but cumbersome, so we will not show it here. The four discrete sectors are given by

$$Z_{ab-}^{O(3),n} = \frac{1}{8} \prod_{i \neq j}^{3} \frac{\theta_{1}(u_{iab}^{-} + u_{jab}^{-})}{\theta_{1}(-z + u_{iab}^{-} + u_{jab}^{-})} \frac{\theta_{1}(\lambda - z + u_{iab}^{-} + u_{jab}^{-})}{\theta_{1}(\lambda + u_{iab}^{-} + u_{jab}^{-})} \prod_{\alpha=1}^{n} \prod_{i=1}^{3} \frac{\theta_{1}(-z + u_{iab}^{-} \pm \xi_{\alpha})}{\theta_{1}(u_{iab}^{-} \pm \xi_{\alpha})}$$

$$= \frac{1}{8} \prod_{i \neq j}^{3} \frac{\theta_{1}(u_{iab}^{-} + u_{jab}^{-})^{2}}{\theta_{1}(\pm z + u_{iab}^{-} + u_{jab}^{-})} \prod_{\alpha=1}^{n} \prod_{i=1}^{3} \frac{\theta_{1}(-z + u_{iab}^{-} \pm \xi_{\alpha})}{\theta_{1}(u_{iab}^{-} \pm \xi_{\alpha})}$$

$$(31)$$

where u_{ab-}^{i} are given in (27).

3.4 $\mathcal{N} = (4,4)$ duality

Before we compare the partitions functions let us comment on their orbifold structure. Being a special case of the (2,2)-orthogonal theories, the $(4,4) - O(k)_{\pm}$ theories can also be obtained by gauging a \mathbb{Z}_2 global symmetry in the SO(k) theory. This symmetry is either charge conjugation, or charge conjugation combined with $(-1)^{F_s}$. The former is the so-called *standard orbifold*, and the latter, *non-standard* orbifold. In terms of the torus partition function this means

$$Z_{T^2}/\mathbb{Z}_2 = \frac{1}{2}(Z_{00} + Z_{10} + Z_{01} + Z_{11}),$$
 (32)

$$Z_{T^2}/\mathbb{Z}_2(-1)^{F_s} = \frac{1}{2}(-Z_{00} + Z_{10} + Z_{01} + Z_{11}).$$
 (33)

In [7], the $O(k)_{\pm}$ theories where then defined according to the parity of N+k such that the three dualities always hold in the same form. For the (4,4) theories since matter comes in hypermultiplets, we define (for any k) the $O(k)_{+}$ theory as the *standard orbifold* of SO(k) and the $O(k)_{-}$ theory as the *non-standard orbifold* of SO(k) (up to an overall sign).

The elliptic genus is then given by

$$Z_{T^2}^{SO(k),n} = 2(Z_{00+}^{O(k),n} + y^{-2n}e^{i\theta}Z_{00-}^{O(k),n}), \tag{34}$$

$$Z_{T^2}^{O(k)_+,n} = \sum_{a,b} (y^{-nb} Z_{ab+}^{O(k),n} + y^{n(b-2)} e^{i\theta} Z_{ab-}^{O(k),n})$$
(35)

$$Z_{T^2}^{O(k)_-,n} = \sum_{a,b} (-1)^{ab+a+b+k} (y^{-nb} Z_{ab+}^{O(k),n} + y^{n(b-2)} e^{i\theta} Z_{ab-}^{O(k),n})$$
 (36)

where the factors of $y=e^{2\pi iz}$ are required for modular invariance. Setting $\theta=\pi$ for the regular theories, we have analytically showen (see appendix B) that the following equalities hold

$$Z_{T^2}^{SO(2),1} = Z_{T^2}^{O(1)_+,1},$$
 (37)

$$Z_{T^2}^{O(2)_+,1} = Z_{T^2}^{SO(1),1},$$
 (38)

$$Z_{T^2}^{O(2)_-,1} = Z_{T^2}^{O(1)_-,1}.$$
 (39)

For the higher rank dualities we have checked numerically (up to order q^5 , where $q = e^{2\pi i \tau}$) that the following partition functions agree

$$Z_{T^2}^{SO(3),2} = Z_{T^2}^{O(2)_+,2},$$
 (40)

$$Z_{T^2}^{O(3)_+,2} = Z_{T^2}^{SO(2),2},$$
 (41)

$$Z_{T^2}^{O(3)_-,2} = Z_{T^2}^{O(2)_-,2},$$
 (42)

confirming the proposed dualities in these cases.

In the limit $z \to 0$ we obtain the Witten index, which is given by

$$I_W^{SO(1),n} = I_W^{O(1)_-,n} = 1, I_W^{O(1)_+,n} = 2,$$
 (43)

$$I_W^{O(2)_+,n} = I_W^{O(2)_-,n} = n, I_W^{SO(2),n} = 2n,$$
 (44)

and

$$I_W^{SO(3),n} = I_W^{O(3)_-,n} = n, I_W^{O(3)_+,n} = 2n.$$
 (45)

3.5 Twisting

⁵ As mentioned in the previous section the The $O(k)_{\pm}$ theories are obtained by gauging the SO(k) theory by a \mathbb{Z}_2 subgroup of its global symmetry that acts as charge conjugation.

⁵The following discussion is parallel for the (2,2) orthogonal theories defined in [7].

Their elliptic genera are therefore given by twisting the SO(k) partition function and summing over the twists. The twisted SO(k) elliptic genus is

$$Z_{T^{2}(\alpha,\beta)}^{SO(k),n} = 2(Z_{\alpha\beta+}^{O(k),n} + y^{-2n}e^{i\theta}Z_{\alpha\beta-}^{O(k),n})$$
(46)

and the sums are given by (35-36).

The resulting orbifold theories $O(k)_{\pm}$ posses in turn a quantum $\widehat{\mathbb{Z}}_2$ - global symmetry which acts non-trivially on the twisted (untwisted) sectors for the standard(non-standard) orbifolds. Their twisted partition function is therefore given by

$$Z_{T^{2}(\alpha,\beta)}^{O(k)_{+,n}} = \sum_{a,b} (y^{-nb} Z_{ab+}^{O(k),n} + y^{n(b-2)} e^{i\theta} Z_{ab-}^{O(k),n}) (-1)^{\alpha b + \beta a}, \tag{47}$$

$$Z_{T^{2}(\alpha,\beta)}^{O(k)_{-},n} = (-1)^{k\alpha\beta} \sum_{a,b} (-1)^{ab+a+b+k} (y^{-nb} Z_{ab+}^{O(k),n} + y^{n(b-2)} e^{i\theta} Z_{ab-}^{O(k),n}) (-1)^{\alpha(b+1)+\beta(a+1)}$$

$$(48)$$

where $(-1)^{k\alpha\beta}$ is a background term.

Setting $\theta = \pi$ we can now write a twisted generalization of the (4, 4)-dualities

$$Z_{T^2(\alpha,\beta)}^{SO(k),n} = Z_{T^2(\alpha,\beta)}^{O(2n-k+1)_+,n},$$
 (49)

$$Z_{T^2(\alpha,\beta)}^{O(k)_+,n} = Z_{T^2(\alpha,\beta)}^{SO(2n-k+1),n},$$
 (50)

$$Z_{T^2(\alpha,\beta)}^{O(k)_{-,n}} = Z_{T^2(\alpha,\beta)}^{O(2n-k+1)_{-,n}}.$$
 (51)

As a consistency check let us gauge the \mathbb{Z}_2 symmetry by summing over the twists. For example, for k = n = 1 we find

$$1/2\sum_{\alpha\beta} Z_{T^2(\alpha,\beta)}^{SO(1),1} = Z_{(0,0)}^{O(1)_{+},1},$$
 (52)

$$1/2\sum_{\alpha\beta}(-1)^{\alpha\beta+\alpha+\beta+1}Z_{T^{2}(\alpha,\beta)}^{SO(1),1} = Z_{(0,0)}^{O(1)-,1},$$
(53)

$$1/2\sum_{\alpha\beta} Z_{T^2(\alpha,\beta)}^{O(1)_+,1} = Z_{(0,0)}^{SO(1),1}, \tag{54}$$

$$1/2\sum_{\alpha\beta}(-1)^{\alpha\beta+\alpha+\beta}Z_{T^{2}(\alpha,\beta)}^{O(1)_{+},1} = Z_{(0,0)}^{O(1)_{-},1},$$
 (55)

$$1/2\sum_{\alpha\beta}(-1)^{\alpha\beta+\alpha+\beta}Z_{T^{2}(\alpha,\beta)}^{O(1)_{-},1} = Z_{(0,0)}^{O(1)_{+},1},$$
(56)

$$1/2\sum_{\alpha\beta}(-1)^{\alpha\beta+\alpha+\beta+1}Z_{T^{2}(\alpha,\beta)}^{O(1)_{-,1}} = Z_{(0,0)}^{SO(1),1},$$
(57)

and for k=2, n=1

$$1/2\sum_{\alpha\beta} Z_{T^2(\alpha,\beta)}^{SO(2),1} = Z_{(0,0)}^{O(2)_{+},1},$$
 (58)

$$1/2\sum_{\alpha\beta}(-1)^{\alpha\beta+\alpha+\beta}Z_{T^{2}(\alpha,\beta)}^{SO(2),1} = Z_{(0,0)}^{O(2)_{-},1},$$
(59)

$$1/2\sum_{\alpha\beta} Z_{T^2(\alpha,\beta)}^{O(2)_{+,1}} = Z_{(0,0)}^{SO(2)_{,1}}, \tag{60}$$

$$1/2\sum_{\alpha\beta}(-1)^{\alpha\beta+\alpha+\beta+1}Z_{T^{2}(\alpha,\beta)}^{O(2)_{+},1} = Z_{(0,0)}^{O(2)_{-},1}, \tag{61}$$

$$1/2\sum_{\alpha\beta}(-1)^{\alpha\beta+\alpha+\beta+1}Z_{T^{2}(\alpha,\beta)}^{O(2)_{-},1} = Z_{(0,0)}^{O(2)_{+},1}, \tag{62}$$

$$1/2\sum_{\alpha\beta} Z_{T^2(\alpha,\beta)}^{O(2)_{-,1}} = Z_{(0,0)}^{SO(2)_{,1}}.$$
 (63)

Denoting the gauging operations $1/2 \sum_{\alpha\beta}$ and $1/2 \sum_{\alpha\beta} (-1)^{\alpha\beta+\alpha+\beta+1}$ by \mathbb{Z}_2 and $\mathbb{Z}_2(-1)^{F_s} \equiv \mathbb{Z}_2'$ respectively and an overall sign by additional "-", we can summarize these relations with the following diagram

$$SO(1) \iff O(2)_{+}$$

$$\downarrow^{\mathbb{Z}'_{2}} \qquad \mathbb{Z}'_{2} \downarrow$$

$$O(1)_{-} \iff O(2)_{-}$$

$$-\mathbb{Z}'_{2} \downarrow^{\mathbb{Z}_{2}} \qquad -\mathbb{Z}'_{2} \downarrow^{\mathbb{Z}_{2}}$$

$$O(1)_{+} \iff SO(2)$$

$$(64)$$

This is similar to Hori's (2, 2)-orthogonal dualities [7], where starting from any duality the other two follow by the two gauging procedures.

4 Conclusions

Using the elliptic genus machinery we investigated the brane based duality conjuncture of 2d (4,4)-orthogonal gauge theories. Successfully testing the $O(1) \leftrightarrow O(2)$ and $O(2) \leftrightarrow O(3)$ cases. Our analysis reviles a similar structure to the (2,2)-dualities found in [7] where there are three theories and three pairs of dualities, all connected by gauging a global Z_2 symmetry.

On the way we have classified the O(k) flat connections on the torus. This has important application for any elliptic genus computation of orthogonal theories when the θ parameter of the theory is non-trivial.

These theories deserve further investigation, in particular it would be interesting to understand the mechanism that lifts the part of coulomb branch associated (in the (2,2) language) with the additional adjoint scalar.

5 Acknowledgements

We thank Hyungchul Kim, Jaemo Park, Shlomo Razamat, and Gabi Zafrir for useful discussions. This work is supported in part by the Israel Science Foundation under grant no. 1390/17.

A Rank one elliptic genus formula

The elliptic genus of a gauge theory with gauge group G of rank one is given by [3]

$$Z_{T^2} = -\frac{1}{|W|} \sum_{u_j \in M_{\text{sing}}^+} \oint_{u=u_j} \frac{i\eta(q)^3}{\theta_1(-z)} \prod_{\alpha \in G} \frac{\theta_1(\alpha u)}{\theta_1(\alpha u - z)} \prod_{\Phi_i} \prod_{\rho \in \mathcal{R}_i} \frac{\theta_1\left(\frac{R_i}{2} - 1\right)z + \rho u + \xi_i\right)}{\theta_1\left(\frac{R_i}{2}z + \rho u + \xi_i\right)}$$
(65)

where |W| is the order of the Weyl group, $\eta(q)$ is the Dedekind eta function

$$\eta(q) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$$
(66)

and θ_1 is the Jacobi theta function

$$\theta_1(\tau|z) = -iq^{\frac{1}{8}}y^{\frac{1}{2}} \prod_{k=1}^{\infty} (1 - q^k)(1 - yq^k)(1 - y^{-1}q^{k-1})$$
(67)

with $q \equiv e^{2\pi i \tau}$ and $y \equiv e^{2\pi i z}$. The first two terms of the integrand are the contribution of gauge multiplet and the double product comes from the matter fields. The integral is evaluated using the Jeffrey-Kirwan pole prescription which for rank one gauge group amounts to taking the poles that are inside the unit circle consistently e.g if $u = z - \frac{a+b\tau}{2} \in M_{\rm sing}^+$ then $u = -z + \frac{a+b\tau}{2}$ does not.

B Playing with Jacobi theta functions

In this appendix we derive the first three dualities $Z_{T^2}^{S/O(2)_{\pm},1} = Z_{T^2}^{S/O(1)_{\pm},1}$ using the known identities for the Jacobi theta functions.

The Jacobi theta functions are defined as follows

$$\theta_1(\tau|z) = -iq^{\frac{1}{8}}y^{\frac{1}{2}} \prod_{k=1}^{\infty} (1 - q^k)(1 - yq^k)(1 - y^{-1}q^{k-1}), \tag{68}$$

$$\theta_2(\tau|z) = q^{\frac{1}{8}} y^{\frac{1}{2}} \prod_{k=1}^{\infty} (1 - q^k) (1 + yq^k) (1 + y^{-1}q^{k-1}), \tag{69}$$

$$\theta_3(\tau|z) = \prod_{k=1}^{\infty} (1 - q^k)(1 + yq^k)(1 + y^{-1}q^{k-1}), \tag{70}$$

$$\theta_4(\tau|z) = \prod_{k=1}^{\infty} (1 - q^k)(1 - yq^k)(1 - y^{-1}q^{k-1}), \tag{71}$$

they are known to satisfy many relations, we list here the ones that we need (we will use the shorthand $\theta_i(\tau|z) \equiv \theta_i(z), \theta_i(\tau|0) \equiv \theta_i$).

Shift symmetries:

$$\theta_1(-z) = -\theta_1(z),\tag{72}$$

$$\theta_1(z+a+b\tau) = (-1)^{a+b} e^{-2\pi i bz - \pi i b^2 \tau} \theta_1(z). \tag{73}$$

$$\theta_{1}(z + \frac{1}{2}) = \theta_{2}(z), \qquad \theta_{4}(z + \frac{1}{2}) = \theta_{3}(z),
\theta_{1}(z + \frac{1+\tau}{2}) = q^{-\frac{1}{8}}y^{-\frac{1}{2}}\theta_{3}(z), \qquad \theta_{4}(z + \frac{1+\tau}{2}) = q^{-\frac{1}{8}}y^{-\frac{1}{2}}\theta_{2}(z),
\theta_{1}(z + \frac{\tau}{2}) = iq^{-\frac{1}{8}}y^{-\frac{1}{2}}\theta_{4}(z), \qquad \theta_{4}(z + \frac{\tau}{2}) = iq^{-\frac{1}{8}}y^{-\frac{1}{2}}\theta_{1}(z).$$
(74)

Addition rules:

$$\theta_1(x \pm z) = \frac{1}{\theta_4^2} (\theta_1(x)^2 \theta_4(z)^2 - \theta_4(x)^2 \theta_1(z)^2), \tag{75}$$

$$\theta_2(x \pm z) = \frac{1}{\theta_4^2} (\theta_2(x)^2 \theta_4(z)^2 - \theta_3(x)^2 \theta_1(z)^2), \tag{76}$$

$$\theta_3(x \pm z) = \frac{1}{\theta_4^2} (\theta_3(x)^2 \theta_4(z)^2 - \theta_2(x)^2 \theta_1(z)^2), \tag{77}$$

$$\theta_4(x \pm z) = \frac{1}{\theta_4^2} (\theta_4(x)^2 \theta_4(z)^2 - \theta_1(x)^2 \theta_1(z)^2). \tag{78}$$

Duplication formulas:

$$\theta_1(2x) = \frac{2\theta_1(x)\theta_2(x)\theta_3(x)\theta_4(x)}{\theta_2\theta_3\theta_4},\tag{79}$$

$$\theta_4(2x) = \frac{1}{\theta_4^3} (\theta_4(x)^4 - \theta_1(x)^4) = \frac{1}{\theta_4^3} (\theta_3(x)^4 - \theta_2(x)^4). \tag{80}$$

Square identities:

$$\theta_1(x)^2 \theta_4^2 = \theta_3(x)^2 \theta_2^2 - \theta_2(x)^2 \theta_3^2, \tag{81}$$

$$\theta_2(x)^2 \theta_4^2 = \theta_4(x)^2 \theta_2^2 - \theta_1(x)^2 \theta_3^2, \tag{82}$$

$$\theta_3(x)^2 \theta_4^2 = \theta_4(x)^2 \theta_3^2 - \theta_1(x)^2 \theta_2^2, \tag{83}$$

$$\theta_4(x)^2 \theta_4^2 = \theta_3(x)^2 \theta_3^2 - \theta_2(x)^2 \theta_2^2. \tag{84}$$

We can now derive the proposed dualities. Starting with $Z_{T^2}^{SO(2),1}=Z_{T^2}^{O(1)_+,1}$ we need to show that

$$\frac{1}{4} \sum_{a,b=0}^{1} \frac{\theta_1(\xi + \frac{a+b\tau}{2} \pm z)}{\theta_1(\xi + \frac{a+b\tau}{2})^2} - \frac{\theta_1(2\xi \pm z)}{\theta_1(2\xi)^2} = 0.$$
 (85)

Using identity (75) we get

$$\frac{1}{4} \sum_{a,b=0}^{1} \left(\frac{\theta_4(z)^2}{\theta_4^2} - \frac{\theta_1(z)^2 \theta_4(\xi + \frac{a+b\tau}{2})^2}{\theta_4^2 \theta_1(\xi + \frac{a+b\tau}{2})^2} \right) - \left(\frac{\theta_4(z)^2}{\theta_4^2} - \frac{\theta_1(z)^2 \theta_4(2\xi)^2}{\theta_4^2 \theta_1(2\xi)^2} \right) = 0, \tag{86}$$

so it is left to show that

$$\frac{1}{4} \sum_{a,b=0}^{1} \frac{\theta_4(\xi + \frac{a+b\tau}{2})^2}{\theta_1(\xi + \frac{a+b\tau}{2})^2} - \frac{\theta_4(2\xi)^2}{\theta_1(2\xi)^2} = 0.$$
 (87)

Using the relations (74-74) on the first term and (79-80) on the second we get

$$\frac{1}{4} \left(\frac{\theta_4(\xi)^2}{\theta_1(\xi)^2} + \frac{\theta_1(\xi)^2}{\theta_4(\xi)^2} + \frac{\theta_3(\xi)^2}{\theta_2(\xi)^2} + \frac{\theta_2(\xi)^2}{\theta_3(\xi)^2} \right) - \frac{\theta_2^2 \theta_3^2}{4\theta_4^4} \frac{(\theta_4(\xi)^4 - \theta_1(\xi)^4)^2}{\theta_1(\xi)^2 \theta_2(\xi)^2 \theta_3(\xi)^2 \theta_4(\xi)^2} = 0$$
 (88)

or

$$(\theta_4(\xi)^4 + \theta_1(\xi)^4)\theta_2(\xi)^2\theta_3(\xi)^2 + (\theta_2(\xi)^4 + \theta_3(\xi)^4)\theta_1(\xi)^2\theta_4(\xi)^2 - \frac{\theta_2^2\theta_3^2}{\theta_4^4}(\theta_4(\xi)^4 - \theta_1(\xi)^4)^2 = 0,$$
(89)

finally using (81-84) we get an equality.

Next we show that $Z_{T^2}^{O(2)_+,1}=Z_{T^2}^{SO(1),1}$. Starting with the LHS, we have the continuous sector

$$Z_{00}^{O(2),1} = \frac{\theta_1(2\xi \pm z)}{\theta(2\xi)^2} \tag{90}$$

and the six discrete sectors which can be written in the following form

$$Z_{ab+}^{O(2),1} = \frac{1}{4} \frac{\theta_1(\xi \pm z)\theta_1(\frac{a+b\tau}{2})^2\theta_1(\xi \pm z + \frac{a+b\tau}{2})}{\theta_1(\xi)^2\theta_1(z + \frac{a+b\tau}{2})^2\theta_1(\xi + \frac{a+b\tau}{2})^2}$$
(91)

$$Z_{ab-}^{O(2),1} = \frac{1}{4} y^{(a+1)(1-b)} \frac{\theta_1(\frac{a+b\tau}{2})^2 \theta_1(\xi \pm z + \frac{1-ab+\tau}{2}) \theta_1(\xi \pm z + \frac{b+a(1-b)\tau}{2})}{\theta_1(z + \frac{a+b\tau}{2})^2 \theta_1(\xi + \frac{1-ab+\tau}{2})^2 \theta_1(\xi + \frac{b+a(1-b)\tau}{2})^2}$$
(92)

placing in (35) and expanding the theta functions such that each contains only one fugacity (using identities (75-80)) we end up with

$$Z_{T^{2}}^{O(2)+,1} = \frac{\theta_{4}(z)^{2}}{\theta_{4}^{2}} - \frac{\theta_{2}^{2}\theta_{3}^{2}\theta_{1}(z)^{2}(\theta_{4}(\xi)^{4} - \theta_{1}(\xi)^{4})^{2}}{4\theta_{4}^{6}\theta_{1}(\xi)^{2}\theta_{2}(\xi)^{2}\theta_{3}(\xi)^{2}\theta_{4}(\xi)^{2}}$$

$$+ \frac{\theta_{2}^{2}\left(\theta_{1}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{4}(\xi)^{2}\theta_{1}(z)^{2}\right)\left(\theta_{2}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{3}(x)^{2}\theta_{1}(z)^{2}\right)}{4\theta_{4}^{4}\theta_{1}(\xi)^{2}\theta_{2}(\xi)^{2}\theta_{2}(z)^{2}}$$

$$- \frac{\theta_{2}^{2}\left(\theta_{4}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{1}(\xi)^{2}\theta_{1}(z)^{2}\right)\left(\theta_{3}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{2}(\xi)^{2}\theta_{1}(z)^{2}\right)}{4\theta_{4}^{4}\theta_{3}(\xi)^{2}\theta_{4}(\xi)^{2}\theta_{2}(z)^{2}}$$

$$+ \frac{\theta_{3}^{2}\left(\theta_{1}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{4}(\xi)^{2}\theta_{1}(z)^{2}\right)\left(\theta_{3}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{2}(\xi)^{2}\theta_{1}(z)^{2}\right)}{4\theta_{4}^{4}\theta_{1}(\xi)^{2}\theta_{3}(\xi)^{2}\theta_{3}(z)^{2}}$$

$$- \frac{\theta_{3}^{2}\left(\theta_{4}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{1}(\xi)^{2}\theta_{1}(z)^{2}\right)\left(\theta_{2}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{3}(\xi)^{2}\theta_{1}(z)^{2}\right)}{4\theta_{4}^{4}\theta_{2}(\xi)^{2}\theta_{4}(\xi)^{2}\theta_{3}(z)^{2}}$$

$$+ \frac{\left(\theta_{1}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{4}(\xi)^{2}\theta_{1}(z)^{2}\right)\left(\theta_{4}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{1}(\xi)^{2}\theta_{1}(z)^{2}\right)}{4\theta_{4}^{2}\theta_{1}(\xi)^{2}\theta_{4}(\xi)^{2}\theta_{4}(z)^{2}}$$

$$- \frac{\left(\theta_{2}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{3}(\xi)^{2}\theta_{1}(z)^{2}\right)\left(\theta_{3}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{2}(\xi)^{2}\theta_{1}(z)^{2}\right)}{4\theta_{4}^{2}\theta_{2}(\xi)^{2}\theta_{3}(\xi)^{2}\theta_{4}(z)^{2}}$$

$$- \frac{\left(\theta_{2}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{3}(\xi)^{2}\theta_{1}(z)^{2}\right)\left(\theta_{3}(\xi)^{2}\theta_{4}(z)^{2} - \theta_{2}(\xi)^{2}\theta_{1}(z)^{2}\right)}{4\theta_{4}^{2}\theta_{2}(\xi)^{2}\theta_{3}(\xi)^{2}\theta_{4}(z)^{2}}$$

For the last step, comparing with the dual side

$$Z_{T^2}^{SO(1),1} = \frac{\theta_4(z)^2}{\theta_4^2} - \frac{\theta_4(x)^2 \theta_1(z)^2}{\theta_4^2 \theta_1(x)^2}$$
(94)

we use *Mathematica* to simplify the expression (93) minus (94) with the assumptions of identities (81-84), resulting with the expected zero.

The third duality $Z_{T^2}^{O(2)_-,1}=Z_{T^2}^{O(1)_-,1}$ is also easy to prove following the same steps.

References

[1] O. Bergman and E. Avraham, "Branes and 2d $\mathcal{N}=(2,2)$ gauge theories with orthogonal and symplectic groups," JHEP **1808**, 023 (2018) doi:10.1007/JHEP08(2018)023 [arXiv:1804.00694 [hep-th]].

- [2] J. H. Brodie, "Two-dimensional mirror symmetry from M theory," Nucl. Phys. B **517**, 36 (1998) doi:10.1016/S0550-3213(97)00755-4 [hep-th/9709228].
- [3] F. Benini, R. Eager, K. Hori and Y. Tachikawa, "Elliptic genera of two-dimensional N=2 gauge theories with rank-one gauge groups," Lett. Math. Phys. **104**, 465 (2014) doi:10.1007/s11005-013-0673-y [arXiv:1305.0533 [hep-th]].
- [4] F. Benini, R. Eager, K. Hori and Y. Tachikawa, "Elliptic Genera of 2d $\mathcal{N}=2$ Gauge Theories," Commun. Math. Phys. **333**, no. 3, 1241 (2015) doi:10.1007/s00220-014-2210-y [arXiv:1308.4896 [hep-th]].
- [5] O. Aharony, N. Seiberg and Y. Tachikawa, "Reading between the lines of four-dimensional gauge theories," JHEP 1308, 115 (2013) doi:10.1007/JHEP08(2013)115 [arXiv:1305.0318 [hep-th]].
- [6] C. Cordova, P. S. Hsin and N. Seiberg, "Global Symmetries, Counterterms, and Duality in Chern-Simons Matter Theories with Orthogonal Gauge Groups," SciPost Phys. 4, 021 (2018) doi:10.21468/SciPostPhys.4.4.021 [arXiv:1711.10008 [hep-th]].
- [7] K. Hori, "Duality In Two-Dimensional (2,2) Supersymmetric Non-Abelian Gauge Theories," JHEP 1310, 121 (2013) doi:10.1007/JHEP10(2013)121 [arXiv:1104.2853 [hep-th]].
- [8] C. Closset, N. Mekareeya and D. S. Park, "A-twisted correlators and Hori dualities," JHEP 1708, 101 (2017) doi:10.1007/JHEP08(2017)101 [arXiv:1705.04137 [hep-th]].
- [9] H. Kim and J. Park, "2D Seiberg-like dualities for orthogonal gauge groups," arXiv:1710.06069 [hep-th].
- [10] J. Kim, S. Kim, K. Lee, J. Park and C. Vafa, "Elliptic Genus of E-strings," 1709, 098 (2017) doi:10.1007/JHEP09(2017)098 [arXiv:1411.2324 [hep-th]].
- [11] E. Witten, "Toroidal compactification without vector structure," JHEP **9802**, 006 (1998) doi:10.1088/1126-6708/1998/02/006 [hep-th/9712028].
- [12] M. F. Atiyah, R. Bott and A. Shapiro, Topology 3, S3 (1964). doi:10.1016/0040-9383(64)90003-5
- [13] E. Witten, "On the conformal field theory of the Higgs branch," JHEP **9707**, 003 (1997) doi:10.1088/1126-6708/1997/07/003 [hep-th/9707093].