

Stochastic Loewner Evolutions, Fuchsian Systems and Orthogonal Polynomials

Igor Loutsenko and Oksana Yermolayeva

Abstract

We find a wide class of Levy-Loewner evolutions for which the value of integral means beta-spectrum $\beta(q)$ at $q = 2$ is the maximal real eigenvalue of a three-diagonal matrix. The second moments of derivatives of corresponding conformal mappings are expressed through solutions of matrix Fuchsian systems with three singular points.

1 Introduction

The Schramm-Loewner Evolution is so far the only model where a multi-fractal spectrum is not trivial and known explicitly (see e.g. [2, 3, 6, 7, 11, 17, 18]). Explicit computation of harmonic spectrum is almost impossible task for deterministic fractals and it is a difficult task even for random fractals. Apart from the case of the Schramm-Loewner evolution, two exact results for the second moment of harmonic measure were obtained in the study of Levy-Loewner Evolution (LLE) [7, 19]. Extending this study, we will present an infinite set of non-trivial examples.

The Bounded Whole-Plane Levy-Loewner evolution (also called exterior whole-plane LLE)¹ is a stochastic process of the growth of the curve out of the origin $z = 0$ in the complex plane z (see Figure 1).

The curve is described by time dependent conformal mapping $z = F(w, t)$ from exterior of the unit circle $|w| > 1$ in the w -plane to the complement of the curve in the z plane. This mapping obeys the Levy-Loewner equation

$$\frac{\partial F(w, t)}{\partial t} = w \frac{\partial F(w, t)}{\partial w} \frac{w + e^{iL(t)}}{w - e^{iL(t)}}, \quad \lim_{w \rightarrow \infty} F'(w, t) = e^t, \quad -\infty < t < \infty, \quad (1)$$

where $L(t)$ is a Levy process and prime denotes the w -derivative. The limit $t \rightarrow -\infty$ corresponds to the origin of the curve.

Here we consider the Levy processes without a drift. Without loss of generality we set

$$\langle L(t) \rangle = 0, \quad (2)$$

where $\langle \rangle$ denote expectation (ensemble average).

The LLE is a conformally invariant stochastic process in the sense that the time evolution is consistent with composition of conformal maps. When $L(t)$ is a continuous function of time, the conformal mapping $z = F(w, t)$ describes growth of a random continuous curve. The only continuous (modulo uniform drift) process of Levy type is the Brownian motion

$$L(t) = \sqrt{\kappa} B(t), \quad \langle (B(t + \tau) - B(t))^2 \rangle = |\tau|, \quad (3)$$

where the positive parameter κ is the “temperature” of the Brownian motion. The stochastic Loewner evolution driven by Brownian motion is called Schramm-Loewner Evolution or SLE_{κ} . Since it describes non-branching planar stochastic curves with a conformally-invariant probability distribution, SLE is a useful tool for description of boundaries of critical clusters in two-dimensional equilibrium statistical mechanics. In this picture, different κ correspond to different classes of models of statistical mechanics (a good introduction to SLE for physicists can be found e.g. in [4, 8] as well as mathematical reviews are given e.g. in [15, 16]).

On the other hand, in the general case, when $L(t)$ is a discontinuous function of time the conformal mapping describes stochastic (infinitely) branching curve.

The unbounded (or interior) version of the whole-plane LLE is an inversion $F(w, t) \rightarrow 1/F(1/w, t)$ of the above bounded version. Obviously, this mapping from an interior of the unit disc in the w -plane to the complement of stochastic curve which grows from infinity towards the origin in the z -plane also satisfies the Loewner equation (1) with different asymptotic conditions given now at $w = 0$

$$F'(w = 0, t) = e^{-t}.$$

¹For a short introduction to the whole-plane LLE see e.g. [19], [14]. A good introduction to chordal LLE can be found in [21, 22]. For a quick introduction to Levy processes see e.g. [1].




Figure 1: Bounded whole-plane LLE: Time dependent mapping F from the exterior of the unit circle in the w -plane to the complement of the curve in z -plane. Point $w = e^{iL(t)}$ on the unit circle is mapped to the growing tip of the curve.

This version of LLE has been studied mainly due to its relationship with the problem of Bieberbach coefficients of conformal mappings [7, 17, 19].

To get the multi-fractal spectrum of LLE one has to find first the so-called “ β -spectrum”²: The integral means $\beta(q)$ -spectrum of the domain is defined through the q th moment of a derivative of conformal mapping at the unit circle (i.e. at $|w| \rightarrow 1$) as follows

$$\beta(q) = \overline{\lim}_{\epsilon \rightarrow 0^\pm} \frac{\log \int_0^{2\pi} \langle |F'(e^{\epsilon+i\varphi})|^q \rangle d\varphi}{-\log \pm \epsilon}, \quad (4)$$

where \pm signs correspond to bounded and unbounded whole-plane LLE (in what follows the upper/lower signs will correspond to the bounded/unbounded version respectively).

To find $\beta(q)$ -spectrum (4) one needs to estimate moments of derivative: One can show that the value

$$\rho = e^{\mp qt} \langle |F'(e^{iL(t)} w, t)|^q \rangle,$$

is time independent and is a function of³ w, \bar{w} and q only [2, 3, 7, 17, 18, 19]. Moreover it satisfies the linear integro-differential equation

$$\mathcal{L}^\pm \rho = 0, \quad (5)$$

where the linear operator \mathcal{L}^\pm equals

$$\mathcal{L}^\pm = -\hat{\eta} + w \frac{w+1}{w-1} \partial_w + \bar{w} \frac{\bar{w}+1}{\bar{w}-1} \partial_{\bar{w}} - \frac{q}{(w-1)^2} - \frac{q}{(\bar{w}-1)^2} + q \mp q. \quad (6)$$

In (6), the linear operator $\hat{\eta}$ acts on functions of w, \bar{w} as follows (for details see e.g. [2], [7] or [19])

$$\hat{\eta} \rho(w, \bar{w}) = \lim_{t \rightarrow 0} \frac{1}{t} \int_0^{2\pi} (\rho(w, \bar{w}) - \rho(e^{i\varphi} w, e^{-i\varphi} \bar{w})) P(\varphi, t) d\varphi,$$

where $P(\varphi, t)$ is the probability density that $L(t) = \varphi$ under condition $L(0) = 0$. One can present operator $\hat{\eta}$ in the more convenient (integro-differential) form as

$$\hat{\eta} \rho(w, \bar{w}) = \frac{\kappa}{2} (w \partial_w - \bar{w} \partial_{\bar{w}})^2 \rho(w, \bar{w}) + \int_{-\pi}^{\pi} (\rho(w, \bar{w}) - \rho(e^{i\varphi} w, e^{-i\varphi} \bar{w})) d\eta(\varphi), \quad d\eta(\varphi) \geq 0 \quad (7)$$

²For introduction to multi-fractal analysis and relationship between different kinds of multi-fractal spectra see e.g. [2], [9], [10], [11], [23].

³ \bar{w} denotes complex conjugate of w

where $d\eta(\varphi)$ is a symmetric Levy measure. In polar coordinates (r, ϕ) , such that $w = re^{i\phi}, \bar{w} = re^{-i\phi}$, the operator $\hat{\eta}$ acts only wrt angular variable ϕ , i.e. $\hat{\eta}$ commutes with r and

$$\hat{\eta}f(\phi) = -\frac{\kappa}{2}\partial_{\phi}^2 f(\phi) + \int_{-\pi}^{\pi} (f(\phi) - f(\phi + \varphi)) d\eta(\varphi)$$

whatever is dependence of f on r . In other words, it acts diagonally on the basis $w^n \bar{w}^m$

$$\hat{\eta}w^n \bar{w}^m = \eta_{n-m} w^n \bar{w}^m, \quad n, m \in \mathbb{Z}. \quad (8)$$

Since we consider only Levy processes without drift (2) (i.e. with Levy measure symmetric wrt reflection $\phi \rightarrow -\phi$), the characteristic coefficients η_n are real, non-negative and symmetric

$$\eta_n = \bar{\eta}_n = \eta_{-n}, \quad \eta_0 = 0.$$

They express through the Levy measure as

$$\eta_n = \frac{\kappa n^2}{2} + \int_{-\pi}^{\pi} (1 - \cos n\phi) d\eta(\phi). \quad (9)$$

The above equation is a particular case of Levy-Khintchine formula for a class of Levy processes we deal with.

The probability density $P(\phi, t) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} e^{-\eta_n t + in\phi}$ is a fundamental solution of the integro-differential equation of the parabolic type on the unit circle

$$\partial_t P(\phi, t) = -\hat{\eta}P(\phi, t), \quad P(\phi, 0) = \delta(\phi).$$

Note that the particular case $d\eta(\phi) = 0$ corresponds to the Schramm-Loewner evolution SLE_{κ} . Here, eq. (5,6) becomes the second-order differential equation with

$$\hat{\eta} = -\frac{\kappa}{2}\partial_{\phi}^2, \quad \eta_n = \frac{\kappa n^2}{2},$$

which allows to find exact form of the multi-fractal spectrum of the SLE_{κ} [2, 3, 11, 18].

In the case of the bounded whole-plane LLE one can use analyticity of $\rho(w, \bar{w})$ at infinity as boundary conditions for linear equation (5,6). Namely (see e.g. [19] for details), in this version of LLE ρ has following asymptotic expansion at $w \rightarrow \infty$

$$\rho = \sum_{i=0, j=0}^{\infty} \frac{\rho_{i,j}}{w^i \bar{w}^j}, \quad \rho_{0,0} = 1$$

with expansion coefficients $\rho_{i,j}$ fixed uniquely by equation (5,6). In other word, once analytic, non-vanishing at infinity solution of (5,6) is found, it gives (modulo constant factor) expectation of the moment of the derivative of conformal mappings of the bounded whole-plane LLE.

Similarly, for the unbounded version of LLE one has to look for non-vanishing analytic solution of the corresponding equation, this time at the origin $w = 0$

$$\rho = \sum_{i=0, j=0}^{\infty} \rho_{i,j} w^i \bar{w}^j, \quad \rho_{0,0} = 1.$$

Similarly to the bounded case, such a solution is unique.

2 Beta Spectrum of the Unbounded Whole-Plane LLE and Fuchsian Systems

Let us try to find the value of β -spectrum at $q = 2$: Representing $\rho(w, \bar{w})$ in the form

$$\rho(w, \bar{w}) = (1-w)(1-\bar{w})\Theta(w, \bar{w}), \quad (10)$$

from (5,6) with $q = 2$ we get

$$-\hat{\eta}[(1-w)(1-\bar{w})\Theta] + (w+1)(\bar{w}-1)w \frac{\partial \Theta}{\partial w} + (\bar{w}+1)(w-1)\bar{w} \frac{\partial \Theta}{\partial \bar{w}} + 3(2w\bar{w} - w - \bar{w})\Theta = 0, \quad (11)$$

where Θ is the series

$$\Theta = \theta_0(\xi) + \sum_{i=1}^{\infty} (w^i + \xi^i w^{-i}) \theta_i(\xi), \quad \xi = r^2 = w\bar{w}. \quad (12)$$

Substituting it into (11), and taking into account the facts that operator $\hat{\eta}$ commutes with $\xi = r^2 = w\bar{w}$ and that, according to (8), $\hat{\eta}[w^i] = \eta_i w^i$, from (11,12) we get a three-term differential recurrence relation for $\theta_i(\xi)$.

$$2\xi(\xi - 1)\theta'_i(\xi) - (\eta_i + i + (\eta_i - i - 6)\xi)\theta_i(\xi) + \xi(\eta_i + i - 2)\theta_{i+1}(\xi) + (\eta_i - i - 2)\theta_{i-1}(\xi) = 0, \quad \theta_{-i}(\xi) = \xi^i \theta_i(\xi) \quad (13)$$

It is easy to see that this recurrence relation has a solution that truncates at $i = N$, i.e. $\theta_i = 0$ for $i \geq N$, when $\eta_N = N + 2$ for some N .

The simplest truncation happens when $N = 1$, i.e. when $\eta_1 = 3$, which corresponds to the case of conjecture by authors of [7] proved by us in [19]. Before studying the general case of an arbitrary N we will consider separately the case of truncation at $N = 2$. Here not only $\beta(2)$, but also $\rho(w, \bar{w})$ can be found explicitly for arbitrary η_i , $i \neq 2$. We have the following

Theorem 1. *For the unbounded whole plane LLE with $\eta_2 = 4$ the integral means β -spectrum has the following value at $q = 2$*

$$\beta(2) = \frac{6 - \eta_1 + \sqrt{\eta_1^2 - 4\eta_1 + 12}}{2}, \quad (14)$$

Proof. The proof is based on explicit computation of $\rho(w, \bar{w})$. It is convenient to re-parametrize $\eta_1 \geq 0$ through β given by lhs of eq.(14), i.e.

$$\eta_1 = \frac{\beta^2 - 6\beta + 6}{2 - \beta}, \quad 2 < \beta < 3 + \sqrt{3}, \quad (15)$$

and introduce new dependent variables f_0, f_1 such that $\theta_i(\xi) = (1 - \xi)^{-\beta} f_i(\xi)$.

In the case of truncation at $N = 2$ (i.e. when $\eta_2 = 4$), system (13) writes as

$$\xi(\xi - 1)f'_0 + (3 - \beta)\xi f_0 - 2\xi f_1 = 0 \quad (16)$$

$$2\xi(\xi - 1)f'_1 - (\eta_1 + 1 + (\eta_1 - 7 + 2\beta)\xi) f_1 + (\eta_1 - 3)f_0 = 0 \quad (17)$$

Then expressing f_1 through f_0, f'_0 into (16) and substituting the result in (17) we get the hypergeometric equation for f_0

$$\xi(\xi - 1)f''_0 + ((a + b + 1)\xi - c) f'_0 + abf_0 = 0$$

where

$$a = \frac{\beta^2 - 5\beta + 8}{2(2 - \beta)}, \quad b = 3 - \beta, \quad c = \frac{\beta^2 - 7\beta + 8}{2(2 - \beta)} \quad (18)$$

Since we are looking for $\rho(w, \bar{w})$ which is non-vanishing and analytic at $w = 0$, from all linearly independent solutions of the above hypergeometric equation we choose the one which is non-vanishing and analytic at $\xi = 0$, i.e.

$$f_0 = {}_2F_1(a, b, c; \xi)$$

Then from (16) it follows that

$$f_1 = \frac{\xi - 1}{2} f'_0 + \frac{3 - \beta}{2} f_0 = \frac{ab}{2c} (\xi - 1) {}_2F_1(a + 1, b + 1, c + 1; \xi) + \frac{3 - \beta}{2} {}_2F_1(a, b, c; \xi)$$

and according to (10,12)

$$\rho(w, \bar{w}) = (1 - w)(1 - \bar{w})(1 - w\bar{w})^{-\beta} (f_0(w\bar{w}) + (w + \bar{w})f_1(w\bar{w})).$$

According to (4,10,12), the value of the β -spectrum equals the blowup rate (also called ‘‘polynomial’’ growth rate) of $\theta_0 - \theta_1 = (1 - \xi)^{-\beta} (f_0(\xi) - f_1(\xi))$ at $\xi \rightarrow 1$. Using the Gauss identity

$${}_2F_1(a, b, c; 1) = \frac{\Gamma(c)\Gamma(c - a - b)}{\Gamma(c - a)\Gamma(c - b)}, \quad \Re(c - a - b) > 0,$$

and taking eqs. (15, 18) into account, it is easy to see that $f_0 - f_1$ is finite and non-vanishing at $\xi = 1$. Therefore, the blowup rate of $\theta_0 - \theta_1$ equals β and is given by lhs of eq.(14). This completes the proof. \square

Remark 1. Similar statement is valid for the bounded version of LLE for the matrix given by (29). We will refer to both statements as Lemma 1.

Proof. It is known (see e.g. [13]) that an asymptotics of a general solution of the Fuchsian system (19) in a neighborhood of singular point $\xi = 1$ has the form

$$\theta \rightarrow \sum_{l=0}^{N-1} (1-\xi)^{-\beta_l} f^{(l)}(C, \xi), \quad \xi \rightarrow 1, \quad (24)$$

where β_l is an eigenvalue of B and $C = \{C_0, \dots, C_{N-1}\}$ are arbitrary constants. If matrix B is diagonalizable and non-resonant⁴, the vector functions $f^{(l)}$ are finite at $\xi = 1$ and $f^{(l)}(C, \xi) = C_l \tilde{f}^{(l)}$, with $\tilde{f}^{(l)}$ being an eigenvector of B corresponding to an eigenvalue β_l . If B is not diagonalizable or/and resonant, $f^{(l)}(C, \xi)$ are polynomials in $x = \log(1-\xi)$ (i.e. they are of finite non-negative integer degrees in x).

From (24) and (12) it follows that the corresponding solution of (11) has the following asymptotics at $\xi \rightarrow 1$ (or $r \rightarrow 1$ in polar coordinates (r, ϕ) , $w = re^{i\phi}$)

$$\Theta(r, \phi) \rightarrow \sum_{l=0}^{N-1} (1-r)^{-\text{Re}\beta_l} \sum_{m=0}^{M_l} G_{lm}(\phi) x^m \cos(x\text{Im}\beta_l - g_{lm}(\phi)), \quad x = \log(1-r^2), \quad r \rightarrow 1,$$

where all functions $G_{lm}(\phi)$ are bounded. As $r \rightarrow 1$, the above finite sum will be dominated by term(s) with the maximal $\text{Re}\beta_l$ and, if there are several of them, one has to choose the one(s) with the maximal M_l among them. In other words

$$\Theta \rightarrow (1-r)^{-\max \text{Re}\beta_l} x^M \mathcal{G}(x, \phi), \quad r \rightarrow 1, \quad (25)$$

where M is some non-negative integer and $\mathcal{G}(x, \phi)$ is a sum of finite number of the cosine terms

$$\mathcal{G}(x, \phi) = \sum_{\alpha_i \in \{\text{Im}\beta_1, \dots, \text{Im}\beta_{N-1}\}} \mathcal{G}_i(\phi) \cos(\alpha_i x - \gamma_i(\phi)). \quad (26)$$

From (25, 26) it follows that at least one eigenvalue with maximal real part has zero imaginary part. Indeed, let us suppose that the above is not true. Then all α_i in (26) are non-zero. Since (26) is the sum of a finite number of the cosine terms with x running along the semi-infinite interval, $\mathcal{G}(x, \phi)$ is not of constant sign on this interval. Therefore, any real solution of (11) that truncates at N in (12), including an analytic at $\xi = 0$ solution we are looking for, oscillates, changing its sign as $\xi \rightarrow 1$. However, an analytic at $\xi = 0$ solution cannot change sign at $\xi < 1$, since both $\rho(w, \bar{w})$ and $(1-w)(1-\bar{w})$ are not negative at $\xi < 1$. Therefore, there exists a real eigenvalue which is not less than the real part of any non-real eigenvalue.

Next, if all real β_l were negative, then (according to (25)) any non-negative solution would have a negative blow-up rate and therefore $\beta(2)$ given by (4) would be negative. This is also a contradiction, since the integral means β -spectrum must be non-negative (see e.g. [20]). Thus, matrix B must have at least one non-negative eigenvalue, and $\beta(2)$ equals to one of non-negative eigenvalues of B . □

By consequence, in the $N = 2$ case all eigenvalues are real⁵ (as confirmed by Theorem 1). Note that here both eigenvalues can be positive, but $\beta(2)$ always equals the maximal one.

Another useful example where $\beta(2)$ and spectrum of B can be found explicitly is the N -truncated unbounded SLE, i.e. the unbounded whole-plane SLE $_{\kappa}$ with $\kappa = 2(N+2)/N^2$ (for details see [18])

$$\eta_n = \frac{N+2}{N^2} n^2, \quad n = 0..N,$$

$$\beta_l = \frac{N+2 - (2N^2 - 3N - 6)l + 2(N+2)l^2}{N^2}, \quad l = 0..N-1. \quad (27)$$

Here all eigenvalues β_l are real, while not all a_n are positive. The value of the integral means β spectrum at $q = 2$ equals the maximal eigenvalue β_{N-1} , i.e. $\beta(2) = \max\{\beta_l, l = 0 \dots N-1\} = (5N-2)/N$ (see [18, 7]).

⁴Matrix is resonant if it has eigenvalues which differ by a non-zero integer

⁵Also, one can check that in the $N = 3$ case all roots of cubic characteristic polynomial are always real for any $\eta_1, \eta_2 \geq 0$, whatever are signs of a_n .




Figure 2: Dependence of $\beta(2)$ of the PLE_λ on the arrival rate $\lambda = N - 2$ of the Poisson process. Numerical results suggest that $\beta(2) \rightarrow 1$ as $\lambda \rightarrow \infty$. Results of numerical computations of the maximal eigenvalue in general case (i.e. case of infinite matrices) is shown separately in the zoomed region by the solid line.

The next proposition considers generic nontrivial $N < 5$ cases

Theorem 3. *For the bounded whole-plane LLE with $\eta_N = N - 2$, where $N < 5$, the value of the integral means β -spectrum at $q = 2$ is the maximal real eigenvalue of the three-diagonal matrix (29).*

Proof. By direct computation, from (22,30) we get

$$P_3 = \beta^3 + (\eta_1 + \eta_2 + 3)\beta^2 + \left(2 + \eta_1 + \frac{5}{4}\eta_2 + \frac{3}{4}\eta_2\eta_1\right)\beta - \frac{1}{4}\eta_2\eta_1 - \frac{3}{4}\eta_2,$$

$$P_4 = \beta^4 + (\eta_2 + \eta_1 + \eta_3 + 4)\beta^3 + \left(\frac{3}{4}\eta_2\eta_1 + \frac{5}{2}\eta_2 + \eta_3\eta_1 + \frac{3}{4}\eta_3\eta_2 + 2\eta_3 + 6 + 2\eta_1\right)\beta^2 +$$

$$+ \left(\frac{1}{2}\eta_3\eta_2\eta_1 + \frac{3}{4}\eta_3\eta_2 + \eta_2 + 4 + \frac{3}{4}\eta_2\eta_1 + 2\eta_1\right)\beta - \frac{3}{4}\eta_3\eta_2 - \frac{1}{4}\eta_3\eta_2\eta_1 - \frac{3}{4}\eta_2 - \frac{1}{4}\eta_2\eta_1.$$

Since $\eta_n > 0$, $n > 0$, the polynomial P_4 has a single negative coefficient. By Descartes' rule of sign it has only one positive root and the rest of roots are negative. The same ⁷ applies to P_3 . By Lemma 1, $\beta(2)$ equals the maximal root of characteristic polynomial. □

Concluding this section we would like to mention another set of examples which are similar to the Hastings-Levitov (HL) models [12], namely, LLE driven by a compound Poisson process with jumps distributed uniformly over the circle (we call them PLE_λ , where λ is the arrival rate of the Poisson process) ⁸. Here, $\eta_n = \lambda = N - 2$, $n > 0$ and the recurrence coefficients a_n , $n = 1 \dots N - 1$ are positive. Therefore all roots of $P_N(\beta)$ are real and simple. We have verified for N up to several hundred that a polynomial $P_N(\beta)$ has the only one change of sign in the sequence of its coefficients. By consequence, in these (verified) models, $\beta(2)$ equals maximal eigenvalue which is the only non-negative eigenvalue (see Figure 2).

4 Conclusions

In the present paper we considered a wide class of the whole-plane LLEs driven by a generic Levy process on a circle, without the drift, restricted by a condition that there exist N for which $\eta_N = N + 2$ in the unbounded

⁷Actually, there is no need to consider P_3 separately since existence of only one non-negative eigenvalue for N automatically implies the same for $N' < N$.

⁸Similarly to the HL model, the PLE_λ is a composition of random elementary “spike” mappings uniformly distributed over the circle. Note, however, that the HL models are not LLEs. LLEs driven by compound Poisson processes have been considered in [14].

version of LLE and $\eta_N = N - 2$ in the bounded version. We showed that the second moment of derivative of the stochastic conformal mapping for the whole-plane LLE is expressed in terms of solution of N -dimensional Fuchsian system with three singular points. The value of the integral means β spectrum at $q = 2$ (i.e. $\beta(2)$) is a non-negative eigenvalue of a three-diagonal matrix. We also were able to show that $\beta(2)$ is actually the maximal eigenvalue for several wide classes of processes.

Therefore, one might try to generalize the Theorem 1, 2 and 3 by dropping the condition of smallness of $\delta\eta_n$ in the Theorem 2. We recall that in the case of Theorems 1, 3 condition of smallness is absent, so we may put forward the following

Conjecture 1. *Let the whole-plane LLE is driven by a Levy process without drift for which there exists an integer N a such that*

- $\eta_N = N + 2$ in the unbounded version of LLE
- $\eta_N = N - 2$ in the bounded version of LLE.

Then, the value of the integral means β -spectrum at $q = 2$ equals the maximal real eigenvalue of the three-diagonal matrix B given by (20) or (29) correspondingly.

Moreover, one can generalize the above conjecture for the case when the truncation conditions do not necessarily hold. Indeed, in this case matrix B is infinite and we can look for a maximal eigenvalue of this infinite matrix. In more detail, one has to take an infinite set of “formally orthogonal” polynomials and look at the sequence of maximal roots (or equally at the sequence of maximal eigenvalues $\beta_{\max}(N)$ of the sub-matrices of dimension N of an infinite matrix B). One expects this sequence to converge to $\beta(2)$ as $N \rightarrow \infty$.

We have checked this hypothesis numerically for SLE_κ (where $\beta(q)$ -spectrum is known for any κ and q) for different κ . Also, for PLE_λ numerical computations suggest that the maximal eigenvalue is a smooth function of λ (see Figure 2). When the N -truncation takes place, the maximal eigenvalue of the infinite matrix equals the one of its sub-matrix of size N , i.e. the sequence $\beta_{\max}(M)$ reaches its limit at $M = N$.

Therefore, one may state the following

Conjecture 2. *Let the whole-plane LLE is driven by a Levy process without drift. Then $\beta(2)$ is a limit of the sequence of maximal real eigenvalues of matrices B_N , where B_N is given by (20) for the unbounded version of LLE or (29) for the bounded version (or, equivalently, the sequence of maximal roots of polynomials set by the recurrence relation (22,23) or (22,30) correspondingly).*

References

- [1] Applebaum D, *Levy Processes - From Probability to Finance and Quantum Groups*, Notices of the AMS, 51 (11): 13361347 (2004)
- [2] D. Beliaev, S. Smirnov, *Harmonic measure and SLE*, Commun. Math. Phys. 290, 577595 (2009).
- [3] D. Beliaev, B. Duplantier, and M. Zinsmeister, *Integral means spectrum of whole-plane SLE*, Comm. Math. Phys. Volume 353, Issue 1, pages 119133, (2017).
- [4] John Cardy, *SLE for theoretical physicists*, Ann.Phys. 318 (2005) 81-118
- [5] T. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, NY, 1978
- [6] B.Duplantier, *Conformally invariant fractals and potential theory*, Phys.Rev.Lett., 84(7): 1363-1367,(2000)
- [7] Bertrand Duplantier, Thi Phuong Chi Nguyen, Thi Thuy Nga Nguyen, Michel Zinsmeister, *The Coefficient Problem and Multifractality of Whole-Plane SLE and LLE*, Annales Henri Poincare, Volume 16, Issue 6, pp 1311 - 1395, (2015)
- [8] Ilya A. Gruzberg, *Stochastic geometry of critical curves, Schramm–Loewner evolutions and conformal field theory*, J. Phys. A: Math. Gen. **39**, no. 41 (2006) 12601–12655.
- [9] Halsey, T. C., Honda, K., Duplantier, B. (1996), *Multifractal dimensions for branched growth*, J.Stat.Phys., 85(5-6), pp 681743

- [10] T.C. Halsey, M.H. Jensen, L.P. Kadanoff, I. Procaccia, and B.I. Shraiman, *Fractal measures and their singularities: the characterization of strange sets*, Phys. Rev. A (3), 33(2): p. 1141, 1986.
- [11] Matthew B. Hastings, *Exact Multifractal Spectra for Arbitrary Laplacian Random Walks*, Phys. Rev. Lett. **88** (2002)
- [12] M.B. Hastings, L.S. Levitov, *Laplacian growth as one-dimensional turbulence*, Physica D, 116 (1998), 244-252
- [13] Yu. Ilyashenko, S. Yakovenko, *Lectures on analytic differential equations*, Graduate Studies in Mathematics, 86. American Mathematical Society, Providence, RI, 2008. xiv+625 pp. ISBN: 978-0-8218-3667-5
- [14] Fredrik Johansson, Alan Sola, *Rescaled Levy-Loewner hulls and random growth*, Bulletin des Sciences Mathematiques, Volume 133, Issue 3, April 2009, Pages 238-256
- [15] Gregory F. Lawler, *Conformally invariant processes in the plane*, Mathematical Surveys and Monographs, **114**, Amer. Math. Soc., Providence, RI, (2005).
- [16] Gregory F. Lawler, *Conformal invariance and 2D statistical physics*, Bull. Amer. Math. Soc. **46** (2009) 35–54.
- [17] Igor Loutsenko, *SLE $_{\kappa}$: correlation functions in the coefficient problem*, J. Phys. A: Math. Theor. 45 275001, 2012,(2012)
- [18] Igor Loutsenko, Oksana Yermolayeva, *On Exact Multi-Fractal Spectrum of the Whole-Plane SLE*, J.Stat. Mech., (2013) doi: 10.1088/1742-5468/2013/04/P4007
- [19] Igor Loutsenko, Oksana Yermolayeva, *New exact results in spectra of stochastic Loewner evolution*, J. Phys. A, 47(16):165202, 15, 2014.
- [20] N. G. Makarov, *Fine structure of harmonic measure*, St. Petersburg Math. J., 10(2):217268, 1999.
- [21] P. Oikonomou, I. Rushkin, I. A. Gruzberg, and L. P. Kadanoff, *Global properties of stochastic Loewner evolution driven by Lévy processes*, J. Stat. Mech. (2008) P01019
- [22] I. Rushkin, P. Oikonomou, L. P. Kadanoff, and I. A. Gruzberg, *Stochastic Loewner evolution driven by Levy processes*, J. Stat. Mech. (2006) P01001
- [23] http://en.wikipedia.org/wiki/Multifractal_system