

On the Existence of a Fixed Spectrum for a Multi-channel Linear System: A Matroid Theory Approach

F. Liu¹ and A. S. Morse¹

Abstract—Conditions for the existence of a fixed spectrum {i.e., the set of fixed modes} for a multi-channel linear system have been known for a long time. The aim of this paper is to reestablish one of these conditions using a new and transparent approach based on matroid theory.

I. INTRODUCTION

The classical “decentralized control” problem considered in [1], [2] focuses on stabilizing or otherwise controlling a $k > 1$ channel linear system of the form

$$\dot{x} = Ax + \sum_{i=1}^k B_i u_i, \quad y_i = C_i x_i \quad (1)$$

Decentralization is enforced by restricting the feedback of each measured signal y_i to only its corresponding control input u_i , possibly through a linear dynamic controller. Wang and Davison [1] were able to show that no matter what these feedback controllers might be, as long as they are finite dimensional and linear time-invariant (LTI), the spectrum of the resulting closed-loop system contains a fixed subset depending only on A , the B_i and the C_i , which they elected to call the set of “fixed modes” of the system. Roughly speaking, the set of fixed modes of (1), henceforth called the “fixed spectrum” of (1), is the the spectrum of A that cannot be shifted by the decentralized output feedback laws $u_i = F_i y_i$, $i \in \{1, 2, \dots, k\}$. That is, if $A \in \mathbb{R}^{n \times n}$, $B_i \in \mathbb{R}^{n \times m_i}$, $C_i \in \mathbb{R}^{l_i \times n}$, the *fixed spectrum* of (1), written Λ_{fixed} , is precisely

$$\Lambda_{\text{fixed}} = \bigcap_{F_i \in \mathbb{R}^{m_i \times l_i}} \Lambda \left(A + \sum_{i=1}^k B_i F_i C_i \right)$$

where $\Lambda(\cdot)$ denotes the spectrum. Since the F_i can be zero, it is clear that the fixed spectrum is a subset of the spectrum of A . It is possible that the fixed spectrum is an empty set, in which case it is said that the system has no fixed spectrum.

Wang and Davison showed that Λ_{fixed} is contained in the closed-loop spectrum of the system which results when any given finite dimensional LTI decentralized controls are applied to (1). Thus Λ_{fixed} must be a stable spectrum if decentralized stabilization is to be achieved. Wang and Davison were also able to show that the stability of Λ_{fixed} is sufficient for decentralized stabilization with linear dynamic controllers. Not surprisingly, the notion of a fixed

spectrum also arises in connection with the decentralized spectrum assignment problem treated in [2], [3]. In particular it is known that a necessary and sufficient condition for “free” assignability of a closed-loop spectrum with finite dimensional LTI decentralized controllers is that there is no fixed spectrum [1], [2]. However, it should be noted that unlike the centralized case, free spectrum assignability in the decentralized case presumes that the overall spectrum admits a suitable partition into a finite number of symmetric sets, the partition being determined by the strongly connected components in a suitably defined graph of (1) [2]. It is clear from the preceding that Λ_{fixed} plays a central role in both the decentralized stabilization and decentralized spectrum assignment problems. Accordingly many characterizations of Λ_{fixed} exist [4]. A certain subset of a fixed spectrum can be eliminated by time-varying decentralized controllers [5], [6], sampling strategies [7], [8], or other techniques [9], [10]. The subset of a fixed spectrum which cannot be eliminated by any (including nonlinear) decentralized controllers are characterized in [11]. An explicit necessary and sufficient matrix-algebraic condition for a complex number λ to be in the fixed spectrum of (1) is derived in [12], using matrix pencils and matrix nets [13]. An equivalent algebraic condition is established in [14], [15]. Equivalent graph-theoretic criteria for the existence of a fixed spectrum are developed in [16]. Frequency domain characterizations of a fixed spectrum are presented in [17]–[22]. In particular, the characterization of a fixed spectrum in [12] is quite fundamental and inspires a number of later work [4], [11], [14]–[16].

Let $\mathbf{k} \triangleq \{1, 2, \dots, k\}$. Suppose $\mathcal{S} = \{i_1, i_2, \dots, i_s\} \subset \mathbf{k}$ with $i_1 < i_2 < \dots < i_s$, the complement of \mathcal{S} in \mathbf{k} is denoted by $\mathbf{k} - \mathcal{S} = \{j_1, j_2, \dots, j_{k-s}\}$ with $j_1 < j_2 < \dots < j_{k-s}$. Let

$$B_{\mathcal{S}} \triangleq [B_{i_1} \ B_{i_2} \ \dots \ B_{i_s}], \quad C_{\mathbf{k}-\mathcal{S}} \triangleq \begin{bmatrix} C_{j_1} \\ C_{j_2} \\ \vdots \\ C_{j_{k-s}} \end{bmatrix}$$

The result in [12] is stated below.

Proposition 1: [12] A k -channel linear system $\{A, B_i, C_i; k\}$ has $\lambda \in \Lambda(A)$ in its fixed spectrum if and only if $\exists \mathcal{S} \subset \mathbf{k}$ such that

$$\text{rank} \begin{bmatrix} \lambda I_n - A & B_{\mathcal{S}} \\ C_{\mathbf{k}-\mathcal{S}} & \mathbf{0} \end{bmatrix} < n \quad (2)$$

Although originally proved with techniques like matrix pencils and matrix nets [12], [13], Proposition 1 reveals an insight that whether a multi-channel linear system has a fixed

*This work was supported by National Science Foundation grant n. 1607101.00 and US Air Force grant n. FA9550-16-1-0290.

¹ F. Liu and A. S. Morse are with the Department of Electrical Engineering, Yale University, New Haven, CT, USA. {fengjiao.liu, as.morse}@yale.edu

spectrum is indeed a combinatorial problem. By exploiting this fact, a new and transparent proof of Proposition 1 using matroid theory is presented in this paper.

Roughly speaking, a matroid is a mathematical structure that generalizes the notion of linear independence in vector spaces. It is a handy tool for studying combinatorial properties of matrices.

II. PROOF

A formal proof of Proposition 1 is given in this section, for which some concepts and three lemmas are needed.

Lemma 1: Let matrices $A \in \mathbb{C}^{n \times n}$, $B \in \mathbb{C}^{n \times m}$, and $C \in \mathbb{C}^{l \times n}$. Then $\text{rank} \begin{bmatrix} A & B \\ C & \mathbf{0} \end{bmatrix} < n$ if and only if $\text{rank}(A + BE + KC) < n$ for any matrices $E \in \mathbb{C}^{m \times n}$ and $K \in \mathbb{C}^{n \times l}$.

Proof of Lemma 1: (Necessity) If $\text{rank} \begin{bmatrix} A & B \\ C & \mathbf{0} \end{bmatrix} < n$, then $\text{rank} \begin{bmatrix} A + BE + KC & B \\ C & \mathbf{0} \end{bmatrix} < n$ for any matrices $E \in \mathbb{C}^{m \times n}$ and $K \in \mathbb{C}^{n \times l}$, as the rank of a matrix remains unchanged under elementary row and column operations. So $\text{rank}(A + BE + KC) < n$ for any E and K .

(Sufficiency) Suppose $\text{rank} \begin{bmatrix} A & B \\ C & \mathbf{0} \end{bmatrix} \geq n$, by elementary column operations, $\exists E \in \mathbb{C}^{m \times n}$ such that $\text{rank} \begin{bmatrix} A + BE \\ C \end{bmatrix} = n$. Similarly, by elementary row operations, $\exists K \in \mathbb{C}^{n \times l}$ such that $\text{rank}(A + BE + KC) = n$. This completes the proof of Lemma 1. ■

Let M be a parameterized matrix whose entries are linear combinations of algebraically independent scalar parameters. The *generic rank* of M , denoted by $\text{grank } M$, is the maximum rank of M that can be achieved as the parameters vary over the entire parameter space. It is generic in the sense that it is achievable by any parameter values in the complement of a proper algebraic set in the parameter space.

For some $d \in \mathbb{N}$, we are given column vectors $w_1, w_2, \dots, w_d \in \mathbb{C}^{n_1}$ and row vectors $r_1, r_2, \dots, r_d \in \mathbb{C}^{1 \times n_2}$. Let $\mathbf{d} \triangleq \{1, 2, \dots, d\}$. Suppose $\mathcal{S} = \{i_1, i_2, \dots, i_s\} \subset \mathbf{d}$ with $i_1 < i_2 < \dots < i_s$, and $\mathbf{d} - \mathcal{S} = \{j_1, j_2, \dots, j_{d-s}\}$ with $j_1 < j_2 < \dots < j_{d-s}$, let

$$w_{\mathcal{S}} = [w_{i_1} \ w_{i_2} \ \dots \ w_{i_s}], \quad r_{\mathbf{d}-\mathcal{S}} = \begin{bmatrix} r_{j_1} \\ r_{j_2} \\ \vdots \\ r_{j_{d-s}} \end{bmatrix} \quad (3)$$

Let $\mathcal{P} = \{(w_i, r_i) \mid i \in \mathbf{d}\}$ be the set of d pairs of vectors. For $\mathcal{I} \subset \mathcal{P}$, let $\{w_i \mid (w_i, r_i) \in \mathcal{I}\}$ be the set of column vectors appearing in \mathcal{I} ; let $\{r_i \mid (w_i, r_i) \in \mathcal{I}\}$ be the set of row vectors appearing in \mathcal{I} . Note that

$$|\mathcal{I}| = |\{w_i \mid (w_i, r_i) \in \mathcal{I}\}| = |\{r_i \mid (w_i, r_i) \in \mathcal{I}\}|$$

where $|\mathcal{I}|$ is the number of vector pairs in \mathcal{I} . A nonempty subset $\mathcal{I} \subset \mathcal{P}$ is *jointly independent* if $\{w_i \mid (w_i, r_i) \in \mathcal{I}\}$ and $\{r_i \mid (w_i, r_i) \in \mathcal{I}\}$ are both linearly independent sets. Let $\mathcal{J}\{w_i, r_i; d\}$ be the set of all jointly independent subsets of \mathcal{P} .

\mathcal{P} . The following proposition summarizes Lemma 1 (derived from the matroid intersection theorem [23]) and Lemma 2 in [24].

Proposition 2: [24] For some $d \in \mathbb{N}$, given column vectors $w_1, w_2, \dots, w_d \in \mathbb{C}^{n_1}$, row vectors $r_1, r_2, \dots, r_d \in \mathbb{C}^{1 \times n_2}$, and algebraically independent scalar parameters p_1, p_2, \dots, p_d , the following equation holds.

$$\begin{aligned} \text{grank} \left(\sum_{i \in \mathbf{d}} w_i p_i r_i \right) &= \max_{\mathcal{I} \in \mathcal{J}\{w_i, r_i; d\}} |\mathcal{I}| \\ &= \min_{\mathcal{S} \subset \mathbf{d}} (\text{rank } w_{\mathcal{S}} + \text{rank } r_{\mathbf{d}-\mathcal{S}}) \end{aligned}$$

Although Proposition 2 was initially developed for real vectors and parameters, the same proof applies to complex vectors and parameters without change. Therefore a proof of Proposition 2 will not be given here. The next lemma extends Proposition 2 to a more general case.

For some $d \in \mathbb{N}$, we are given complex matrices W_1, W_2, \dots, W_d , each of which has n_1 rows and an arbitrary number of columns, and complex matrices R_1, R_2, \dots, R_d , each of which has an arbitrary number of rows and n_2 columns. Suppose $\mathcal{S} = \{i_1, i_2, \dots, i_s\} \subset \mathbf{d}$ with $i_1 < i_2 < \dots < i_s$, and $\mathbf{d} - \mathcal{S} = \{j_1, j_2, \dots, j_{d-s}\}$ with $j_1 < j_2 < \dots < j_{d-s}$, let

$$W_{\mathcal{S}} = [W_{i_1} \ W_{i_2} \ \dots \ W_{i_s}], \quad R_{\mathbf{d}-\mathcal{S}} = \begin{bmatrix} R_{j_1} \\ R_{j_2} \\ \vdots \\ R_{j_{d-s}} \end{bmatrix}$$

For each $i \in \mathbf{d}$, let $\text{col}(W_i)$ denote the set of column vectors of W_i and let $\text{row}(R_i)$ denote the set of row vectors of R_i . Let $\mathcal{P} = \{(w, r) \mid w \in \text{col}(W_i), r \in \text{row}(R_i), i \in \mathbf{d}\}$ be the set of all vector pairs taken from matrices with the same indices. For $\mathcal{I} \subset \mathcal{P}$, let $\{w \mid (w, r) \in \mathcal{I}, r \in \text{row}(R_{\mathbf{d}})\}$ be the set of column vectors appearing in \mathcal{I} , with any common vectors repeated; let $\{r \mid (w, r) \in \mathcal{I}, w \in \text{col}(W_{\mathbf{d}})\}$ be the set of row vectors appearing in \mathcal{I} , with any common vectors repeated. That is, if a column vector w appears exactly n_w times in \mathcal{I} , the set $\{w \mid (w, r) \in \mathcal{I}, r \in \text{row}(R_{\mathbf{d}})\}$ contains n_w copies of w . Note that

$$\begin{aligned} |\mathcal{I}| &= |\{w \mid (w, r) \in \mathcal{I}, r \in \text{row}(R_{\mathbf{d}})\}| \\ &= |\{r \mid (w, r) \in \mathcal{I}, w \in \text{col}(W_{\mathbf{d}})\}| \end{aligned}$$

A nonempty subset $\mathcal{I} \subset \mathcal{P}$ is *jointly independent* if $\{w \mid (w, r) \in \mathcal{I}, r \in \text{row}(R_{\mathbf{d}})\}$ and $\{r \mid (w, r) \in \mathcal{I}, w \in \text{col}(W_{\mathbf{d}})\}$ are both linearly independent sets. Let $\mathcal{J}\{W_i, R_i; d\}$ be the set of all jointly independent subsets of \mathcal{P} .

Lemma 2: For some $d \in \mathbb{N}$, given complex matrices W_1, W_2, \dots, W_d , each of which has n_1 rows and an arbitrary number of columns, complex matrices R_1, R_2, \dots, R_d , each of which has an arbitrary number of rows and n_2 columns, and parameterized matrices P_1, P_2, \dots, P_d of appropriate sizes, whose entries are algebraically independent thus are modeled by distinct parameters, the following equation

holds.

$$\begin{aligned} \text{grank} \left(\sum_{i \in \mathbf{d}} W_i P_i R_i \right) &= \max_{\mathcal{I} \in \mathcal{J}\{W_i, R_i; d\}} |\mathcal{I}| \\ &= \min_{S \subset \bar{\mathbf{d}}} (\text{rank } W_S + \text{rank } R_{\bar{\mathbf{d}}-S}) \end{aligned}$$

As Proposition 2 is proved directly from the matroid intersection theorem, it is natural to ask whether Lemma 2 can be proved in the same way. Unfortunately, the answer is no. In order to do that, we may need a more general notion of matroid which allows the rank of an element to be greater than 1.

Proof of Lemma 2: For each $t \in \mathbf{d}$, suppose W_t is of size $n_1 \times \alpha_t$, R_t is of size $\beta_t \times n_2$, and P_t is of size $\alpha_t \times \beta_t$.

Note that $W_t P_t R_t = \sum_{i=1}^{\alpha_t} \sum_{j=1}^{\beta_t} w_t^i p_t^{ij} r_t^j$, where $w_t^i \in \mathbb{C}^{n_1}$ is the i th column of W_t , p_t^{ij} is the ij th entry of P_t , and $r_t^j \in \mathbb{C}^{1 \times n_2}$ is the j th row of R_t . Recall that $\mathcal{P} = \{(w, r) \mid w \in \text{col}(W_t), r \in \text{row}(R_t), t \in \mathbf{d}\}$, now let $\bar{d} = |\mathcal{P}| = \sum_{t \in \mathbf{d}} \alpha_t \beta_t$.

Let the entries of the P_t , $t \in \mathbf{d}$, be denoted by p_1 through $p_{\bar{d}}$ and let the vector pairs in \mathcal{P} be labeled 1 through \bar{d} , i.e., $\mathcal{P} = \{(w_i, r_i) \mid i \in \bar{\mathbf{d}}\}$, where $\bar{\mathbf{d}} \triangleq \{1, 2, \dots, \bar{d}\}$. So $\mathcal{J}\{w_i, r_i; \bar{d}\} = \mathcal{J}\{W_i, R_i; d\}$. Thus

$$\sum_{t \in \mathbf{d}} W_t P_t R_t = \sum_{t=1}^d \sum_{i=1}^{\alpha_t} \sum_{j=1}^{\beta_t} w_t^i p_t^{ij} r_t^j = \sum_{i \in \bar{\mathbf{d}}} w_i p_i r_i$$

By Proposition 2,

$$\begin{aligned} \text{grank} \left(\sum_{i \in \mathbf{d}} W_i P_i R_i \right) &= \text{grank} \left(\sum_{i \in \bar{\mathbf{d}}} w_i p_i r_i \right) \\ &= \max_{\mathcal{I} \in \mathcal{J}\{w_i, r_i; \bar{d}\}} |\mathcal{I}| \\ &= \max_{\mathcal{I} \in \mathcal{J}\{W_i, R_i; d\}} |\mathcal{I}| \end{aligned}$$

As \mathcal{P} contains all possible combinations of W_t 's columns and R_t 's rows for each $t \in \mathbf{d}$, it is not hard to see that $\forall S \subset \bar{\mathbf{d}}, \forall t \in \mathbf{d}$, either all columns of W_t appear in w_S or all rows of R_t appear in $r_{\bar{\mathbf{d}}-S}$. By Proposition 2, there exists $S_0 \subset \bar{\mathbf{d}}$ such that $\text{rank } w_{S_0} + \text{rank } r_{\bar{\mathbf{d}}-S_0} = \min_{S \subset \bar{\mathbf{d}}} (\text{rank } w_S + \text{rank } r_{\bar{\mathbf{d}}-S})$. Now if all columns of W_t appear in w_{S_0} , remove rows of R_t from $r_{\bar{\mathbf{d}}-S_0}$ if any; else remove columns of W_t from w_{S_0} if any. Repeat the removal process for all $t \in \mathbf{d}$, and denote the resultant matrices by $w_{S_0}^-$ and $r_{\bar{\mathbf{d}}-S_0}^-$ respectively. Then $\exists S_1 \subset \mathbf{d}$ such that $w_{S_0}^- = W_{S_1}$ up to a permutation of columns and $r_{\bar{\mathbf{d}}-S_0}^- = R_{\bar{\mathbf{d}}-S_1}$ up to a permutation of rows. So

$$\begin{aligned} &\min_{S \subset \bar{\mathbf{d}}} (\text{rank } w_S + \text{rank } r_{\bar{\mathbf{d}}-S}) \\ &\leq \min_{S \subset \bar{\mathbf{d}}} (\text{rank } W_S + \text{rank } R_{\bar{\mathbf{d}}-S}) \\ &\leq \text{rank } W_{S_1} + \text{rank } R_{\bar{\mathbf{d}}-S_1} \\ &= \text{rank } w_{S_0}^- + \text{rank } r_{\bar{\mathbf{d}}-S_0}^- \\ &\leq \text{rank } w_{S_0} + \text{rank } r_{\bar{\mathbf{d}}-S_0} \\ &= \min_{S \subset \bar{\mathbf{d}}} (\text{rank } w_S + \text{rank } r_{\bar{\mathbf{d}}-S}) \end{aligned} \quad (4)$$

It implies that

$$\begin{aligned} \text{grank} \left(\sum_{i \in \mathbf{d}} W_i P_i R_i \right) &= \text{grank} \left(\sum_{i \in \bar{\mathbf{d}}} w_i p_i r_i \right) \\ &= \min_{S \subset \bar{\mathbf{d}}} (\text{rank } w_S + \text{rank } r_{\bar{\mathbf{d}}-S}) \quad \text{By Proposition 2} \\ &= \min_{S \subset \mathbf{d}} (\text{rank } W_S + \text{rank } R_{\bar{\mathbf{d}}-S}) \quad \text{By (4)} \end{aligned}$$

This completes the proof of Lemma 2. \blacksquare

Let $M \in \mathbb{C}^{n_1 \times n_2}$ be a complex matrix of rank $t \leq \min\{n_1, n_2\}$. By rank decomposition, there exist a full column rank matrix $W \in \mathbb{C}^{n_1 \times t}$ and a full row rank matrix $R \in \mathbb{C}^{t \times n_2}$ such that $M = WR$. In another word, $M = \sum_{i=1}^t w_i r_i$, where $w_i \in \mathbb{C}^{n_1}$ is the i th column of W and $r_i \in \mathbb{C}^{1 \times n_2}$ is the i th row of R . Let $\mathbf{t} \triangleq \{1, 2, \dots, t\}$. It is easy to see that $\{w_i \mid i \in \mathbf{t}\}$ and $\{r_i \mid i \in \mathbf{t}\}$ are both linearly independent sets.

Lemma 3: Suppose a complex matrix $M_{n_1 \times n_2} = \sum_{i=1}^t w_i r_i$, where $w_i \in \mathbb{C}^{n_1}$ are column vectors, $r_i \in \mathbb{C}^{1 \times n_2}$ are row vectors, $\{w_1, w_2, \dots, w_t\}$ and $\{r_1, r_2, \dots, r_t\}$ are both linearly independent sets. For some $d \geq t$, given column vectors $w_{t+1}, w_{t+2}, \dots, w_d \in \mathbb{C}^{n_1}$, row vectors $r_{t+1}, r_{t+2}, \dots, r_d \in \mathbb{C}^{1 \times n_2}$, and algebraically independent scalar parameters p_1, p_2, \dots, p_d , the following equation holds.

$$\text{grank} \left(M + \sum_{i=t+1}^d w_i p_i r_i \right) = \text{grank} \left(\sum_{i=1}^d w_i p_i r_i \right) \quad (5)$$

Proof of Lemma 3: By the definition of generic rank, the left-hand side of equation (5) is less than or equal to the right-hand side, so it suffices to prove the other direction. There exists $S_0 \subset \mathbf{d}$ such that $\mathcal{I}_0 = \{(w_i, r_i) \mid i \in S_0\}$ is a jointly independent subset of $\{(w_i, r_i) \mid i \in \mathbf{d}\}$ with the maximum cardinality, i.e., $\max_{\mathcal{I} \in \mathcal{J}\{w_i, r_i; d\}} |\mathcal{I}| = |\mathcal{I}_0|$.

First, it is claimed that equation (5) holds when $\mathbf{d} - \mathbf{t} \subset S_0$. Now suppose the claim is true and consider the general case when $\mathbf{d} - \mathbf{t} \not\subset S_0$. Let $S_1 = S_0 \cap (\mathbf{d} - \mathbf{t})$. We have

$$\begin{aligned} &\text{grank} \left(\sum_{i=1}^d w_i p_i r_i \right) \\ &= |\mathcal{I}_0| = \text{grank} \left(\sum_{i \in S_0} w_i p_i r_i \right) \quad \text{By Proposition 2} \\ &= \text{grank} \left(\sum_{i \in \mathbf{t} \cup S_1} w_i p_i r_i \right) \quad \text{By the definition of } \mathcal{I}_0 \\ &= \text{grank} \left(M + \sum_{i \in S_1} w_i p_i r_i \right) \quad \text{By the claim} \\ &\leq \text{grank} \left(M + \sum_{i=t+1}^d w_i p_i r_i \right) \end{aligned}$$

So equation (5) holds for the general case.

Next, the claim will be proved. Now assume $\mathbf{d} - \mathbf{t} \subset S_0$. Let $\mathbf{t}_1 = S_0 \cap \mathbf{t}$. Let $\mathbf{t}_2 = \mathbf{t} - \mathbf{t}_1 = \mathbf{d} - S_0$. As $\{w_i \mid i \in \mathbf{t}\}$

and $\{r_i | i \in \mathbf{t}\}$ are both linearly independent sets, and by the definition of \mathcal{S}_0 , it is clear that each vector in $\{w_i | i \in \mathbf{t}_1\}$ (respectively, $\{r_i | i \in \mathbf{t}_1\}$) is linearly independent with the vectors in $\{w_i | i \in \mathbf{d} - \mathbf{t}_1\}$ (respectively, $\{r_i | i \in \mathbf{d} - \mathbf{t}_1\}$). So

$$\text{grank} \left(\sum_{i=1}^d w_i p_i r_i \right) = |\mathbf{t}_1| + \text{grank} \left(\sum_{i \in \mathbf{d} - \mathbf{t}_1} w_i p_i r_i \right) \quad (6)$$

Suppose the generic rank of $\left(\sum_{i \in \mathbf{t}_2} w_i r_i + \sum_{i=t+1}^d w_i p_i r_i \right)$ is admissible by $p_i = \bar{p}_i \in \mathbb{C}$ for $i \in \mathbf{d} - \mathbf{t}$, then

$$\begin{aligned} & \text{grank} \left(M + \sum_{i=t+1}^d w_i p_i r_i \right) \\ & \geq \text{rank} \left(\sum_{i \in \mathbf{t}_1} w_i r_i + \sum_{i \in \mathbf{t}_2} w_i r_i + \sum_{i=t+1}^d w_i \bar{p}_i r_i \right) \\ & = \text{rank} \left(\sum_{i \in \mathbf{t}_1} w_i r_i \right) + \text{rank} \left(\sum_{i \in \mathbf{t}_2} w_i r_i + \sum_{i=t+1}^d w_i \bar{p}_i r_i \right) \\ & = |\mathbf{t}_1| + \text{grank} \left(\sum_{i \in \mathbf{t}_2} w_i r_i + \sum_{i=t+1}^d w_i p_i r_i \right) \end{aligned}$$

As $|\mathcal{S}_0| = |\mathbf{t}_1| + d - t$, by equation (6), $\text{grank} \left(\sum_{i \in \mathbf{d} - \mathbf{t}_1} w_i p_i r_i \right) = d - t$. Without loss of generality, assume $\mathbf{t}_1 = \{1, 2, \dots, t_1\}$. It is easy to see that

$$\mathcal{Q} \triangleq \left\{ (p_{t_1+1}, p_{t_1+2}, \dots, p_d) \in \mathbb{C}^{d-t_1} \mid \text{rank} \left(\sum_{i \in \mathbf{d} - \mathbf{t}_1} w_i p_i r_i \right) = d - t \right\}$$

is an open set in \mathbb{C}^{d-t_1} , because its complement is a closed set. Let $p_{u,v} \in \mathbb{C}^{d-t_1}$ be the vector whose first $t - t_1$ entries are $u \in \mathbb{C}$ and last $d - t$ entries are $v \in \mathbb{C}$, i.e., $p_i = u$ for $i \in \mathbf{t}_2$ and $p_j = v$ for $j \in \mathbf{d} - \mathbf{t}$. Clearly $p_{0,1} \in \mathcal{Q}$, which implies that $\exists \delta > 0$ such that $p_{\delta,1} \in \mathcal{Q}$ thus $p_{1, \frac{1}{\delta}} \in \mathcal{Q}$. That

is, $\text{grank} \left(\sum_{i \in \mathbf{t}_2} w_i r_i + \sum_{i=t+1}^d w_i p_i r_i \right) \geq d - t$. Therefore,

$$\begin{aligned} & \text{grank} \left(M + \sum_{i=t+1}^d w_i p_i r_i \right) \\ & \geq |\mathbf{t}_1| + d - t = |\mathcal{S}_0| \\ & = \text{grank} \left(\sum_{i=1}^d w_i p_i r_i \right) \end{aligned}$$

The claim is proved. This completes the proof. \blacksquare

Corollary 1: Suppose a complex matrix $M_{n_1 \times n_2} = \sum_{i=1}^t w_i r_i$, where $w_i \in \mathbb{C}^{n_1}$ are column vectors, $r_i \in \mathbb{C}^{1 \times n_2}$ are row vectors, $\{w_1, w_2, \dots, w_t\}$ and $\{r_1, r_2, \dots, r_t\}$ are both linearly independent sets. For some $d \geq t$, given complex matrices $W_{t+1}, W_{t+2}, \dots, W_d$, each of which

has n_1 rows and an arbitrary number of columns, complex matrices $R_{t+1}, R_{t+2}, \dots, R_d$, each of which has an arbitrary number of rows and n_2 columns, algebraically independent scalar parameters p_1, p_2, \dots, p_t , and parameterized matrices $P_{t+1}, P_{t+2}, \dots, P_d$ of appropriate sizes, whose entries are algebraically independent thus are modeled by distinct parameters, the following equation holds.

$$\begin{aligned} & \text{grank} \left(M + \sum_{i=t+1}^d W_i P_i R_i \right) \\ & = \text{grank} \left(\sum_{i=1}^t w_i p_i r_i + \sum_{i=t+1}^d W_i P_i R_i \right) \end{aligned}$$

Proof of Corollary 1: For each $s \in \mathbf{d} - \mathbf{t}$, suppose W_s is of size $n_1 \times \alpha_s$, R_s is of size $\beta_s \times n_2$, and P_s is of size $\alpha_s \times \beta_s$. Then

$$\begin{aligned} & \text{grank} \left(M + \sum_{s=t+1}^d W_s P_s R_s \right) \\ & = \text{grank} \left(M + \sum_{s=t+1}^d \sum_{i=1}^{\alpha_s} \sum_{j=1}^{\beta_s} w_s^i p_s^{ij} r_s^j \right) \\ & = \text{grank} \left(\sum_{s=1}^t w_s p_s r_s + \sum_{s=t+1}^d \sum_{i=1}^{\alpha_s} \sum_{j=1}^{\beta_s} w_s^i p_s^{ij} r_s^j \right) \\ & = \text{grank} \left(\sum_{s=1}^t w_s p_s r_s + \sum_{s=t+1}^d W_s P_s R_s \right) \end{aligned}$$

where $w_s^i \in \mathbb{C}^{n_1}$ is the i th column of W_s , p_s^{ij} is the ij th entry of P_s , and $r_s^j \in \mathbb{C}^{1 \times n_2}$ is the j th row of R_s . \blacksquare

Proof of Proposition 1: (Sufficiency) If $\exists \mathcal{S} \subset \mathbf{k}$ such that (2) holds, by Lemma 1,

$$\text{rank} (\lambda I_n - A - B_{\mathcal{S}} E_{\mathcal{S}} - E_{\mathbf{k}-\mathcal{S}} C_{\mathbf{k}-\mathcal{S}}) < n$$

for any matrices $E_{\mathcal{S}} \in \mathbb{R}^{m_{\mathcal{S}} \times n}$ and $E_{\mathbf{k}-\mathcal{S}} \in \mathbb{R}^{n \times l_{\mathbf{k}-\mathcal{S}}}$, where $m_{\mathcal{S}} \triangleq \sum_{i \in \mathcal{S}} m_i$ and $l_{\mathbf{k}-\mathcal{S}} \triangleq \sum_{j \in \mathbf{k}-\mathcal{S}} l_j$. It implies that

$$\text{rank} \left(\lambda I_n - A - \sum_{i \in \mathbf{k}} B_i F_i C_i \right) < n$$

for any matrices $F_i \in \mathbb{R}^{m_i \times l_i}$, $i \in \mathbf{k}$. Thus λ is in the fixed spectrum of system $\{A, B_i, C_i; k\}$.

(Necessity) If λ is in the fixed spectrum of system $\{A, B_i, C_i; k\}$,

$$\text{rank} \left(\lambda I_n - A - \sum_{i \in \mathbf{k}} B_i F_i C_i \right) < n$$

for any matrices $F_i \in \mathbb{R}^{m_i \times l_i}$, $i \in \mathbf{k}$. Suppose $\text{rank} (A - \lambda I_n) = t < n$, then there exist $w_i \in \mathbb{C}^n$ and $r_i \in \mathbb{C}^{1 \times n}$, $i \in \mathbf{t}$, such that $\{w_i | i \in \mathbf{t}\}$ and $\{r_i | i \in \mathbf{t}\}$ are both linearly independent sets, and $A - \lambda I_n = \sum_{i=1}^t w_i r_i$. By Corollary 1,

$$\text{grank} \left(\sum_{i=1}^t w_i p_i r_i + \sum_{j=1}^k B_j F_j C_j \right) < n$$

By Lemma 2, $\exists \mathcal{S}_1 \subset \mathbf{t}, \mathcal{S}_2 \subset \mathbf{k}$ such that

$$\text{rank} [w_{\mathcal{S}_1} \ B_{\mathcal{S}_2}] + \text{rank} \begin{bmatrix} r_{\mathbf{t}-\mathcal{S}_1} \\ C_{\mathbf{k}-\mathcal{S}_2} \end{bmatrix} < n$$

So

$$\text{rank} [w_{\mathcal{S}_1} \ B_{\mathcal{S}_2}] \begin{bmatrix} r_{\mathcal{S}_1} \\ E_{\mathcal{S}_2} \end{bmatrix} + \text{rank} [w_{\mathbf{t}-\mathcal{S}_1} \ E_{\mathbf{k}-\mathcal{S}_2}] \begin{bmatrix} r_{\mathbf{t}-\mathcal{S}_1} \\ C_{\mathbf{k}-\mathcal{S}_2} \end{bmatrix} < n$$

for any matrices $E_{\mathcal{S}_2}$ and $E_{\mathbf{k}-\mathcal{S}_2}$ of appropriate sizes. As $A - \lambda I_n = w_{\mathcal{S}_1} r_{\mathcal{S}_1} + w_{\mathbf{t}-\mathcal{S}_1} r_{\mathbf{t}-\mathcal{S}_1}$, it implies that

$$\text{rank} (A - \lambda I_n + B_{\mathcal{S}_2} E_{\mathcal{S}_2} + E_{\mathbf{k}-\mathcal{S}_2} C_{\mathbf{k}-\mathcal{S}_2}) < n$$

for any matrices $E_{\mathcal{S}_2}$ and $E_{\mathbf{k}-\mathcal{S}_2}$. By Lemma 1, (2) holds for $\mathcal{S}_2 \subset \mathbf{k}$. ■

III. CONCLUDING REMARKS

This paper utilizes matroid theory to simplify the proof of the algebraic condition derived in [12] for the existence of a fixed spectrum in a multi-channel linear system.

ACKNOWLEDGEMENT

The authors would like to thank Brian D. O. Anderson for suggesting the problem which this paper is addressed.

REFERENCES

- [1] S.-H. Wang and E. J. Davison, "On the stabilization of decentralized control systems," *IEEE Trans. Autom. Control*, vol. 18, no. 5, pp. 473–478, 1973.
- [2] J. P. Corfmat and A. S. Morse, "Decentralized control of linear multivariable systems," *Automatica*, vol. 12, no. 5, pp. 479–495, 1976.
- [3] M. Vidyasagar and N. Viswanadham, "Algebraic characterization of decentralized fixed modes and pole assignment," in *Proc. IEEE Conf. Decision Control*, Orlando, FL, USA, 1982, pp. 501–505.
- [4] Z. Gong and M. Aldeen, "On the characterization of fixed modes in decentralized control," *IEEE Trans. Autom. Control*, vol. 37, no. 7, pp. 1046–1050, 1992.
- [5] B. D. O. Anderson and J. B. Moore, "Time-varying feedback laws for decentralized control," *IEEE Trans. Autom. Control*, vol. 26, no. 5, pp. 1133–1139, 1981.
- [6] J. L. Willems, "Time-varying feedback for the stabilization of fixed modes in decentralized control systems," *Automatica*, vol. 25, no. 1, pp. 127–131, 1989.
- [7] U. Ozguner and E. J. Davison, "Sampling and decentralized fixed modes," in *Proc. Amer. Control Conf.*, Boston, MA, USA, 1985, pp. 257–262.
- [8] J. L. Willems, "Elimination of fixed modes in decentralized systems by means of sampling," *Syst. & Control Lett.*, vol. 10, no. 1, pp. 1–8, 1988.
- [9] J. Lavaei and A. G. Aghdam, "Elimination of fixed modes by means of high-performance constrained periodic control," in *Proc. IEEE Conf. Decision Control*, San Diego, CA, USA, 2006, pp. 4441–4447.
- [10] S. S. Stanković and D. D. Šiljak, "Stabilization of fixed modes in expansions of lti systems," *Syst. & Control Lett.*, vol. 57, no. 4, pp. 365–370, 2008.
- [11] Z. Gong and M. Aldeen, "Stabilization of decentralized control systems," *J. Math. Systems Estim. Control*, vol. 7, pp. 111–114, 1997.
- [12] B. D. O. Anderson and D. J. Clements, "Algebraic characterization of fixed modes in decentralized control," *Automatica*, vol. 17, no. 5, pp. 703–712, 1981.
- [13] B. D. O. Anderson and H.-M. Hong, "Structural controllability and matrix nets," *Int. J. Control*, vol. 35, no. 3, pp. 397–416, 1982.
- [14] Z. Gong and M. Aldeen, "Equivalence of characterizations of fixed modes," *Syst. & Control Lett.*, vol. 22, no. 1, pp. 61–69, 1994.
- [15] W. Q. Liu, K. L. Teo, and W. Y. Yan, "Comments on the characterizations of fixed modes for decentralized control systems," *J. Franklin Inst.*, vol. 333, no. 1, pp. 31–40, 1996.
- [16] K. Reinschke, "Graph-theoretic characterization of fixed modes in centralized and decentralized control," *Int. J. Control*, vol. 39, no. 4, pp. 715–729, 1984.
- [17] H. Seraji, "On fixed modes in decentralized control systems," *Int. J. Control*, vol. 35, no. 5, pp. 775–784, 1982.
- [18] B. D. O. Anderson, "Transfer function matrix description of decentralized fixed modes," *IEEE Trans. Autom. Control*, vol. 27, no. 6, pp. 1176–1182, 1982.
- [19] M. Vidyasagar and N. Viswanadham, "Algebraic characterization of decentralized fixed modes," *Syst. & Control Lett.*, vol. 3, no. 2, pp. 69–72, 1983.
- [20] X. Xie and Y. Yang, "Frequency domain characterization of decentralized fixed modes," *IEEE Trans. Autom. Control*, vol. 31, no. 10, pp. 952–955, 1986.
- [21] X. Xu, Y. Xi, and Z. Zhang, "Further studies on algebraic characterization of fixed modes," in *Proc. IEEE Conf. Decision Control*, Athens, Greece, 1986, pp. 2195–2199.
- [22] J. Kong and J. H. Seo, "Graph-theoretic characterization of fixed modes in frequency domain," *Automatica*, vol. 32, no. 7, pp. 1057–1060, 1996.
- [23] K. Murota, *Matrices and Matroids for Systems Analysis*. Berlin, Germany: Springer-Verlag, 2010.
- [24] F. Liu and A. S. Morse, "A graphical characterization of structurally controllable linear systems with dependent parameters," *IEEE Trans. Autom. Control*, 2019, to appear.